
Article

Distribution of dissolved nitrogen compounds in the water column of a meromictic subarctic lake

Taisiya Y. Vorobyeva¹, Anna A. Chupakova¹, Artem V. Chupakov¹, Svetlana A. Zabelina¹, Olga Y. Moreva¹, and Oleg S. Pokrovsky^{1,2*}

¹ N. Laverov Federal Center for Integrated Arctic Research of the Ural Branch of the Russian Academy of Sciences, 23 Severnoi Dviny Emb., 163000 Arkhangelsk, Russia ; vtais@yandex.ru

² Géosciences Environnement Toulouse (GET), UMR 5563, CNRS-OMP-Université Toulouse, 14 Avenue Edouard Belin, 31400 Toulouse, France; oleg.pokrovsky@get.omp.eu

* Correspondence: oleg.pokrovsky@get.omp.eu

Abstract: In order to better understand the biogeochemical cycle of nitrogen in meromictic lakes, which can serve as a model for past aquatic environments, we measured dissolved concentrations of nitrate, nitrite, ammonium and organic nitrogen in deep (39 m maximal depth) subarctic Lake Svetloe (NW Russia). The lake is a rare type of freshwater meromictic water body with high concentrations of methane, ferrous iron, manganese and low concentrations of sulfates and sulfides in the monimolimnion. In the oligotrophic mixolimnion, the concentration of mineral forms of nitrogen decreased in summer compared to winter, likely due to phytoplankton bloom. The decomposition of the bulk of organic matter occurs under microaerophilic/anaerobic conditions of the chemocline and accompanied by the accumulation of nitrogen in the form of N-NH₄ in the monimolimnion. We revealed a strong relationship between methane and nitrogen cycles in the chemocline and monimolimnion horizons. The nitrate concentrations in Lake Svetloe varied in the range from 9 to 13 μM throughout the water column. This fact is rare for meromictic lakes, where nitrate concentrations up to 13 μM are found in the monimolimnion zone down to the bottom layers. We hypothesize, in accord with available data for other stratified lakes, that under conditions of high concentrations of manganese and ammonium at the boundary of redox conditions and below, anaerobic nitrification with the formation of nitrates occurs. Overall, most of organic matter in Lake Svetloe undergoes biodegradation essentially under microaerophilic/anaerobic conditions of the chemocline and the monimolimnion. Consequently, the manifestation of the biogeochemical nitrogen cycle is expressed in these horizons in the most vivid and complex relationship with other cycles of elements.

Keywords: nitrogen cycle; ammonium; nitrate; nitrite; stratification; boreal; meromictic lake

1. Introduction

Nitrogen is the most important nutrient, ranking fourth among biophilic elements after carbon, oxygen, and hydrogen. The biogeochemical nitrogen cycle is one of the main ones in the biosphere [1-6]. The nitrogen cycle is an interconnected chain of reactions for the transformation of various forms of nitrogen compounds [7], the main role in which belongs to microorganisms [8-15]. The concentration of nitrogen compounds determines the water body biological productivity to a large extent. The change of the composition and the ratio of the various forms concentrations of nitrogen compounds indicate the direction of the dominant biological and biogeochemical processes, including the processes of self-purification of water bodies [2]. In addition to assessing water quality, information on the content and distribution of various forms of nitrogen is important for nutrient balance and the water chemical composition [16].

Because the nitrogen cycle in aquatic systems is strongly dependent on redox conditions, chemically stratified water bodies are particularly important for biogeochemistry of N in continental settings. Biogeochemical processes in meromictic water bodies are very

interesting from the point of view of limnology and aquatic ecology, because lakes of this type exhibit peculiar features: there is no water mixing between different layers and the anaerobic zone persists throughout the whole year. There are 3 zones in the structure of meromictic lakes: mixolimnion – a zone in which convective and wind mixing of water occurs; monimolimnion, which is an anaerobic zone, a non-mixing layer that does not have contact with the atmosphere, and a chemocline which is a layer of an abrupt change of hydrochemical characteristics at the boundary of the mixolimnion and the monimolimnion, where a complex microbial community is usually formed [17,18].

Of greatest interest in water bodies of this type are the processes occurring in the chemocline and monimolimnion of lakes. The oxygen minimum zone (OMZ), formed in the chemocline, is a site of intensive nitrogen turnover due to the influx of organic material from the overlying water layers and the formation of a large amount of organic matter in the chemocline zone. The OMZ contains unique microbial communities that use alternative electron acceptors for respiration. The OMZs conditions provide an almost complete nitrogen (N) cycle. Remineralization can occur both due to the reduction of nitrate to nitrite, and can be associated with non-assimilatory reduction of nitrate to ammonium, whereas the formation of gaseous nitrogen can occur due to heterotrophic denitrification and anammox. Many microorganisms inhabiting the oxygen minimum zones are capable of performing various functions in the nitrogen and other cycles of elements [19].

Nitrifying bacteria (both ammonium and nitrite oxidants) are present throughout the entire oxygen minimum zone. Even in waters with lowest oxygen content, chemoautotrophic activity and oxidation of nitrogen compounds are usually detected in situ. The greatest abundance of nitrifying bacteria and their greatest activity are found in the gradient region at the upper boundary of the oxygen minimum [19, 20]. A close relationship is found between nitrification and denitrification in the oxygen minimum zone where the rapid cycling of intermediate products of the nitrogen cycle, including nitrate, nitrite, and nitrogen oxide (I) occurs [20].

An increase in exports from mixolimnion and the formation of a large amount of organic matter in the chemocline (hence, respiration within this zone) leads to the depletion of dissolved oxygen, followed by other main alternative electron acceptors (nitrate, sulfate, etc.). Some of the sulfide/sulfur oxidizers can use $\text{NO}_3^-/\text{NO}_2^-$ as electron acceptors and release gaseous N_2 and/or N_2O , while the released energy is used to fix inorganic carbon via so-called chemolithoautotrophic denitrification. The latter process can also be associated with hydrogenotrophy, methanotrophy, and iron oxidation [19].

Microorganisms of the non-biogeochemical nitrogen cycle are also capable of influencing its transformation under the specific conditions of meromictic lakes. Microorganisms of the methane cycle (including both methanogenesis and methane oxidation) play an important role in the binding of global carbon and nitrogen cycles in microaerophilic and anoxic environments [21-26]. In addition to the fact that methanotrophic bacteria use methane as a source of carbon and energy, they also modify the nitrogen cycle, especially at the boundary of the aerobic – anaerobic interaction (chemocline zone) or just above or below it [27]. Methanotrophic bacteria can assimilate nitrogen as ammonia or nitrate and can compete with nitrifying bacteria for ammonia and oxygen [26, 28, 29].

Studies of small Red/Ox-stratified boreal lakes [24] have demonstrated that methanotrophs of boreal lakes can associate methane oxidation with NO_x^- reduction under hypoxic conditions. Recent studies of shallow, seasonally stratified subalpine lake [30] have shown that NH_4^+ , being the main nutrient in this aquatic ecosystem, have a positive correlation with methane oxidizing bacteria (MOB). The latter can metabolize NH_4^+ for growth. Investigations of the oxygen-stratified Lake Fohnsee (southern Germany) have shown that anaerobic methane oxidation, denitrification, and anammox can simultaneously occur in the anoxic water column [31]. Analysis of gas taken from various anaerobic environments, where a significant amount of organic matter accumulates and decomposes with the release of methane, demonstrated that free nitrogen of biological origin can constitute up to 30% of all the released gas [7]. A relationship between methane oxidation

and nitrate reduction during hypoxia has recently been shown for Gammaproteobacterium *Methylomonas denitrificans*, attributing to this species a previously overlooked role in coupling carbon and nitrogen cycles [32].

The methane cycle in Lake Svetloe is fairly well known [33–38]. The microbial communities participating in the methane cycle are represented by several genera of methanogens and methanotrophs typical of freshwater lakes [35]. Methane oxidation was found in both oxygen and anoxic conditions with a maximum in chemocline. Methanotrophic bacteria *Methylobacter* sp. and methylophilic *Methylotenera* sp., *Methylophilus* sp. showed similar profiles of relative abundance throughout the epilimnion, chemocline, and hypolimnion of Lake Svetloe [35, 36]. However, in contrast to fairly good knowledge of carbon and methane cycle in Lake Svetloe, the knowledge of nitrogen cycle remains highly limited.

This work presents the first data on the spatiotemporal dynamics of the concentrations of organic and inorganic nitrogen forms in the subarctic meromictic Lake Svetloe (NW Russia). The lake is a rare type of freshwater meromictic water bodies with high concentrations of dissolved methane and ferrous iron, manganese, and low concentrations of sulfates and sulfides in the hypolimnion [34, 36, 39, 40]. Iron-rich and sulfur-depleted meromictic lakes with conditions suitable for photoferrotrophy are considered modern analogues of the ancient Archean ocean [21, 41]. The high transparency of the waters, low water color, and the distribution of the photic layer throughout the depth create favorable conditions for the development of phytoplankton and the specific microbial community of the chemocline [33, 36]. Our specific objectives were to (i) assess vertical and seasonal dynamics of dissolved nitrogen forms and (ii) relate the spatial pattern of these concentrations to already known cycles of carbon, oxygen, methane, iron, manganese in the lake, taken into account the microbiological control on the N cycle.

2. Study Site and Methods

2.1 Site description

Lake Svetloe is located in the northern part of the boreal zone of European Russia (N 65°04.98', E 41°06.26'), 65 km NNE of Arkhangelsk and its watershed is not subjected to any direct anthropogenic influence. The maximum depth of the lake is 39 meters (Figure 1).




Figure 1. 3D model and bathymetric scheme of Lake Svetloe (scale bar in meters)

Two pronounced relief depressions conditionally divide the lake into two parts with maximum depths of 39 and 20 m (Figure 1). In accordance with the meromictic status, three layers in the water column of the lake can be distinguished: 1) mixolimnion (from 0

to 20 meters), subject to convective mixing throughout the year and exhibits aerobic conditions 2) chemocline, which is a transitional zone at a depth of 20–24 m, where microaerophilic conditions are formed; in this zone, oxygen is produced by cyanobacteria and is actively consumed by other microorganisms [36], and 3) monimolimnion (from 25 meters to the bottom) which is an anaerobic layer [36, 39]. Lake Svetloe is characterized by a predominance of autochthonous dissolved organic matter with a low content of dissolved organic carbon (83.3 to 333.3 μM , [39, 40]) and high water transparency (12 to 16 m Secchi depth).

The water in the chemocline of Lake Svetloe has faint pink color of varying intensity, due to the development of phototrophic communities inhabiting at the boundary of the aerobic and anaerobic zones. Studies of phototrophic bacteria in the communities of the chemocline zone have demonstrated that the dominant are oxygenic phototrophic cyanobacteria of the genus *Synechococcus* (maximum development at a depth of 23 meters) [33, 36], which have a positive chemotaxis to nitrogen sources [42–44] and are capable to fix molecular nitrogen [45].

2.2 Sampling and Analyses

Sampling was carried out from 2010 to 2016, and included 30 survey campaigns during all hydrological seasons. Sampling was carried out over the entire water column from the surface to the bottom, with a step of 1 to 6 m. Water samples were taken at the deepest point of the lake from a PVC boat from May to October, and from the ice in winter (November to April) using a pre-cleaned polycarbonate horizontal water sampler (Aquatic Research Co, ID, USA).

The nitrogen forms (N-NO₂, N-NO₃, N-NH₄, TN) determinations were based on colorimetric assays [40, 46]. The indophenol blue method was used to measure ammonium (N-NH₄), relative error up to 12%, detection limit 1 μM . A spectrophotometric method employing sulfanilamide and N-(1-naphthyl)-ethylenediamine dihydrochloride was used for the analysis of nitrite (N-NO₂), relative error up to 18%, detection limit 0.03 μM . A spectrophotometric method employing sodium salicylate been chosen for the determination of nitrate (N-NO₃), relative error up to 18%, detection limit 7 μM . Total dissolved organic nitrogen (Norg) was evaluated from the difference between the total dissolved nitrogen (TN, persulfate oxidation, relative error up to 12%, detection limit 18 μM) and the total dissolved inorganic nitrogen (DIN, or the sum of nitrite, nitrate and ammonium nitrogen). All concentrations for the nitrogen compounds are given in micro mole per liter (μM) of nitrogen.

3. Results

3.1. Main hydrochemical characteristics

The main hydrological parameters of the lake are presented elsewhere [34, 39, 40]. Seasonal water temperature variation occurs to a depth of 22 meters; further below, the water temperature decreases with a gradient of about 0.7 °C/m and below 27 meters, the water temperature is in the range of 3.5–3.6 °C all year round. Specific conductivity of the surface horizons varies ranges from 150 to 250 $\mu\text{S/cm}$, whereas in the bottom horizons, it ranges from 340 to 380 $\mu\text{S/cm}$. In the mixolimnion, the pH varies from 7 to 8.4 (maximum values during periods of mass phytoplankton bloom); in the chemocline, the pH varies from 6.8 to 8; in the monimolimnion, a narrower range of variation is observed (from 6.8 to 7.6). A general decrease in pH is observed with depth. At depths of 24–25 m, there is a slight increase in pH values associated with the development of anoxygenic bacteria or cyanobacteria.

The dissolved oxygen concentration in the summer period in the thermocline (3–7 m) reaches its maximum values (over 375 μM). In the lower layers, it gradually decreases to the minimum values in the chemocline. In winter, a high oxygen content is noted throughout the mixolimnion with an abrupt increase above the chemocline. In the chemocline at depths of 21 to 24 m, a layer with microaerophilic conditions is formed,

which persists throughout the year. The absence of oxygen from a depth of 24-25 m down to the bottom was noted. In the chemocline layers, the sign of the redox potential changes. At the depth of 22-24 meters, according to estimates (our data and [36]), Eh has a negative value, -107-(-168) mV.

The lake is a meromictic waterbody of iron-manganese type. The sulfide concentration is lower than that of iron and manganese, which is quite rare among meromictic lakes [39]. The most characteristic intervals (25% -75%) of the concentrations of iron and manganese (dissolved fraction $<0.45 \mu\text{m}$) in the zones of mixolimnion, chemocline and monimolimnion are equal, respectively: Fe – 0.01-0.028, 0.6-41.7, 111-144 μM ; Mn – 0.006-0.028, 12-43, 52-54 μM . A sharp increase in the concentration of Fe and Mn also occurs when the Red/Ox conditions change.

Lake Svetloe exhibits low concentration of dissolved organic carbon (DOC) (the minimum and maximum values were 68 μM and 358 μM , respectively; the average value over the entire water column was $167 \pm 64 \mu\text{M}$). The maximum values of the DOC concentration in the mixolimnion were observed in the surface layers during the phytoplankton bloom. The minimum and maximum value of dissolved inorganic carbon (DIC) was 1725 μM and 5704 μM , respectively; the average value over the entire water column was $3407 \pm 130 \mu\text{M}$.

3.2. Nitrogen compounds

The distribution of the N-NH_4 concentration in the water column of Lake Svetloe is characterized by a sharp increase in the chemocline during all seasons (Figure 2(a), Figure 3). The maximal concentration reaches in average 144 to 177 μM in the bottom layer. In the mixolimnion, N-NH_4 concentrations rarely exceeded 1.2 μM . In summer, the concentrations of N-NH_4 in the surface layers were lower (up to 0.7 μM) than in winter.




Figure 2. Distribution of (a) ammonium, (b) nitrite nitrogen and (c) nitrate nitrogen in the water column of Lake Svetloe (in μM) averaged over 4 seasons during 2010 to 2016 years. Middle point – median; box value – percentiles (25%-75%); whisker value – min-max.




Figure 3. Seasonal dynamics of ammonium and the amount of nitrite and nitrate nitrogen in the water column of Lake Svetloe (in μM). Winter (November-April), spring (May), summer (June-August), autumn (September-October).

The distribution of N-NO_3 during all seasons demonstrated a slight decrease in concentration in the chemocline zone with a subsequent increase in the deeper layers (Figure 2(c), Figure 3). The concentration of N-NO_3 in the monimolimnion did not exceed the values achieved in the mixolimnion and varied in the range 9-13 μM . The TN concentration in the mixolimnion varied within the range of 9 to 23 μM (Figure 4).




Figure 4. Various nitrogen forms distribution in the water column of Lake Svetloe (in μM). Middle point – median; whisker value – percentiles (25%-75%).

There was a sharp increase in TN in the chemocline (by an order of magnitude). The TN concentration reached 293 μM near the bottom layer. The share of Norg in the

mixolimnion varied from 16 to 75% of TN, and in monimolimnion it ranged from 12 to 95% of TN (Figure 4)

The vertical distribution of mineral forms of nitrogen (Figure 3) demonstrated that the N-NO₃ form always predominated in the mixolimnion of Lake Svetloe. The predominance of N-NH₄ is observed from a depth of 22 meters; the content of N-NO₂ in all seasons was very low and varied within the range of 0-1.4 μM, which is 0.1-4% of the sum of inorganic forms of nitrogen (Figure 2(b))

4. Discussion

Comparative analysis of inorganic forms of nitrogen concentration in some meromictic lakes (Table 1) showed that there is a similar tendency of an increase in the content of ammonium from mixolimnion to monimolimnion with a sharp increase in N-NH₄ concentration in the chemocline. In the vertical distribution of nitrate concentration, there are similar features between lakes Svetloe (this study) and Pavin [48]. In both lakes, a decrease in the concentration of nitrate nitrogen in the chemocline zone is observed. In contrast to the compared meromictic lakes, the concentration of nitrates in the monimolimnion in the Svetloe Lake remains at the level of the mixolimnion. A similar dynamics is observed only in the meromictic Lake Zug, where nitrate is present in the anoxic hypolimnion [47].

Table 1. The content of mineral forms of nitrogen in Lake Svetloe (25%-75%) and other meromictic lakes in the world. Concentrations are given in μM.

Lake	N-NH ₄			N-NO ₂			N-NO ₃			Reference
	mix	chem	monim	mix	chem	monim	mix	chem	monim	
Lake Svetloe	0.5- 1.2	13-91	144-177	0-0.07	0.3- 0.8	0.6-1.4	5-9	5-8	9-13	this research
Lake Matano	up to 20	20-300	200-300	< 0,1	all layers < 0,1 only 90-100 m 0.1-0.2					[21]
Lake Pavin	-	-	max 389	-	-	-	max 32	-	-	[48]
Lake La Cruz	1.1-12.8	122.2	up to 3000	0.04-0.11	0	0	5-20	0	0	[18, 49]
Kabuno Bay	-	-	-	up to 1	N-(NO ₂ +NO ₃) Up to 0.5 in dry season up to 1.5			up to 0.5		[50]
Lake Kuznechikha	1.4-3.5	17.8-50	83-121	-	-	-	-	-	-	[51]
Char lake	-	-	-	-	-	-	< 0.7	-	< 0.7	[52]
Lake Chernoe	-	-	up to 265	-	-	-	-	-	-	[53]
Hall Lake	0	-	305	-	-	-	13.3	-	4.4	[18, 54]
Lake Zug			10	-	-	-	20- 25	17-18	1-5	[47]

The surface horizon of Lake Svetloe (0.5-3 m) is almost always distinguished by an increased number of heterotrophic bacteria, including ammonifying ones, and a slight excess of ammonium concentrations in comparison with the lower layers of the epilimnion, which may indicate a more active ammonification. By the content of nitrogen compounds, the mixolimnion layer can be characterized as oligotrophic, which is also confirmed by the values of the isotopic composition of suspended organic carbon: for the upper oxygen layer, the values δ¹³C_{org} of -27.2 to -30.2 ‰ are common for phytoplankton of oligotrophic freshwater lakes [36]. The microbial communities of the mixolimnion of Lake Svetloe are

typical of those for aerobic lake environments [55] and are mainly represented by microbial groups participating in the initial stages of organic matter decomposition [35, 36]. Measurements of the rate of dark CO₂ assimilation indicate a low activity of heterotrophic bacterioplankton [35, 36].

Aerobic nitrification of ammonia to nitrite and nitrate is a major process in the global nitrogen cycle. Low concentrations of N-NH₄ and high concentrations of nitrates in the mixolimnion, except for the periods of the peak of phytoplankton developments, when the concentration of nitrates can decrease below 1.6 μM. This testifies active processes of nitrification which proceed to the final compound of nitrate in the water of Lake Svetloe. Low N-NO₂ concentrations in the mixolimnion and a large amount of N-NO₃ in this layer may indicate that nitrification processes proceed to the final stage in the presence of a sufficient amount of oxygen in the mixolimnion.

The peak in the development of phytoplankton communities (maximum oxygenic phototrophic activity [35, 36]) is most often observed at a depth of 6-10 m, which is also evidenced by an increase in pH and dissolved oxygen concentration at this depth. There is also a slight increase in the concentration of organic nitrogen and a decrease of the inorganic nitrogen concentration, notably of nitrates.

The concentration of dissolved oxygen starts to decrease already at 16-17 m, reaching the minimum at 20-21 m. As a result, the intensity of aerobic processes sharply decreases and not completely mineralized organic matter enters the chemocline. In the lake mixolimnion, the suspended organic matter content and the total number of microorganisms are significantly lower than in the chemocline and monimolimnion [35, 36].

The largest amount of organic matter is formed in the chemocline zones. The major peak of microbial processes in the chemocline coincides with the local maximum of carbon isotope fractionation [35, 36]. Oxygenic phototrophic cyanobacteria, highly abundant in the chemocline, provide specific conditions for the functioning of the ecosystem. For example, the first peak of oxygenic photosynthesis is observed in the mixolimnion of the lake, and the second peak of oxygenic and anoxygenic photosynthesis is observed in the chemocline [35, 36]. In accord with previous works [56], we believe that the presence of constant stratification in such water bodies creates a "trap" for nutrients in the anaerobic zone. The organic matter formed in the chemocline enters the anaerobic layers, monimolimnion, where the anaerobic ammonification takes place.

Given that the majority of organic matter in Lake Svetloe undergoes biodegradation essentially under microaerophilic/anaerobic conditions of the chemocline and monimolimnion, the manifestation of the biogeochemical nitrogen cycle is expressed in these horizons in the most vivid and complex relationship with other cycles of elements. Due to production of a significant amount of organic matter in the chemocline [36] and additional accumulation of OM in the chemocline linked to a sharp increase in the vertical density gradient [18], a sharp increase of ammonium ions (up to 200 times) concentration occurs within this relatively thin water layer. A sharp increase in the concentration of ammonium ions in the chemocline of the lake in comparison with the mixolimnion indicates intensive ammonification processes. As a result, at the boundary of aerophilic/microaerophilic/anaerobic conditions, aerobic and facultative anaerobic ammonifiers are likely to develop intensively.

The chemocline zone can be considered as a very strong biological filter of methane. Indeed, in this zone, the proliferating methanotrophic bacteria [35, 36, 57] use nitrogen in the form of ammonia/ammonium or nitrate in the process of methane oxidation. Methanotrophic bacteria can assimilate nitrogen as ammonia or nitrate and can compete with nitrifying bacteria for ammonia and oxygen [28]. It is possible that the nitrifying bacteria development in the chemocline of Lake Svetloe is limited due to the active process of methane oxidation, since simultaneous nitrification and methane oxidation were observed only while maintaining relatively high levels of ammonium and oxygen [27]. Use of nitrate and/or suppression of the nitrification process consistent with a slight decrease in nitrate concentrations in the chemocline (Figure 2).

The oxidation of ammonia is carried out by both bacteria and archaea Crenarchaeota or Thaumarchaeota [58, 59], which can play a dominant role in the oxidation of ammonia. The distribution of ammonium-oxidizing bacteria (AOB) and ammonium-oxidizing archaea (AOA) along the water column and among the seasons is influenced by environmental factors, such as oxygen concentration [60, 61], ammonia concentration [62, 63], pH, concentrations of nitrites, nitrates, and phosphates [64].

Under conditions of low oxygen concentration, ammonium-oxidizing archaea are highly stable [65-67]. In addition, they are probably not influenced by methanotrophic bacteria, as in AOB. The presence of archaea Thaumarchaeota in the oxygen minimum zone [35] at the depth 17-22 m is consistent with sufficient ammonium concentration and the presence of oxygen. As is known, under anaerobic conditions, ammonification is slow and does not lead to significant ammonia/ammonium production. Therefore, a sharp increase in the concentration of ammonium in the monimolimnion is not observed.

There is another pathway for the formation of ammonium, the so-called dissimilatory nitrate reduction to ammonium (DNRA) process, also known as nitrate / nitrite ammonification [19, 68]. The processes of nitrate reduction to N_2 and NH_3 can occur simultaneously in water bodies. In natural environments, dissimilated nitrate reduction occurs under high ratio of ammonium nitrogen to nitrate nitrogen [7]. In Lake Svetloe, the ratio of ammonium to nitrate strongly increases from a depth of 24 meters (Figure 5), which may indicate the predominance of dissimilatory nitrate reduction over denitrification under anaerobic conditions.




Figure 5. The ratio of the concentrations of ammonium and nitrate nitrogen in the water column of Lake Svetloe. Middle point – median; box value – percentiles (25%-75%); whisker value – min-max.


The monimolimnion remains a fairly stable environment. The decomposition of the main fraction of organic matter, which occurs under the anaerobic conditions of monimolimnion, is accompanied by the accumulation of nitrogen in the $N-NH_4$ form, as evidenced by high concentrations of $N-NH_4$ in all layers regardless of season. In the monimolimnion of lakes, nitrate is usually present in minimal concentrations, but for Lake Svetloe, rather high concentrations are observed up to bottom sediments (Figure. 2).

Nitrification usually does not occur under anaerobic conditions, whereas the denitrification processes are quite pronounced. Most likely, oxygen is still present in the chemocline and monimolimnion of Lake Svetloe. Microbiological studies of Lake Svetloe [35-37] reported a high relative abundance of cyanobacteria in the chemocline zone. Cyanobacteria were also found in the monimolimnion down to the bottom sediments, but in a lower relative abundance. Oxygen produced by photosynthetic cyanobacteria can be used by nitrifying bacteria, ammonium-oxidizing bacteria or ammonium-oxidizing archaea, bacteria of the Nitrospirae family, found in small amounts in the chemocline of Lake Svetloe

[35], representatives of which carry out the complete oxidation of ammonium into nitrate (comammox).

Comammox was discovered and described relatively recently [69, 70]. After oxygen, the nitrate ion is the next preferred electron acceptor for respiration [71]. Moreover, it is the nitrate form of nitrogen that is the first preferred alternative electron acceptor among all nitrogen compounds. Consequently, nitrate ions are primarily subjected to assimilation / dissimilation under anaerobic conditions. The presence of nitrates down to the bottom horizons may indicate, as mentioned above, the presence of oxygen, which promotes the formation of nitrates.

It is also possible that under conditions of high concentrations of manganese and ammonium, anaerobic occurs with the formation of nitrates according to the reaction [72, 73]:



Usually, the oxidation of Mn to form Mn particles occurs where Mn (II) and oxygen are present. Manganese oxides fall below the redox boundary, where there are very few sulfides acting as a reducing agent, but sufficient amount of ammonium [47, 74]. Manganese reduction can occur under oxidized conditions [75, 76]; therefore, oxidation and reduction can occur simultaneously in the same water layer. For meromictic Lake Zug, it has been suggested that ammonium oxidation in the presence of manganese can be a source of nitrates thus rising their concentration up to 18 μM in the monimolimnion [47]. However, we could not unambiguously identify the driver of manganese oxidation in the anaerobic horizons of the Svetloe Lake water column. It should be noted that zero measured concentration value does not mean complete absence of a component. Oxygen can be found in concentrations below the detection limit of the method, for example, in a form not measurable by this method (eg peroxide or free radical). There may be an established balance of redox processes as a result of which the formed oxygen is immediately consumed in the reaction. As such, the scenario described above is quite possible.

Microorganisms associated with the process of nitrate/nitrite-dependent anaerobic oxidation of methane (ANME-2d) [35, 37], that is, archaea of ANME 2d clusters, including '*Candidatus Methanoperedens nitroreducens*' and bacteria of the NC10 type ('*Candidatus Methyloirabilis oxyfera*') are absent or found in very small numbers in Lake Svetloe. Apparently, nitrate / nitrite-dependent anaerobic oxidation of methane is minimal is low [21, 25].

A slight increase in nitrite in the bottom layers can occur due to the oxidation of ammonia to nitrite. In the horizons 27-35 m of Lake Svetloe, ammonia-oxidizing archaea Thaumarchaeota of the genus *Nitrosopumilus* are abundant, constituting up to 6.6% of all 16S rRNA gene sequences [35, 37]. Ammonia-oxidizing archaea are ubiquitous in marine, freshwater and terrestrial ecosystems. They are now considered to be a significant contributor to the carbon and nitrogen cycle. Representatives of this genus are autotrophs that receive energy from aerobic oxidation of ammonia to nitrite and fixation of inorganic carbon [64, 77, 78].

The permanent redoxcline and simultaneous presence of ammonia and nitrite provide stable conditions for anaerobic ammonia oxidation ("anammox") [79]. Anammox is the anaerobic oxidation of ammonium with nitrite, resulting in the formation of molecular nitrogen N_2 . Although anammox is an anaerobic process, oxygen does not appear to completely suppress it up to concentration of $\sim 13.5 \mu\text{M}$ [80]. Anammox-conducting bacteria live in many natural ecosystems usually under conditions of limited ammonia and make a significant contribution to the nitrogen cycle [81]. The anammox reaction uses NH_4^+ и NO_2^- in a stoichiometric ratio of about 1: 1 [19]. In lake ecosystems, this process has been suggested for sediments [82] or for the chemocline of several meromictic lakes: Lake Tanganyika [83], Lake Rassnitz (Germany; [84]) and Lake Lugano (Switzerland, Italy; [85]). However, for Lake Svetloe, the anammox process is the least likely, since the bacteria Planctomycetes that carry out this process [81, 86] have not been found in the lake monimolimnion [35]. It is possible that high ammonium content and low nitrite concentrations in chemocline and monimolimnion are limiting factors for their development [19, 81]. It is

also not excluded that in Lake Svetloe, anammox is additionally inhibited by methanol [80], formed during methane oxidation [87].

5. Conclusions

Analysis of the data on the spatio-temporal dynamics of the nitrogen forms concentrations in the low sulphide, ferro-manganese meromictic subarctic lake demonstrated the predominance of anaerobic processes in the nitrogen cycle. When the Red/Ox conditions change, the dominant form of mineral nitrogen changes from oxidized (NO_3) to reduced (NH_4). N-NO_3 form always predominated in the mixolimnion of Lake Svetloe, and seasonal fluctuations in nitrogen concentration were associated with phytoplankton consumption. Below depth of 22 m, the N-NH_4 predominated, whereas the concentration of N-NO_2 never exceeded 4% of the sum of in-organic forms of nitrogen. In the horizons of the chemocline and monimolimnion, there was a close relationship between the biogeochemical cycles of methane and nitrogen. This is indirectly confirmed by the local minimum of the nitrate concentration in the chemocline, where the methanotrophic bacteria could use N-NH_4 or N-NO_3 for the methane oxidation.

A peculiar feature of the nitrogen cycle in Lake Svetloe is high concentrations of nitrate in the anaerobic waters of monimolimnion. This phenomenon is possibly a consequence of the iron-manganese type of meromixia of the waterbody. Under conditions of high concentrations of manganese and ammonium, an anaerobic oxidation of ammonium with the formation of nitrates could occur, where MnO_2 acts as an oxidizing agent. This process is described in the literature for other meromictic bodies. The functioning of the nitrogen cycle in the meromictic lake Svetloe reflects the climatic features of the subarctic (long glacial season), geochemical factors of the catchment (low DOC content and high transparency), the type of meromixia of the lake (high concentrations of Fe and Mn), and the features of the carbon cycle (methanogenesis/methane oxidation). Further, more comprehensive analysis of these relationships should provide necessary information on the functioning of the ecosystem, and the biogeochemical nitrogen cycle provides such an opportunity. Further work is required in this direction, in particular, on the assessment of the content of the gas component of the nitrogen cycle such as N_2 and N_2O .

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Table S1: The various forms of nitrogen concentration and other physicochemical characteristics of the water column of Lake Svetloe in different seasons of the year. All concentrations are given in μM (for the nitrogen compounds μM nitrogen in L).

Author Contributions: Conceptualization, T.Y. Vorobyeva and O.S. Pokrovsky.; methodology A.A. Chupakova.; software, A.V. Chupakov; validation, O.S. Pokrovsky, S.A. Zabelina; formal analysis, A. A. Chupakova.; investigation, A.A. Chupakova, A.V. Chupakov, O.Y. Moreva; writing—original draft preparation, T.Y. Vorobyeva, A.A. Chupakova, A.V. Chupakov; writing—review and editing, O.S. Pokrovsky and S.A. Zabelina; supervision, O.S. Pokrovsky; project administration, T.Y. Vorobyeva.; funding acquisition, T.Y. Vorobyeva All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the State Task AAAA-A18-118012390200-5

Acknowledgments: We are grateful to Klimov S.I. for hydrological studies of Lake Svetloe.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Campbell, N.A. *Biology*, 2nd ed.; The Benjamin/Cummings Publishing Company: Redwood City, Calif., 1990.
2. Bashkin, V.N. *Biogeochemistry*; Scientific world: Moscow, Russia, 2004; 584 p. (in Russian)

3. Downing, Ja.; Mccauley, E. The nitrogen - phosphorus relationship in lakes. *Limnology And Oceanography* **1992**, 37(5), 936–945. doi: 10.4319/lo.1992.37.5.0936.
4. Burt T.P. Nitrogen Cycle. In *Encyclopedia of Ecology*, 2nd ed. Fath, B. Eds.; Elsevier, 2013; Volume 4, 2013, pp. 135–142 doi: 10.1016/B978-0-12-409548-9.00944-1
5. Gruber, N.; Galloway, J. An Earth-system perspective of the global nitrogen cycle. *Nature* **2008**, 451, 293–296. doi: 10.1038/nature06592
6. Mackenzie, F.; Ver, L.M.; Sabine, C.; Lane, M.; Lerman, A. C, N, P, S global biogeochemical cycles and modeling of global change. In *Interactions of C, N, P and S Biogeochemical Cycles and Global Change*. NATO ASI Series (Series I: Global Environmental Change); Wollast, R., Mackenzie, F.T., Chou, L. Eds.; Springer: Heidelberg, Berlin, 1993; V 4. P. 1–61. doi: 10.1007/978-3-642-76064-8_1
7. Kuznetsov, S.I.; Dubinina, G.A. *Methods for studying aquatic microorganisms*; Science: Moscow, Russia, 1989; 285 p. (in Russian)
8. Cabello, P.; Roldán, M.D.; Castillo, F.; Moreno-Vivián, C. Nitrogen Cycle. In *Encyclopedia of Microbiology*, 3rd ed. 2009, pp. 299–321. doi:10.1016/B978-012373944-5.00055-9
9. Kuypers, M.; Marchant, H.; Kartal, B. The microbial nitrogen-cycling network. *Nat Rev Microbiol* **2018**, 16, 263–276. doi: 10.1038/nrmicro.2018.9.
10. Pajares, S.; Ramos, R. Processes and Microorganisms Involved in the Marine Nitrogen Cycle: Knowledge and Gaps. *Front. Mar. Sci.* **2019**, 6, 739. doi: 10.3389/fmars.2019.00739.
11. Hayatsu, M.; Tago, K.; Saito, M. Various players in the nitrogen cycle: Diversity and functions of the microorganisms involved in nitrification and denitrification. *Soil Science & Plant Nutrition* **2008**, 54, 33–45. doi: 10.1111/j.1747-0765.2007.00195.x
12. Stein, L.Y.; Klotz, M.G. The nitrogen cycle. *Curr. Biol.*, **2016**, 26 (3), R94–R 98. doi: 10.1016/j.cub.2015.12.021
13. Zehr, J. P.; Ward, B. B. Nitrogen cycling in the ocean: new perspectives on processes and paradigms. *Appl. Environ. Microbiol.* **2002**, 68, 1015–1024. Doi: 10.1128/aem.68.3.1015-1024.2002
14. Kuznetsov, S.I.; Saralov, A.I.; Nazina, T.N. *Microbiological processes of carbon and nitrogen circulation in lakes*; Science: Moscow, Russia, 1985; 216 p. (in Russian)
15. Gorlenko, V.M.; Dubinina, G.A.; Kuznetsov, S.I. *Ecology of aquatic microorganisms*; Science: Moscow, Russia, 1977; 289 p. (in Russian)
16. Camargo, J.A.; Alonso, Á. Ecological and toxicological effects of inorganic nitrogen pollution in aquatic ecosystems: A global assessment. *Environ Int.* **2006**, 32 (6), 831–849. doi: 10.1016/j.envint.2006.05.002
17. Stewart, K.M.; Walker, K.F.; Likens, G.E. *Meromictic Lakes*. In *Encyclopedia of Inland Waters*; Likens, G.E., Eds; Academic Press, 2009; P. 589–602. doi: 10.1016/B978-012370626-3.00027-2
18. Gulati, R.D.; Zadereev, E.S.; Degermendzhi, A.G. (eds.) *Ecology of Meromictic Lakes*; Springer International Publishing, Cham, Switzerland, 2017. doi: 10.1007/978-3-319-49143-1
19. Lam, P.; Kuypers M. M.M. Microbial nitrogen cycling processes in oxygen minimum zones. *Ann Rev Mar Sci.* **2011**, 3, 317–345. doi: 10.1146/annurev-marine-120709-142814
20. Ward, B.B.; Glover, H.E.; Lipschultz, F. Chemoautotrophic activity and nitrification in the oxygen minimum zone off Peru // *Deep Sea Research Part A. Oceanographic Research Papers* **1989**, 36, 1031–1051. doi: 10.1016/0198-0149(89)90076-9
21. Crowe, S.A.; Katsev, S.; Leslie, K.; Sturm, A.; Magen, C.; Nomosatryo, S.; Pack, M.A.; Kessler, J.D.; Reeburgh, W.S.; Roberts, J.A.; Gonzalez, L.; Haffner, G.D.; Mucci, A.; Sundby, B.; Fowle, D.A. The methane cycle in ferruginous Lake Matano. *Geobiology* **2011**, 9(1), 61–78. doi: 10.1111/j.1472-4669.2010.00257.x
22. Oshkin, I.Y.; Beck, D.A.; Lamb, A.E.; Tchesnokova, V.; Benuska, G.; McTaggart, T.L.; Kalyuzhnaya, M.G.; Dedysh, S.N.; Lidstrom, M.E.; Chistoserdova, L. Methane-fed microbial microcosms show differential community dynamics and pinpoint taxa involved in communal response. *ISME J* **2015**, 9, 1119–1129. doi: 10.1038/ismej.2014.203.
23. Chistoserdova, L. Methylophiles in natural habitats: current insights through metagenomics. *Appl Microbiol Biotechnol.* **2015**, 99 (14), 5763–5779. doi: 10.1007/s00253-015-6713-z.
24. Rissanen, A.J.; Saarenheimo, J.; Tirola, M.; Peura, S.; Aalto, S.L.; Karvinen, A.; Nykanen, H.; Gammaproteobacterial methanotrophs dominate methanotrophy in aerobic and anaerobic layers of boreal lake waters. *Aquat Microb Ecol.* **2018**, 81, 257–276. doi: 10.3354/ame01874
25. He, Z.; Zhang, Q.; Feng, Y.; Luo, H.; Pan, X.; Gadd, G.M. Microbiological and environmental significance of metal-dependent anaerobic oxidation of methane. *Sci Total Environ.* **2018**, 610–611, 759–768. doi:10.1016/j.scitotenv.2017.08.140
26. Cabrol, L.; F. Thalasso, L.; Gandois, A.; Sepulveda-Jauregui, K.; Martinez-Cruz, R.; Teisserenc, N.; Tananaev, A.; Tveit, M.M.; Svenning, M.; Barret M. Anaerobic oxidation of methane and associated microbiome in anoxic water of Northwestern Siberian lakes. *Sci. Total Environ.* **2020**, 736, Article 139588. doi: 10.1016/j.scitotenv.2020.139588
27. Knowles, R. Denitrifiers associated with methanotrophs and their potential impact on the nitrogen cycle. *Ecological Engineering* **2005**, 24(5), 441–446. doi: 10.1016/j.ecoleng.2005.01.001
28. Megm, S. R.; Knowles R. Active methanotrophs suppress nitrification in a humisol. *Biol Fert Soils* **1987**, 4, 205–212. doi: 10.1007/BF00270942
29. Raghoebarsing, A.A.; Pol, A.; van de Pas-Schoonen, K.T.; Smolders, A.J.P.; Ettwig, K.F.; Rijpstra, W. I. C.; Schouten S.; Damsté, J.S.S., Op den Camp, H.J.M.; Jetten, M.S.M.; Strous M. A microbial consortium couples anaerobic methane oxidation to denitrification. *Nature* **2006**, 440, 918–921. doi: 10.1038/nature04617.

30. Guggenheim, C.; Freimann, R.; Mayr, M.J.; Beck, K.; Wehrli, B.; Bürgmann, H. Environmental and Microbial Interactions Shape Methane-Oxidizing Bacterial Communities in a Stratified Lake // *Front. Microbiol.* **2020**, 11:579427. doi: 10.3389/fmicb.2020.579427
31. Einsiedl, F.; Wunderlich, A.; Sebiló, M.; Coskun, Ö.K.; Orsi W.D.; Mayer B. Biogeochemical evidence of anaerobic methane oxidation and anaerobic ammonium oxidation in a stratified lake using stable isotopes. *Biogeosciences* **2020**, 17, 5149–5161. doi: 10.5194/bg-17-5149-2020
32. Kits, K.D.; Klotz, M.G.; Stein, L.Y. Methane oxidation coupled to nitrate reduction under hypoxia by the Gammaproteobacterium *Methylomonas denitrificans*, sp. nov. type strain FJG1: denitrifying metabolism in *M. denitrificans* FJG1. *Environ Microbiol.* **2015**, 17, 3219–3232. doi: 10.1111/1462-2920.12772.
33. Zabelina S.A.; Kompantseva E.I.; Chupakov A.V.; Ershova A.A. Seasonal dynamics of photosynthetic pigments in a freshwater meromictic lake. Biodiagnostics of the state of natural and natural-technogenic systems: Proceedings of the XII All-Russian scientific-practical conference with international participation. Book 1. (Kirov, 2–3 December 2014). Publishing house of LLC "Vesi": Kirov, Russia, 2014. P. 119–123. (In Russian)
34. Kokryatskaya, N. M.; Chupakov A.V.; Titova, K.; Chupakova, A.A.; Zabelina, S.A.; Moreva, O.Yu.; Neverova, N.V.; Zhibareva, T.A. Hydrological and hydrochemical characteristics of the iron-manganese meromictic freshwater lake Svetloe (Arkhangelsk Region). *J. Sib. Fed. Univ. Biol.* **2019**, 12(2), 147–159. doi: 10.17516/1997-1389-0036.
35. Kallistova, A.; Kadnikov, V.; Rusanov, I.; Kokryatskaya, N.; Beletsky, A.; Mardanov, A.; Savvichev, A.; Ravin, N.; Pimenov, N. Microbial communities involved in aerobic and anaerobic methane cycling in a meromictic ferruginous subarctic lake. *Aquatic Microbial Ecology* **2019**, 82(1), 1–18. doi: <https://doi.org/10.3354/ame01878>
36. Savvichev, A.S.; Rusanov, I.I.; Zakharova, E.E.; Veslopolova, E.F.; Lunina, O.N.; Sigalevich, P.A.; Pimenov, N.V.; Gorlenko, V.M.; Kokryatskaya, N.M.; Zabelina, S.A.; Patutina, E.O.; Bumazhkin, B.K.; Gruzdev, D.S.; Kuznetsov, B.B. Microbial processes of the carbon and sulfur cycles in an ice-covered, iron-rich meromictic lake Svetloe (Arkhangelsk region, Russia). *Environ Microbiol.* **2017**, 19, 659–672. doi: 10.1111/1462-2920.13591.
37. Kadnikov, V.V.; Mardanov, A.V.; Beletsky, A.V.; Ravin, N.V.; Savvichev, A.S.; Merkel, A.Y.; Pimenov, N.V. Microbial communities involved in the methane cycle in the near-bottom water layer and sediments of the meromictic subarctic lake svetloe. *Antonie van Leeuwenhoek* **2019**, 112 (12), 1801–1814. doi: 10.1007/s10482-019-01308-1.
38. Ershova, A.A.; Vorobyeva, T.Ya.; Moreva, O.Yu.; Chupakov, A.V.; Zabelina, S.A.; Neverova, N.V. Hydrochemical and microbiological research of a nitrogen cycle in freshwater meromictic lake Svetloe (the Arkhangelsk region). *Reg Environ.* **2015**, (5), 44–50. (in Russian)
39. Chupakov, A.V.; Pokrovsky, O.S.; Shirokova, L.S.; Vorobyeva, T.Ya.; Zabelina, S.A.; Kokryatskaya, N.M.; Moreva, O.Yu.; Ershova, A.A.; Shorina, N.V.; Klimov, S.I. Hydrochemical characteristics of meromictic freshwater lake Svetloe (Arkhangelsk region). *Vestnik of Northern (Arctic) Federal University. Series Natural Sciences* **2013**, 1, 20–31 (in Russian)
40. Chupakov, A.V.; Chupakova, A.A.; Moreva, O.Yu.; Shirokova, L.S.; Zabelina, S.A.; Vorobieva, T.Y.; Klimov, S.I.; Brovko, O.S.; Pokrovsky, O.S. Allochthonous and autochthonous carbon in deep, organic rich and organic-poor lakes of the European Russian subarctic. *Boreal Environment Research* **2017**, 22, 213–230.
41. Canfield, D.E.; Rosing, M.T.; Bjerrum, C. Early anaerobic metabolisms. *Philos Trans R Soc Lond B Biol Sci* **2006**, 361, 1819–1836.
42. Scanlan, D.J.; West, N.J. Molecular ecology of the marine cyanobacterial genera *Prochlorococcus* and *Synechococcus*. *FEMS Microbiology Ecology* **2002**, 40, 1–12. doi: 10.1111/j.1574-6941.2002.tb00930.x.
43. Partensky, F.; Blanchot, J.; Vaultot, D. Differential distribution and ecology of *Prochlorococcus* and *Synechococcus* in oceanic waters: a review. In *Marine cyanobacteria*. no. NS 19. *Bulletin de l'Institut Océanographique Monaco*; Charpy, L., Larkum, A.W.D., Eds.; Musee océanographique : Monaco, **1999**; V NS 19.P. 457–475.
44. Willey, J.M.; Waterbury, J.B. Chemotaxis toward Nitrogenous Compounds by Swimming Strains of Marine *Synechococcus* spp. *Appl Environ Microbiol.* **1989**, 55, 1888–1894. doi: 10.1128/aem.55.8.1888-1894.1989
45. Steunou, A.S. Jensen, S.I.; Brecht, E.; Becraft, E.D.; Bateson, M.M.; Kilian, O.; Bhaya, D.; Ward, D.M.; Peters, J.W.; Grossman, A.R.; Köhl, M. Regulation of *nif* gene expression and the energetics of N₂ fixation over the diel cycle in a hot spring microbial mat. *ISME J.* **2008**, 2, 364–378. doi: 10.1038/ismej.2007.117.
46. Koroleff, F. Total and organic nitrogen. In: *Methods for seawater analysis*; Grasshoff K., Ehrhardt M., Kremling K., Eds.; Verlag Chemie Weinheim, **1983**; P. 162–168.
47. Oswald, K.; Milucka, J.; Brand, A.; Hach, P.; Littmann, S.; Wehrli, B.; Kuypers, M.M.M.; Schubert, C.J. Aerobic gammaproteobacterial methanotrophs mitigate methane emissions from oxic and anoxic lake waters. *Limnol Oceanogr.* **2016**, 61, S101–S118. doi: 10.1002/lno.10312.
48. Biderre-Petit, C.; Taib, N.; Gardon, H.; Hochart, C.; Debroas, D. New insights into the pelagic microorganisms involved in the methane cycle in the meromictic Lake Pavin through metagenomics, *FEMS Microbiol Ecol.* **2019**, 95 (3). doi: 10.1093/femsec/fiy183
49. Rodrigo, M.; Miracle, M.; Vicente, E. The meromictic Lake La Cruz (Central Spain). Patterns of stratification. *Aquat. sci.* **2001**, 63, 406–416. doi: 10.1007/s00027-001-8041-x
50. Llíros, M.; Garcá-Armisen, T.; Darchambeau, F.; Morana, C.; Triadó-Margarit, X.; Inceolu, Ö.; Borrego, C. M.; Bouillon, S.; Servais, P.; Borges, A. V.; Descy, J. P.; Canfield, D. E.; Crowe, S. A. Pelagic photoferrotrophy and iron cycling in a modern ferruginous basin. *Sci Rep.* **2015**, 5, 13803. doi: 10.1038/srep13803.

51. Umanskaya, M.V.; Tarasova, N.G.; Gorbunov, M.Y. Phototrophic plankton of siderotrophic meromictic Lake Kuznechikha (Republic Mari El, Russia). *Inland Water Biol* **2017**, *10*, 158–167 doi: 10.1134/S1995082917020158
52. Hove, P.V.; Belzile, C.; Gibson, J.A.E.; Vincent, W.F. Coupled landscape–lake evolution in High Arctic Canada. *Canadian Journal of Earth Sciences* **2006**, *43* (5), 533–546. doi:10.1139/e06-003
53. Stepanova, A.B.; Sharafutdinova, G.F.; Voyakina, E.Yu. Hydrochemical features of Valaam Island small lakes. *Scientific notes of RSHU*. **2010**, *12*, 97–110. (In Russian).
54. Culver, D.A. Biogenic meromixis and stability in a soft-water lake. *Limnology and Oceanography* **1977**, *22*(4), 667–686. doi: 10.4319/lo.1977.22.4.0667
55. Newton, R.J.; Jones, S.E.; Eiler, A.; McMahon, K.D.; Bertilsson, S. A guide to the natural history of freshwater lake bacteria. *Microbiol Mol Biol Rev* **2011**, *75*, 14–49. doi: 10.1128/MMBR.00028-10.
56. Melack, J.M., Jellison, R. Limnological conditions in Mono Lake: contrasting monomixis and meromixis in the 1990s. *Hydrobiologia* **1998**, *384*, 21–39 doi: 10.1023/A:1003352511328
57. Zabelina, S.A.; Pokrovsky, O.S.; Klimov, S.I.; Chupakov, A.V.; Kokryatskaya, N.M.; Vorobyova, T.Ya . The first results of the methane concentration in the stratified lakes of the River Svetlaya (the catchment area of the White Sea). In *Geology of seas and oceans*; Lisitzin, A.P., Politova, N.V., Shevchenko, V.P., Eds. Proc 20th Int Sci Conf on Marine Geology, 2013, 3. GEOS, Moscow, P. 156–160 (In Russian)
58. Brochier-Armanet, C., Boussau, B., Gribaldo, S., Forterre, P. Mesophilic crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. *Nat. Rev. Microbiol.* **2008**, *6*, 245–252. doi:10.1038/nrmicro1852
59. Hugoni, M.; Etien, S.; Bourges, A.; Lepe`re, C.; Domaizon, I.; Mallet, C.; Bronner, G.; Debroas, D.; Mary, I. Dynamics of ammonia-oxidizing Archaea and Bacteria in contrasted freshwater ecosystems. *Res Microbiol.* **2013**, *164*, 360–370. doi: 10.1016/j.resmic.2013.01.004.
60. Lliros, M.; Gich, F.; Plasencia, A.; Auguet, J.C.; Darchambeau, F.; Casamayor, E.O.; Descy, J.P.; Borrego, C. Vertical distribution of ammonia-oxidizing crenarchaeota and methanogens in the epipelagic waters of Lake Kivu (Rwanda-Democratic Republic of the Congo). *Appl. Environ. Microbiol.* **2010**, *76*, 6853–6863. doi: 10.1128/AEM.02864-09.
61. Pouliot, J., Galand, P.E., Lovejoy, C., Vincent, W.F. Vertical structure of archaeal communities and the distribution of ammonia monooxygenase A gene variants in two meromictic High Arctic lakes. *Environ. Microbiol.* **2009**, *11*, 687–699. doi: 10.1111/j.1462-2920.2008.01846.x.
62. Hatzenpichler, R.; Lebedeva, A.; Spieck, E.; Stoecker, K.; Richter, A.; Daims, H.; Wagner, M. A moderately thermophilic ammoniaoxidizing crenarchaeote from a hot spring. *Proc. Natl. Acad. Sci. U.S.A* **2008**, *105*, 2134–2139. doi: 10.1073/pnas.0708857105.
63. Park, H.D.; Wells, G.F.; Bae, H.; Criddle, C.S.; Francis, C.A. Occurrence of ammonia-oxidizing archaea in wastewater treatment plant bioreactors. *Appl Environ Microb.* **2006**, *72*, 5643–5647. doi: 10.1128/AEM.00402-06.
64. Erguder, T.H.; Boon, N.; Wittebolle, L.; Marzorati, M.; Verstraete, W. Environmental factors shaping the ecological niches of ammonia-oxidizing archaea. *FEMS Microbiol. Rev.* **2009**, *33*, 855–869. doi: 10.1111/j.1574-6976.2009.00179.x.
65. Lam, P.; Jensen, M.M.; Lavik, G.; McGinnis, D.F.; Muller, B.; Schubert, C.J.; Amann, R.; Thamdrup, B.; Kuypers, M.M.M. Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea. *Proc Natl Acad Sci USA* **2007**, *104*, 7104–7109. doi: 10.1073/pnas.0611081104.
66. Beman, J.M.; Popp, B.N.; Francis, C.A. Molecular and biogeochemical evidence for ammonia oxidation by marine Crenarchaeota in the Gulf of California. *ISME J.* **2008**, *2*(4), 429–441. doi: 10.1038/ismej.2007.118.
67. Molina, V.; Belmar, L.; Ulloa, O. High diversity of ammonia-oxidizing archaea in permanent and seasonal oxygen-deficient waters of the eastern South Pacific // *Environ Microbiol.* **2010**, *12*, 2450–2465. doi: 10.1111/j.1462-2920.2010.02218.x.
68. Simon, J. Enzymology and bioenergetics of respiratory nitrite ammonification. *FEMS Microbiol. Rev.* **2002**, *26*, 285–309. doi: 10.1111/j.1574-6976.2002.tb00616.x.
69. Daims, H.; Lebedeva, E.V.; Pjevac, P.; Han, P.; Herbold, C.; Albertsen, M.; Jehmlich, N.; Palatinszky, M.; Vierheilig, J.; Bulaev, A.; Kirkegaard, R.H.; Bergen, M.V.; Rattei, T.; Bendinger, B.; Nielsen, P.H.; Wagner, M. Complete nitrification by *Nitrospira* bacteria. *Nature* **2015**, *528* (7583), 504–509. doi:10.1038/nature16461.
70. Kits, K.D.; Sedlacek, Ch. J.; Lebedeva, E. V.; Han, P.; Bulaev, A.; Pjevac, P.; Daebeler, A.; Romano, S.; Albertsen, M. Kinetic analysis of a complete nitrifier reveals an oligotrophic lifestyle. *Nature* **2017**, *549* (7671), 269–272. doi:10.1038/nature23679
71. Froelich, P.N.; Klinkhammer, G.P.; Bender, M.L.; Luedtke, N.A.; Heath, G.R.; Cullen, D.; Dauphin, P.; Hammond, D.; Hartman, B.; Maynard, V. Early oxidation of organic matter in pelagic sediments of eastern equatorial Atlantic: suboxic diagenesis. *Geochimica et Cosmochimica Acta* **1979**, *43*, 1075–1090. doi: 10.1016/0016-7037(79)90095-4.
72. Luther, G.W.; Sundby, B.; Lewis, B.L.; Brendel, P.J.; Silverberg, N. Interactions of manganese with the nitrogen cycle: Alternative pathways to dinitrogen. *Geochim. Cosmochim. Ac.* **1997**, *61*, 4043–4052. doi: 10.1016/S0016-7037(97)00239-1.
73. Hulth, S.; Aller, R.C.; Gilbert, F. Coupled anoxic nitrification/manganese reduction in marine sediments. *Geochimica et Cosmochimica Acta* **1999**, *63*, 49–66. doi: 10.1016/S0016-7037(98)00285-3
74. Jones, C.; Crowe, S.A.; Sturm, A.; Leslie, K.L.; MacLean, L.C.W.; Katsev, S.; Henny, C.; Fowle, D.A.; Canfield, D.E. Biogeochemistry of manganese in ferruginous Lake Matano, Indonesia. *Biogeosciences* **2011**, *8*, 2977–2991. doi: 10.5194/bg-8-2977-2011.
75. Bratina, B.J.; Stevenson, B.S.; Green, W.J.; Schmidt, T.M. Manganese reduction by microbes from oxic regions of the Lake Vanda (Antarctica) water column. *Appl. Environ. Microbiol.* **1998**, *64*, 3791–3797. doi: 10.1128/AEM.64.10.3791-3797.1998

76. Canfield, D.E.; Kristensen, E.; Thamdrup, B. Aquatic Geomicrobiology. In: Adv. Mar. Biol.; Southward, A.J., Tyler, P.A., Young, C.M., Fuiman, L.A., Eds.; Elsevier: San Diego, 2005; 640 pp.
77. Konneke, M.; Bernhard, A.E.; de la Torre, J.R.; Walker, C.B.; Waterbury, J.B.; Stahl, D.A.; Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* **2005**, *437*, 543–546. doi: 10.1038/nature03911.
78. Walker, C.B.; de la Torre, J.R.; Klotz, M.G.; Urakawa, H.; Pinel, N.; Arp, D.J.; Brochier-Armanet, C.; Chain, P.S.J.; Chan, P.P.; Gollabgir, A.; Hemp, J.; Hmgler, M.; Karr, E.A.; Kwnneke, M.; Shin, M.; Lawton, T.J.; Lowe, T.; Martens-Habbena, W.; Sayavedra-Soto, L.A.; Lang, D.; Sievert, S.M.; Rosenzweig, A.C.; Manning, G.; Stahl, D.A. Nitrosopumilus maritimus genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proc Natl Acad Sci USA* **2010**, *107*(19), 8818–8823. doi: 10.1073/pnas.0913533107.
79. Mulder, A.; van de Graaf, A.A.; Robertson, L.A.; Kuenen, J.G. Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed reactor. *FEMS Microbiol Ecol* **1995**, *16*, 177–184. doi: 10.1016/0168-6496(94)00081-7.
80. Jensen, M.M.; Thamdrup, B.; Dalsgaard T. Effects of specific inhibitors on anammox and denitrification in marine sediments. *Appl. Environ. Microbiol.* **2007**, *73*, 3151–3158. doi: 10.1128/AEM.01898-06.
81. Kartal, B.; Kuypers, M.M.M.; Lavik, G.; Schalk, J.; Op den Camp, H.J.M.; Jetten, M.S.; Strous, M. Anammox bacteria disguised as denitrifiers: nitrate reduction to dinitrogen gas via nitrite and ammonium. *Environ. Microbiol.* **2007**, *9*, 635–642. doi: 10.1111/j.1462-2920.2006.01183.x.
82. Hu, B-I; Shen, L-D.; Xu, X-Y.; Zheng, P. Anaerobic ammonium oxidation (anammox) in different natural ecosystems. *Biochem Soc Trans.* **2011**, *39*, 1811–1816. doi: 10.1042/BST20110711.
83. Schubert, C.J.; Durisch-Kaiser, E.; Wehrli, B.; Thamdrup, B.; Lam, P.; Kuypers, M.M.M. Anaerobic ammonium oxidation in a tropical freshwater system (Lake Tangayika). *Environ Microbiol.* **2006**, *8*, 1857–1863. doi: 10.1111/j.1462-2920.2006.01074.x.
84. Hamersley, M.R.; Woebken, D.; Boehrer, B.; Schultze, M.; Lavik, G.; Kuypers, M.M.M. Water column anammox and denitrification in a temperate permanently stratified lake (Lake Rassnitzer, Germany). *Syst Appl Microbiol.* **2009**, *32*, 571–582. doi: 10.1016/j.syapm.2009.07.009
85. Wenk, C.B.; Zopfi, J.; Brees, J.; Veronesi, M.; Niemann, H.; Lehmann, M.F. Community N and O isotope fractionation by sulphide-dependent denitrification and anammox in a stratified lacustrine water column. *Geochim Cosmochim Acta* **2014**, *125*, 551–563. doi: 10.1016/j.gca.2013.10.034
86. Strous, M.; Pelletier, E.; Mangenot, S.; Rattei, T.; Lehner, A.; Taylor, M.; Horn, M.; Daims, H.; Bartol-Mavel, D.; Wincker, P.; Barbe, V.; Fonknechten, N.; Vallenet, D.; Segurens, B.; Schenowitz-Truong, C.; Médigue, C.; Collingro, A.; Snel, B.; Dutilh, B.; Op den Camp, H.; van der Drift, C.; Cirpus, I.; van de Pas-Schoonen, K.; Harhangi, H.; van Niftrik, L.; Schmid, M.; Keltjens, J.; van de Vossenberg, J.; Kartal, B.; Meier, H.; Frishman, D.; Huynen, M.; Mewes, H-W.; Weissenbach, J.; Jetten, M.; Wagner, M.; Le Paslier, D. Deciphering the evolution and metabolism of an anammox bacterium from a community genome. *Nature* **2006**, *440*(7085), 790–794. doi: 10.1038/nature04647.
87. Anthony, C. The biochemistry of methylotrophs; Academic Press: London, 1982.