

## TITLE PAGE

# COMPARISON OF GROUND REACTION FORCES BETWEEN COMBAT BOOTS AND SPORTS SHOES

**Authors:** Rodrigo Rico Bini<sup>1</sup>, Daniel D. Kilpp<sup>2</sup>, Pedro D. Júnior<sup>2</sup>, Adriane D. Muniz<sup>2</sup>

<sup>1</sup> *La Trobe Rural Health School, Bendigo Campus, Flora Hill, Australia*

<sup>2</sup> *Escola de Educação Física do Exército, Centro de Capacitação Física do Exército, Rio de Janeiro, Brazil.*

## Name and address for correspondence

Dr. Rodrigo Rico Bini, La Trobe Rural Health School, College of Science, Health and Engineering, La Trobe University | Flora Hill Campus, Bendigo, Victoria, 3550, AUSTRALIA; Phone (61) 3 5444 7466; Email: [r.bini@latrobe.edu.au](mailto:r.bini@latrobe.edu.au)

## ABSTRACT

It is unclear whether military shoes (combat boots and sports shoes) attenuate loading rate or affect force transfer during walking. Therefore, this study compared ground reaction forces (GRF) related to impact and force transfer between combat boots, military sports shoes and running shoes. Ten army recruits walked over a walkway with two force plates embedded. GRF were measured when walking barefoot (for data normalization) and with combat boots, military sports shoes and running shoes. Loading rate, first and second peak forces and push-off rate of force were computed along with temporal analysis of waveforms. Reduced loading rate was observed for the running shoe compared to the combat boot ( $p = 0.02$  and  $d = 0.98$ ) and to the military sports shoe ( $p = 0.04$  and  $d = 0.92$ ). The running shoe elicited a smaller second peak force than the combat boot ( $p < 0.01$  and  $d = 0.83$ ). Walking with military shoes and combat boots led to larger force transfer than running shoes potentially due to harder material used in midsole composition (i.e. styrene-butadiene rubber). These results could lead to a potentially larger risk of injuries while long duration walking in military shoes and boots compared to traditional running shoes.

**Keywords:** gait, loading rate, ethylene-vinyl acetate, styrene-butadiene rubber

## Introduction

Physical training is a key part of the job requirements of defense personnel, which involve marching, jumping and walking with large external loads. Several studies indicated that external loads can contribute to an increased likelihood of injuries during operational tasks [1]. Aerobic and strength training are also prescribed to improve general fitness [1]. During these activities, recruits use various types of shoes, from military boots to regular sports shoes [2]. However, it has been observed that the type of footwear used by military personnel could also increase overuse injury risk, particularly when walking with combat boots [3,4].

It has been long established that shoes can affect ground reaction forces (e.g. altering vertical loading rate) during running and walking [5-7] due to the characteristics of the shoe midsole [8] and due to shoe-ground interaction [9]. Among the variables assessed from ground reaction forces, loading rate (i.e. rate of vertical force increments at the initial stance phase) has been a key variable because it relates positively to the velocity at which ground reaction forces are absorbed by the musculoskeletal system [10,11]. Therefore, large loading rates lead to faster transfer of force and less time for the soft tissues to accommodate the load [12], which could lead to overuse injuries. In addition, push-off rate of force (i.e. rate of force decrement late in the stance phase) can indicate how rapidly the forces are applied to propel the body forward during motion [5]. Large rate of force decrement could also lead to overuse injuries given the increased force transferred through the metatarsal heads [13]. Therefore, shoe design could play a role in alleviating force transfer through the foot by improving the cushioning characteristics of shoe midsole.

For army recruits, boots used during marching and other activities have been shown to reduce ankle range of motion [2] without differences in loading rate or push-off rate of force

compared to a generic running shoe [5]. Although these findings may show that military boots do not impact force transmission, Pasis et al. [5] tested subjects in their existing shoes without reporting the shoes' characteristics (e.g. material of the shoe midsole or shoe design), which limit implications from their findings. Military boots have been shown to increase Achilles tendon force [14] and knee load [15] compared to running shoe with EVA (Ethylene-vinyl acetate) midsole. However, differences to a shoe with mixed EVA and rubber, as commonly observed in running shoes [13,16], has not been assessed in terms of loading rate or force transfer. This is critical to provide data that could support the improvements in design of military shoes, in order to reduce injury marks in army recruits [1]. Moreover, the temporal analysis of ground reaction forces is critical because it allows for detecting differences in external forces which are not always captured when analyzing zero-dimensional data [i.e. peaks and means; 17].

Therefore, the aim of this study was to compare ground reaction forces between combat boots, sports shoes designed for military training and running shoes during walking gait. The option of walking was based on the large proportion of walking activities performed by military personnel [i.e. 60-70% of physical activity; 4,18]. The assessment of a sports shoe designed for military training was based on the use of the same type of midsole compared to the combat boot, which should allow for differences in shape between boots and shoes to be further explored. The hypothesis of this study was that the combat boot and the sports shoe would have larger loading rates and push-off rates of force compared to the running shoe.

## Methods

### *Participants*

This study design was cross-sectional with randomized crossover trials between types of shoes. Ten army recruits ( $28 \pm 1.8$  years of age,  $75 \pm 3.9$  kg of body mass and  $177 \pm 6.6$  cm of height) volunteered to take part in the study. This sample size was estimated using a statistical package (Biostat 5.3, Instituto Mamirauá, Brazil), for a change in mean in a cross-over design assuming Type I and II errors at 5% and 20%, respectively. Before taking part in the study, all participants read and signed an informed consent form with details of the study approved by the local committee of ethics in research with humans (approval code: 46243715.4.0000.5235).

### *Shoe specifications*

Combat boots, military sports shoes and running shoes (size 43 Europe – see Figure 1), with specifications as described in Table 1 were used in this study. For measurements of shoe mass, a force plate was utilized (resolution of 9 g, FP4060-10, Bertec Corporation, USA). Midsole stiffness were collected from testing conducted by manufacturers, following local guidelines for shoe assessments [19].



**Figure 1.** Military sports shoe (A), combat boot (B) and running shoe (C).

**Table 1.** Specifications of the combat boot, military sports shoe and the running shoe.

Shoe type	Mass (g)	Midsole height (cm)	Shoe structure and midsole properties	Insole
Combat boot (LV Distribuidora LTDA, Brazil)	550	Forefoot: 270 Height: 380	11 holes cording SBR midsole (~65 Shore A stiffness)	EVA
Military sports shoe (LV Distribuidora LTDA, Brazil)	350	Forefoot: 280 Height: 380	5 holes cording SBR midsole (~65 Shore A stiffness)	EVA
Running shoe (Prorunner 17, Mizuno)	260	Forefoot: 210 Height: 350	5 holes cording Carbon sole with 30% EVA (~45 Shore A for EVA stiffness)	EVA

EVA - Ethylene-vinyl acetate, SBR - styrene-butadiene rubber


### *Data collection*

Before gait analysis, participants' height and body mass were measured using a weight scale (Prix 2098PP, Toledo, Brazil). After that, all participants were familiarized with the gait analysis procedures, which involved walking barefoot over a wooden walkway with 11 meters of length. At the center of the walkway, a pair of force plates (FP4060-10, Bertec Corporation, USA) were embedded to enable measurements of the ground reaction forces. During familiarization trials, participants were instructed to modulate their gait pattern (via changes in stride length and cadence) in order to walk at  $5 \pm 0.25$  km/h. For assessment and feedback on walking speed, two measurements were taken by two experienced technicians, based on the time taken to cover a distance of 4.3 m (i.e. length of the force plates at the wooden walkway) [20].

After familiarization, participants performed ten trials of barefoot walking at the target speed and with a stride length that enabled a full contact of the foot in each force plate (one for each foot). After that, in random order, they performed walking trials using the combat boot, the military sports shoe and the running shoe. Ground reaction forces were collected during all trials using the software from the force plates (Digital Acquire 4.0.12.411, Bertec Corporation, USA) at a sampling rate of 1kHz (per channel). Time for covering the 4.3 meters distance was recorded, using two time-watches (WT058, DLKSports, Brazil), in a spreadsheet for later computation of mean walking speed between trials and for provision of feedback on walking speed to participants during each trial.

### Data analyses

Heel strike and toe-off were detected using a 20 N threshold [21] in order to determine stance phases (i.e. contact time) for each foot. After that, first and second peaks were determined from the vertical component of the ground reaction force waveforms for each foot. The slope of 10-90% of first peak force was used to calculate instantaneous loading rate (after heel strike) and the slope of 90-10% of the second peak force enabled to calculate push-off rate (before toe-off, Figure 2). Force data was time-normalised to 100 samples using heel strike and toe-off to enable temporal statistical analysis of data. All data processing was conducted using custom made scripts (Matlab, MathWorks Inc, Natick, USA).



**Figure 2.** Illustration of the first (FPF) and second (SPF) peak force used to determine a range of 10-90% of each peak force. These ranges were then used to determine instantaneous loading rate (after heel strike) and push-off rate of force (before toe-off).

### *Statistical analyses*

Vertical ground reaction force data were normalized to outcomes taken from the barefoot trials in order to reduce non-uniformity distribution of the data, then converted into means from both force plates. Mean and standard deviation of the first and second peak forces, contact time (s), loading rate (% bodyweight/s) and push-off rate of force (% bodyweight/s) were computed. Differences between shoes were assessed using repeated measures ANOVA followed by post-hoc analysis (with Holm's correction) when main effects were observed. Magnitude of differences were rated using Cohen's effect sizes [22]. Substantial differences between shoes were assumed when  $p < 0.05$  and Cohen's  $d > 0.80$ . A statistical package (JASP, Version 0.13.1, University of Amsterdam, The Netherlands) was used for statistical analysis and differences were determined significant when  $p < 0.05$ .

For analysis of differences between boots and shoes in terms of temporal patterns, one-dimensional force data were analysed by repeated measures ANOVAs using the SPM method (Pataky et al., 2013). Pairwise comparisons were performed using paired samples t-tests with Bonferroni correction, to protect from Type I error. Critical t-thresholds were determined at  $\alpha = 0.05$  (Pataky et al., 2016). SPM analyses were implemented in Matlab (R2015, The Mathworks, USA) using the spm1d toolbox (<http://www.spm1d.org>).

### **Results**

The control of timing at the central section of the walkway secured similarities in walking speed between trials ( $p = 0.24$ , Table 2). There was a reduced loading rate for the running shoe compared to the combat boot ( $p = 0.02$  and  $d = 0.98$ ) and compared to the military sports shoe ( $p = 0.04$  and  $d = 0.92$ ). In addition, the running shoe elicited a smaller second peak force than the combat boot ( $p < 0.01$  and  $d = 0.83$ ). There was also a trend for




lower second peak force for the military sports shoe compared to the combat boot ( $p < 0.01$  and  $d = 0.69$ ). These results are shown in Table 2.

**Table 2.** mean (SD) gait speed, loading rate, first and second peak forces and push-off rate of force for walking trials with combat boot, military sports shoe and running shoes.

	Combat boot	Military sports shoe	Running shoe
Gait speed (km/h)	5.09 $\pm$ 0.04	5.11 $\pm$ 0.08	5.11 $\pm$ 0.05
Contact time (s)	0.67 $\pm$ 0.02	0.67 $\pm$ 0.03	0.67 $\pm$ 0.03
Loading rate (% barefoot)	19 $\pm$ 3	20 $\pm$ 5	16 $\pm$ 2* <sup>#</sup>
1 <sup>st</sup> peak force (% barefoot)	106 $\pm$ 3	105 $\pm$ 3	105 $\pm$ 3
2 <sup>nd</sup> peak force (% barefoot)	104 $\pm$ 1	101 $\pm$ 2	101 $\pm$ 1*
Push-off rate of force (% barefoot)	90 $\pm$ 12	83 $\pm$ 11	86 $\pm$ 10

\* Indicates difference to combat boot and <sup>#</sup> Indicates difference to military sports shoe when  $p < 0.05$  and  $d > 0.80$ .

Main effects were detected by the SPM-ANOVA for the vertical ground reaction force between 73-78% of the stance but differences in post hoc test were only observed between the combat boot and the running shoe at 73-74% of the stance (Figure 3).



**Figure 3.** A) Average vertical GRF data. B) ANOVA Footwear main effect trajectory. The horizontal dotted lines indicate the critical random field theory threshold of  $\alpha < 0.05$ . As the SPM {F} line crosses the dotted line above, a statistical difference was found. C) T-test comparison between military shoe vs. sports shoe. D) T-test comparison between combat boot shoe vs. running shoe. E) T-test comparison between military shoe vs. combat boot.

## Discussion

Although research on shoe midsole material has been covered in many studies, the assessment of shoes used by military recruits has received less attention compared to sports shoes [5,14,15,23]. These studies were generally limited to the comparison of combat boots with users' own shoes [5] or custom made shoes [15,24]. The main finding from the present study was that combat boots and military sports shoes presented reduced cushioning properties compared to a commercially available running shoe. This information is novel

because it indicates that a shoe with a midsole merging EVA and carbon rubber (i.e. running shoe) provides improved cushioning compared to the SBR midsole used in combat boots and in the military sports shoe. A secondary finding was that a trend was observed towards larger second peak force for the combat boot compared to the military shoe ( $p < 0.01$  and  $d = 0.69$ ). This element indicates that limitations in ankle range of motion from the combat boot could play a role given SBR midsole was common to combat boots and the military sports shoe.

Shoe midsole has been shown critical to reduce plantar pressure [13], Achilles tendon loading rate [14] and knee loads [15,25] during walking and running. Therefore, the choice for a proper shoe seems to help reducing musculoskeletal loading. Indeed, the running shoe presented better cushioning profile (reduced loading rate) compared to the combat boot and to the military sports shoe potentially because it has 30% of EVA associated with carbon rubber at the midsole. Differently, the military shoe and the combat boot had a SBR midsole (~65 Shore A stiffness) which is harder compared to the EVA (~45 Shore A stiffness). Contrary, recent comparison of combat boots indicated that boots with harder midsole were associated with less loading rate [17,26]. In a different line, Pasis et al. [5] observed similar loading rates for combat boots compared to users' running shoe, which could be linked to differences in experimental design. In the present study, a brand-new shoe was used rather than the users' shoes, which should present altered midsole properties from long term use [27]. A second difference involves the assessment of ground reaction forces using overground walking (present study) rather than treadmill walking, which would influence walking gait style. A third difference was that loading rate was normalized, for each shoe in the present study, by the individual's loading rate during barefoot trials in order to reduce the variance from individual's gait profile [28]. However, further studies may be required to fully examine other gait changes related to the use of shoes and boots with different design (e.g. kinematics of boots and shoes with the same midsole structure).

The second part of stance phase is related to the propulsion of the body towards the swing phase [29]. Therefore, the trend for the larger second peak force when walking with the combat boot could be associated with an attempt to mitigate the limited ankle plantar flexion [2]. This could require larger force production from ankle plantar flexor muscles to develop similar joint power. This was partially highlighted by Sinclair et al. [14], who observed increased loading rate at the Achilles tendon when recruits ran with combat boots rather than with running shoes. In addition, larger second peak forces may lead to increased pressure at the metatarsal heads [13], increasing the risk of injuries in the forefoot. However, more data is needed to ascertain if larger forces could be associated with larger pressure at the forefoot when using combat boots.

Although differences in loading rate and second peak force were observed between combat boots compared to running shoes, contact time during stance was unaffected by footwear. This result suggests that stride length was potentially similar between shoes due to walking speed control. For overground walking, the control of speed is critical because, different from treadmill walking, larger degrees of freedom are available during overground walking, which could affect ground reaction forces. Therefore, previously observed increases in activation of peroneous longus, tibialis anterior and rectus femoris when walking with combat boots could be related to participants attempting to optimize force transfer through the ankle joint and control joint kinematics given the restriction in ankle motion [23]. We also did not observe substantial differences between shoes in terms of temporal patterns (i.e. force waveforms), which could be associated with the prescribed walking speed. This is somehow surprising because SPM analysis have been shown sensitive to determine differences between boots with SBR vs. polyurethane midsole [17]. It may be that combat boots and military shoes with same type of midsole may have subtle differences that are not detectable by the SPM analysis. The fact that the SPM analysis was conducted using data normalized by body weight

rather than the barefoot trials could also add differences to the outcomes. Future studies could explore new ways to normalize ground reaction force data in order to use these waveforms for temporal comparisons.

The present study was limited to some extent. The absence of measurements of muscle activation and joint kinematics limited the combined analysis of joint control and ground reaction forces, as conducted in other studies [2,23]. The use a different insole for the running shoe, compared to the combat boot and the military shoe may have impacted our results, although all insoles were built from EVA. In addition, data from Pasis et al. [5] showed that EVA and polyurethane insoles did not affect loading rate or push-off rate of force. Therefore, it is unclear if insole type has largely influenced our results.

## **Conclusions**

In conclusion, combat boots and military sports shoes presented larger loading rate than running shoes during walking, which was thought to be potentially due to harder material used in the midsole (i.e. SBR). Combat boots also elicited increased second peak force, which was possibly associated with limited ankle range of motion. Combat boots used by army recruits do not optimize load transmission and may lead, in a long-term perspective, to greater injury risk.

**Supplementary Materials:** None.

**Funding:** This research received no external funding.

**Acknowledgements:** Thanks are given to participants who volunteered for this study.

**Author contribution:** Conceptualization, R.B, D.K. and P.J.; Methodology, R.B, D.K. and P.J.; Software, R.B. and A.M.; Formal Analysis, R.B. and A.M.; Investigation, R.B, D.K, P.J. and A.M.; Resources, R.B. and A.M.; Data Curation, R.B. and A.M.; Writing – Original Draft Preparation, R.B.; Writing – Review & Editing, R.B, D.K, P.J. and A.M.; Visualization, X.X.; Supervision, R.B.; Project Administration, R.B.; Funding Acquisition, N/A.

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Ethics Committee of the Centro Universitario Augusto Motta/UNISUAM (protocol code CAE: 46243715.4.0000.5235 approved on 14 October 2015).

**Data Availability Statement:** Data will be made available upon reasonable request to the corresponding author.

**Conflict of interest statement:** The authors declare no existing conflict of interest for the content of the article.

## References

1. Knapik, J.J.; Reynolds, K.L.; Harman, E. Soldier load carriage: historical, physiological, biomechanical, and medical aspects. *Military Medicine* **2004**, *169*, 45-56, doi:<http://publications.amsus.org/doi/pdf/10.7205/MILMED.169.1.45>.
2. Schulze, C.; Lindner, T.; Woitge, S.; Schulz, K.; Finze, S.; Mittelmeier, W.; Bader, R. Influence of footwear and equipment on stride length and range of motion of ankle,

- knee and hip joint. *Acta of Bioengineering and Biomechanics* **2014**, *16*, 45-51, doi:[www.actabio.pwr.wroc.pl/Vol16No4/5.pdf](http://www.actabio.pwr.wroc.pl/Vol16No4/5.pdf).
3. Bullock, S.H.; Jones, B.H.; Gilchrist, J.; Marshall, S.W. Prevention of Physical Training–Related Injuries: Recommendations for the Military and Other Active Populations Based on Expedited Systematic Reviews. *American Journal of Preventive Medicine* **2010**, *38*, S156-S181, doi:<http://dx.doi.org/10.1016/j.amepre.2009.10.023>.
  4. Kaufman, K.R.; Brodine, S.; Shaffer, R. Military training-related injuries: Surveillance, research, and prevention. *American Journal of Preventive Medicine* **2000**, *18*, 54-63, doi:[http://dx.doi.org/10.1016/S0749-3797\(00\)00114-8](http://dx.doi.org/10.1016/S0749-3797(00)00114-8).
  5. Pasis, P.; Hanley, B.; Havenetidis, K.; Bissas, A. Cypriot and greek army military boot cushioning: ground reaction forces and subjective responses. *Military Medicine* **2013**, *178*, e493-497, doi:10.7205/MILMED-D.
  6. Sepúlveda, G.; Flores, F.L.; Gurgel, J.L.; Porto, F.; Espinosa, G.; Gonçalves, F.S.; Fontoura, A.; Russomano, T. Relação de custo-benefício de calçados esportivos de Corrida quanto à capacidade de absorção de impacto. In Proceedings of the XII Congresso Brasileiro de Biomecanica, Águas de São Pedro, 2007; pp. 1671-1676.
  7. Hamill, J. Evaluating sport shoes using ground reaction force data. In Proceedings of the 14 International Symposium on Biomechanics in Sports, 1996; pp. 111-119.
  8. Pereira, G.; Avila, A.O.V.; Palhano, R. Vertical ground reaction force analysis during gait with unstable shoes. *Fisioterapia em Movimento* **2015**, *28*, 459-466, doi:10.1590/0103-5150.028.003.AO04.
  9. Nigg, B.M. The role of impact forces and foot pronation: a new paradigm. *Clinical Journal of Sports Medicine* **2001**, *11*, 2-9, doi:[http://journals.lww.com/cjsportsmed/Fulltext/2001/01000/The\\_Role\\_of\\_Impact\\_Forces\\_and\\_Foot\\_Pronation\\_A.2.aspx](http://journals.lww.com/cjsportsmed/Fulltext/2001/01000/The_Role_of_Impact_Forces_and_Foot_Pronation_A.2.aspx).

10. Crowell, H.P.; Davis, I.S. Gait retraining to reduce lower extremity loading in runners. *Clinical Biomechanics* **2011**, *26*, 78-83, doi:10.1016/j.clinbiomech.2010.09.003.
11. Zifchock, R.A.; Davis, I.; Hamill, J. Kinetic asymmetry in female runners with and without retrospective tibial stress fractures. *Journal of Biomechanics* **2006**, *39*, 2792-2797, doi:10.1016/j.jbiomech.2005.10.003.
12. Furey, M.J. Joint Lubrication. In *Biomechanics: Principles and Applications*, Bronzino, J.D., Ed.; CRC Press: 2007; pp. 73-97.
13. Wiegerinck, J.I.; Boyd, J.; Yoder, J.C.; Abbey, A.N.; Nunley, J.A.; Queen, R.M. Differences in plantar loading between training shoes and racing flats at a self-selected running speed. *Gait & Posture* **2009**, *29*, 514-519, doi:10.1016/j.gaitpost.2008.12.001.
14. Sinclair, J.; Taylor, P.J.; Atkins, S. Influence of running shoes and cross-trainers on Achilles tendon forces during running compared with military boots. *Journal of the Royal Army Medical Corps* **2015**, *161*, 140-143, doi:10./jramc-2014-000308.
15. Sinclair, J.; Taylor, P.J.; Atkins, S. Effects of new military footwear on knee loading during running. *Footwear Science* **2015**, *7*, 165-171, doi:10.1080/19424280.2015.1066879.
16. Even-Tzur, N.; Weisz, E.; Hirsch-Falk, Y.; Gefen, A. Role of EVA viscoelastic properties in the protective performance of a sport shoe: computational studies. *Bio-medical materials and engineering* **2006**, *16*, 289-299, doi:<http://content.iospress.com/articles/bio-medical-materials-and-engineering/bme429>.
17. Mara de Souza Muniz, A.; Sizenando, D.; Lobo, G.; Neves, E.B.; Gonçalves, M.; Marson, R.; Palhano, R.; Menegaldo, L.; Bini, R.R. Effects from loaded walking with polyurethane and styrene-butadiene rubber midsole military boots on kinematics and



- external forces: A statistical parametric mapping analysis. *Applied Ergonomics* **2021**, 94, 103429, doi:<https://doi.org/10.1016/j.apergo.2021.103429>.
18. Orr, R.M.; Johnston, V.; Coyle, J.; Pope, R. Reported Load Carriage Injuries of the Australian Army Soldier. *Journal of Occupational Rehabilitation* **2015**, 25, 316-322, doi:10.1007/s10926-014-9540-7.
  19. Técnicas, A.-A.B.d.N. Construction inferior of the shoes - Bottom, soles and materials for this purpose - Shore A and D hardness test. *ABNT NBR 14454:2007* **2007**, 5.
  20. Kim, H.-j.; Park, I.; Lee, H.j.; Lee, O. The reliability and validity of gait speed with different walking pace and distances against general health, physical function, and chronic disease in aged adults. *J Exerc Nutrition Biochem* **2016**, 20, 46-50, doi:10.20463/jenb.2016.09.20.3.7.
  21. Zeni, J.A., Jr.; Richards, J.G.; Higginson, J.S. Two simple methods for determining gait events during treadmill and overground walking using kinematic data. *Gait and Posture* **2008**, 27, 710-714, doi:10.1016/j.gaitpost.2007.07.007.
  22. Cohen, J. *Statistical power analysis for the behavioral sciences*, 2 ed.; Routledge Academic: Hillsdale, NJ, 1988; Volume 1, p. 590.
  23. Schulze, C.; Lindner, T.; Schulz, K.; Finze, S.; Kundt, G.; Mittelmeier, W.; Bader, R. The influence in airforce soldiers through wearing certain types of army-issue footwear on muscle activity in the lower extremities. *The Open Orthopaedics Journal* **2011**, 5, 302-306, doi:10.2174/1874325001105010302.
  24. Sinclair, J.; Naemi, R.; Chockalingam, N.; Taylor, P.J.; Shore, H. The effects of shoe temperature on the kinetics and kinematics of running. *Footwear Science* **2015**, 7, 173-180, doi:10.1080/19424280.2015.1084389.
  25. Erhart, J.C.; Dyrby, C.O.; D'Lima, D.D.; Colwell, C.W.; Andriacchi, T.P. Changes in in vivo knee loading with a variable-stiffness intervention shoe correlate with changes

- in the knee adduction moment. *Journal of Orthopaedic Research* **2010**, 28, 1548-1553, doi:10.1002/jor.21183.
26. Muniz, A.M.S.; Bini, R.R. Shock attenuation characteristics of three different military boots during gait. *Gait & Posture* **2017**, 58, 59-65, doi:<https://doi.org/10.1016/j.gaitpost.2017.07.037>.
27. Kong, P.W.; Candelaria, N.G.; Smith, D.R. Running in new and worn shoes: A comparison of three types of cushioning footwear. *British Journal of Sports Medicine* **2009**, 43, 745-749.
28. Revill, A.L.; Perry, S.D.; Michelle Edwards, A.; Dickey, J.P. Variability of the impact transient during repeated barefoot walking trials. *Journal of Biomechanics* **2008**, 41, 926-930, doi:10.1016/j.jbiomech.2007.11.002.
29. Vaughan, C.L.; Davis, B.L.; O'Connor, J.C. *Gait Analysis Laboratory: An Interactive Book & Software Package. Diskette 2*; Human Kinetics Publ.: 1992.