

Article

“Motorcycle ambulance” the policy to promote health and sustainable development in large cities

Korakot Apiratwarakul ¹, Takaaki Suzuki ², Ismet Celebi ³, Somsak Tiamkao ⁴, Vajarabhongsa Bhudhisawasdi ¹, Chatkane Pearkao ⁵ and Kamonwon Ienghong ^{1,*}

¹ Department of Emergency Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand; korakot@kku.ac.th (K.A.); md221@kku.ac.th (V.B.)

² Department of Emergency and Critical Care Medicine, University of Tsukuba Hospital, Tsukuba 305-8577, Japan; takaaki1986@hotmail.com

³ Department of Paramedic, Gazi University, Ankara 06500, Turkey; ismetcelebi@gazi.edu.tr

⁴ Department of Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand; somtia@kku.ac.th

⁵ Department of Adult Nursing, Faculty of Nursing, Khon Kaen University, Khon Kaen 40002, Thailand; porpea@kku.ac.th

* Correspondence: kamonwan@kku.ac.th; Tel.: +66-0433-66869

Abstract: Motorcycles can be considered a new form of smart vehicle when taking into account their small and modern structure and due to the fact that nowadays they are used in the new role of ambulance to rapidly reach emergency patients in large cities with traffic congestion. However, there is no study regarding the measuring of access time for motorcycle ambulances in large cities of Thailand. Therefore, this study aims to compare access times to patients between motorcycle ambulances and conventional ambulances, including analysis of the use of AED installed on motorcycle ambulances to contribute to the sustainable development of public health policies. In two hundred and seventy-one operations, it was found that the times from dispatch to vehicle (ambulance and motorlance) being en route and times from the 1669 center (center for command-and-control EMS in Thailand) call receipt to arrival on scene (response time) of motorcycle ambulances was shorter than that of a conventional ambulance and the use of AED on a motorcycle ambulance can increase the chances of survival for patients with cardiac arrest outside the hospital in public places where AEDs are not available and in all cases where access to an AED was less than eight minutes.

Keywords: emergency health services; emergency medical services; motorcycles; health policy; cardiac arrest; response time; two-wheelers; ambulances

1. Introduction

Motorcycles used as ambulances (motorlance) have been shown to greatly increase access to emergency patients. A study in many major cities around the world with traffic congestion found that motorlance access took less time than access with the traditional ambulance [1-3]. Due to its structure as a small vehicle, it can more easily access small alleys as well as being able to pass through gridlock traffic in confined areas. Motorlances must have a specific design, structure, color, and equipment installed inside [4,5]. The color used for the vehicle must be a medical emergency color code RAL 1016 sulfur yellow, which is the color that reflects the most clearly on the traffic route both for people with typical vision and also for those with color blindness problems. Reflective stickers are installed on the sides of the motorlance to increase the visibility of other vehicles, which boosts the level of safety during the operation. A wind shield is thought to prevent wind and dust that may blow onto the driver and includes the installation of a siren signal. As for drivers, they must pass a safety training course in the driving and best practices of

operation of the motorlance. Each operation consists of two members, the driver is an emergency medical technician (EMTs) or advanced emergency medical technicians (AEMTs), the passenger is a doctor or nurse [6-8]. Global positioning system (GPS) navigation is used to reach emergency patients in the shortest possible time [9]. In the future, there is an action plan for using the medical telemedicine system to pass on patient information from the scene to the hospital emergency room. In addition, the motorlances that are currently in design and manufacturing stages are also equipped with an automated external defibrillator (AED) on the vehicle [10,11]. Most of the motorcycle ambulances will be used in the role of speeding up patient access and initial assessment, but there are no studies on the use of motorlance AED (Table 1). Therefore, the researchers are interested in exploring the possible changes to public health policy by assigning motorlances with AED to reach patients in large cities. It has the potential to play an important role in the development of sustainable health care policies.

Table 1. Previous studies about motorlance in SCOPUS database

Authors	Main objective	AED study
Dennis et al. (2008) [12]	develop motorcycle-based ambulance protocols	no
Hofman et al. (2008) [6]	compare the access times of a traditional ambulance and motorlance	no
Peyravi et al. (2009) [13]	arrival time of EMS to the scene	no
Wilson et al. (2013) [14]	analyze vehicle time in transport	no
Ssebunya et al. (2016) [15]	delivered to health facilities supplied in pregnancy	no
Apiratwarakul et al. (2019) [16]	arrival time of EMS to the scene	no
Alaofe et al. (2020) [17]	assess the effect of emergency transportation interventions on the outcome of labor and delivery in LMICs	no
Apiratwarakul et al. (2020) [18]	motorlance operation time during the COVID-19 pandemic	no
Alarcon-Bernal et al. (2020) [19]	compare the access times of a traditional ambulance and motorlance	no
Apiratwarakul et al. (2020) [20]	compare the access times of a traditional ambulance and motorlance during major sporting events	no

EMS: Emergency medical services; LMICs: low-and middle-income countries; COVID-19: Coronavirus disease 2019.

AED in Thailand have been prescribed to be installed in public areas such as bus stations and shopping malls, but it has been found that there are still very few compared to the proportion of the area which may be due to the high price of the device, about three thousand US dollars per device [21,22]. Although there is a law in Thailand that requires an AED device to be available to be used in first aid. That means the general public can use this tool when seeing a patient undergoing an out-of-hospital cardiac arrest. Another problem that can be found is the use of an AED for people without knowledge and expertise, resulting in people in the community who do not dare to use such a tool which can be a serious problem, especially in Thailand [23-25]. Thailand does not specify or mandate a formal course for the use of an AED but rather it is considered basic knowledge from the primary to secondary school level, the rate of use of such machines there are few. Together with the installation of an AED in public areas, there are not enough numbers [26,27]. From the information of Khon Kaen Province, it was found that in Mueang District

which has a population of more than four hundred thousand people in an area of 953.4 square kilometers there are less than fifty AED devices in public areas.

Khon Kaen Provinces located in the northeastern region and has a population of about two million people. This city was grouped in the MICE provinces that are important to the country's economic development [28]. It has important geography as a gateway connecting the route to Indochina. The important activities of MICE consist of M = Meetings, I = Incentives, C = Conventions, and E = Exhibitions. All the activities involve a large number of people including the public health system that is directly and indirectly involved [29-31]. Public health standards require large gatherings to include first aid units and AEDs. Due to various limitations, there are very few AEDs to compare with the size of Khon Kaen province. Therefore, the idea of using motorlances to solve such shortage problems has been put forward

The aim of this study was to analyze the duration of accessing the patient by motorlance and develop the pattern of using an AED on a motorlance.

2. Materials and Methods

2.1. Study Design and Setting

A cross-sectional study was conducted on all motorlance operations in emergency medical services (EMS) at Srinagarind Hospital (medical school), Thailand, from January 2019 to December 2020. This hospital has an average of two thousand EMS operations per year. The area of study was in Mueang District, Khon Kaen Province, one of the groups in the designated MICE provinces.

2.2. Study Participants

All data regarding patients in the period of study treated with both traditional ambulance and motorlance delivery to access patients were collected. Cases with incomplete data and those that went out of EMS operation and did not find a patient were excluded from this study.

2.3. Data Collection

Data were recorded using a national standard operation record form for Thailand EMS consisting of demographic data (age, gender), operation time (activation time, response time), type of patients (trauma, non-trauma), type of first procedure on scene for all patients, time of AED access, and disposition of patients at hospital.

2.4. Definitions

Activation time was defined as times from dispatch to vehicle (traditional ambulance and motorlance) being en route. Response time was defined as times from 1669 center (center for command-and-control EMS in Thailand) call receipt to arrival on scene. AED access time was defined as time from dispatch to AED arrival on scene.

2.5. Study Size

The sample size was calculated based on the following formula [32]: $n = Z^2 [p(1 - P)]/d^2$. The variables in the formula are defined as follows: n = sample size; Z-score = 1.96 with confidence level 95%; P = expected proportion; and d = precision (margin of error). The estimate for P was made using data from a previously published study [16], we determined that a sample size of 271 would be required. Statistical analysis was performed with Khon Kaen University license (SPSS Inc., Chicago, IL, USA) by IBM SPSS for Windows version 26.0. Categorical data were presented as percentage, and continuous data were presented using mean and standard deviation. Univariate analysis was performed using a two-sample t-test for numerical data and Chi-squared test for comparing data between the groups

2.6. Ethical Considerations

Ethical approval was provided by the Khon Kaen University Ethics Committee for Human Research (HE641369). Requirement for informed consent from the patients was waived since patient confidentiality protection had been guaranteed, as patients were not identified by name, but by a unique study number.

2.7. Motorcycle ambulance

Motorlance modified from Honda New Forza 300 models (Figure 1). The width of the model is 75.30-centimeter, length 216.60-centimeter, height 118.90-centimeter, seat height 71.60-centimeter, ground clearance 13.90 centimeter and has a net weight of 192 kilogram. The fuel used was gasohol 91, 95, or E20 and fuel tank capacity was 11.5 liters. The three device boxes were installed in the rear of the motorlance, one box on the left and right sides, and one box behind the passenger seat. The box attached to the side of the motorlance was where the AED equipment was stowed for easy access. In addition, an airway management device and equipment related to stopping bleeding were prepared. The reflective stickers are installed to increase safety during operation along with emergency and sirens with radio communication in the vehicle.



Figure 1. The motorcycle ambulance model in Srinagarind Hospital, Thailand

3. Results

Two hundred seventy-one motorlance operations were examined over a two-year period, the characteristics of which are shown in Table 2. A total of 52.4% (N = 142) of the patients were male. The age group of the patients was mostly more than 60 years of age (32.2%). Operations time was most commonly performed between afternoon shifts (4 PM to 12 AM) 55.7%. Trauma patients accounted for 56.1% of all cases in the period of study. The level of operation was advanced (with a doctor) in 22.9% of cases and basic (without a doctor) in 77.1% of cases.

Table 2. Characteristics of the subjects with motorlance services (N=271)

Feature	Category	Frequency	Percent
Gender	Male	142	52.4
	Female	129	47.6
Age group	< 20	3	1.1
	20-29	12	4.4
	30-39	25	9.2
	40-49	64	23.6
	50-59	80	29.5
	≥ 60	87	32.2
Operation time (shift)	8 AM to 4 PM (Morning)	92	34.0
	4 PM to 12 AM (Afternoon)	151	55.7
	12 AM to 8 AM (Night)	28	10.3
Type of patients	Trauma	152	56.1
	Non-trauma	119	43.9
Level of operations	With doctor	62	22.9

Without doctor	209	77.1
----------------	-----	------

The average times from dispatch to vehicle (motorlance and traditional ambulance) being en route (activation time) for motorlance and ambulance in afternoon shift were 0.59 minutes and 1.45 minutes, respectively ($p = 0.004$; Table 3). The average motorlance response time in the afternoon shift was 6.12 minutes and ambulance response time was 9.10 minutes at the same shift.

Table 3. Comparing services time between motorlance and ambulance

Time	Shift	Motorlance (min) N=271		Ambulance (min) N=1860		p-Value
		Mean	SD	Mean	SD	
Activation time	Morning	0.57	0.12	1.12	0.15	0.010 *
	Afternoon	0.59	0.15	1.45	0.12	0.004 *
	Night	1.02	0.22	1.55	0.20	0.014 *
Response time	Morning	5.42	1.20	8.26	1.30	< 0.001 *
	Afternoon	6.12	1.40	9.10	1.20	< 0.001 *
	Night	5.52	2.02	8.20	1.50	< 0.001 *

SD = Standard Deviation. Min = minute. * Statistical significance

In the period of study forty-six patients used AED in motorlance, the characteristics of whom are shown in Table 4. 54.3% of patients were male with most of those falling in the age group of more than 60 years ($N = 25/46$). The afternoon shift (39.1%) was the most common time for the operation of AEDs used in motorlances to occur. Almost all of the motorlance operations (97.8%), were found to have no access to AED equipment installed in public areas. The average time from dispatch to AED arrival on scene (AED access time) was 5.02 minutes. After AED was used, the status of the patients when they arrived at the hospital was mostly in continuous resuscitation (95.7%).

Table 4. Characteristics of the subjects with AED used in motorlance (N=46)

Feature	Category	Frequency	Percent
Gender	Male	25	54.3
	Female	21	45.7
Age group	40-49	9	19.6
	50-59	12	26.1
	≥ 60	25	54.3
Operation time (shift)	Morning	12	26.1
	Afternoon	18	39.1
	Night	16	34.8
AED in scene area	Yes	1	2.2
	No	45	97.8
AED access time	Mean \pm SD (min)	5.02 \pm 0.45	
Status when arrive hospital	Continue CPR	44	95.7
	Death	2	4.3

AED = automated external defibrillator. SD = Standard Deviation. Min = minute. CPR = Cardiopulmonary resuscitation

4. Discussion

This study examines the use of motorcycles in the sustainable development of health policies. There are two main objectives: firstly, compare the duration of access to emergency patients between motorcycles with the traditional ambulance model and

second, we analyzed the use of the AED on the motorlance to reach emergency patients in areas without accessible AED in public places.

In the emergency operation with motorcycles, it was found that the population studied were similar in number of males and females. This is consistent with previous studies that found that emergency patients for the most part had no gender differences [3,5,7,8,16].

In terms of the age group who use emergency motorcycle services, it was found that the group that primarily used was mostly over sixty years old, followed by the fifty-to-fifty-nine-year age group which may be caused by data gathering between the years 2019-2020. It was during the outbreak of the COVID-19 virus. Most of the elderly do not want to go to the hospital or are restricted to travel by government policies and may be lacking regular medications, loss of a follow-up, and waited until the service is needed through the EMS [33,34]. Another reason may be from the hospital policy to send an emergency motorcycle out to check first emergency patients in case there is a risk of contracting COVID-19 in which case they will arrange an ambulance with appropriate protective equipment to continue to take care of patients [16].

As for the operating shift, most of the operations took place in the afternoon shifts. This is consistent with previous studies [35-37]. The main reason for this is the pattern of family relationships in Thai society that dictates that normally children of working age will go to work outside the accommodation, while parents will stay at home and take care of their grandchildren [38-40]. This kind of pattern results in the elderly having symptoms of illness and having to wait for their children to return to meet their parents at home after that request assistance in the EMS setting. Therefore, most operations take place in the afternoon shift, which is the time after work.

For Thailand's EMS, there are personnel in operation consisting of a doctor, nurse, paramedic, AEMTs and EMTs. The operation has two levels including advanced and basic levels. In the advanced operation level, there must be a doctor or nurse participating in the operation. An advanced-level deployment will occur when the patient's symptoms are notified and initial assessment is of critically ill patients, the command center will order the operation. In the case of urgent symptoms or when immediate treatment is not necessary, they will be ordered to operate at the basic level of operations [41-44].

We also found that when comparing the time in EMS from an emergency motorcycle with the ambulance in all shifts, activation time of the motorlance was shorter than the ambulance which is consistent with previous studies [45,46]. In this study ambulance and motorlance parking spots were in the same area. Therefore, the shorter activation time of motorlance is due to the lower number of operational personnel (only two operational personnel). Unlike an ambulance which requires four to five personnel to operate, making the length of time it takes for personnel to arrive at the vehicle different. However, according to EMS standards, the proper activation time should be less than two minutes which shows the readiness of personnel to carry out operations, with both models of emergency vehicles using less than two minutes. In terms of response time, motorlances took a shorter time to arrive at the scene than an ambulance. This is consistent with previous studies which point out one of the main benefits of a motorlance in that it is able to move in a small street or in traffic jams better [5,7,9,16,47,48]. Most of the areas in this study were conducted within the university campus and surroundings which have different conditions of traffic routes. The university road has two to four traffic lanes but in rush hour there will be a lot of vehicles. In areas around the university, small alleyways and unmarked 1 lane roads make up a majority of the grid, which in turn makes the direction giving and route to the patient complicated. Motorlances are clearly more advantageous in these types of environments. However, the period of EMS time (activation and response time) of motorlances and ambulances were shorter than the previous study [16], possibly due to the study period during the epidemic of the COVID-19 virus when traffic conditions are more flexible than usual, along with a number of

students and staff working from home causing the number of vehicles on the traffic route to decrease resulting in access to patients faster than normal operation [39,42,45].

In addition, the use of an AED on emergency motorcycles will be used in the event of a notification that there is an emergency patient who is unconscious, collapses or stops breathing. Due to limitations of the installation of AEDs in public places, this study applied the new health policy of bringing an AED to be installed on an emergency motorcycle to solve two problems. Firstly, the lack of publicly installed AEDs and secondly, the lack of expertise in using the equipment. It was found that the length of time to reach the patient is less than eight minutes which is the standard assigned to the emergency medical unit to reach out-of-hospital cardiac arrest patients to increase the chance of survival [22,24,27]. The study also found that almost all areas that reach emergency patients do not have AED equipment installed in public places which corresponds to the nature of residence. Most of them live in one house or in an allotted village which is far from the area where an AED is located, such as a bus station or shopping mall. The modification of health policy will increase chances of survival of the people in the community [49,50]. As can clearly be seen from the study, almost all of the patients that are accessed with an AED-equipped emergency motorcycle have a chance to continue CPR at the hospital. The health sustainable development policy with the use of an emergency motorcycle plays an important role in reducing access time to patients. In addition, the installation of the AED on the motorlance increases the chance of survival which is considered a worthwhile investment [51-53]. In countries with limited budgets, installing AEDs in public areas is an important policy that both the government and private sectors should pay attention to and jointly develop, but during the waiting period for development, the application of available resources by bringing an emergency motorcycle to install an AED would be beneficial to people in the community, especially patients with out-of-hospital cardiac arrest. In addition, the role of emergency motorcycles differs depending on the context of each country such as in European countries an emergency motorcycle will be used for assessment of disaster situations to assess if the area is safe. This includes early access to assess the resources needed at the scene. In Korea, an emergency motorcycle was used to reach out-of-hospital cardiac arrest patients to increase chances of survival. In African countries motorlances were used to bring pregnant women to undergo antenatal care at community health centers, including taking pregnant women to deliver their babies at the hospital. Not surprisingly, motorlances are significantly cheaper to acquire and operate when compared with ambulances in the form of regular cars. Therefore, it is a new challenge in applying motorcycles to perform duties in the development of public health policies especially in large urban areas with traffic congestion. Limitations of the vehicle found that emergency motorcycles are unable to operate in conditions of severe rain, poor visibility or during severe wind breaks because it may be dangerous to the operator. It was also found that at a distance of more than ten kilometers, the operation with an emergency motorcycle takes a similar amount of time to reach the patient as an ambulance. Therefore, in such cases, the use of an ambulance is more beneficial in terms of its ability to transport patients.

Limitation of the study was data collection gathered from a single EMS which may look at the study population differently from other agencies. Therefore, information should be collected from many study agencies.

5. Conclusions

The motorlance plays an important role in reaching patients quickly, both in terms of time after receiving the notification call to vehicle (activation time) as well as traveling from the hospital to the scene (response time). It takes a shorter time than a normal model of ambulance. In addition, the installation of AEDs on emergency motorcycles increases the opportunity for people with cardiac arrest outside the hospital to access prompt treatment and this means a higher chance of revival. This is an important policy in public health for sustainable development in large city areas.

6. Future work

There may be some modifications to the vehicle configuration from a motorcycle to another form of three-wheel vehicle (Tuk Tuk) which is unique to Thailand, larger than a motorcycle but smaller than a car. Tuk Tucs can solve emergency situations that motorcycles cannot, such as operating during heavy rain or strong winds. The Tuk Tuk can install rainproof plastic to prevent rain, resulting in a greater ability to perform and create a better environment for the operator and emergency patients. Further developments such as was telemedicine systems may be installed such as a real time camera recording so that EMS members or doctors who are stationed at the hospital can see the patient's condition to be assessed promptly and deliver a more timely treatment, including possibly installing a navigation system from the hospital to the scene by using the shortest travel time.

Author Contributions: Conceptualization, K.A., K.I. and C.P.; methodology, K.A. and T.S.; validation, K.A. and I.C.; formal analysis, K.A. and K.I.; investigation, K.A. and K.I.; resources, S.T. and V.B.; data curation, K.A. and K.I.; writing—original draft preparation, K.A. and K.I.; writing—review and editing, K.A., K.I. and C.P.; visualization, K.A.; supervision, S.T. and V.B.; project administration, K.A. and K.I.; All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Research and Graduate Studies, Khon Kaen University, Thailand. (RP64-7/002)

Acknowledgments: The authors would like to express our sincere gratitude to Josh Macknick for acting as an English consultant.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Peleg, K.; Pliskin, J.S. A geographic information system simulation model of EMS: reducing ambulance response time. *Am. J. Emerg. Med.* **2004**, *22*, 164-170.
2. Bahadori, M.; Ravangard, R.; Teymourzadeh, E. Development of Emergency Medical Services (EMS) in Iran: Components of Access. *Int. J. Collab. Res. Intern. Med. Public Health.* **2012**, *4*, 387-394.
3. Soares-Oliveira, M.; Egipto, P.; Costa, I.; Cunha-Ribeiro, L.M. Emergency motorcycle: has it a place in a medical emergency system? *Am. J. Emerg. Med.* **2007**, *25*, 620-622.
4. Jafari, M.; Shakeri, K.; Mahmoudian, P.; Bathaei, S.A. Innovation in the use of motor ambulance for prehospital emergency care. *J. Educ. Health Promot.* **2019**, *8*, 38.
5. Kiefe, C.C.; Soares-Oliveira, M. Medical emergency motorcycles: are they safe? *Eur. J. Emerg. Med.* **2008**, *15*, 40-42.
6. Hofman, J.J.; Dzmadzi, C.; Lungu, K.; Ratsma, E.Y.; Hussein, J. Motorcycle ambulances for referral of obstetric emergencies in rural Malawi: Do they reduce delay and what do they cost? *Int. J. Gynaecol. Obstet.* **2008**, *102*, 191-197.
7. Riley, P. Motorcycle medics. *JEMS.* **2000**, *25*, 32-41.
8. Pons, P.T.; Haukoos, J.S.; Bludworth, W.; Cribley, T.; Pons, K.A.; Markovchick, V.J. Paramedic response time: does it affect patient survival? *Acad. Emerg. Med.* **2005**, *12*, 594-600.
9. Chinprasatsak, S.; Satthapong, S.; Kotruchin, P.; Maporn, K. Telegraphic medicine systems improve medical diagnosis in pre-hospital settings: A pilot study in a tertiary care hospital. *J. Med. Assoc. Thai.* **2017**, *100*, 686-691.
10. Bigdeli, M.; Khorasani-Zavareh, D.; Mohammadi, R. Pre-hospital care time intervals among victims of road traffic injuries in Iran. A cross-sectional study. *BMC Public Health.* **2010**, *10*, 1-7.
11. Vukmir, R.B. Survival from prehospital cardiac arrest is critically dependent upon response time. *Resuscitation.* **2006**, *6*, 229-234.
12. Dennis, R.A. Low cost ambulance for developing countries. In *2008 5th IET Seminar on Appropriate Healthcare Technologies for Developing Countries.* **2008** May 21, 1-7.
13. Peyravi, M.R.; Toubaei, F.; Pourmohammadi, K. The efficiency of motorlance in comparison with ambulance in Shiraz, southern Iran. *Iran. Red Crescent Med. J.* **2009**, *11*, 330-333.
14. Wilson, A.; Hillman, S.; Rosato, M.; Skelton, J.; Costello, A.; Hussein, J.; MacArthur, C.; Coomarasamy, A. A systematic review and thematic synthesis of qualitative studies on maternal emergency transport in low-and middle-income countries. *Int. J. Gynaecol. Obstet.* **2013**, *122*, 192-201.
15. Ssebunya, R.; Matovu, J.K. Factors associated with utilization of motorcycle ambulances by pregnant women in rural eastern Uganda: a cross-sectional study. *BMC PREGNANCY CHILDB.* **2016**, *16*, 1-11.
16. Apiratwarakul, K.; Ienghong, K.; Mitsungrern, T.; Kotruchin, P.; Phungoen, P.; Bhudhisawasdi, V. Use of a motorlance to deliver emergency medical services; a prospective cross sectional study. *Arch. Acad. Emerg. Med.* **2019**, *7*, e48.
17. Ehiri, J.; Alaofè, H.; Asaolu, I.; Chebet, J.; Esu, E.; Meremikwu, M. Emergency transportation interventions for reducing adverse pregnancy outcomes in low-and middle-income countries: a systematic review protocol. *Syst. Rev.* **2018**, *7*, 1-9.

18. Apiratwarakul, K.; Ienghong, K.; Bhudhisawasdi, V.; Gaysonsiri, D.; Tiamkao, S. Response Times of Motorcycle Ambulances during the COVID-19 Pandemic. *Open Access Maced. J. Med. Sci.* **2020**, *8*, 526-529.
19. Alarcón-Bernal, Z.E.; Aceves-García, R.; Rojas-Arce, J.L. Location of Bases for Prehospital Services. In *Data Analysis and Optimization for Engineering and Computing Problems* **2020**, 73-84, Springer, Cham.
20. Apiratwarakul, K.; Ienghong, K.; Gaysonsiri, D.; Buranasakda, M.; Bhudhisawasdi, V.; Tiamkao, S. Role of motorcycle-based ambulance (motorlance) in major sporting events. *J. Med. Assoc. Thai.* **2020**, *103*, 15-17.
21. Allan, K.S.; Jefkins, T.T.; O'Neil, E.; Dorian, P.; Lin, S. Mandating Training Is Not Enough: The State of Cardiopulmonary Resuscitation and Automated External Defibrillator Training in Ontario Schools. *Can. J. Cardiol.* **2021**, *3*, 822-826.
22. Lewko, J.; Milewski, M.; Kowalewska, B.; Jankowiak, B.; Sokołowska, G.; Milewski, R. Is Cluster Analysis the Appropriate Statistical Method for Planning the Optimal Locations for Automated External Defibrillators? *Stud. Log. Gramm. Rhetor.* **2020**, *64*, 7-14.
23. Smereka, J.; Aluchna, M.; Aluchna, A.; Szarpak, L. Preparedness and attitudes towards medical emergencies in the dental office among Polish dentists. *Int. Dent. J.* **2019**, *69*, 321-328.
24. Smith, C.M.; Griffiths, F.; Fothergill, R.T.; Vlaev, I.; Perkins, G.D. Identifying and overcoming barriers to automated external defibrillator use by GoodSAM volunteer first responders in out-of-hospital cardiac arrest using the Theoretical Domains Framework and Behaviour Change Wheel: A qualitative study. *BMJ Open.* **2020**, *10*, e034908.
25. O'Callaghan, P.A.; Swampillai, J.; Stiles, M.K. Availability of automated external defibrillators in Hamilton, New Zealand. *N. Z. Med J.* **2019**, *132*, 75-82.
26. Rishipathak, P.; Bhandari, M.; Hinduja, A. Retention of Cognitive Skills Acquired in Heartsaver First Aid CPR AED Course of American Heart Association (AHA), USA among Non-medical Students in Pune city. *Indian J. Public Health Res. Dev.* **2019**, *10*, 742-744.
27. Post, E.G.; Schaefer, D.A.; Biese, K.M.; Cadmus-Bertram, L.A.; Watson, A.M.; McGuine, T.A.; Brooks, M.A.; Bell, D.R. A comparison of emergency preparedness between high school coaches and club sport coaches. *J. Athl. Train.* **2019**, *54*, 1074-1082.
28. Tunming, P.; Chaigasem, T.; Siri Wong, P. The increasing of potential in tourism logistics supply chain to Khon Kaen ME city, Thailand. *Afr. J. Hosp. Tour. Leis.* **2019**, *8*, 1-12.
29. Pathumporn, J.; Nakapaksin, S. Accommodation management for integrated sustainable tourism management. *Int. J. Sustain. Dev. Plan.* **2016**, *11*, 236-244.
30. Kantawateera, K.; Naipinit, A.; Sakolnakorn, TP.; Kroeksakul, P. Tourist transportation problems and guidelines for developing the tourism industry in Khon Kaen, Thailand. *Asian Soc. Sci.* **2015**, *11*, 89-95.
31. Maneenetr, T.; Naipinit, A.; Tran, T.H. Guidelines to promote local community participation in developing agrotourism: A case study of Ban Mor Village, Sam Sung District, Khon Kaen Province, Thailand. *Asian Soc. Sci.* **2014**, *10*, 178-186.
32. Al-Wathinani, A.M.; Schwebel, D.C.; Al-Nasser, A.H.; Alrugaib, A.K.; Al-Suwaidan, H.I.; Al-Rowais, S.S.; AlZahrani, A.N.; Abushryei, R.H.; Mobrad, A.M.; Alhazmi, R.A.; et al. The Prevalence of Risky Driving Habits in Riyadh, Saudi Arabia. *Sustainability.* **2021**, *13*, 7338.
33. Thornton, J. Covid-19: A&E visits in England fall by 25% in week after lockdown. *BMJ.* **2020**, *369*, m1401.
34. Apiratwarakul, K.; Kongudom, N.; Kotruchin, P.; Phungoen, P. Visits to the emergency department during the 2020 COVID-19 outbreak in Thailand. *J. Med. Assoc. Thai.* **2021**, *104*, S1-S4.
35. Faramand, Z.; Frisch, S.O.; Martin-Gill, C.; Landis, P.; Alrawashdeh, M.; Al-Robaidi, K.A.; Callaway, C.W.; Al-Zaiti, S.S.; Diurnal, weekly and seasonal variations of chest pain in patients transported by emergency medical services. *Emerg. Med.* **2019**, *36*, 601-607.
36. Maudet, L.; Sarasin, F.; Dami, F.; Carron, P.N.; Pasquier, M. Emergency medical services: COVID-19 crisis. *Rev. Med. Suisse.* **2020**, *16*, 810-814.
37. Croda, J.H.; Garcia, L.P. Immediate health surveillance response to COVID-19 epidemic. *Epidemiol. Serv. Saude.* **2020**, *29*, e2020002.
38. Thanapop, S.; Thanapop, C. Work ability of Thai older Workers in Southern Thailand: a comparison of formal and informal sectors. *BMC Public Health.* **2021**, *21*, 1218.
39. Suratana, S.; Tamornpark, R.; Apidechkul, T.; Srichan, P.; Mulikaburt, T.; Wongnuch, P.; Kitchanapaibul, S.; Yeemard, F.; Udpplong, A. Impacts of and survival adaptations to the COVID-19 pandemic among the hill tribe population of northern Thailand: A qualitative study. *PLoS One.* **2021**, *16*, e0252326.
40. Bhuanantanondh, P.; Buchholz, B.; Arphorn, S.; Kongtip, P.; Woskie, S. The Prevalence of and Risk Factors Associated with Musculoskeletal Disorders in Thai Oil Palm Harvesting Workers: A Cross-Sectional Study. *Int. J. Environ. Res. Public Health.* **2021**, *18*, 5474.
41. Pattanarattanamolee, R.; Sanglun, R.Y.; Nakahara, S. Community-Based First Responder Network in Rural Thailand: A Case Study of Out-of-Hospital Cardiac Arrest. *Prehosp. Disaster Med.* **2021**, *36*, 234-236.
42. Prommoon, P.; Phibalsak, T.; Netwachirakul, J.; Mekthat, M.; Jitpi boon, W.; Sangthong, R. Epidemiological Characteristics of Traffic and Non-traffic Injuries and Quality of Emergency Medical Services in Southern Thailand. *J. Health Sci. Med. Res.* **2021**, *39*, 273-282.
43. Pochaisan, O.; Pattanarattanamolee, R.; Pongphuttha, W.; Chadbunchachai, W.; Nakahara, S. Development of an emergency medical services system in Thailand: Roles of the universal health coverage and the national lead agency. *Emerg. Med. Australas.* **2021**, *33*, 1-3.

44. Riyapan, S.; Chantanakomes, J.; Ruangsomboon, O.; Chaisirin, W.; Limsuwat, C.; Prapruetkit, N.; Chakorn, T.; Monsomboon, A. Assessment of prehospital management of patients transported to a Thai University Hospital. *Siriraj Med. J.* **2020**, *72*, 287-295.
45. Alanazy, A.R.; Fraser, J.; Wark, S. Organisational factors affecting emergency medical services' performance in rural and urban areas of Saudi Arabia. *BMC Health Serv. Res.* **2021**, *21*, 1-8.
46. Kelker, H.; Yoder, K.; Musey, P.; Harris, M.; Johnson, O.; Sarmiento, E.; Vyas, P.; Henderson, B.; Adams, Z.; Welch, J.L. Prospective study of emergency medicine provider wellness across ten academic and community hospitals during the initial surge of the COVID-19 pandemic. *BMC Emerg. Med.* **2021**, *21*, 36.
47. Alassaf, W.; Al Hamid, S.; Kentab, O.; Al Otaibi, A.; Al Mufareh, B. EMS operation in a female restricted university, Saudi prospective. *Int. J. Emerg. Med.* **2021**, *14*, 16.
48. Walker, K.J.; Jiarpakdee, J.; Loupis, A.; Tantithamthavorn, C.; Joe, K.; Ben-Meir, M.; Akhlaghi, H.; Hutton, J.; Wang, W.; Stephenson, M.; et al. Predicting Ambulance Patient Wait Times: A Multicenter Derivation and Validation Study. *Ann Emerg Med.* **2021**, *78*, 113-122.
49. Sekendiz, B. Incidence, bystander emergency response management and outcomes of out-of-hospital cardiac arrest at exercise and sport facilities in Australia. *Emerg. Med. Australas.* **2021**, *33*, 100-106.
50. Pemberton, K.; Bosley, E.; Franklin, R.C.; Watt, K. Epidemiology of pre-hospital outcomes of out-of-hospital cardiac arrest in Queensland, Australia. *Emerg. Med. Australas.* **2019**, *31*, 821-829.
51. Katipoglu, B.; Madziala, M.A.; Evrin, T.; Gawlowski, P.; Szarpak, A.; Dabrowska, A.; Bialka, S.; Ladny, J.R.; Szarpak, L.; Konert, A.; et al. How should we teach cardiopulmonary resuscitation? Randomized multi-center study. *Cardiol. J.* **2021**, *28*, 439-445.
52. Abelairas-Gómez, C.; Schroeder, D.C.; Carballo-Fazanes, A.; Böttiger, B.W.; López-García, S.; Martínez-Isasi, S.; Rodríguez-Núñez, A. KIDS SAVE LIVES in schools: cross-sectional survey of schoolteachers. *Eur. J. Pediatr.* **2021**, *180*, 2213-2221.
53. Nakagawa, N.K.; Oliveira, K.M.; Lockey, A.; Semeraro, F.; Aikawa, P.; Macchione, M.; Carvalho-Oliveira, R.; Gouvêa, G.B.; Boaventura, A.P.; Maiworm, A.I.; et al. Effectiveness of the 40-minute handmade manikin program to teach hands-on cardiopulmonary resuscitation at school communities. *Am J Cardiol.* **2021**, *139*, 126-130.