

Barriers to access of healthcare services for rural women – applying gender lens on TB in a rural district of Sindh, Pakistan

Shifa Salman Habib¹, Wafa Zehra Jamal^{1,*}, Syed Mohammad Asad Zaidi¹, Junaid-ur-Rehman Siddiqui², Hira Mustafa Khan², Jacob Creswell³, Sri Chand⁴, Anna Versfeld³

¹Community Health Solutions, Karachi, Pakistan

²Aahung, Karachi, Pakistan

³The Stop TB Partnership, Geneva, Switzerland

⁴Pakistan Tuberculosis Program

* correspondence: wafa.zj@yahoo.com, 00923002889961, ORCID:0000-0002-3509-5260

Abstract

Background: Women in rural districts of Pakistan face numerous barriers to healthcare, rendering gender-responsive health programming important, including for Tuberculosis (TB). This study was conducted to assess the general understanding of TB and of access to healthcare for women, as a first step towards implementation of a gender responsive TB programme in TandoAllayar, a rural district of Pakistan.

Methods: A total of 36 participants were interviewed. The focus group discussion guide comprised of questions on: (1) family/household dynamics (2) community norms (3) healthcare system (4) women's access to healthcare (5) TB Awareness, and (6) women's access to TB Care.

Results: Limited autonomy in household financial decision-making, disapproval of unassisted travel, long travel time, lack of prioritization of spending on women's health and inadequate presence of female health providers, were identified as barriers to access of healthcare for women, higher in younger women. Facilitators to access of TB care included a reported lack of TB-related stigma, moderate knowledge about TB disease, and broad understanding of tuberculosis as a curable disease. Other suggested facilitators include health facilities closer to the villages and the availability of higher quality services.

Conclusion: Significant barriers are faced by women in access to TB care in rural districts of Pakistan. Programme implementers in high burden countries, should shift towards improved gender-responsive TB programming.

Keywords: tuberculosis, patriarchy, gender barriers, women

Introduction:

Gender is an important determinant of health, particularly in low-and-middle income countries (LMICs). [1,2] Pakistani society is highly patriarchal, gender-based disparities are common between women and men in health, education, income opportunities, employment opportunities, personal security, control over assets and participation in the politics [3,4]. Pakistan ranks 151st on the World Economic Forum's Global Gender Gap Report, third from last on the list. [5] Women face numerous, well-documented barriers to healthcare, including limited decision-making powers [6,7], household care responsibilities, restrictions on travelling alone, and the prioritisation of male family members' health. [8] This context renders gender-responsive health programming particularly important. Yet there is dearth of initiatives that explore the barriers for women to tuberculosis care.

Globally, tuberculosis (TB) is one of the top six causes of mortality among women aged 15–49 years. [9] Pakistan has the fifth highest burden of TB in the world and an estimated 42% of cases remain undiagnosed. [10] In line with global patterns, men experience a higher burden of TB disease than women in Pakistan. However, the cumulative reported male-female (M:F) ratio for TB notifications, at 1.1, [11] is one of the lowest in the world, indicating a proportionally high burden of TB in women. (Globally males have twice the TB case notifications compared to females. [12]) There are also wide discrepancies in the M:F ratios between provinces and regions in Pakistan, suggesting the impact of socio-cultural and economic factors on TB infection and notifications. [13] Women of reproductive age (15- 34 years) also have a marginally higher burden of TB than men in the same age category. [14]

The Stop TB Partnership's TB REACH Wave 7 funding, supported by Global Affairs Canada, provided funding to projects that integrated a women's empowerment approach into their TB programming. With this support Community Health Solutions (CHS), a social enterprise based in Pakistan, implemented a project in Tando Allayar, a rural district of the Sindh Province. The Sindh Province has the country's highest TB prevalence rate (454 per 100,000) and over two thirds 67.3% of the population people live under the poverty line. [15] Almost half of children aged under five are reportedly stunted or short for their age and a quarter are severely stunted. [16]

According to unpublished NTP surveillance data, district Tando Allayar has one of the lowest recorded case detection rates in Pakistan (29%) and females account for 46% of case notification. Health services are limited. Only 29.1% of ever married women in Tando Allayar complete four or more antenatal care visits. [17] The population of 575,720 is served by limited and poorly resourced health facilities including 14 Basic Health Units, three Rural Health Centers and one 100-bed tertiary-care hospital. [18] There are two TB Basic Management Units for TB in the public sector, marked by frequent staff-absenteeism and

limited work hours. Hyderabad (40 km) and Kotri (50 km) are the closest large cities with public and private hospitals and facilities for TB care.

Noting the role that TB programming has in shifting harmful gender norms [13], TB REACH Wave 7 funded projects were required to incorporate aspects of women's empowerment. Supported by this wave of funding, CHS developed a Rural Tele-Healthcare Hub at Tando Allayar district. The Hub offered high quality TB diagnostics and treatment services to patients, including the use of digital chest x-rays and a microbiology laboratory and notified patients to the National TB Program. The Hub also linked under-employed female doctors in major cities to rural women via tele-medicine, supported by Sehat Kahani, a private technology startup providing telehealth solutions. [19] Onsite, a Lady Health Visitor (LHV) provided maternal and child health services and initiated treatment for people with TB, under supervision of the online physicians. The Hub was linked with direct outreach for TB awareness, screening and treatment into communities through an existing network of over 200 women volunteers, through a pioneering approach for community-mobilization developed by National Rural Support Programme (NRSP). [20] Under this approach, rural women are encouraged to advocate for their rights through village-based community organizations and participate in implementation of development initiatives. Our project was the first adaptation of this approach for TB. Through these female volunteers, households are mapped, screened and persons with presumptive TB are identified and linked to the hub with a travel voucher.

In order to validate the design and ensure commensurate implementation of a gender responsive TB programme, the first step was the implementation of an assessment of general understanding of TB and of access to healthcare for women at the intervention site. This paper outlines the findings of this study and provides useful insights into key considerations for gender responsive care in rural settings of Pakistan.

Methods:

Study Design

This study reports on exploratory focus group discussions (FGD), conducted using semi structured interview guides, to understand the barriers and facilitators of access to general healthcare and TB services. Conceptually we have drawn on the socioecological model (SEM) in the design and analysis phases for this study. [21] While the SEM identifies five nested, or hierarchical levels of influence; for our study we adapted the model to focus on the following four levels: individual, interpersonal, environmental/structural and community (Figure 1)




Figure 1: An adaptation of the socioecological model by McLeroy et al.

Participants

Women from three groups were included: younger lay community women aged 18 to 25 years; older lay community women aged 26 to 49 years; and community resource persons (CRPs) from the National Rural Support Programme (NRSP), the largest rural upliftment program in Pakistan, working with 3.6 million poor households. CRPs are rural women who are recruited into community organizations and provided with trainings on leadership, communication, and community support. They engage in a range of development initiatives, including healthcare provision. [22] The inclusion of CRPs and lay women was to allow for understanding the differences in perspectives that may arise from differential exposure and training. The lay community were divided into age-group categories to encourage participation among peer groups and to investigate the presence of age-related differences in barriers to care. CRPs were

recruited through the NRSP district support office and the lay community women in both groups were, in turn, identified and recruited through the CRPs through a mix of convenience and snowball sampling.

There were 12 women in each of these 3 groups (36 women in total) and total of six FGDs, using semi-structured questionnaires, were conducted, two within each of the three groups of women. The FGDs with the CRPs were carried out at the NRSP office whereas the discussions with the lay community women were carried out in the houses of CRPs. Each FGD lasted approximately one hour. Baseline demographic data was collected. Discussions covered household decision-making and family dynamics; perceived gender norms in the community; the concept of health and common health issues for women; inequities of existing healthcare system; knowledge, attitudes and perceptions around TB; and adequacy of local healthcare system for TB treatment. CRPs were also asked about their opinions about the need for a health programme focused on women in Tando Allayar and their motivation for being involved in such program as volunteers.

Data management and analysis

All the FGDs were conducted by a trained researcher, experienced in moderating focus groups discussions in a variety of settings. The FGDs were audio-recorded and transcribed in the local language by two research associates, who were both skilled in transcription. Transcripts were then translated into English language for the purpose of analysis. Data were stored electronically in password-protected encrypted computers. Data were manually analysed with deductive analysis based on the SEM. Additional inductive analysis using a grounded approach was used to ascertain other emerging themes and sub-themes during the process of transcription and analysis. The coding framework was refined through a series of meetings between the researchers as it was being developed.

Ethical considerations

Ethical approval for the study (Reference IRD_IRB_2020_03_007) was obtained from the Institutional Review Board (IRB) of Interactive Research & Development, registered with the Department of Health and Human Services, USA. Participation was voluntary, and all participants were informed about their right to withdraw at any time, or to refuse to answer any questions. Verbal consent was obtained from all participants, as was appropriate in a context of low rates of literacy and they were ensured of strict confidentiality. Women were transported to the study site by vehicles owned by NRSP. Refreshments were provided but no remuneration was provided to the participants. Research was discussed with and

approved by the authorities from Provincial Tuberculosis Programme and community leaders from NRSP prior to FGD initiation.

Results

Interpretation of the results that follow should be considered in light of participant characteristics. Marriage and children were among the reported characteristics as in Pakistan these are important markers of a woman's stage of life which may be a determinant of her views. The participant median age 33 years. The majority (29) of the women were married and 7 were unmarried at the time of FGD nor were they previously married. Most (27) of the married women were mothers, with the number of children ranging from one to nine, one participant was pregnant for the first time. Among the lay community women, a majority (70%) of the participants were working as labourers. A few participants also reported as having independent sources of income through cosmetic and sartorial work.

Individual level analysis

General health awareness and health seeking:

Health was defined by participants as the absence of illness and linked it with diet, hygiene, and use of medicines for illness. The common health problems identified were hepatitis, diabetes, hypertension, diarrhoea, TB and pneumonia.

"There is sugar and blood pressure. If someone with blood pressure issues consumes sugar, then they immediately become diabetic. Besides that, there is fever, cough, TB; these problems have become more prevalent."

-Lay community woman, 26 to 49 years group

Participants identified anaemia, diabetes, urinary tract infections (UTIs), menstrual cramps and TB as problems particularly facing women. Furthermore, participants also reported a recent increase in incidences of neonatal mortality, though reasons for this were not suggested.

"Many women lose their first child. Often the new-born's heart doesn't work or stops working after a few days."

-Lay community woman, 18 to 25 years group

TB was understood to be a contagious disease by the study participants and also described as "common", easily transmissible, and treatable. Nearly all participants either knew, or knew of, someone affected by TB, currently or previously.

“TB is a common disease and it is curable. I’ve heard when one person has TB in the household then it spreads to the entire family. If the person has children, then the children may also get it.”

-Community Resource Person

“My mother had TB and then my three-year old daughter caught it as well. Now they are complete[ly] fine after completion of treatment.”

-Community Resource Person

All participants were generally aware of the common symptoms of TB and identified persistent cough, bloody cough, breathing difficulty, fatigue, fever, and weight loss as being associated with TB. While some participants were able to identify bacteria as the cause of TB and were also knowledgeable about airborne transmission, more than half of the participants believed that TB could be caused by drinking cold drinks, eating oily foods, smoke and dust, and sexual intercourse. In terms of diagnosis, participants largely spoke of clinical evaluation by a doctor’s consultation and did not exhibit knowledge of the diagnostic tests that may be required. Participants reported being aware of a treatment “course” for TB, with some participants referring to a six-month time span. At the same time, participants reported first using traditional remedies (“*totkay*”) such as use of locally available herbs, self-medication, or going to local *hakims* for medical care. Going to the doctor in the city was resorted to after these methods had failed, generally when the cough persists for over a month or other symptoms such as fever or haemoptysis develop. Discussions centred on pulmonary TB and participants did not report any perceived difference in symptoms between men and women.

Overall, the CPRs demonstrated better knowledge and awareness about TB, its cause, prevention, and treatment, likely due to their experience as frontline workers in public health related projects.

Interpersonal level analysis

Highly patriarchal societal norms were described as infusing all aspects of life. Participants reported that major decisions such as expenditure on housing, children’s education and healthcare were controlled by men. Income earned by women themselves was also reported to be managed by their husbands. Women’s decision-making was found to be limited to routine household management such as cooking

meals, a realm where men were not reported to be involved. Experiences of greater female autonomy was only reported for households in *exceptional circumstances* whereby the husband was medically incapacitated.

“My husband says that children should be engaged in active labour work and earn money while I say that they should go to school.”

-Lay community woman, 18 to 25 years group

Men were perceived as the ones who were prioritized for receiving education, responsible for financially supporting the household, and responsible for performing physical labour. Spending over young boys was also considered more rewarding as they are believed to provide for the family as they grow up and become financially active. Girls were described to face neglect and not considered worthy of healthcare expense because they eventually leave the family with marriage.

Other barriers to TB care access reported for women were the same as women experience for general healthcare. The demands of household chores mean that women cannot rest or travel to seek care. Permission and accompaniment from a male partner or household member are required for women to travel, and this is not always provided, and women cannot necessarily cover the costs of the required journey.

“Men can stay alone in the hospital while there should be someone with the woman. When a woman goes to the hospital, her husband or brother goes with her. When the breadwinner of the house goes with the woman then everyone in the family gets worried. There are all kinds of resources in Kotri, but because of these reasons, women don't go.”

-lay community woman, 26 to 49 years group

“If we need to travel to the city for any purpose, we must seek our husband's permission and travel with a male family member. If the husband does not give us permission, we do not go”

-lay community woman, 26 to 49 years group

Similarly, women also felt that they could not seek care due to social responsibilities as they were the primary care takers for households, children and the elderly. It was also reported that their health was often deprioritized over health of the children, husbands, or brothers.

“If the son falls ill, he is sent to the hospital immediately but if the girl falls ill, she is told to take a tablet to make it better.”

-Lay community woman, 26 to 49 years group

As working women, the CRPs reportedly exercised relatively greater freedom with regards to their expenses, mobility, and personal decision-making compared to lay community women. However, they too reported depending on their husbands and elder male figures of the family for long-distance travelling, larger expenses, and high-impact decisions related to work, household, children and healthcare.

“If there is a big issue, for example, if there is a need to travel to Hyderabad or Karachi due to sickness, we obviously cannot travel without our husband or brother-in-law. We are not independent in doing such big things, but can take care of smaller issues ourselves.”

-Community Resource Person

Community level analysis

Women’s social functioning was largely restricted to the household, the fields, and tending to children. However, these restrictions were found to reduce with age, with younger women facing a much more restrictive environment than older women.

“ You know that ‘maahol’ (environment) is not suitable for girls that’s why we do not send them out of the house unless needed. Small kids roam in the streets all day, but we do not send our daughters outside as they grow slightly older”

-lay community woman, 26 to 49 years group

Participants identified and recognized the value of education and vocational skills in gaining financial autonomy and in turn attain larger control over their health and well-being.

“If a woman has vocational skills then she will never have to beg anyone for money.”

-Lay community woman, 26 to 49 years group

“ Women who are “khud-mukhtaar” (self-reliant) can take better care of their own health and well-being. They buy groceries and food items for the house and hence are free to consume those whichever way they like”

-Lay community woman, 18 to 25 years group

Participants were also cognisant of the challenges that women face in attaining education and skills. However, with the increasing inflation and economic hardship it was reported that some men are more accepting of women taking jobs for remuneration.

Environmental/ structural level analysis

A key barrier to health services emphasised in all focus groups was the lack of medical facilities in the village. The existing structure of healthcare system was described as inadequate to deal with the reported health challenges at the village-level. Participants reported that people travel to nearby cities including Kotri (50km), Hyderabad (40 km) and Karachi (200 km), to seek testing and treatment for TB since there was no clinic or doctor available in the village. The lack of facilities was also linked to poor individual-level knowledge about TB and delayed treatment seeking.

Transportation was a major limitation as public buses were either not available or delayed. Even when public transport was accessible and available, due to long routes and frequent stops it was challenging to reach the city for treatment and many patients were described as having suffered critically during the journey. Private transport was expensive and scarcely available. In addition, these challenges were further exacerbated for women due to their inability to travel alone and lack of financial autonomy.

Individuals' experiences in the hospital or clinic were reportedly highly dependent on the patients' socioeconomic class. At a village level, when medicines are available, they are reportedly provided only to influential individuals, who have either resource to offer as bribes or are connected to key resource persons at the health facility or the government. Impoverished patients were reportedly often neglected even if they have a *sehat* voucher (a coupon or a pass given to underprivileged households by the local government that allows them to avail healthcare services free of cost from public hospitals). Moreover, participants preferred visiting private facilities as it was easier to receive treatment compared to a public facility where doctors and staff were often inattentive, and it was difficult to receive services without references.

“Everything is done on referential requests [bribes]; if you have a reference then you immediately get the treatment or else you are made to wait and often the doctor leaves and you have to come again. Therefore, we don't go to Civil (Hospital) and prefer to go to private facilities.”

-lay community woman, 18 to 25 years group

To counter barriers faced by women, participants suggested that a well-equipped health facility, with quality service delivery at a reasonable distance from the villages around Tando Allayar city should be established.

“All we want is that there should be a hospital built in our neighbourhood so we can go there for our health issues”

They further highlighted that the healthcare providers, at these facilities, should be empathetic, free of prejudice, and committed to treating patients without any monetary incentive. Social injunctions on contact between men and women inhibit women seeking care from male healthcare providers and few participants suggested the idea of having female healthcare providers locally referred to as “lady doctors” at these facilities. Some participants also proposed the idea of financial aid for patients belonging to lower socio-economic status; this financial aid would include coverage for treatment and transport to and from the health facility.

CRPs had similar concerns regarding access to healthcare as that of lay community women. They stressed on the need for affordable and reliable health services and facilities in Tando Allayar at a reasonable distance from the surrounding villages. The participants strongly advocated for the need of household and community-level health awareness campaigns.

“ how can they go to the doctor; transport is an issue, cost of treatment is an issue, having small kids to take care of at home is an issue...”

-Community Resource Person

Discussion

At an individual level our findings about barriers to TB care access in the Sindh Province reflect those noted about healthcare in general for women in Pakistan. [23,24] Aligned with Khan and colleagues [25] we noted that at an individual level biomedical TB treatment seeking was inhibited by limited knowledge and initial use of traditional healthcare systems. Like Shaik and colleagues [26] we noted that the care roles played by women and the demands of household chores undermined their ability to leave the household space to seek care. We echo others in noting that the costs of care, limited finances, and

women's lack of independent income all inhibit care access [8, 27]; and in finding that the household prioritization of men and boys, and women's reliance on male-decision making undermines women's health autonomy [26, 28].

At a community level, we noted that the inter-personal norms, are very largely representative of social-cultural norms in the region. In addition, the lack of local healthcare facilities, and prohibitions on women – particularly young women being independent or moving about independently – all hamper treatment access. Finally, the difficulty attaining the skills that would allow for independent income generation also pose a barrier to women's ability to change their dependent situations.

At an environmental/structural level, the distances women must travel to access health services coupled with the lack of available, inexpensive transport makes accessing biomedical care difficult. The poor quality of public services encourages women to seek care in private healthcare facilities, where TB diagnosis is less likely, and testing and treatment expensive. Along with others [28, 29] we note that in the context of the social prohibition on men and women spending time together the lack of female healthcare providers undermines women's care access. Importantly, we further found care access depends on social standing, the utility of the health vouchers under different social protection schemes is limited, and the use of bribes is ubiquitous. Given that bribing is a social practice that occurs between men, may be suggesting an additional (and very largely unspoken) way in which women do not control their own healthcare decisions and access.

The structure of our research was locally unique in that our inclusion, and differentiation, of lay people and CRPS showed that even limited engagement in community work made a difference not only to health knowledge, but also in women's self-efficacy and their ability to attain charge of their life and health decisions. Our separation of age-cohorts, where age was used as a proxy for life-stage, further suggested that gender plays a greater limiting role on younger and unmarried women than it does on older women. Young women may be in need of additional support for gaining TB knowledge and accessing TB services. We also, to our surprise, found very low levels of TB-related stigma reported. This may be because the women in the focus groups had generally not been affected by TB themselves, but we have no reason to doubt the veracity of this claim, which was repeated across the three separate focus group discussions.

A range of facilitators to care access also emerged from our work. Already present facilitators included a reported lack of TB-related stigma, some (though limited) knowledge about TB infection and disease, and a broad-based understanding of tuberculosis as a curable disease. Facilitators that were suggested to

improve healthcare access included health facilities closer to the villages and the availability of better services. Improved services would have healthcare providers that demonstrate empathy, and serve everyone in need equally, irrespective of social standing and ability to pay bribes. It is notable, all the changes mentioned by participants that would facilitate care access were at an environmental/structural level, but no changes were suggested at a community or individual level. This may suggest that women do not see possibilities for change at a personal or community level and that basic gender equity awareness and advocacy will likely to be a starting point of any women's empowerment work.

While we have presented these barriers and facilitators as set in particular SEM realms, we note that each realm impacts the other. As others [30, 31] have noted, patriarchal social norms prevalent in the Pakistani society at large, especially in rural areas shape healthcare access. We found this at all levels. For example, individual levels barriers (such as lack of individual access to finances), are shaped by community norms in which women do not earn an independent income. Environmental/structural barriers, such as a lack of female doctors, are, in turn, shaped by the norm that women do not work independently. Despite this interplay, we believe that the conceptual separation is useful, for it highlights entry points for interventions to target for change processes.

This research further validated our approach to TB programming which was tailored to be gender sensitive in terms of service delivery mechanism, service providers, and the target population.

Additional areas which could be tackled to improve women's access to TB care include interventions that seek to shift the gender relation norms that undermine women's decision making and autonomy. Microfinance and vocational training may serve as potential approaches to increasing women's ability to generate income and secure livelihoods. The *Benazir Income Support Programme* (BISP), under which females from households identified below the poverty threshold receive unconditional cash transfers from the government, has shown to contribute to women's empowerment. Further research is needed to assess its impact on health outcomes, including for TB.

Conclusion

National TB programs should be informed by and respond to the ways in which gender effects individual, household and community practices that may increase risk of TB and influence health seeking behavior. Gender roles and specific needs will have to be considered when designing and implementing these programs to ensure that health systems serve to address gender inequalities and advance health outcomes equitably [32]. Pakistan is unusual in that the M:F TB burden ration, at 1.1 is one of the lowest

of any countries. Our paper has shown that a contributor to this may be the multiple layers of barriers women face accessing healthcare. We further showed that young women, who locally demonstrate a higher TB burden than men of the same cohort, were reported to experience greater challenges in TB care access than older women. This demonstrates a need for additional focus on younger women. This should include work that not only focused on environmental/structural changes, but also that focuses on women's empowerment at the individual and community level, for ultimately, it is these changes that will lead to systemic improvement that goes beyond the successful treatment of TB, and generates shifts towards a more gender-equitable world.

In showing that current systems that seek to support marginalized people to access healthcare, such as social protection schemes, may be critical, but are currently failing to do enough. Our work shows that more needs to be done to counter the class-related stigma and research into how this can be successfully implemented is needed. At the same time, our work confirms much of what has already been demonstrated in Pakistan. This suggests that while a gender-analysis should ideally be undertaken before any project design is set out, where this is not possible, programme implementers can, as we did, already shift towards improved gender-responsive programming by drawing on the available literature.

Limitations

The study was carried out in a single rural district in Sindh and may not be generalizable to urban parts of the country and rural parts of other provinces where there may be cultural differences. Furthermore, the interviews were conducted only among women, none of whom were affected by TB themselves; therefore, the study did not capture in-depth information about men's perspectives on barriers to healthcare for women, or the specific barriers to TB care.

Declarations

Ethical approval

An ethical approval was granted by the Institutional Review Board (IRB) at Interactive Research and Development (IRD)(IRD_IRB_2020_03_007). The IRB is registered with the U.S. Department of Health and Human Services (DHHS), Office for Human Research Protections. An informed consent was obtained from the participants.

Consent for publication

Not applicable.

Availability of data and materials

The dataset used for analysis during the current study is available from the corresponding author on reasonable request.

Conflict of interest:

The authors have no conflicts of interest to declare.

Funding

The implementation of the project was funded by Stop TB Partnership's TB REACH Wave 7, however, no funding was available for data analysis & interpretation, manuscript write up and publication of this study.

Authors contribution:

S.S.H., S.M.A.Z. and W.Z.J. were involved in conception of the study, finalizing the study design. W.Z.J and S.S.H conducted the literature review. J.R.S. and H.M.K were responsible for conducting the focused group discussions and data compilation in the form of transcripts. S.S.H and W.Z.J drafted the manuscript. J.C., A.V reviewed the drafts critically and finalized the manuscript. All authors reviewed and approved the final version to be published.

Acknowledgements

We would like to acknowledge the efforts of Elsbet Lodenstein from KIT university Netherlands for her contributions in the earlier stages of development of the manuscript concept. Also PTP Sindh, DHO, and district teams for their support during the implementation stages of the project. We acknowledge Sara Khurram and Iffat Agha from Sehat Kahani, Nabeela Shahid from NRSP and Aisha Ijaz from Aahung for their valuable contribution in this project.

References:

1. Fikree FF, Pasha O. Role of gender in health disparity: the South Asian context. *Bmj*. 2004 Apr 1;328(7443):823-6.

2. Iqbal N, Gkiouleka A, Milner A, Montag D, Gallo V. Girls' hidden penalty: analysis of gender inequality in child mortality with data from 195 countries. *BMJ global health*. 2018 Oct 1;3(5):e001028.
3. Chauhan K. Patriarchal Pakistan: women's representation, access to resources, and institutional practices. In *Gender Inequality in the Public Sector in Pakistan 2014* (pp. 57-87). Palgrave Macmillan, New York.
4. Ali SM. Gender and health care utilisation in Pakistan. *The Pakistan Development Review*. 2000 Oct 1:213-34.
5. Global Gender Gap Report 2020 [Internet]. World Economic Forum; 2020. Available from: http://www3.weforum.org/docs/WEF_GGGR_2020.pdf
6. Mumtaz Z, Salway S, Waseem M, Umer N. Gender-based barriers to primary health care provision in Pakistan: the experience of female providers. *Health policy and planning*. 2003 Sep 1;18(3):261-9.
7. Mahmood N. Women's role in domestic decision-making in Pakistan: implications for reproductive behaviour. *The Pakistan development review*. 2002 Jul 1:121-48.
8. Bari F. Women in Pakistan: Country briefing paper. Asian Development Bank; 2000.
9. TUBERCULOSIS IN WOMEN [Internet]. World Health Organization; 2019. Available from: https://www.who.int/tb/publications/tb_women_factsheet.pdf?ua=1
10. Qadeer E, Fatima R, Yaqoob A, Tahseen S, Ul Haq M, Ghafoor A, Asif M, Straetemans M, Tiemersma EW. Population based national tuberculosis prevalence survey among adults (> 15 years) in Pakistan, 2010–2011. *PloS one*. 2016 Feb 10;11(2):e0148293.
11. Global tuberculosis report 2019. Geneva: World Health Organization; 2019. Licence: CC BY-NC-SA 3.0 IGO
12. Global tuberculosis report 2020. Geneva: World Health Organization; 2020. Licence: CC BY-NC-SA 3.0 IGO
13. Khan, M. S., M. S. Khan, R. Hasan, and P. Godfrey-Faussett. "Unusual sex differences in tuberculosis notifications across Pakistan and the role of environmental/structural factors." *East Mediterr Health J* 19, no. 9 (2013): 821-5.
14. Codlin AJ, Khowaja S, Chen Z, Rahbar MH, Qadeer E, Ara I, McCormick JB, Fisher-Hoch SP, Khan AJ. Gender differences in tuberculosis notification in Pakistan. *The American journal of tropical medicine and hygiene*. 2011 Sep 1;85(3):514-7.
15. United Nations Development Programme. Multidimensional Poverty in Pakistan [Internet]. Available from: <https://www.undp.org/content/dam/pakistan/docs/MPI/Multidimensional%20Poverty%20in%20Pakistan.pdf>
16. Asim M, Nawaz Y. Child malnutrition in Pakistan: evidence from literature. *Children*. 2018 May;5(5):60.
17. Nutrition Profile - District Tando Allahyar [Internet]. 2021 [cited 5 July 2021]. Available from: http://www.rspn.org/wp-content/uploads/2019/03/Nutrition-Profile_Tando-Allahyar.pdf
18. National population data (District wise). [Internet]. National TB Control Program; 2021. Available from: http://ntp.gov.pk/ntp-old/uploads/pi/DISTRICTS_POPULATION_2015.pdf
19. Ltd. G. Sehat Kahani | The Story of Health [Internet]. Sehat Kahani. [cited 19 July 2021]. Available from: <https://sehatkahani.com/>
20. National Rural Support Programme [Internet]. Nrsp.org.pk. [cited 19 July 2021]. Available from: <https://nrsp.org.pk/>
21. Golden SD, Earp JA. Social ecological approaches to individuals and their contexts: twenty years of health education & behavior health promotion interventions. *Health education & behavior*. 2012 Jun;39(3):364-72.
22. Program for improved nutrition in Sindh (PINS) [Internet]. 2021. Available from: <http://www.rspn.org/wp-content/uploads/2020/05/Consolidated-Report-of-CRPs-on-WASH-Activities-SBCC-Toolkit.pdf>

23. Khan A. Mobility of women and access to health and family planning services in Pakistan. *Reproductive health matters*. 1999 Jan 1;7(14):39-48.
24. Country Gender Profile: Islamic Republic of Pakistan [Internet]. Japan International Cooperation Agency (JICA) Japan Development Service Co., Ltd (JDS); 2015. Available from: https://www.jica.go.jp/english/our_work/thematic_issues/gender/background/c8h0vm0000anjqj6-att/pakistan_2015.pdf
25. Khan A, Shaikh BT, Baig MA. Knowledge, awareness, and health-seeking behaviour regarding tuberculosis in a rural district of Khyber Pakhtunkhwa, Pakistan. *BioMed research international*. 2020 Apr 21;2020.
26. Shaikh BT, Hatcher J. Health seeking behaviour and health service utilization in Pakistan: challenging the policy makers. *Journal of public health*. 2005 Mar 1;27(1):49-54.
27. Riaz A, Zaidi S, Khawaja AR. Perceived barriers to utilizing maternal and neonatal health services in contracted-out versus government-managed health facilities in the rural districts of Pakistan. *International journal of health policy and management*. 2015 May;4(5):279.
28. Panezai S, Ahmad MM, Saqib SE. Factors affecting access to primary health care services in Pakistan: a gender-based analysis. *Development in practice*. 2017 Aug 18;27(6):813-27.
29. Rizvi N, Nishtar S. Pakistan's health policy: appropriateness and relevance to women's health needs. *Health policy*. 2008 Dec 1;88(2-3):269-81.
30. Bustamante-Gavino MI, Rattani S, Khan K. Women's Empowerment in Pakistan—Definitions and Enabling and Disabling Factors: A Secondary Data Analysis. *Journal of Transcultural Nursing*. 2011 Apr;22(2):174-81.
31. Rizvi N, Khan KS, Shaikh BT. Gender: shaping personality, lives and health of women in Pakistan. *BMC women's health*. 2014 Dec;14(1):1-8.
32. Tuberculosis Information Note [Internet]. The Global Fund; 2019 [cited 1 June 2021]. Available from: https://www.theglobalfund.org/media/4762/core_tuberculosis_infonote_en.pdf