

## **WNT/ $\beta$ -catenin signaling promotes TGF- $\beta$ -mediated activation of human cardiac fibroblasts by enhancing IL-11 production**

Edyta Działo<sup>1</sup>, Marcin Czepiel<sup>1</sup>, Karolina Tkacz<sup>1</sup>, Maciej Siedlar<sup>1</sup>, Gabriela Kania<sup>2</sup>, Przemysław Błyszczuk<sup>1,2#</sup>

<sup>1</sup>Department of Clinical Immunology, Jagiellonian University Medical College, Cracow, Poland

<sup>2</sup>Center of Experimental Rheumatology, Department of Rheumatology, University Hospital Zurich/University of Zurich, Switzerland

# correspondence:

Przemysław Błyszczuk, PhD

Department of Clinical Immunology,

Institute of Paediatrics,

Jagiellonian University Medical College,

Wielicka 265, 30-663, Cracow, Poland

Phone: +48-12-658-24-86

E-mail: [przemyslaw.blyszczuk@uj.edu.pl](mailto:przemyslaw.blyszczuk@uj.edu.pl)

**Abstract**

Cardiac fibrosis is a pathological process associated with development of heart failure. TGF- $\beta$  and WNT signaling have been implicated in pathogenesis of cardiac fibrosis, however little is known about molecular cross-talk between these two pathways. The aim of this study was to examine the effect of exogenous canonical WNT3a and non-canonical WNT5a in TGF- $\beta$ -activated human cardiac fibroblasts. We found that WNT3a and TGF- $\beta$  induced  $\beta$ -catenin-dependent response, whereas WNT5a prompted AP-1 activity. TGF- $\beta$  triggered profibrotic signature in cardiac fibroblasts, and co-stimulation with WNT3a or co-activation of the  $\beta$ -catenin pathway with GSK3 $\beta$  inhibitor CHIR99021 enhanced collagen I and fibronectin production and development of active contractile stress fibers. In the absence of TGF- $\beta$ , neither WNT3a nor CHIR99021 exerted profibrotic response. On a molecular level, in TGF- $\beta$ -activated fibroblasts WNT3a enhanced phosphorylation of TAK1 and production and secretion of IL-11 but showed no effect on Smad pathway. Neutralization of IL-11 activity with the blocking anti-IL-11 antibody effectively reduced profibrotic response of cardiac fibroblasts activated with TGF- $\beta$  and WNT3a. In contrast to canonical WNT3a, co-activation with non-canonical WNT5a suppressed TGF- $\beta$ -induced production of collagen I. In conclusion, WNT/ $\beta$ -catenin signaling promotes TGF- $\beta$ -mediated fibroblast-to-myofibroblast transition by enhancing IL-11 production. Thus, the uncovered mechanism broadens our knowledge on molecular basis of cardiac fibrogenesis and defines novel therapeutic targets for fibrotic heart diseases.

**Keywords:** cardiac fibroblasts, WNT3a, WNT5a,  $\beta$ -catenin, TGF- $\beta$  signaling, IL-11, cardiac fibrosis

**Abbreviations:** ECM (extracellular matrix), TGF- $\beta$  (transforming growth factor-beta),  $\alpha$ SMA (alpha smooth muscle actin),  $\gamma$ SMA (gamma smooth muscle actin), FN (fibronectin), IL-11 (interleukin-11), AP-1 (activator protein-1), TAK1 (TGF- $\beta$ -activated kinase 1), ERK1/2 (extracellular signal-regulated kinase 1/2), JNK (c-Jun N-terminal kinase), TCF (T-cell factor), LEF (lymphoid enhancer factor), END1 (endothelin 1), PDGF2 (platelet-derived growth factor), FGF2 (fibroblasts growth factor 2)

## 1. Introduction

Cardiac fibrosis refers to excessive accumulation of stromal cells and extracellular matrix (ECM) components in the myocardium and is often associated with cardiac pathologies. On the one hand, fibrotic scar develops following myocardial infarction to support tissue integrity and regeneration. On the other hand, hypertension, cardiac hypertrophy, local and systemic inflammation, co-morbidities, certain medications and aging may cause pathological tissue remodeling affecting cardiac function [1]. Increased stiffness of cardiac tissue and disturbed propagation of electrical impulses are direct outcomes of fibrosis, and these changes may lead to heart failure and arrhythmias [2]. Consequently, patients with persistent cardiac fibrosis show longer hospitalization and increased mortality [3].

ECM-producing cardiac fibroblasts represent cell type responsible for the maintenance of physiological architecture of the heart muscle. Once activated, cardiac fibroblasts play a central role in fibrotic processes in the heart. Activated cardiac fibroblasts show uncontrolled proliferation and ECM overproduction and may switch to myofibroblast phenotype. Myofibroblasts combine high secretory potential with phagocytic and contractile properties [4,5]. Alpha-smooth muscle actin ( $\alpha$ SMA)-positive stress fibers represent the most characteristic phenotypic feature of myofibroblasts in vitro. Cardiac injury or extracardiac triggers can activate fibroblast-to-myofibroblast transition. On a molecular level, this activation is mediated by profibrotic signaling molecules and/or mechanical stress [1,4,6].

Transforming growth factor-beta (TGF- $\beta$ ) represents the most prominent profibrotic cytokine, which plays a key role in cardiac fibrosis. TGF- $\beta$  triggers phenotypic switch of cardiac fibroblast into myofibroblasts, and the disruption or blockade of TGF- $\beta$  signaling was shown to protect from cardiac fibrosis in animal models [7]. Following binding of an active form of TGF- $\beta$  to the transmembrane TGF- $\beta$  type II receptor, the TGF- $\beta$  type I receptor is recruited and activated. Canonical TGF- $\beta$  receptor signal transduction involves Smad proteins and results in transcriptional response of Smad-responsive target genes [8]. In addition to Smad-dependent signaling, TGF- $\beta$  activates pathways independently of Smads. TGF- $\beta$ -activated kinase 1 (TAK1), phosphatidylinositol-3-kinase/Akt, RhoA-ROCK axis and mitogen-activated protein kinases p38, c-Jun N-terminal kinase (JNK) and extracellular signal-regulated kinase 1/2 (Erk1/2) represent examples of Smad-independent TGF- $\beta$  signaling [9]. Despite of pivotal role of TGF- $\beta$  in cardiac fibrosis, clinical use of direct or indirect TGF- $\beta$  inhibitors is currently limited due to problems with toxicity [10,11]

TGF- $\beta$  upregulates production of ECM components and multiple signaling molecules. Recent studies demonstrated that TGF- $\beta$  controls production of profibrotic WNTs and IL-11 in cardiac fibroblasts [12,13]. In humans, the family of WNT ligands represents 19 proteins. Extracellular WNTs bind to transmembrane receptor complexes and activate  $\beta$ -catenin-dependent and  $\beta$ -catenin-independent downstream mechanisms [14,15]. In a canonical response, stimulation with certain WNTs, like WNT3a, leads to stabilisation and nuclear translocation of  $\beta$ -catenin resulting in activation of the T-cell factor (TCF) and the lymphoid enhancer factor (LEF) transcription factors [16]. Other WNTs, like WNT5a,

can trigger transcriptional response independently of  $\beta$ -catenin. Such non-canonical WNT response involves activation of Erk1/2 and JNK signaling pathways regulating activity of the downstream activator protein 1 (AP-1) transcription factor, but also RhoA-ROCK axis [17,18]. Thus, non-canonical WNTs and TGF- $\beta$  control the same signaling pathways. Furthermore, TGF- $\beta$  was shown to stimulate production and secretion of canonical WNTs and to activate  $\beta$ -catenin in cardiac fibroblasts [19,20]. Noteworthy, animal studies specifically pointed to the importance of WNT/ $\beta$ -catenin pathway in cardiac fibrosis [21]

IL-11 represents pleiotropic cytokine induced by respiratory viruses such rhinovirus and parainfluenza as well as diverse profibrotic factors including TGF- $\beta$ , fibroblasts growth factor 2 (FGF2), endothelin 1 (EDN1), and platelet-derived growth factor (PDGF) [22,23]. Expression of IL-11 is strongly correlated with fibroblasts activation and associated with cardiac fibrosis.

Human cardiac fibroblasts-to-myofibroblasts transitions and development of cardiac fibrosis in mice were shown to be dependent on IL-11 signaling [13]. IL-11 plays an important profibrotic role not only in the cardiac tissue but also in the lung, kidney and liver [22,24]. A growing body of evidence points to IL-11 as a promising therapeutic target candidate in fibrotic disease. So far, however, little was known about molecular regulation of IL-11 production.

Herein, by using human cardiac fibroblasts we studied interplay between TGF- $\beta$  and canonical WNT3a and non-canonical WNT5a in fibrotic response. Our results indicate that WNT/ $\beta$ -catenin specifically controls profibrotic IL-11 in TGF- $\beta$ -dependent fibroblast-to-myofibroblast transition.

## Results

### *Canonical and non-canonical WNT responses in human cardiac fibroblasts*

Before investigating the effect of WNT signaling on differentiation of human cardiac fibroblasts, we addressed their responsiveness to exogenous WNT3a and WNT5a. In this approach, we used conditional medium of L-WNT3a, L-WNT5a (referred later as WNT3a and WNT5a, respectively) and of control L cell lines (used as controls for WNT3a and WNT5a) as well as TGF- $\beta$ , which is known to induce endogenous WNT production. Nuclear translocation of  $\beta$ -catenin represents the most characteristic feature of the canonical Wnt signaling. As expected, both WNT3a and TGF- $\beta$ , but not WNT5a, induced nuclear translocation of  $\beta$ -catenin in cardiac fibroblasts (Fig. 1A). In line with this finding, treatment with WNT3a or TGF- $\beta$  induced expression of  $\beta$ -catenin-responsive genes *AXIN2* and *TCF7* (Fig. 1B) confirming activation of canonical WNT pathway. For activation of non-canonical WNT pathway, we analyzed expression of *ATF2* and *JUN* and measured activity of the AP-1 transcription factor by using human cardiac fibroblasts transfected with the AP-1-reporter system (AP-1-hCF). WNT5a induced expression of *ATF2* and *JUN* in cardiac fibroblasts (Fig. 1B). Furthermore, we observed significantly upregulated AP-1 activity in AP-1-hCFs following treatment with WNT5a or TGF- $\beta$  (Fig. 1C). Collectively, our data confirmed that human cardiac fibroblasts differentially respond to WNT3a and WNT5a, and that TGF- $\beta$  could effectively activate canonical and non-canonical WNT pathways in these cells.

### *WNT3a and WNT5a differentially regulate gene expression in TGF- $\beta$ -activated cardiac fibroblasts*

TGF- $\beta$  plays a pivotal role in fibroblast activation. By employing a full genome transcriptomics, we analyzed transformation of human cardiac fibroblasts caused by TGF- $\beta$  signaling. Differential gene expression analysis demonstrated that cardiac fibroblasts 72h after treatment with TGF- $\beta$  showed deregulated expression of 313 genes including genes associated with cell differentiation (GO:0030154, 124 genes,  $p \leq 0.01$ ), cellular component organization (GO:0051128, 92 genes  $p \leq 0.01$ ), and extracellular matrix biology (GO:0031012, 41 genes,  $p \leq 0.01$ ). In particular, TGF- $\beta$  induced expression of collagens (*COL1A1*, *COL5A1*, *COL16A1*, *COL11A1*) and other extracellular matrix proteins involved in fibrotic response including *POSTN*, *FNI*, *TNC*, *VCAN* (Supp. Fig. 1).

WNTs are typically produced by activated cells and act as secondary triggers, therefore the effect of exogenous WNTs was studied in TGF- $\beta$ -activated fibroblasts additionally stimulated with WNT3a or WNT5a. Stimulation with WNT3a resulted in deregulation of 124 genes in TGF- $\beta$ -treated fibroblasts (Suppl. Fig. 2). Pathway analysis confirmed activation of WNT signaling (GO:0016055, 12 genes,  $p$  value  $\leq 0.0001$ ) and pointed to induction of cell differentiation (GO:0030154, 26 genes,  $p$  value  $\leq 0.0001$ ) and cellular component organization (GO:0051128, 13 genes,  $p$  value  $\leq 0.001$ ). WNT3a specifically up-regulated *ACTA2* (encoding alpha smooth muscle actin;  $\alpha$ SMA), *ACTG2* (encoding gamma smooth muscle actin;  $\gamma$ SMA) and *VCL* (encoding vinculin), genes involved in stress fibers

formation, as well as *LTBP1* (encoding latent-TGF- $\beta$ -binding protein 1) which regulates TGF- $\beta$  bioavailability (Fig. 2A). In contrast to profound effect of WNT3a, treatment with WNT5 upregulated expression of only 2 and downregulated 21 genes in TGF- $\beta$ -activated cells (Fig. 2B). Altogether, these results pointed to the relevance of WNT canonical arm in enhancing TGF- $\beta$ -mediated profibrotic response in cardiac fibroblasts.

#### *WNT/ $\beta$ -catenin signaling promotes TGF- $\beta$ -dependent fibroblast-to-myofibroblast transition*

Our previous data showed that WNT3a stimulated production of  $\alpha$ SMA and type I collagen in TGF- $\beta$ -activated fibroblasts [25]. Following these results and the described above transcriptomics data, we measured formation of  $\alpha$ SMA-positive stress fibers by immunocytochemistry. WNT3a in the absence of TGF- $\beta$  failed to increase the number of  $\alpha$ SMA-positive cardiac fibroblasts, whereas in TGF- $\beta$ -activated cells WNT3a effectively boosted formation of  $\alpha$ SMA-positive fibers (Fig. 3A) – a characteristic feature of myofibroblasts. Stress fibers are the major contractile structures in the cell, therefore in the next step we measured functional contraction of cardiac fibroblasts in a collagen gel. Obtained data confirmed that WNT3a enhanced contractile properties of TGF- $\beta$ -activated fibroblasts (Fig. 3B).

Next, we asked whether a specific pharmacological activation of the  $\beta$ -catenin-dependent pathway with GSK3 $\beta$  inhibitor CHIR99021 could promote myofibroblast phenotype in TGF- $\beta$ -activated cells. Treatment with CHIR99021 indeed up-regulated expression of *ACTA2*, *FNI* and *COL1A1* (Fig. 3C), as well as increased the corresponding intracellular protein levels of  $\alpha$ SMA, fibronectin (Fig. 3D) and the amount of secreted pro-collagen I  $\alpha$ 1 (Fig. 3E) in TGF- $\beta$ -activated fibroblasts. Of note, CHIR99021 in non-activated cells failed to trigger the myofibroblast signature.

Transcriptomics data suggested that WNT5a was not associated with profibrotic response. In fact, treatment with WNT5a suppressed TGF- $\beta$ -induced expression of *ACTA2* and *COL1A1* (Fig. 4A). On a protein level, however, we found no difference in  $\alpha$ SMA levels between cells treated with TGF- $\beta$  and with TGF- $\beta$  and WNT5a (Fig. 4B). Instead, WNT5a suppressed secretion of pro-collagen I  $\alpha$ 1 in TGF- $\beta$ -activated fibroblasts (Fig. 4C). Collectively, these data indicate that specific activation of the WNT/ $\beta$ -catenin pathway promotes fibroblast-to-myofibroblast transition induced by TGF- $\beta$ .

#### *WNT3a boosts TAK1 signaling and IL-11 production in activated cardiac fibroblasts*

In the next step, we aimed to identify molecular mechanism by which WNT3a enhances profibrotic TGF- $\beta$  response. In the canonical response TGF- $\beta$  activates Smad-dependent signaling. In contrast to TGF- $\beta$ , WNT3a did not induce phosphorylation of Smad2 and co-stimulation with WNT3a and TGF- $\beta$  showed similar activation of Smad2 to stimulation with TGF- $\beta$  alone (Fig. 5A). Treatment with TGF- $\beta$  activates not only Smad-dependent, but also non-canonical Smad-independent pathways, such as TAK1 signaling. Even though WNT3a alone failed to activate TAK1, we observed that in the

presence of TGF- $\beta$ , WNT3a effectively enhanced phosphorylation of TAK1 (Fig. 5B). These results suggest that WNT3a is implicated in strengthening the non-canonical arm of the TGF- $\beta$  signaling in cardiac fibroblasts.

Recently, IL-11 has been implicated in profibrotic TGF- $\beta$  response in cardiac fibroblasts [13], therefore we addressed whether WNT3a could regulate IL-11. We found that although WNT3a alone did not regulate production and secretion of IL-11, it enhanced IL-11 production in TGF- $\beta$ -activated cells (Fig. 5C). As TGF- $\beta$  induces WNT secretion in profibrotic response, we studied how blockade of WNT secretion with the porcupine inhibitor WNT-C59 affected TGF- $\beta$ -induced IL-11 production. We found that treatment with WNT-C59 reduced IL-11 levels in TGF- $\beta$ -activated cardiac fibroblasts (Fig. 5D). These results indicate that canonical WNT signaling controls TGF- $\beta$ -dependent IL-11 production.

#### *IL-11 mediates profibrotic WNT3a-mediated response in activated cardiac fibroblasts*

Our data suggested that WNT3a-dependent production of IL-11 might contribute to profibrotic response in cardiac fibroblasts. To address this hypothesis, we blocked IL-11 activity with anti-IL-11 neutralizing antibody in cardiac fibroblasts activated with TGF- $\beta$  and WNT3a and analyzed their fibrotic profile. Indeed, neutralizing anti-IL11 antibody effectively suppressed transcription of profibrotic genes *ACTA2*, *COL1A1*, *FNI* and *ACTG2* in cells co-stimulated with TGF- $\beta$  and WNT3a (Fig. 6A). In line with regulation of *ACTA2* and *FNI* transcripts, inhibition of IL-11 reduced protein level of  $\alpha$ SMA and fibronectin (Fig. 6B). Furthermore, inhibition of IL-11 in TGF- $\beta$ -activated and WNT3a-boosted cardiac fibroblasts reduced secretion of pro-collagen I  $\alpha 1$  (Fig. 6C) and suppressed their contractile properties (Fig. 6D). Collectively, obtained data confirmed profibrotic activity of IL-11 produced by cardiac fibroblasts activated with TGF- $\beta$  and WNT3a.

## Discussion

Activation of cardiac fibroblasts and their transition into myofibroblast phenotype plays a key role in cardiac fibrosis. TGF- $\beta$  and WNT ligands have been identified as important profibrotic factors involved in this process, however little is known about molecular cross-talk between these two pathways in cardiac fibrogenesis. Our data showed that specific activation of WNT/ $\beta$ -catenin pathway (either with WNT3a or with GSK3 $\beta$  inhibitor CHIR99021) promoted fibroblast-to-myofibroblast transition, mainly by enhancing TGF- $\beta$ -induced IL-11 production. In particular, WNT/ $\beta$ -catenin signaling specifically promoted formation of functional contractile stress fibers – an important feature of mature myofibroblasts. Previously, we showed that cardiac fibroblasts produced profibrotic WNTs upon treatment with TGF- $\beta$  [25]. Here, we confirmed that also exogenous canonical WNT3a could promote fibrotic processes in cardiac fibroblasts. This is not surprising, because WNT ligands are molecules that exert their signaling functions through the extracellular space. Of note, WNT secretion is a tightly regulated process resulting in controlled secretion either of soluble WNT proteins or WNTs incorporated into extracellular vesicles [25,27].

Previous studies highlighting importance of WNT/ $\beta$ -catenin pathway in cardiac fibrosis were commonly based on loss-of-function approaches [12,20]. Our findings not only complement published results by gain-of-function data, but also point to the involvement of multiple profibrotic pathways in fibroblast-to-myofibroblast transition. In fact, inability of WNT3a to trigger fibrotic response in non-activated fibroblasts (i.e. in the absence of TGF- $\beta$ ) was an important finding of this study. On a molecular level, WNT3a unlike TGF- $\beta$  was unable to activate Smad pathway. In contrast to our findings, treatment of mouse cardiac fibroblasts with recombinant WNT3a (in the absence of any other stimulus) activated myofibroblast phenotype by stimulating TGF- $\beta$  production and activating Smads [28]. These differences might result from differential activation status of the fibroblasts due to culture conditions, isolation procedure and/or cellular origin. Less likely, it might originate from the difference in WNT/ $\beta$ -catenin signaling between mouse and human cells. Nevertheless, results of both studies point to cooperative activity of TGF- $\beta$  and WNT3a in cardiac fibroblast-to-myofibroblast transition. We think that efficient phenotypic switch to myofibroblasts requires co-activation of multiple molecular pathways including Smad,  $\beta$ -catenin and TAK1 in cardiac fibroblasts. This multiple pathway co-activation hypothesis is further strengthened by the data showing that blockade of one of these pathways protects cardiac fibroblasts from activation and the heart from fibrosis [7,29,30].

In our experiments, WNT3a showed no effect on activation of Smad pathway, but enhanced TAK1 signaling in the presence but not in the absence of TGF- $\beta$ . These findings may suggest that once Smad signaling is on, output of  $\beta$ -catenin pathway regulates profibrotic response in cardiac fibroblasts. It should be noted however, that in addition to Smads, TGF- $\beta$  activates non-classical response dependent on mitogen-activated protein kinases, such as Erk1/2, the JNKs and the p38 isoforms [9]. Regulation of these non-classical TGF- $\beta$  pathways by WNT3a in cardiac fibroblasts remains, however, unknown. Regardless of the need for activation of the classical (Smad-dependent) or non-classical arm of TGF- $\beta$



response,  $\beta$ -catenin pathway seems to play a central role in regulating the degree of myofibroblast differentiation and thereby fibrosis. This study identifies TAK1 pathway as a molecular target of  $\beta$ -catenin in activated fibroblasts. Interestingly, previous data showed that inhibition of TAK1 signaling blocked the activation of  $\beta$ -catenin pathway in TGF- $\beta$ -activated cells [29]. In the light of both findings, a  $\beta$ -catenin-TAK1 feedback loop mechanism could be suggested. At this point, one needs to mention that TGF- $\beta$  activates  $\beta$ -catenin and our recent study indicated critical involvement of endogenous angiotensin II signaling in this process. Interestingly, cells lacking angiotensin II receptor 1 showed impaired TAK1 activation following TGF- $\beta$  treatment [31]. All these data point to the  $\beta$ -catenin-TAK1 positive feedback loop as a novel mechanism in TGF- $\beta$ -activated cells.

Our study identified canonical WNT as a new regulator of profibrotic IL-11 in TGF- $\beta$ -activated cardiac fibroblasts. Previous studies showed that IL-11 is a downstream effector of TGF- $\beta$ -mediated fibrosis in the heart, lungs, aorta and kidney [13,22,24]. In line with previous findings, we observed that blockade of IL-11 suppressed profibrotic signature in activated myofibroblasts. As in case of profibrotic response, WNT3a in the absence of TGF- $\beta$  was insufficient to regulate IL-11. We consider that IL-11 is directly regulated by  $\beta$ -catenin signaling at transcriptional level, but its expression requires co-activation of other TGF- $\beta$ -dependent pathways. Of note, IL-11 regulates Erk1/2 and thereby accelerates one of the non-classical arms of TGF- $\beta$  signaling. Collectively, these data underline a superior role of TGF- $\beta$  in cardiac fibrosis and the complexity of the downstream mechanisms. The  $\beta$ -catenin-TAK1 axis triggering IL-11 production seem to represent a major molecular mechanism of TGF- $\beta$ -dependent fibrosis. Regulation of IL-11 by  $\beta$ -catenin pathway has not been reported in fibrosis, however cooperation of TGF- $\beta$  and WNT in regulating IL-11 was observed in intestinal tumor model [32].

In opposite to profibrotic activity of the canonical WNT3a, a non-canonical WNT5a suppressed fibroblast-to-myofibroblast transition induced by TGF- $\beta$ . It should be noticed that WNT5a does not activate  $\beta$ -catenin pathway and therefore is unable to promote myofibroblast phenotype. In line with our findings, anti-fibrotic but also anti-angiogenic effects of WNT5a have been also reported in a model of TGF- $\beta$ -dependent peritoneal injury [33]. Furthermore, published data demonstrated WNT5a controlling IL-6 production in human cardiac fibroblasts [25,34]. It seems that WNT5a is switching the phenotype of cardiac fibroblasts from profibrotic towards proinflammatory. Interestingly, despite IL-11 and WNT5a show opposite effects on activation of cardiac fibroblast, Erk1/2 pathway was identified as a key downstream signaling for both of them [13,35]. It suggests that Erk1/2 might act pro- or anti-fibrotic depending on the presence of specific co-activators or the status of activated cells, but any other alternative mechanism cannot be excluded.

A growing body of evidence indicates elevated levels of multiple WNT ligands in the affected heart [12,30]. Thus, a balance between canonical and non-canonical WNT signaling might be decisive for the fate of activated cardiac fibroblasts in the injured heart and eventually for development of cardiac fibrosis. Accumulating data strongly indicate that the canonical  $\beta$ -catenin pathway is critically involved in pathogenesis of cardiac fibrosis. Herein, by using a model of human cardiac fibroblasts we elucidated

an interplay between by TGF- $\beta$ , WNT/ $\beta$ -catenin and IL-11 and proposed a novel mechanism controlling cardiac fibrosis. This study adds molecular knowledge to our understanding of profibrotic mechanisms in the heart, which is essential for development of effective cardioprotective therapies targeting cardiac fibrosis.

## 2. Materials and Methods

### *Cell cultures and stimulations*

Primary fetal human cardiac were originally obtained from Cell Applications (Cell Applications, San Diego, USA). L- WNT3a (overexpressing WNT3a), L- WNT5a (overexpressing WNT5a) and L (control line for L- WNT3a and L- WNT5a) cell lines were obtained from ATCC (ATCC, Manassas, USA). The AP-1 reporter cardiac fibroblasts were described previously [25]. Cells were cultured in the Dulbecco's modification of Eagle medium (DMEM, Corning, New York, USA) supplemented with 10% FBS (EURx, Gdansk, Poland), 1:100 penicillin/streptomycin, 100 mM nonessential amino acids (all Corning, New York, USA) and 50 mM  $\beta$ -mercaptoethanol (Sigma-Aldrich, Taufkirchen, Germany). L- WNT3a and L- WNT5a cell lines were selected with medium containing G418 (Sigma-Aldrich). Human cardiac fibroblasts were cultured up to 12 passages. Cells were passaged using standard protocol with 0.25% trypsin (Corning) and seeded 24 h before experiments. All cells used in this study were mycoplasma-free. Cell culture supernatants from L-, L-WNT3a, and L- WNT5a cultured for 48 h were collected, filtered (pore size 0.2  $\mu$ m, Carl Roth, Karlsruhe, Germany) and used fresh for cardiac fibroblasts stimulation. Cells were stimulated with 10 ng/mL of TGF- $\beta$  (Peprotech, Rocky Hill, USA) or with 1  $\mu$ M CHIR99021 (CHIR, Tocris, Bristol, UK) in the presence or absence of 10 ng/mL TGF- $\beta$  for 72 h. To block WNT secretion cells were treated with 0.1  $\mu$ M Wnt-C59 (Tocris) with or without 10 ng/mL TGF-  $\beta$  for 24 h. Neutralization of IL-11 was performed with 0.5  $\mu$ g/mL anti-IL-11 antibody (clone: 226226, R&D systems, Minneapolis, USA) for 72 h.

### *Immunofluorescence*

After stimulation, cells were fixed with 4% paraformaldehyde for 5 min at room temperature, permeabilized with 0.01% Triton X-100 for 5 min, and blocked with 5% BSA in PBS for 20 min. Specimens were incubated with primary mouse anti- $\beta$ -catenin (1:500, clone CAT-5H10, Thermo Fisher Scientific, Waltham, USA) anti- $\alpha$ SMA antibody (1:200, clone 1A4, Biolegend, San Diego, USA) at room temperature, followed by the secondary AlexaFluor555 goat anti-mouse antibody (1:8000) and phalloidin (1:1000, all Thermo Fisher Scientific) at room temperature for 1 h. Hoechst (1:1000, Thermo Fisher Scientific) was used to label nuclei. Immunofluorescence was analyzed using Leica DM5500 B fluorescence microscope with DFC365 FX camera (Leica Microsystems, Wetzlar, Germany).

### *Quantitative RT-PCR*

Total RNA was extracted with Trizol Reagent (Thermo Fisher Scientific). 150ng or 250ng of total RNA was used to synthesize complementary DNA using NG dART RT Kit (EURx, Gdansk, Poland). Quantitative real-time PCR was performed using SYBR Green PCR Master Mix (EURx) and oligonucleotides complementary to transcripts of the analyzed genes using the Quant Studio 7 Real-Time PCR system (Applied Biosystems, Foster City, USA). The following oligonucleotides were used in this study: *AXIN2*: 5'-TAACCCCTCAGAGCGATGGA-3' and 5'-

CCTCCTCTCTTTTACAGCAGGG-3', *TCF7*: 5'-CCAAGAATCCACCACAGGAGG-3' and 5'-GCCTAGAGCACTGTCATCGG-3', *JUN*: 5'-GAGCTGGAGCGCCTGATAAT-3' and 5'-CCCTCCTGCTCATCTGTCAC-3', *ATF2*: 5'-GTGAATTCTGCCAGGCAATAC-3' and 5'-CCTCGTTGGTAAAACGCTGG-3', *ACTA2*: 5'-GACAATGGCTCTGGGCTCTGTAA-3' and 5'-ATGCCATGTTCTATCGGGTACTT-3', *IL11*: 5'-TCG AGT TTC CCC AGA CCC TC-3' and 5'-GAA TCC AGG TTG TGG TCC CC-3', *ACTG2*: 5'-GCAAGCGAGGGATCCTAACTC-3' and 5'-AGGAGTGGTGCCAGATCTTCTC-3' *COL1A1*: 5'-CAGCCGCTTCACCTACAGC-3' and 5'-TTTTGTATTCAATCACTGTCTTGC-3', *FNI*: 5'-TGCCACTGTTCTCCTACGTGG-3' and 5'-GGAGAATTCAAGTGTGACCCTCATG-3', *VCL*: 5'-AATGGTCCAGCAAGCCGGG-3' and 5'-TCGTCTTCCATGTCAGGGGG-3', *GAPDH*: 5'-GGGAAGCTTGTCATCAATGGA-3' and 5'-TCTCGCTCCTGGAAGATGGT-3'. Transcript levels of *GAPDH* were used as endogenous reference and relative gene expression was calculated using the  $2^{-\Delta\Delta Ct}$  method.

#### Luciferase Assay

Luciferase expression in the AP-1 reporter fibroblasts was assessed using the Bright-Glo Luciferase Assay System (Promega, Madison, USA) according to the manufacturer's protocol. Bioluminescence was measured by SynergyHT plate reader (BioTek, Winooski, USA) or Wallac Victor2 1420 Multilabel counter (Perkin-Elmer, Waltham, USA).

#### RNA-seq analysis

Total RNA from three biological independent experiments of cardiac fibroblasts were sent to Genewiz for preparation RNA-Seq libraries. Libraries were run with Illumina, strand-specific RNA-Seq with PolyA selection and Illumina NovaSeq 2x150 bp sequencing, 10M read pairs. Raw reads were mapped to the GRCh38 reference genome using STAR aligner v.2.5.2b. NSEMBL using the STAR aligner v.2.5.2b. Unique gene hit counts were calculated by using feature Counts from the Subread package v.1.5.2. After extraction of gene hit counts, the gene hit counts table was used for downstream differential expression analysis. Comparison of gene expression between the groups was performed using DESeq2. The Wald test was used to generate *p* values and log2 fold changes. Genes with an adjusted *p* value < 0.05 and absolute log2 fold change > 1 were called as differentially expressed genes for each comparison. Differentially expressed probes from respective groups are highlighted in the heatmap plot. Heatmap was generated using online tool: <https://software.broadinstitute.org/morpheus/>. The raw values were transformed, scaled with red representing upregulated genes and blue representing downregulated genes. The GO analyses were performed using online tool: [https://tools.dice-database.org/GO\\_net/](https://tools.dice-database.org/GO_net/) for human species.

#### Western Blotting

For protein expression analysis, a conventional SDS–PAGE followed by protein blotting strategy was performed as described previously [25]. The following antibodies were used: rabbit anti-GAPDH (1:5000, clone GA1R, Thermo Fisher Scientific), rabbit anti- $\alpha$ SMA (1:800, clone 1A4, Biolegend), mouse anti-fibronectin (1:700, polyclonal, Abcam), rabbit anti -TAK (polyclonal 1:1000), rabbit anti-phospho-TAK (polyclonal, 1:1000), anti-Smad2 (clone D43B4, 1:1000) anti-phospho-Smad2 (clone 138D4, 1:1000, all Cell Signaling Technology, Danvers, USA). Protein abundance was normalized to GAPDH levels. Results were analyzed with ImageJ software (NIH, Bethesda, USA) according to method for Western Blot densitometry band quantification through image analysis software with background subtraction [26].

### *ELISA*

Type I collagen was measured in cell culture supernatants using Human Procollagen I alpha 1 DuoSet ELISA (R&D Systems) and IL-11 was measured in cell culture supernatants using IL-11 Human ELISA kit (Thermo Fisher Scientific) according to the manufacturer's protocol.

### *Collagen Contraction Assay*

After stimulations cells were trypsinized, counted and mixed with Collagen Gel Working Solution prepared according to manufacturer's protocol (Cell Biolabs Inc, San Diego, USA). The solution was added to 24-well plated and allowed to polymerize for 1h at 37°C. Fresh growth media was added to the solidified collagen gels and plates were returned to the incubator. After 24h the surface of contraction was released and contraction was monitored over the next 24 hours. Next the surface area of contracted gels was imaged using ChemiDoc instrument (Bio-Rad) and measured using ImageJ software (NIH).

### *Statistical Analysis*

Experimental results are presented as mean $\pm$  SEM. Raw data normally distributed and data with homogenous variance (Brown–Forsythe test) were compared using a one-way ANOVA followed by Dunn's post-hoc test or two-way ANOVA followed by Tukey's post-hoc test for the indicated comparisons. All analyses were computed using GraphPad Prism 9.1.1 software. Differences were considered as statistically significant for  $p < 0.05$ .

### **Author contributions**

Designed research: P.B. Performed research: E.D., M.C., K.T. Analyzed data: E.D., G.K., P.B. Writing – original draft, E.D. and P.B.; Writing – review & editing: G.K. Supervision: M.S., G.K., P.B.

### **Acknowledgements**

This work has been supported by the National Science Centre (Poland), grant 2014/14/E/NZ5/00175.

### **Disclosures**

None declared.

**Figure legends****Figure 1. Activation of canonical and non-canonical WNT pathways in human cardiac fibroblasts.**

Human cardiac fibroblasts were stimulated with WNT3a, WNT5a or TGF- $\beta$ . Panel (A) illustrates cellular localizations of  $\beta$ -catenin in fibroblasts 30 min after indicated treatment. DAPI stains cell nuclei. Arrows indicate nuclear localization of  $\beta$ -catenin. Data are representant of three independent experiments. Bar=50 $\mu$ m. Panel (B) represents expression of canonical (*AXIN2* and *TCF7*) and non-canonical (*JUN* and *ATF2*) WNT responsive genes 1 h after indicated treatment. Panel (C) shows quantification of AP-1 activity in AP-1-hCF reporter cardiac fibroblasts following 24h treatment ( $n=6$ ). Graphs show means  $\pm$  SEM,  $p$  values computed using one-way ANOVA followed by Dunnett's multiple comparisons test, \*  $p < 0.05$  (post-hoc test vs. control).

**Figure 2. RNAseq analysis of TGF- $\beta$ -activated fibroblasts following WNT3a or WNT5a treatment.**

Cardiac fibroblasts were stimulated with WNT3a or WNT5a in the presence of TGF- $\beta$  for 72 h. Panel (A) shows a heat map of fibrosis-related genes differentially expressed (fold change $>2.0$  and  $p < 0.05$ ) in fibroblasts treated with WNT3a/TGF- $\beta$  versus TGF- $\beta$  alone ( $n=3$ ). All dysregulated genes of WNT3a/TGF- $\beta$  versus TGF- $\beta$  are presented in the Supp Fig. 1. Panel (B) represents a heat map of all differentially expressed genes (fold change $>2.0$  and  $p < 0.05$ ) in cardiac fibroblasts treated with WNT5a/TGF- $\beta$  versus TGF- $\beta$  alone ( $n=3$ ). GO categories: Cell differentiation (GO:0030154), Regulation of cellular component organization (GO:0051128), Extracellular matrix (GO:0031012), WNT signaling pathway (GO:0016055) and Muscle Contraction (GO:0006936).

**Figure 3. WNT/ $\beta$ -catenin enhances profibrotic response of TGF- $\beta$ -activated cardiac fibroblasts.**

Panel (A) illustrates representative immunocytochemistry staining of  $\alpha$ SMA, F-actin and cell nuclei (DAPI) of cardiac fibroblasts stimulated with WNT3a and/or TGF- $\beta$  for 72 h. Data are representant of three biological independent experiments. Bar=100  $\mu$ m. Panel (B) shows quantification of gel contraction (72 h,  $n=3$ ) and representative image of collagen gel containing cardiac fibroblasts pre-treated as indicated. Panels (C-E) describe profibrotic effect of GSK3 $\beta$  inhibitor CHIR99021 (CHIR) in the presence or absence of TGF- $\beta$  in cardiac fibroblasts. Expression of profibrotic genes after 72 h of stimulation is presented in panel (C) ( $n=4$ ). Panel (D) shows representative immunoblots and the respective quantifications of  $\alpha$ SMA and fibronectin after 72 h of stimulation (all  $n=4$ ) and panel (E) indicates pro-collagen I  $\alpha 1$  level in supernatants ( $n=6$ ) stimulated for 5 days. Bars and whiskers represent means  $\pm$  SEM. Main effect of WNT3a/CHIR, TGF- $\beta$  and their interaction were tested using two-way ANOVA followed by Tukey's multiple comparisons test, when the  $p$ -value for interaction is  $\leq 0.1$ ; \* $p < 0.05$  (post-hoc TGF- $\beta$  vs. TGF- $\beta$ +WNT3a or CHIR vs. TGF $\beta$ +CHIR).

**Figure 4. WNT5a reverses profibrotic response of TGF- $\beta$ -activated cardiac fibroblasts.**

Panel (A) shows expression of profibrotic genes ( $n=8$ ) and panel (B) illustrates representative immunoblots with

the respective quantifications of  $\alpha$ SMA ( $n=4$ ) and fibronectin ( $n=4$ ) in cardiac fibroblasts stimulated with WNT5a and/or TGF- $\beta$  for 72 h. Panel (C) shows pro-collagen I  $\alpha 1$  levels measured in supernatants of cardiac fibroblasts treated as indicated for 5 days ( $n=5$ ). Bars and whiskers represent means  $\pm$ SEM. Main effect of WNT5a, TGF- $\beta$  and their interaction were tested using two-way ANOVA followed by Tukey's multiple comparisons test, when the  $p$ -value for interaction is  $<0.05$ ,  $*p < 0.05$  (post-hoc TGF- $\beta$  vs. TGF- $\beta$ +WNT5a).

**Figure 5. WNT3a enhances TAK1 activation and IL-11 production in TGF- $\beta$ -activated fibroblasts.** Representative immunoblots and the respective quantification of p-SMAD2/SMAD2 (A) ( $n=4$ ) and p-TAK1/TAK1 (B) ( $n=4$ ) in cardiac fibroblasts stimulated with Wnt3a and/or TGF- $\beta$  for 1 h. Panel (C) indicates *IL11* gene expression (24 h  $n=6$ ) and panel (D) secreted IL-11 protein level in supernatants (72 h  $n=5$ ) of cardiac fibroblast stimulated as indicated. Panel (E) shows secreted IL11 proteins in supernatants ( $n=4$ ) of cardiac fibroblast stimulated with or without TGF- $\beta$  for 24h in the presence or absence WNT secretion blocker WNT-C59. Bars and whiskers represent means  $\pm$ SEM. Main effect of WNT3a or WNT-C59, TGF- $\beta$  and their interaction were tested using two-way ANOVA followed by Tukey's multiple comparisons test, when the  $p$ -value for interaction is  $\leq 0.1$ ;  $*p < 0.05$ ,  $*p < 0.05$  (post-hoc TGF- $\beta$  vs. TGF- $\beta$ +WNT3a or TGF- $\beta$  vs. TGF- $\beta$ +WNT-C59)

**Figure 6. IL-11 contributes to profibrotic response enhanced by WNT3a.** Human cardiac fibroblast were cultured with TGF- $\beta$  and stimulated with WNT3a in the presence or absence of anti-IL-11 neutralizing antibody for 72 h. Panel (A) presents mRNA levels of profibrotic genes after indicated treatment. *ACTA2*, *COL1A1*, *FNI* ( $n=10$ ) and *ACTG2*, *VCL* ( $n=6$ ). Panel (B) shows quantification of  $\alpha$ SMA and fibronectin ( $n=4$ ) and the representative immunoblots. Pro-collagen I  $\alpha 1$  levels measured in supernatants ( $n=6$ ) are presented in panel (C). Panel (D) shows quantification of gel contraction ( $n=3$ ) containing cardiac fibroblasts pre-treated as indicated. Bars and whiskers represent means  $\pm$ SEM. Main effect of WNT3a-TGF- $\beta$ , anti-IL11 antibody and their interaction were tested using two-way ANOVA followed by Tukey's multiple comparisons test, when the  $p$ -value for interaction is  $\leq 0.1$ ;  $*p < 0.05$  (post-hoc TGF- $\beta$ +WNT3a vs. TGF- $\beta$ +WNT3a+anti- IL11).



## References

1. Humeres, C.; Frangogiannis, N.G. Fibroblasts in the Infarcted, Remodeling, and Failing Heart. *JACC Basic to Transl. Sci.* **2019**, *4*, 449–467, doi:10.1016/j.jacbts.2019.02.006.
2. González, A.; Schelbert, E.B.; Díez, J.; Butler, J. Myocardial Interstitial Fibrosis in Heart Failure: Biological and Translational Perspectives. *J. Am. Coll. Cardiol.* **2018**, *71*, 1696–1706, doi:10.1016/j.jacc.2018.02.021.
3. Kato, S.; Saito, N.; Kirigaya, H.; Gytoku, D.; Iinuma, N.; Kusakawa, Y.; Iguchi, K.; Nakachi, T.; Fukui, K.; Futaki, M.; et al. Prognostic significance of quantitative assessment of focal myocardial fibrosis in patients with heart failure with preserved ejection fraction. *Int. J. Cardiol.* **2015**, *191*, 314–319, doi:10.1016/j.ijcard.2015.05.048.
4. van Putten, S.; Shafieyan, Y.; Hinz, B. Mechanical control of cardiac myofibroblasts. *J. Mol. Cell. Cardiol.* **2016**, *93*, 133–142, doi:10.1016/j.yjmcc.2015.11.025.
5. Nakaya, M.; Watari, K.; Tajima, M.; Nakaya, T.; Matsuda, S.; Ohara, H.; Nishihara, H.; Yamaguchi, H.; Hashimoto, A.; Nishida, M.; et al. Cardiac myofibroblast engulfment of dead cells facilitates recovery after myocardial infarction. *J. Clin. Invest.* **2017**, *127*, 383–401, doi:10.1172/JCI83822.
6. Gibb, A.A.; Lazaropoulos, M.P.; Elrod, J.W. Myofibroblasts and fibrosis: Mitochondrial and metabolic control of cellular differentiation. *Circ. Res.* **2020**, *127*, 427–447, doi:10.1161/CIRCRESAHA.120.316958.
7. Kuwahara, F.; Kai, H.; Tokuda, K.; Kai, M.; Takeshita, A.; Egashira, K.; Imaizumi, T. Transforming growth factor- $\beta$  function blocking prevents myocardial fibrosis and diastolic dysfunction in pressure-overloaded rats. *Circulation* **2002**, *106*, 130–135, doi:10.1161/01.CIR.0000020689.12472.E0.
8. Massagué, J. TGF $\beta$  signalling in context. *Nat Rev Mol Cell Biol* **2014**, *13*, 616–630, doi:10.1038/nrm3434.TGF.
9. Zhang, Y.E. Non-Smad signaling pathways of the TGF- $\beta$  family. *Cold Spring Harb. Perspect. Biol.* **2017**, *9*, 1–18, doi:10.1101/cshperspect.a022129.
10. Sweeney, M.; Corden, B.; Cook, S.A. Targeting cardiac fibrosis in heart failure with preserved ejection fraction: mirage or miracle? *EMBO Mol. Med.* **2020**, *12*, 1–26, doi:10.15252/emmm.201910865.
11. Fang, L.; Murphy, A.J.; Dart, A.M. A clinical perspective of anti-fibrotic therapies for cardiovascular disease. *Front. Pharmacol.* **2017**, *8*, 1–8, doi:10.3389/fphar.2017.00186.
12. Działo, E.; Tkacz, K.; Błyszczuk, P. Crosstalk between the TGF- $\beta$  and WNT signalling pathways during cardiac fibrogenesis. *Acta Biochim. Pol.* **2018**, *65*, 341–349, doi:10.18388/abp.2018\_2635.
13. Schafer, S.; Viswanathan, S.; Widjaja, A.A.; Lim, W.W.; Moreno-Moral, A.; DeLaughter, D.M.; Ng, B.; Patone, G.; Chow, K.; Khin, E.; et al. IL-11 is a crucial determinant of

- cardiovascular fibrosis. *Nature* **2017**, 552, 110–115, doi:10.1038/nature24676.
14. Moon, R.T.; Kohn, A.D.; De Ferrari, G. V.; Kaykas, A. WNT and  $\beta$ -catenin signalling: Diseases and therapies. *Nat. Rev. Genet.* **2004**, 5, 691–701, doi:10.1038/nrg1427.
  15. Garriock, R.J.; Warkman, A.S.; Meadows, S.M.; D'Agostino, S.; Krieg, P.A. Census of vertebrate Wnt genes: Isolation and developmental expression of *Xenopus* Wnt2, Wnt3, Wnt9a, Wnt9b, Wnt10a, and Wnt16. *Dev. Dyn.* **2007**, 236, 1249–1258, doi:10.1002/dvdy.21156.
  16. Valenta, T.; Hausmann, G.; Basler, K. The many faces and functions of  $\beta$ -catenin. *EMBO J.* **2012**, 31, 2714–2736, doi:10.1038/emboj.2012.150.
  17. Bengoa-Vergniory, N.; Kypta, R.M. Canonical and noncanonical Wnt signaling in neural stem/progenitor cells. *Cell. Mol. Life Sci.* **2015**, 72, 4157–4172, doi:10.1007/s00018-015-2028-6.
  18. Seidenberg, J.; Stellato, M.; Hukara, A.; Ludewig, B.; Klingel, K.; Distler, O.; Błyszczuk, P.; Kania, G. The ap-1 transcription factor fosl-2 regulates autophagy in cardiac fibroblasts during myocardial fibrogenesis. *Int. J. Mol. Sci.* **2021**, 22, 1–19, doi:10.3390/ijms22041861.
  19. Błyszczuk, P.; Müller-Edenborn, B.; Valenta, T.; Osto, E.; Stellato, M.; Behnke, S.; Glatz, K.; Basler, K.; Lüscher, T.F.; Distler, O.; et al. Transforming growth factor- $\beta$ -dependent Wnt secretion controls myofibroblast formation and myocardial fibrosis progression in experimental autoimmune myocarditis. *Eur. Heart J.* **2017**, 38, 1413–1425, doi:10.1093/eurheartj/ehw116.
  20. Xiang, F.L.; Fang, M.; Yutzey, K.E. Loss of  $\beta$ -catenin in resident cardiac fibroblasts attenuates fibrosis induced by pressure overload in mice. *Nat. Commun.* **2017**, 8, doi:10.1038/s41467-017-00840-w.
  21. Lal, H.; Ahmad, F.; Zhou, J.; Yu, J.E.; Vagnozzi, R.J.; Guo, Y.; Yu, D.; Tsai, E.J.; Woodgett, J.; Gao, E.; et al. Cardiac fibroblast glycogen synthase kinase-3 $\beta$  regulates ventricular remodeling and dysfunction in ischemic heart. *Circulation* **2014**, 130, 419–430, doi:10.1161/CIRCULATIONAHA.113.008364.
  22. Ng, B.; Dong, J.; Viswanathan, S.; D'Agostino, G.; Widjaja, A.A.; Lim, W.W.; Ko, N.S.J.; Tan, J.; Chothani, S.P.; Huang, B.; et al. IL-11 is a therapeutic target in idiopathic pulmonary fibrosis. *bioRxiv* **2018**, 1–15, doi:10.1101/336537.
  23. Einarsson, O.; Geba, G.P.; Zhu, Z.; Landry, M.; Elias, J.A. Interleukin-11: Stimulation in vivo and in vitro by respiratory viruses and induction of airways hyperresponsiveness. *J. Clin. Invest.* **1996**, 97, 915–924, doi:10.1172/JCI118514.
  24. Corden, B.; Adami, E.; Sweeney, M.; Schafer, S.; Cook, S.A. IL-11 in cardiac and renal fibrosis: Late to the party but a central player. *Br. J. Pharmacol.* **2020**, 177, 1695–1708, doi:10.1111/bph.15013.
  25. Działo, E.; Rudnik, M.; Koning, R.I.; Czepiel, M.; Tkacz, K.; Baj-Krzyworzeka, M.; Distler, O.; Siedlar, M.; Kania, G.; Błyszczuk, P. WNT3a and WNT5a transported by exosomes

- activate WNT signaling pathways in human cardiac fibroblasts. *Int. J. Mol. Sci.* **2019**, *20*, 1–15, doi:10.3390/ijms20061436.
26. Gallo-Oller, G.; Ordoñez, R.; Dotor, J. A new background subtraction method for Western blot densitometry band quantification through image analysis software. *J. Immunol. Methods* **2018**, *457*, 1–5, doi:10.1016/j.jim.2018.03.004.
  27. Gross, J.C.; Chaudhary, V.; Bartscherer, K.; Boutros, M. Active Wnt proteins are secreted on exosomes. *Nat. Cell Biol.* **2012**, *14*, 1036–1045, doi:10.1038/ncb2574.
  28. Carthy, J.M.; Garmaroudi, F.S.; Luo, Z.; McManus, B.M. Wnt3a induces myofibroblast differentiation by upregulating TGF- $\beta$  signaling through SMAD2 in a  $\beta$ -catenin-dependent manner. *PLoS One* **2011**, *6*, doi:10.1371/journal.pone.0019809.
  29. Blyszczuk, P.; Müller-Edenborn, B.; Valenta, T.; Osto, E.; Stellato, M.; Behnke, S.; Glatz, K.; Basler, K.; Lüscher, T.F.; Distler, O.; et al. Transforming growth factor- $\beta$ -dependent Wnt secretion controls myofibroblast formation and myocardial fibrosis progression in experimental autoimmune myocarditis. *Eur. Heart J.* **2016**, *6*, 8–9, doi:10.1093/eurheartj/ehw116.
  30. Zhao, Y.; Wang, C.; Wang, C.; Hong, X.; Miao, J.; Liao, Y.; Zhou, L.; Liu, Y. An essential role for Wnt/ $\beta$ -catenin signaling in mediating hypertensive heart disease. *Sci. Rep.* **2018**, *8*, 1–14, doi:10.1038/s41598-018-27064-2.
  31. Czepiel, M.; Diviani, D.; Jaźwa-Kusior, A.; Tkacz, K.; Rolski, F.; Smolenski, R.T.; Siedlar, M.; Eriksson, U.; Kania, G.; Blyszczuk, P. Angiotensin II receptor 1 controls profibrotic Wnt/ $\beta$ -catenin signalling in experimental autoimmune myocarditis. *Cardiovasc. Res.* **2021**, 1–12, doi:10.1093/cvr/cvab039.
  32. Labbé, E.; Lock, L.; Letamendia, A.; Gorska, A.E.; Gryfe, R.; Gallinger, S.; Moses, H.L.; Attisano, L. Transcriptional cooperation between the transforming growth factor- $\beta$  and Wnt pathways in mammary and intestinal tumorigenesis. *Cancer Res.* **2007**, *67*, 75–84, doi:10.1158/0008-5472.CAN-06-2559.
  33. Padwal, M.; Liu, L.; Margetts, P.J. The role of WNT5A and Ror2 in peritoneal membrane injury. *J. Cell. Mol. Med.* **2020**, *24*, 3481–3491, doi:10.1111/jcmm.15034.
  34. Abraitte, A.; Vinge, L.E.; Askevold, E.T.; Lekva, T.; Michelsen, A.E.; Ranheim, T.; Alfsnes, K.; Fiane, A.; Aakhus, S.; Lunde, I.G.; et al. Wnt5a is elevated in heart failure and affects cardiac fibroblast function. *J. Mol. Med.* **2017**, *95*, 767–777, doi:10.1007/s00109-017-1529-1.
  35. Abraitte, A.; Lunde, I.G.; Askevold, E.T.; Michelsen, A.E.; Christensen, G.; Aukrust, P.; Yndestad, A.; Fiane, A.; Andreassen, A.; Aakhus, S.; et al. Wnt5a is associated with right ventricular dysfunction and adverse outcome in dilated cardiomyopathy. *Sci. Rep.* **2017**, *7*, 1–10, doi:10.1038/s41598-017-03625-9.