

Review

# Overview Of Approaches to Improve Rhizoremediation of Petroleum Hydrocarbon-Contaminated Soils

Fahad Alotaibi <sup>1,2</sup>, Mohamed Hijri <sup>1,3</sup> and Marc St-Arnaud <sup>1,\*</sup>

<sup>1</sup> Biodiversity Centre, Institut de Recherche en Biologie Végétale, Université de Montréal and Jardin botanique de Montréal, Montréal, QC, Canada

<sup>2</sup> Department of Soil Science, King Saud University, Riyadh 11564, Saudi Arabia

<sup>3</sup> African Genome Center, Mohammed VI Polytechnic University (UM6P), 43150 Ben Guerir, Morocco

\* Correspondence: marc.st-arnaud@umontreal.ca

**Abstract:** Soil contamination with petroleum hydrocarbons (PHCs) has become a global concern and has resulted from the intensification of industrial activities. This has created a serious environmental issue; therefore, there is a need to find solutions, including application of efficient remediation technologies or improvement of current techniques. Rhizoremediation is a green technology that has received global attention as a cost-effective and possibly efficient remediation technique for PHC-polluted soil. Rhizoremediation refers to the use of plants and their associated microbiota to clean up contaminated soils, where plant roots stimulate soil microbes to mineralize organic contaminants to H<sub>2</sub>O and CO<sub>2</sub>. However, this multipartite interaction is complicated because many biotic and abiotic factors can influence microbial processes in the soil, making the efficiency of rhizoremediation unpredictable. This review reports the current knowledge of rhizoremediation approaches that can accelerate the remediation of PHC-contaminated soil. Recent approaches discussed in this review include 1) selecting plants with desired characteristics suitable for rhizoremediation; 2) exploiting and manipulating the plant microbiome by using inoculants containing plant growth-promoting rhizobacteria (PGPR) or hydrocarbon-degrading microbes, or a combination of both types of organisms; 3) enhancing the understanding of how the host–plant assembles a beneficial microbiome, and how it functions, under pollutant stress. A better understanding of plant–microbiome interactions could lead to successful use of rhizoremediation for PHC-contaminated soil in the future.


**Keywords:** Phytoremediation; PGPR; hydrocarbon-degrading bacteria; *Salix*; contaminated soils; Alkanes; PAHs.

## 1. Introduction

Industrial activities, including mining and extraction of oil and gas, as well as chemical inputs into agricultural production systems, have led to different degrees of environmental contamination worldwide. Petroleum hydrocarbons (PHCs) are among the major pollutants that can pose a serious environmental threat. PHC products have adversely affected various ecosystems, causing disturbing damage to natural habitats with serious economic consequences [1].

PHCs are heterogeneous organic mixtures composed of carbon and hydrogen atoms arranged in varying structural configurations and have different physical and chemical properties [2]. These compounds consist mainly of hydrocarbons and fewer numbers of other non-hydrocarbon constituents such as nitrogen, oxygen, and sulfur [3,4]. They are broadly classified into two major fractions: aliphatic hydrocarbons and aromatic hydrocarbons (Figure 1). Prior to processing, PHCs are composed, on average, of ~57% aliphatic hydrocarbons, ~29% aromatic hydrocarbons, and ~14% asphaltenes and other polar compounds containing nitrogen, oxygen, and sulfur [5]. Aliphatic hydrocarbons include both

linear or branched-chain hydrocarbons, which may be unsaturated (alkenes and alkynes) or saturated (alkanes) [6]. Aromatic hydrocarbons include monocyclic (i.e., benzene, toluene, phenol, etc.) and polycyclic aromatic hydrocarbons (PAHs) (Figure 1). PHCs are the most common pollutants in soil and ground water worldwide. The ever-increasing dependency of modern society on fuel for energy generation in many vital sectors such as electricity, heat, industry, and transportation has resulted in the extensive exploitation of PHCs [2]. Although environmental transition actions have been taken in many countries, dependency on petroleum will last for some decades, contributing to organic pollution risks.



**Figure 1** Schematic diagram showing the classification of PHCs.

Soil contamination with PHCs is an international issue, and the magnitude of soil pollution is hard to quantify. For example, in Australia, around 80,000 sites are estimated to be contaminated by PHCs [7], whereas in Canada around 22,000 federal-owned sites are identified as being contaminated by PHCs [8]. In Europe, PHC contamination was observed in at least 342,000 sites [9]. These organic contaminants also pose serious health risks to humans and other organisms in addition to their adverse impact on the soil microflora, leading to environmental quality degradation. For instance, some aromatic substances such as BTEX and PAHs are notorious mutagens and carcinogens that can enter our food chain together with lipophilic compounds [10], and they have been linked with probable causes of bladder, kidney, liver, lung, and skin cancers. This explains the growing concern with these contaminants and the urgent need to use all possible means to protect the environment and to find the appropriate techniques to remediate polluted soils.

Various chemical, physical, and thermal conventional techniques have been used to remediate soils contaminated with PHCs. These conventional methods, which can contain, destroy, or separate the pollutants, include a wide range of both in situ and ex situ cleanup technologies such as asphalt batching, biopiles, chemical oxidation, excavation, hydrolysis, incineration, photolysis, pump and treat, multi-phased slurry reactors, soil vapor extraction, soil washing, and thermal desorption. However, these methods have

particular limitations. First, their cost is often prohibitive; for example, it can cost between US \$480 and 813 per m<sup>3</sup> for extraction [11]. Second, chemical procedures only work for specific organic compounds, and they most often destroy soil microbial communities. Third, these methods do not often result in a complete degradation of the pollutants [2,12]. Finally, PHC-contaminated soil contains numerous classes and types of toxic organic compounds, which make the choice of the proper method a challenging task. Hence, phytoremediation is a more recent and promising green-biotechnology that is perceived as an environmentally friendly, more cost-effective, and less destructive approach to cleanup contaminants in the environment.

## 2. Phytoremediation

Phytoremediation is a remediation technique that relies on the ability of plants and their associated microbiomes to accumulate, degrade, sequester, or stabilize harmful environmental contaminants [13,14]. Over the past two decades, the deployment of plants (and their associated microbiomes) to remediate a wide spectrum of inorganic and organic pollutants in soil and water environments has been carried out. This technique has been applied to remediate various types of pollutants such as chlorinated solvents [15], explosives [16], heavy metals [17], landfill leachates [18], pesticides [19], PHCs [20], radionuclides [21], and salts [22]. Although phytoremediation is still very much in its infancy, its application has been adopted by a growing number of companies. For example, the phytoremediation market has grown continuously at a rapid rate, with an estimated value of US \$32.2 billion in 2016 and is expected to reach US \$65.7 billion by 2025 [23].

Despite the broad public acceptance and it being an environmentally friendly approach, phytoremediation still remains a marginal option for *in situ* soil remediation [24]. As any other technique, phytoremediation has its own advantages and disadvantages (Table 1).

**Table 1.** Advantages and disadvantages of phytoremediation.

Advantages	Disadvantages
Has gained broad public acceptance	Slow (5 to 100 years)
Requires less maintenance effort, and as a result, little financial input is needed	Effectiveness is limited by plant root depth, as plant must reach the contaminants
Cost-effective since it is a naturally driven process that is powered by solar energy	Efficiency of phytoremediation depends on various environmental conditions, including soil physicochemical properties
Environmentally friendly approach	Plant selected for phytoremediation must tolerate toxicity
Minimum site disturbance	Unpredictable performance
Faster than natural attenuation	

Adapted and modified from [13,26,27,32].

Phytoremediation efficiency is dependent on many factors, including plant selection [25]; environmental parameters such as nutrient status, contaminant concentration, and bioavailability; soil pH, etc. [26], in addition to the composition and activity of plant-associated microbiomes. Plants and their associated microbiomes facilitate pollutant uptake from the environment via different processes, including degradation, extraction, stabilization, transformation, and volatilization [13,14]. The type of plant and pollutants plus the environmental conditions are key factors for determining the way in which phytoremediation techniques can be applied. Generally, phytoremediation technologies are divided into five different categories (Table 2). The phytoremediation method suitable for petroleum hydrocarbon-contaminated soil is called rhizoremediation [27], which is defined as the breakdown of organic pollutants by using plants and their root-associated microbiomes.

**Table 2.** Phytoremediation mechanisms whereby plants remediate polluted soils.


Category	Mechanism	Target Pollutants	Region of activity	References
Phytoextraction	Uptake and concentrate contaminants within plant tissue	Metals, radionuclides	Plant tissue	[14, 33]
Phytostabilization	Immobilization of contaminants in the root zone	Primarily metals	Root	[34]
Phytotransformation	Various enzymes work to convert pollutants into more stable and less toxic forms	Chlorinated solvents, ammunition waste, herbicides, mono-aromatic hydrocarbons	Plant tissue	[34]
Phytovolatilization	Uptake and evapotranspiration	Volatile organics (TCE, toluene, MTBE),	Plant tissue	[14]
Rhizoremediation	Breakdown of organic pollutants by using plants and root-associated microbiomes	PHCs, pesticides	Root	[27]

Adapted and modified from [14, 27, 33, 34].

Rhizoremediation of PHCs is facilitated through a process known as the 'rhizosphere effect' [28], in which plants exude a variety of organic compounds into their root-surrounding zone (the rhizosphere), resulting in an increase in abundance and activity of certain rhizospheric microbes, which in turn can degrade or metabolize hydrocarbon contaminants [28,29]. Understanding the plant–microbiome partnerships, and the underlying processes that govern and control PHC degradation, is a priority challenge in rhizoremediation research nowadays [2,30,31].

### 3. The rhizosphere Microbiome

By definition, the rhizosphere refers to the narrow zone of nutrient-rich soil in close proximity to plant roots and influenced greatly by plant exudates [35]. The rhizosphere is a hot spot for a myriad of organisms, including algae, archaea, arthropods, bacteria, fungi, nematodes, protozoa, and viruses [36], and it has been estimated that one gram of fresh roots contains up to  $10^{11}$  microbial cells representing more than 30000 prokaryotic species [37]. The structure of the rhizosphere microbiome depends on many factors such as soil type, environmental factors, the period of the year, plant development stage, and plant species and genotypes [37,38]. The rhizosphere microbiome is part of the larger root microbiome that also includes the rhizoplane microbiome and root interior microbiome (endophytes) [39,40] (Figure 2). The rhizosphere microbiome is one of the most complex habitats on our planet, and microbial functions occurring within the rhizosphere have critical influences on plant growth and productivity, soil fertility, carbon sequestration, and degradation of environmental contaminants [37].



**Figure 2.** Model of the root microbiome.

Plant roots exude a myriad of organic substances into the surrounding soil, comprising both low molecular weight organic compounds (amino acids, organic acids, sugars, phenolics, secondary metabolites, etc.) and high molecular weight organic compounds (polysaccharides, proteins, etc.). It has been estimated that 6-21% of photosynthetically fixed carbon in plants is released through root systems [7]. Therefore, root exudates are the major driver in shaping the rhizosphere microbiome. This countless and steady release of fixed carbon compounds into the rhizosphere, a process referred to as the rhizosphere effect, increases the activity and abundance of the rhizosphere microbial community compared to nearby bulk soil [40-42]. The magnitude of bacterial density in the rhizosphere is 10 to 1000 times higher than that in adjacent bulk soil; however, microbial community diversity in the rhizosphere is generally lower than that of bulk soils since rhizodeposition selectively enhances specific microbial taxa [37,43]. Although recruitment of the rhizosphere microbiome by plants is strongly dependent on the structure and composition of the bulk soil microbiota [44], different plant genotypes were found to select for different rhizosphere microbiomes [45], inferring that differential recruitment of beneficial microbiomes is also dependent on the genetic variation across plant species [44,46].

In addition to shaping the microbial communities in the rhizosphere, root exudates have other functions that benefit the plant itself. Through root exudation, plants can change the soil physico-chemical properties, contributing to nutrient assimilation, reducing the growth of competitor plant species, increasing the abundance of certain beneficial microbes, and regulating the microbiome composition in the rhizosphere [7,30,47,48].

The important role of root exudation, secreted by plants growing in PHC-contaminated soils, as facilitators of hydrocarbon rhizoremediation has been recognized recently [30,49]. This microbial process can function through different mechanisms. First, root exudates include degradable low molecular weight organic compounds such as carbohydrates, amino acids, and organic acids, all of which are readily available energy and nutrient sources for microbial utilization, stimulating the proliferation of microbial biomass and activities [27]. For instance, the addition of sugar and amino acids into soils causes an instant response (within 1 h) in microbial respiration [50]. Additionally, compounds essential in plant nutrient acquisition secreted by roots, such as enzymes (e.g., acid phosphatases) and chelating agents (phytosiderophores), provide microbial communities in the rhizosphere with a source of nutrients [49]. Second, plant root exudation can enhance PHC degradation by emitting a wide range of enzymes such as cytochrome P450

monooxygenases, dehalogenases, dioxygenase, laccases, and peroxidases [7,29,51]. Plant-secreted enzymes play a key role in the oxidation of PHCs [52], and the initial attack on the pollutant itself is primarily performed by soil microbial enzymes [52]. Third, secondary metabolites released by the plant roots, such as flavonoids and phenols, are analogous to many organic pollutants, thus increasing the abundance and activity of microbial communities equipped with genes relevant to degradation of organic pollutants within the rhizosphere, even in unpolluted soils [53]. Fourth, root-released exudates have been shown to increase the availability of organic pollutants for microbial metabolisms [29]. For example, Gao et al. [54] reported that the availability of phenanthrene and pyrene increased in the soil after the addition of citric acid and oxalic acid.

Considering the above-mentioned role of root exudates, the rhizosphere is hypothesized to be a suitable niche for rhizoremediation of PHC-contaminated soil [30]. Additionally, the rhizosphere is one of the environmental niches that is conducive to horizontal gene transfer (HGT) [55]. HGT is a mechanism used by bacterial communities to adapt to the presence of organic contaminants in their environments [57]. Bacteria may acquire genetic information from either closely related or phylogenetically distinct taxa in the community by HGT via different routes such as plasmids and transposons [57]. Several studies have reported that plasmids were shown to help bacterial communities adapt to environmental pollution stress [56,57].

To overcome the limitations and improve the efficiency of rhizoremediation, current research trends focus on several auxiliary strategies such as (1) selecting plants with desired characteristics suitable for rhizoremediation (such as increased contaminant tolerance or production of vigorous root system and shoot biomass [25], but also abilities to form symbiotic interactions with microorganisms), (2) exploiting and manipulating the plant microbiome by using inoculants containing plant growth-promoting rhizobacteria (PGPR) or hydrocarbon-degrading microbes or their combination [30], and (3) enhancing the understanding of the mechanisms through which host plants assemble a beneficial microbiome, and how it functions, under pollutant stress [31].

### 3.1. Plant Selection

Since the beginning of phytoremediation research, many plant species have been tested for their potential to enhance rhizoremediation of PHCs [58]. Plants enhance the degradation of PHCs principally by the unique properties of the plant itself and by providing optimal conditions for microbial proliferation in the rhizosphere [59]. In general, selection of plants suitable for rhizoremediation of PHCs should be based on the following criteria: tolerance to a broad range of PHCs, speed of growth, root morphology, ability to grow in many soil types, and the root exudate profile [59,60,61]. Additionally, plants should not be selected based solely on the contaminant uptake efficiency; their ability to stimulate microbial activity and abundance also should be considered [62,63].

Plants that have been used thus far in rhizoremediation span a wide range of families. Grasses (annual ryegrass, tall fescue) and other herbaceous crops (Indian mustard, sunflower), legumes (alfalfa, clover), and woody trees (hybrid poplars, willows), among others, have shown a high potential in the rhizoremediation of soil contaminated with PHCs [62,64].

Grasses have been studied extensively regarding their potential to facilitate the rhizoremediation of PHCs-impacted soil [61]. Grasses are often chosen for rhizoremediation applications because of their fast growth, high tolerance to PHCs, extensive fibrous root systems, large root surface area, and deeper root penetration into the soil matrix to depths of up to 3 m [60,64]. These unique characteristics of grass root systems allow microbial colonization and establishment of abundant microbial populations. For example, bacterial populations found in the rhizosphere of goosegrass (*Eleusine indica*) cultivated in PHC contaminated soil were 72 times more abundant than bacterial populations observed in the nearby uncultivated soil [65].

Legumes have also been tested for their potential to enhance rhizoremediation of PHC-contaminated soil [62]. The remarkable ability of legumes to form symbiotic relationships with the N-fixing rhizobia is of great importance in PHC-contaminated soil, which is characterized by low nutrient availability and high C/N ratio [64]. In addition, some legumes species such as alfalfa have a deep-rooting system that can penetrate highly compacted soil layers and create soil macropore spaces, thus increasing oxygenation of the soil matrix and consequently promoting microbial degradation [62].

Trees, such as willows (*Salix*), have also gained attention regarding their potential to improve rhizoremediation of PHC-polluted soil [2]. Willows are attractive for rhizoremediation of PHCs because they are easy to propagate, exhibit extremely fast growth in low-fertility soils, have high tolerance to several stressful environments, produce large biomass, and generate widespread deep-rooting systems [66]. Additionally, the large diversity of willows (~350-500 genetically distinct species), with a wide range of tolerance to various environmental conditions, facilitates selection of the most appropriate species suitable for a particular environment [66]. Compelling evidence has been reported about the use of willows for the rhizoremediation of soils polluted with organic contaminants, including PHCs [67-69].

More recently, promising approaches including the screening and identification of native plants grown on PHC-contaminated soil have been used. For example, Pérez-Jaramillo et al. [70] proposed a “back to the roots” framework that involves surveying indigenous plants and associated microbiomes, and their native habitats, to identify plants and microbial traits with the goal to restore associations that may have been diluted during plant domestication [70]. In fact, using native plant species in rhizoremediation offers many advantages over non-native species, including minimizing the potential of introducing alien species that can become invasive and disturb local flora and fauna [71], in addition to the fact that indigenous plants are more genetically diverse and more adapted to a wide range of climatic conditions compared to other plants currently chosen for rhizoremediation [72]. Following this approach, Desjardins et al. [73] described plants growing spontaneously in highly petroleum-contaminated decantation basins of a former petrochemical plant in Varennes (southern Québec, Canada) and identified three plants species (*Alisma triviale*, *Eleocharis obtusa*, and *Panicum capillare*) that were tolerant to PAHs and PHCs. Additionally, Lee et al. [74] studied the diversity of arbuscular mycorrhizal fungal communities of native plant species grown in highly PHC-contaminated soil and identified *Rhizophagus* as a key PHC-tolerant genus. Arbuscular mycorrhizal fungi (AMF) are obligate symbionts with approximately 80% of plant species on earth [75]. In agricultural settings, AMF are known for their plant growth-promoting effects including improved plant uptake of mineral nutrients, in particular phosphorus [76]. Additionally, AMF were shown to enhance plant tolerance toward several biotic and abiotic stresses such as nutrient deficiencies, plant pathogens, drought, salinity, and contaminants [76-80]. Therefore, AMF have more recently gained attention regarding their use in phytoremediation of soils contaminated with different pollutants such as heavy metals and PHCs [74,81,82].

Plants can sequester some PHCs into their root tissues and then transport them into shoots and leaves where they can be volatilized into the atmosphere or stored in plant cells such as in the vacuole or cell wall [58,83,84]. This process of sequestration and transportation of foreign chemicals inside plant tissues is analogous to that in the mammalian liver and has been dubbed “green liver” [85]. Plants also can degrade or transform organic pollutants into less toxic forms via their enzymatic machinery [13,86]. Additionally, due to their metabolic versatility, plants can adapt and confront many unfavorable stressful conditions, such as PHC contamination, where they can modify their own physiology and metabolism by synthesizing a variety of defensive proteins, regulatory enzymes, and metabolites [87]. However, plant growth under stress, such as PHC contamination, is expected to be lower than it would be under optimal conditions [87]. Therefore, plant growth may be positively enhanced by the presence of plant growth-promoting rhizobacteria (PGPR) that are able to alleviate stresses in plants via many mechanisms such as reducing soil nutrient deficiencies (fixing nitrogen, solubilizing phosphorus, and

enhancing iron uptake), synthesizing plant hormones, suppressing ethylene production via 1-aminocyclopropane-1-carboxylate (ACC) deaminase activity [87,88], and degrading a broad range of PHCs [2].

### 3.2. Exploiting and manipulating the plant microbiome through inoculation

#### 3.2.1. Plant Growth-Promoting Rhizobacteria (PGPR)

Bacteria are the predominant group within the soil microbiome community. It has been estimated that one gram of soil contains around  $10^8$  - $10^9$  bacterial cells [89] representing tens of thousands of different species [37]. The capacity of bacteria to utilize a wide range of many compounds as nutrient and energy sources, and their diverse metabolism, make them ideal associates in plant–microbiome partnerships [37]. Among bacterial communities dwelling in the rhizosphere are PGPR. PGPR are free-living and beneficial soil-borne bacteria associated with the root microbiome, enhancing plant growth and development by direct and indirect means [90-92]. The direct means by which PGPR may promote plant growth occur through fixing atmospheric N, increasing nutrient acquisition such as phosphorous, stimulating plant growth by producing different phytohormones, sequestration of iron by synthesis of siderophores, and alleviating stresses in plants by producing 1-aminocyclopropane-1-carboxylate (ACC) deaminase [90-92]. Indirect means of growth stimulation occur through biocontrol activities of PGPR against many plant phytopathogens via different mechanisms, including production of antimicrobial metabolites such as siderophores, antibiotics, and bacteriocins as well as induced systemic resistance (ISR) in plants [93,94].

A plethora of bacterial genera such as *Acetobacter*, *Acinetobacter*, *Arthrobacter*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Burkholderia*, *Caulobacter*, *Chromobacterium*, *Flavobacterium*, *Enterobacter*, *Erwinia*, *Herbaspirillum*, *Klebsiella*, *Micrococcus*, *Paenibacillus*, *Rhizobium*, *Pseudomonas*, *Serratia*, *Stenotrophomonas*, *Streptomyces*, *Variovorax*, and *Xanthomonas* have been shown to stimulate plant growth and development [91,92]. These phylogenetically diverse bacterial group have wide spectrum plant growth-promoting capabilities, and they can be categorized as biocontrol, biofertilizer, and phytostimulation agents [92,93,95,96].

##### 3.2.1.1. Enhanced nutrient acquisition (biofertilizer)

A major mechanism used by PGPR to stimulate crop growth and development is biofertilization. Several mineral nutrients such as nitrogen, phosphorus, and iron can be limited in the soil, thus limiting plant growth and development [92]. Nitrogen is the most liming factor for crop growth, although the geosphere contains  $1.6 \times 10^{17}$  t, most of which is found in the atmosphere with an estimated  $3.86 \times 10^{15}$  t [97]. Nitrogen (N<sub>2</sub>) represents around 78% of the atmosphere, and it is inaccessible to all plants and other eukaryotic life. Biological nitrogen fixation (BNF) is a process carried out by a few adapted prokaryotic diazotroph, that possess the enzyme nitrogenase, which catalyzes the reduction of N<sub>2</sub> to ammonia, a form of N utilized by plants [98]. Diazotrophic bacteria can be classified according to the degree of intimacy with plants: symbiotic N-fixing bacteria such as rhizobia, associative N-fixing bacteria such as *Azospirillum* spp., and free-living N-fixing bacteria such as *Azotobacter* spp. [92]. The efficiency and significant contribution of BNF from PGPR is well documented for several crops such as legumes, sugarcane, and grasses [98-100].

The other major nutrient limiting plant growth is phosphorus (P). Although soil often has abundant quantities of P (~0.05% w/w), only a small fraction of this P (~0.1%) is readily available for plant uptake [101,102]. Low availability of P in soils is due to the fact that the majority of soil-bounded P is present in insoluble form [102]. Plants can take-up P in two soluble forms, either as monobasic (H<sub>2</sub>PO<sub>4</sub><sup>-</sup>) or dibasic (HPO<sub>4</sub><sup>2-</sup>) ions [103]. A subset of bacteria, known as phosphate solubilizing bacteria (PSB), can influence the availability of P [104,105]. PSB are commonly found in the rhizosphere of plants and encompass genera such as *Azotobacter*, *Bacillus*, *Bradyrhizobium*, *Burkholderia*, *Enterobacter*, *Ralstonia*, *Rhizobium*, *Rhodococcus*, *Paenibacillus*, *Pseudomonas*, and *Serratia* [92,101]. These PSB can

solubilize insoluble forms of P to plant-available forms through different mechanisms such as the secretion of organic acids, siderophores, protons, hydroxyl ions, and CO<sub>2</sub>, as in the case of inorganic P solubilization [105,106]. These bacteria also produce many extracellular enzymes such as acid and alkaline phosphatases and phytases that stimulate organic P mineralization [104,105,107].

Iron is another essential plant nutrient that plays a key role in plant growth and development. Despite its abundance in the soil, most of iron is present in insoluble forms, mainly as ferric hydroxide [108]. Plant roots prefer to take-up the reduced form of iron, the ferrous (Fe<sup>2+</sup>) ion compared to the ferric (Fe<sup>3+</sup>) ion [92,109]. Siderophores are low-molecular-weight iron-chelating agents that are produced by many soil bacteria and fungi under stressed low iron conditions [110]. Bacterial produced-siderophores can enhance plant growth by enhancing plant iron nutrition through binding Fe<sup>3+</sup> and render it available for reduction to Fe<sup>2+</sup> [92,95]. Apart from improving plant iron nutrition, siderophores also stimulate plant growth indirectly via suppressing plant pathogen activities in the rhizosphere by depriving pathogens of Fe<sup>2+</sup> required for their cellular growth and development, thus lowering the probability of plant disease [94]. Additionally, siderophore-producing bacteria were shown to play an important role in enhancing plant growth in heavy metal-contaminated soils by alleviating heavy metal toxicity [111,112]. Several reports indicated that microbial siderophores bind and form stable compounds with other heavy metals such as Al, Cd, Cu, Pb, and Zn [95,111-113].

In addition to Pseudomonads, which synthesize high-affinity Fe<sup>3+</sup>-binding siderophores [114], several other PGPR are capable of producing siderophores including *Azospirillum* spp., *Azotobacter* spp., *Bacillus* spp., *Klebsiella* spp., *Nocardia* spp., *Paenibacillus* spp., *Pantoea* spp., *Serratia* spp., and *Streptomyces* spp. [92,95,110].

The role of PGPR in solubilizing and oxidizing other essential plant nutrients, such as potassium, sulfur, and micronutrients, and their effects on plant growth-promotion are far less studied compared to N, P, and Fe. For more information regarding the role of PGPR in providing these elements to plants, the reader can consult recent publications on this topic [95,115].

### 3.2.1.2. Plant growth regulation (phytostimulation)

Other direct modes of action employed by PGPR to stimulate plant growth and development is through production of phytohormones [96]. Various PGPR are known to synthesize different classes of phytohormones, including abscisic acid (ABA), auxins, ethylene, gibberellins, and cytokinins [92,96]. These phytohormones are responsible for many processes in plants during their different development stages. Auxins, for example, are responsible for cell enlargement and cell division, root initiation, increased fruit development, and leaf senescence [116]. Cytokinins on the other hand play a major role in the promotion of cell division and root hair formation, initiation and expansion of shoots and other plant parts, and decreased root growth [117]. Phytohormone gibberellins take part in regulating seed dormancy and germination, speeding up fruit and flowering processes, and modifying plant morphology, particularly stem elongation [118,119]. When produced at low concentration, the plant growth regulator ethylene is involved in many plant growth stages including stimulation of seed germination, formation and elongation of roots, and fruit and leaf maturation [120]. Finally, the phytohormone ABA plays main roles in seed development and maturation and mediating stomatal opening [119]. The most studied phytohormone, to date, produced by PGPR is auxin indole-3-acetic acid (IAA), which is involved in enhancing root growth and root length as well as formation and proliferation of lateral root hairs [116]. IAA-synthesizing PGPR include bacterial genera such as *Aeromonas*, *Agrobacterium*, *Alcaligenes*, *Azospirillum*, *Bradyrhizobium*, *Comamonas*, *Bacillus*, *Bradyrhizobium*, *Burkholderia*, *Enterobacter*, *Rhizobium*, and *Pseudomonas* [92,109,121].

### 3.2.1.3. Reduction of plant ethylene (stress alleviating)

Ethylene production by plants at low concentrations can be beneficial, as mentioned above. However, when produced at high concentrations, it can stunt plant growth and development by inhibiting root growth [122]. In response to various biotic and abiotic stressor conditions, plants synthesize different enzymes, metabolites, and stress proteins to alleviate the adverse effects of stress [123]; of particular interest is ethylene. Once plants encounter stress such as flooding, drought, or presence of toxic compounds, plant growth is inhibited because the ethylene precursor, 1-aminocyclopropane-1-carboxylate, is induced [124]. However, certain PGPR can hinder ethylene biosynthesis via production of 1-aminocyclopropane-1-carboxylate deaminase (ACCD) that cleaves the ethylene precursor ACC into alpha-ketobutyrate and ammonia [87,125], thus balancing ethylene levels and reducing its adverse impact on plant growth [125].

The beneficial roles of PGPR-containing ACCD have been studied in plants grown under different stress conditions such as drought [126], waterlogging [127], high salinity [128], and heavy metal contamination [129]. Several PGPR are known for their production of ACCD such as *Achromobacter* spp., *Azotobacter* spp., *Bacillus* spp., *Enterobacter* spp. *Herbaspirillum* spp., *Ochrobactrum* spp., *Pseudomonas* spp., and *Serratia* spp. [95,126-129].

### 3.2.2. Plant growth-promoting rhizobacteria-assisted phytoremediation

Over the last few decades, the immense interest in exploiting PGPR as a biofertilizer or biocontrol agent in agriculture has resulted in the development of successful commercial inoculants in many parts of the world, including Canada, Europe, and the United States [130-132]. Considerable research investigations have been conducted over the last decade to utilize PGPR in bioremediation of heavy metal-contaminated soils by promoting plant growth and also ameliorating the phytostabilization or phytoextraction efficiency [133-135]. However, the utilization of PGPR in rhizoremediation of PHC-contaminated soil is new and represents a large, untapped potential [2,30]. Recent reports of PGPR-assisted phytoremediation of PHC-contaminated soil and its host plants are summarized in Table 3

**Table 3.** Examples of the use of PGPR in rhizoremediation of PHC-contaminated soil.

Plant	Contaminants	Conditions	Bacteria	Role of PGPR	Reference
Ryegrass ( <i>Lolium perenne</i> )	Diesel	Greenhouse	<i>Pantoea</i> sp. strain BTRH79	ACC deaminase activity	[136]
<i>Cytisus striatus</i> and <i>Lupinus luteus</i>	Diesel	Greenhouse	<i>Bradyrhizobium japonicum</i> strain ER33	IAA, organic acids	[137]
<i>Cytisus striatus</i> and <i>Lupinus luteus</i>	Diesel	Greenhouse	<i>Streptomyces costaricanus</i> strain RP92	IAA, siderophores, organic acids	[137]
Leguminous plants and pastures	Oil refinery sludge	Field trial	<i>Pseudomonas putida</i> strain BIRD-1	P-solubilization, Siderophores, IAA	[138]
Tall fescue ( <i>Festuca arundinacea</i> )	Aliphatic hydrocarbons	Field trial	PGPR consortia	N/A	[139]

### 3.2.3. Hydrocarbon-degrading bacteria

#### 3.2.3.1. Ecology and diversity of PHC-degrading bacteria

The fate of most PHCs encountered in the terrestrial environment is degradation and/or biotransformation by soil bacteria. These bacteria are heterotrophs thus, they utilize PHC compounds as nutrient and energy sources for their cellular growth and development. PHC-degrading bacteria are widespread in nature and have been found in Arctic and Antarctic soils [140], aquatic environments [141], and pristine environments [142]. The abundance of PHCs in the environment maintains the degradation potential within most bacterial communities [143]. Interestingly, certain obligate hydrocarbonoclastic bacteria (OHCb) such as *Alcanivorax* spp., *Cycloclasticus* spp., *Marinobacter* spp., *Oleispira* spp., *Planomicrombium* spp., and *Thallassolituus* spp. are found undetectable or in low abundance in unpolluted environments; however, they prevail after PHC pollution occurs [141,144,145].

Over the last few decades, many bacterial species have been isolated and identified from various terrestrial and aquatic environments [141,144,146]. Some of these bacteria can utilize a wide spectrum of PHC compounds; for example, the bacterial strain *Dietzia* sp. DQ12-45-1b could grow on many *n*-alkanes (C6–C40) and other monoaromatic and polycyclic aromatic hydrocarbons as the sole carbon source and energy [147]. To date, more than 79 bacterial genera that can degrade PHCs have been isolated and identified [144,145], such as *Achromobacter*, *Acinetobacter*, *Alkanindiges*, *Alteromonas*, *Arthrobacter*, *Bacillus*, *Burkholderia*, *Dietzia*, *Enterobacter*, *Kocuria*, *Marinobacter*, *Mycobacterium*, *Nocardia*, *Pseudomonas*, *Rhodococcus*, *Streptomyces*, and *Variovorax* [145,146,148]. As different bacteria vary in their catalytic enzyme activity, no single bacterial species can break down the entire PHC fraction completely [145,146]; therefore, their effectiveness in remediating PHC-polluted sites also varies widely [145].

#### 3.2.3.2. Alkane-degrading bacteria

Alkanes are saturated hydrocarbons that can be further classified as branched (*isoo*-alkanes), cyclic (*cyclo*-alkanes), or linear (*n*-alkanes) [149]. Although many living organisms such as bacteria, plants, and green algae produce alkanes [150,151], the main source of alkanes in terrestrial environments comes from PHC contamination, as alkanes are the main constituent of crude oil and natural gas [149,152]. Bacterial alkane degradation is of great significance for the bioremediation of PHC-contaminated soil as well as for microbial enhanced oil recovery [153]. Bacteria metabolize alkanes under both aerobic and anaerobic conditions [149]. Most bacteria degrade alkanes aerobically; therefore, aerobic degradation will be discussed hereafter.

Aerobic degradation of alkanes starts with terminal or sub-terminal incorporation of oxygen atoms ( $O_2$ ) into the hydrocarbon substrate by an alkane hydroxylase enzyme [152]. Alkane hydroxylases (AHs) are a class of several specific enzymes that insert  $O_2$  into the hydrocarbons to initiate degradation [154]. Depending on the chain length of the alkane substrate, there are different enzymes classes that carry out the oxygenation of hydrocarbons [154] (Table 4). For example, bacteria degrading short-chain alkanes (C2-C4) have enzymes related to methane monooxygenases, while bacterial strains degrading medium-chain alkanes (C5-C20) usually contain alkane 1-monooxygenase and soluble cytochrome P450 enzymes, and bacterial strains degrading long-chain alkanes ( $>C20$ ) contain several recently discovered types of AHs such as flavin-binding monooxygenase and thermophilic flavin-dependent monooxygenase [149,152,154,155] (Table 4).

**Table 4.** Examples of alkane-degrading genes, enzymes, and their bacterial source.

Enzyme class	Substrate range	Gene	Bacterial species
Soluble methane Monooxygenase	C1-C8	<i>mmoX</i>	<i>Gordonia, Methylococcus, Methylosinus, Methylocystis, Methylomonas, Methylocella.</i>
Particulate methane Monooxygenase	C1-C5	<i>pmoC</i>	<i>Methylococcus, Methylosinus, Methylocystis, Methylobacter, Methylomonas, Methylomicrobium, Nocardioides.</i>
Alkane 1-monooxygenase	C10-C20	<i>alkB</i>	<i>Acinetobacter, Alcanivorax, Burkholderia, Mycobacterium, Pseudomonas, Rhodococcus.</i>
Soluble cytochrome P450	C5-C16	<i>CYP153</i>	<i>Acinetobacter, Alcanivorax, Caulobacter, Mycobacterium, Rhodococcus, Sphingomonas.</i>
Flavin-binding monooxygenase	C20-C36	<i>Alma</i>	<i>Alcanivorax, Marinobacter, Acinetobacter.</i>
Thermophilic flavin-dependent monooxygenase	C10-C36	<i>LadA</i>	<i>Geobacillus thermodenitrificans NG80-2</i>

Adapted and modified from [149, 152, 154, 155].

Among the above-mentioned (AHs) enzyme systems, alkane 1-monooxygenase (encoded by *alkB*) is the most common found in alkanes degrading  $\alpha$ -,  $\beta$ -, and  $\gamma$ -*Proteobacteria* and high G+C content Gram-positive bacteria [149,152,154,155]. The substrates for AlkB-harboring bacteria comprise alkanes ranging from C10 to C16 [154]; however, some AlkB-harboring *Actinobacteria* such as *Dietzia* sp. and *Gordonia* sp. can degrade alkanes with chain lengths up to C32 [156,157]. Another bacterial AH enzyme system for degradation of short- and medium-chain substrates is cytochrome P450 hydroxylase of the CYP153 family, which is frequently found in alkane-degrading bacteria lacking the AlkB enzyme [158,159]. It is common that bacterial strains contain more than one alkB homologous gene, as in the case of *Rhodococcus* strain Q15, which contains at least four alkane 1-monooxygenases [160]. Additionally, several bacterial strains have more than one AH system, as has been shown in *Dietzia* sp. strain DQ12-45-1b, which has AlkB and CYP153 systems co-existing together [159]. The co-existence of more than one AH system in bacteria can expand its ability to degrade a wider alkane range [158,159]. *AlkB* and *CYP153* genes are commonly assessed to determine the degradation potential of bacterial communities in PHC-impacted soil and water environments [161,164].

### 3.2.3.3. Polycyclic aromatic hydrocarbon-degrading bacteria

The other major fraction of PHCs is polycyclic aromatic hydrocarbons (PAHs). PAHs are ubiquitous in nature. They have two or more aromatic benzene rings in their structure [165]. PAHs are found in nature as a byproduct of many biogeochemical and biological processes as well as incomplete combustion of woods, coal, and gasoline [166]. However, the main entry source of PAHs in the environment is industrial activities related to the petroleum and gas industry [165,166]. Due to their electrochemical stability, high

persistence in terrestrial environments, bio-accumulative behaviors, and their “multi-faceted disease-causing” effects (carcinogenic, mutagenic, teratogenic), the United States Environmental Protection Agency (U.S. EPA), as well as agencies in many other countries, has listed 16 PAH compounds as priority pollutants [165,167].

The main principle of PAH biodegradation, mediated by aerobic bacteria, involves activation and subsequent cleavage of the thermodynamically stable benzene ring in PAH substrates [166]. Under aerobic conditions, the initial step is the hydroxylation of the benzene ring by dioxygenase enzymes, resulting in the formation of *cis*-dihydrodiols. After this step, *cis*-dihydrodiols are further dehydrogenated, via the action of dehydrogenase enzymes, to form several dihydroxylated intermediates. Subsequently, these diol intermediates are cleaved by intradiol or extradiol ring-cleaving dioxygenase enzymes, leading to the formation of central intermediates such as protocatechuates and catechols that can be further metabolized to acetone, succinate, or pyruvate, which then enter the tricarboxylic acid cycle (TCA) cycle [168-171].

PAHs are broadly classified into low-molecular-weight (LMW PAHs) compounds with fewer than three rings (<3 rings), and high-molecular-weight (HMW PAHs) compounds with more than three rings (>3 rings) [166]. LMW PAHs such as naphthalene, phenanthrene, anthracene, fluorene, acenaphthene, and acenaphthylene are found in high quantities in PHC-contaminated sites, and diverse bacterial communities have the ability to utilize these LMW PAHs as the sole carbon and energy source [170,171]. LMW PAH-degrading bacteria are ubiquitous in nature, and the isolation, degradation potential, and elucidation of catabolic pathways, enzymatic machineries, and genetic regulations within these bacteria are well documented [169-171]. For example, several bacterial genera are well known for their high efficiency to degrade LMW PAHs such as *Acinetobacter*, *Comamonas*, *Novosphingobium*, *Ochrobactrum*, *Ralstonia*, *Rhodococcus*, *Pseudomonas*, *Sphingomonas*, *Sphingobium*, and *Staphylococcus* [169-171].

Unlike LMW PAHs, HMW PAHs, due to their stable physicochemical structures, do not biodegrade easily; however, research on bacterial degradation of HMW PAHs has advanced significantly over the last two decades [171,172]. Several HMW PAH bioavailability-enhancing strategies and adaptation mechanisms have been identified and include biofilm formation, cell surface hydrophobicity, low requirements for energy and O<sub>2</sub> for cell growth and maintenance, high substrate uptake affinity, production of biosurfactants, and wide substrate utilization patterns [169,172,173]. More importantly, the functions of diverse, versatile catabolic genes involved in HMW PAH degradation and enzymatic activities, as well as their regulation mechanisms, have been discovered in various HMW PAH-degrading bacteria [172]. Table 5 lists genes and enzymes involved in both LMW and HMW PAH biodegradation.

**Table 5.** Examples of PAH-degrading genes, enzymes, and their bacterial source.

Enzyme.	Gene	Bacterial source	Reference
Naphthalene dioxygenase	<i>Nah</i>	<i>Pseudomonas putida</i> strain G7	[174]
Phenanthrene dioxygenase	<i>phnAc</i>	<i>Burkholderia</i> sp. strain RP007	[175]
Pyrene dioxygenase	<i>nidA</i>	<i>Mycobacterium</i> sp. strain PYR-1	[176]
Extradiol dioxygenase	<i>phdF</i>	<i>Mycobacterium</i> sp. strain SNP11	[177]
Catechol 1,2-dioxygenase	<i>C12O</i>	<i>Pseudomonas</i> sp. strain EST1001	[178]

Although most of the HMW PAH-degrading bacteria described so far belong to the *Actinobacteria* phylum, representing genera such as *Arthrobacter*, *Corynebacterium*, *Dietzia*, *Gordonia*, *Microbacterium*, *Mycobacterium*, *Nocardia*, *Rhodococcus*, and *Streptomyces* [1,171], a variety of non-actinomycete bacterial genera such as *Achromobacter*, *Burkholderia*, *Pseudomonas*, *Sphingomonas*, *Sphingobium*, and *Stenotrophomonas* have been reported as well [169,171].

#### 4. Enhancing the understanding of mechanisms through which host plants assemble a beneficial microbiome, and how it functions, under pollutant stress

A challenging problem facing studies of the microbiome in many disciplines is the fact that the majority of microbial taxa are resistant to cultivation using current culture-dependent techniques. However, over the past decade or so, the advancement of next-generation sequencing and bioinformatics has paved the way to enhance our understanding of the structure, function, and composition of microbial communities in different habitats and environmental conditions, including PHC-contaminated soil [179].

##### 4.1. High-throughput amplicon sequencing

Studies of the rhizosphere microbiome in natural and agricultural settings have generated most of our knowledge about host plant selection processes and plant–microbiome interactions taking place in the rhizosphere and how plants recruit different microbiota from surrounding environments [31,180]. For example, previous studies using 16S rRNA amplicon sequencing revealed that microbial communities in the rhizosphere and adjunct bulk soils are different; the recruitment of rhizosphere microbiota by plants is strongly dependent on the structure and composition of the bulk soil microbiome [44], and different plant genotypes were found to select for different rhizosphere microbiomes [45].

Over the past few years, several experiments have been conducted to optimize phytoremediation systems and improve their efficacy using high-throughput sequencing approaches. For example, Bell et al. [67] used high-throughput 454-pyrosequencing of bacterial 16S rRNA genes and the fungal internal transcribed spacer (ITS) region to compare the community structure and composition of the rhizosphere microbiome of native and non-native *Salix* cultivars across uncontaminated and PHC-contaminated soil. Their results indicated higher fungal sensitivity to PHC contamination than that found for bacterial communities. Additionally, certain fungal class (*Pezizomycetes*) reacted differently following plant introduction to soils [67], implying the importance of plant species selection in phytoremediation with regard to their impact on plant-associated microbiomes [31]. Similarly, Hassan et al. [82] used 454-pyrosequencing of the AMF 18S rDNA gene to examine how rhizospheric AMF communities are shaped within the rhizosphere of 11 *Salix* cultivars introduced across non-contaminated and PHC-contaminated soil. While PHC contamination levels had a strong impact on AMF community structure, *Salix* planting increased the abundance of several AMF families [82], inferring that AMF, possibly due to opportunistic associations with the plant, are involved in plant adaptation to PHC contamination [31].

Tardif et al. [181] amplified the bacterial 16S rRNA gene and fungal ITS regions using Ion Torrent sequencing in order to characterize the variations between plant

compartments (bulk soil, rhizosphere soil, roots, and stems) in the microbiome of two *Salix* cultivars growing under three PHC contamination levels at a former petrochemical site. PHC contamination was found to be the main factor shaping not only the rhizosphere but also the root and stem microbiome structure [181]. Additionally, the presence of the plant offered a protective buffer zone against PHC pollution in the rhizosphere and other plant tissues, subsequently minimizing the severe effects of PHC contamination on the microbiome composition, as compared with adjunct bulk soil [181]. Finally, increasing PHC contamination caused a shift in the microbial community composition, favoring beneficial microbiome communities such as putative PHC-degraders and PGPR [181].

In a recent study, Mitter et al. [182] used high-throughput Illumina MiSeq amplicon sequencing of the 16S rRNA gene to characterize the bacterial root microbiome associated with annual barley and sweet clover growing in an oil sands reclamation site. Results confirmed that, consistent with previous reports, the rhizosphere compartment produced the strongest differentiation of the root microbiome community structure [44,182,183]; for example, *Proteobacteria* was the predominant phyla in the endosphere microbiome, whereas phyla such as *Acidobacteria* and *Gemmatimonadetes* were restricted only to the rhizosphere microbiome [182]. Additionally, host plants play a major role in shaping the root microbiome community structure [182], implying plants have the ability to select for specific soil microbiota [182].

#### 4.2. Metatranscriptomics

Metatranscriptomics refers to the study of mRNA expression profiles of complex microbial communities within natural environments [184]. Unlike metagenomics, metatranscriptomics can enhance our understanding about how microbiome functions can be altered due to PHC contamination [184]. In the context of phytoremediation, metatranscriptomics has been used to study plant-associated microbial activities in PHC-contaminated soil. For example, in a greenhouse experiment, Yergeau et al. [53] compared the expression of functional genes in the rhizosphere and bulk soil of willow plants growing in contaminated and uncontaminated soil using a metatranscriptomics approach. Combined selective pressure of the pollutants and rhizosphere resulted in an increased expression of genes related to competition, such as antibiotic resistance and biofilm formation, in the contaminated rhizosphere [53]. Additionally, genes related to PHC degradation were more expressed in polluted soils [53].

More recently, Yergeau et al. [69] sequenced the rhizosphere metatranscriptome of four willow species and the plant root metatranscriptome for two willow species growing in PHC-contaminated and non-contaminated soil at a former petroleum refinery site. The abundance of transcripts for many microbial taxa and functions were significantly higher in contaminated rhizosphere soil for *Salix eriocephala*, *S. miyabeana*, and *S. purpurea*, compared to the rhizosphere of *S. caprea* [69]. The root metatranscriptomes of two willow cultivars were compared, showing that plant transcripts were mostly influenced by willow species, while microbial transcripts primarily responded to contamination level [69].

Page et al. [185] used a transcriptomics-based approach to identify microbes involved in willow-microbes PHC degradation systems. Enhanced expression of the four genes related to PHC degradation was observed within the bacterial orders *Actinomycetales*, *Rhodospirillales*, *Burkholderiales*, *Alteromonadales*, *Solirubrobacteriales*, *Caulobacterales*, and *Rhizobiales*, implying that members of these microbial taxa are active participants in the willow-microbes association [185]. Information obtained from metatranscriptomics studies on complex systems, such as plants and their associated highly diverse microbial communities, growing in PHC-contaminated soil could help optimize phytoremediation and enhance their use [53,69,184,185].

#### 4.3. Genome sequencing

Due to the decrease in cost and difficulty over the past decade, sequencing, assembly, and annotation of bacterial genomes is becoming a relatively common practice in many fields of microbiology, including environmental microbiology. By sequencing the entire bacterial genome, valuable information can be obtained such as isolate identification, finding important bacterial traits, life style, ecological adaptation, genetic structure, and metabolic pathways.

Over the past few years, many complete and draft genome sequences of bacterial strains, with versatile abilities to degrade PHCs, have been published and are available in public databases [167]. The genome sequences of different PHC-degrading bacterial strains provide structures for sets of genes, operons, and degradative pathways responsible for remediation of PHC-contaminated environments [167]. Some of these bacterial genomes and their importance in rhizoremediation are listed in Table 6.

Zhao et al. [186] reported the complete genome sequence of *Sphingobium yanoikuyae* strain B1 that has versatile abilities to degrade various PHCs pollutants, such as biphenyl, naphthalene, phenanthrene, toluene, and anthracene. The 5,200,045 bp genome of this bacterium contains 35 dioxygenases or putative dioxygenases genes, including catechol 1,2-dioxygenase, biphenyl 2,3-dioxygenase, and biphenyl-2,3-diol 1,2-dioxygenase [186]. Additionally, the genome of *S. yanoikuyae* strain B1 contains 48 ABC transporter-related genes and 82 TonB-dependent receptors, which may be involved in PAH transportation [186]. Such valuable information can provide clues about the genetic versatility of *Sphingobium* strains and the mechanisms of PAHs biodegradation, which might potentially aid in rhizoremediation applications [186].

**Table 6.** Recent genomes of bacterial strains capable of degrading PHCs.

Bacterial strain	Importance in bioremediation	Isolation source	PGPR features	Genome size	Reference
<i>Pseudomonas veronii</i> strain VI4T1	degradation of aromatic and aliphatic hydrocarbons	long-term oil field-polluted soil	IAA, siderophore	7.15 Mb	[187]
<i>Pseudomonas</i> sp. strain VI4.1	degradation of aromatic and aliphatic hydrocarbons	long-term oil field-polluted soil	IAA, siderophore	7.3 Mb	[187]
<i>Halomonas</i> sp. strain G11	degradation of alkanes and polyaromatic hydrocarbons	hypersaline sediment	Salt-tolerance, biosurfactant production	3.96 Mb	[188]
<i>Pseudomonas aeruginosa</i> strain DN1	fluoranthene degradation	petroleum-contaminated soil	N/D	6.6 Mb	[189]
<i>Alcaligenes aquatilis</i> strain BU33N	degradation of n-alkanes and phenanthrene	hydrocarbon polluted sediments	biosurfactant production; heavy metals resistance	3.8 Mb	[190]
<i>Gordonia paraffinivorans</i> strain MTZ052	degradation of n-hexadecane	composting pile	N/D	4.8 Mb	[191]
<i>Gordonia sihwensis</i> strain MTZ096	degradation of n-hexadecane	composting pile	N/D	3.9 Mb	[191]
<i>Klebsiella pneumoniae</i> strain AWD5	degradation of xenobiotic compounds	PAH-contaminated soil	siderophore production	4.8 Mb	[192]
<i>Bacillus licheniformis</i> strain VSD4	degradation of diesel fuel	leaves of <i>Hedera helix</i> plants growing at a high-traffic city center	IAA, siderophore	4.19 Mb	[193]
<i>Pseudomonas putida</i> strain BS3701	degradation of crude oil and PAHs	soil contaminated with coke by-product waste	N/D	6.3 Mb	[194]

## 5. Concluding remarks and future perspectives

In spite of the remarkable progress detailed above, rhizoremediation remains a marginal choice for *in situ* soil decontamination. Given the important role of the rhizosphere microbiome in phytoremediation, future efforts to optimize this technology should include (i) selection of the right plant host, which can alter the function of the rhizosphere microbiome to benefit rhizoremediation activities. Special emphasis should be focus on selecting native plants that show tolerance toward PHCs. Using such plants could offer economic and environmentally sustainable solutions to remediate PHC-contaminated soil. (ii) Modern microbial ecology omics-tools should be used not only to better understand the structure and function of the rhizosphere microbiome associated with plants but also to recommend more efficient management strategies and predict the clean-up time of rhizoremediation. (iii) Large-scale field experiments the effect of novel microbiome inocula combining PGPR and hydrocarbon-degrading bacteria. (iv) The complicity of the rhizosphere environment and the influence of many biotic and abiotic factors on the composition and function of rhizosphere microbiome should be taken into account, which might subsequently affect rhizoremediation efficiency. Therefore, it would be important to characterize biotic and abiotic parameters in PHC-contaminated sites prior to application of rhizoremediation strategies.

**Author Contributions:** Writing, FA. MH and MSA supervised the work, edited, and revised the draft of the manuscript. All authors approved submission of the manuscript.

**Funding:** We acknowledge funding for this work by the GenoRem Project, which is primarily financed by Genome Canada and Genome Quebec, and by NSERC discovery grants to MSA and MH. FA was also supported by a grant from King Saud University. The GenoRem project contains several industrial partners, but these partners have in no way influenced or modified this manuscript

## References

1. Brzeszcz, J.; Kaszycki, P., Aerobic bacteria degrading both n-alkanes and aromatic hydrocarbons: an undervalued strategy for metabolic diversity and flexibility. *Biodegradation* **2018**, *29* (4), 359-407.
2. Gkorezis, P.; Daghio, M.; Franzetti, A.; Van Hamme, J. D.; Sillen, W.; Vangronsveld, J., The Interaction between Plants and Bacteria in the Remediation of Petroleum Hydrocarbons: An Environmental Perspective. *Front Microbiol* **2016**, *7*, 1836.
3. Wang, Z., Fingas, M., Yang, C., Christensen, J. H., Crude Oil and Refined Product Fingerprinting: Principles. In *Environmental Forensics: Contaminant Specific Guide*, Morrison, R. D., Murphy, B. L., Ed. Academic Press: New York, 2006; pp 339-407.
4. Farrell-Jones, J., *Petroleum hydrocarbons and polycyclic aromatic hydrocarbons*. Blackwell Publishing CRC Press: New York: 2003.
5. Tissot, B. P.; Welte, D. H., *Petroleum formation and occurrence*. Springer Science & Business Media: 2013.
6. Pandey, P.; Pathak, H.; Dave, S., Microbial ecology of hydrocarbon degradation in the soil: a review. *Research Journal of Environmental Toxicology* **2016**, *10* (1), 1.
7. Hoang, S. A.; Lamb, D.; Seshadri, B.; Sarkar, B.; Choppala, G.; Kirkham, M. B.; Bolan, N. S., Rhizoremediation as a green technology for the remediation of petroleum hydrocarbon-contaminated soils. *J Hazard Mater* **2021**, *401*, 123282.
8. Secretariat Federal Contaminated Sites Inventory. <https://www.tbs-sct.gc.ca/fcsi-rscf/home-accueil-eng.aspx> (accessed August 18, 2020).
9. Panagos, P.; Van Liedekerke, M.; Yigini, Y.; Montanarella, L., Contaminated sites in Europe: review of the current situation based on data collected through a European network. *J Environ Public Health* **2013**, *2013*, 158764.
10. Henner, P.; Schiavon, M.; Morel, J.-L.; Lichtfouse, E., Polycyclic aromatic hydrocarbon (PAH) occurrence and remediation methods. *Analisis* **1997**, *25* (9-10), M56-M59.
11. Inoue, Y.; Katayama, A., Two-scale evaluation of remediation technologies for a contaminated site by applying economic input-output life cycle assessment: risk-cost, risk-energy consumption and risk-CO<sub>2</sub> emission. *J Hazard Mater* **2011**, *192* (3), 1234-42.

12. Yerushalmi, L.; Rocheleau, S.; Cimpoia, R.; Sarrazin, M.; Sunahara, G.; Peisajovich, A.; Leclair, G.; Guiot, S. R., Enhanced biodegradation of petroleum hydrocarbons in contaminated soil. *Journal of Soil Contamination* **1998**, *7* (1), 37-51.
13. Pilon-Smits, E., Phytoremediation. *Annu Rev Plant Biol* **2005**, *56*, 15-39.
14. Salt et al., Phytoremediation. *Annual review of plant biology* **1998**, *49* (1), 643-668.
15. Aken, B. V.; Doty, S. L., Transgenic plants and associated bacteria for phytoremediation of chlorinated compounds. *Biotechnology and Genetic Engineering Reviews* **2009**, *26* (1), 43-64.
16. Panz, K.; Miksch, K., Phytoremediation of explosives (TNT, RDX, HMX) by wild-type and transgenic plants. *Journal of Environmental Management* **2012**, *113*, 85-92.
17. Leguizamo, M. A. O.; Gómez, W. D. F.; Sarmiento, M. C. G., Native herbaceous plant species with potential use in phytoremediation of heavy metals, spotlight on wetlands—a review. *Chemosphere* **2017**, *168*, 1230-1247.
18. Jerez Ch, J. A.; Romero, R. M., Evaluation of Cajanus cajan (pigeon pea) for phytoremediation of landfill leachate containing chromium and lead. *International journal of phytoremediation* **2016**, *18* (11), 1122-1127.
19. Olette, R.; Couderchet, M.; Biagiotti, S.; Eullaffroy, P., Toxicity and removal of pesticides by selected aquatic plants. *Chemosphere* **2008**, *70* (8), 1414-1421.
20. Newman, L. A.; Reynolds, C. M., Phytodegradation of organic compounds. *Current opinion in Biotechnology* **2004**, *15* (3), 225-230.
21. Sharma, S.; Singh, B.; Manchanda, V., Phytoremediation: role of terrestrial plants and aquatic macrophytes in the remediation of radionuclides and heavy metal contaminated soil and water. *Environmental Science and Pollution Research* **2015**, *22* (2), 946-962.
22. Devi, S.; Nandwal, A.; Angrish, R.; Arya, S.; Kumar, N.; Sharma, S., Phytoremediation potential of some halophytic species for soil salinity. *International journal of phytoremediation* **2016**, *18* (7), 693-696.
23. Transparency Market Research: Bioremediation technology & services market to reach valuation of ~US\$ 20 BN by 2030: TRANSPARENCY MARKET RESEARCH. <https://www.transparencymarketresearch.com/pressrelease/bioremediation-technology-services-market.htm> (accessed September 18, 2020).

24. Mench, M.; Lepp, N.; Bert, V.; Schwitzguébel, J.-P.; Gawronski, S. W.; Schröder, P.; Vangronsveld, J., Successes and limitations of phytotechnologies at field scale: outcomes, assessment and outlook from COST Action 859. *Journal of Soils and Sediments* **2010**, *10* (6), 1039-1070.
25. Wenzel, W. W., Rhizosphere processes and management in plant-assisted bioremediation (phytoremediation) of soils. *Plant and Soil* **2009**, *321* (1), 385-408.
26. Vangronsveld, J.; Herzig, R.; Weyens, N.; Boulet, J.; Adriaensen, K.; Ruttens, A.; Thewys, T.; Vassilev, A.; Meers, E.; Nehnevajova, E.; van der Lelie, D.; Mench, M., Phytoremediation of contaminated soils and groundwater: lessons from the field. *Environ Sci Pollut Res Int* **2009**, *16* (7), 765-94.
27. Kuiper, I.; Lagendijk, E. L.; Bloemberg, G. V.; Lugtenberg, B. J., Rhizoremediation: a beneficial plant-microbe interaction. *Molecular plant-microbe interactions* **2004**, *17* (1), 6-15.
28. Anderson, T. A.; Guthrie, E. A.; Walton, B. T., Bioremediation in the rhizosphere. *Environmental Science & Technology* **1993**, *27* (13), 2630-2636.
29. Martin, B. C.; George, S. J.; Price, C. A.; Ryan, M. H.; Tibbett, M., The role of root exuded low molecular weight organic anions in facilitating petroleum hydrocarbon degradation: current knowledge and future directions. *Sci Total Environ* **2014**, *472*, 642-53.
30. Correa-García, S.; Pande, P.; Séguin, A.; St-Arnaud, M.; Yergeau, E., Rhizoremediation of petroleum hydrocarbons: a model system for plant microbiome manipulation. *Microb Biotechnol* **2018**, *11* (5), 819-832.
31. Thijss, S.; Sillen, W.; Rineau, F.; Weyens, N.; Vangronsveld, J., Towards an Enhanced Understanding of Plant-Microbiome Interactions to Improve Phytoremediation: Engineering the Metaorganism. *Front Microbiol* **2016**, *7*, 341.
32. Arthur, E. L.; Rice, P. J.; Rice, P. J.; Anderson, T. A.; Baladi, S. M.; Henderson, K. L.; Coats, J. R., Phytoremediation—an overview. *Critical Reviews in Plant Sciences* **2005**, *24* (2), 109-122.
33. Ali, H.; Khan, E.; Sajad, M. A., Phytoremediation of heavy metals--concepts and applications. *Chemosphere* **2013**, *91* (7), 869-81.
34. Behera, K. K., Phytoremediation, transgenic plants and microbes. In *Sustainable agriculture reviews*, Springer: 2014; pp 65-85.

35. Philippot, L.; Raaijmakers, J. M.; Lemanceau, P.; van der Putten, W. H., Going back to the roots: the microbial ecology of the rhizosphere. *Nat Rev Microbiol* **2013**, *11* (11), 789-99.
36. Raaijmakers, J. M.; Paulitz, T. C.; Steinberg, C.; Alabouvette, C.; Moënne-Locoz, Y., The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant and soil* **2009**, *321* (1), 341-361.
37. Berendsen, R. L.; Pieterse, C. M.; Bakker, P. A., The rhizosphere microbiome and plant health. *Trends Plant Sci* **2012**, *17* (8), 478-86.
38. Berg, G.; Smalla, K., Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiol Ecol* **2009**, *68* (1), 1-13.
39. Gaiero, J. R.; McCall, C. A.; Thompson, K. A.; Day, N. J.; Best, A. S.; Dunfield, K. E., Inside the root microbiome: bacterial root endophytes and plant growth promotion. *Am J Bot* **2013**, *100* (9), 1738-50.
40. Bakker, P. A.; Berendsen, R. L.; Doornbos, R. F.; Wintermans, P. C.; Pieterse, C. M., The rhizosphere revisited: root microbiomics. *Front Plant Sci* **2013**, *4*, 165.
41. Kuzyakov, Y.; Blagodatskaya, E., Microbial hotspots and hot moments in soil: Concept & review. *Soil Biology and Biochemistry* **2015**, *83*, 184-199.
42. Smalla, K.; Wieland, G.; Buchner, A.; Zock, A.; Parzy, J.; Kaiser, S.; Roskot, N.; Heuer, H.; Berg, G., Bulk and rhizosphere soil bacterial communities studied by denaturing gradient gel electrophoresis: plant-dependent enrichment and seasonal shifts revealed. *Appl Environ Microbiol* **2001**, *67* (10), 4742-51.
43. Loepmann, S.; Blagodatskaya, E.; Pausch, J.; Kuzyakov, Y., Substrate quality affects kinetics and catalytic efficiency of exo-enzymes in rhizosphere and detritusphere. *Soil Biology and Biochemistry* **2016**, *92*, 111-118.
44. Bulgarelli, D.; Rott, M.; Schlaeppi, K.; Ver Loren van Themaat, E.; Ahmadinejad, N.; Assenza, F.; Rauf, P.; Huettel, B.; Reinhardt, R.; Schmelzer, E.; Peplies, J.; Gloeckner, F. O.; Amann, R.; Eickhorst, T.; Schulze-Lefert, P., Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* **2012**, *488* (7409), 91-5.
45. Lundberg, D. S.; Lebeis, S. L.; Paredes, S. H.; Yourstone, S.; Gehring, J.; Malfatti, S.; Tremblay, J.; Engelbrektson, A.; Kunin, V.; Del Rio, T. G.; Edgar, R. C.; Eickhorst, T.; Ley,

- R. E.; Hugenholtz, P.; Tringe, S. G.; Dangl, J. L., Defining the core *Arabidopsis thaliana* root microbiome. *Nature* **2012**, *488* (7409), 86-90.
46. Jambon, I.; Thijs, S.; Weyens, N.; Vangronsveld, J., Harnessing plant-bacteria-fungi interactions to improve plant growth and degradation of organic pollutants. *Journal of Plant Interactions* **2018**, *13* (1), 119-130.
47. Vieira, S.; Sikorski, J.; Dietz, S.; Herz, K.; Schrumpf, M.; Bruelheide, H.; Scheel, D.; Friedrich, M. W.; Overmann, J., Drivers of the composition of active rhizosphere bacterial communities in temperate grasslands. *Isme j* **2020**, *14* (2), 463-475.
48. Vives-Peris, V.; de Ollas, C.; Gómez-Cadenas, A.; Pérez-Clemente, R. M., Root exudates: from plant to rhizosphere and beyond. *Plant Cell Rep* **2020**, *39* (1), 3-17.
49. Rohrbacher, F.; St-Arnaud, M., Root exudation: the ecological driver of hydrocarbon rhizoremediation. *Agronomy* **2016**, *6* (1), 19.
50. Jones, D.; Murphy, D., Microbial response time to sugar and amino acid additions to soil. *Soil Biology and Biochemistry* **2007**, *39* (8), 2178-2182.
51. Gao, Y.; Yang, Y.; Ling, W.; Kong, H.; Zhu, X., Gradient distribution of root exudates and polycyclic aromatic hydrocarbons in rhizosphere soil. *Soil Science Society of America Journal* **2011**, *75* (5), 1694-1703.
52. Muratova, A.; Dubrovskaya, E.; Golubev, S.; Grinev, V.; Chernyshova, M.; Turkovskaya, O., The coupling of the plant and microbial catabolisms of phenanthrene in the rhizosphere of *Medicago sativa*. *J Plant Physiol* **2015**, *188*, 1-8.
53. Yergeau, E.; Sanschagrin, S.; Maynard, C.; St-Arnaud, M.; Greer, C. W., Microbial expression profiles in the rhizosphere of willows depend on soil contamination. *Isme j* **2014**, *8* (2), 344-58.
54. Gao, Y.; Ren, L.; Ling, W.; Gong, S.; Sun, B.; Zhang, Y., Desorption of phenanthrene and pyrene in soils by root exudates. *Bioresour Technol* **2010**, *101* (4), 1159-65.
55. van Elsas, J. D.; Bailey, M. J., The ecology of transfer of mobile genetic elements. *FEMS Microbiol Ecol* **2002**, *42* (2), 187-97.
56. Sentchilo, V.; Mayer, A. P.; Guy, L.; Miyazaki, R.; Green Tringe, S.; Barry, K.; Malfatti, S.; Goessmann, A.; Robinson-Rechavi, M.; van der Meer, J. R., Community-wide plasmid gene mobilization and selection. *Isme j* **2013**, *7* (6), 1173-86.

57. Top, E. M.; Springael, D., The role of mobile genetic elements in bacterial adaptation to xenobiotic organic compounds. *Curr Opin Biotechnol* **2003**, *14* (3), 262-9.
58. Khan, S.; Afzal, M.; Iqbal, S.; Khan, Q. M., Plant-bacteria partnerships for the remediation of hydrocarbon contaminated soils. *Chemosphere* **2013**, *90* (4), 1317-32.
59. Chaudhry, Q.; Blom-Zandstra, M.; Gupta, S.; Joner, E. J., Utilising the synergy between plants and rhizosphere microorganisms to enhance breakdown of organic pollutants in the environment. *Environ Sci Pollut Res Int* **2005**, *12* (1), 34-48.
60. Aprill, W.; Sims, R. C., Evaluation of the use of prairie grasses for stimulating polycyclic aromatic hydrocarbon treatment in soil. *Chemosphere* **1990**, *20* (1-2), 253-265.
61. Gaskin, S. E.; Bentham, R. H., Rhizoremediation of hydrocarbon contaminated soil using Australian native grasses. *Sci Total Environ* **2010**, *408* (17), 3683-8.
62. Hall, J.; Soole, K.; Bentham, R., Hydrocarbon phytoremediation in the family Fabaceae--a review. *Int J Phytoremediation* **2011**, *13* (4), 317-32.
63. Reynolds, C.; Wolf, D.; Gentry, T.; Perry, L.; Pidgeon, C.; Koenen, B.; Rogers, H.; Beyrouty, C., Plant enhancement of indigenous soil micro-organisms: A low-cost treatment of contaminated soils. *Polar Record* **1999**, *35* (192), 33-40.
64. Frick, C.; Germida, J.; Farrell, R. In *Assessment of phytoremediation as an in-situ technique for cleaning oil-contaminated sites*, technical seminar on chemical spills, Environment Canada; 1998: 1999; pp 105a-124a.
65. Lu, M.; Zhang, Z.; Sun, S.; Wei, X.; Wang, Q.; Su, Y., The use of goosegrass (*Eleusine indica*) to remediate soil contaminated with petroleum. *Water, Air, & Soil Pollution* **2010**, *209* (1), 181-189.
66. Kuzovkina, Y. A.; Volk, T. A., The characterization of willow (*Salix L.*) varieties for use in ecological engineering applications: co-ordination of structure, function and autecology. *Ecological Engineering* **2009**, *35* (8), 1178-1189.
67. Bell, T. H.; El-Din Hassan, S.; Lauron-Moreau, A.; Al-Otaibi, F.; Hijri, M.; Yergeau, E.; St-Arnaud, M., Linkage between bacterial and fungal rhizosphere communities in hydrocarbon-contaminated soils is related to plant phylogeny. *Isme j* **2014**, *8* (2), 331-43.

68. de Cácer, D. A.; Martín, M.; Karlson, U.; Rivilla, R., Changes in bacterial populations and in biphenyl dioxygenase gene diversity in a polychlorinated biphenyl-polluted soil after introduction of willow trees for rhizoremediation. *Appl Environ Microbiol* **2007**, *73* (19), 6224-32.
69. Yergeau, E.; Tremblay, J.; Joly, S.; Labrecque, M.; Maynard, C.; Pitre, F. E.; St-Arnaud, M.; Greer, C. W., Soil contamination alters the willow root and rhizosphere metatranscriptome and the root-rhizosphere interactome. *Isme j* **2018**, *12* (3), 869-884.
70. Pérez-Jaramillo, J. E.; Mendes, R.; Raaijmakers, J. M., Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Mol Biol* **2016**, *90* (6), 635-44.
71. Timmis, K. N.; Pieper, D. H., Bacteria designed for bioremediation. *Trends in biotechnology* **1999**, *17* (5), 201-204.
72. Brown, R. W., *Revegetation of an alpine mine disturbance: Beartooth Plateau, Montana*. US Department of Agriculture, Forest Service, Intermountain Forest & Range ...: 1976; Vol. 206.
73. Desjardins, D.; Nissim, W. G.; Pitre, F. E.; Naud, A.; Labrecque, M., Distribution patterns of spontaneous vegetation and pollution at a former decantation basin in southern Québec, Canada. *Ecological engineering* **2014**, *64*, 385-390.
74. Lee, S. J.; Kong, M.; St-Arnaud, M.; Hijri, M., Arbuscular Mycorrhizal Fungal Communities of Native Plant Species under High Petroleum Hydrocarbon Contamination Highlights Rhizophagus as a Key Tolerant Genus. *Microorganisms* **2020**, *8* (6).
75. Brundrett, M., Mycorrhizas in natural ecosystems. *Advances in ecological research* **1991**, *21*, 171-313.
76. Roy-Bolduc, A.; Hijri, M., The use of mycorrhizae to enhance phosphorus uptake: a way out the phosphorus crisis. *J. Biofertil. Biopestic* **2011**, *2* (104), 1-5.
77. Miransari, M., Hyperaccumulators, arbuscular mycorrhizal fungi and stress of heavy metals. *Biotechnol Adv* **2011**, *29* (6), 645-53.
78. Porcel, R.; Aroca, R.; Ruiz-Lozano, J. M., Salinity stress alleviation using arbuscular mycorrhizal fungi. A review. *Agronomy for Sustainable Development* **2012**, *32* (1), 181-200.
79. Smith, S. E.; Read, D. J., *Mycorrhizal symbiosis*. Academic press: 2008.
80. St-Arnaud, M.; Vujanovic, V., Effects of the Arbuscular Mycorrhizal Symbiosis on Plant Diseases and Pests. In *Mycorrhizae in Crop Production*, Hamel, C., Plenquette, C., Ed. CRC Press: London, 2007; pp 67-123.

81. Hassan, S. E.; Hijri, M.; St-Arnaud, M., Effect of arbuscular mycorrhizal fungi on trace metal uptake by sunflower plants grown on cadmium contaminated soil. *New biotechnology* **2013**, *30* (6), 780-787.
82. Hassan Sel, D.; Bell, T. H.; Stefani, F. O.; Denis, D.; Hijri, M.; St-Arnaud, M., Contrasting the community structure of arbuscular mycorrhizal fungi from hydrocarbon-contaminated and uncontaminated soils following willow (*Salix* spp. L.) planting. *PLoS One* **2014**, *9* (7), e102838.
83. Mc Farlane, C.; Trapp, S., *Plant contamination: modeling and simulation of organic chemical processes*. CRC Press: 1994.
84. Reichenauer, T. G.; Germida, J. J., Phytoremediation of organic contaminants in soil and groundwater. *ChemSusChem* **2008**, *1* (8-9), 708-17.
85. Sandermann, H., Jr., Plant metabolism of xenobiotics. *Trends Biochem Sci* **1992**, *17* (2), 82-4.
86. Singer, A. C., Advances in development of transgenic plants for remediation of xenobiotic pollutants. In *Phytoremediation and Rhizoremediation: Theoretical Background*, Mackova, M., Dowling, D., Macek, T., Ed. Springer: Dordrecht, 2006; pp 5–21.
87. Glick, B. R.; Stearns, J. C., Making phytoremediation work better: maximizing a plant's growth potential in the midst of adversity. *Int J Phytoremediation* **2011**, *13 Suppl 1*, 4-16.
88. Hardoim, P. R.; van Overbeek, L. S.; Elsas, J. D., Properties of bacterial endophytes and their proposed role in plant growth. *Trends Microbiol* **2008**, *16* (10), 463-71.
89. Rughöft, S.; Herrmann, M.; Lazar, C. S.; Cesarz, S.; Levick, S. R.; Trumbore, S. E.; Küsel, K., Community Composition and Abundance of Bacterial, Archaeal and Nitrifying Populations in Savanna Soils on Contrasting Bedrock Material in Kruger National Park, South Africa. *Front Microbiol* **2016**, *7*, 1638.
90. Richardson, A. E.; Barea, J.-M.; McNeill, A. M.; Prigent-Combaret, C., Acquisition of phosphorus and nitrogen in the rhizosphere and plant growth promotion by microorganisms. *Plant and soil* **2009**, *321* (1), 305-339.
91. Schlaeppi, K.; Bulgarelli, D., The plant microbiome at work. *Mol Plant Microbe Interact* **2015**, *28* (3), 212-7.
92. Vessey, J. K., Plant growth promoting rhizobacteria as biofertilizers. *Plant and soil* **2003**, *255* (2), 571-586.

93. Bloemberg, G. V.; Lugtenberg, B. J., Molecular basis of plant growth promotion and biocontrol by rhizobacteria. *Curr Opin Plant Biol* **2001**, *4* (4), 343-50.
94. Haas, D.; Défago, G., Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nat Rev Microbiol* **2005**, *3* (4), 307-19.
95. dos Santos, R. M.; Diaz, P. A. E.; Lobo, L. L. B.; Rigobelo, E. C., Use of Plant Growth-Promoting Rhizobacteria in Maize and Sugarcane: Characteristics and Applications. *Frontiers in Sustainable Food Systems* **2020**, *4* (136).
96. Lugtenberg, B.; Kamilova, F., Plant-growth-promoting rhizobacteria. *Annu Rev Microbiol* **2009**, *63*, 541-56.
97. Stevens, C. J., Nitrogen in the environment. *Science* **2019**, *363* (6427), 578-580.
98. Boddey, R. M.; De Oliveira, O.; Urquiaga, S.; Reis, V.; De Olivares, F.; Baldani, V.; Döbereiner, J., Biological nitrogen fixation associated with sugar cane and rice: contributions and prospects for improvement. In *Management of biological nitrogen fixation for the development of more productive and sustainable agricultural systems*, Springer: 1995; pp 195-209.
99. Dobbelaere, S.; Vanderleyden, J.; Okon, Y., Plant growth-promoting effects of diazotrophs in the rhizosphere. *Critical reviews in plant sciences* **2003**, *22* (2), 107-149.
100. Sessitsch, A.; Howieson, J.; Perret, X.; Antoun, H.; Martinez-Romero, E., Advances in Rhizobium research. *Critical Reviews in Plant Sciences* **2002**, *21* (4), 323-378.
101. Alori, E. T.; Glick, B. R.; Babalola, O. O., Microbial Phosphorus Solubilization and Its Potential for Use in Sustainable Agriculture. *Front Microbiol* **2017**, *8*, 971.
102. Stevenson, F. J.; Cole, M. A., *Cycles of soils: carbon, nitrogen, phosphorus, sulfur, micronutrients*. 2nd Edition ed.; John Wiley & Sons: New York, USA, 1999.
103. Glass, A. D., *Plant mineral nutrition. An introduction to current concepts*. Jones and Bartlett Publishers, Inc.: 1989.
104. Kim, K.; Jordan, D.; McDonald, G., Effect of phosphate-solubilizing bacteria and vesicular-arbuscular mycorrhizae on tomato growth and soil microbial activity. *Biology and fertility of soils* **1997**, *26* (2), 79-87.
105. Rodríguez, H.; Fraga, R., Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnology advances* **1999**, *17* (4-5), 319-339.

106. Sharma, S. B.; Sayyed, R. Z.; Trivedi, M. H.; Gobi, T. A., Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils. *Springerplus* **2013**, *2*, 587.
107. Jorquera, M. A.; Crowley, D. E.; Marschner, P.; Greiner, R.; Fernández, M. T.; Romero, D.; Menezes-Blackburn, D.; De La Luz Mora, M., Identification of  $\beta$ -propeller phytase-encoding genes in culturable Paenibacillus and Bacillus spp. from the rhizosphere of pasture plants on volcanic soils. *FEMS Microbiol Ecol* **2011**, *75* (1), 163-72.
108. Zhang, X.; Zhang, D.; Sun, W.; Wang, T., The Adaptive Mechanism of Plants to Iron Deficiency via Iron Uptake, Transport, and Homeostasis. *Int J Mol Sci* **2019**, *20* (10).
109. Weyens, N.; van der Lelie, D.; Taghavi, S.; Newman, L.; Vangronsveld, J., Exploiting plant-microbe partnerships to improve biomass production and remediation. *Trends Biotechnol* **2009**, *27* (10), 591-8.
110. Crowley, D. E.; Kraemer, S. M., Function of siderophores in the plant rhizosphere. In *The Rhizosphere, Biochemistry and Organic Substances at the Soil-Plant Interface*, CRC Press Boca Raton, FL, USA: 2007; pp 73-109.
111. Rajkumar, M.; Ae, N.; Prasad, M. N.; Freitas, H., Potential of siderophore-producing bacteria for improving heavy metal phytoextraction. *Trends Biotechnol* **2010**, *28* (3), 142-9.
112. Sessitsch, A.; Kuffner, M.; Kidd, P.; Vangronsveld, J.; Wenzel, W. W.; Fallmann, K.; Puschenreiter, M., The role of plant-associated bacteria in the mobilization and phytoextraction of trace elements in contaminated soils. *Soil Biology and Biochemistry* **2013**, *60*, 182-194.
113. Gururani, M. A.; Upadhyaya, C. P.; Baskar, V.; Venkatesh, J.; Nookaraju, A.; Park, S. W., Plant growth-promoting rhizobacteria enhance abiotic stress tolerance in Solanum tuberosum through inducing changes in the expression of ROS-scavenging enzymes and improved photosynthetic performance. *Journal of plant growth regulation* **2013**, *32* (2), 245-258.
114. Sharma, A.; Johri, B.; Sharma, A.; Glick, B., Plant growth-promoting bacterium *Pseudomonas* sp. strain GRP3 influences iron acquisition in mung bean (*Vigna radiata* L. Wilzeck). *Soil Biology and Biochemistry* **2003**, *35* (7), 887-894.

115. Mitter, E. K.; Tosi, M.; Obregón, D.; Dunfield, K. E.; Germida, J. J., Rethinking Crop Nutrition in Times of Modern Microbiology: Innovative Biofertilizer Technologies. *Frontiers in Sustainable Food Systems* **2021**, 5 (29).
116. McSteen, P., Auxin and monocot development. *Cold Spring Harb Perspect Biol* **2010**, 2 (3), a001479.
117. Salisbury, F. B., The role of plant hormones. In *Plant–Environment Interactions*, Wilkinson, R. E., Ed. Marcel Dekker: 1994; pp 39–81.
118. Martínez, C.; Espinosa-Ruiz, A.; Prat, S., Gibberellins and plant vegetative growth. *Annual Plant Reviews online* **2018**, 285-322.
119. Shu, K.; Zhou, W.; Chen, F.; Luo, X.; Yang, W., Abscisic Acid and Gibberellins Antagonistically Mediate Plant Development and Abiotic Stress Responses. *Front Plant Sci* **2018**, 9, 416.
120. Abts, W.; Vandenbussche, B.; De Proft, M. P.; Van de Poel, B., The Role of Auxin-Ethylene Crosstalk in Orchestrating Primary Root Elongation in Sugar Beet. *Front Plant Sci* **2017**, 8, 444.
121. Patten, C. L.; Glick, B. R., Bacterial biosynthesis of indole-3-acetic acid. *Can J Microbiol* **1996**, 42 (3), 207-20.
122. Vacheron, J.; Desbrosses, G.; Bouffaud, M. L.; Touraine, B.; Moënne-Loccoz, Y.; Muller, D.; Legendre, L.; Wisniewski-Dyé, F.; Prigent-Combaret, C., Plant growth-promoting rhizobacteria and root system functioning. *Front Plant Sci* **2013**, 4, 356.
123. Li, J.; Sun, J.; Yang, Y.; Guo, S.; Glick, B. R., Identification of hypoxic-responsive proteins in cucumber roots using a proteomic approach. *Plant Physiol Biochem* **2012**, 51, 74-80.
124. Li, Q.; Saleh-Lakha, S.; Glick, B. R., The effect of native and ACC deaminase-containing *Azospirillum brasilense* Cd1843 on the rooting of carnation cuttings. *Can J Microbiol* **2005**, 51 (6), 511-4.
125. Glick, B. R., Modulation of plant ethylene levels by the bacterial enzyme ACC deaminase. *FEMS Microbiol Lett* **2005**, 251 (1), 1-7.
126. Sandhya, V.; Ali, S. Z.; Grover, M.; Reddy, G.; Venkateswarlu, B., Effect of plant growth promoting *Pseudomonas* spp. on compatible solutes, antioxidant status and plant growth of maize under drought stress. *Plant Growth Regulation* **2010**, 62 (1), 21-30.
127. Ali, S.; Kim, W. C., Plant Growth Promotion Under Water: Decrease of Waterlogging-Induced ACC and Ethylene Levels by ACC Deaminase-Producing Bacteria. *Front Microbiol* **2018**, 9, 1096.

128. Mayak, S.; Tirosh, T.; Glick, B. R., Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiol Biochem* **2004**, *42* (6), 565-72.
129. Belimov, A. A.; Safronova, V. I.; Sergeyeva, T. A.; Egorova, T. N.; Matveyeva, V. A.; Tsyganov, V. E.; Borisov, A. Y.; Tikhonovich, I. A.; Kluge, C.; Preisfeld, A.; Dietz, K. J.; Stepanok, V. V., Characterization of plant growth promoting rhizobacteria isolated from polluted soils and containing 1-aminocyclopropane-1-carboxylate deaminase. *Can J Microbiol* **2001**, *47* (7), 642-52.
130. Backer, R.; Rokem, J. S.; Ilangumaran, G.; Lamont, J.; Praslickova, D.; Ricci, E.; Subramanian, S.; Smith, D. L., Plant Growth-Promoting Rhizobacteria: Context, Mechanisms of Action, and Roadmap to Commercialization of Biostimulants for Sustainable Agriculture. *Front Plant Sci* **2018**, *9*, 1473.
131. Bashan, Y.; de-Bashan, L. E.; Prabhu, S.; Hernandez, J.-P., Advances in plant growth-promoting bacterial inoculant technology: formulations and practical perspectives (1998–2013). *Plant and soil* **2014**, *378* (1), 1-33.
132. Owen, D.; Williams, A. P.; Griffith, G. W.; Withers, P. J., Use of commercial bio-inoculants to increase agricultural production through improved phosphorous acquisition. *Applied Soil Ecology* **2015**, *86*, 41-54.
133. Khan, M. S.; Zaidi, A.; Wani, P. A.; Oves, M., Role of plant growth promoting rhizobacteria in the remediation of metal contaminated soils. *Environmental chemistry letters* **2009**, *7* (1), 1-19.
134. Oleńska, E.; Małek, W.; Wójcik, M.; Świecicka, I.; Thijs, S.; Vangronsveld, J., Beneficial features of plant growth-promoting rhizobacteria for improving plant growth and health in challenging conditions: A methodical review. *Sci Total Environ* **2020**, *743*, 140682.
135. Ma, Y.; Prasad, M. N.; Rajkumar, M.; Freitas, H., Plant growth promoting rhizobacteria and endophytes accelerate phytoremediation of metalliferous soils. *Biotechnol Adv* **2011**, *29* (2), 248-58.
136. Arslan, M.; Afzal, M.; Amin, I.; Iqbal, S.; Khan, Q. M., Nutrients can enhance the abundance and expression of alkane hydroxylase CYP153 gene in the rhizosphere of ryegrass planted in hydrocarbon-polluted soil. *PLoS One* **2014**, *9* (10), e111208.
137. Balseiro-Romero, M.; Gkorezis, P.; Kidd, P. S.; Van Hamme, J.; Weyens, N.; Monterroso, C.; Vangronsveld, J., Use of plant growth promoting bacterial strains to improve *Cytisus striatus* and

- Lupinus luteus development for potential application in phytoremediation. *Sci Total Environ* **2017**, *581-582*, 676-688.
138. Pizarro-Tobías, P.; Niqui, J. L.; Roca, A.; Solano, J.; Fernández, M.; Bastida, F.; García, C.; Ramos, J. L., Field trial on removal of petroleum-hydrocarbon pollutants using a microbial consortium for bioremediation and rhizoremediation. *Environ Microbiol Rep* **2015**, *7* (1), 85-94.
139. Hou, J.; Liu, W.; Wang, B.; Wang, Q.; Luo, Y.; Franks, A. E., PGPR enhanced phytoremediation of petroleum contaminated soil and rhizosphere microbial community response. *Chemosphere* **2015**, *138*, 592-8.
140. Whyte, L. G.; Schultz, A.; Beilen, J. B.; Luz, A. P.; Pellizari, V.; Labbé, D.; Greer, C. W., Prevalence of alkane monooxygenase genes in Arctic and Antarctic hydrocarbon-contaminated and pristine soils. *FEMS Microbiol Ecol* **2002**, *41* (2), 141-50.
141. Yakimov, M. M.; Timmis, K. N.; Golyshin, P. N., Obligate oil-degrading marine bacteria. *Curr Opin Biotechnol* **2007**, *18* (3), 257-66.
142. Afzal, M.; Yousaf, S.; Reichenauer, T. G.; Sessitsch, A., Ecology of alkane-degrading bacteria and their interaction with the plant. *Molecular microbial ecology of the rhizosphere* **2013**, *2*, 975-989.
143. Johnsen, A. R.; Karlson, U., PAH degradation capacity of soil microbial communities--does it depend on PAH exposure? *Microb Ecol* **2005**, *50* (4), 488-95.
144. Tremblay, J.; Yergeau, E.; Fortin, N.; Cobanli, S.; Elias, M.; King, T. L.; Lee, K.; Greer, C. W., Chemical dispersants enhance the activity of oil- and gas condensate-degrading marine bacteria. *Isme j* **2017**, *11* (12), 2793-2808.
145. Xu, X.; Liu, W.; Tian, S.; Wang, W.; Qi, Q.; Jiang, P.; Gao, X.; Li, F.; Li, H.; Yu, H., Petroleum Hydrocarbon-Degrading Bacteria for the Remediation of Oil Pollution Under Aerobic Conditions: A Perspective Analysis. *Front Microbiol* **2018**, *9*, 2885.
146. Varjani, S. J., Microbial degradation of petroleum hydrocarbons. *Bioresour Technol* **2017**, *223*, 277-286.
147. Wang, X. B.; Chi, C. Q.; Nie, Y.; Tang, Y. Q.; Tan, Y.; Wu, G.; Wu, X. L., Degradation of petroleum hydrocarbons (C6-C40) and crude oil by a novel *Dietzia* strain. *Bioresour Technol* **2011**, *102* (17), 7755-61.

148. Sarkar, P.; Roy, A.; Pal, S.; Mohapatra, B.; Kazy, S. K.; Maiti, M. K.; Sar, P., Enrichment and characterization of hydrocarbon-degrading bacteria from petroleum refinery waste as potent bioaugmentation agent for in situ bioremediation. *Bioresour Technol* **2017**, 242, 15-27.
149. Rojo, F., Degradation of alkanes by bacteria. *Environ Microbiol* **2009**, 11 (10), 2477-90.
150. Post-Beittenmiller, D., Biochemistry and molecular biology of wax production in plants. *Annu Rev Plant Physiol Plant Mol Biol* **1996**, 47, 405-430.
151. Schirmer, A.; Rude, M. A.; Li, X.; Popova, E.; del Cardayre, S. B., Microbial biosynthesis of alkanes. *Science* **2010**, 329 (5991), 559-62.
152. Ji, Y.; Mao, G.; Wang, Y.; Bartlam, M., Structural insights into diversity and n-alkane biodegradation mechanisms of alkane hydroxylases. *Front Microbiol* **2013**, 4, 58.
153. Nie, Y.; Chi, C. Q.; Fang, H.; Liang, J. L.; Lu, S. L.; Lai, G. L.; Tang, Y. Q.; Wu, X. L., Diverse alkane hydroxylase genes in microorganisms and environments. *Sci Rep* **2014**, 4, 4968.
154. van Beilen, J. B.; Funhoff, E. G., Alkane hydroxylases involved in microbial alkane degradation. *Appl Microbiol Biotechnol* **2007**, 74 (1), 13-21.
155. Wang, W.; Shao, Z., Enzymes and genes involved in aerobic alkane degradation. *Front Microbiol* **2013**, 4, 116.
156. Bihari, Z.; Szvetnik, A.; Szabó, Z.; Blastyák, A.; Zombori, Z.; Balázs, M.; Kiss, I., Functional analysis of long-chain n-alkane degradation by *Dietzia* spp. *FEMS Microbiol Lett* **2011**, 316 (2), 100-7.
157. Lo Piccolo, L.; De Pasquale, C.; Fodale, R.; Puglia, A. M.; Quatrini, P., Involvement of an alkane hydroxylase system of *Gordonia* sp. strain SoCg in degradation of solid n-alkanes. *Appl Environ Microbiol* **2011**, 77 (4), 1204-13.
158. van Beilen, J. B.; Funhoff, E. G.; van Loon, A.; Just, A.; Kaysser, L.; Bouza, M.; Holtackers, R.; Röthlisberger, M.; Li, Z.; Witholt, B., Cytochrome P450 alkane hydroxylases of the CYP153 family are common in alkane-degrading eubacteria lacking integral membrane alkane hydroxylases. *Appl Environ Microbiol* **2006**, 72 (1), 59-65.
159. Nie, Y.; Liang, J. L.; Fang, H.; Tang, Y. Q.; Wu, X. L., Characterization of a CYP153 alkane hydroxylase gene in a Gram-positive *Dietzia* sp. DQ12-45-1b and its "team role" with alkW1 in alkane degradation. *Appl Microbiol Biotechnol* **2014**, 98 (1), 163-73.

160. Whyte, L. G.; Smits, T. H.; Labb  , D.; Witholt, B.; Greer, C. W.; van Beilen, J. B., Gene cloning and characterization of multiple alkane hydroxylase systems in *Rhodococcus* strains Q15 and NRRL B-16531. *Appl Environ Microbiol* **2002**, *68* (12), 5933-42.
161. Wang, L.; Wang, W.; Lai, Q.; Shao, Z., Gene diversity of CYP153A and AlkB alkane hydroxylases in oil-degrading bacteria isolated from the Atlantic Ocean. *Environ Microbiol* **2010**, *12* (5), 1230-42.
162. Yousaf, S.; Andria, V.; Reichenauer, T. G.; Smalla, K.; Sessitsch, A., Phylogenetic and functional diversity of alkane degrading bacteria associated with Italian ryegrass (*Lolium multiflorum*) and Birdsfoot trefoil (*Lotus corniculatus*) in a petroleum oil-contaminated environment. *J Hazard Mater* **2010**, *184* (1-3), 523-532.
163. Yergeau, E.; Sanschagrin, S.; Beaumier, D.; Greer, C. W., Metagenomic analysis of the bioremediation of diesel-contaminated Canadian high arctic soils. *PLoS One* **2012**, *7* (1), e30058.
164. Long, H.; Wang, Y.; Chang, S.; Liu, G.; Chen, T.; Huo, G.; Zhang, W.; Wu, X.; Tai, X.; Sun, L.; Zhang, B., Diversity of crude oil-degrading bacteria and alkane hydroxylase (alkB) genes from the Qinghai-Tibet Plateau. *Environ Monit Assess* **2017**, *189* (3), 116.
165. Varjani, S. J.; Gnansounou, E.; Pandey, A., Comprehensive review on toxicity of persistent organic pollutants from petroleum refinery waste and their degradation by microorganisms. *Chemosphere* **2017**, *188*, 280-291.
166. Pe  rez-Pantoja, D., Gonza  lez, B., Pieper, D.H. , Aerobic Degradation of Aromatic Hydrocarbons. In *Handbook of Hydrocarbon and Lipid Microbiology*, Timmis, K. N., Ed. Springer-Verlag Berlin Heidelberg: 2010; pp 800-837.
167. Kotoky, R.; Rajkumari, J.; Pandey, P., The rhizosphere microbiome: Significance in rhizoremediation of polycyclic aromatic hydrocarbon contaminated soil. *J Environ Manage* **2018**, *217*, 858-870.
168. Cerniglia, C. E., Biodegradation of polycyclic aromatic hydrocarbons. *Biodegradation* **1992**, *3* (2-3), 351-368.
169. Peng, R. H.; Xiong, A. S.; Xue, Y.; Fu, X. Y.; Gao, F.; Zhao, W.; Tian, Y. S.; Yao, Q. H., Microbial biodegradation of polycyclic aromatic hydrocarbons. *FEMS Microbiol Rev* **2008**, *32* (6), 927-55.

170. Mallick, S.; Chakraborty, J.; Dutta, T. K., Role of oxygenases in guiding diverse metabolic pathways in the bacterial degradation of low-molecular-weight polycyclic aromatic hydrocarbons: a review. *Crit Rev Microbiol* **2011**, *37* (1), 64-90.
171. Ghosal, D.; Ghosh, S.; Dutta, T. K.; Ahn, Y., Current State of Knowledge in Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *Front Microbiol* **2016**, *7*, 1369.
172. Kweon, O., Kim, S-J., Cerniglia, C.E., An Update on the Genomic View of Mycobacterial High-Molecular-Weight Polycyclic Aromatic Hydrocarbon Degradation. In *Aerobic Utilization of Hydrocarbons, Oils and Lipids, Handbook of Hydrocarbon and Lipid Microbiology*, Rojo, F., Ed. Springer International Publishing AG: 2018; pp 1-16.
173. Wick, L. Y.; Ruiz de Munain, A.; Springael, D.; Harms, H.; de, M. A., Responses of *Mycobacterium* sp. LB501T to the low bioavailability of solid anthracene. *Appl Microbiol Biotechnol* **2002**, *58* (3), 378-85.
174. Simon, M. J.; Osslund, T. D.; Saunders, R.; Ensley, B. D.; Suggs, S.; Harcourt, A.; Suen, W. C.; Cruden, D. L.; Gibson, D. T.; Zylstra, G. J., Sequences of genes encoding naphthalene dioxygenase in *Pseudomonas putida* strains G7 and NCIB 9816-4. *Gene* **1993**, *127* (1), 31-7.
175. Laurie, A. D.; Lloyd-Jones, G., The phn genes of *Burkholderia* sp. strain RP007 constitute a divergent gene cluster for polycyclic aromatic hydrocarbon catabolism. *J Bacteriol* **1999**, *181* (2), 531-40.
176. Khan, A. A.; Wang, R. F.; Cao, W. W.; Doerge, D. R.; Wennerstrom, D.; Cerniglia, C. E., Molecular cloning, nucleotide sequence, and expression of genes encoding a polycyclic aromatic ring dioxygenase from *Mycobacterium* sp. strain PYR-1. *Appl Environ Microbiol* **2001**, *67* (8), 3577-85.
177. Pagnout, C.; Frache, G.; Poupin, P.; Maunit, B.; Muller, J. F.; Férand, J. F., Isolation and characterization of a gene cluster involved in PAH degradation in *Mycobacterium* sp. strain SNP11: expression in *Mycobacterium smegmatis* mc(2)155. *Res Microbiol* **2007**, *158* (2), 175-86.
178. Kivisaar, M.; Kasak, L.; Nurk, A., Sequence of the plasmid-encoded catechol 1,2-dioxygenase-expressing gene, pheB, of phenol-degrading *Pseudomonas* sp. strain EST1001. *Gene* **1991**, *98* (1), 15-20.
179. Hiraoka, S.; Yang, C. C.; Iwasaki, W., Metagenomics and Bioinformatics in Microbial Ecology: Current Status and Beyond. *Microbes Environ* **2016**, *31* (3), 204-12.

180. Quiza, L.; St-Arnaud, M.; Yergeau, E., Harnessing phytomicrobiome signaling for rhizosphere microbiome engineering. *Front Plant Sci* **2015**, *6*, 507.
181. Tardif, S.; Yergeau, É.; Tremblay, J.; Legendre, P.; Whyte, L. G.; Greer, C. W., The Willow Microbiome Is Influenced by Soil Petroleum-Hydrocarbon Concentration with Plant Compartment-Specific Effects. *Front Microbiol* **2016**, *7*, 1363.
182. Mitter, E. K.; de Freitas, J. R.; Germida, J. J., Bacterial Root Microbiome of Plants Growing in Oil Sands Reclamation Covers. *Frontiers in Microbiology* **2017**, *8* (849).
183. Ofek-Lalzar, M.; Sela, N.; Goldman-Voronov, M.; Green, S. J.; Hadar, Y.; Minz, D., Niche and host-associated functional signatures of the root surface microbiome. *Nat Commun* **2014**, *5*, 4950.
184. Bell, T. H.; Joly, S.; Pitre, F. E.; Yergeau, E., Increasing phytoremediation efficiency and reliability using novel omics approaches. *Trends Biotechnol* **2014**, *32* (5), 271-80.
185. Pagé, A. P.; Yergeau, É.; Greer, C. W., Salix purpurea Stimulates the Expression of Specific Bacterial Xenobiotic Degradation Genes in a Soil Contaminated with Hydrocarbons. *PLoS One* **2015**, *10* (7), e0132062.
186. Zhao, Q.; Hu, H.; Wang, W.; Peng, H.; Zhang, X., Genome Sequence of *Sphingobium yanoikuyae* B1, a Polycyclic Aromatic Hydrocarbon-Degrading Strain. *Genome Announc* **2015**, *3* (1).
187. Imperato, V.; Portillo-Estrada, M.; McAmmond, B. M.; Douwen, Y.; Van Hamme, J. D.; Gawronski, S. W.; Vangronsveld, J.; Thijs, S., Genomic Diversity of Two Hydrocarbon-Degrading and Plant Growth-Promoting *Pseudomonas* Species Isolated from the Oil Field of Bóbrka (Poland). *Genes (Basel)* **2019**, *10* (6).
188. Neifar, M.; Chouchane, H.; Najjari, A.; El Hidri, D.; Mahjoubi, M.; Ghedira, K.; Naili, F.; Soufi, L.; Raddadi, N.; Sghaier, H.; Ouzari, H. I.; Masmoudi, A. S.; Cherif, A., Genome analysis provides insights into crude oil degradation and biosurfactant production by extremely halotolerant *Halomonas desertis* G11 isolated from Chott El-Djerid salt-lake in Tunisian desert. *Genomics* **2019**, *111* (6), 1802-1814.
189. He, C.; Li, Y.; Huang, C.; Chen, F.; Ma, Y., Genome Sequence and Metabolic Analysis of a Fluoranthene-Degrading Strain *Pseudomonas aeruginosa* DN1. *Front Microbiol* **2018**, *9*, 2595.

- 
190. Mahjoubi, M.; Aliyu, H.; Cappello, S.; Naifer, M.; Souissi, Y.; Cowan, D. A.; Cherif, A., The genome of *Alcaligenes aquatilis* strain BU33N: Insights into hydrocarbon degradation capacity. *PLoS One* **2019**, *14* (9), e0221574.
191. Silva, N. M.; de Oliveira, A.; Pegorin, S.; Giusti, C. E.; Ferrari, V. B.; Barbosa, D.; Martins, L. F.; Morais, C.; Setubal, J. C.; Vasconcellos, S. P.; da Silva, A. M.; de Oliveira, J. C. F.; Pascon, R. C.; Viana-Niero, C., Characterization of novel hydrocarbon-degrading *Gordonia paraffinivorans* and *Gordonia sihwensis* strains isolated from composting. *PLoS One* **2019**, *14* (4), e0215396.
192. Rajkumari, J.; Singha, L. P.; Pandey, P., Draft Genome Sequence of *Klebsiella pneumoniae* AWD5. *Genome Announc* **2017**, *5* (5).
193. Stevens, V.; Thijs, S.; McAmmond, B.; Langill, T.; Van Hamme, J.; Weyens, N.; Vangronsveld, J., Draft Genome Sequence of *Bacillus licheniformis* VSD4, a Diesel Fuel-Degrading and Plant Growth-Promoting Phyllospheric Bacterium. *Genome Announc* **2017**, *5* (11).
194. Filonov, A.; Delegan, Y.; Puntus, I.; Valentovich, L.; Akhremchuk, A.; Evdokimova, O.; Funtikova, T.; Zakhrova, M.; Akhmetov, L.; Vetrova, A.; Titok, M., Complete Genome Sequence of *Pseudomonas putida* BS3701, a Promising Polycyclic Aromatic Hydrocarbon-Degrading Strain for Bioremediation Technologies. *Microbiol Resour Announc* **2020**, *9* (40).