

Article

Genome-wide Investigation and Expression Profiles of Heat Shock Transcription Factor (HSF) Gene Family in Maize (*Zea mays* L.) under Different Growth Stages and Abiotic Stress Conditions

Saqlain Haider ^{1,*}, Shazia Rehman ², Yumna Ahmad ¹, Ali Raza ^{3,*}, Javaria Tabassum ⁴, Talha Javed ^{5,6}, Hany S. Osman ⁷ and Tariq Mahmood ¹

¹ Plant Biochemistry and Molecular Biology laboratory, Department of Plant Sciences, Quaid-i-Azam University, Islamabad 45320, Pakistan; yumna789ahmad@gmail.com (Y.A.); tmahmood.qau@gmail.com (T.M.)

² University Institute of Biochemistry and Biotechnology, PMAS, Arid Agriculture University, Rawalpindi, Pakistan; shaziarehman7@gmail.com

³ Fujian Provincial Key Laboratory of Crop Molecular and Cell Biology, Oil Crops Research Institute, Center of Legume Crop Genetics and Systems Biology/College of Agriculture, Fujian Agriculture and Forestry University (FAFU), Fuzhou, Fujian 350002, China;

⁴ State Key Laboratory of Rice Biology, China National Rice Research Institute, Chinese Academy of Agricultural Science (CAAS), Hangzhou-310000, China; javaria.tabassum@outlook.com

⁵ College of Agriculture, Fujian Agriculture and Forestry University, Fuzhou-350002, China; mta-haj@fafu.edu.cn

⁶ Department of Agronomy, University of Agriculture Faisalabad-38000, Pakistan;

⁷ Department of Agricultural Botany, Faculty of Agriculture, Ain Shams University, Hadeyek Shubra, Cairo 11241, Egypt; hany_osman1@agr.asu.edu.eg

* Correspondence: shaider@bs.qau.edu.pk (S.H.); alirazamughal143@gmail.com (A.R.)

Abstract: Heat shock transcription factors (HSFs) participate in regulating many environmental stress responses and biological processes in plants. Maize (*Zea mays* L.) is a major cash crop that is grown worldwide. However, the growth and yield of maize are affected by several adverse environmental inputs. Therefore, investigating the factors that regulate maize growth and development and resistance to abiotic stress is an essential task for developing stress-resilient maize varieties. Thus, a comprehensive genome-wide identification analysis was performed to identify HSFs in the maize genome. The current study identified 25 *ZmHSFs*, randomly distributed throughout the maize genome. Phylogenetic analysis revealed that *ZmHSFs* are divided into three classes and 13 sub-classes. Gene structure and protein motif analysis supported the results obtained through the phylogenetic analysis. Domain analysis showed the DNA-binding domain to be the most conserved region of *ZmHSFs*. Segmental duplication is shown to be responsible for the expansion of *ZmHSFs*. Most of the *ZmHSFs* are localized inside the nucleus, and the *ZmHSFs* which belong to the same group show similar physio-chemical properties. The 3D structures revealed comparable conserved *ZmHSFs* protein structures. RNA-seq analysis revealed a major role of class A HSFs including, *ZmHSFA-1a* and *ZmHSFA-2a* in all the maize growth stages, i.e., seed, vegetative, and reproductive development. Furthermore, *ZmHSFs* displayed an obvious spatiotemporal expression. Under abiotic stress conditions (heat, drought, cold, UV, and salinity), members of class A and B *ZmHSFs* are induced. Gene ontology (GO) annotation analysis indicated a major role of *ZmHSFs* in resistance to environmental stress and regulation of primary metabolism. Further, the protein-protein interaction analysis showed that *ZmHSFs* interact with several molecular chaperons and major stress-responsive proteins. To summarize, this study provides novel insights for functional studies on the *ZmHSFs* in maize breeding programs.

Keywords: abiotic stress; HSFs; genomics; gene ontology; maize breeding; protein 3D structures

1. Introduction

In recent years, there has been an increasing trend to focus on the responses of plants to abiotic stresses due to global climate change [1]. Particularly, research has been focused on plant heat stress (HS) tolerance mechanisms, as higher temperatures adversely affect all the growth phases of plants [2,3]. Although plants are susceptible to HS throughout their lifespan, reproductive tissues are specifically characterized by vulnerability to HS [4,5]. Heat waves are expected to become more frequent in the future, and atmospheric temperature is on course to rise ~4 °C by the end of this century [6,7]. HS negatively affects plant morphology, physiology, and growth in diverse ways [5,8]. HS alters plasma membrane fluidity, creates proteotoxic stress, causes the overproduction of reactive oxygen species (ROS), and dismantles cellular organization by inducing the collapse of cytoskeleton apparatus [3,8-10]. Nonetheless, plants possess an efficient system that allows them to perceive the environmental stimulus, activate the signaling pathways, and alter their gene expression to ensure survival [9,11,12]. A major step in this process is the activation of stress-inducible genes, the expression of which is controlled by transcription factors (TFs). TFs represent a group of regulatory proteins that control the expression pattern of genes under various developmental and stressful conditions [13,14]. Several TFs families including heat shock transcription factors (HSFs), WRKY (named due to conserved WRKYGQK motif), v-myb avian myeloblastosis viral oncogene homolog (MYB), *Petunia* NAM, *Arabidopsis* ATAF1/2 and CUC2 (NAC), and dehydration responsive-element binding transcriptional activator (DREB), etc., have been shown to positively regulate HS-responsive gene expression and improve plant HS tolerance [14,15]. Among them, the most comprehensively analyzed family is HSFs, the members of which are considered the master regulators of plant HS-response (HSR) and are also involved in regulating plant responses to other abiotic and biotic stress conditions [12,16]. Since the first HSF gene was identified in yeast [17], the HSF gene family has been characterized in several plant species [18,19]. The pioneering study by Nover et al. [20] allowed the identification of the HSF gene family in various plant species, including essential crop plants [19]. The Heatster database (<http://www.cibiv.at/services/hsf>) currently holds 848 HSF sequences from 33 different plant species.

Generally, HSFs are divided into three classes, i.e., A, B, and C, based on phylogenetic analysis and structural properties [21]. All the HSFs contain two highly conserved domains, the DNA-binding domain (DBD), which binds with “heat-shock elements” (5'-nGAAnnTTCn-3') present in regulatory sequences of target genes, through helix-turn-helix (HTH) motif, and oligomerization domain (OD), which has a bipartite heptad repeat pattern of the hydrophobic-associated region (HR-A/B) and is responsible for the trimerization of HSFs [20,22]. Based on the linker length between the HR-A/B region, HSFs are classified into different classes. The linker length comprises 21 amino acid residues in the case of class A and 7 for class C HSFs. On the other hand, HSFs of class B, lack any insertion. The classification of HSFs is also supported by the length of the linker between DBD and OD, 9 to 39 amino-acid for class A, 50 to 78 for class B, and 14 to 49 for class C [20-22]. The HSFs of class A are transcriptional activators, and class B are repressors [20,21]. However, in tomato, the HSF1 possesses both co-activator and repressor functions [23,24]. The class C HSFs are activators like class A [25]. However, they lack activator peptide motif (AHA), and thus cannot initiate transcription on their own [19,20]. The AHA motif is present towards the C-terminal of HSFs and is specific to class A HSFs [26]. In addition to these domains, HSFs also possess nuclear localization signal (NLS), and some contain nuclear export signal (NES) [27]. The NLS and NES are responsible for nuclear import and export of HSFs, an essential step in cellular functioning [27].

The function of HSFs as the master regulators of plant HSR has been demonstrated mainly in *Arabidopsis* and tomato [28,29]. In tomato, the overexpression of *HSFA1a*, improved plant thermotolerance, while co-suppression mutants were susceptible to HS [28]. The mutant plants and their fruits were characterized by extreme sensitivity to HS. *HSFA1a*, *HSFA2*, and *HSFB1* control the fundamental HSF network in tomato [30]. The role of the master regulator is shared by *HSFA1a*, *HSFA1b*, and *HSFA1d* in *Arabidopsis* [29]. In *hsfa1a/b/d* triple mutants, expression of TFs and chaperons was severely hampered

under HS, while the expression of several genes was down-regulated even under normal growth conditions [29]. *HSFA2* is responsible for the extension of acquired thermotolerance (AT) in *Arabidopsis* [31]. In *hsfa2* mutants, the duration of AT was compromised, and the expression of HS-inducible genes was down-regulated. Lämke et al. reported that *HSFA2* promotes the sustained activation of several HS-memory genes through methylation of the target genomic region [32]. The transcripts of *HSFA2* are undetectable under normal conditions. However, after HS, the *HSFA2* becomes the most strongly induced HSF in plants [18,33]. Yoshida et al. reported that the overexpression of *HSFA3* improves plant thermotolerance, while the T-DNA mutants showed reduced thermotolerance [34]. Lin et al. reported that *HSFA2*, *HSFA4a*, and *HSFA7a* are essential for HSR and cytoplasmic protein response. HSFs have also been characterized in several crop plants [35]. It was reported that *OsHSFA4a* and its homolog in wheat *TaHSFA4a*, confers cadmium tolerance to plants [36]. The expression of *TaHSFA2-10* is induced in response to HS, oxidative stress, salicylic acid, and its overexpression improve plant thermotolerance [37]. In addition, HSFs are also involved in the regulation of growth and development in plants [16]. For example, *HSFA9* is expressed specifically during embryogenesis and maturation in the seeds of *Arabidopsis* [38]. Albihlal et al. reported that in *Arabidopsis*, at least 85 development-associated genes are controlled by *HSFA1b* [39]. Authors proposed that the *HSFA1b* allows plants to adjust growth and development under continuously varying environments by transducing external stimuli to stress-associated and development-related genes.

HSF gene family has been characterized in several plant species, including *Arabidopsis thaliana* [20], *Oryza sativa* [40], *Zea mays* [41], *Glycine max* [42], *Populus trichocarpa* [43], *Solanum lycopersicum* [44], *Brachypodium distachyon* [45], and *Triticum aestivum* [46]. However, the role of HSFs in plant growth and development and in responses to stresses other than HS, is not very well understood in maize. Computational biology approaches provide a convenient and reliable platform upon which further wet-lab studies could be carried out. Here, we perform an extensive in-silico analysis of maize HSFs to gain better insights into the functional role of maize HSFs in growth and development and tolerance to multiple abiotic stresses.

2. Materials and Methods

2.1. Sequence Retrieval

The protein sequences of *Zea mays* HSFs were extracted from PLAZA 4.5 (https://bioinformatics.psb.ugent.be/plaza/versions/plaza_v4_5_monocots/) and Ensembl Plants (<https://plants.ensembl.org/index.html>) databases. For this, the BLASTP search was performed using the *Arabidopsis* (AT4G17750, AT2G26150, AT5G03720, AT4G36990, and AT3G24520) and *Zea mays* (GRMZM2G115456, GRMZM2G002131, GRMZM2G086880) HSFs against maize genome Zm-B73-REFERENCE-NAM-5.0, as a query to obtain a putative list of HSFs in the maize genome. These sequences were checked for the presence of DNA binding domain (DBD) and oligomerization domain (OD) through EMBL-EBI employing the hidden Markov model (HMM) (<https://www.ebi.ac.uk/Tools/hmmer/search/phmmer>) and SMART (<http://smart.embl-heidelberg.de/>) tools [47]. The coiled-coil structure, which is a property of OD were predicted using MARCOIL [48]. After carefully analyzing the sequence, the proteins that lacked DBD and/or OD (HR-A/B region) were removed. In addition, the redundant proteins that were the product of a single gene were also discarded from further analysis. Finally, a total of 25 maize HSFs genes were selected for further analysis.

2.2. Sequence Analysis

The coding sequence (CDS) and genomic DNA sequences were retrieved from the maize genome database (MaizeGDB) (<https://www.maizegdb.org/>). The fundamental data such as gene chromosomal location, position, strand position, the total number of transcripts, and intron and exon number was also retrieved from MaizeGDB. The physio-

chemical properties of HSFs were predicted using ExPASy website (<https://www.expasy.org/>). Sub-cellular localization was predicted using two online tools, i.e., WoLFPSORT [49] and CELLO server.2.5 [50]. The conserved protein domains were identified using the MEME suite (<https://meme-suite.org/meme/tools/meme>) [51]. Twenty conserved motifs were identified using default parameters. Using CDS, the intron-exon structure of maize HSF genes was analyzed through GSDS 2.0 (<http://gsds.gao-lab.org/>) [52].

2.3. Sequence Alignment and Phylogenetic Analysis

The protein sequences of *Arabidopsis thaliana*, *Oryza sativa*, *Brachypodium distachyon*, and *Sorghum bicolor* were aligned with *Zea mays* HSFs through Clustal W [53]. An unrooted phylogenetic tree was constructed using the Molecular Evolution Genetic Analysis X (MEGA X) tool [54]. The tree was constructed via the Neighbor-joining method with 1000 bootstrap values.

2.4. Gene Duplication and Evolutionary Analysis

Gene duplication events were investigated by following two parameters: (1) the length of alignable sequence cover > 80% of the longer gene; and (2) the similarity of the aligned regions > 70% [55,56]. To analyze the molecular evolutionary rates of duplicated gene pairs, the non-synonymous substitution (Ka) and synonymous substitution (Ks) ratio were calculated using Ka/Ks calculation tool (<http://services.cbu.uib.no/tools/kaks>). Based on a rate of 6.1×10^{-9} substitutions per site per year, the divergence time (T) was calculated as $T = Ks / (2 \times 6.1 \times 10^{-9}) \times 10^6$ million years ago (Mya) for HSF genes [57]. The duplicated HSF genes were connected using Tbttools [58].

2.5. Chromosomal Distribution and Expression Profiling of HSF Genes

Based on their initial positions on the maize genome, the HSF genes were named, and a chromosomal graph was constructed using Tbttools. The RNA-seq data utilized in the current study was retrieved from maize MaizeGDB (https://qteller.maizegdb.org/genes_by_name_B73v4.php). For this, the comprehensive maize gene expression analysis performed by Stelpflug et al. was downloaded [59]. The present study included the analysis of the growth, tissue, and stress-specific expression of maize HSFs. The transcript per million values was normalized using the log-2 base, and a heat map was constructed using Tbttools.

2.6. Protein 3D Structure, Network Interaction, and Gene Ontology Analysis

The 3-dimensional (3D) structure of maize HSFs were predicted through AlphaFold (<https://www.alphafold.ebi.ac.uk/>) [60]. The protein interaction network analysis was performed using the STRING database (<https://string-db.org/>) [61]. The network interaction file was downloaded and visualized using Cytoscape V. 3.8.2 (<https://cytoscape.org/>) [62]. Gene ontology annotation analysis was performed by uploading the gene IDs of ZmHSFs to the GENE ONTOLOGY RESOURCE (<http://geneontology.org/>) [63].

3. Results


3.1. Identification and Chromosomal Distribution of Maize HSFs

With the availability of the genomic sequences of the number of plant species, including maize, it is now possible to obtain the protein sequences of all the HSF members. In the present study, a total of 25 HSFs were identified from the maize genome (Figure 1; Table 1). All the HSF proteins were surveyed for the presence of DBD and OD through EMBL-EBI, employing HMM. Furthermore, SMART was used to search the HSF-DBD to check the accuracy of the results. After discarding redundant sequences, 25 ZmHSFs were selected for analysis. These HSFs were named based on their chromosomal locations (*ZmHSF-01* to *ZmHSF-25*) (Table S1). The characteristics of maize HSF genes have been presented in Table 1. All the HSFs were mapped on the chromosomes of maize (Figure 1). The maize genome was shown to possess HSF genes on all of its chromosomes, though

the number of HSFs between different chromosomes varied considerably. Chromosome 1 had a maximum of 6 HSFs genes, whereas a single HSF gene copy was localized on each of chromosomes 4, 6, and 10. On the other hand, chromosomes 2, 3, 7, and 9 harbors two gene copies each, while three gene copies were recognized on each of chromosomes 5 and 8. Except for *ZmHSF01*, *ZmHSF02*, *ZmHSF12*, *ZmHSF16*, and *ZmHSF23*, all the other HSF genes were present on the lower arm of the chromosomes.

Figure 1. The chromosomal position of HSF genes in maize. The scale on the left side represents megabases (Mb). The chromosome number has been indicated at the middle of each bar.

Table 1. Characteristics of identified maize HSF genes.



ZmHSF-13	GRMZM2G105348_T01	5	150228697-150230232	-	1	1536	1185	2	1
ZmHSF-14	GRMZM2G179802_T02	5	162047221-162053475	-	3	6261	1920	2	1
ZmHSF-15	GRMZM2G059851_T01	5	168263116-168266561	+	2	3446	1527	2	1
ZmHSF-16	AC206165.3_FGT007	6	159812447-159814105	-	1	1659	1410	2	1
ZmHSF-17	GRMZM2G125969_T01	7	9712577-9716596	-	3	4020	1284	2	1
ZmHSF-18	GRMZM2G139535_T01	7	125896425-125900126	+	2	3702	897	2	1
ZmHSF-19	GRMZM2G165272_T01	7	142270318- 142272208	+	1	1891	1185	2	1
ZmHSF-20	AC205471.4_FGT003	8	124710390-124712153	+	1	1764	1341	2	1
ZmHSF-21	GRMZM2G086880_T01	8	144887947-144889308	+	1	1364	1248	2	1
ZmHSF-22	GRMZM2G118453_T01	8	173979796-173982203	-	1	2407	1870	2	1
ZmHSF-23	GRMZM2G173090_T02	9	59460696-59462007	+	2	1473	1357	2	1
ZmHSF-24	GRMZM2G026742_T01	9	146029209-146032662	-	4	4461	1840	5	4
ZmHSF-25	GRMZM2G301485_T01	10	137124859-137126235	-	2	1376	1300	2	1

3.2. Phylogenetic Analysis and Classification of Maize HSFs

In present study, the evolutionary relationship among AtHSFs, OsHSFs, SbHSFs, BdHSFs, and ZmHSFs were explored. A total of 118 HSFs were divided into three classes and 15 sub-classes based on the phylogenetic tree (Figure 2; Table S2). Variation in HSF gene number was observed among different plant species and between sub-groups (Table 2). For example, *Arabidopsis thaliana* contains 21 HSFs (15 HSFAs, 5 HSFBs, and 1 HSFC), the *Oryza sativa* possess 25 HSFs (13 HSFAs, 8 HSFBs, and 4 HSFCs), *Zea mays* harbors 25 HSFs in its genome (15 HSFAs, 7 HSFBs, and 3 HSFCs), *Brachypodium distachyon* 24 HSFs (14 HSFAs, 7 HSFBs, and 4 HSFCs), while *Sorghum bicolor* contains 23 HSFs (12 HSFAs, 7 HSFBs, and 4 HSFCs) (Table 2). Results indicated that maize HSFs were closed to sorghum HSFs. Similarly, HSFs of rice were closer to *Brachypodium* HSFs, which is in line with the botanical classification among monocots. In contrast to *Arabidopsis*, sub-class A1 contains less HSFs in monocots (Table 2). On the other hand, sub-class A2 appears to have expanded in monocots. There was no ortholog of *Arabidopsis* HSFA9, HSFB3 in monocots that might reflect the loss of these sub-groups in family Poaceae (Figure 2, Table 2). Contrarily, this could also signify the gain of these sub-groups in dicots. A higher number of class C HSFs was observed in monocots. The AtHSFA2, AtHSFC1 did not align with sub-class A2, C1 in the present study, which is in alignment with the results reported by Lin et al. [41].




Figure 2. A neighbor-joining phylogenetic tree was constructed after aligning protein sequences of *Arabidopsis thaliana*, *Oryza sativa*, *Brachypodium distachyon*, *Sorghum bicolor*, and *Zea mays*. Overall, 21 AtHSFs (marron circle), 25 OsHSFs (turquoise rhombus), 23 SbHSFs (purple rectangular), 24 BdHSFs (yellow triangle), and 25 ZmHSFs (red square) were divided into 3 classes and further into 15 sub-classes.

Table 2. Distribution of HSF genes in different sub-classes in selected plant species.

Sub-class	<i>Arabidopsis</i> AtHSFs	<i>Zea mays</i> ZmHSFs	<i>Oryza sativa</i> OsHSFs	<i>Brachypodium distachyon</i> BdHSFs	<i>Sorghum bicolor</i> SbHSFs
A1	4	2	1	1	1
A2	1	2	3	3	3
A3	1	1	1	1	1
A4	2	3	2	2	1
A5	1	1	1	1	1

A6	2	2	2	2	2
A7	2	2	2	2	2
A8	1	2	1	1	1
A9	1	0	0	0	0
B1	1	2	1	1	1
B2	2	4	3	3	3
B3	1	0	0	0	0
B4	1	1	4	3	3
C1	1	2	2	2	2
C2	0	1	2	2	2
Total	21	25	25	24	23

3.3. Gene Duplication Analysis and Evolutionary Rate Calculation

Gene duplications play an important role in evolution as duplications cause genes to generate gene families [64]. In fact, it has been proposed that tandem and segmental duplications have been the primary driving source of evolution as these events lead to expansion of gene families, and generation of proteins with novel functions [65]. Tandem duplication involves the duplication of two or more genes located on the same chromosome, while segmental duplication refers to the phenomenon when genes belonging to the same clade but located on different chromosomes are duplicated [66]. In the present study, a total of 18 (18/25; 72 %) maize HSF genes were shown to be duplicated (Table 3). Further, only one pair of a gene (*ZmHSF-01/Zm-HSF-04*) appeared to be tandemly duplicated, which was recognized on chromosome number 1 (Figure 3). The rest of duplicated genes were all segmentally duplicated, with eight different clusters containing 16 genes. These genes were recognized on chromosomes 1- 9. Moreover, the molecular evolutionary rate of tandemly and segmentally duplicated HSF genes was calculated to gain insights into the selective constraints on the duplicated HSF genes. The ratio of Ka and Ks substitution rate is an effective method to investigate the selective constraint among duplicated gene pairs [67]. Hence, in the present study, Ka, Ks, and Ka/Ks values for each pair of the paralogous gene were calculated (Table 3). In principle, the value of Ka/Ks < 1 signifies the purifying selection (negative selection), Ka/Ks > 1 signifies positive selection, and Ka/Ks = 1 means neutral selection [57]. Here, 18 *ZmHSF* genes were shown to be duplicated. The Ka/Ks ratio for duplicated *ZmHSF* genes ranged from 0.3415 to 0.7572. All the HSF genes in the present study have Ka/Ks value < 1.

The outcome suggests that these genes were under strong purifying selection pressure by natural selection during the course of evolution. Further, the divergence periods of tandemly and segmentally duplicated *ZmHSF* genes were estimated to range from 5.96 to 38.04 with an average of 20.89 million years ago (MYA). Some paralogous gene pairs (*ZmHSF-02/ZmHSF-24*, *ZmHSF-09/ZmHSF21*, and *ZmHSF-16/ZmHSF20*) appeared to be recently duplicated (Table 3). The grass species are estimated to have diverged around 56-73 MYA [68,69]. In the present analysis, all the HSF genes in maize appeared to have duplicated after the divergence of grass species. Further, most of the HSF genes in maize are paralogs, and it can be concluded that duplication events (primarily segmental) played a significant role in the expansion of the HSF gene family in maize.

Table 3. Duplicated gene pairs, non-synonymous substitution rate (Ka), synonymous substitution rate (Ks), non-synonymous to synonymous substitution rate ratio (Ka/Ks), estimated time of duplication event in a million years ago (MYA), and mode of gene duplication.

Duplicated genes	Ka	Ks	Ka/Ks	Estimated time (MYA)	Mode of duplication
ZmHSF-01/ZmHSF-04	0.1837	0.3733	0.4921	30.59	Tandem
ZmHSF-02/ZmHSF-24	0.0248	0.0727	0.3415	5.96	Segmental
ZmHSF-03/ZmHSF-11	0.2497	0.3726	0.6702	30.54	Segmental
ZmHSF-05/ZmHSF-17	0.1810	0.3250	0.5570	26.63	Segmental
ZmHSF-06/ZmHSF-15	0.2772	0.4600	0.6026	37.70	Segmental
ZmHSF-07/ZmHSF-19	0.1538	2.2641	0.5143	21.64	Segmental
ZmHSF-08/ZmHSF-18	0.1814	0.2395	0.7572	19.63	Segmental
ZmHSF-09/ZmHSF-21	0.0389	0.0839	0.4644	6.87	Segmental
ZmHSF-16/ZmHSF20	0.0733	0.1040	0.7049	8.52	Segmental




Figure 3. Circular illustration of the maize genome. The paralogous HSF genes have been connected with grey lines. The purple lines on top of the chromosomal bar represent the position of HSFs.

3.4. Gene Structure and Protein Motif Analysis

To investigate the structural relationship among the HSF genes, the intron-exon organization of all the targeted HSFs was analyzed using GSDS software. The intron-exon structure and number play a key role in the evolution of gene families [70,71]. The gene structure analysis was in line with the phylogenetic relationship among maize HSFs (Figure 4). In general, the intron and exon numbers were shown to be highly consistent. Particularly, 92% (23/25) HSFs contain only one intron except for HSF-02 and HSF-24 (Figure 4). Similarly, HSF-02 and HSF-24 were shown to contain 3 and 5 exons. In contrast, the rest of the HSF genes contained two exons. Further, 17 HSFs contained 5' UTR and 3' UTR. The HSF genes belonging to the same class and sub-class showed a similar intron-exon

pattern in terms of intron number, exon length, intron phase, and overall gene length (Figure 4).

MEME was used to identify the conserved motifs/regions responsible for DNA-binding, oligomerization, nuclear localization, nuclear export, and biological activation of HSFs (Figure 5). In total, 20 motifs designated as motifs 1-20 were identified among maize HSF proteins (Table 4). The highly conserved DBD is represented by motif 1, 2, and 4. Motif 3 corresponds to OD of class A and C HSFs. The HR-A core region of OD is represented motif 5 and is present in all HSFs. OD of class B HSFs is depicted by motif 7. Motif 6 constitutes the NLS of class A HSFs and is present in eight members. The AHA motif is shown by motif 8 and is present in 11 class A HSFs. The NES of class A HSFs is represented by motif 16. Motif 15 constitutes the NLS of Class B2 HSFs. Certain HSFs also possess class-specific conserved motifs i.e., ZmHSF-02 and ZmHSF-24 contain motif 10 and 13, HSFs of subclass B1 harbor motif 14, sub-class C1 HSFs motif 19, the sub-class B2 HSFs motif 11, and sub-class A4 members (ZmHSF-16, ZmHSF-20) contain motif 18. Motif 9 and 12 are present in the same members of class A HSFs. Four members of class A HSFs contain motif 17. Motif18 is found in two members of class A HSFs. Finally, motif 20 is found in members of sub-class A2 and A6 HSFs.




Figure 4. The intron-exon structure of maize HSF genes. A black line represents the introns. Exons by a yellow rectangle and 5' UTR and 3' UTR by pink wedges. The gene structure of *Zea mays* HSFs was in alignment with the phylogenetic relationship.




Figure 5. Conserved motifs in maize HSFs as identified by MEME. The motifs were identified by using full-length protein sequences of ZmHSFs. Name of HSFs and p-values have been indicated on the left side. Each motif is represented by a unique color, as indicated at the bottom.

Table 4. Conserved motif sequence of *Zea mays* HSFs.

Motif	Consensus sequence
1	LPKYFKHNNFSSFVRQLNTYGFRKVPDRWEFANEGFLRGQKHLLKNIHR
2	PFLTKTYEMVDDPATDAVVSWSGAAGNSFV
3	LLAELVRLRQZQSTSEQLQALERRLQGMEQRQQQMMAFLA
4	VWBPAEFARDL
5	GLEEEIERLKRDKAL
6	MQNPDFLRQLVQQQEKSKELEDAINKKRR
7	LARELAQMRKLCNNILLMSKYADTQQPD
8	VNDDFWEZFLTE
9	DGGPVDDSEAAGGGGQIIKYQPIPEAAKQPLPKNLAFDSS
10	MPMDVEMASNNVGTDFSTGNDFTDTSALCEWDDMDIFGGELEHILQQPEQ
11	QSWPIYRPRPVYHPLRACNG
12	KPSQDGPSPQPPVKTAPGPENIEIGKY
13	TIEDYGYDRPWLEQDCQMEAQQNCKNPQY
14	KSVKLFVLLKDAARKRGRCEEAAASERPIKMIRIGEPWIGVPSSGPGRC
15	RLFGVSIGRKMRD
16	DNVBQLTEQMGYLSSANH
17	RKPIHSHSPQTQ
18	VDMCSDTTTGDTSQDETTSETGGSHGPAK
19	CCISMGGEDHR
20	PRPMEGLHDVGPP

3.5. Domain Analysis and Physio-chemical Properties

The modular structure and the functional domains of HSFs have been studied and described extensively [22]. The HSF-type DBD was highly conserved and consisted of approximately 100 amino acids (Figure 6). The locations of DBD and OD were predicted using SMART and MARCOIL (Table 5). The DBD of most maize HSFs was located at the beginning of the N-terminal. Few exceptions were ZmHSF-03, ZmHSF-10, ZmHSF-14, and ZmHSF-15. As expected, the linker length between DBD and OD of HSBs was larger than HSFAs and HSFs.

The physio-chemical properties of HSFs such as amino-acid length, Mw, and pI were investigated using Expasy (Table 5). In addition, the amino acid composition of each group was analyzed using the online tool CoPId (Table S3). The amino acid length of class A HSF ranged from 350 (ZmHSF-23) to 528 (ZmHSF-14), for class B, 298 (ZmHSF-08) to 414 (ZmHSF-03) and in class C, the amino acid length ranged from 257 (ZmHSF-13) to 348 (ZmHSF-21). The pI of class A HSFs ranged from 4.70 (ZmHSF-17) to 8.87 (ZmHSF-10). For class B it varied from 5.00 (ZmHSF-19) to 9.53 (ZmHSF-18), and for class C, pI ranged from 5.85 (ZmHSF-13) to 8.09 (ZmHSF-21). For class A, the Mw varied from 38.154 (ZmHSF-23) to 58.138 (ZmHSF-14), for class B it ranged from 33.258 (ZmHSF-18) to 44.381 (ZmHSF-03). While for class C HSFs, Mw ranged from 27.836 (ZmHSF-13) to 37.409 (ZmHSF-21). In general, the pI of class A HSFs was in acidic ranges except for ZmHSF-10, which was shown to be in a slightly basic range. The pI of class B was in slightly acidic and basic ranges. Finally, for class C the pI was in similar ranges as class B HSFs. The average amino acid composition of ZmHSFs ranged from 1.1 (cysteine) to 10.0 (alanine) (Figure 7). The average amino acid composition of class A HSFs ranged from 0.8 (cysteine) to 8.7 (alanine). In contrast, the average amino acid composition of class B ranged from 1.3 (tryptophan) to 12.6 (alanine). Finally, the class C amino acid composition ranged from 1.1 (tryptophan) to 11.2 (alanine).




Figure 6. Multiple sequence alignment of maize HSFs. The highly conserved DBD consists of four antiparallel β -stranded sheets ($\beta 1$, $\beta 2$, $\beta 3$, $\beta 4$) and a bundle of three α -helices ($\alpha 1$, $\alpha 2$, $\alpha 3$) which form the secondary structure. Rectangular boxes represent α -helices while square boxes represent β -stranded sheets.

Table 5. The physio-chemical properties of maize HSFs proteins including group, protein ID, domains, amino acid physio-chemical properties, and sub-cellular localization.

Gene	Group	Protein ID	DBD	OD	Amino acid	pI	Molecular weight	Localization CELLO	Localization WoLFPSORT
ZmHSF-01	A-6b	GRMZM2G165972_P02	44-133	164-184	384	5.30	43268.46	Nucleus	Nucleus
ZmHSF-02	A-8a	GRMZM2G118485_P02	51-140	170-190	417	5.09	46816.21	Nucleus	Nucleus
ZmHSF-03	B-2d	GRMZM2G164909_P01	72-161	231-260	414	7.14	44381.82	Nucleus	Nucleus
ZmHSF-04	A-6a	GRMZM2G010871_P01	46-135	165-206	357	5.03	40550.58	Nucleus	Nucleus
ZmHSF-05	A-2a	GRMZM2G132971_P01	40-129	149-204	359	5.57	40587.23	Nucleus	Nucleus
ZmHSF-06	A-1a	GRMZM2G115456_P01	59-148	176-203	527	5.11	56724.50	Nucleus	Nucleus
ZmHSF-07	B-4a	GRMZM2G088242_P01	22-112	216-243	394	7.81	41742.83	Nucleus	Nucleus
ZmHSF-08	B-1a	GRMZM2G002131_P01	18-107	165-192	298	9.13	32270.31	Nucleus	Nucleus
ZmHSF-09	C-1c	GRMZM2G089525_P01	27-116	168-188	331	5.94	35883.69	Nucleus	Nucleus
ZmHSF-10	A-7a	GRMZM2G005815_P01	140-231	260-280	462	8.87	50970.67	Chloroplast	Nucleus
ZmHSF-11	B-2b	GRMZM2G098696_P01	37-126	196-216	370	5.89	39562.03	Nucleus	Nucleus
ZmHSF-12	A-3	GRMZM2G384339_P01	32-121	149-176	497	5.06	54104.58	Nucleus	Nucleus
ZmHSF-13	C-2a	GRMZM2G105348_P01	15-104	135-171	257	5.85	27836.98	Nucleus	Nucleus
ZmHSF-14	A-5	GRMZM2G179802_P02	69-159	177-225	528	5.57	58138.65	Nucleus	Nucleus
ZmHSF-15	A-1b	GRMZM2G059851_P01	76-165	190-231	508	4.96	56061.66	Nucleus	Nucleus
ZmHSF-16	A-4b	AC206165.3_FGP007	25-114	135-155	469	5.41	51623.79	Nucleus	Nucleus
ZmHSF-17	A-2b	GRMZM2G125969_P01	52-141	169-210	375	4.70	42043.73	Nucleus	Nucleus
ZmHSF-18	B-1b	GRMZM2G139535_P01	20-109	169-189	298	9.53	32258.35	Nucleus	Nucleus
ZmHSF-19	B-2c	GRMZM2G165272_P01	45-134	207-234	394	5.00	41468.10	Nucleus	Nucleus
ZmHSF-20	A-4d	AC205471.4_FGP003	13-102	123-143	446	5.15	49718.46	Nucleus	Nucleus
ZmHSF-21	C-1a	GRMZM2G086880_P01	35-124	173-193	348	8.09	37409.50	Nucleus	Nucleus
ZmHSF-22	A-4a	GRMZM2G118453_P01	10-99	120-175	433	5.30	48829.92	Nucleus	Nucleus
ZmHSF-23	A-7b	GRMZM2G173090_P02	51-140	164-205	350	4.95	38154.66	Cytoplasm	Cytoplasm
ZmHSF-24	A-8b	GRMZM2G026742_P01	51-140	170-190	407	4.97	45317.63	Nucleus	Nucleus
ZmHSF-25	B-2a	GRMZM2G301485_P01	51-140	170-190	318	5.89	39562.03	Nucleus	Nucleus




Figure 7. The amino acid composition of maize HSFs.

3.6. Proteins Structure and Sub-cellular Localization of Maize HSFs

The protein structures of maize HSFs were predicted through Alphafold. The predicted models were downloaded to view their 3D structure (Figure 8). The highly conserved DBD is represented by α -helices and β -sheets. The OD can be seen to be linked with DBD through linker residues. Most of maize HSFs were predicted to be localized inside the nucleus (Table 5). Exceptions were ZmHSF-10 and ZmHSF-23 (HSFA7a and HSFA7b). This indicates that class 7 HSFs might possess distinct properties.





Figure 8. Protein structure of maize HSFs. The prediction confidence level is presented at the bottom. For detailed information about proteins, see Table 5.

3.7. Expression Profiles of *Zea mays* HSFs during Different Developmental Stages

The expression patterns of *Zea mays* HSF genes were investigated during different growth stages and across different time scales. Although gene expression pattern is not always directly related to protein abundance, the transcriptome profiles can still provide

insights into the probable role of genes in particular biological processes [72]. The transcriptome data utilized in the present study was downloaded from the maize genome database [59]. We analyzed HSF profiles in 3 different growth stages (seed, vegetative, and reproductive) and across 20 different tissues (embryo, endosperm, whole seed, primary root, tap root, whole root, stem and shoot apical meristem, immature leaves, tip of stage 2 leaf, mature leaf tissue, pooled leaves, topmost leaves, vegetative meristem and surrounding tissues, immature tassel, meiotic tassel, anthers, mature pollen, mature female spikelet, pre-pollination cob, immature cob, and silks) (Table S4). Furthermore, the expression patterns were investigated at different timescales in a particular developmental stage to get an overview of the spatiotemporal expression of HSFs.

The *ZmHSF-04*, *ZmHSF-05*, and *ZmHSF-06* are shown to be the most highly induced HSFs across all the tissues during different growth stages (Figure 9 A,B,C). During seed developmental stages, *ZmHSF-02*, *ZmHSF-03*, *ZmHSF-04*, *ZmHSF-07*, *ZmHSF-08*, *ZmHSF-09*, *ZmHSF-13*, *ZmHSF-15*, *ZmHSF-19*, *ZmHSF-20*, *ZmHSF-23*, and *ZmHSF-25* are up-regulated (Figure 9A). 8 HSFs showed very little or no expression at all. Interestingly, the transcripts of all *ZmHSFs* completely disappear 12 days after pollination. During vegetative stage, the *ZmHSF-01*, *ZmHSF-02*, *ZmHSF-09*, *ZmHSF-13*, *ZmHSF-14*, *ZmHSF-15*, *ZmHSF-19*, *ZmHSF-23*, and *ZmHSF-25* are up-regulated (Figure 9B). This indicate these genes might play putative regulate role in these tissues and control vegetative growth. *ZmHSF-22* and *ZmHSF-24* showed no expression during vegetative stage. The highest expression of *ZmHSFs* was observed in root tissues. Similarly, at reproductive stage across different tissues, *ZmHSF-02*, *ZmHSF-04*, *ZmHSF-13*, *ZmHSF-15*, *ZmHSF-19*, and *ZmHSF-25* are up-regulated (Figure 9C). Interestingly, the transcripts of most HSFs except for *ZmHSF-02*, *ZmHSF-04*, *ZmHSF-05*, *ZmHSF-06*, *ZmHSF-16*, and *ZmHSF-25* completely disappear in pollen tissues. *ZmHSF-17* showed almost no expression in any tissue during reproductive stage. Most of the HSFs were highly expressed in silk tissues.






Figure 9. Expression profiles of maize HSFs in different seed (A), vegetative (B), and reproductive (C) tissues based on transcriptome data. The heat map was constructed using Tbttools. The color bar at the top right represents the normalized TPM values (log-transformed fold change values). Green represents higher, red lower, and yellow, medium transcript values.

3.8. Expression Pattern of *Zea mays* HSFs under Abiotic Stresses

Maize growth, development, and yield is adversely affected by several abiotic stresses [73]. Therefore, to examine maize HSFs expression under different abiotic stress events, RNA-seq data was analyzed and a heat map was constructed (Figure 10; Table S5). In response to HS, members of class A and B HSFs showed the highest expression. Interestingly, under the drought stress, only the transcript of *ZmHSF-05* was moderately over-expressed. Under cold stress, 3 members of class A HSFs (*ZmHSF-05*, *ZmHSF-06*, and *ZmHSF-16*) showed relatively higher expression. Moderate expression of *ZmHSF-05*, *ZmHSF-06*, *ZmHSF-04*, and *ZmHSF-19* was observed under UV stress. Under salinity

stress, *ZmHSF-05* and *ZmHSF-14* showed the highest expression. Under abiotic stress conditions, *ZmHSF-01*, *ZmHSF-02*, *ZmHSF-05*, *ZmHSF-06*, *ZmHSF-07*, *ZmHSF-09*, *ZmHSF-12*, *ZmHSF-13*, *ZmHSF-14*, *ZmHSF-15*, *ZmHSF-19*, *ZmHSF-23*, and *ZmHSF-25* are up-regulated (Figure 10). However, some HSFs were only induced by a particular stress. For example, higher transcripts of *ZmHSF-24* are only detected after HS treatment. Similarly, *ZmHSF-16* is slightly overexpressed after cold treatment. While some HSFs showed almost no expression under any stress condition. These include *ZmHSF-03*, *ZmHSF-08*, *ZmHSF-10*, *ZmHSF-11*, *ZmHSF-17*, *ZmHSF-18*, *ZmHSF-20*, *ZmHSF-21*, and *ZmHSF-22* (Figure 10).




Figure 10. Expression profiles of maize HSFs under different abiotic stress conditions. Color bar at the top right represents the normalized TPM values (log-transformed fold change values). Green represents higher, red lower, and yellow, medium transcript values.

3.9. Functional Annotation of Maize HSFs

HSFs have been reported to play a major role not only under stressful conditions but also in plant growth and development [16,72]. Therefore, the regulatory functions of maize HSFs were predicted through GO annotation investigation based on the biological process (BP), molecular function (MF), and cellular component (CC) classes (Figure 11; Table S6). The BP annotation analysis indicated that maize HSFs are mainly involved in cellular response to heat (GO:0034605), response to heat (GO:0009408), response to temperature stimulus (GO:0009266), Regulation of transcription by RNA polymerase II (GO:0006357), response to abiotic stimulus (GO:0009628), cellular response to stress (GO:0033554), regulation of RNA biosynthetic process (GO:2001141), etc. (Figure 11). With regards to MF annotation analysis, it was revealed that maize HSFs are mostly involved in RNA polymerase II cis-regulatory region sequence-specific DNA binding (GO:0000978), cis-regulatory region sequence-specific DNA binding (GO:0000987), RNA polymerase II transcription regulatory region sequence-specific DNA binding (GO:0000977), transcription regulatory region nucleic acid binding (GO:0001067), transcription cis-regulatory region binding (GO:0000976), sequence-specific double-stranded DNA binding (GO:1990837), double-stranded DNA binding (GO:0003690), etc. (Figure 11). The CC annotation study showed that ZmHSFs are majorly involves in the nucleus (GO:0005634), intracellular membrane-bounded organelle (GO:0043231), membrane-bounded organelle (GO:0043227), intracellular organelle (GO:0043229), organelle

(GO:0043226), etc. (Figure 11). To conclude, the GO annotation study confirms the role of maize HSFs in regulating abiotic stresses and plant metabolism.




Figure 11. Gene ontology enrichment analysis. The figure depicts the predicted role of HSFs in biological processes, molecular functions, and cellular components.

3.10. Protein-Protein Interaction Network Analysis

The protein network interaction analysis can help understand the biological functions and mechanisms of protein function [74]. Since both the RNA-seq data and GO annotation analysis suggested the role of HSFs in stress conditions and normal growth, we performed network analysis to predict the interacting partners of ZmHSFs (Table S7). The results showed that maize HSFs interact not only with themselves but also with a range of proteins with well know functions in cellular growth and stress responses (Figure 12). For example, HSFs were shown to interact with molecular chaperons HSP101, HSP82 (belongs to HSP90 family), HSBP-2, and DnaJ-like protein (belongs to the HSP40 family). It was reported that HSP101 and HSA32 interact with each other and promote acquired ther-motolerance in *Arabidopsis* [75]. The HSP82 was reported to be induced by higher temper-atures. A higher concentration of HSP82 is required for normal cellular growth in yeast at higher temperatures [76]. Gu et al. reported that maize HSBP-2 and HSFA2 interact with each other and modulate raffinose biosynthesis [77]. HSFA2 was shown to bind to the

promoter sequence of HSBP-2 and activate its expression. Higher raffinose synthesis improved HS tolerance of *Arabidopsis thaliana*. The DnaJ-like proteins are molecular co-chaperones that interact with HSP70s and control protein homeostasis [78]. DnaJ proteins have been reported to play a critical role in plant growth, development, and HS tolerance [78-80]. ZmHSFs also interact with two major proteins, i.e., multi-protein bridging factor 1c (MBF1c) and DREB2A. Both these proteins have been shown to accumulate under diverse abiotic stress conditions. DREB2A is a major protein, and its overexpression improves plant HS, drought stress, cold stress, etc., tolerance [81]. MBF1c is a transcriptional co-activator that modulates the expression of DREB2A, some HSFs, and phytohormones [3]. Interestingly, MBF1c is necessary for basal thermotolerance but not for acquired thermotolerance [82]. In addition, MBF1c is also shown to be required for plant developmental responses [83].

Maize HSFs are also predicted to interact with SUMO proteins. SUMOylation is a post-translational phenomenon where SUMO proteins are covalently attached and detached to target proteins [84]. This process affects several biological processes inside the cell, including transcriptional regulation of gene expression, apoptosis, programmed cell death, cellular response to stress, stability of proteins, etc. [84]. Rytz et al. reported that SIZ1, a SUMO protein, targets multiple TFs, chromatin remodelers, transcriptional co-activators/repressors connected to abiotic and biotic stress responses [85]. This suggests maize HSFs may also be SUMOylated under diverse biological conditions and stress responses. To conclude, PPI analysis was in alignment with the RNA-seq and GO annotation analysis which indicated that HSFs of *Zea mays* play an important role not only in abiotic stress conditions but also in maize growth and metabolism.




Figure 12. Protein-Protein network of maize HSFs. The line connecting two proteins represents that an interaction exists between them.

4. Discussion

Maize (*Zea mays*) is a major cereal crop that is widely cultivated worldwide for food, feed, fiber, and fuel. Maize is also considered a model plant for basic and applied research in plant science [86]. Unraveling the factors regulating the growth and stress resistance would contribute significantly to the development of climate-smart, stress-resilient maize cultivars with higher agricultural productivity. The sequencing of maize genome (B73 inbred line) in 2009 opened the plethora of opportunities to identify, analyze, and characterize stress-associated genes in maize [87]. To provide food security in the scenario of

climate change and ever-growing world population, it is imperative to understand the molecular mechanisms behind plant stress resistance and explore genetic resources associated with the higher crop yield [3,15]. HSFs have been identified in several plant species, including important crops. The *Arabidopsis thaliana*, *Oryza sativa*, *Zea mays*, *Glycine max*, *Populus trichocarpa*, *Solanum lycopersicum*, *Brachypodium distachyon*, *Sorghum bicolor*, and *Triticum aestivum* contain 21, 25, 25, 38, 28, 26, 24, 23, and 61 HSFs in their genomes [20,40-45,74,88]. Following the sequencing of several plants, it is found out that the number of HSFs may be independent of the genome size [74]. For example, *Arabidopsis thaliana* (135 Mb) contains 21 HSFs [20,89], while *Medicago truncatula* (375 Mb) harbors 15 HSFs [43,90]. Similarly, 25 HSFs are found in *Oryza sativa* (430 Mb) [40,91] and an equal number of HSFs are also present in *Zea mays* (2.4 Gb) [41,87]. Even though the HSF gene family was previously characterized by Lin et al. [41], our work differs from theirs in multiple aspects. Their research was mostly restricted to the identification and classification of ZmHSFs. On the other hand, this comprehensive study particularly focused on the evolutionary analysis, expression profiling, GO, and PPI networks to explore the probable regulatory role played by ZmHSFs under benign and stress conditions.

The distribution of the HSF gene family in maize was analyzed by constructing a chromosomal map (Figure 1). The fact that all the chromosomes harbor at least one HSF gene suggests that the most recent common ancestor of *Zea mays* has HSF genes distributed widely in its genome. Phylogenetic analysis indicated maize HSFs are divided into three classes and further into 13 sub-classes which is consistent with the HSF class number observed in other monocots. For example, the HSFs of *Oryza sativa*, *Brachypodium distachyon*, *Sorghum bicolor*, and *Triticum aestivum* are also divided into three classes and 13 sub-classes [40,45,46,88]. Despite that, differences among HSF numbers were observed in different sub-classes between monocots (Figure 2, Table 2). For example, compared to rice, *Brachypodium*, and *Sorghum*, the sub-family B4, A2, and C2 contain less HSF members in maize. On the other hand, the sub-class A1, A4, A8, B1, and B2 is expanded in *Zea mays* (Figure 2, Table 2). Gene duplications generate new genes and provide novel possibilities for evolutionary success [92]. In the present analysis, 9 pair of ZmHSFs were shown to be paralogs (Table 3, Figure 3). The results indicate that segmental duplication events have played a major role in the expansion of the HSF gene family in maize. An increase in gene regulatory repertoire such as transcriptional regulators, developmental regulators, signal transducers, etc., is a prerequisite for the evolution of complex systems in different organisms [22,93]. Since the gene duplication events result doubling of a single gene which cannot account for such large expansions, it has been suggested that whole-genome duplication (WGD) events have been instrumental in expanding the regulatory repertoire of plants [92]. It is assumed that the *Arabidopsis* genome went under two rounds of WGD in the past 60-70 million years [94,95] and that more than 90% increase in regulatory genes have been caused by duplication events in *Arabidopsis* in the past 150 million years [94]. This suggests that an increase in the HSF gene members in plants accounts for WGDs. Additionally, segmental duplications occur in gene families, which evolve at a slower rate [65]. It is thought that the increase or decrease in exon number play an important role in the evolution of a gene family [96]. Therefore, we investigated the number and distribution of introns and exons in ZmHSF genes. Our results showed that the ZmHSF genes contain 2 exons and 1 intron except for *ZmHSF-02* and *ZmHSF-24* (Figure 4; Table 1). Moreover, the length and position of exons and introns were well conserved in the same sub-classes but varied considerably between different sub-classes.

The previous investigations showed that HSFs play an important role in plant growth [39]. Therefore, we investigated the tissue-specific expression of ZmHSFs in twenty different developmental tissues using RNA-seq data (Figure 9A, B, C). Several genes showed an enhanced expression that reflects their role under various developmental stages. In particular, *ZmHSF-05* (A-2a) and *ZmHSF-06* (A-1a) were highly expressed almost across all the growth phases. The *hsfa1abde* quadruple mutants displayed abnormal phenotype and growth retardation, implying HSF1s is also involved in developmental processes [29]. Interestingly, *HSFA2* could rescue the developmental defects of

hsfa1abde quadruple mutants [97]. This further supports the result obtained from our analysis and provides a strong base for further wet-lab studies to characterize the function of *ZmHSF-05* and *ZmHSF-06* in plant growth and development. Similarly, HSFs have been reported to play a key role in plant acclimation to abiotic stress conditions. Kumar et al. reported that *TaHSFA6e* modulates tolerance of wheat to HS and drought stress during pollination and grain filling stages [98]. Yokotani et al. reported that *OsHSFA2e* improves Arabidopsis tolerance to HS and salinity stress by activating the expression of HSPs [99]. Thus, the expression patterns of ZmHSFs were evaluated under abiotic stress conditions. Most of ZmHSFs displayed stress-specific expression with some HSFs showing up-regulation only under particular stress events (Figure 10). Jiang et al. reported that *ZmHSF-04* improves plant tolerance to HS, salinity stress and increases the sensitivity to abscisic acid [100]. Similarly, *ZmHSF-12* overexpression improves plant basal thermotolerance and AT [101]. These results are in line with our analysis which showed a higher transcript accumulation of this TFs under respective stress conditions (Figure 10).

The PPI analysis indicated that maize HSFs interact with molecular chaperones and stress-associated proteins (Figure 12). Molecular chaperones are present inside the cells and are constitutively expressed under normal conditions or are induced under specific developmental stages or stress conditions [102]. These chaperons perform various functions under physiological conditions inside the cells, such as signaling, folding and stabilization, translocation, and degradation of proteins [11,102]. Under harsh environmental conditions, molecular chaperones act as powerful buffers to limit protein misfolding/unfolding and prevent protein aggregate formation that might be otherwise toxic to plant cells [25]. Here, the ZmHSFs was shown to interact with chaperons belonging to different families (HSP101, HSP90, HSP40) and genes with a well-known role in thermotolerance (HSA32, HSP82, HSBP-2). DREB2A is a major transcriptional activator that functions downstream of HSFA1s dependent transcriptional cascade in Arabidopsis thaliana [3,12,15,29]. Similarly, MBF1c is a major protein that is characterized by its role in the regulation of abiotic stress responses and growth in plants [3,82,83]. SUMO proteins are attached to their target proteins and modify their biological activities under various physiological and stress conditions [84]. In Arabidopsis, SUMOylation has been proposed as one of the molecular mechanisms that are responsible for the activation of HSFA1s [12]. Many HSFs in Arabidopsis such as HSFA1d, HSFA2, and HSFB2B have the potential to be SUMOylated [103]. In tomato, the knockout of SIZI (a SUMO ligase) reduces plant thermotolerance [104]. Here, ZmHSFs was shown to interact with all these proteins (DREB2A, MBF1c, and SUMO proteins), which further confirms their role in the regulation of abiotic stress response.

Taken together, our present analysis provides strong support for the positive role of HSFs in the growth and development of maize by the regulation of primary metabolism. Furthermore, HSFs of maize interact with the major stress-responsive proteins and confer abiotic stress resistance.

5. Conclusion

Here, we identified a total of 25 HSFs from the maize genome through genome-wide investigation analysis. To gain a better understanding of phylogenetic analysis, gene structure and conserved protein motif analysis, gene duplication and evolutionary analysis, domain analysis and physio-chemical properties, protein 3-D structure, GO and PPI network analysis, and expression profiles of ZmHSFs under various developmental stages and stress conditions were studied. The results indicate ZmHSFs play a major role in plant growth and stress responses. Therefore, additional studies are recommended to confirm the proposed role of HSFs in maize growth, development, and response to numerous environmental cues.

Supplementary Materials: The following supplementary material is available online. Table S1: Protein sequences of *Zea mays* HSFs; Table S2: The protein sequences of *Arabidopsis thaliana*, *Oryza sativa*, *Brachypodium distachyon*, and *Sorghum bicolor* HSFs; Table S3: The class specific amino acid composition of *Zea mays* HSFs; Table S4: TPM values of *Zea mays* HSFs under different growth stages across

various tissues; Table S5: TPM values of *Zea mays* HSFs under multiple abiotic stress conditions; Table S6: GO annotation file of *Zea mays* HSFs; Table S7: Protein-protein interaction network file of *Zea mays* HSFs.

Author Contributions: Conceptualization, S.H. and T.M.; methodology, S.H.; software, S.H.; formal analysis, S.H., S.R., Y.A.; resources, A.R.; writing—original draft preparation, S.H., S.R., Y.A., A.R., J.T., and T.J.; writing—review and editing, A.R., H.S.O.; supervision, T.M.; funding acquisition, A.R., and H.S.O. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Nguyen, H.C.; Lin, K.H.; Ho, S.L.; Chiang, C.M.; Yang, C.M. Enhancing the abiotic stress tolerance of plants: from chemical treatment to biotechnological approaches. *Physiol. Plant.* **2018**, *164*, 452–466.
2. Lippmann, R.; Babben, S.; Menger, A.; Delker, C.; Quint, M. Development of wild and cultivated plants under global warming conditions. *Curr. Biol.* **2019**, *29*, R1326–R1338.
3. Haider, S.; Iqbal, J.; Naseer, S.; Yaseen, T.; Shaukat, M.; Bibi, H.; Ahmad, Y.; Daud, H.; Abbasi, N.L.; Mahmood, T. Molecular mechanisms of plant tolerance to heat stress: current landscape and future perspectives. *Plant Cell Rep.* **2021**, 1–25.
4. Chaturvedi, P.; Wiese, A.J.; Ghatak, A.; Závěská Drábková, L.; Weckwerth, W.; Honys, D. Heat stress response mechanisms in pollen development. *New Phytol.* **2021**.
5. Raza, A. Metabolomics: a systems biology approach for enhancing heat stress tolerance in plants. *Plant Cell Rep.* **2020**, 1–23.
6. Stocker, T. *Climate change 2013: the physical science basis: Working Group I contribution to the Fifth assessment report of the Intergovernmental Panel on Climate Change*; Cambridge university press: 2014.
7. Russo, S.; Marchese, A.F.; Sillmann, J.; Immé, G. When will unusual heat waves become normal in a warming Africa? *Environ. Res. Lett.* **2016**, *11*, 054016.
8. Hasanuzzaman, M.; Nahar, K.; Alam, M.; Roychowdhury, R.; Fujita, M. Physiological, biochemical, and molecular mechanisms of heat stress tolerance in plants. *Int. J. Mol. Sci.* **2013**, *14*, 9643–9684.
9. Hayes, S.; Schachtschabel, J.; Mishkind, M.; Munnik, T.; Arisz, S.A. Hot topic: Thermosensing in plants. *Plant, Cell Environ.* **2021**.
10. Zahra, N.; Shaukat, K.; Hafeez, M.B.; Raza, A.; Hussain, S.; Chaudhary, M.T.; Akram, M.Z.; Kakavand, S.N.; Saddiq, M.S.; Wahid, A. Physiological and Molecular Responses to High, Chilling, and Freezing Temperature in Plant Growth and Production: Consequences and Mitigation Possibilities. *Harsh Environment and Plant Resilience: Molecular and Functional Aspects* **2021**, 235.
11. Mittler, R.; Finka, A.; Goloubinoff, P. How do plants feel the heat? *Trends Biochem. Sci.* **2012**, *37*, 118–125.
12. Ohama, N.; Sato, H.; Shinozaki, K.; Yamaguchi-Shinozaki, K. Transcriptional regulatory network of plant heat stress response. *Trends Plant Sci.* **2017**, *22*, 53–65.
13. Jin, J.; Tian, F.; Yang, D.-C.; Meng, Y.-Q.; Kong, L.; Luo, J.; Gao, G. PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Res.* **2016**, gkw982.
14. Raza, A.; Tabassum, J.; Kudapa, H.; Varshney, R.K. Can omics deliver temperature resilient ready-to-grow crops? *Crit. Rev. Biotechnol.* **2021**, 1–24.
15. Haider, S.; Iqbal, J.; Naseer, S.; Shaukat, M.; Abbasi, B.A.; Yaseen, T.; Zahra, S.A.; Mahmood, T. Unfolding molecular switches in plant heat stress resistance: A comprehensive review. *Plant Cell Rep.* **2021**, 1–24.
16. András, N.; Pettkó-Szandtner, A.; Szabados, L. Diversity of plant heat shock factors: regulation, interactions, and functions. *J. Exp. Bot.* **2021**, *72*, 1558–1575.
17. Sorger, P.K.; Pelham, H.R. Yeast heat shock factor is an essential DNA-binding protein that exhibits temperature-dependent phosphorylation. *Cell* **1988**, *54*, 855–864.
18. Scharf, K.-D.; Heider, H.; Hoffeld, I.; Lyck, R.; Schmidt, E.; Nover, L. The tomato Hsf system: HsfA2 needs interaction with HsfA1 for efficient nuclear import and may be localized in cytoplasmic heat stress granules. *Mol. Cell. Biol.* **1998**, *18*, 2240–2251.
19. Guo, M.; Liu, J.-H.; Ma, X.; Luo, D.-X.; Gong, Z.-H.; Lu, M.-H. The plant heat stress transcription factors (HSFs): structure, regulation, and function in response to abiotic stresses. *Front. Plant Sci.* **2016**, *7*, 114.
20. Nover, L.; Bharti, K.; Döring, P.; Mishra, S.K.; Ganguli, A.; Scharf, K.-D. Arabidopsis and the heat stress transcription factor world: how many heat stress transcription factors do we need? *Cell stress & chaperones* **2001**, *6*, 177.
21. von Koskull-Döring, P.; Scharf, K.-D.; Nover, L. The diversity of plant heat stress transcription factors. *Trends in plant science* **2007**, *12*, 452–457.
22. Scharf, K.-D.; Berberich, T.; Ebersberger, I.; Nover, L. The plant heat stress transcription factor (Hsf) family: structure, function and evolution. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms* **2012**, *1819*, 104–119.

23. Bharti, K.; von Koskull-Doering, P.; Bharti, S.; Kumar, P.; Tintschl-Koerberitz, A.; Treuter, E.; Nover, L. Tomato heat stress transcription factor HsfB1 represents a novel type of general transcription coactivator with a histone-like motif interacting with the plant CREB binding protein ortholog HAC1. *The Plant Cell* **2004**, *16*, 1521-1535.
24. Fragkostefanakis, S.; Simm, S.; El-Shershaby, A.; Hu, Y.; Bublak, D.; Mesihovic, A.; Darm, K.; Mishra, S.K.; Tschiersch, B.; Theres, K. The repressor and co-activator HsfB1 regulates the major heat stress transcription factors in tomato. *Plant, cell & environment* **2019**, *42*, 874-890.
25. Jacob, P.; Hirt, H.; Bendahmane, A. The heat-shock protein/chaperone network and multiple stress resistance. *Plant biotechnology journal* **2017**, *15*, 405-414.
26. Czarnecka-Verner, E.; Yuan, C.-X.; Scharf, K.-D.; Englich, G.; Gurley, W.B. Plants contain a novel multi-member class of heat shock factors without transcriptional activator potential. *Plant molecular biology* **2000**, *43*, 459-471.
27. Kotak, S.; Larkindale, J.; Lee, U.; von Koskull-Döring, P.; Vierling, E.; Scharf, K.-D. Complexity of the heat stress response in plants. *Current opinion in plant biology* **2007**, *10*, 310-316.
28. Mishra, S.K.; Tripp, J.; Winkelhaus, S.; Tschiersch, B.; Theres, K.; Nover, L.; Scharf, K.-D. In the complex family of heat stress transcription factors, HsfA1 has a unique role as master regulator of thermotolerance in tomato. *Genes & development* **2002**, *16*, 1555-1567.
29. Yoshida, T.; Ohama, N.; Nakajima, J.; Kidokoro, S.; Mizoi, J.; Nakashima, K.; Maruyama, K.; Kim, J.-M.; Seki, M.; Todaka, D. Arabidopsis HsfA1 transcription factors function as the main positive regulators in heat shock-responsive gene expression. *Molecular Genetics and Genomics* **2011**, *286*, 321-332.
30. Hahn, A.; Bublak, D.; Schleiff, E.; Scharf, K.-D. Crosstalk between Hsp90 and Hsp70 chaperones and heat stress transcription factors in tomato. *The Plant Cell* **2011**, *23*, 741-755.
31. Chang, Y.-y.; Liu, H.-c.; Liu, N.-y.; Chi, W.-t.; Wang, C.-n.; Chang, S.-h.; Wang, T.-t. A heat-inducible transcription factor, HsfA2, is required for extension of acquired thermotolerance in Arabidopsis. *Plant physiology* **2007**, *143*, 251-262.
32. Lämke, J.; Brzezinka, K.; Altmann, S.; Bäurle, I. A hit-and-run heat shock factor governs sustained histone methylation and transcriptional stress memory. *The EMBO journal* **2016**, *35*, 162-175.
33. Nishizawa, A.; Yabuta, Y.; Yoshida, E.; Maruta, T.; Yoshimura, K.; Shigeoka, S. Arabidopsis heat shock transcription factor A2 as a key regulator in response to several types of environmental stress. *The Plant Journal* **2006**, *48*, 535-547.
34. Yoshida, T.; Sakuma, Y.; Todaka, D.; Maruyama, K.; Qin, F.; Mizoi, J.; Kidokoro, S.; Fujita, Y.; Shinozaki, K.; Yamaguchi-Shinozaki, K. Functional analysis of an Arabidopsis heat-shock transcription factor HsfA3 in the transcriptional cascade downstream of the DREB2A stress-regulatory system. *Biochemical and biophysical research communications* **2008**, *368*, 515-521.
35. Lin, K.-F.; Tsai, M.-Y.; Lu, C.-A.; Wu, S.-J.; Yeh, C.-H. The roles of Arabidopsis HSFA2, HSFA4a, and HSFA7a in the heat shock response and cytosolic protein response. *Botanical studies* **2018**, *59*, 1-9.
36. Shim, D.; Hwang, J.-U.; Lee, J.; Lee, S.; Choi, Y.; An, G.; Martinoia, E.; Lee, Y. Orthologs of the class A4 heat shock transcription factor HsfA4a confer cadmium tolerance in wheat and rice. *The Plant Cell* **2009**, *21*, 4031-4043.
37. Guo, X.-l.; Yuan, S.-n.; Zhang, H.-n.; Zhang, Y.-y.; Zhang, Y.-j.; Wang, G.-y.; Li, Y.-q.; Li, G.-l. Heat-response patterns of the heat shock transcription factor family in advanced development stages of wheat (*Triticum aestivum* L.) and thermotolerance-regulation by TaHsfA2-10. *BMC plant biology* **2020**, *20*, 1-18.
38. Kotak, S.; Vierling, E.; Baumbach, H.; Koskull-Doering, P.v. A novel transcriptional cascade regulating expression of heat stress proteins during seed development of Arabidopsis. *The Plant Cell* **2007**, *19*, 182-195.
39. Albihlal, W.S.; Obomighie, I.; Blein, T.; Persad, R.; Chernukhin, I.; Crespi, M.; Bechtold, U.; Mullineaux, P.M. Arabidopsis HEAT SHOCK TRANSCRIPTION FACTOR A1b regulates multiple developmental genes under benign and stress conditions. *Journal of experimental botany* **2018**, *69*, 2847-2862, doi:10.1093/jxb/ery142.
40. Guo, J.; Wu, J.; Ji, Q.; Wang, C.; Luo, L.; Yuan, Y.; Wang, Y.; Wang, J. Genome-wide analysis of heat shock transcription factor families in rice and Arabidopsis. *Journal of genetics and genomics* **2008**, *35*, 105-118.
41. Lin, Y.-X.; Jiang, H.-Y.; Chu, Z.-X.; Tang, X.-L.; Zhu, S.-W.; Cheng, B.-J. Genome-wide identification, classification and analysis of heat shock transcription factor family in maize. *BMC genomics* **2011**, *12*, 1-14.
42. Chung, E.; Kim, K.-M.; Lee, J.-H. Genome-wide analysis and molecular characterization of heat shock transcription factor family in Glycine max. *Journal of genetics and genomics* **2013**, *40*, 127-135.
43. Wang, F.; Dong, Q.; Jiang, H.; Zhu, S.; Chen, B.; Xiang, Y. Genome-wide analysis of the heat shock transcription factors in *Populus trichocarpa* and *Medicago truncatula*. *Molecular biology reports* **2012**, *39*, 1877-1886.
44. Yang, X.; Zhu, W.; Zhang, H.; Liu, N.; Tian, S. Heat shock factors in tomatoes: genome-wide identification, phylogenetic analysis and expression profiling under development and heat stress. *PeerJ* **2016**, *4*, e1961.
45. Wen, F.; Wu, X.; Li, T.; Jia, M.; Liu, X.; Li, P.; Zhou, X.; Ji, X.; Yue, X. Genome-wide survey of heat shock factors and heat shock protein 70s and their regulatory network under abiotic stresses in *Brachypodium distachyon*. *PloS one* **2017**, *12*, e0180352.
46. Duan, S.; Liu, B.; Zhang, Y.; Li, G.; Guo, X. Genome-wide identification and abiotic stress-responsive pattern of heat shock transcription factor family in *Triticum aestivum* L. *BMC genomics* **2019**, *20*, 1-20.
47. Letunic, I.; Copley, R.R.; Schmidt, S.; Ciccarelli, F.D.; Doerks, T.; Schultz, J.; Ponting, C.P.; Bork, P. SMART 4.0: towards genomic data integration. *Nucleic acids research* **2004**, *32*, D142-D144.

48. Delorenzi, M.; Speed, T. An HMM model for coiled-coil domains and a comparison with PSSM-based predictions. *Bioinformatics* **2002**, *18*, 617-625.
49. Horton, P.; Park, K.-J.; Obayashi, T.; Fujita, N.; Harada, H.; Adams-Collier, C.; Nakai, K. WoLF PSORT: protein localization predictor. *Nucleic acids research* **2007**, *35*, W585-W587.
50. Yu, C.S.; Chen, Y.C.; Lu, C.H.; Hwang, J.K. Prediction of protein subcellular localization. *Proteins: Structure, Function, and Bioinformatics* **2006**, *64*, 643-651.
51. Bailey, T.L.; Boden, M.; Buske, F.A.; Frith, M.; Grant, C.E.; Clementi, L.; Ren, J.; Li, W.W.; Noble, W.S. MEME SUITE: tools for motif discovery and searching. *Nucleic acids research* **2009**, *37*, W202-W208.
52. Hu, B.; Jin, J.; Guo, A.-Y.; Zhang, H.; Luo, J.; Gao, G. GSDS 2.0: an upgraded gene feature visualization server. *Bioinformatics* **2015**, *31*, 1296-1297.
53. Thompson, J.D.; Higgins, D.G.; Gibson, T.J. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic acids research* **1994**, *22*, 4673-4680.
54. Kumar, S.; Stecher, G.; Li, M.; Knyaz, C.; Tamura, K. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Molecular biology and evolution* **2018**, *35*, 1547-1549.
55. Gu, Z.; Cavalcanti, A.; Chen, F.-C.; Bouman, P.; Li, W.-H. Extent of gene duplication in the genomes of *Drosophila*, nematode, and yeast. *Molecular biology and evolution* **2002**, *19*, 256-262.
56. Yang, S.; Zhang, X.; Yue, J.-X.; Tian, D.; Chen, J.-Q. Recent duplications dominate NBS-encoding gene expansion in two woody species. *Molecular Genetics and Genomics* **2008**, *280*, 187-198.
57. Lynch, M.; Conery, J.S. The evolutionary fate and consequences of duplicate genes. *science* **2000**, *290*, 1151-1155.
58. Chen, C.; Chen, H.; He, Y.; Xia, R. TBtools, a toolkit for biologists integrating various biological data handling tools with a user-friendly interface. *BioRxiv* **2018**, 289660.
59. Stelplflug, S.C.; Sekhon, R.S.; Vaillancourt, B.; Hirsch, C.N.; Buell, C.R.; de Leon, N.; Kaeppler, S.M. An expanded maize gene expression atlas based on RNA sequencing and its use to explore root development. *The plant genome* **2015**, *9*.
60. Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Žídek, A.; Potapenko, A. Highly accurate protein structure prediction with AlphaFold. *Nature* **2021**, 1-11.
61. Szklarczyk, D.; Gable, A.L.; Nastou, K.C.; Lyon, D.; Kirsch, R.; Pyysalo, S.; Doncheva, N.T.; Legeay, M.; Fang, T.; Bork, P. The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic acids research* **2021**, *49*, D605-D612.
62. Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N.S.; Wang, J.T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Research* **2003**, *13*, 2498-2504.
63. Consortium, G.O. The gene ontology resource: 20 years and still GOing strong. *Nucleic acids research* **2019**, *47*, D330-D338.
64. Jiang, M.; Chu, Z. Comparative analysis of plant MKK gene family reveals novel expansion mechanism of the members and sheds new light on functional conservation. *Bmc Genomics* **2018**, *19*, 1-18.
65. Cannon, S.B.; Mitra, A.; Baumgarten, A.; Young, N.D.; May, G. The roles of segmental and tandem gene duplication in the evolution of large gene families in *Arabidopsis thaliana*. *BMC plant biology* **2004**, *4*, 1-21.
66. Liu, Y.; Jiang, H.; Chen, W.; Qian, Y.; Ma, Q.; Cheng, B.; Zhu, S. Genome-wide analysis of the auxin response factor (ARF) gene family in maize (*Zea mays*). *Plant Growth Regulation* **2011**, *63*, 225-234.
67. Rehman, S.; Jørgensen, B.; Aziz, E.; Batool, R.; Naseer, S.; Rasmussen, S.K. Genome wide identification and comparative analysis of the serpin gene family in brachypodium and barley. *Plants* **2020**, *9*, 1439.
68. Gaut, B.S. Evolutionary dynamics of grass genomes. *New phytologist* **2002**, *154*, 15-28.
69. Initiative, I.B. Genome sequencing and analysis of the model grass *Brachypodium distachyon*. *Nature (London)* **2010**, *463*, 763-768.
70. Song, J.; Zeng, L.; Chen, R.; Wang, Y.; Zhou, Y. In silico identification and expression analysis of superoxide dismutase (SOD) gene family in *Medicago truncatula*. *3 Biotech* **2018**, *8*, 1-12.
71. Xu, X.; Yang, Y.; Liu, C.; Sun, Y.; Zhang, T.; Hou, M.; Huang, S.; Yuan, H. The evolutionary history of the sucrose synthase gene family in higher plants. *BMC plant biology* **2019**, *19*, 1-14.
72. Fragkostefanakis, S.; Röth, S.; Schleiff, E.; SCHARF, K.D. Prospects of engineering thermotolerance in crops through modulation of heat stress transcription factor and heat shock protein networks. *Plant, cell & environment* **2015**, *38*, 1881-1895.
73. De Cuyper, C.; Struk, S.; Braem, L.; Gevaert, K.; De Jaeger, G.; Goormachtig, S. Strigolactones, karrikins and beyond. *Plant, cell & environment* **2017**, *40*, 1691-1703.
74. Ye, J.; Yang, X.; Hu, G.; Liu, Q.; Li, W.; Zhang, L.; Song, X. Genome-wide investigation of heat shock transcription factor family in wheat (*Triticum aestivum* L.) and possible roles in anther development. *International journal of molecular sciences* **2020**, *21*, 608.
75. Wu, T.-y.; Juan, Y.-t.; Hsu, Y.-h.; Wu, S.-h.; Liao, H.-t.; Fung, R.W.; Charng, Y.-y. Interplay between heat shock proteins HSP101 and HSA32 prolongs heat acclimation memory posttranscriptionally in *Arabidopsis*. *Plant Physiology* **2013**, *161*, 2075-2084.
76. Borkovich, K.; Farrelly, F.; Finkelstein, D.; Taulien, J.; Lindquist, S. hsp82 is an essential protein that is required in higher concentrations for growth of cells at higher temperatures. *Molecular and cellular biology* **1989**, *9*, 3919-3930.

77. Gu, L.; Jiang, T.; Zhang, C.; Li, X.; Wang, C.; Zhang, Y.; Li, T.; Dirk, L.M.; Downie, A.B.; Zhao, T. Maize HSFA 2 and HSBP 2 antagonistically modulate raffinose biosynthesis and heat tolerance in Arabidopsis. *The Plant Journal* **2019**, *100*, 128-142.
78. Pulido, P.; Leister, D. Novel DNAJ-related proteins in Arabidopsis thaliana. *New Phytologist* **2018**, *217*, 480-490.
79. Kong, F.; Deng, Y.; Wang, G.; Wang, J.; Liang, X.; Meng, Q. LeCDJ1, a chloroplast DnaJ protein, facilitates heat tolerance in transgenic tomatoes. *Journal of integrative plant biology* **2014**, *56*, 63-74.
80. Kong, F.; Deng, Y.; Zhou, B.; Wang, G.; Wang, Y.; Meng, Q. A chloroplast-targeted DnaJ protein contributes to maintenance of photosystem II under chilling stress. *Journal of experimental botany* **2014**, *65*, 143-158.
81. Mizoi, J.; Shinozaki, K.; Yamaguchi-Shinozaki, K. AP2/ERF family transcription factors in plant abiotic stress responses. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms* **2012**, *1819*, 86-96.
82. Suzuki, N.; Bajad, S.; Shuman, J.; Shulaev, V.; Mittler, R. The transcriptional co-activator MBF1c is a key regulator of thermotolerance in Arabidopsis thaliana. *Journal of Biological Chemistry* **2008**, *283*, 9269-9275.
83. Jaimes-Miranda, F.; Chávez Montes, R.A. The plant MBF1 protein family: a bridge between stress and transcription. *Journal of experimental botany* **2020**, *71*, 1782-1791.
84. Hay, R.T. SUMO: a history of modification. *Molecular cell* **2005**, *18*, 1-12.
85. Rytz, T.C.; Miller, M.J.; McLoughlin, F.; Augustine, R.C.; Marshall, R.S.; Juan, Y.-t.; Charng, Y.-y.; Scalf, M.; Smith, L.M.; Vierstra, R.D. SUMOylome profiling reveals a diverse array of nuclear targets modified by the SUMO ligase SIZ1 during heat stress. *The Plant Cell* **2018**, *30*, 1077-1099.
86. Strable, J.; Scanlon, M.J. Maize (Zea mays): a model organism for basic and applied research in plant biology. *Cold spring harbor protocols* **2009**, *2009*, pdb. emo132.
87. Schnable, P.S.; Ware, D.; Fulton, R.S.; Stein, J.C.; Wei, F.; Pasternak, S.; Liang, C.; Zhang, J.; Fulton, L.; Graves, T.A. The B73 maize genome: complexity, diversity, and dynamics. *science* **2009**, *326*, 1112-1115.
88. Nagaraju, M.; Sudhakar Reddy, P.; Anil Kumar, S.; K. Srivastava, R.; Kavi Kishor, P.; Rao, D.M. Genome-wide scanning and characterization of Sorghum bicolor L. heat shock transcription factors. *Current genomics* **2015**, *16*, 279-291.
89. Kaul, S.; Koo, H.L.; Jenkins, J.; Rizzo, M.; Rooney, T.; Tallon, L.J.; Feldblyum, T.; Nierman, W.; Benito, M.-I.; Lin, X. Analysis of the genome sequence of the flowering plant Arabidopsis thaliana. *nature* **2000**, *408*, 796-815.
90. Young, N.D.; Debellé, F.; Oldroyd, G.E.; Geurts, R.; Cannon, S.B.; Udvardi, M.K.; Benedito, V.A.; Mayer, K.F.; Gouzy, J.; Schoof, H. The Medicago genome provides insight into the evolution of rhizobial symbioses. *Nature* **2011**, *480*, 520-524.
91. Yu, J.; Hu, S.; Wang, J.; Wong, G.K.-S.; Li, S.; Liu, B.; Deng, Y.; Dai, L.; Zhou, Y.; Zhang, X. A draft sequence of the rice genome (Oryza sativa L. ssp. indica). *science* **2002**, *296*, 79-92.
92. Van de Peer, Y.; Maere, S.; Meyer, A. The evolutionary significance of ancient genome duplications. *Nature Reviews Genetics* **2009**, *10*, 725-732.
93. Solé, R.V.; Fernandez, P.; Kauffman, S.A. Adaptive walks in a gene network model of morphogenesis: insights into the Cambrian explosion. *arXiv preprint q-bio/0311013* **2003**.
94. Maere, S.; De Bodt, S.; Raes, J.; Casneuf, T.; Van Montagu, M.; Kuiper, M.; Van de Peer, Y. Modeling gene and genome duplications in eukaryotes. *Proceedings of the National Academy of Sciences* **2005**, *102*, 5454-5459.
95. Jiao, Y.; Wickett, N.J.; Ayyampalayam, S.; Chanderbali, A.S.; Landherr, L.; Ralph, P.E.; Tomsho, L.P.; Hu, Y.; Liang, H.; Soltis, P.S. Ancestral polyploidy in seed plants and angiosperms. *Nature* **2011**, *473*, 97-100.
96. Rose, A. Intron-mediated regulation of gene expression. *Nuclear pre-mRNA processing in plants* **2008**, 277-290.
97. Liu, H.-c.; Charng, Y.-y. Common and distinct functions of Arabidopsis class A1 and A2 heat shock factors in diverse abiotic stress responses and development. *Plant physiology* **2013**, *163*, 276-290.
98. Kumar, R.R.; Goswami, S.; Singh, K.; Dubey, K.; Rai, G.K.; Singh, B.; Singh, S.; Grover, M.; Mishra, D.; Kumar, S. Characterization of novel heat-responsive transcription factor (TaHSFA6e) gene involved in regulation of heat shock proteins (HSPs)—A key member of heat stress-tolerance network of wheat. *Journal of biotechnology* **2018**, *279*, 1-12.
99. Yokotani, N.; Ichikawa, T.; Kondou, Y.; Matsui, M.; Hirochika, H.; Iwabuchi, M.; Oda, K. Expression of rice heat stress transcription factor OsHsfA2e enhances tolerance to environmental stresses in transgenic Arabidopsis. *Planta* **2008**, *227*, 957-967.
100. Jiang, Y.; Zheng, Q.; Chen, L.; Liang, Y.; Wu, J. Ectopic overexpression of maize heat shock transcription factor gene ZmHsf04 confers increased thermo and salt-stress tolerance in transgenic Arabidopsis. *Acta Physiologiae Plantarum* **2018**, *40*, 1-12.
101. Li, G.; Zhang, Y.; Zhang, H.; Zhang, Y.; Zhao, L.; Liu, Z.; Guo, X. Characteristics and regulating role in thermotolerance of the heat shock transcription factor ZmHsf12 from Zea mays L. *Journal of Plant Biology* **2019**, *62*, 329-341.
102. Banti, V.; Mafessoni, F.; Loret, E.; Alpi, A.; Perata, P. The heat-inducible transcription factor HsfA2 enhances anoxia tolerance in Arabidopsis. *Plant Physiology* **2010**, *152*, 1471-1483.
103. Miller, M.J.; Barrett-Wilt, G.A.; Hua, Z.; Vierstra, R.D. Proteomic analyses identify a diverse array of nuclear processes affected by small ubiquitin-like modifier conjugation in Arabidopsis. *Proceedings of the National Academy of Sciences* **2010**, *107*, 16512-16517.
104. Zhang, S.; Wang, S.; Lv, J.; Liu, Z.; Wang, Y.; Ma, N.; Meng, Q. SUMO E3 ligase SIZ1 facilitates heat tolerance in tomato. *Plant and Cell Physiology* **2018**, *59*, 58-71.