

Article

Prevalence of Health Problems Targeted by the National School-based Screening Program among Primary School Students in Saudi Arabia, 2019

Manal Matar Al Daajani^{1*}, Dina Mohammed Al-Habib², Mona Hamed Ibrahim^{2,3}, Nora Abdulrhman Al Shewair², Yahya Mohammad Faghih², Abrar Abdulazeem Alzaher², Amjad Fawzi Alfaleh^{2,4}, Khaled Ibrahim Alabdulkareem^{5,6}

1 Public Health Department, Health Affairs, Jeddah, Kingdom of Saudi Arabia (KSA).

2 General Administration of School Health, Ministry of Health, Riyadh, (KSA).

3 Department of Community, Environmental and Occupational Medicine, Faculty of Medicine, Zagazig University, Egypt.

4 General Directorate of Medical Consultation, Ministry of Health, Riyadh, (KSA).

5 Assistant Prof. of family medicine. College of Medicine, Al Imam Mohammad Ibn Saud Islamic University (IMSIU), (KSA).

6 Assistant Deputy Minister for Primary Health Care, Assisting Deputyship for Primary Health Care, MOH, (KSA).

* Correspondence: manal-m-daajani@hotmail.com; Tel.: 00966553200352

Abstract: The Ministry of Health (MOH) in the Kingdom of Saudi Arabia (KSA) established a National School-based Screening Program (NSBSP) for health screening of school-going children. Students from specific grades were systematically screened for several health problems including obesity, visual and auditory problems, dental cavities, scoliosis, and attention-deficit/hyperactivity disorder (ADHD). This cross-sectional study aimed to determine the prevalence of these health problems among primary school students based on secondary data obtained from the NSBSP. We included 444,259 screened school children from the 1st and 4th grades of 50% of the selected schools (both private and public) across the KSA during the academic year 2018-2019. Among them, the most prevalent health problems identified were dental cavities (38.7%), eye refractory errors (10.9%), and overweight and obesity (10.5%); the less prevalent problems included ADHD (2.81%), auditory problems (0.6%), and scoliosis (0.48%). A greater prevalence of most health problems was observed in girls than boys. The NSBSP successfully aided the detection of health conditions with high and low prevalence among primary school students in the KSA, and thus, identification of health problems of specific concern. Implementation of effective school health services for the prevention, early detection, diagnosis, and treatment of these health problems is imperative.

Keywords: Health Status; National Survey; KSA; School Health; Students

1. Introduction

Schools play a very important role in childhood development, education, and psychological and social health, which are fundamental for the quality of life and future of students. Evidence confirms that healthier students are better learners [1]. Health barriers to learning, such as vision and hearing deficits, and dental pain exert adverse effects on gaining information process, attendance of the student, academic pattern, and development [2]. Barriers to learning are believed to negatively affect the future financial stability and social life of students. Nearly 80% of learning processes occur through visual tasks, such as writing, reading, and using computers. Untreated visual problems affect students' ability to read [3], while uncorrected auditory problems are associated with higher rates of dysfunction in speech, language, poor performance on educational tasks, social and emotional difficulties, and lower self-esteem [4]. Dental pain and dental cavities lead to school absences, sleep disturbances, and unaccomplished homework [5]. Several health problems are prevalent in school-age children. The aim of health screening programs is to

increase the early detection of health problems and promote prevention of accompanying physical, mental, and social consequences.

The Saudi Ministry of Health (MOH) established the National School-based Screening Program (NSBSP) in 2018-201. The program targeted first and fourth grade students of primary schools, first grade students of intermediate schools, and first grade students of secondary schools. This program targeted a set of health problems, including weight-related health problems, visual and auditory problems, dental cavities, smoking, scoliosis, depression, and attention-deficit/hyperactivity disorder (ADHD). One of the objectives of the program was to create a database for recording health problems and to assess the efficacy of appropriate health interventions to prevent these health problems.

ADHD is the most common mental disorder diagnosed in children and usually begins in childhood and becomes prominent in later school years. The global prevalence of ADHD ranges from 2% to 18% [6]. ADHD screening is encouraged in the early childhood to promote early diagnosis and symptom management to improve the functioning of children and their families [6]. Scoliosis is an abnormal lateral curvature of the spine, as defined by the American Association of Neurological Surgeons [7]. It is often diagnosed in childhood or early adolescence. Scoliosis screening at early ages is important to detect abnormalities and promptly launch conservative treatment in order to avoid necessitating invasive surgery later in life [7]. Obesity is a global health issue among schoolchildren. In 2016, the World Health Organization (WHO) reported that more than 340 million school-aged children were obese [8]. It is well known that childhood obesity continues into adulthood and is a risk factor for several chronic health conditions later in life [9]. Similarly, other diseases and health problems that arise in childhood affect students' health and impact their quality of life and future opportunities. This study aimed to determine the prevalence of health problems targeted by the NSBSP among primary school students in the KSA in 2019.

2. Materials and Methods

This is a cross-sectional study based on a secondary analysis of data obtained from the NSBSP. This study was conducted by the General Administration of School Health (GASH) during the academic year 2018-2019. Data analysis was performed on students from both private and public primary schools from the 20 health regions of the KSA. We included screening data from the first and fourth grades of primary schools in the KSA. The total number of participants was 444,259, excluding students with incomplete data who were excluded from the analysis ($n = 3,009$).

NSBSP is a routine surveillance health program organized and implemented by the GASH at the MOH. Well-trained health teams participated in the screening program. Each health team comprised physicians, nurses, and dentists under the direct supervision of GASH officials. Following examination and screening of students, data obtained during screening were entered into the electronic system of the health screening program. The collected data included sociodemographic characteristics (age, sex, school grade, and region), immunization status, body mass index, visual acuity, dental cavities, hearing problems, scoliosis, and ADHD.

To assess immunization status, health teams inquired about the completion of pre-school immunizations as specified by the National Advisory Committee on Immunization. The Saudi immunization schedule includes the oral polio vaccine; measles, mumps and rubella vaccine; varicella; diphtheria tetanus vaccine; and whole-cell pertussis vaccine (or diphtheria, tetanus, and pertussis vaccine for those 7 years or older). Moreover, the health team physician assessed meningococcal vaccine and influenza vaccine by examining the student's immunization schedule card [10].

Health teams nurses calculated the body mass index (BMI) after measuring the weight and height of each student. BMI data were plotted on Saudi's sex-specific percentile charts. BMI was categorized into four groups: underweight (less than 5th percentile); normal weight (between 5th percentile and 85th percentile); overweight (between 85th

and 95th percentile); and obese (more than 95th percentile) [11]. Health teams examined visual acuity and tested students' near vision using Snellen charts. Students who scored $\leq 6/12$ were considered to be myopic. Dentists assessed the students for the presence of dental cavities and reported them in the electronic records. In cases in which the teacher suspected that a student had a hearing problem, the health teams examined hearing using an audiometer. The present analysis includes hearing examination data of the 1st grade students only. We considered the normal hearing range 1000-4000 Hz at an intensity of 25 dB [12]. Health team physicians assessed scoliosis in 4th grade students using Adam's forward bending test [7]. ADHD was assessed in suspected 1st grade students using the Vanderbilt ADHD Rating Scale based on assessment forms received from parents and teachers.

2.1. Statistical Analysis

Data analysis was performed using SPSS version 21 (IBM, Armonk, NY, USA). The data were cleaned for incomplete or missing variables. Qualitative data are presented as frequencies and percentages, and quantitative data are represented as mean \pm standard deviation (SD). Pearson's chi-square test was used to analyze the categorical variables. A two-sided significance test was used, with a p value <0.05 .

2.2. Ethical considerations

Approval to conduct this study was obtained from the Institutional Review Board (IRB) of the MOH prior to study execution.

3. Results

The total number of students enrolled in this study was 444,259, of which 53.4% were girls and 46.6% were boys, as shown in Table 1. The students' ages ranged from 6 to 14 years, and the mean age was $(8.06 \pm 1.7$ years). The majority of the students had completed preschool vaccinations (95.2%). Dental cavities, myopia, and obesity/overweight were the most prevalent health problems for primary school students, as they were observed in 38.7%, 10.9%, and 10.5% of screened students, respectively (Table 2). ADHD, scoliosis, and auditory health problems were prevalent in only 2.8%, 0.48%, and 0.7% of screened students, respectively.

Table 1. The frequency distribution of students in relation to educational grade and sex.

Parameter	N = 444259	%
Age (Years)		
Mean \pm SD	8.06 ± 1.7	
Range	6-14	
Student's grade		
First grade	219,218	49.3
Fourth grade	225,041	50.7
Sex		
Boys	207,109	46.6
Girls	237,150	53.4

Table 2. The prevalence of health problems among screened primary school students in KSA (N = 444259).

Health problem	N	%
Not completed immunization*	10,521	4.80%

Eye refractory errors	48,313	10.90%
Overweight and obesity	46,658	10.50%
Overweight	28,430	6.40%
Obesity	18,228	4.10%
Dental caries	172,113	38.70%
Scoliosis and/or kyphosis**	1,086	0.48%
ADHD disorders ***	878	2.81%
Attention deficit (AD)	422	1.35%
Hyperactivity disorder (HD)	165	0.53%
Both (AD plus HD)	291	0.93%
Auditory problems ***	109	0.7%
Right ear	17	0.1%
Left ear	20	0.1%
Both ears	72	0.47%

* Only 1st grade students (N= 219,218).

** Only 4th grade students screened (N=225,041).

*** Students suspected by parents and teachers only in 1st grade (number of students suspected with ADHD suspected students: 31,210, and with auditory problem: 15,426).

ADHD: Attention-Deficit/Hyperactivity Disorder

Health problems that were significantly higher among girls than boys were scoliosis (0.7% vs. 0.2%), error of refraction (12.1% vs. 9.4%), obesity and overweight (11.3% vs. 9.6%), dental cavities (41.5% vs. 35.6%), and ADHD (1.2% vs. 0.7%). However, incomplete vaccination status and auditory problems were significantly higher among boys than girls (5.2% vs. 4.4%) and (1.0% vs. 0.6%), respectively (p = 0.002) (Table 3).

Table 3. The association of health problems with sex of screened primary school students (N = 444259)

Health problem	Sex		P value
	Boys N = 207,109	Girls N = 237,150	
Incomplete Vaccination status *	5,354 (5.2%)	5,167 (4.4%)	<0.001
Scoliosis and/or kyphosis **	216 (0.2%)	870 (0.7%)	<0.001
Eye refractory errors	19,528 (9.4%)	28,785 (12.1%)	<0.001
Obesity and overweight	19,896 (9.6%)	26,762 (11.3%)	<0.001
Overweight	11,947 (5.8%)	16,483 (7%)	<0.001
Obesity	7,949 (3.8%)	10,279 (4.3%)	<0.001
Dental caries	73,701 (35.6%)	98,412 (41.5%)	<0.001
ADHD disorders ***	353 (1.1%)	525 (1.2%)	0.002
Auditory problems ****	52 (1.0%)	57 (0.6 %)	0.002

* Only 1st grade students (N= 219,218; boys =102,490, girls = 116,728).

** Only 4th grade students screened (N = 225,041; boys =104,619, girls = 120,422).

*** Students suspected by parents and teachers only in 1st grade. (N = 31,210 for ADHD problems; boys, 14,142, girls= 17,068).

**** N= 15,426 for auditory problems; boys = 5,186, girls = 10,240) Students suspected by parents and teachers only in 1st grade.

Dental cavities were highly prevalent among primary school students in Jazan (78.1%), Najran (77.9%), Tabouk (75.0%), and Makkah (72.8%). ADHD was prevalent in Al-Baha (51.5%), and obesity and overweight were prevalent in the northern borders (44.7%), and Al-Jouf (38.5%). Errors of refraction were prevalent in the eastern region, Al-Jouf, and Al-Hasa (32.5%, 31.5%, and 30.4%, respectively). The highest prevalence of auditory problems was observed among students in the eastern region (11.7%), while scoliosis prevalence was highest among students in Makkah (1.9%) and the Eastern Region (1.1%). Of all regions, incomplete preschool vaccination was highest among students in Riyadh (24.2%) and Jeddah (19.6%), as shown in Table 4.

Table 4. Distribution of health problems of screened primary school students according to KSA health regions.

Region	Total screened	Obesity & overweig ht N (%)	Errors of refraction N (%)	Dental caries N (%)	Scoliosis N* (%)	ADHD N** (%)	Auditory problems N** (%)	Incomplete immunization N*** (%)
Riyadh	68,032	12,725 (18.7)	11,882 (17.5)	42,955 (63.1)	277 (0.4)	160 (1.7)	33 (0.85)	2,549 (24.2%)
Jeddah	30,291	4,384 (14.5)	4,007 (13.2)	21,773 (71.9)	77 (0.4)	49 (1.5)	1 (0.1)	2,067 (19.6%)
Eastern Region	28,133	6,724 (23.9)	9,139 (32.5)	11,981 (42.6)	212 (1.1)	70 (12.0)	7 (11.7)	986 (9.4%)
Makkah	25,841	2,781 (10.8)	3,927 (15.2)	18,805 (72.8)	326 (1.9)	1 (100.0)	1 (100.0)	953 (9.1%)
Qassim	17,913	3,168 (17.7)	3,641 (20.3)	10,977 (61.3)	29 (0.2)	94 (6.7)	4 (3.5)	670 (6.4%)
Jazan	10,953	1,303 (11.9)	996 (9.1)	8,552 (78.1)	34 (0.3)	54 (2.7)	14 (0.9)	141 (1.3%)
Hail	7,510	1,474 (19.6)	689 (9.2)	5,300 (70.6)	14 (0.1)	26 (6.9)	7 (2.7)	344 (3.3%)
Al-Hasa	17,510	2,868 (16.4)	5,320 (30.4)	9,061 (51.7)	20 (0.2)	229 (4.9)	12 (0.3)	54 (0.5%)
Aseer	8,602	1,880 (21.9)	1,505 (17.5)	5,148 (59.8)	19 (0.2)	46 (2.7)	4 (0.3)	443 (4.2%)

Tabouk	11,715	1,219 (10.4)	1,626 (13.9)	8,789 (75.0)	5 (0.5)	70 (17.5)	6 (4.8)	920 (8.7%)
Al-Taif	6,778	927 (13.7)	1,081 (15.9)	4,702 (69.4)	32 (0.4)	34 (2.6)	2 (1.8)	701 (6.7%)
Medina	9,787	1,593 (16.3)	1,115 (11.4)	7,070 (72.2)	6 (0.1)	3 (0.5)	0 (0.0)	41 (0.4%)
Najran	3,568	454 (12.7)	330 (9.2)	2,780 (77.9)	3 (0.1)	0 (0.0)	1 (1.9)	262 (2.5%)
Al-Jouf	2,986	1,151 (38.5)	940 (31.5)	889 (29.8)	2 (0.0)	4 (3.4)	0 (0.0)	269 (2.6%)
Bisha	2,334	518 (22.2)	182 (7.8)	1,623 (69.5)	4 (0.2)	4 (0.3)	3 (0.6)	17 (0.2%)
Northern Borders	1,475	659 (44.7)	89 (6.0)	708 (48.0)	7 (0.2)	3 (0.2)	9 (1.0)	57 (0.5%)
Hafr Al- Batin	4,190	1,052 (25.1)	171 (4.1)	2,957 (70.6)	7 (0.2)	0 (0.0)	3 (0.4)	9 (0.1%)
Al-Baha	4,289	518 (12.1)	769 (17.9)	2,981 (69.5)	4 (0.1)	17 (51.5)	0 (0.0)	16 (0.2%)
Al-Qunfudhah	4,544	661 (14.5)	776 (17.1)	3,087 (67.9)	6 (0.2)	14 (3.2)	0 (0.0)	22 (0.2%)
Al-Qurayyat	68,032	12,725 (18.8)	11,882 (17.5)	42,955 (63.1)	2 (0.1)	0 (0.0)	2 (100.0)	0 (0.0%)

* Only 4th grade students (N = 225,041)

** Students suspected by parents and teachers in each region.

*** Only 1st grade students (N=219,218)

4. Discussion

This study presents the nationwide prevalence of health problems in school-aged children identified by the screening program. The data presented was collected from first- and fourth-grade Saudi primary school students (n = 444,259) during the academic year 2018-2019. This dataset explored variations in the prevalence of the screened conditions across the 20 regions of the KSA. The large sample size and numerous health problems screened for by this program add strength and uniqueness to this study. The prevalence of health problems ranged from 0.48% to 38.7%, which highlights the importance of screening and early detection of health problems, allowing early management. The proportion of completely immunized for schoolchildren has risen from 75% in 1993 to 95.2% in the current study [13, 14]. This could be attributed to the tremendous efforts of the Saudi MOH in providing childhood vaccinations that are accessible at primary healthcare centers on a daily basis. This effort is further enhanced by MOH endeavors to increase vaccination adherence. MOH, through its vaccination reminder services, contributes to raising awareness and positively affects the attitude of parents towards vaccination, increasing childhood vaccination [15, 16]. A recent systematic review conducted in the KSA

demonstrated overall positive attitudes and awareness among parents regarding childhood vaccination [17].

Our study showed that the prevalence of overweight and obesity among school students was 6.4% and 4.1%, respectively, which is lower than rates reported by other local studies, where the prevalence of overweight and obesity was 10.7–14.4% and 7.9%–13.9%, respectively [18,19]. Our results are also lower than those reported in the National Growth Study conducted by Al Mouzan et al. in KSA in 2010, where the prevalence of overweight and obesity in schoolchildren aged 5–12 years was 19.6% and 7.9%, respectively [20]. However, several BMI percentile standards are used to define overweight/obesity among the pediatric population, namely the Center for Disease Control (CDC) standards, World Health Organization (WHO) standards, and country-specific standards [21]. Most previous studies have employed the CDC or WHO standards. Moreover, Al-Mendalawi has affirmed the possible overestimation of the prevalence of overweight and obesity among Saudi children when using standards other than country-specific growth charts [21]. In 2016, Al Mouzan et al. established a new reference for the evaluation of nutrition and growth specific values for Saudi school-age children and adolescents, which provides a more accurate clinical assessment [11]. This chart was used to screen primary school students in the current study. A study conducted in the KSA in 2019 found that increasing physical education time in schools was positively associated with an increase in physical activity. The study concluded that Saudi school students did not meet the recommended level of physical activity [22]. This highlights the importance of engaging children in physical education and activity, given its role in reducing the prevalence of overweight and obesity [23].

The prevalence of eye refractory errors (myopia) was 10.9% in our study. This is similar to the findings from a local study, which reported a prevalence of 9% among primary school students aged 6–14 years [24]. However, according to the literature, there are huge variations in the prevalence of myopia among school students. A prevalence of 3.3% was reported among schoolchildren aged 6–7 years old in Ireland [25], 14.9% among 6–15 year old school students in Iran [26], and 35.8% in 6–8-year-old children in China [27]. Myopia was as high as 63.1%–78.6% in 6–8-year-old children in Japan [28]. The discrepancy in the definition of myopia and different screening methods played a significant role in this interstudy variation. Furthermore, in our study, the prevalence of myopia was found to be higher among females (12.1% girls vs. 9.4% boys), which is consistent with other local studies [24]. In contrast, other international studies have found that myopia is higher among boys [29, 30].

In the current study, the prevalence of auditory problems was 0.7% among first-grade students. This prevalence is lower than that found in international studies, where the prevalence is 8.1% in Canada [31], 19.6% in Brazil [32], and 27.2% in Kyrgyzstan [33]. However, in the latter studies, the screening threshold was 20 dB, while in the current study, it was at 25 dB, resulting in fewer cases detected. Additionally, only students suspected of hearing issues were screened. The assessment was based on the specific signs identified by teachers. Failure to recognize students led to their exclusion from screening which may have resulted in a lower prevalence.

Dental pain and cavities are global issues and may lead to student absences from school and sleep disturbances, hence affecting academic achievement [5]. The prevalence of dental cavities in this study was 38.7%. This is much lower than that reported in a systematic review conducted in the KSA in 2013, which reported a prevalence of 70% based on data obtained from children's primary dentists. However, the review included 26 studies conducted between 1988 and 2010 among different age groups [34], which could explain the variation in the results obtained. It is well known that first permanent molars usually erupt during the first or second grade (6–7 years old), while adolescents at age 13 usually have 28 permanent teeth. [35] Even though the current study included only first- and fourth-grade students, the screening for dental cavities included permanent teeth. Additionally, girls typically erupt permanent teeth much earlier than boys, explaining the

higher prevalence of dental cavities in girls compared to boys (41.5% vs. 35.6%) [36, 37]. A local study conducted among schoolchildren aged 6–9 years in Riyadh showed that the prevalence of dental cavities was 14.4% [38]. Furthermore, a worldwide systematic review (1995–2019) reported that the prevalence of dental cavities in children's permanent teeth was 53.8% [39].

The overall prevalence of scoliosis and/or kyphosis was 0.48% among fourth-grade students in this study. This is lower than that reported in two systematic reviews conducted in China and Australia, where the prevalence of scoliosis among students was 1.02% and 1.1%, respectively. Nonetheless, these studies included older children, in whom the prevalence was usually higher [40]. In the present study, the prevalence of scoliosis among girls was higher than that among boys (0.7% vs. 0.2%). This is in line with the results of previous studies that reported that scoliosis was higher among females [41, 42]. A study conducted among school boys aged 10–14 years in Al-Kharj reported a prevalence of 0.08% in adolescent idiopathic scoliosis [42], which is lower than the prevalence reported in our study among boys. However, the current study employed a larger sample size, and also included screening for kyphosis.

ADHD is one of the most common mental disorders that usually begins in childhood and becomes more prominent during school years [6]. Worldwide, studies of ADHD among school-aged children revealed a prevalence between 2% and 18% [6]. In KSA, different studies have been conducted and have revealed variations in the overall rates of ADHD, ranging from 2.7% to 7.4% [43–45]. This is in agreement with the overall prevalence reported in the current study (2.8%). In contrast to previous studies included in a systematic review of ADHD in Arab countries [46], our study found that ADHD was more prevalent in girls. This finding could be attributed to the teachers' own judgment in the selection of students suspected of ADHD, which was merely based on signs and symptoms. Anastopoulos et al. reported that the sex of the informant could play a clinically significant role in assessing ADHD in children, as female teachers and parents' ratings of symptoms were significantly higher than those of male teachers and parents [47].

One limitation of this study was the unavailability of sociodemographic data in the school health-screening program. The inclusion of such data could allow for a more meaningful understanding of the determinants of health issues. Another limitation was that screening for ADHD and hearing issues was only carried out in selected students in the first grade, which could obscure the true prevalence of these conditions. Involving all students in both the first and fourth grades could allow for better assessment of the prevalence of ADHD and hearing issues.

5. Conclusions

The prevalence of health problems among primary school students ranged from 0.48% to 38.7%. The most common health problems were dental cavities, error of refraction, obesity, and being overweight. These problems were significantly higher among girls than among boys. Appropriate school health programs should be implemented in primary schools in the KSA to address and prevent these problems.

Author Contributions: **Author Contributions:** Conceptualization and methodology and writing—original draft preparation, writing—review and editing were done by Manal Al Daajani, Dina Al-Habib, Nora Al Shewair, Yahya Fagihi, Abrar Alzaher, Amjad Alfaleh; software, validation, formal analysis and data curation were done by Mona Ibrahim, Khaled Alabdulkareem; resources and funding acquisition done by Manal Al Daajani; Supervision and project administration were done by Manal Al Daajani, Mona Ibrahim and Khaled Alabdulkareem.

All authors have read and agreed to the published version of the manuscript." Please turn to the CRediT taxonomy for the term explanation. Authorship must be limited to those who have contributed substantially to the work reported.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board of Central Committee of Research, MOH, KSA (protocol code 2019-0103M Date: 28/7/2019).

Informed Consent Statement: Participant consent was waived as this is study was based on the secondary data of the national survey with involvement of large number of students. Written approval to publish this paper has been obtained from the General Directorate for Research and Studies (GDRS), MOH, KSA (Date: 24/8/2021).

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to privacy concerns.

Acknowledgments: We acknowledge the contribution of every screening health team member in the schools, the administrators, and the schoolteachers across the KSA, as well as the administrators, managers, and assistants at the Directorate General for School Health at MOH. We are very thankful to regional school health administrations for their efforts to collect data

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Basch, C., 2011. Healthier Students Are Better Learners: A Missing Link in School Reforms to Close the Achievement Gap. *Journal of School Health*, 81(10), pp.593-598.
2. Gracy, D., Fabian, A., Roncaglione, V., Savage, K. and Redlener, I., 2017. Health Barriers to Learning: The Prevalence and Educational Consequences in Disadvantaged Children. [online] Childrenshealthfund.org. Available at: <<https://www.childrenshealthfund.org/wp-content/uploads/2017/01/Health-Barriers-to-Learning.pdf>> [Accessed 2 August 2020].
3. Ying, G. and Quinn, G., 2016. Uncorrected hyperopia and preschool early literacy: results of the Vision in Preschoolers-Hyperopia In Preschoolers (VIP-HIP) Study. *Journal of American Association for Pediatric Ophthalmology and Strabismus*, 20(4), p.e11.
4. Lieu, J., Tye-Murray, N., Karzon, R. and Piccirillo, J., 2010. Unilateral Hearing Loss Is Associated With Worse Speech-Language Scores in Children. *PEDIATRICS*, 125(6), pp.e1348-e1355.
5. Seirawan, H., Faust, S. and Mulligan, R., 2012. The Impact of Oral Health on the Academic Performance of Disadvantaged Children. *American Journal of Public Health*, 102(9), pp.1729-1734.
6. Polanczyk, G., de Lima, M., Horta, B., Biederman, J. and Rohde, L., 2007. The Worldwide Prevalence of ADHD: A Systematic Review and Metaregression Analysis. *American Journal of Psychiatry*, 164(6), pp.942-948.
7. Aans.org. 2021. Scoliosis – Symptoms, Diagnosis and Treatment. [online] Available at: <<https://www.aans.org/en/Patients/Neurosurgical-Conditions-and-Treatments/Scoliosis>> [Accessed 4 September 2021].
8. Who.int. 2021. Obesity and overweight. [online] Available at: <<https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>> [Accessed 4 September 2021].
9. Biro, F. and Wien, M., 2010. Childhood obesity and adult morbidities. *The American Journal of Clinical Nutrition*, 91(5), pp.1499S-1505S.

10. Ministry of Health Saudi Arabia. 2021. Saudi Immunization Schedule. [online] Available at: <<https://www.moh.gov.sa/en/HealthAwareness/EducationalContent/HealthTips/Documents/Immunization-Schedule.pdf>> [Accessed 18 September 2020].
11. El Mouzan, M., Al Salloum, A., Alqurashi, M., Al Herbish, A. and Al Omar, A., 2016. The LMS and Z scale growth reference for Saudi school-age children and adolescents. *Saudi Journal of Gastroenterology*, 22(4), p.331.
12. Gates, G. and Hoffman, H., 2011. What the Numbers Mean: An Epidemiological Perspective on Hearing. [online] NIDCD. Available at: <<https://www.nidcd.nih.gov/health/statistics/what-numbers-mean-epidemiological-perspective-hearing>> [Accessed 11 April 2020].
13. Rashid, A., 1993. Childhood Immunization Status Related to Social and Educational Status of Parents in a Peripheral Northern Town of Saudi Arabia. *Annals of Saudi Medicine*, 13(4), pp.335-339.
14. Ministry of Health, Saudi Arabia. 2019. Vaccinations Description. [online] Available at: <<https://www.moh.gov.sa/en/eServices/Pages/VaccinationsDescription.aspx>> [Accessed 4 September 2021].
15. Alshammari, T., Subaiea, G., Hussain, T., Moin, A. and Yusuff, K., 2018. Parental perceptions, attitudes and acceptance of childhood immunization in Saudi Arabia: A cross sectional study. *Vaccine*, 36(1), pp.23-28.
16. Alsubaie, S., Gosadi, I., Alsaadi, B., Albacker, N., Bawazir, M., Bin-Daud, N., Almanie, W., Alsaadi, M. and Alzamil, F., 2019. Vaccine hesitancy among Saudi parents and its determinants. *Saudi Medical Journal*, 40(12), pp.1242-1250.
17. Alabadi, M. and Aldawood, Z., 2020. Parents' Knowledge, Attitude and Perceptions on Childhood Vaccination in Saudi Arabia: A Systematic Literature Review. *Vaccines*, 8(4), p.750.
18. Al-Hussaini, A., Bashir, M., Khormi, M., AlTuraiki, M., Alkhamis, W., Alrajhi, M. and Halal, T., 2019. Overweight and obesity among Saudi children and adolescents: Where do we stand today?. *Saudi Journal of Gastroenterology*, 25(4), p.229.
19. Al Shaikh A, Farahat F, Abaalkhail B, Kaddam I, Aseri K, Al Saleh Y, Al Qarni A, Al Shuaibi A, Tamimi W. Prevalence of Obesity and Overweight among School-Aged Children in Saudi Arabia and Its Association with Vitamin D Status. *Acta Biomed*. 2020 Nov 3;91(4):e2020133. doi: 10.23750/abm.v91i4.10266. PMID: 33525269; PMCID: PMC7927516.
20. El Mouzan, M., Foster, P., Al Herbish, A., Al Salloum, A., Al Omer, A., Qurachi, M. and Kecojevic, T., 2010. Prevalence of overweight and obesity in Saudi children and adolescents. *Annals of Saudi Medicine*, 30(3), pp.203-208.
21. Al-Mendalawi, M., 2019. Is the prevalence of overweight/obesity overestimated among Saudi children and adolescents? *Saudi Journal of Gastroenterology*, 25(6), p.0.

22. Aljuhani and Sandercock, 2019. Contribution of Physical Education to the Daily Physical Activity of Schoolchildren in Saudi Arabia. *International Journal of Environmental Research and Public Health*, 16(13), p.2397.
23. Aliss, E., Sutaih, R., Kamfar, H., Alagha, A. and Marzouki, Z., 2020. Physical activity pattern and its relationship with overweight and obesity in Saudi children. *International Journal of Pediatrics and Adolescent Medicine*, 7(4), pp.181-185.
24. Al Wadaani, F., Amin, T., Ali, A. and Khan, A., 2012. Prevalence and pattern of refractive errors among primary school children in Al Hassa, Saudi Arabia. *Global Journal of Health Science*, 5(1).
25. Harrington, S., Stack, J., Saunders, K. and O'Dwyer, V., 2018. Refractive error and visual impairment in Ireland schoolchildren. *British Journal of Ophthalmology*, 103(8), pp.1112-1118.
26. Norouzirad, R., Hashemi, H., Yekta, A., Nirouzad, F., Ostadioghaddam, H., Yazdani, N., Dadbin, N., Javaherforoushzadeh, A. and Khabazkhoob, M., 2015. The prevalence of refractive errors in 6- to 15-year-old schoolchildren in Dezful, Iran. *Journal of Current Ophthalmology*, 27(1-2), pp.51-55.
27. Zhou J, Ma Y, Ma J, Zou Z, Meng X, Tao F, Luo C, Jing J, Pan D, Luo J, Zhang X, Wang H, Zhao H. [Prevalence of myopia and influencing factors among primary and middle school students in 6 provinces of China]. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2016 Jan;37(1):29-34. Chinese. doi: 10.3760/cma.j.issn.0254-6450.2016.01.006. PMID: 26822639.
28. Yotsukura, E., Torii, H., Inokuchi, M., Tokumura, M., Uchino, M., Nakamura, K., Hyodo, M., Mori, K., Jiang, X., Ikeda, S., Kondo, S., Negishi, K., Kurihara, T. and Tsubota, K., 2019. Current Prevalence of Myopia and Association of Myopia With Environmental Factors Among Schoolchildren in Japan. *JAMA Ophthalmology*, 137(11), p.1233.
29. Norouzirad, R., Hashemi, H., Yekta, A., Nirouzad, F., Ostadioghaddam, H., Yazdani, N., Dadbin, N., Javaherforoushzadeh, A. and Khabazkhoob, M., 2015. The prevalence of refractive errors in 6- to 15-year-old schoolchildren in Dezful, Iran. *Journal of Current Ophthalmology*, 27(1-2), pp.51-55.
30. Singh, N., James, R., Yadav, A., Kumar, R., Asthana, S. and Labani, S., 2019. Prevalence of Myopia and Associated Risk Factors in Schoolchildren in North India. *Optometry and Vision Science*, 96(3), pp.200-205.
31. Feder, K., Michaud, D., McNamee, J., Fitzpatrick, E., Ramage-Morin, P. and Beauregard, Y., 2017. Prevalence of Hearing Loss Among a Representative Sample of Canadian Children and Adolescents, 3 to 19 Years of Age. *Ear & Hearing*, 38(1), pp.7-20.
32. Nunes, A., Balen, S., Souza, D. and Barbosa, I., 2019. Prevalence of Hearing Loss and Associated Factors in School-Age Individuals in an Urban Area of Northeast Brazil. *International Archives of Otorhinolaryngology*, 24(03), pp.e330-e337.

33. Skarżyński, P., Świerniak, W., Gos, E., Pierzyńska, I., Walkowiak, A., Cywka, K., Wołujewicz, K. and Skarżyński, H., 2020. Results of hearing screening of school-age children in Bishkek, Kyrgyzstan. *Primary Health Care Research & Development*, 21.
34. Al Agili, D., 2013. A systematic review of population-based dental caries studies among children in Saudi Arabia. *The Saudi Dental Journal*, 25(1), pp.3-11.
35. Ada.org. 2006. Tooth eruption. The permanent teeth. [online] Available at: <https://www.ada.org/~/media/ada/publications/files/patient_58.ashx> [Accessed 4 February 2020].
36. Bayrak, S., Sen Tunc, E., Tuglu, N. and Acikgoz, A., 2012. Timing of Permanent Teeth Eruption in Turkish Children. *Journal of Clinical Pediatric Dentistry*, 37(2), pp.207-211.
37. Diamanti, J. and Townsend, G., 2003. New standards for permanent tooth emergence in Australian children. *Australian Dental Journal*, 48(1), pp.39-42.
38. Alshihha, S., Alwayli, H., Hattan, M., Alfraih, Y., Alamri, A. and Aldossary, M., 2018. Prevalence of Dental Caries and Fissure Sealants in the First Permanent Molars among Male Children in Riyadh, Kingdom of Saudi Arabia. *International Journal of Clinical Pediatric Dentistry*, 11(5), pp.365-370.
39. Kazeminia, M., Abdi, A., Shohaimi, S., Jalali, R., Vaisi-Raygani, A., Salari, N. and Mohammadi, M., 2020. Dental caries in primary and permanent teeth in children's worldwide, 1995 to 2019: a systematic review and meta-analysis. *Head & Face Medicine*, 16(1).
40. Zhang, H., Guo, C., Tang, M., Liu, S., Li, J., Guo, Q., Chen, L., Zhu, Y. and Zhao, S., 2015. Prevalence of Scoliosis Among Primary and Middle School Students in Mainland China. *Spine*, 40(1), pp.41-49.
41. Baroni, M., Sanchis, G., Assis, S., Santos, R., Pereira, S., Sousa, K. and Lopes, J., 2015. Factors Associated With Scoliosis in Schoolchildren: a Cross-Sectional Population-Based Study. *Journal of Epidemiology*, 25(3), pp.212-220.
42. Elshazly, FA., Ahmed, AR., Mahmoud, WS., Ahmed, AS., 2014. Screening Study for Early Detection of Scoliosis in School Children in Al-kharj City in KSA. *World Applied Sciences Journal*, 31(6), pp.993-997.
43. Alqahtani, M., 2010. Attention-deficit hyperactive disorder in school-aged children in Saudi Arabia. *European Journal of Pediatrics*, 169(9), pp.1113-1117.
44. Alhraiwil, N., Ali, A., Househ, M., Al-Shehri, A. and El-Metwally, A., 2015. Systematic review of the epidemiology of attention deficit hyperactivity disorder in Arab countries. *Neurosciences*, 20(2), pp.137-144.
45. Homidi M., Obaidat Y., and Hamaidi D. 2013. Prevalence of attention deficit and hyperactivity disorder among primary school students in Jeddah city, KSA. *Life Science Journal*. 10(3), pp. 280-285.
46. Alhraiwil, N., Ali, A., Househ, M., Al-Shehri, A. and El-Metwally, A., 2015. Systematic review of the epidemiology of attention deficit hyperactivity disorder in Arab countries. *Neurosciences*, 20(2), pp.137-144.

47. Anastopoulos, A., Beal, K., Reid, R., Reid, R., Power, T. and DuPaul, G., 2018. Impact of child and informant gender on parent and teacher ratings of attention-deficit/hyperactivity disorder. *Psychological Assessment*, 30(10), pp.1390-1394.