

Bioassay-guided Identification of Bioactive Compounds from *Senna alata* L. against Methicillin-resistant *Staphylococcus aureus*

Suwaiba Muhammad Ladan ¹, Yusuf Wada ^{2,3,*}, Mustapha Mohammed ^{4,5}, Sani Ibrahim ⁶, Kabir Musa Yusuf ⁷, Olayeni Stephen Olonitola ⁸, Mubarak Hussaini Ahmad ⁹, Sagir Mustapha ^{9,10}, Zaidah Abdul Rahman ^{2,11} and Abubakar Sha'ban ^{4,5,*}

¹ Department of Chemical Science, Federal University of Kashere, Gombe, 0182, Gombe, Nigeria; suwaibana@gmail.com

² Department of Medical Microbiology and Parasitology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, 16150, Kelantan, Malaysia;

³ Department of Zoology, Faculty of Life Sciences, Ahmadu Bello University, Zaria, 810107, Kaduna, Nigeria

⁴ School of Pharmaceutical Sciences, Universiti Sains Malaysia, Minden, 11800, Pulau Pinang, Malaysia; mohammedmmrx@gmail.com, abuushaaban2@gmail.com

⁵ Department of Clinical Pharmacy and Pharmacy Practice, Ahmadu Bello University, Zaria, 810107, Kaduna, Nigeria

⁶ Department of Biochemistry, Ahmadu Bello University, Zaria, 810107, Kaduna, Nigeria; sanibro2000@yahoo.com

⁷ Department of Pharmacognosy and Drug Development, Ahmadu Bello University, Zaria, 810107, Kaduna, Nigeria; drashrafamjad921@gmail.com

⁸ Department of Microbiology, Ahmadu Bello University, Zaria, 810107, Kaduna, Nigeria; olonisteve@gmail.com

⁹ Department of Pharmacology and Therapeutics, Ahmadu Bello University, Zaria, 810107, Kaduna, Nigeria; mubarakhussainiahmad@gmail.com, maimunat001@gmail.com

¹⁰ Department of Pharmacology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, 16150, Kelantan, Malaysia

¹¹ Hospital Universiti Sains Malaysia, Universiti Sains Malaysia, Kubang Kerian, 16150, Kelantan, Malaysia; drzaidah@usm.my

* Correspondence: abuushaaban2@gmail.com (AS); wadayusuf34@gmail.com (YW); mohammedmmrx@gmail.com (MM); suwaibana@gmail.com (SML)

Abstract: *Senna alata* (Linn) Roxb. plant is widely used to manage various infections in folkloric medicine. Methicillin-resistant *Staphylococcus aureus* (MRSA) infection continues to be a major global public health problem. This study aims to investigate the bioactive components of *S. alata* leaves active against MRSA. The leaves of *S. alata* were sequentially extracted and fractionated using standard methods and screened for activities against MRSA. The diethyl ether active thin layer chromatography (TLC) spot was subjected to infrared (IR) and gas chromatography-mass spectroscopic (GC-MS) studies. The aqueous extract and diethyl ether fraction of *S. alata* leaves elicited the highest activity against the MRSA. The GC-MS analysis of the fraction produced 15 eluates; only the sub-fraction 13 was effective. The TLC analysis of the sub-fraction 13 revealed three spots; only the second spot produced activity. The GC-MS result of the spot showed six peaks. The spectral results for peak 3 match the data from the IR study suggestive of 9-octadecenoic acid methyl ester. *Senna alata* leaves possess bioactive compounds closely related to 9-octadecenoic acid methyl ester with potent antibacterial activity against MRSA.

Keywords: bioassay; chromatography; folkloric medicine; methicillin-resistant *Staphylococcus aureus*; *Senna alata*; spectroscopy

1. Introduction

Infectious diseases are among the top causes of global mortality, morbidity and disability [1,2]. The pathogens such as *Staphylococcus aureus* form a serious burden on health due to the related infections, including pneumonia, endocarditis, and sepsis [3,4]. The methicillin-resistant *Staphylococcus aureus* (MRSA) is one of the most critical multidrug-resistant (MDR) pathogens mostly encountered in hospitals. [5–7]. The global incidence of MRSA infection has increased the burden on the health care system and growing death over the years. [8]. Besides, the MRSA infection still extends to the community population. [7]. Due to the frequent emergence of antimicrobial resistance, limited options are available for treating MRSA-related infections [9]. The ineffectiveness of the available antibacterial agents necessitates the search for new compounds for use against resistant bacteria. [10].

Plants have contributed to the general health and well-being of the populace for a long time [11]. About 75% of the people globally depend on traditional medicine for their basic health requirements [12]. Many therapeutic compounds employed in orthodox medicine originate from plants [12]. Plants synthesize various secondary metabolites with different chemical diversity and biological actions [13]. Plants and their secondary metabolites have been reported to serve as sources of antimicrobial agents [14–17]. For years, attention has been shifted to investigate phytochemical compounds with potential antibacterial activity, particularly against MDR bacteria [18]. Given the roles of medicinal plants in curtailing infections disease, there is a need to continue the search for novel therapeutic agents to pave the way for the discovery of new therapeutic agents [19,20].

The plant *Senna alata* (Linn) Roxb. synonymous with *Cassia alata* (Leguminosae), is commonly distributed in Asia, Brazil, Australia, and all-over African nations, including Egypt, Somalia and Nigeria [21]. The plant is a tropically erect annual herb that grows up to 2-5 meters in high. It possesses big, leathery and slippery compound leaves bilateral that usually fold at night [22]. It has a pod fruit with small and square-shaped seeds [23]. In English, this medicinally important plant is called candle bush, craw-craw, Acapulco or ringworm bush plant [21]. The local names of the plant are *Asunwon oyinbo* in Yoruba, *Nelkhi* in Igbo, and *Hantsi* in the Hausa languages of Nigeria [24]. The plant is used as herbal preparation to manage hepatitis, gastroenteritis, constipation, dermatitis, eczema, jaundice, diarrhea [25], and bacterial infections [26]. The plant's leaves, stems, and roots have been ingested as a decoction to treat wounds, respiratory tract, and skin infections in Northern Nigeria [25]. Also, in Cameroon, the leaves, stems and bark of *S. alata* were documented to be used to treat gastroenteritis and skin infections [21]. Its leaves, bark and stem are effective against intestinal parasitosis and syphilis as a decoction in China, Philippines and India [27]. Keeping in view the ethnopharmacological indication and promising therapeutic properties of *S. alata* for treating infectious ailments, this study was conducted to screen the leaves extracts of *S. alata* for activity against MRSA and identify the bioactive compounds likely responsible for the activity.

2. Materials and Methods

2.1. Plant collection and identification

The whole plant *Senna alata* was sourced from Jaji village, along Kaduna-Zaria Road, Kaduna State of Nigeria. The authentication of the plant material was conducted at the Herbarium section of the Biological Science Department, Ahmadu Bello University (ABU), Zaria, Nigeria. The herbarium specimen voucher number obtained was 2421.

2.2. Extraction and fractionation

The extraction and fractionation were conducted based on the procedures used by Kupchan et al. (1973) [29]. The fresh leaves of the *S. alata* were kept to dry under shade until a uniform was attained. The dried leaves were powdered using mortar and pestle into fine particles, packaged in a tightly closed container, and kept in a cool and dry environment. The Soxhlet apparatus exhaustively extracted one kilogram (1 kg) of the dried

powdered *S. alata* leaves with 2.5 liters of petroleum ether (60-80 grades). The marc of the petroleum ether extraction was then extracted with 2.5 liters absolute methanol with the aid of the Soxhlet apparatus. The marc of the methanolic extraction was dried and percolated with 1000 ml deionized water for 24-hours for the aqueous extraction. The extracts (petroleum ether, methanolic, and aqueous) were concentrated separately on a water bath set at 45 0C. The values for percentage yield of the extracts were calculated in relation to the formula below:

$$\text{Percentage yield (\%)} = \frac{\text{Weight of the crude extract (g)}}{\text{Weight of the powdered plant material (g)}} \times 100$$

One hundred grams (100 g) of the crude aqueous extract was suspended in water (500 ml) and successively partitioned with ethyl-acetate (5 × 500 ml) and diethyl ether (5 × 500 ml) to form the corresponding fractions. The ethyl-acetate and diethyl ether were evaporated to dryness while the residual aqueous was removed by placing the fraction on a water bath set at 50°C. The fractions were stored in a container separately before use. Each fraction was made into solutions with distilled water for each experiment. The percentage yield for each fraction was determined according to the formula above.

2.3. Phytochemical studies

Qualitative and quantitative phytochemical investigations were done on the crude aqueous leaves extract of *S. alata* to test the occurrence of phytochemical constituents in the extract according to the procedure outlined by Sofowora (1993) [30].

2.4. Test organism

The procedure previously used by Devillers et al. (1989) [32] was adopted. The methicillin-resistant *Staphylococcus aureus* (MRSA) was sub-cultured on sterile Nutrient Agar (NA) plates and incubated at 37 0C for 48-hours before each antibacterial testing.

2.5. Preparation of the test organism (MRSA)

The test organism used for the study was a standard strain of MRSA (ATCC33591) obtained from the Department of Veterinary Public Health, Ahmadu Bello University, Zaria, Kaduna, Nigeria.

2.6. Antibacterial screening of *Senna alata* leaves extracts and fractions against MRSA by Agar well diffusion method

The various extracts (petroleum ether, methanol and aqueous) and the aqueous fractions (ethyl acetate, diethyl ether and aqueous) of *S. alata* leaves were investigated for antibacterial effect against MRSA by agar well diffusion procedure as reported by Devillers et al. (1989) [32]. In this method, a 24-hour culture of MRSA was suspended in a sterile bottle that contained nutrient broth. Gradually, normal saline was added to obtain turbidity identical to Marcfland standard 0.5, which corresponds to about 108 cells/ml. This was then diluted to produce 106 cells/ml that were used in the experiment. Then 1 ml of the test organism (106 cells/ml) was inoculated into Petri plates (90 mm diameter) for the antibacterial susceptibility test. Then wells (6 mm diameter and 4 mm deep) were punched in the agar using a sterile cork borer. The wells were bored such that not to be closer than 15 mm to the edge of the plate and enough apart to overcome overlapping the inhibition zones. Some of the wells were filled with 25 mg/ml of either one extract or fractions, while some were filled up with one of the solvents as control. An adequate time was given to allow the material to diffuse considerably into each of the media. The plates were then turned upside down and kept in an incubator at 36 0C for 48-hours. Subsequently, the various inhibition zones were observed as an index of antibacterial activity.

2.6.1. Determination of the zone of inhibition

Following the incubation, the plates were checked for antibacterial activities. The compounds of the discs diffused through the medium and formed a concentration

gradient. Hence, the compounds of the wells formed in clear round zones of inhibition, which were measured with naked eyes using a transparent scale.

2.7. Thin-layer chromatographic analysis of diethyl ether fraction obtained from Senna alata crude aqueous leaves extract

The most active fraction (diethyl ether) of aqueous extract of the *S. alata* leaves was investigated for the active compound using thin-layer chromatography (TLC). The TLC analysis was done on pre-coated silica gel TLC plates (60 F254) and subsequently developed by two solvent systems of different polarities; n-butanol: acetic acid: water (6:1:2) and chloroform: acetic acid (9:1). The crude diethyl ether fraction was dissolved in methanol and spotted on the pre-coated G60 F254 TLC plates and developed in each solvent system. The various spots were detected using the following detecting agents; iodine vapor, ammonia vapor, ultraviolet (UV) light at 366 nm and UV in ammonia. The number of spots, colors, and retardation factors (R_f) values for each spot were calculated. The various spots of the crude diethyl ether fraction were investigated for activity against the MRSA by the agar overlay protocol

2.8. Antibacterial screening of various TLC spots obtained from diethyl ether fraction of Senna alata crude aqueous leaves extract against MRSA by Agar overlay method

Three (3) milliliters of nutrient broth in a sterile capped tube were inoculated with MRSA. The inoculated bottle was incubated in an incubator at 37 °C for 48-hrs. The developed TLC plates were aseptically placed in sterile Petri dishes. About 15 ml of melted top NA was measured using a pipette into three (3) different sterile capped bottles and allowed to cool to about 45 °C. To each bottle containing the melted NA, about 0.8 ml of the Nutrient broth containing MRSA was pipetted and vortexed to mix the test organism. The content of each bottle; melted top agar containing-MRSA was then poured onto each of the Petri dishes containing the developed TLC plates and control plates containing the unspotted TLC plates developed in the solvent system; chloroform: acetic acid (9:1). The plates were gently tilted back and forth to have an even distribution of the test organism, then allowed to gel fully and incubated at 37 °C for 48-hours. Subsequently, the various inhibition zones were measured as an index of antibacterial activity.

2.8.1. Determination of the zone of inhibition

Following the incubation, the plates were checked for antibacterial activities. The compounds of the well diffused through the medium and formed a concentration gradient. Thus, the compounds of the plates resulted in a clear zone of inhibition for a particular organism and were measured by naked eyes using a transparent scale.

2.8. Column chromatographic analysis of the diethyl ether fraction obtained from the Senna alata crude aqueous leaf extract

The most active fraction (diethyl ether) was chromatographed over a silica gel column (mesh size: 70-230) and successively eluted with solvents to increase polarity using n-hexane and ethyl-acetate. The eluates were then applied onto a pre-coated TLC card of silica gel 60G254 (thickness 0.1mm) plates and eluted with methanol: chloroform (4:1). [33] The eluates with the same R_f values were pooled together. The column eluted with n-hexane: ethyl acetate (40:60) afforded the most active eluate.

2.9. Gas chromatography-mass spectrometry of the active spot (Spot 2)

The TLC active spot (spot 2) was analyzed with a gas chromatography-mass spectrometry (GC-MS) (GC-MS-QP2010PLUS) analyzer. The data were obtained on an elite-1 (100% dimethyl polysiloxane) column (30 0.25 mm 1 μmdf). The carrier gas used was helium (99.99%) with a flow rate of 1ml/min in the split mode (10:1). An aliquot of 2 μl of ethanol solution of the sample was injected into the column with the injector temperature at 250°C. The gas chromatography (GC) oven temperature was started at 110°C and held

for 2min. The temperature was elevated to 200°C at the rate of 10°C/min without holding. Holding was allowed at 280°C for 9 min with a program rate of 5°C/min. The temperature of the injector and detector were set at 250°C and 280°C, respectively. Ion source temperature was maintained at 200°C. The mass spectrum of compounds in the samples was obtained by electron ionization at 70 eV, and the detector was operated in scan mode from 45-450 atomic mass units (amu). A scan interval of 0.5 seconds and fragments from 45 to 450 Da was maintained. The total running time was 36 minutes.

2.9. Infra-red spectroscopy of the active spot (spot 2)

The infra-red (IR) spectroscopic analysis of the active spot was carried out on the KBr disc using Fourier transform (FT) Spectrophotometer.

3. Results

3.1. Percentage yield of the *Senna alata* leaves extracts

The sequential extraction of one kilogram (1 kg) of the powdered *Senna alata* leaves with petroleum ether, methanol and water yielded 150.2 g, 104.03 g and 204.57 g representing an extractive value of 15.02%, 10.40% and 20.46%, respectively.

3.2. Percentage yield fractions per 100 g of *S. alata* crude aqueous leaves extract

The partitioning of 100 g of the *S. alata* crude aqueous extract with ethyl-acetate, diethyl ether, and the aqueous fraction yielded 1.68 g, 1.83 g, 2.45 g corresponding to 16.80%, 18.30% and 24.50% of the starting material, respectively.

3.3. Qualitative phytochemicals analysis of the aqueous leaves extract of *S. alata*

The qualitative phytochemical analysis of the crude aqueous leaves extract of *S. alata* indicated alkaloids, phenolic compounds, tannins, carbohydrates, reducing sugar, polyuronides, coumarins, saponins, steroids, triterpenoids, flavonoids, resins, cyanophores, cardiac glycosides and anthraquinones. However, carotenoids were absent.

3.4. Quantitative phytochemical analysis of the crude aqueous leaves extract of *S. alata*

The quantitative phytochemical investigation of the aqueous leaves extract of *S. alata*, showed that 10g of the plant material contained 1.27 g \pm 0.00 (12.7% \pm 0.01) of flavonoids, 0.64 g \pm 0.01 (12.8% \pm 0.01) of alkaloids, 0.004 g \pm 0.00 (0.9% \pm 0.00) of cyanogenic glycosides, 0.30 g \pm 0.02 (1.48% \pm 0.02) saponins and 0.28 g \pm 0.01 (9.6% \pm 0.01) of tannins.

3.5. Antibacterial activities of leaves extract of *S. alata*

The methanol and aqueous leaves extract of *S. alata* elicited remarkable inhibition against MRSA. However, the petroleum ether extract was not active against the bacterial isolate. The MRSA was resistant to the standard antibacterial agent used (oxacillin). The aqueous and methanol extracts exhibited the highest (14.5 mm \pm 0.01) and lowest (11mm \pm 0.11) zone of inhibition, respectively, against the test organism. The outcomes of the antibacterial screening of the petroleum ether, methanol and aqueous extracts of *S. alata* leaves against MRSA are shown in Table 1.

Table 1. Antibacterial activities of leaves extract of *Senna alata* against MRSA.

Extract	Concentration (mg/ml)	Zone of inhibition (mm)
Petroleum ether	25	0.00 \pm 0.00
Methanol	25	11.0 \pm 0.11
Aqueous	25	14.5 \pm 0.01
Oxacillin	25	0.00 \pm 0.00

The values are the mean of two measurements across each zone of inhibition and in duplicates measured in millimeters (mm). The zero values indicate no inhibition.

3.6. Antibacterial activities of fractions obtained from the crude aqueous extracts of *S. alata* on MRSA

The diethyl ether fraction inhibited the MRSA (14.00 ± 0.03 mm). However, the ethyl-acetate and aqueous fractions were ineffective against the bacterial isolate. The output of antibacterial effects of the ethyl-acetate, diethyl ether and the crude aqueous leaves extract of *S. alata* against MRSA are presented in Table 2.

Table 2. Antibacterial activities of fractions from the crude aqueous extract of *S. alata* on MRSA.

Fraction	Concentration (mg/ml)	Zone of inhibition (mm)
Ethyl-acetate	25	0.00 ± 0.00
Diethyl ether	25	14.00 ± 0.03
Aqueous	25	0.00 ± 0.00

The values are the mean of two measurements across each zone of inhibition and in duplicates measured in millimeters (mm). The zero values indicate no inhibition.

3.7. Antibacterial screening of the various silica gel column eluates of diethyl ether fraction obtained from the crude aqueous leaves extract of *S. alata* on MRSA

The result indicated that only sub-fraction 13 of the diethyl ether fraction obtained by eluting the column with n-hexane: ethyl acetate (40:60) produced antibacterial activity (15.00 mm ± 0.01) against the MRSA. The antibacterial effect of the various column eluates of the most active fraction (diethyl ether) is presented in Table 3.

Table 3. Antibacterial screening of the various silica gel columns eluates of the diethyl ether fraction obtained from the crude aqueous leaves extracts of *S. alata* on MRSA.

Sub-fraction	Concentration (mg/ml)	Zone of inhibition (mm)
1	200	0.00 ± 0.00
2	200	0.00 ± 0.00
3	200	0.00 ± 0.00
4	200	0.00 ± 0.00
5	200	0.00 ± 0.00
6	200	0.00 ± 0.00
7	200	0.00 ± 0.00
8	200	0.00 ± 0.00
9	200	0.00 ± 0.00
10	200	0.00 ± 0.00
11	200	0.00 ± 0.00
12	200	0.00 ± 0.00
13	200	15.00 ± 0.01
14	200	0.00 ± 0.00
15	200	0.00 ± 0.00

The values are the mean of two measurements across each zone of inhibition and in duplicates measured in millimeters (mm). The zero values indicate no inhibition.

3.8. Antibacterial screening of the various spots of sub-fraction 13 of the diethyl ether fraction obtained from aqueous leaves extract of *S. alata* on MRSA

The result of TLC analysis of the active column eluates (sub-fraction 13) revealed three (3) different spots when the plate was developed with solvent system chloroform: methanol (4:1). The antibacterial screening of the spots against the MRSA revealed that only spot 2 produced activity against the MRSA (15.00 ± 0.05 mm). The antibacterial activity of the various spot of sub-fraction 13 on the MRSA is shown in Table 4.

Table 4. Antibacterial Screening of the different spots of sub-fraction 13 obtained from the diethyl ether fraction of aqueous leaves extracts of *S. alata* on MRSA.

Spot	Concentration (mg/ml)	Zone of inhibition (mm)
1	100	0.00 ± 0.00

2	100	15.00 ± 0.05
3	100	0.00 ± 0.00

The values are the mean of two measurements across each zone of inhibition and in duplicates measured in millimeters (mm). The zero values indicate no inhibition.

3.9. Thin layer chromatographic studies of diethyl ether fraction obtained from crude aqueous leaves extract of *S. alata*

Different colors were observed on exposing the TLC plate of the active diethyl ether sub-fraction (fraction 13) of the crude aqueous leaves extract of *S. alata* to different detecting mediums. Under direct light, bright yellow (Spot 1) was observed. When this plate was exposed to ammonia vapor, two spots were observed: yellow (Spot 1) and pink (Spot 2). Nine yellow Spots were observed when the TLC plate was exposed to iodine vapor. These spots changed to various colors when observed under UV light in the presence and absence of ammonia vapor. Viewing the plate under UV light, in the absence of ammonia revealed 9 variedly colored spots: spot 1 purple (Rf=96), spot 2 blue (Rf=94), spot 3 light purple (Rf=80), spot 4 orange (Rf=73), spot 5 purple (Rf=66), spot 6 blue-green (Rf=45), spot 7 light purple (Rf=38), spot 8 purple (Rf=28) and spot 9 purple-green (Rf=19). However, when the plate was exposed to UV light, in the presence of ammonia, spot 1 changed from purple color to blue color, spot 3 changed to pink color, while spots 2, 4, 5, 6, 7, 8, 9 appeared as orange, blue-green, green, light purple, light purple, light blue and green respectively.

3.10. Rf values and color of fraction 13 of diethyl ether fraction obtained from crude aqueous leaves extract of *S. alata* when viewed under different detecting medium

Thin-layer chromatographic analysis of fraction 13 of the diethyl ether showed three spots with Rf values of 45, 54, and 93. All three spots turned yellow with iodine vapor. Spot 2 turned pink with ammonia vapor and appeared yellow under UV light (366 nm), while spot 3 turned light blue under the UV light, as shown in Table 5.

Table 5. Rf values and color of TLC Spots of fraction 13 obtained from diethyl ether column eluates of *S. alata* crude aqueous leaves extract when viewed under different detecting mediums.

Spots	R _f x 100	DL	Iodine vapor	Ammonia vapor	Ultra-violet (UV)	UV/Ammonia
1	45	ND	ND	LP	LY	LP
2	54	ND	Y	P	Y	P
3	93	ND	Y	ND	ND	B

Y: Yellow, P: Pink, LB: Light blue, ND: not detected, LB: Light pink.

3.11. Infra-red spectroscopic analysis of TLC spot 2 of the active column eluate of fraction 13 obtained from *S. alata* diethyl ether column eluate

The FT-IR spectrum of TLC Spot 2 from sub-fraction 13 of diethyl ether column eluates was obtained as KBr disk and shown in figure 1, which contains seven (7) vibrational bands assigned to their respective functional groups as shown in Table 10. The prominent bands are assignable to O-H (3320 cm⁻¹), C-H stretching vibration (2922 cm⁻¹), conjugated C=O stretching vibration (1697 cm⁻¹), conjugated C=C stretching vibration (1648 cm⁻¹), C-H bending vibration (1514 cm⁻¹), C-O stretching vibration (1033 cm⁻¹), and C-H out of plane bending vibration (670 cm⁻¹) bonded to a carbon-carbon double bond (C=C).




Figure 1. Infrared spectrum of Spot 4 of fraction 13 obtained from diethyl ether column eluates

Table 6. Infrared bands and assignments of the active TLC Spot (spot 2) of fraction 13 obtained from diethyl ether column eluates for *S. alata*.

Bands	Wave number (cm ⁻¹)	Assignment
1	3320	O-H Stretching
2	2922	C-H Stretching
3	1697	C=O Stretching
5	1514	C-H Bending
6	1033	C-O Stretching
7	670	C-H Bending
1	3320	O-H Stretching

3.12. Gas chromatography-mass spectrometry (GC-MS) of spot 2 obtained from column fraction 13 obtained from the *S. alata* diethyl ether column eluate

Total ion chromatogram (TIC) of TLC spot 2 of fraction 13 from the diethyl ether column eluates shown in Figure 2 indicated that the most prominent peak (peak 3) eluted at retention time 22.10 minutes belongs to TLC Spot 2 of fraction 13. The other peaks in the TIC correspond to impurities present in the eluate, which were insignificant except

peak 2. The electron impact mass spectrometry (EIMS) spectrum of spot 4 of sub-fraction 13 exhibited prominent peaks that are shown in Figure 3, which indicated a molecular ion peak at m/z 296[M]⁺ with the following significant fragment ion peaks at m/z 264 [C₁₉H₃₆]⁺, 222 [C₁₆H₃₀]⁺, 180 [C₁₃H₂₄]⁺, 137 [C₁₂H₁₇]⁺, 112 [C₈H₁₆]⁺, 98 [C₇H₁₂]⁺, 84 [C₆H₁₂]⁺, 83 [C₆H₁₁]⁺, 69 [C₅H₉]⁺, 55 [C₄H₇]⁺, 41 [C₃H₅]⁺ and 40 [C₃H₄]⁺. The fragment ion at m/z 264 corresponding to C₁₉H₃₆⁺ was formed from the loss of two oxygen atoms (2O) from the fragmentation of the molecular ion at m/z 296 [M]⁺. The fragment ion at m/z 84 [C₆H₁₂]⁺ indicated the loss of 180 mass [C₁₃H₂₄]⁺ ion. Fragment ion loss at m/z 137 is accompanied by a loss of a carbon-carbon double bond (C=C), while fragment ion at m/z 55 is the base peak ion.

4. Discussion

Plant-derived products play a crucial role in exploring active biomolecules to treat infectious diseases, currently causing serious health challenges globally, especially in developing nations [34]. Therefore, it is essential to intensify research on natural products to establish the secondary metabolites responsible for activity against infectious diseases [35,36]. The current study elucidates the antibacterial potentials of various extracts and bioactive components of the aqueous leaves extract of *Senna alata* against Methicillin-resistant *Staphylococcus aureus* (MRSA). A compound related to 9-octadecenoic acid methyl ester was found to have potent antibacterial activity against MRSA.

Based on the current investigation, the aqueous extract of the plant produced the highest extractive and could contain the highest amount of secondary metabolites present in the plant. Our findings are in line with the work of Saito et al. (2012) [38], where the 100g of milled *S. alata* leaves produced 25.4% aqueous extract. However, the work of Faruq et al. (2010) [40] has shown that the methanol extract produced the highest yield.

Interestingly, this investigation has shown that the crude aqueous extract of *S. alata* exerted the highest antibacterial effects against the bacteria. Other studies by Adedayo et al. (2001), Modi et al. (2012), Saito et al. (2012), Somchit et al. (2003) [25],[43],[38],[42] demonstrated the inhibitory effect of aqueous extract of *S. alata* against *S. aureus*, which corroborates with the outcome of this study. On the contrary, a previous report by Faruq et al. (2010) [40] has shown that the *S. aureus* was resistant to aqueous leaf extract of *S. alata*. In addition, the antibacterial activity elicited by the methanol extract of *S. alata* against the MRSA in this work agrees with the findings of Hazni et al. (2008) [45] and Sermakkani & Thangapandian (2012) [47]. The difference in the antibacterial activities produced by the methanol and aqueous extracts of *S. alata* leaves in the current work might be related to the secondary metabolites varying degrees of solubility in categories of solvents based on the polarity, which also corroborates with the higher extraction yield in the aqueous extract [48].

Bioassay-guided fractionation is an effective procedure to discover novel therapeutic agents via obtaining active fractions or isolated bioactive agents. In this method, each fraction is usually investigated for biological activity, and subsequently, the most active portions are eventually fractionated for further evaluation [49]. The chemical contents and potential mechanisms of activity of bioactive compounds usually vary in different parts of the plants, as well as a difference in the solubility of the secondary metabolites in different solvents [50]. In this study, the crude aqueous leaves extract of *S. alata* afforded ethyl-acetate and diethyl ether fractions. However, only the diethyl ether fraction elicited an antibacterial activity related to the higher amount of the bioactive agents in the fraction, as shown by its higher extractive value than the ethyl-acetate fraction.

Phytochemical determination gives an overview of the possible category of the plants' secondary metabolites and their quantity in a particular fraction which could guide the isolation strategy of the bioactive components.[51] Based on this research, the aqueous extract of *S. alata* leaves revealed saponins, alkaloids, emodin, tannins, steroids, anthraquinones, and flavonoids that could be associated with the antibacterial effects. The outcome agrees with the previous research by El-Mahmood & Doughari (2008) and Sule et

al. (2011) [54]. Generally, the anti-infective action of plants is related to the chemical compounds including tannins, phenols, saponins, steroids, alkaloids, flavonoids and many other compounds via various processes [55]. Polyphenol agents such as tannins, flavonoids, and alkaloids possess antimicrobial effects [56,57]. Flavonoids, including quercetin and kaempferol have antimicrobial actions by inhibiting the action of the bacterial enzymes [55,58,59]. The antimicrobial action of flavonoids is also associated with their capability to produce a complex with soluble, extracellular proteins as well as bacterial cell walls, whereas tannins inhibit microbial adhesions, transport proteins, and enzymes [60,61]. Besides, flavonoids destabilize microbial cell membranes [62]. Therefore, these phytochemical compounds in *S. alata* could be responsible for its antimicrobial activities.

The observation in the current work that only sub-fraction 13 of the column fractions of the diethyl ether fraction of the crude aqueous *S. alata* leaves extract showed activity concurs with the work of Faruq et al. (2010) [40], in which only one of the column fractions elicited antibacterial activity. Furthermore, a single spot was observed to possess bioactivity following the active diethyl ether column fraction to silica gel TLC analysis. The bioactive spot (Spot 2) of the sub-fraction 13 obtained from the diethyl ether fraction of the crude aqueous leaves extract of *S. alata* being the only spot effective against the test organism possesses a hydroxyl group (-OH) in its structure as shown in its IR spectrum assignable to the vibrational band at 3320 cm⁻¹, which is broad and of low intensity. This is typical of the -OH group found in fatty acids, and it belongs to -OH bonded to a carboxyl group carbon atom. The presence of the carboxyl group in the structure of spot 2 of the sub-fraction 13 has been confirmed by the GC-MS spectral data. The loss of two oxygen atoms (2O) from the fragmentation of the molecular ion at m/z 296 to form the fragment ion peak at m/z 264 indicated that this loss of oxygen atom emanates from a carboxyl group, which is typical of fatty acids methyl ester [63].

The FT-IR spectrum of spot 2 of sub-fraction 13 also showed a carbon-carbon double bond (C=C) due to the stretching vibration band at 1684 cm⁻¹ (Table 10). The presence of this bond was confirmed in the GC-MS spectral data because the fragmentation of the fragment ion at m/z 137 is accompanied by a loss of carbon-carbon double bond (C=C) to produce the fragment ion at m/z 112. The orientation of the C=C bond is cis configuration. This is confirmed by the presence of C-H out-of-plane bending vibration (670 cm⁻¹) of an olefinic bond since the bending vibration at 950-970 cm⁻¹ (characteristic of trans configuration) is absent in the IR spectrum of spot 2 of the sub-fraction 13. Furthermore, the C-H bending vibration at 1514 cm⁻¹ results from the presence of a terminal methyl (CH₃) group in the eluate. The GC-MS spectral data confirm this because the fragment ion at m/z 83 loses a methyl group and one hydrogen atom to form the fragment ion at m/z 69. This is a diagnostic of unsaturated fatty acid with a terminal methyl group [64].

In this work, the IR spectra data of spot 2 of the sub-fraction 13 agrees with that of cis-Octadecenoic acid methyl ester [65]. In addition, the entire fragment ions in the GC-MS spectra data of spot 2 of the active sub-fraction13, including the molecular ion and base peak ion, are in line with the data reported for methyl 9-octadecenoate by Yayli et al. (2001) [67]. Therefore, based on the data presented in this work and the literature, and by comparing with the library search result, spot 2 of the active sub-fraction 13 was determined to be 9-octadecenoic acid methyl ester (E). Besides, the molecular formula of the active spot 2 of the sub-fraction13 was deduced as (C₁₉H₃₅O₂) based on the information obtained from its molecular ion at m/z 296; this was supported by the findings of Igwe & Onwu (2015) [69] that the oleic acid was 75.03 of total essential oil of *S. alata* leaf extract using GC-MS. Besides, Rahman et al. (2006) [71] have shown that oleic acids are among the major fatty acids present in *S. alata*, confirming the presence of fatty acids in the *S. alata* leaves in the current work. Many researchers, including Ogunwande et al. (2010) [73], Adiana & Mazura (2011) [75] as well as Thenmozhi & Rajan (2015) [77], have previously used Fourier transform infrared (FT-IR) spectroscopy and GC-MS to analyze the various phytocompounds.

The antibacterial action of the active spot 2, which seems to be a fatty acid methyl ester based on the library search and spectral analysis, is in line with the findings of Igwe

& Onwu (2015) [69], who reported the antibacterial effects of essential oil from *S. alata* leaves extract against *S. aureus*. Similar reports by Kabara et al. (1972) [79], Knapp & Melly (1986) [81], Farrington et al. (1992) [83] and Sun et al. (2003) [85] have shown that long-chain unsaturated fatty acids, including oleic acids found naturally are effective bactericidal against pathogens including MRSA. In addition, Zheng et al. (2005) [87] in their work reported the effectiveness of unsaturated fatty acid esters and unsaturated fatty acids esters derivatives against *S. aureus* and MRSA. However, the precise mechanism of antibacterial action is precisely unknown. These reports support the observation made on the antibacterial activity of the identified compound against MRSA in the present study, which could explain the traditional applications of *S. alata* in treating bacterial infections.

5. Conclusions

The *Senna alata* leaves possess varying degrees of antibacterial activities against MRSA. The antimicrobial activity of the plant against MRSA was likely due to the 9-Octadecenoic acid methyl ester (E) component of the plant. In addition, 9-Octadecenoic acid methyl ester (E) could serve as a lead compound for further development to discover novel therapeutic agents as an alternative treatment for drug-resistant bacterial infections. Therefore, we recommend further spectroscopic studies like a proton (¹H) and Nuclear magnetic resonance (NMR) to ascertain the structure of the identified compound. Also, pharmacological and toxicity studies on the identified phytochemical compound should be conducted to establish the molecular basis of its antibacterial action and safety profile.

Author Contributions: Conceptualization, S.M.L., S.I., K.M.Y. and O.S.O.; methodology, Y.W., M.M., M.H.A. and S.M.; software, Y.W., M.M., M.H.A. and S.M.; validation, S.M.L., A.S. and Z.A.; formal analysis, S.M.L., Y.W., M.M., M.H.A. and S.M.; investigation, S.M.L., S.I., K.M.Y. and O.S.O.; resources, S.M.L., Y.W., M.M. and A.S.; data curation, S.A., M.H.A. and Z.A.; writing—original draft preparation, S.M.L., S.I., K.M.Y., O.S.O. and A.S.; writing—review and editing, Y.W., M.M., M.H.A., S.M. and Z.A.; supervision, S.I., K.M.Y. and O.S.O.; project administration, Y.W. M.M., M.H.A. and S.M.; funding acquisition, M.M. and A.S. All authors have read and agreed to the published version of the manuscript.

Funding: Please add: This research received no external funding.

Data Availability Statement: All data related to the manuscript are presented in the manuscript. The raw data can be obtained from the corresponding author based on reasonable request.

Acknowledgments: The authors are thankful to the entire academic and non-academic staff of the Department of Pharmacognosy and Drug Development, Ahmadu Bello University, Zaria, Kaduna, Nigeria, for their support during the conduct of the study. The USM Fellowship covers YW.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Etame, R. M. E.; Mouokeu, R. S.; Poundeu, F. S. M.; Voukeng, I. K.; Cidjeu, C. L. P.; Tiabou, A. T.; Yaya, A. J. G.; Ngane, A. N. R.; Kuiate, J. R.; Etoa, F. X. Effect of fractioning on antibacterial activity of n-butanol fraction from *Enantia chlorantha* stem bark methanol extract. *BMC Complementary and Alternative Medicine* 2019, 19, 1–7.
2. Nair, J. J.; Wilhelm, A.; Bonnet, S. L.; Staden, J. Van Antibacterial constituents of the plant family Amaryllidaceae. *Bioorganic & Medicinal Chemistry Letters* 2017, 27, 4943–4951.
3. Kale, P.; Dhawan, B. The changing face of community-acquired methicillin-resistant *Staphylococcus aureus*. *Indian Journal of Medical Microbiology* 2016, 34, 275–285.
4. David, M. Z.; Daum, R. S. Community-Associated Methicillin-Resistant *Staphylococcus aureus*: Epidemiology and Clinical Consequences of an Emerging Epidemic. *Clinical microbiology reviews* 2010, 23, 616–687.
5. Gibbons, S. Phytochemicals for Bacterial Resistance – Strengths, Weaknesses and Opportunities. *Planta medica* 2008, 74, 594–602.
6. Dadashi, M.; Nasiri, M. J.; Fallah, F.; Owlia, P.; Hajikhani, B.; Emameini, M.; Mirpour, M. Methicillin-Resistant *Staphylococcus aureus* (MRSA) in Iran: A Systematic Review and Meta-analysis. *Journal of global antimicrobial resistance* 2017, 12.

7. David, M. Z.; Daum, R. S. Community-Associated Methicillin-Resistant *Staphylococcus aureus*: Epidemiology and Clinical Consequences of an Emerging Epidemic. *Clinical microbiology reviews* 2010, 23, 616–687.
8. Dadashi, M.; Nasiri, M. J.; Fallah, F.; Owlia, P.; Hajikhani, B.; Emameini, M.; Mirpour, M. Methicillin-Resistant *Staphylococcus aureus* (MRSA) in Iran: A Systematic Review and Meta-analysis. *Journal of global antimicrobial resistance* 2017, 12.
9. Omoshaba, E. O.; Ojo, O. E.; Oyekunle, M. A.; Sonibare, A. O.; Adebayo, A. O. Methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from raw milk and nasal swabs of small ruminants in Abeokuta, Nigeria. *Tropical Animal Health and Production* 2020.
10. Xie, Y.; Yang, W.; Tang, F.; Chen, X.; Ren, L. Antibacterial Activities of Flavonoids: Structure- Activity Relationship and Mechanism. *Current Medicinal Chemistry* 2015, 22, 132–149.
11. Nair, J. J.; Wilhelm, A.; Bonnet, S. L.; Staden, J. Van Antibacterial constituents of the plant family Amaryllidaceae. *Bioorganic & Medicinal Chemistry Letters* 2017, 27, 4943–4951.
12. Ahmad, M. H.; Jatau, A. I.; Khalid, G. M.; Alshargi, O. Y. Traditional uses, phytochemistry, and pharmacological activities of *Cochlospermum tinctorium* A. Rich (Cochlospermaceae): a review. *Future Journal of Pharmaceutical Sciences* 2021, 7, 1–13.
13. Atanasov, A. G.; Waltenberger, B.; Pferschy-Wenzig, E. M.; Linder, T.; Wawrosch, C.; Uhrin, P.; Temml, V.; Wang, L.; Schwaiger, S.; Heiss, E. H.; et al. Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology Advances* 2015, 33, 1582–1614.
14. Slobodníková, L.; Fialová, S.; Rendeková, K.; Kovář, J. Antibiofilm Activity of Plant Polyphenols. *Molecules* 2016, 21, 1–15.
15. Barbieri, R.; Coppo, E.; Marchese, A.; Daglia, M.; Sobarzo-Sanchez, S.; Nabavi, S. F.; Nabavi, S. M. Phytochemicals for human disease: An update on plant-derived compounds antibacterial activity. *Microbiological Research* 2016, 196, 44–68.
- Bajalan, I.; Rouzbahani, R.; Pirbalout, A. G.; Maggi, F. Chemical composition and antibacterial activity of Iranian *Lavandula x hybrida*. *Chemistry & Biodiversity* 2017, 14, e1700064.
16. Gibbons, S. Phytochemicals for Bacterial Resistance – Strengths, Weaknesses and Opportunities. *Planta medica* 2008, 74, 594–602.
17. Barbieri, R.; Coppo, E.; Marchese, A.; Daglia, M.; Sobarzo-Sanchez, S.; Nabavi, S. F.; Nabavi, S. M. Phytochemicals for human disease: An update on plant-derived compounds antibacterial activity. *Microbiological Research* 2016, 196, 44–68.
18. Ahmad, M. H.; Zezi, A. U.; Bola, A. S.; Alhassan, Z.; Mohammed, M.; Danraka, R. N. Mechanisms of antidiarrhoeal activity of methanol leaf extract of *Combretum hypopilinum* diels (combretaceae): Involvement of opioidergic and (α 1 and β) -adren-ergic pathways. *Journal of Ethnopharmacology* 2020, 269, 113750.
19. Oladeji, O. S.; Adelowo, F. E.; Oluyori, A. P.; Bankole, D. T. Ethnobotanical Description and Biological Activities of *Senna alata*. *Evidence-based Complementary and Alternative Medicine* 2020, 2580259.
20. Oladeji, O. S.; Adelowo, F. E.; Oluyori, A. P.; Bankole, D. T. Ethnobotanical Description and Biological Activities of *Senna alata*. *Evidence-based Complementary and Alternative Medicine* 2020, 2580259.
21. Yakubu, M. T.; Musa, I. F. Effects of post-coital administration of alkaloids from *Senna alata* (Linn. Roxb) leaves on some fetal and maternal outcomes of pregnant rats. *Journal of reproduction & infertility* 2012, 13, 211–217.
22. Adedayo, O.; Anderson, W. A.; Snieckus, V.; Patil, P. A. Phytochemistry and Antibacterial Activity of *Senna alata* Flower. *Pharmaceutical Biology* 2001, 39, 408–412.
23. Yakubu, M. T.; Musa, I. F. Effects of post-coital administration of alkaloids from *Senna alata* (Linn. Roxb) leaves on some fetal and maternal outcomes of pregnant rats. *Journal of reproduction & infertility* 2012, 13, 211–217.
24. Adedayo, O.; Anderson, W. A.; Snieckus, V.; Patil, P. A. Phytochemistry and Antibacterial Activity of *Senna alata* Flower. *Pharmaceutical Biology* 2001, 39, 408–412.
25. Somchit, M. N.; Reezal, I.; Nur, I. E.; Mutalib, A. R. In vitro antimicrobial activity of ethanol and water extracts of *Cassia alata*. *Journal of Ethnopharmacology* 2003, 84, 1–4.
26. Igoli, J. O.; Igwue, I. C.; Igoli, N. P. Traditional medicinal practices among the Iggede people of Nigeria. *Journal of Herbs, Spices and Medicinal Plants* 2013, 10, 1–10.
27. Kupchan, S. M.; Britton, R. W.; Ziegler, M. F.; Sigel, C. W. Bruceantin, a new potent antileukemic simaroubolide from *Brucea antidysenterica*. *Journal of Organic Chemistry* 1973, 38, 178–179.
28. Kupchan, S. M.; Britton, R. W.; Ziegler, M. F.; Sigel, C. W. Bruceantin, a new potent antileukemic simaroubolide from *Brucea antidysenterica*. *Journal of Organic Chemistry* 1973, 38, 178–179.
29. Sofowora, A. *Medicinal Plants and Traditional Medicine in Africa*; 2nd Editio.; Spectrum Books Ltd: Ibadan, Nigeria, 1993.
30. Devillers, J.; Steiman, R.; Seigle-Murandi, F. The usefulness of the agar-well diffusion method for assessing chemical toxicity to bacteria and fungi. *Chemosphere* 1989, 19, 1693–1700.
31. Devillers, J.; Steiman, R.; Seigle-Murandi, F. The usefulness of the agar-well diffusion method for assessing chemical toxicity to bacteria and fungi. *Chemosphere* 1989, 19, 1693–1700.
32. Rahman, M. S.; Ali, M. Y.; Ali, M. U. In vitro screening of two flavonoids compounds isolated from *cassia alata* leaves for fungicidal activities. *Journal of biological sciences* 2008, 16, 139–142.
33. Cragg, G. M.; Newman, D. J. Biodiversity: A continuing source of novel drug leads *. *Pure and applied chemistry* 2005, 77, 7–24.
34. Voukeng, I. K.; Nganou, B. K.; Sandjo, L. P.; Celik, I.; Beng, Veronique, P.; Tane, P.; Kuete, V. Antibacterial activities of the methanol extract, fractions and compounds from *Elaeophorbia drupifera* (Thonn.) Stapf . *BMC Complementary and Alternative Medicine* 2017, 17, 1–9.

35. Etame, R. M. E.; Mouokeu, R. S.; Poundeu, F. S. M.; Voukeng, I. K.; Cidjeu, C. L. P.; Tiabou, A. T.; Yaya, A. J. G.; Ngane, A. N. R.; Kuiate, J. R.; Etoa, F. X. Effect of fractioning on antibacterial activity of n-butanol fraction from *Enantia chlorantha* stem bark methanol extract. *BMC Complementary and Alternative Medicine* 2019, 19, 1–7.
36. Saito, S. T.; Trentin, D. da S.; Macedo, A. J.; Pungartnik, C.; Gosmann, G.; Silveira, J. D. D.; Guecheva, T. N.; Henriques, J. A. P.; Brendel, M. Bioguided Fractionation Shows *Cassia alata* Extract to Inhibit *Staphylococcus epidermidis* and *Pseudomonas aeruginosa* Growth and Biofilm Formation. *Evidence-based Complementary and Alternative Medicine* 2012, 1–13.
37. Saito, S. T.; Trentin, D. da S.; Macedo, A. J.; Pungartnik, C.; Gosmann, G.; Silveira, J. D. D.; Guecheva, T. N.; Henriques, J. A. P.; Brendel, M. Bioguided Fractionation Shows *Cassia alata* Extract to Inhibit *Staphylococcus epidermidis* and *Pseudomonas aeruginosa* Growth and Biofilm Formation. *Evidence-based Complementary and Alternative Medicine* 2012, 1–13.
38. Faruq, Z. U.; Rahman, U. A.; Bello, M.; Obianke, M.; Atiku, F. A. Antibacterial activity of the active Component of *Cassia alata* (Linn) Leaves. *Nigerian Journal of Basic and Applied Research* 2010, 18, 97–100.
39. Faruq, Z. U.; Rahman, U. A.; Bello, M.; Obianke, M.; Atiku, F. A. Antibacterial activity of the active Component of *Cassia alata* (Linn) Leaves. *Nigerian Journal of Basic and Applied Research* 2010, 18, 97–100.
40. Modi, C.; Mody, S.; Patel, H.; Dudhatra, G.; Kumar, A.; Awale Madhavi Herbal antibacterials: a review. *Journal of Intercultural Ethnopharmacology* 2012, 1, 52–61.
41. Somchit, M. N.; Reezal, I.; Nur, I. E.; Mutalib, A. R. In vitro antimicrobial activity of ethanol and water extracts of *Cassia alata*. *Journal of Ethnopharmacology* 2003, 84, 1–4.
42. Modi, C.; Mody, S.; Patel, H.; Dudhatra, G.; Kumar, A.; Awale Madhavi Herbal antibacterials: a review. *Journal of Intercultural Ethnopharmacology* 2012, 1, 52–61.
43. Hazni, H.; Ahmad, N.; Hitotsuyanagi, Y.; Takeya, K.; Choo, C.-Y. Phytochemical Constituents from *Cassia alata* with Inhibition against Methicillin-Resistant *Staphylococcus aureus* (MRSA). *Planta Medica* 2008, 74, 1802–1805.
44. Hazni, H.; Ahmad, N.; Hitotsuyanagi, Y.; Takeya, K.; Choo, C.-Y. Phytochemical Constituents from *Cassia alata* with Inhibition against Methicillin-Resistant *Staphylococcus aureus* (MRSA). *Planta Medica* 2008, 74, 1802–1805.
45. Sermakkani, M.; Thangapandian, V. GC-MS Analysis of *cassia italica* Leaf Methanol Extract. *Asian Journal of Pharmaceutical and Clinical Research* 2012, 5, 90–94.
46. Sermakkani, M.; Thangapandian, V. GC-MS Analysis of *cassia italica* Leaf Methanol Extract. *Asian Journal of Pharmaceutical and Clinical Research* 2012, 5, 90–94.
47. Mabotja, M. B.; Venter, S. L.; Plooy, C. P. Du; Kudanga, T.; Amoo, S. O. Inhibitory and Antibacterial Activities of Spineless Cactus Pear Cultivars. *Plants* 2021, 10, 1–14.
48. Rufatto, C. L.; Luchtenberg, P.; Garcia, C.; Thomassigny, C.; Bouttier, S.; Dumas, F.; Henriques, J. A. P.; Roeschy-Ely, M.; Moura, S. Brazilian red propolis: Chemical composition and antibacterial activity determined using bioguided fractionation. *Microbiological Research* 2018, 74–82.
49. Lutfullah, G.; Shah, A.; Ahmad, K.; Haider, J. Phytochemical screening, antioxidant and antibacterial properties of *daphne mucronata*. *Journal of Traditional Chinese Medicine* 2019, 39, 764–771.
50. Bhatnagar, M.; Sarkar, N.; Gandharv, N.; Apang, O.; Singh, S.; Ghosal, S. Evaluation of antimycobacterial, leishmanicidal and antibacterial activity of three medicinal orchids of Arunachal Pradesh, India. *BMC Complementary and Alternative Medicine* 2017, 17, 1–10.
51. El-Mahmood, A. M.; Doughari, J. H. Phytochemical screening and antibacterial evaluation of the leaf and root extracts of *Cassia alata* Linn. *African Journal of Pharmacology and Pharmacognosy* 2008, 2, 124–129.
52. Sule, W. F.; Okonko, I. O.; Omo-Ogun, S.; Nwanze, J. C.; Ojezele, M. O.; Ojezele, O. J.; Alli, J. A.; Soyemi, E. T.; Olaonipekun, T. O. Phytochemical properties and in-vitro antifungal activity of *Senna alata* Linn. crude stem bark extract. *Journal of Medicinal Plants Research* 2011, 5, 176–183.
53. El-Mahmood, A. M.; Doughari, J. H. Phytochemical screening and antibacterial evaluation of the leaf and root extracts of *Cassia alata* Linn. *African Journal of Pharmacology and Pharmacognosy* 2008, 2, 124–129.
54. Sule, W. F.; Okonko, I. O.; Omo-Ogun, S.; Nwanze, J. C.; Ojezele, M. O.; Ojezele, O. J.; Alli, J. A.; Soyemi, E. T.; Olaonipekun, T. O. Phytochemical properties and in-vitro antifungal activity of *Senna alata* Linn. crude stem bark extract. *Journal of Medicinal Plants Research* 2011, 5, 176–183.
55. Mogana, R.; Adhikari, A.; Tzar, M. N.; Ramliza, R.; Wiart, C. Antibacterial activities of the extracts, fractions and isolated compounds from *Canarium patentinervium* Miq. against bacterial clinical isolates. *BMC Complementary and Alternative Medicine* 2020, 20, 1–11.
56. Vu, T. T.; Kim, H.; Tran, V. K.; Vu, H. D.; Hoang, T. X.; Han, W. J.; Choi, Y. H.; Jang, K. S.; Choi, G. J.; Kim, J. Antibacterial activity of tannins isolated from *Sapium baccatum* extract and use for control of tomato bacterial wilt. *PLoS ONE* 2017, 12, 1–12.
57. Cushnie, T. P. T.; Lamb, A. J. Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents* 2005, 26, 343–356.
58. Cushnie, T. P. T.; Lamb, A. J. Recent advances in understanding the antibacterial properties of flavonoids. *International Journal of Antimicrobial Agents* 2011, 38, 99–107.
59. Sharma, A.; Rawat, S.; Verma, R.; Vishwavaidyalaya, R. D. Identification of antibacterial component of some Indian medicinal plants against *Klebsiella pneumoniae*. *International Journal of Pharmacy and Pharmaceutical Sciences* 2010, 2, 123–127.

-
60. Mogana, R.; Adhikari, A.; Tzar, M. N.; Ramliza, R.; Wiart, C. Antibacterial activities of the extracts , fractions and isolated compounds from *Canarium patentinervium* Miq . against bacterial clinical isolates. *BMC Complementary and Alternative Medicine* 2020, 20, 1–11.
 61. Sharma, A.; Rawat, S.; Verma, R.; Vishwavidyalaya, R. D. Identification of antibacterial component of some Indian medicinal plants against *Klebsiella pneumoniae*. *International Journal of Pharmacy and Pharmaceutical Sciences* 2010, 2, 123–127.
 62. Oyugi, D. A.; Ayorinde, F. O.; Gugssa, A.; Allen, A.; Izevbigie, E. B.; Eribo, B.; Anderson, W. A. Biological activity and mass spectrometric analysis of *Vernonia amygdalina* fractions. *Journal of Bioscience and Technology* 2011, 2, 287–304.
 63. Oyugi, D. A.; Ayorinde, F. O.; Gugssa, A.; Allen, A.; Izevbigie, E. B.; Eribo, B.; Anderson, W. A. Biological activity and mass spectrometric analysis of *Vernonia amygdalina* fractions. *Journal of Bioscience and Technology* 2011, 2, 287–304.
 64. Kaneshiro, T.; Marr, A. G. Cis-9, 10-Methylene hexadecanoic acid from phospholipids of *Escherichia coli*. *The Journal of Biological Chemistry* 1961, 236, 2615–2619.
 65. Yayli, N.; Kiran, Z.; Seymen, H.; Genc, H. Characterization of Lipids and Fatty Acid Methyl Ester Contents in Leaves and Roots of *Crocus vallicola*. *Turk Journal of Chemistry* 2001, 25, 391–395.
 66. Yayli, N.; Kiran, Z.; Seymen, H.; Genc, H. Characterization of Lipids and Fatty Acid Methyl Ester Contents in Leaves and Roots of *Crocus vallicola*. *Turk Journal of Chemistry* 2001, 25, 391–395.
 67. Igwe, O. U.; Onwu, F. K. Leaf essential oil of *Senna alata* Linn from Southeast Nigeria and its Antimicrobial activity. *International Journal of Research in Pharmacy and Chemistry* 2015, 5, 27–33.
 68. Igwe, O. U.; Onwu, F. K. Leaf essential oil of *Senna alata* Linn from Southeast Nigeria and its Antimicrobial activity. *International Journal of Research in Pharmacy and Chemistry* 2015, 5, 27–33.
 69. Rahman, M. S.; Ali, M. Y.; Ali, M. U.; Moynul, A. J. M. Studies on the Lipid and Glyceride Compositions of *Cassia alata* Seed Oil. *Bangladesh Journal of Scientific and Industrial Research* 2006, 41, 83–88.
 70. Rahman, M. S.; Ali, M. Y.; Ali, M. U.; Moynul, A. J. M. Studies on the Lipid and Glyceride Compositions of *Cassia alata* Seed Oil. *Bangladesh Journal of Scientific and Industrial Research* 2006, 41, 83–88.
 71. Ogunwande, I. A.; Flamini, G.; Cioni, P. L.; Omikorede, O.; Azeez, R. A.; Ayodele, A. A.; Kamil, Y. O. Aromatic Plants growing in Nigeria: Essential Oil Constituents of *Cassia alata* (Linn.) Roxb . and *Helianthus annuus* L. *Record of Natural Products* 2010, 4, 211–217.
 72. Ogunwande, I. A.; Flamini, G.; Cioni, P. L.; Omikorede, O.; Azeez, R. A.; Ayodele, A. A.; Kamil, Y. O. Aromatic Plants growing in Nigeria: Essential Oil Constituents of *Cassia alata* (Linn.) Roxb . and *Helianthus annuus* L. *Record of Natural Products* 2010, 4, 211–217.
 73. Adiana, M. A.; Mazura, M. P. Study on *Senna alata* and its different extracts by Fourier transform infrared spectroscopy and two-dimensional correlation infrared spectroscopy. *Journal of Molecular Structure* 2011, 991, 84–91.
 74. Adiana, M. A.; Mazura, M. P. Study on *Senna alata* and its different extracts by Fourier transform infrared spectroscopy and two-dimensional correlation infrared spectroscopy. *Journal of Molecular Structure* 2011, 991, 84–91.
 75. Thenmozhi, S.; Rajan, S. GC-MS analysis of bioactive compounds in *Psidium guajava* leaves. *Journal of Pharmacognosy and Phytochemistry* 2015, 3, 162–166.
 76. Thenmozhi, S.; Rajan, S. GC-MS analysis of bioactive compounds in *Psidium guajava* leaves. *Journal of Pharmacognosy and Phytochemistry* 2015, 3, 162–166.
 77. Kabara, J. J.; Swieczkowski, D. M.; Conley, A. J.; Truant, J. P. Fatty Acids and Derivatives as Antimicrobial Agents Antimicrobial Agents. *Antimicrobial agents and chemotherapy* 1972, 2, 23–28.
 78. Kabara, J. J.; Swieczkowski, D. M.; Conley, A. J.; Truant, J. P. Fatty Acids and Derivatives as Antimicrobial Agents Antimicrobial Agents. *Antimicrobial agents and chemotherapy* 1972, 2, 23–28.
 79. Knapp, H. R.; Melly, M. A. Bactericidal Effects of Polyunsaturated Fatty Acids. *Journal of Infectious Diseases* 1986, 154, 84–94.
 80. Knapp, H. R.; Melly, M. A. Bactericidal Effects of Polyunsaturated Fatty Acids. *Journal of Infectious Diseases* 1986, 154, 84–94.
 81. Farrington, M.; Brenwald, N.; Haines, D.; Walpole, E. Resistance to desiccation and skin fatty acids in outbreak strains of methicillin-resistant *Staphylococcus aureus*. *Journal of medical microbiology* 1992, 36, 56–60.
 82. Farrington, M.; Brenwald, N.; Haines, D.; Walpole, E. Resistance to desiccation and skin fatty acids in outbreak strains of methicillin-resistant *Staphylococcus aureus*. *Journal of medical microbiology* 1992, 36, 56–60.
 83. Sun, C. Q.; Connor, C. J. O.; Robertson, A. M. Antibacterial actions of fatty acids and monoglycerides against *Helicobacter pylori*. *FEMS Immunology & Medical Microbiology* 2003, 36, 9–17.
 84. Sun, C. Q.; Connor, C. J. O.; Robertson, A. M. Antibacterial actions of fatty acids and monoglycerides against *Helicobacter pylori*. *FEMS Immunology & Medical Microbiology* 2003, 36, 9–17.
 85. Zheng, C. J.; Yoo, J.; Lee, T.; Cho, H.; Kim, Y.-H.; Kim, W.-G. Fatty acid synthesis is a target for antibacterial activity of unsaturated fatty acids. *FEBS letters* 2005, 579, 5157–5162.
 86. Zheng, C. J.; Yoo, J.; Lee, T.; Cho, H.; Kim, Y.-H.; Kim, W.-G. Fatty acid synthesis is a target for antibacterial activity of unsaturated fatty acids. *FEBS letters* 2005, 579, 5157–5162.