
Review

Advances on Greener Asymmetric Synthesis of Antiviral Drugs via Organocatalysis

Everton M. da Silva¹, Hérica D. A. Vidal¹ and Arlene G. Corrêa^{1,*}

¹ Centre of Excellence for Research on Sustainable Chemistry, Department of Chemistry, Federal University of São Carlos, 13565-905, São Carlos, SP, Brazil

* Correspondence: agcorrea@ufscar.br

Abstract: Viral infections inflict many serious human diseases, being responsible for remarkably high mortality rates. In this sense, both the academy and the pharmaceutical industry are continuously searching for new compounds with antiviral activity, and in addition, face the challenge of developing greener and more efficient methods to synthesize these compounds. This becomes even more important with drugs possessing stereogenic centers as highly enantioselective processes are required. In this minireview, the advances achieved to improve synthetic routes efficiency and sustainability of important commercially antiviral chiral drugs are discussed, highlighting the use of organocatalytic methods.

Keywords: antivirals; green chemistry; selectivity; asymmetric synthesis; organocatalysis

1. Introduction

The increasing prevalence of diverse microbial infections as well as the emergence and re-emergence of viral epidemics with high morbidity and mortality rates are a major public health threat. Despite the persistent production of antiviral drugs and vaccines in the global market, viruses remain one of the leading causes of deadly human diseases. Effective control of viral diseases, caused particularly by Zika, dengue, herpes simplex, human immunodeficiency (HIV), and Ebola virus, remain promising goals amidst the mutating viral strains [1].

In the last decades, it is estimated that an average of 3 million deaths per year worldwide are caused by different viral pathogens, in special hepatitis, AIDS, and influenza [2]. More recently, the coronavirus disease (COVID-19) caused by SARS-CoV-2 is creating tremendous human suffering, and, to date, no effective drug is available to directly treat the disease. According to the World Health Organization, globally, between January 2020 and September 2021, there have been more than 234 million confirmed cases of COVID-19, and 4.8 million deaths [3].

In this sense, both the academy and the pharmaceutical industry are continuously searching for new compounds with antiviral activity [4]. Emphasis should focus on orally available drugs for outpatient use, if possible, and on identifying combination therapies that combat viral and immune-mediated pathologies, extend the effectiveness of therapeutic windows, and reduce drug resistance. As the emergence of viral diseases is expected to

accelerate, proactive programs to develop broadly active family-specific and cross-family antiviral therapeutics will be key to prepare for future disease [5,6].

Chiral molecules have played a prominent role in the development of new drugs, as individual stereoisomers may exhibit marked differences in pharmacodynamic, pharmacokinetic, and toxicological properties [7]. Therefore, there is currently considerable interest in fully characterizing and examining both enantiomers in the early stages of new drug development [8]. In this regard, the pharmaceutical industries either exploit the chiral switching practice from already marketed racemates or develop *de novo* enantiomerically pure compounds [9].

One important step in the therapeutical is the synthesis of drug candidates under development, their critical intermediates as well as the active pharmaceutical ingredients (APIs), since it encompasses the search for routes with high efficacy, good yields, and less impact in the environment [10]. Furthermore, the synthesis of APIs requires large-scale analysis of the process, thus, using less toxic reagents, greener solvents, more efficient methods, and waste reduction is essential for the sustainability of the production [11].

The three pillars of asymmetric catalysis are bio-, metal- and organocatalysis [12]. The biological catalysts, such as isolated enzymes or whole cells, are increasingly used on an industrial scale and are particularly favored for hydrolytic reactions [13]. The rapid development of organometallic chemistry and catalysis has made numerous breakthrough contributions to organic synthesis; however, such widespread use of metal compounds has highlighted the significance of environmental and toxicity issues [14]. In this perspective, a recent trend in the chemical science and pharmaceutical industry is the dismissal of catalytic metal complexes and their replacement with organocatalysts. The efficacy and synthetic versatility of asymmetric organocatalysis [15,¹⁶] and the use of a plethora of highly effective small-molecule organocatalysts have enriched the field of organic synthesis, including chiral proline derivatives, *N*-heterocyclic carbenes, chiral thioureas, and Brønsted acids as well as phase-transfer catalysts (PTC), such as the quaternary ammonium salts derived from cinchona alkaloids [17].

Moreover, in the context of green and sustainable syntheses, continuous flow chemistry offers exceptional opportunities to accelerate, integrate, simplify, scale-up and automatize chemical reactions. Early this year, Ötvös and Kappe have reviewed the flow chemistry-based approaches for the synthesis of chiral APIs and their advanced stereogenic intermediates, covering the utilization of biocatalysis, organometallic catalysis and organocatalysis [18].

A significant number of antiviral agents used in clinical practice are amino acids, short peptides, or peptidomimetics, and among them are several HIV protease inhibitors (e. g. lopinavir, atazanavir) [19], HCV protease inhibitors (e. g. grazoprevir, glecaprevir) [20], and HCV NS5A protein inhibitors (e. g. daclatasvir, pibrentasvir) [21]. Recently, Skwarecki et al. [22] reviewed the synthesis of these antiviral agents with special attention in the synthetic aspects of non-proteinogenic amino acid components.

In Table 1, we present selected chiral antiviral drugs, being the anti-HIV agents the major examples [23], highlighting the asymmetric strategy employed as key step to achieve their total synthesis. It is worth noting that most of the synthetic routes uses chiral pools, including amino acids, as starting materials (entries 3, 4, 6, 12, 13, 15-17). Chiral resolution using either enzymes or metal catalysts has also been successfully employed (entries 1, 2, 8, 9, 14) and only on few cases enantioselective organocatalyzed methods have been employed (entries 10 and 11).






In this minireview, the advances achieved to improve synthetic routes efficiency and sustainability of important commercially antiviral chiral drugs are discussed. The organocatalytic methods were selected to be presented in more detailed and the proposed mechanism pathways for the introduction of the stereogenic centers are highlighted.







2. Synthesis of Efavirenz







Efavirenz (**1**) is a reverse transcriptase inhibitor drug used in the treatment of infections caused by the human immunodeficiency virus (HIV) type I. Approved in 1998 by the FDA, it has been used in combination with other drugs for the treatment of AIDS [24].


Pierce and co-workers [25] reported in 1998 an enantioselective synthesis of efavirenz in nine steps with 49% overall yield. The key step involves an enantioselective addition of the cyclopropylacetylide **5** to trifluoromethyl ketone **3** in the presence of the additive (1*R*,2*S*)-*N*-pyrrolidinylnorephedrine (**4**) (Scheme 1). Initially, 4-chloroaniline (**2**) was transformed in the ketoaniline **3** with 75% yield over 6 steps. The enantioselective alkynylation occurred with the formation of the chiral complex, obtained by the reaction of a mixture of chiral ligand **4** and cyclopropyl acetylene (**5**) with *n*-butyllithium at -10 to 0 °C in a THF-toluene-hexane mixture, then it was cooled below -50 °C and the ketoaniline was added. The enantioselectivity of this transformation is dependent on the *N*-protecting group of compound **3**. The mixture of acetylide alkoxide generated at low temperature exists as a complex mixture of species. A stable aggregate has been fully characterized as the 2:2 symmetric cubic tetramer, and it was observed that optimal reaction conversion and enantioselectivity require strict control of reagent stoichiometry and reaction conditions. The amino alcohol **7** was obtained with 91% yield and 95.5% *ee*, then was converted into benzoxazinone and the deprotection occurs by ceric ammonium nitrate (CAN) giving efavirenz (**1**) in 72% yield. This route is only suitable for small-scale synthesis since an equal amount of *p*-methoxy benzaldehyde is formed and not efficiently removed upon crystallization, and a significant quantity of cerium salt was generated as a waste, being a concern for the scale-up. Although the chiral ligand **4** was employed in equimolar amount, it could be recovered in 98% yield and recycled.

Table 1. Selected chiral antiviral drugs

Entry	API	Chemical structure	Treatment or indications and mode of action	Enantioselective synthesis: chiral pool or key step
1	Amprenavir		AIDS, HIV protease inhibitor	a) Co(salen) catalyzed kinetic resolution of 2-(1-azido-2-phenylethyl)oxirane [26] b) from <i>L</i> -acid malic and <i>L</i> -phenylalanine [27]
2	Abacavir		AIDS and hepatitis B virus (HBV), reverse transcriptase inhibitor	a) chiral resolution using γ -lactamase [28] b) Oppolzer-sultam as auxiliary then asymmetric rhodium-catalyzed coupling using Josiphos 003-1 [29]
3	Cobicistat		AIDS, human CYP3A proteins inhibitor	chiral amino acids as starting materials [30,31]
4	Daclatasvir		chronic hepatitis C virus (HCV) genotype 3, inhibitor of the NS5A	continuous flow synthesis with chiral amino acids as starting materials [32]
5	Dolutegravir		AIDS, integrase strand transfer inhibitor	(<i>R</i>)-3-amino-1-butanol as starting material [33]


6	Elvitegravir		AIDS, HIV integrase inhibitor	(S)-valinol as starting material [34]
7	Emtricitabine		AIDS and HBV, reverse transcriptase inhibitor	L-menthol as chiral auxiliary [35]
8	Indinavir		AIDS, HIV-1 protease inhibitor	a) (R)-glycidol as starting material [36] b) from indanone via enantioselective ester hydrolysis with <i>Rhizopus oryzae</i> [37] c) (1S,2R)-cis-aminoindanol as starting material [38]
9	Lamivudine		AIDS and HBV, reverse transcriptase inhibitor	a) enzymatic dynamic kinetic resolution [39] b) (S)-lactic acid as starting material [40] c) L-menthol as chiral auxiliary [41]
10	Maraviroc		AIDS, antagonist of the CCR5 receptor	a) (S)-tert-butanefulfinamide as chiral auxiliary [42] b) aza-Michael/hemiacetal reaction using the Hayashi-Jørgensen organocatalyst [43]
11	Nelfinavir		AIDS, HIV protease inhibitor	a) Ti catalyzed asymmetric aminolysis of epoxide [44] b) starting from chiral sulfoxide [45] c) bromocyclization of bisallylic catalyzed by (S)-BINAP [46]

12	Ritonavir		AIDS, HIV protease inhibitor	chiral amino acids as starting materials [47,48]
13	Saquinavir		AIDS, HIV protease inhibitor	chiral amino acids as starting materials [49]
14	Simeprevir		HCV, NS3/4A protease inhibitor	a) chiral resolution using pig liver esterase [50] b) chiral resolution using cinchonidine [51]
15	Telaprevir		HCV, NS3/4A protease inhibitor	lipase mediated desymmetrization of 1,2-cyclopentanedimethanol and chiral amino acids [52]
16	Tenofovir alafenamide		AIDS and HBV, nucleotide reverse transcriptase inhibitor	(R)-9-[2-(phosphonomethoxy) propyl]adenine (PMPA) [53] from <i>L</i> -threonine and <i>L</i> -alanine as starting materials [54]
17	Valacyclovir		Herpes, prodrug converted to acyclovir, a DNA polymerase inhibitor	from <i>L</i> -valine [55]



Scheme 1. First total synthesis of efavirenz (**1**).


In 2011, Shibata and co-workers [56] reported the first synthesis of efavirenz based on a metal-free, enantioselective trifluoromethylation of alkynyl ketone (**8**) with trimethyl(trifluoromethyl)-silane (Me_3SiCF_3). This transformation is different from the previously reported methods, because an organocatalytic based on a cinchonidine derivative **10** was used, however with only 50% *ee*. The enantioselectivity was improved (up to 80% *ee*) in 2014 by the same group [57], using a quinine-based catalyst **11** bearing a butyloxy group (Scheme 2). Nevertheless, the organocatalytic process was still less efficient than the organometallic enantioselectivity approach. Thus, in 2016, Shibata and co-workers [58] reported the use of a quaternary ammonium salt of cinchona alkaloids having an alkynyl group (**12**). Structurally, the best catalyst used has only the different ethynyl groups, so the 3D conformation of the transition state (TS) is similar to the others. However, the catalyst has a closed structure between the ethynyl group and the quinoline ring, and the alkenyl ketones can be stabilized by a π - π interaction with the quinoline ring, and the methoxy group helps in positioning the ketones by a steric interaction. Some modifications in the conditions of this reaction were necessary, but the trifluoromethylation using a novel catalyst occurred with 88% yield and 93% *ee*. With the trifluoro methylated compound in hand, the synthesis of efavirenz was completed in 2 steps. Reduction of the nitro group to aniline was obtained in 83% yield and 91% *ee*. The aniline was treated with 4-nitrophenyl chloroformate in presence of KHCO_3 to obtain efavirenz (**1**) in 89% yield. Using *n*-hexane/ CH_2Cl_2 by single recrystallization, the enantiopurity was increased to 99%.



Scheme 2. Organocatalyzed trifluoromethylation of alkynyl ketone 8.**3. Synthesis of Oseltamivir**

Tamiflu, the phosphate of (-)-oseltamivir (**13**), is the most efficient drug for the treatment and prevention of flu. The mechanism of action fundamentally involves inhibition of the neuraminidase enzyme of type A and type B viruses [59].


The first total synthesis of the phosphate of (-)-oseltamivir was described in 1998 by Rohloff and co-workers (Scheme 3) [60]. The synthesis is based on two natural products, (-)-quinic acid (**14**) and (-)-shikimic acid (**15**), which makes it difficult to be employed on an industrial scale quickly and cheaply [61]. The two routes pass for the same intermediate, the 3,4-pentyline ketal **16**. In the route of the quinic acid (**14**), the 3,4-pentyline ketal **16** was formed in 4 steps, in 29% yield, and in the route of (-)-shikimic acid (**15**), the 3,4-pentyline ketal **16** was obtained in 3 steps in 80% yield. The 3,4-pentyline ketal **16** was transformed in pentyl ether, and converted in a epoxide which was converted in the azido alcohol, and finally in the aziridine **17** with 50% yield over 4 steps. The aziridine was converted in the azido amine that was directly acylated. Finally, the azide was reduced to amine and the salt of oseltamivir.H₃PO₄ (**13**) was isolated after 4 steps with 26% overall yield. The final product was obtained in 11 steps and 10.4% yield from (-)-shikimic acid (**15**) and in 12 steps in 3.7% yield from (-)-quinic acid (**14**).



Scheme 3. First total synthesis of phosphate of (-)-oseltamivir (**13**).


The syntheses based on the natural products presented low yields, in addition to numerous reaction steps that required several purification processes. With this in mind, in 2009, Ishikawa and co-workers [62] reported a high-yielding synthesis of oseltamivir via three one-pot operations. The chiral cyclohexane was built based on organocatalyzed reaction and the process minimized chemical waste generation and time-consuming on the purifications. Diphenylprolinol silyl ether (**20**) was an effective organocatalyst, promoting an asymmetric reaction with excellent enantioselectivities. In the first step, the organocatalyst provides the Michael adduct with alkoxyaldehyde (**18**) and nitroalkane (**19**) in a quantitative yield and 96% *ee* (Scheme 4). Then, a Michael reaction between the nitroalkane and vinyl phosphonate **21** occurs, and the formed phosphonate undergoes an intramolecular Horner–Wardsworth–Emmons reaction with the formyl group to generate ethyl cyclohexenecarboxylate. Treatment of the mixture with *p*-toluenethiol in the presence of Cs₂CO₃ gives the product **22** in 70% yield. With compound **22** in hands, occurs the deprotection, followed by the conversion of carboxylic acid in

a acyl azide **23** in 3 steps. This azide is submitted to a domino Curtius rearrangement, followed by reduction of the NO₂ to the amine. A retro-Michael reaction of the thiol afforded oseltamivir, which was obtained in 82% yield from compound **22** after purification by acid/base extraction. In summary, the efficiency of the Hayashi-Jørgensen organocatalyst was described in the total enantioselective synthesis of (-)-oseltamivir, that was carried out in 9 reactions, in three one-pot operations, and just one purification by chromatographic column with 57% overall yield.



Scheme 4. Asymmetric organocatalyzed synthesis of oseltamivir (**13**).

Although this route was very efficient, the reaction time was quite long. Thus, based on previous reports published in the literature [62,63], Hayashi and Ogasawara described in 2016 a time economical total synthesis of (-)-oseltamivir in 60 minutes (Scheme 5) [64]. In the first step, the Michael reaction of nitroalkene **24** and α -alkoxyaldehyde **18** proceeds in the presence of diphenylprolinol silyl ether **25** and Schreiner's thiourea **26** affording product **27** that was not isolated. The organocatalyst was key for the generation of enamine, and the thiourea activated the nitroalkene via hydrogen bonding. The cyclohexene was formed via a domino Michael and Horner-Wadsworth-Emmons reactions. Then, after protonation with TMSCl, epimerization using TBAF and reduction of nitro group in presence of Zn, the desired compound **13** was obtained after purification by column chromatography in 15% yield, via a one-pot procedure. In summary, the total synthesis of oseltamivir was described in a single reactor over five steps, being the route optimized in not only yield and selectivity but also in reaction time.




Scheme 5. Time economy synthesis of oseltamivir (13).

4. Synthesis of Zanamivir

Zanamivir (28) is used in the treatment of influenza type A and B, and was approved in the United States in 1999 [65]. Zanamivir inactivates the influenza viral neuraminidase, an enzyme responsible for cleaving sialic acid residues in newly formed viruses. Such inhibition results in virus aggregation on the host cell surface, which limits the extent of infection and accelerates recovery from the disease [66].

The first scalable synthesis of zanamivir was described by Chandler and co-workers [67]. The route started with acetylneuraminic acid (29), which was converted into a methyl ester, and the hydroxy groups were acetylated (Scheme 6). Using TMSOTf, the oxazoline was formed, and selectively converted in the azide 30. The acetate protecting group was removed by a catalytic amount of NaOMe to gain water solubility, following hydrolysis of the methyl ester with aqueous TEA to give the triethylammonium salt. This salt was hydrogenated in presence of Lindlar catalyst to give the amine. Then, addition to the aminoiminomethanesulfonic acid with the amino acid, the guanidine functionality was introduced. The crystalline zanamivir (28) was isolated by ion-exchange chromatography. Several synthetic steps were likely to be performed on a gram scale. Most of the intermediates were isolated by recrystallization, however, in one case, desalting and freeze-drying are also necessary, and the use of the ion-exchange chromatography to obtain the pure product. The overall yield of the 9-step synthesis was only 8.3%.




Scheme 6. First scalable synthesis of zanamivir.

Due to the high cost of sialic acid, and considering the need for production on a global scale, in 2014, Tian and co-workers [68] developed a total synthesis of zanamivir, in 13 linear steps,

starting from *D*-araboascorbic acid (Scheme 7). Initially, the (*Z*)-*tert*-butyl (2-nitro vinyl)carbamate (**34**) was prepared in 3 steps from nitromethane with 72% yield. The organocatalyzed Michael addition of compound **32** in acetone was investigated, and the best conditions were achieved using thiourea catalyst **33** in toluene and benzoic acid as an additive. The reaction produced compound **34** in 72% yield and 98% *ee* after recrystallization of the crude product from ethyl acetate/petroleum ether (1:10).

With compound **34** in hand, the Henry reaction with aldehyde **35** was investigated. The aldehyde **35** was synthesized from *D*-araboascorbic acid [69]. Based on previous reports for *anti*-selective addition, a combination of CuBr₂ and the proline-derived catalyst **37** was used, and after spontaneous cyclization and subsequent dehydration with thionyl chloride and pyridine, selective formation of the *anti*-product **36** was observed with 60% yield, together with the formation of the undesired product in 8% yield. By observing the possible transition states, the volume of the protective group interferes with the reaction, which proceeds favorably by the proposed transition state, due to the repulsion between the protective group and the carbamate side chain (the only formation of the non-selective product was observed when the TBS was used). Therefore, smaller protection groups (MOM and Me) were used, thus favoring the formation of the *anti*-selective product. Then, the hydroxyl protecting groups of **36** were changed in order to obtain the carboxylic acid via oxidation of the methyl group. The protecting groups were all removed with HCl and finally, after guanidination with **38**, lead to the formation of zanamivir with single recrystallization. The overall yield was 18% and the synthetic route offers advantages due to the cost of the starting materials, being a cheap and accessible alternative for the manufacture of this anti-influenza drug.




Scheme 7. Total synthesis of zanamivir starting from *D*-araboascorbic acid.

5. Synthesis of Letemovir

Letermovir (**39**) is a drug approved by the FDA in 2017 for the treatment of infections caused by the human cytomegalovirus (HCMV) [70]. Severe, acute HCMV infections most commonly involve symptomatic infection of the gastrointestinal tract, liver, and central nervous system; hematological manifestations; and eye, lung, or arterial or venous thrombosis. The viral terminase complex is highly specific to herpesviruses, since no cellular protein carrying out its function has been identified in mammalian cells, making this viral complex a good target for antivirals. Letermovir targets pUL56 of the HCMV terminase complex [71].

Humphrey and co-workers [72] proposed an asymmetric synthesis for obtaining **39** with high enantioselectivity (Scheme 8). The total synthesis of letermovir (**39**) was carried out in eight steps from commercially available starting materials, and the key step was an aza-Michael reaction catalyzed by a cinchonidine-based phase transfer catalyst (PTC) **40** producing about 1t of product. The authors suggest some factors that interfere in enantioselectivities such as stirring rate, concentration, and equivalents of K_3PO_4 aqueous solution, and the counterions Br^- and PO_4^{3-} show low results. In terms of Green Chemistry, the method shows an excellent enantiomeric excess (99.6% *ee*) using a cinchonidine-based organocatalytic under mild conditions with an overall yield > 60% in eight steps, providing letermovir (**39**) on a large scale. Based on previous work [73], the TS was proposed by theoretical studies showing that the use of the catalyst promotes (*S*)-stereodiscrimination in the ring closure as a result of the interaction between catalyst/substrate, hydrogen bonding (O-H), substrate conformation, π -stacking interactions, and interactions with the benzyl- CF_3 group of the catalyst.




Scheme 8. Synthesis of letermovir (**39**) using organocatalysts **40** or **44**.

Futhermore, the same group reported the used of chiral bistriflamides **44** as organocatalyst to promote the enantioselective cyclization for the synthesis of letermovir (**39**). According to the mechanism pathway, supported by density functional theory (DFT) calculations and kinetics studies, the bistriflamide **44** acts as a Brønsted acid (Scheme 8). Initially, using **42** occurs the protonation of nitrogen by bistriflamide **44**, followed by tautomerization and cyclization to produce guanidine **43**. The catalyst promotes a double donation of hydrogens, so the TS is formed by the tautomerization of the starting material. This method showed high yields (96%) and a good enantiomeric ratio (84:16 *er*), being applied in large-scale for the synthesis of letermovir (**39**), demonstrating efficiency with reduced drawbacks when compared with the PTC-catalysed method.

6. Synthesis of Ruxolitinib

Ruxolitinib (**45**) was the first drug used in the treatment of myelofibrosis and polycythemia vera (PCV), approved by the FDA in 2011. Ruxolitinib acts specifically in the inhibition of protein tyrosine janus kinases (JAK) 1 and 2, promoting a decrease in inflammation and an inhibition of cellular proliferation [74].


One of the key parts of the structure of this drug is the chiral pyrazole ring, which defines stereochemistry. Lin and co-workers [75] developed the synthesis of ruxolitinib via aza-Michael addition, based on studies by Jørgensen [76] with proline-derived organocatalysts. According to the proposed mechanism, the addition by the *Re* face is favored with 94% *ee*, while the *Si* face is protected by the catalyst. This methodology presents two possibilities, both using cyclopentanecarbaldehyde (**46**) as starting material. In the first route (Scheme 9A), the aldehyde **46** reacted with compound **49** catalyzed by **47**, furnishing *R*-**51** in 80% yield and 90% *ee*. In the next two steps, **45** is obtained. In the second synthetic route (Scheme 9B), compound **46** reacted with 4-bromo-1H-pyrazole (**50**) catalyzed by **48**, and the product *R*-**52** is obtained with 85% yield and 84% *ee*. The next steps are conversion of the aldehyde in nitrile and **45** is obtained with 53% yields. In both routes, the TS was proposed based on Jørgensen's study where an addition on the *Re* face is favored, so in reactions using the more steric hindered organocatalysts, higher enantioselectivities were observed. The authors do not present information on the recovery of the catalysts nor perform tests on large scales, although the presented methodologies have good prospects in terms of green chemistry through the use of organocatalysts and an efficient asymmetric route.



Scheme 9. Synthesis of ruxolitinib (**45**) using a proline-derived organocatalyst.

Hayld and co-workers described the synthesis of ruxolitinib (**45**) [77] based on strategies developed by their group on the enantioselectivity of pyrazoles using a rhodium/JoSPOPhos (**53** and **54**) catalyst system (Scheme 10), with a high atom economy. Initially, with 4-bromopyrazole (**56**) and cyclopentylalene (**55**) in the rhodium-catalyzed coupling, it was possible to obtain the product **57** in the gram scale, with 95% yield and 90% *ee*. Then, after five steps, **45** is obtained with 53% overall yield. The authors do not present detailed information about the mechanism, it is

only shown that upon tautomerism of the pyrazole ring, and the use of **54** it is possible to obtain only the desired product, and also that the use of pyridinium *p*-toluenesulfonate (PPTS) plays a fundamental role in selectivity.




Scheme 10. Rhodium-catalyzed asymmetric synthesis of ruxolitinib (**45**).

7. Synthesis of Remdesivir

Remdesivir (**58**), a prodrug of the parent nucleoside GS441524 [78], was initially developed for the treatment of Ebola, however, due to the pandemic caused by SARS-CoV-2 after clinical tests, was approved in 2020 by the FDA to be used in severe cases of the Covid-19 [79]. The active form of remdesivir acts as a nucleoside analog and inhibits the RNA-dependent RNA polymerase (RdRp) of coronaviruses [80].

The main challenge in the synthesis of remdesivir is to establish the stereogenic center in the phosphorus atom [81,82]. The first- and second-generation syntheses of remdesivir (**58**) are shown in Scheme 11. In both synthetic routes, the same starting material **61** is employed, which is submitted to phosphorylation with the formation of compounds **62** and **63**, respectively. After the coupling with nucleobases (**59** or **60**), in the first generation, the two isomers formed are separated using chiral HPLC [83]. In the second-generation synthesis occurs a selective nucleophilic displacement, where there is no racemic production after phosphorylation, so only the *S* enantiomer is obtained upon recrystallization, using hydrochloric acid. Although it is possible to obtain the enantioenriched product **58**, the phosphorylation occurs with the lowest yield, so improvements were needed in this step of the synthesis.

Wang and co-workers [84] developed a synthesis of remdesivir based on the use of imidazole organocatalysts. A kinetic study was carried out evaluating the velocity constants (*k*) (Scheme 12), where it was observed that the key point of using the catalyst is that it can promote the *k*₁/*k*₂ racemization and that it is enantioselective in the formation of product **58**. The first test used compounds **62** and **60** without any catalyst, observing only traces of the product with a diastereoisomeric ratio (*dr*) of 1.1:1 (*Sp:Rp*), then achiral catalysts were used and *N*-methyl imidazole (NMI) presented the best result with 1.5:1 *dr*. Continuing the study, chiral bicyclic imidazole catalysts, modified with benzyl, acetyl ester, *tert*-butyl carbonate, and carbamates were used, and the catalyst **64** modified with adamantinyl group showed the best results with a ratio 11.2:1 *dr*. In addition, 2,6-lutidine (**65**) was used as a base at low temperature (-40°C), resulting in an increase of the diastereoselectivity to 21.8:1 and that the catalyst concentration does not interfere in the enantioselectivity, in cases of using a lower concentration, a longer reaction time is necessary.




Scheme 11. First- and second-generation synthesis of remdesivir (**58**).


In the mechanistic proposal, it was initially suggested a transition state involving a simultaneous action of phosphoryl chloride and nucleoside by the catalyst. Theoretical studies were carried out and showed that the Sp configuration was favored in 3.5 kcal. mol, in accordance with the experimental data. Finally, they explored this method in large-scale reaction (10 g), employing 10 mol% catalysts **64** at 40°C , and the products were obtained with a ratio of 21.2:1 *dr*, the Sp enantiomer was obtained with 89% yield and > 99:1 *dr* (85% yield by recrystallization and 4% yield of residual mother liquor by preparative HPLC). The authors emphasize the atom economy of the method and synthetic efficiency. Moreover, the catalyst was recovered for later use without any damage, it could be reused in subsequent cycles, and the high yield obtained in the gram scale reaction shows its potential application in the industry.

In 2021, Gannedi and co-workers [85] studied a strategy for obtaining the stereogenic center of the phosphorus atom via a one-pot synthesis also employing an imidazole-derived catalyst (Scheme 13). Several bicyclic chiral imidazole derivatives were tested, but catalyst **67** with phenyl group showed the best result, with 25:1 *dr* and 97% yield. After being defined that catalyst **67** was the most effective, further optimization was carried out in terms of reaction time, temperature, and catalyst loading. The authors concluded that -20°C for 24 h was ideal, since the increase in reaction time caused a decrease in catalytic activity, and reduction in temperature to -40°C promoted a discrepant increase in selectivity, and a decrease in temperature to -78°C no reaction was observed. Concerning catalyst **67**, the decrease in percentage promoted a decrease in selectivity, and extrapolation in the amount of 100% molar catalyst promoted a 42:1 *dr*, however, to avoid further purification of diester, the catalyst loading of 20 mol% was maintained. Some tests to increase the reaction scale were carried out, thus on the 1g scale, after the first step with the substrates **62** and **60**, catalyst **67**, *p*-TSA, and methanol were added in the same flask

reaction (one-pot) at room temperature, resulting in the removal of the isopropylidene group and affording a mixture of **58** with its diastereoisomer, which was purified by recrystallization with 73% yield and 99.6:0.6 *dr*. On the 10 g scale, 70% yield and 99.3:0.7 *dr* of the desired product were obtained and the catalyst was recovered in 83% yield.



Scheme 12. Synthesis of remdesevir (**58**) using the imidazole organocatalyst **64**.



Scheme 13. One-pot synthesis of remdesevir (**58**) using the imidazole-derived catalyst **67**.

The reaction pathway was inspired by the mechanism of histidine-dependent glucose 6-hydrolysis catalyzed by glucose 6-phosphate phosphatase (Scheme 13). It can be seen that the imidazole residues of His119 and His176 may act as a proton donor and receptor, based on this that the catalyst **67** was proposed with similar chemical structure. Furthermore, it was observed that the catalyst configuration directly influences the product stereochemistry, that is, a catalyst

with (S)-configuration favored the (S)-product. The proposed transition state was based on Wang's study [83], which uses similar imidazole organocatalysts.

8. Conclusions

Viral infections inflict many serious human diseases, being responsible for remarkably high mortality rates. In this sense, both the academy and the pharmaceutical industry are continuously searching for new compounds with antiviral activity, and in addition, face the challenge of developing greener and more efficient methods to synthesize these compounds. In this minireview, the developments in asymmetric synthesis of antivirals based on green methods were summarized. The approaches using organocatalysts in the key steps were emphasized including the proposed mechanism to these enantioselective transformations. Although the yield and enantioselectivity of these reactions are in general good to excellent, the catalyst loading is still high (5-10 mol%) and recovery and recycle require tedious steps of purification. Moreover, one-pot reactions, which reduce the generation of waste from purification steps, as well as continuous flow synthesis are paving the way for scale-up. These methods are specially important for developing countries, enabling local drug manufacturing with the consequence of improving access to medicines.

Author Contributions: conceptualization, A.G.C.; investigation, E.M.S. and H.D.A.V.; writing—original draft preparation, E.M.S. and H.D.A.V.; writing—review and editing, A.G.C.; funding acquisition, A.G.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by FAPESP (grants 2013/07600-3 and 2014/50249-8), GlaxoSmithKline, CAPES (Finance Code 001), and CNPq (grants 429748/2018-3 and 302140/2019-0).

Data Availability Statement: Data sharing not applicable. No new data were created or analyzed in this study.

Conflicts of Interest: The authors declare no conflict of interest.

ORCID

Arlene G. Corrêa <https://orcid.org/0000-0003-4247-2228>

References

1. Sagaya, J.R.; Khusro, A.; Agastian, P.; Alfarhan, A.; Al-Dhabi, N.A.; Arasu, M.V.; Rajagopal, R.; Barcelo, D.; Al-Tamini, A. Emerging paradigms of viral diseases and paramount role of natural resources as antiviral agents. *Sci. Total Environ.* **2021**, *759*, 143539.
2. Ma, Y.; Frutos-Beltran, E.; Kang, D.; Pannecouque, C.; Clercq, E.D.; Menendez-Arias, L.; Liu, X.; Zhan, P. Medicinal chemistry strategies for discovering antiviral effective against drug-resistant viruses. *Chem. Soc. Rev.* **2021**, *50*, 4514-4540.
3. WHO coronavirus (COVID-19) Dashboard. World Health Organization, 2021. Available in: <https://covid19.who.int>, accessed on October 5, 2021.
4. Clercq, E.D. Antivirals and antiviral strategies. *Nat. Rev. Microbiol.* **2004**, *2*, 704-720.
5. Megank, R.M.; Baric, R.S. Developing therapeutic approaches for twenty-first-century emerging infectious viral diseases. *Nat. Med.* **2021**, *27*, 401-410.

6. Zhong, Y.L.; Yasuda, N.; Li, H.; McLaughlin, M.; Tschäen, D. Process Chemistry in Antiviral Research. *Top. Curr. Chem.* **2016**, *374*, 77.
7. Nguyen, L.A.; He, H., Pham-Huy, C. Chiral Drugs. An Overview. *Int. J. Biomed. Sci.* **2006**, *2*, 85-100.
8. Abram, M.; Jakubiec, M.; Kaminski, K. Chirality as an Important Factor for the Development of New Antiepileptic Drugs. *ChemMedChem.* **2019**, *14*, 1744-1761.
9. Calcaterra, A.; D'Acquarica, I. The market of chiral drugs: Chiral switches versus de novo enantiomerically pure compounds. *J. Pharmac. Biomed. Anal.* **2018**, *147*, 323-340.
10. Eastgate, M.D.; Schmidt, M.A.; Fandrick, K.R. On the design of complex drug candidate syntheses in the pharmaceutical industry. *Nat. Rev. Chem.* **2017**, *1*, 0016.
11. Rogers, L.; Jensen, K.F. Continuous manufacturing – the Green Chemistry promise?. *Green Chem.* **2019**, *21*, 3481-3498.
12. Reisinger, C.M.; Pan, S.C.; List, B. New Concepts for Catalysis. *Organocatalysis* **2008**, *2*, 35-64.
13. Patel, R.N. Biocatalysis for synthesis of pharmaceuticals. *Bioorg. Med. Chem.* **2018**, *26*, 1252-1274.
14. Egorova, K.S.; Ananikov, V.P. Toxicity of Metal Compounds: Knowledge and Myths, *Organometallics* **2017**, *36*, 4071-4090.
15. Han, B.; He, X.H.; Liu, Y.Q.; He, G.; Peng, C.; Li, J.L. Asymmetric organocatalysis: an enabling technology for medicinal chemistry. *Chem. Soc. Rev.* **2021**, *50*, 1522-1586.
16. Moreira, N. M.; Martelli, L. S. R.; Corrêa, A. G. Asymmetric organocatalyzed synthesis of coumarin derivatives. *Beilstein J. Org. Chem.*, **2021**, *17*, 1952-1980.
17. Holland, M.C.; Gilmour, R. Deconstructing covalent organocatalysis. *Angew. Chem. Int. Ed.* **2015**, *54*, 3862-3871.
18. Otvos, S.B.; Kappe, C.O. Continuous flow asymmetric synthesis of chiral active pharmaceutical ingredients and their advanced intermediates. *Green Chem.* **2021**, *23*, 6117-6138.
19. Ghosh, A.K.; Bilcer, G.; Schiltz, G. Syntheses of FDA Approved HIV Protease Inhibitors. *Synthesis* **2001**, 2203-2229.
20. Ganta, N.M.; Gedda, G.; Rathnakar, B.; Satyanarayana, M.; Yamajala, B.; Ahsan, M.J.; Jadav, S.S.; Balaraju, T. A review on HCV inhibitors: Significance of non-structural polypeptides. *Eur. J. Med. Chem.* **2019**, *164*, 576-601.
21. Voight, E.A.; Greszler, S.N.; Hartung, J.; Ji, J.G.; Klix, R.C.; Randolph, J.T.; Shelat, B.H.; Waters, J.E.; DeGoey, D.A. Desymmetrization of pibrentasvir for efficient prodrug synthesis. *Chem. Sci.* **2021**, *12*, 10076-10082.
22. Skwarecki, A.S.; Nowak, M.G.; Milewska, M.J. Amino Acid and Peptide-Based Antiviral Agents. *ChemMedChem*. DOI 10.1002/cmdc.202100397
23. Mandala, D.; Thompson, W.A.; Watts, P. Synthesis routes to anti-HIV drugs. *Tetrahedron* **2016**, *72*, 3389-3420.
24. Adkins, J.C.; Noble, S. Efavirenz. *Drugs.* **1998**, *56*, 1055-1064.
25. Pierce, M.E.; Parsons, R.L.; Radesca, L.; Lo, Y.S.; Silverman, S.; Moore, J.R.; Islam, O.; Choudhury, A.; Fortunak, J.M.D.; Nguyen, D.; Luo, C.; Morgan, S.J.; Davis, W.P.; Confalone, P.N.; Chen, C.; Tillyer, R.D.; Frey, L.; Tan, L.; Xu, F.; Zhao, D.; Thompson, A.S.; Corley, E.G.; Grabowski, E.J.J.; Reamer, R.; Reider, P.J. Synthesis of Efavirenz via Asymmetric Alkynylation. *J. Org. Chem.* **1998**, *63*, 8536-8543.
26. Gadakh, S.K.; Reddy, R.S.; Sudalai, A. Enantioselective synthesis of HIV protease inhibitor amprenavir via Co-catalyzed HKR of 2-(1-azido-2-phenylethyl)oxirane. *Tetrahedron Asymmetry.* **2012**; *23*:898-903.
27. Honda Y, Katayama S, Kojima M, Suzuki T, Kishibata N, Izawa K. New approaches to the industrial synthesis of HIV protease inhibitors. *Org. Biomol. Chem.* **2004**, *2*, 2061-2070.

28. Evans, C.T.; Roberts, S.M.; Shoberu, K.A.; Sutherland, A.G. Potential use of carbocyclic nucleosides for the treatment of AIDS, chemoenzymatic synthesis of the enantiomers of carbovir. *J. Chem. Soc. Perkin Trans 1*. **1992**, 589-592.
29. Thieme, N.; Breit, B. Enantioselective and Regiodivergent Addition of Purines to Terminal Allenes: Synthesis of Abacavir. *Angew. Chem. Int. Ed.* **2017**, *56*, 1520-1524.
30. Xu, L.; Liu, H.; Murray, B.P.; Callebaut, C.; Lee, M.S.; Hong, A.; Strickley, R.G.; Tsai, L.K.; Stray, K.M.; Wang, Y.; Rhodes, G.R.; Desai, M.C. Cobicistat (GS-9350): A Potent and Selective Inhibitor of Human CYP3A as a Novel Pharmacoenhancer. *ACS Med. Chem. Lett.* **2010**, *1*, 209-213.
31. Xu, L.H.; Liu, H.T.; Hong, A.; Vivian, R.; Murray, B.P.; Callebaut, C.; Choi, Y.C.; Lee, M.S.; Chau, J.; Tsai, L.K.; Stray, K.M.; Strickley, R.G.; Wang, J.H.; Tong, L.; Swaminathan, S.; Rhodes, G.R.; Desai, M.C. Structure-activity relationships of diamine inhibitors of cytochrome P450 (CYP) 3A as novel pharmacoenhancers. Part II: P2/P3 region and discovery of cobicistat (GS-9350). *Bioorg. Med. Chem. Lett.* **2014**, *24*, 995-999.
32. Rana, A.; Mahajan, B.; Ghosh, S.; Srihari, P.; Singh, A.K. Integrated multi-step continuous flow synthesis of daclatasvir without intermediate purification and solvent exchange. *React. Chem. Eng.* **2020**, *5*, 2109-2114.
33. Sankareswaran, S.; Mannam, M.; Chakka, V.; Mandapati, S.R.; Kumar, P. Identification and Control of Critical Process Impurities: An Improved Process for the Preparation of Dolutegravir Sodium. *Org. Process Res. Dev.* **2016**, *20*, 1461-1468.
34. Radl, S.; Stach, J.; Pisa, O.; Cinibulk, J.; Havlicek, J.; Zajicova, M.; Pekareka, T. An Improved Synthesis of Elvitegravir. *J. Heteroc. Chem.* **2016**, *53*, 1738-1749.
35. Mandala, D.; Watts, P. An Improved Synthesis of Lamivudine and Emtricitabine. *ChemistrySelect* **2017**, *2*, 1102-1105.
36. Askin, D.; Eng, K.K.; Rossen, K.; Purick, R.M.; Wells, K.M.; Volante, R.P.; Reider, P.J. Highly diastereoselective reaction of a chiral, non-racemic amide enolate with (S)-glycidyl tosylate. Synthesis of the orally active HIV-1 protease inhibitor L-735,524. *Tetrahedron Lett.* **1994**, *35*, 673-676.
37. Demir, A.S.; Hamamci, H.; Doganel, F.; Ozgul, E. Chemoenzymatic synthesis of (1S,2R)-1-amino-2-indanol, a key intermediate of HIV protease inhibitor, indinavir. *J. Mol. Catal. B Enzym.* **2000**, *9*, 157-161.
38. Cheng, Y.; Lu, Z.; Chapman, K.T.; Tata, J.R. Solid Phase Synthesis of Indinavir and Its Analogues. *J. Comb. Chem.* **2000**, *2*, 445-446.
39. Hu, L.; Schaufelberger, F.; Zhang, Y.; Ramstrom, O. Efficient asymmetric synthesis of lamivudine via enzymatic dynamic kinetic resolution. *Chem. Commun.* **2013**, *49*, 10376-10378.
40. Snead, D.R.; Mcquade, D.T.; Ahmad, S.; Krack, R.; Stringham, R.W.; Burns, J.M.; Abdiaj, I.; Gopalsamuthiram, V.; Nelson, R.C.; Gupton, B.F. An Economical Route to Lamivudine Featuring a Novel Strategy for Stereospecific Assembly. *Org. Process Res. Dev.* **2020**, *24*, 1194-1198.
41. Aher, U.P.; Srivastava, D.; Jadhav, H.S.; Singh, G.P.; Jayashree, B.S.; Shenoy, G.G. Large-Scale Stereoselective Synthesis of 1,3-Oxathiolane Nucleoside, Lamivudine, via ZrCl₄-Mediated N-Glycosylation. *Org. Process Res. Dev.* **2020**, *24*, 387-397.
42. Zhu, Y.J.; Li, H.Y.; Lin, K.L.; Wang, B.; Zhou, W.C. A novel and efficient asymmetric synthesis of anti-HIV drug maraviroc. *Synth. Commun.* **2019**, *49*, 1721-1728.
43. Zhao, G.L.; Lin, S.; Korotvicka, A.; Deiana, L.; Kullberg, M.; Cordova, A. Asymmetric Synthesis of Maraviroc (UK-427,857). *Adv. Synth. Catal.* **2010**, *352*, 2291-2298.
44. Inaba, T.; Yamada, Y.; Abe, H.; Sagawa, S.; Cho, H. *J. Org. Chem.* **2000**, *65*, 1623-1628.
45. Raghavan, S.; Krishnaiah, V.; Sridhar, B. Asymmetric Synthesis of the Potent HIV-Protease Inhibitor, Nelfinavir. *J. Org. Chem.* **2010**, *75*, 498-501.

46. Nagao, Y.; Hisanaga, T.; Utsumi, T.; Egami, H.; Kawato, Y.; Hamashima, Y. Enantioselective Synthesis of Nelfinavir via Asymmetric Bromocyclization of Bisallylic Amide. *J. Org. Chem.* **2018**, *83*, 7290–7295.
47. Kempf, D.J.; Sham, .H.L.; Marsh, K.C.; Flentge, C.A.; Betebenner, D.; Green, B.E.; McDonald, E.; Vasavanonda, S.; Saldivar, A.; Wideburg, N.E.; Kati, W.M.; Ruiz, L.; Zhao, C.; Fino, L.; Patterson, J.; Molla, A.; Plattner, J.J.; Norbeck, D.W. Discovery of Ritonavir, a Potent Inhibitor of HIV Protease with High Oral Bioavailability and Clinical Efficacy. *J. Med. Chem.* **1998**, *41*, 602–617.
48. Ramu, E.; Rao, B.V. A short approach to the synthesis of the ritonavir and lopinavir core and its C-3 epimer via cross metathesis. *Tetrahedron Asymmetry*. **2009**, *20*, 2201–2204.
49. Göbring, W.; Gokbale, S.; Hilpert, H.; Roessler, F.; Schlageter, M.; Vogt, P. Synthesis of the HIV-Proteinase Inhibitor Saquinavir: A Challenge for Process Research. *Chimia*. **1996**, *50*, 532–537.
50. Rosenquist, A.; Samuelsson, B.; Johansson, P.O.; Cummings, M.D.; Lenz, O.; Raboisson, P.; Simmen, K.; Vendeville, S.; de Kock, H.; Nilsson, M.; Horvath, A.; Kalmeijer, R.; de la Rosa, G.; Beumont-Mauviel, M. Discovery and Development of Simeprevir (TMC435), a HCV NS3/4A Protease Inhibitor. *J. Med. Chem.* **2014**, *57*, 1673–1693.
51. Horváth, A.; Depré, D.; Vermeulen, W.A.A.; Wuyts, S.L.; Harutyunyan, S.R.; Binot, G.; Cuypers, J.; Couck, W.; Van Den Heuvel, D. Ring-Closing Metathesis on Commercial Scale: Synthesis of HCV Protease Inhibitor Simeprevir. *J. Org. Chem.* **2019**, *84*, 4932–4939.
52. Moni, L.; Banfi, L.; Basso, A.; Carcone, L.; Rasparini, M.; Riva, R. Ugi and Passerini Reactions of Biocatalytically Derived Chiral Aldehydes: Application to the Synthesis of Bicyclic Pyrrolidines and of Antiviral Agent Telaprevir. *J. Org. Chem.* **2015**, *80*, 3411–3428.
53. Derstine, B.P.; Tomlin, J.W.; Peck, C.L.; Dietz, J.P.; Herrera, B.T.; Cardoso, F.S.P.; Paymode, D.J.; Yue, A.C.; Arduengo, A.J.; Opatz, T. An Efficient Synthesis of Tenofovir (PMPA): A Key Intermediate Leading to Tenofovir-Based HIV Medicines. *Org. Process Res. Dev.* **2020**, *24*, 1420–1427.
54. Yang, B.; Xie, H.M.; Rani, K.R.; Gant, Y.J. Efficient Synthesis and Resolution of Tenofovir Alafenamide. *Lett. Org. Chem.* **2018**, *15*, 10–14.
55. Vetukuri, P.R.V.N.K.V.; Vedantham, R.; Mathad, V.T.; Padi, P.R.; Ramasamy, V.A. A Concise Route to Valacyclovir Hydrochloride. *Helv. Chim. Acta*. **2011**, *94*, 592–596.
56. Kawai, H.; Kitayama, T.; Tokunaga, E.; Shibata, N. A New Synthetic Approach to Efavirenz through Enantioselective Trifluoromethylation by Using the Ruppert–Prakash Reagent. *Eur. J. Org. Chem.* **2011**, 5959–5961.
57. Okuso, S.; Kawai, H.; Yasuda, Y.; Sugita, Y.; Kitayama, T.; Tokunaga, E.; Shibata, N. Asymmetric Synthesis of Efavirenz via Organocatalyzed Enantioselective Trifluoromethylation. *Asian J. Org. Chem.* **2014**, *3*, 449–452.
58. Okuso, S.; Hirano, K.; Yasuda, Y.; Tanaka, J.; Tokunaga, E.; Fukaya, H.; Shibata, N. Alkynyl Cinchona Catalysts affect Enantioselective Trifluoromethylation for Efavirenz under Metal-Free Conditions. *Org. Lett.* **2016**, *18*, 5568–5571.
59. Silva, F.C. Síntese Total do (-)-Oseltamivir (Tamiflu®) por Reações do Tipo Dominó. *Rev. Virt. Quim.* **2009**, *1*, 87–90.
60. Rohloff, J.C.; Kent, K.M.; Postich, M.J.; Becker, M.W.; Chapman, H.H.; Kelly, D.E.; Lew, W.; Louie, M.S.; McGee, L.R.; Prisbe, E.J.; Schlitz, L.M.; Yu, R.H.; Zhang, L. Practical Total Synthesis of the Anti-Influenza Drug GS-4104. *J. Org. Chem.* **1998**, *63*, 4545–4550.
61. Abrecht, S.; Harrington, P.; Iding, H.; Karpf, M.; Trussardi, R.; Wirz, B.; Zutter, U. The Synthetic Development of the Anti-Influenza Neuraminidase Inhibitor Oseltamivir Phosphate (Tamiflu®): A Challenge for Synthesis & Process Research. *Chimia*. **2004**, *58*, 621–629.

-
62. Ishikawa, H.; Suzuki, T.; Hayashi, Y. High-Yielding Synthesis of the Anti-Influenza Neuramidase Inhibitor (-)-Oseltamivir by Three "One-Pot" Operations. *Angew. Chem. Int. Ed.* **2009**, *48*, 1304–1307.
 63. Ishikawa, H.; Suzuki, T.; Orita, H.; Uchamaru, T.; Hayashi, Y. High-Yielding Synthesis of the Anti-Influenza Neuraminidase Inhibitor (-)-Oseltamivir by Two "One-Pot" Sequences. *Chem. Eur. J.* **2010**, *16*, 12616–12626.
 64. Hayashi, Y.; Ogasawara, S. Time Economical Total Synthesis of (-)-Oseltamivir. *Org. Lett.* **2016**, *18*, 3426–3429.
 65. Chapple, K.J.; Hendrick, A.E.; McCarthy, M.W. Zanamivir in the treatment and prevention of influenza. *Ann. Pharmacother.* **2000**, *34*, 798–801.
 66. Ryan, D.M.; Ticehurst, J.; Dempsey, M.H.; Penn, C.R. Inhibition of influenza virus replication in mice by GG167 (4-guanidino-2,4-dideoxy-2,3-dehydro-N-acetylneuraminic acid) is consistent with extracellular activity of viral neuraminidase (sialidase). *Antimicrob. Agents Chemother.* **1994**, *38*, 2270–2275.
 67. Chandler, M.; Bamford, M.J.; Conroy, R.; Lamont, B.; Patel, B.; Patel, V.K.; Steeples, I.P.; Storer, R.; Weir, N.G.; Wright, M.; Williamson, C. Synthesis of the potent influenza neuraminidase inhibitor 4-guanidino Neu5Ac2en. X-Ray molecular structure of 5-acetamido-4-amino-2,6-anhydro-3,4,5-trideoxy-D-erythro-L-glucosonic acid. *J. Chem. Soc.* **1995**, 1173–1180.
 68. Tian, J.; Zhong, J.; Li, Y.; Ma, D. Organocatalytic and Scalable Synthesis of the Anti-Influenza Drugs Zanamivir, Laninamivir, and CS-8958. *Angew. Chem. Int. Ed.* **2014**, *53*, 1–5.
 69. André, C.; Bolte, J.; Demuynck, C. Syntheses of l-threose and d-erythrose analogues modified at position 2. *Tetrahedron Asymmetry.* **1998**, *9*, 1359–1367.
 70. Humprey, G.R.; Dalby, S.M.; Andreani, T.; Xiang, B.; Luzung, M.R.; Song, Z.J.; Shevlin, M.; Christensen, M.; Belyk, K.M.; Tschaen, D.M. Synthesis of Letemovir by an Asymmetric Aza-Michael Reaction. *Org. Process Res. Dev.* **2016**, *20*, 1097–1103.
 71. Krishna, B.A.; Wills, M.R.; Sinclair, J.H. Advances in the treatment of cytomegalovirus. *Brit. Med. Bull.* **2019**, *131*, 5–7.
 72. Chung, C.K.; Liu, Z.; Lexa, K.; Andreani, T.; Xu, Y.; Ji, Y.; DiRocco, D.A.; Humphrey, G.R.; Ruck, R.T. Asymmetric Hydrogen Bonding Catalysis for the Synthesis of Dihydroquinazoline-containing Antiviral, Letemovir. *J. Am. Chem. Soc.* **2017**, *31*, 10637–10640.
 73. Bandini, M.; Bottoni, A.; Eichholzer, A.; Miscione, G.P.; Stenta, M. Asymmetric Phase-Transfer-Catalyzed Intramolecular N-Alkylation of Indoles and Pyrroles: A Combined Experimental and Theoretical Investigation. *Chem. Eur. J.* **2010**, *16*, 12462–12473.
 74. Coricello, A.; Mesiti, F.; Lupia, A.; Maruca, A.; Alcaro, S. Inside Perspective of the Synthetic and Computational Toolbox of JAK Inhibitors: Recent Updates. *Molecules.* **2020**, *25*, 3321.
 75. Lin, Q.; Meloni, D.; Pan, Y.; Xia, M.; Rodgers, J.; Shepard, S.; Li, M.; Galya, L.; Metcalf, B.; Yue, T.Y.; Liu, P.; Zhou, J. Enantioselective Synthesis of Janus Kinase Inhibitor INCB018424 via an Organocatalytic Aza-Michael Reaction. *Org. Lett.* **2009**, *11*, 1999–2002.
 76. Diner, P.; Nielsen, M.; Marigo, M.; Jorgensen, K.A. Enantioselective Organocatalytic Conjugate Addition of N Heterocycles to α,β -Unsaturated Aldehydes. *Angew. Chem. Int. Ed.* **2007**, *46*, 1983–1987.
 77. Haydl, A.M.; Xu, K.; Breit, B. Regio- and Enantioselective Synthesis of N-Substituted Pyrazoles by Rhodium-Catalyzed Asymmetric Addition to Allenes. *Angew. Chem. Int. Ed.* **2015**, *54*, 1–6.
 78. Warren, T.K.; Jordan, R.; Bavari, S. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature.* **2013**, *531*, 381–385.
 79. US Food and Drug Administration. Remdesivir EUA letter of authorization, 2020. Available: <https://www.fda.gov/media/137564/download> [Accessed 1 August 2021].

-
80. Kokic, G.; Hillen, H.S.; Tegunov, D.; Dienemann, C.; Seitz, F.; Schmitzova, J.; Farnung, L.; Siewert, A.; Höbartner, C.; Cramer, P. Mechanism of SARS-CoV-2 polymerase stalling by remdesivir. *Nat. Commun.* **2021**, *12*, 279.
 81. Oka, N.; Wada, T. Stereocontrolled synthesis of oligonucleotide analogs containing chiral internucleotidic phosphorus atoms. *Chem. Soc. Rev.* **2011**, *40*, 5829–5843.
 82. Yi-Ren, Z.; Kui, L.; Jin-Sheng, Y.; Jian, Z. Recent Advances in Catalytic Asymmetric Synthesis of *P*-Chiral Phosphine Oxides. *Acta Chim. Sin.* **2020**, *78*, 193–216.
 83. Siegel, D.; Hui, H.C.; Doerffler, E.; Clarke, M.O.; Chun, K.; Zhang, L.; Neville, S.; Carra, E.; Lew, W.; Ross, B.; Wang, Q.; Wolfe, L.; Jordan, R.; Soloveva, V.; Knox, J.; Perry, J.; Perron, M.; Stray, K.M.; Barauskas, O.; Feng, J.Y.; Xu, Y.; Lee, G.; Rheingold, A.L.; Ray, A.S.; Bannister, R.; Strickley, R.; Swaminathan, S.; Lee, W.A.; Bavari, S.; Cihlar, T.; Lo, M.K.; Warren, T.K.; Mackman, R.L. Discovery and Synthesis of a Phosphoramidate Prodrug of a Pyrrolo[2,1-*f*][triazin-4-amino] Adenine C-Nucleoside (GS-5734) for the Treatment of Ebola and Emerging Viruses. *J. Med. Chem.* **2017**, *60*, 1648–1661.
 84. Wang, M.; Zhang, L.; Huo, X.; Zhang, Z.; Yuan, Q.; Li, P.; Chen, J.; Zou, Y.; Wu, Z.; Zhang, W. Catalytic Asymmetric Synthesis of the anti-COVID-19 Drug Remdesivir. *Angew. Chem. Int. Ed.* **2020**, *59*, 20814–20819.
 85. Gannedi, V.; Kumar, B.K.; Reddy, S.N.; Ku, C.C.; Wong, C.H.; Hung, S.C. Practical Remdesivir Synthesis through One-Pot Organocatalyzed Asymmetric (*S*)-*P*-Phosphoramidation. *J. Org. Chem.* **2021**, *86*, 4977–4985.