

Type of the Paper: Original Article

Bacterial communities associated with the cycling of non-starch polysaccharides and phytate in aquaponics systems

Daniel Menezes-Blackburn^{1,*}, Nahad Al-Mahrouqi^{1,†}, Buthaina Al-Siyabi¹, Adhari Al-Kalbani¹, Ralf Greiner², Sergey Dobretsov^{3,4}

¹ Department of Soils, Water and Agricultural Engineering, Sultan Qaboos University, PO Box 34, Al-Khoud 123, Sultanate of Oman

² Department of Food Technology and Bioprocess Engineering, Max Rubner-Institut, Federal Research Institute of Nutrition and Food, Karlsruhe, Germany

³ Department of Marine Science and Fisheries, Sultan Qaboos University, PO Box 34, Al-Khoud 123, Sultanate of Oman

⁴ Centre of Excellence in Marine Biotechnology, Sultan Qaboos University, PO Box 50, Al-Khoud 123, Sultanate of Oman

[†] Co-first authorship – the first two authors contributed equally to this manuscript.

* Correspondence: danielblac@squ.edu.om; phone: + 968 2414 3668, ORCID: 0000-0002-8142-9655

Abstract: Aquaponics are efficient systems that associate aquatic organisms' production and plants by recirculating water and nutrients between aquaculture and hydroponic tanks. In this study, we have characterised the bacterial communities in the fresh water aquaponics system that can mineralise polysaccharides and phytate by producing carbohydrate degrading enzymes and phytases, by 16S rRNA gene sequencing and in vitro culture techniques. Around 20% of the operational taxonomic units (OTUs) identified were previously reported to carry fibre-degrading enzymes putative genes, namely β -glucanase (1%), xylanase (5%) or cellulases (17%). Ten % of the OTUs were previously reported to carry putative genes of phytases with different catalytic mechanisms, namely β -propeller (6%), histidine acid phytases (3%) and protein tyrosine phytase (<1%). Thirty-eight morphologically different bacteria were isolated from biofilms accumulated in fish and plant compartments, and identified to belong to the Bacilli class. Among these, seven could produce xylanase, 8 produced β -glucanase, 14 produced cellulase, and 11 isolates could secrete amylases. In addition, *Staphylococcus* sp. and *Rosellomorea* sp. could produce consistent extracellular phytate-degrading activity. The PCR amplification of β -propeller genes both in environmental samples and in the isolates obtained showed that this is the most ecologically relevant phytase type in the aquaponics systems used. In summary, the aquaponics system is abundant with bacteria carrying enzymes responsible for plant-nutrient mineralisation.

Keywords: aquaponics, metabarcoding, bacterial communities, carbohydrases, phytases

1. Introduction

Aquaponics is a sustainable closed-loop recirculating aquaculture system (RAS), combined with hydroponic crops, where the integrated production of plants and aquatic animals can be achieved (Buzby et al. 2016; Delaide et al. 2016; Goddek et al. 2016). There is a variety of salt and fresh water fish, invertebrates and plant species that can be cultivated in aquaponics systems, including tilapia, catfish, finfish, flatfish, shrimp, sea urchin, and several types of plants, like culinary herbs, lettuce, algae, and fruiting crops (Rakocy et al. 2016).

Aquaponics systems have recently received high interest from the general public due to their ability to produce fish and vegetables in small areas simultaneously, drastically improving water management compared to when hydroponics and aquaculture are carried out independently (Al-Hafedh et al. 2008; Bandi et al.). Additionally, it reduces the cost of effluent treatment and the environmental impact of water nutrient pollution resulting from agriculture and aquaculture production activities (Buzby et al. 2016; Turkmen and Guner 2010).

Water and biofilm microorganisms present in aquaponics systems have an essential role in the nutrient cycling from fish fecal matter and their bioavailability to plants (Kasozzi et al. 2020; Munguia-Fragozo et al. 2015). In aquaponics systems, heterotrophic microorganisms use undigested fish feed as an organic energy source, while chemo-litho-autotrophic nitrifiers utilise dissolved ammonia instead (Rurangwa and Verdegem 2015). These microorganisms play an important role in converting fish organic wastes into valuable plant nutrients (e.g. PO_4^- , NH_4^+ , NO_3^-) which can then be uptaken by plants in their ionic forms, eliminating the need for plant fertiliser inputs (Tyson et al. 2011). Aquaponics systems display diverse microniches for the growth of microbial populations (Leonard et al. 2000; Munguia-Fragozo et al. 2015). Due to different types of cultivated fish and plants, characteristics of water, design of fish and plant compartments, nutrient composition and temperature of effluents, diverse microbial communities carry out the cycling of various nutrients on this system (Buzby et al. 2016; Schmutz et al. 2017).

On the other hand, non-starch polysaccharides (NSP) and phytate are among the main challenges when plant-based fish feed is used (Council ; Francis et al. 2001; Menezes-Blackburn and Greiner 2014). Plant-based feeds have a high phytate content, and fish cannot efficiently utilise phytate (da Silva Cerozi and Fitzsimmons 2017). This leads to the excretion of phytate into the water, resulting in nutrient pollution, algal growth and hypoxic environments (Liebert and Portz 2005). However, water microorganisms carrying extracellular phytases can mineralise phytate present in the fish feed (Hill et al. 2007; Jorquera et al. 2008). Fish diets can also be supplemented with phytase to increase phosphorus availability and decrease phytate anti-nutrient activity (Hien et al. 2015).

Fish feeds are rich in indigestible fibre that can only be degraded after excretion by an array of water and biofilm microbes, making the aquaponics systems an interesting environment to study bacterial strains that produce non-starch polysaccharides (NSP) degrading enzymes. Fish lack the enzymes capable of hydrolysing the NSP β -glycosidic bonds (Krogdahl et al. 2005). Similarly to phytases, exogenous carbohydrases can be supplemented with plant-based fish feed to increase NSP digestibility, increase feed energy value and decrease NSP anti-nutrient activity (Menezes-Blackburn and Greiner 2014). Both phytate and NSP degrading enzymes are highly valued as feed additives, and aquaponics systems represent a possible source of new enzymes with improved properties compared to the ones available in the market (Menezes-Blackburn and Greiner 2014). However, most microbial studies in aquaponics systems focused on ammonia and nitrite oxidiser bacteria, or on the presence of pathogens (Munguia-Fragozo et al. 2015).

The present study aimed to investigate the diversity of bacterial communities associated with phytate and NSP mineralisation in an aquaponics system using MiSeq 16S rRNA V4 gene sequencing, together with isolation and the screening of bacterial strains capable of secreting cellulase, xylanase, β -glucanase and phytase enzymes.

2. Materials and Methods

2.1 Aquaponics system and samples collection

An outdoor aquaponics system was installed at Sultan Qaboos University, Muscat, Oman (23.5896° N, 58.1735° E). It was composed of three compartments: one fish tank with 4 Nile tilapia (*Oreochromis niloticus*, average length 20cm) connected to a hydroponic unit (plant compartment), and a water treatment unit. The water from the fish tank is used to grow lettuce (*Lactuca sativa*) hydroponically. Each aquaponics unit contained ten lettuce plants floating in the surface of the water in the plant compartment. The treatment unit is used to maintain the water quality in an optimal range. It had a sand filter that removed solids from the water, a biological filter (Bactoballs®) that removed ammonia, and a pump that pumped the water back to the fish tank. The total volume of each aquaponics unit was 600 L and the flow rate used was 1 L min⁻¹. The outdoor aquaponics system was shaded to reduce natural sunlight, and had an average light intensity of 100

$\mu\text{mol m}^{-2} \text{ s}^{-1}$ during the experiment. There were three replicated identical aquaponics units used in this study independently sampled and analysed.

The biofilm/water samples were collected on 23rd January 2020, and the aquaponics systems ran for one month before sampling. Biofilms accumulated on the walls of fish and plant compartments were scraped using a sterile spatula. Biofilms samples were collected in multiple locations of each compartment. Three aquaponics units were sampled independently. Biofilms samples suspended with site water from the corresponding compartment in a sterile 15 ml centrifuge tube (Falcon) were homogenised to obtain a representative sample. For the plant compartment samples, biofilms associated with the root of lettuce were also collected. Then, collected biofilm samples were freshly assayed.

In addition, 50 ml water samples from the fish tank and plant compartments were collected at the end of the experiment and water samples were preserved in a freezer (-20°C) for further analysis. Additionally, water quality was analysed. Water pH and water electrical conductivity were determined. The elemental composition of water was analysed by ICP OES (Thermo Scientific, UK). The temperature and light in both compartments were monitored using HOBO Pendant® loggers. Light intensity was measured at the bottom of the compartment in lux and converted to $\mu\text{mol m}^{-2} \text{ s}^{-1}$. Results from the background water analysis are displayed in Table 1.

Table 1- Average temperature ($T^{\circ}\text{C}$), light intensity ($\mu\text{mol m}^{-2} \text{ s}^{-1}$), pH, electrical conductivity (EC, mS m^{-1}) and metal elemental concentration (mg L^{-1}) in the water samples collected from the fish and plant compartments ($n=3$) of the studied aquaponics system.

Compartment	T ($^{\circ}\text{C}$)	light	pH	EC	P	K	Ca	Mg	Na	Cu	Zn
Fish	29.2	0.24	6.1	2.4	25	33.9	81.8	63.1	198	0.3	<0.01
Plant	29.7	0.19	5.8	2.9	28	34.5	83.8	65.2	183	0.5	<0.01

2.2 Prokaryote diversity through 16S rRNA gene sequencing

The total genomic DNA was extracted from each of the six biofilm and water samples using a Purelink™ microbiome DNA purification kit (Thermo Fisher Scientific, UK) following the manufacturer’s protocol. Then, the concentration and purity of DNA were measured using a NanoDrop™ (Thermo Fisher Scientific, UK) spectrophotometer. The DNA extracts were sent to Molecular Research (Shallowater, Texas, USA) for bacterial 16S (515-806) amplicon diversity assays using bTEFAP® PGM/S5. The V4 variable region of the bacterial 16S rRNA gene was amplified using 515F forward and 806R reverse primers with a barcode on the forward primer during 35 cycles of PCR using the HotStarTaq Plus Master Mix Kit (Qiagen, USA). The following conditions were used: 94°C for 3 mins, followed by 35 cycles of 94°C for 30 sec, 53°C for 40 sec and 72°C for 1 min, thereafter a final elongation step at 72°C for 5 mins was performed. After amplification, PCR products were checked in 2% agarose gel to determine the success of amplification and the relative intensity of the bands. Samples were purified using calibrated Ampure XP beads. Pooled in equimolar ratios, the purified PCR products were sequenced using Illumina MiSeq platform in MR DNA (www.mrdnalab.com, Shallowater, TX, USA) following the manufacturer’s guidelines.

The sequence data were processed using the MR DNA ribosomal and functional gene analysis pipeline and in-house built software (<http://www.mrdnafreesoftware.com>, MR DNA, Shallowater, TX). Sequences were depleted of barcodes and primers. Sequences with ambiguous base calls or with homopolymer runs exceeding 6bp or short sequences < 150bp were removed. Sequences were denoised, and operational taxonomic units (OTUs) were defined clustering at 3% divergence (97% similarity) followed by singleton

sequences and chimeras removal. Final OTUs were taxonomically classified using BLASTn against a curated database derived from RDP II and NCBI (www.ncbi.nlm.nih.gov).

2.2.1. OTUs reported to carry phytases and carbohydrases enzymes

The assigned taxa from the 16S rRNA sequences obtained through the microbial diversity analysis were compared to the bacterial species reported in Pfam database (<http://pfam.xfam.org/>) to carry different types of phytases and carbohydrases. The following protein families were selected and used in this analysis: beta propeller phytase (BPP) (phytase (PF02333)), histidine acid phosphatase (HAP) (His_Phosphatase_2 (PF00328)), Cellulase N-terminal Ig-like domain (cellulase) (CelD_N (PF02927)), protein tyrosine phosphatase (PTP) (PTPlike_phytase (PF14566)), Beta-1,3-glucanase (β -glucanase) (Glyco_hydro_64 (PF16483)) and Carbohydrate family 9 binding domain-like (xylanase) (CB_M91 (PF06452)). This cross-referencing was used as soft evidence for the presence and abundance of these enzymes across the samples. The bacteria screening for the studied enzymes were taken as strong evidence of the presence and abundance of these enzymes in the aquaponics system.

2.3 Isolation and identification of bacterial strains

Bacterial strains were isolated from both fish and plant compartments by enrichment of one gram of each biofilm sample in 20 ml of water from a corresponding compartment, 20 ml compartment water with sterile 3% wheat bran (natural source of phytate and fibres), and minimal media M9 (Pardee et al. 1959). The mixture was incubated at 37°C for three days. Subsequently, 100 μ l from each enrichment was submitted to serial dilution, inoculated on nutrient agar and incubated at 30 °C for one day. Pure colonies were streaked clean in fresh nutrient agar plates separately and grown for another day at 30 °C. The isolated strains were conserved in 25% glycerol-nutrient broth for subsequent assays.

Bacterial genomic DNA was extracted from all strains for further PCR assays using a HiPurA™ Kit (HiMedia, Mumbai, India). All isolated strains positive for extracellular enzyme activities (19 strains) were identified by 16S rRNA gene sequencing. For this purpose, the universal primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-TACGGYTACCTTGTTACGACTT-3') (Heuer et al. 1997) were used with a Hi-Chrom PCR Master Mix (HiMedia, Mumbai, India) according to the manufacturer's instructions. The PCR reactions were performed using 4 min hot start at 96 °C, followed by 30 cycles of 30 s at 94 °C, 30 s at 57 °C and 1 min at 72 °C, and a final extension step at 72 °C for 10 min. Amplicons were purified and both forward and reverse sequenced at the MacroGen sequencing service (MacroGen Inc., Korea). The sequences were aligned, trimmed and compared to the closest sequence (highest ID from NCBI GenBank) and the cleaned consensus sequences were deposited in NCBI GenBank (Table 2).

Table 2. Lab isolation name, length of the 16S rRNA gene sequence, nucleotide sequence length of the strains obtained, NCBI GenBank sequence with the highest identity with the isolate’s 16S rRNA gene, assigned strain name and GenBank accession number of the deposited 16S rRNA sequence.

Isolation ID	Length (bp)	NCBI Genbank closest match				Assigned Strain name	Accession N.
		Seq. ID	Identity	Q.Cover	Species		
SWAM_2Aq	1359	MT539995.1	100.00%	100%	<i>Bacillus subtilis</i>	<i>Bacillus subtilis</i> 2Aq	OK626679
SWAM_6Aq	1318	MT229334.1	98.79%	100%	<i>Bacillus velezensis</i>	<i>Bacillus velezensis</i> 6Aq	OK626680
SWAM_7Aq	1287	KM922586.1	98.45%	100%	<i>Bacillus subtilis</i>	<i>Bacillus subtilis</i> 7Aq	OK626681
SWAM_8Aq	1299	KM922586.1	98.00%	100%	<i>Bacillus subtilis</i>	<i>Bacillus subtilis</i> 8Aq	OK626682
SWAM_10Aq	1317	MF521557.1	93.06%	100%	<i>Bacillus cereus</i>	<i>Bacillus cereus</i> 10Aq	OK626683
SWAM_13Aq	1395	MN330286.1	99.63%	97%	<i>Bacillus subtilis</i>	<i>Bacillus subtilis</i> 13Aq	OK626684
SWAM_15Aq	1405	MK859953.1	99.15%	100%	<i>Bacillus licheniformis</i>	<i>Bacillus licheniformis</i> 15Aq	OK626685
SWAM_16Aq	1327	MW578436.1	91.40%	99%	<i>Bacillus licheniformis</i>	<i>Bacillus</i> sp.16Aq	OK626686
SWAM_17Aq	1422	MH005066.1	91.82%	99%	<i>Bacillus infantis</i>	<i>Bacillus</i> sp.17Aq	OK626687
SWAM_18Aq	1416	MT122066.1	99.36%	100%	<i>Bacillus tequilensis</i>	<i>Bacillus tequilensis</i> 18Aq	OK626688
SWAM_21Aq	1446	CP035288.1	93.05%	99%	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus</i> sp. 21Aq	OK626689
SWAM_22Aq	1347	LS974830.1	93.87%	100%	<i>Rossellomorea marisflavi</i>	<i>Rossellomorea</i> sp. 22Aq	OK626690
SWAM_23Aq	1380	MN704425.1	98.04%	99%	<i>Bacillus stercoris</i>	<i>Bacillus subtilis</i> 23Aq	OK626691
SWAM_26Aq	1414	MT111029.1	98.46%	100%	<i>Bacillus subtilis</i>	<i>Bacillus tequilensis</i> 26Aq	OK626692
SWAM_28Aq	1427	MT326233.1	98.95%	100%	<i>Bacillus licheniformis</i>	<i>Bacillus licheniformis</i> 28Aq	OK626693
SWAM_29Aq	1371	MT611946.1	98.53%	100%	<i>Bacillus cereus</i>	<i>Bacillus cereus</i> 29Aq	OK626694
SWAM_33Aq	1433	CP053934.1	99.02%	100%	<i>Bacillus thuringiensis</i>	<i>Bacillus thuringiensis</i> 33Aq	OK626695
SWAM_36Aq	1423	KF933659.1	93.74%	100%	<i>Rossellomorea aquimaris</i>	<i>Rossellomorea</i> sp. 36Aq	OK626696
SWAM_37Aq	1423	MK942526.1	89.88	100%	<i>Bacillus subtilis</i>	<i>Bacillus</i> sp. 37Aq	OK626697

2.4 Screening for Carbohydrases producing bacteria

2.4.1 Screening for non-starch polysaccharides degrading bacteria

Bacterial isolates were inoculated onto xylanase screening agar and β -glucanase screening agar plates, which constituted of 5 g L⁻¹ xylan (or 5 g L⁻¹ β -glucan for β -glucanase screening agar), 2 g L⁻¹ yeast extract, 0.5 g L⁻¹ NaCl, 0.1 g L⁻¹ CaCl₂, 5 g L⁻¹ peptone, 0.5 g L⁻¹ MgSO₄·7H₂O and 20 g L⁻¹ agar dissolved in distilled water (Subajini et al. 2012) and incubated for 48 h at 37 °C. Then, plates were flooded with 0.1% Congo red (Sigma) solution and washed with 1 M NaCl (Sigma) solution. The presence of a clear zone (halo) indicated the isolate's extracellular β -glucanase enzymatic activity. The halo and colony diameters were recorded as semiquantitative evidence of the β -glucanase activity expressed by each strain.

For the screening of cellulase producing bacteria, carboxymethyl cellulose (CMC) agar medium was used containing 5 g/L carboxymethyl cellulose, 1 g L⁻¹ NaNO₃, 1 g L⁻¹ K₂HPO₄, 1 g L⁻¹ KCl, 0.5 g L⁻¹ MgSO₄, 0.5 g L⁻¹ yeast extract and 15 g L⁻¹ agar dissolved in distilled water (Maravi and Kumar 2020). A loopful of bacterial suspension was spotted onto agar and incubated at 37 °C for 48 h. Similarly to the β -glucanase assays, plates were stained by flooding with 0.1% Congo red then washed using 1 M NaCl solution, and the clearance zone diameter was recorded.

2.4.2 Screening for starch degrading bacteria

Similarly to cellulases and β -glucanases assays, amylase screening agar plates were used to assess the ability of the bacteria strains to produce extracellular amylases (Fooladi and Sajjadian 2010). The amylase screening agar medium contained 10 g of starch, 2 g of yeast extract, 5 g of peptone, 0.5 g of MgSO₄, 0.5 g of NaCl, 0.15 g of CaCl₂, 2 g agar, and 1 litre of water. The plates were incubated for 48 h at 37 °C. Then, then flooded with Gram's iodine solution, and the clearance zones (mm) and diameter of colonies (mm) were recorded in triplicate.

2.5 Screening for Phytase producing bacteria

Bacterial isolates were screened for phytase production using Phytase Screening Medium (PSM) (Demirkan et al., 2014) and M9 Minimal medium with phytate. The M9 broth minimal medium with phytate contained 0.4% Na-phytate (sterile filtered), 0.1% NH₄Cl, 0.012% MgSO₄ and 0.00147% CaCl₂. The PSM medium consisted of 20 g L⁻¹ Glucose, 2 g L⁻¹ CaCl₂, 5 g L⁻¹ NH₄NO₃, 0.5g L⁻¹ MgSO₄, 0.5 g L⁻¹ KCl, 0.01 g L⁻¹ FeSO₄, and 4 g L⁻¹ Na-phytate (sterile filtered). Isolates were inoculated in both media and incubated at 37 °C for 14 days. Subsamples were collected daily and analysed for soluble inorganic phosphate concentration using the malachite green method (Ohno and Zibilske 1991). The isolates that showed a significant increase in the extracellular phosphorus concentration were re-assayed in the corresponding medium for another 14 days at 37 °C for confirmation of the results. The bacterial growth was also measured daily by OD at 600 nm. The uninoculated corresponding sterile media was used as a control.

2.6 Detection of phytase genes by PCR amplification using degenerate primers

The presence of phytase genes was examined using PCR assays using degenerate primers of two different classes (BPP and PTP). This assay was performed for both the environmental genomic DNA extracted from bio-film/water samples and the genomic DNA extracted from the bacterial isolates. For BPP phytases, two different primer pairs were used according to Huang et al. (2009): BPP-F (5'-GACGCAGCCGA YGAYCCNCGNITNTGG-3') and BPP-R (5'-CAGGSCGCANRTCIACRTTRTT-3'). The PCR conditions were: 4 min hot start at 95°C, followed by eight cycles of 95°C for 30 s, 57°C (decreasing by 1°C after each cycle) for 30 s, and 72°C for 30 s, followed by 27 cycles of 95°C for 30 s, 48°C for 30 s, and 72°C for 30 s and then a final extension at 72°C for 5 min. For the second BPP primer pair, DP1 (5'-GAY GCI GCI GAY GAY CCI GC-3') and DP2 (5'-TCR TAY TGY TCR AAY TCIC-3') primers were used according to Tye et al. (2002). Amplification was carried out for 30 cycles of 94 °C for 45 s, 50 °C for 45 s, and 72 °C for 1 min. For the PTP phytase gene amplification, the primers CPhy-F (5'-GTGGACCTRCGRMAR-GARWCICA -3') and CPhy-R (5' GTCCGACCATTGCCTGCTCARTGRAMRTGIADCCA-3') were used according to Huang et al. (2011). The PCR conditions were: 95 °C for 4 min, 10 cycles of 94 °C for 30 s, 58 °C for 30 s (decreasing 0.5 °C for each cycle), and 72 °C for 30 s, followed by 27 cycles of 94 °C for 30 s, 52 °C for 30 s, and 72 °C for 30 s and a final extension step at 72 °C for 10 min.

2.7 Statistical analyses

The data were analysed using one way ANOVA and Tukey's test, $p \leq 0.05$ was used as the significance level. Data calculations, manipulation, average, standard deviation, and correlation analysis were performed using Microsoft Office Excel 2016. Krona (Ondov et al. 2011) was used to build HTML interactive hierarchical microbial diversity graphics, allowing for the visualisation of changes in microbial community composition. PAST4 (Hammer et al. 2001) was used for calculating microbial diversity indexes, and JMP13 statistical software was used for ANOVA and PCA interpretation of the effect of the treatments on microbial parameters. Graphia 2.0 was used to build a correlation network of OUTs with Pearson's correlation coefficients above 0.95 (Köhler et al. 2006), clustered by Markov Cluster Algorithm (MCL, granularity 1.1).

3. Results

3.1 16S rRNA diversity of bacterial communities in fish and plant compartments


Due to the high data variability among replicates from the three studied independent Aquaponics systems, the bacteria community composition was not statistically different between fish and plant compartments. In total, 149,197 bacterial sequences and 1,655 OTUs were obtained from the fish compartments samples and 115,182 sequences and 1,424 OTUs from samples that originated from plant compartments. The Shannon diversity index (H), Dominance (D), Evenness (EH) were very similar for fish (n=3), plant (n=3) and fish+plant compartments (n=6) as shown in Table 03. This was observed both when the analysis was performed using the sum of the number of sequences

of OTUs from individual samples, and also when averaged (\bar{x}) from individual samples. It is noteworthy that even though the diversity indexes were not different between fish and plant compartments, 527 OTUs were uniquely occurring in plant compartments (27% of total OTUs), and 296 OTUs were only found in fish compartments (15% of total OTUs). This evidences that these two environments are very different in their bacterial community composition.

Table 03. Shannon diversity index (H), Dominance (D), Evenness (EH) of total prokaryotic operational taxonomic units (OTUs) for three independent demonstrative aquaponics systems. Samples were grouped by fish (n=3), plant (n=3) and fish+plant compartments (n=6) followed by average (\bar{x}) and standard error ($\sigma\bar{x}$). Analysis was performed using PAST4 software.

	Fish			Plant			Fish+Plant		
	\bar{x}	$\sigma\bar{x}$		\bar{x}	$\sigma\bar{x}$		\bar{x}	$\sigma\bar{x}$	
Shannon H	5.96	5.17	0.21	5.90	5.23	0.18	6.24	5.20	0.12
Dominance D	0.01	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.00
Evenness E_H	0.24	0.25	0.06	0.26	0.27	0.02	0.26	0.26	0.03

Only a small proportion of Archaea (0.1 to 0.2%) was present in the samples. In fish tanks, 82% of Archaea belonged to phyla Thaumarchaeota and 18% to Euryarchaeota (class Thermoplasmata). The dominant genera included *Candidatus Nitrososphaera*, *Cenarchaeum*, and *Methanomassiliicoccus*. In the plant compartment samples, all Archaea belonged to the Thaumarchaeota phylum with dominant genera *Nitrososphaera*, *Candidatus Nitrososphaera*, and *Candidatus Nitrosoarchaeum*.



Bacterial communities were dominated by Proteobacteria and Bacteroidetes Phyla (Figure 1). The Proteobacteria phylum accounted for 53% of total sequences in the fish tanks and 49% of total reads in the plant compartment. The Bacteroidetes phylum was more abundant in the fish compartment (24%) than in the plant compartment (15%). Among Proteobacteria, Alphaproteobacteria was the most dominant class (23-24%, Figure 1), followed by Gammaproteobacteria, Betaproteobacteria and Deltaproteobacteria. The phylum Bacteroidetes was largely represented by the Sphingobacteriia, Flavobacteriia and Cytophagia classes (Figure 1). The phyla Firmicutes, Nitrospirae, Planctomycetes, and Actinobacteria were less abundant (Figure 1). Notably, the relative abundance of the phyla Nitrospirae, Planctomycetes Verrucomicrobia, and Actinobacteria was two-fold higher in the plant compartments than in fish compartments. Similarly, the classes Bacilli, Clostridia, Fusobacteria, and Chloroflexi were present in both compartments but were around two-folds more abundant in the plant than in the fish compartment.

3.2 Multivariate correlation network clustering of bacteria taxa

The correlation network clustering of bacteria OTUs using Markov Cluster Algorithm (MCL, granularity 1.1) showed an outcome of six well-defined clusters (Figure 2). Most of the prokaryote taxa were represented in clusters 1 and 2. Clusters 1 and 3 were 'central' clusters (linking to all other clusters), and Cluster 5 was the most detached among the 6 clusters, connected only with clusters 1 and 3. Cluster 1 was very diverse, represented mainly by Sphingobacteriia, Erysipelotrichia and Bacteroidetes classes. In addition, cluster 1 was extremely rich with species that were reported to carry BPP phytase enzyme and different carbohydrases such as xylanase, β -glucanase and cellulases (see section 3.3). Cluster 3 was dominated by Euryarchaeota, Oscillatoriothymiceae, Fibrobacteres, Holophagae and Thermoplasmata phyla. Cluster 5 largely contains microbes belonging to Acidobacteria and Chloroflexi phyla. Moreover, most OTUs reported harbour PTP phytase genes were represented by cluster 2, which was most abundant in plant compartments. The OTUs most representative from fish compartments (the highest in abundance) were predominantly contained in clusters 1, 3 and 5, while the OTUs most representative plant compartments belonged primarily to clusters 2, 4 and 6.




Figure. 2. Correlation network clustering of operation taxonomic units (OTUs) present in fish and plant compartments of an aquaponics system. Figure generated using Graphia software ($r \geq 0.95$) and OUT data from V4 16S rRNA. Fish compartment OTUs were best represented by clusters 1, 3 and 5, whereas plant compartment OTUs were best represented by clusters 2, 4 and 6.

3.3 Diversity and abundance of microbes reported to carry putative phytases and carbohydrases enzymes genes

The outcome OTUs from the V4 16 rRNA diversity analysis was coss-compared to the bacterial species reported to harbour phytase or carbohydrase genes in the Pfam database (<http://pfam.xfam.org/>, Table 03). Given the currently available knowledge, this analysis represents the abundance and diversity of bacteria potentially carrying the putative genes for the studied enzymes (here deemed ‘potential producers’). Therefore, these data must be taken as soft evidence and only as an attempt to extract valuable information from the 16S diversity analysis pertaining to the scope of the objectives of this study. Besides reporting the total number of OTUs of ‘potential enzyme producers’, Table 03 also illustrates their relative abundance and the average and standard error per sample. The high standard error observed highlights that the high variability of the microbial communities between independent Aquaponics systems prevents drawing definitive conclusions when comparing fish and plant compartments. Among the observed OTUs in the aquaponics system used, the relative abundance of OTUs from potential carbohydrases producers was 2.8-fold higher than the ones reported to carry phytase putative genes (Table 3; 23% carbohydrases and 8% phytases). The fish tank samples showed a higher relative abundance of OTUs from potential carbohydrase producers compared to the samples from the plant compartments (26% and 19%, correspondingly). The number of putative phytase sequences was similar in both compartments. The relative abundance of OUTs potentially carrying BPP phytase genes was around 6% for both compartments, twice the relative abundance observed for HAP phytases (3% of sequences). Bacterial OTUs potentially harbouring PTP putative phytase genes were less abundant than for other types of phytases, corresponding to <1% of the reads in both compartments. The number of OTUs of bacteria reported to carry putative genes of different types of carbohydrases, such as cellulase, β -glucanase and xylanase, were more abundant in the fish compartment compared to the plant compartment. Among carbohydrases, the relative abundance of bacterial OTUs of potential cellulase producers was more abundant in plant compartments (18%) than in fish compartments (23%).

Table 03. Number of total operational taxonomic units (OTUs) of fish (n=3), plant (n=3) and fish+plant compartments (n=6) in three independent demonstrative aquaponics systems, their relative abundance with respect the total amount of sequences (%), the average of the number of OTUs (\bar{x}) and standard error ($\sigma\bar{x}$) of prokaryotes previously reported to carry putative Phytases (β -propeller BPP (PF02333); Histidine acid phytases HAP(PF00328); protein tyrosine phytases PTP (PF14566)) and Carbohydrases putative genes (B-glucanase (PF16483); Xylanase (PF06452); Cellulases (PF02927)). Analysis was performed by comparing the OTUs 16S rRNA diversity analys with a curated database extracted from PFAM (<http://pfam.xfam.org/>).

	Fish				Plant				Fish+ plant			
		%*	\bar{x}	$\sigma\bar{x}$		%*	\bar{x}	$\sigma\bar{x}$		%*	\bar{x}	$\sigma\bar{x}$
All	1655	100	794	245	1424	100	733	188	1951	100	763	139
BPP	76	6	39	13	68	5	33	12	90	6	36	8
HAP	46	3	21	8	31	3	17	4	54	3	19	4
PTP	6	<1	2	2	10	<1	3	2	12	<1	3	1
All phytases	98	9	50	17	87	8	41	14	120	8	46	10

B-glucanase	28	2	15	3	23	1	9	4	31	1	12	3
Xylanase	108	6	55	18	77	4	34	12	117	5	44	11
Cellulases	220	18	116	32	220	23	116	32	250	17	116	20
All carbohydrases	232	26	120	34	199	19	103	21	266	23	112	18

* % of total sequences from all three replicates or relative abundance.

3.4 Phytase, Xylanase, β -glucanase, Cellulase and Amylase producing bacterial isolates

Thirty-eight bacterial isolates were obtained from the fish tanks and plant compartments of the outdoor aquaponics system, and among them, 19 produced at least one of the extracellular enzyme activities looked for (Table 4). Only the strains with enzyme activities were identified by 16S sequencing. Most of them belonged to the *Bacillus* genus. Five strains (2Aq, 7Aq, 8Aq, 13Aq, 26Aq) were identified as *Bacillus subtilis*, two as *Bacillus cereus* and another two as *Bacillus tequilensis*. Besides the *Bacillus* genus, *Staphylococcus* and *Rosellomorea* were also found (Table 2). The four isolates *B. subtilis* 2Aq and 7Aq, *B. velezensis* 6Aq, and *B. tequilensis* 18Aq were capable of producing xylanase, β -glucanase, cellulase and amylase (Table 4). The carbohydrases looked for were differentially expressed by different isolates, based on the different diameters of the clear zone on specific agar minimal media. Moreover, *B. subtilis* 8Aq and 13Aq showed the ability to produce β -glucanase, cellulase and amylase but not xylanase. On the other hand, three isolates *Bacillus* sp. 16Aq, *B. subtilis* 2Aq and *B. licheniformis* 28Aq produced only cellulases, whereas *B. cereus* 10Aq and *Rosellomorea* sp. 36Aq could produce the only amylase. In total, 7, 8, 14 and 11 isolates could secrete extracellular xylanase, β -glucanase, cellulase and amylase, respectively.

Table 4. Bacterial isolates from fish and plant compartments of the aquaponics system with ability to produce extracellular phytase (increase extracellular phosphate concentration in PSM medium), xylanase (XSA, xylanase screening agar clear zone in mm), β -glucanase (BSA, β -glucanase screening agar clear zone in mm), cellulase (CSA, cellulase screening agar clear zone in mm) and amylase (ASA, amylase screening agar clear zone in mm). '+' indicates significant extracellular enzyme activity and '-' no extracellular enzyme activity was detected.

Source	Strain	Phy	XSA	BSA	CSA	ASA
Fish	<i>Bacillus subtilis</i> 2Aq	-	24	14	25	26
Fish	<i>Bacillus velezensis</i> 6Aq	-	26	24	16	18
Fish	<i>Bacillus subtilis</i> 7Aq	-	25	27	18	14
Fish	<i>Bacillus subtilis</i> 8Aq	-	-	26	20	14
Fish	<i>Bacillus cereus</i> 10Aq	-	-	-	-	20
Fish	<i>Bacillus subtilis</i> 13Aq	-	-	49	27	20
Fish	<i>Bacillus licheniformis</i> 15Aq	-	26	-	15	-
Fish	<i>Bacillus</i> sp.16Aq	-	-	-	12	-
Plant	<i>Bacillus</i> sp.17Aq	-	-	33	-	-
Plant	<i>Bacillus tequilensis</i> 18Aq	-	23	46	26	22
Plant	<i>Staphylococcus</i> sp. 21Aq	+	-	-	-	-

Plant	<i>Rossellomorea</i> sp. 22Aq	-	-	-	25	-
Plant	<i>Bacillus subtilis</i> 23Aq	-	16	25	-	-
Plant	<i>Bacillus tequilensis</i> 26Aq	-	36	-	25	30
Plant	<i>Bacillus licheniformis</i> 28Aq	-	-	-	13	-
Plant	<i>Bacillus cereus</i> 29Aq	-	-	-	16	-
Plant	<i>Bacillus thuringiensis</i> 33Aq	-	-	-	16	21
Plant	<i>Rossellomorea</i> sp. 36Aq	+	-	-	-	13
Plant	<i>Bacillus</i> sp. 37Aq	-	-	-	15	10

All strains grown in phytate supplemented M9 medium didn't show any release of phosphorus. However, when cultured in PSM medium for one week, seven different strains were capable of increasing the phosphorus concentration in the growth medium. Nevertheless, the initial assays were somewhat ambiguous, and these seven isolates that initially showed a significant increase in phosphorus concentrations were re-assayed for another 14 days of cultivation. Only two phytase producing bacteria were confirmed: *Staphylococcus* sp. 21Aq and *Rossellomorea* sp. 36Aq were able to consistently increase the extracellular concentration of phosphorus after eight days of cultivation in PSM medium. These isolates exhibited high phosphorus release capabilities, sustaining over 20 mg P L⁻¹ from days 8 to day 14 of cultivation (Figure 3). To confirm the presence of phytase genes, different phytase degenerate primers BPP (for BPP), DP (for BPP) and Cphy (for PTP) were used. Results showed that these strains showed positive amplification of the BPP primers, thus indicating that these bacteria very likely produce β -propeller phytases.




Figure 3. Change in the phosphorus concentration (mg L⁻¹) in PSM medium (with respect to uninoculated control) after inoculation with *Staphylococcus* sp. 21Aq or *Rossellomorea* sp. 36Aq strains. The *Bacillus* sp. 37Aq strain with no extracellular phytate hydrolysis ability was used as the negative control illustration.

4. Discussion

Aquaponics are commonly proposed efficient recirculating aquatic systems that combine the production of aquatic organisms (fish) and plants with the re-use of water (Tyson et al. 2011). In order to function properly, aquaponics systems are relay on proper organic matter mineralisation and nutrient cycling. These processes are heavily reliant on water

and microbial biofilms. Thus, it is necessary to study microbes and their enzymes associated with aquaponics systems. Nitrogen transformations are carried out by both aerobic and anaerobic heterotrophic microorganisms (Joo et al. 2005). Various microniches exist within recirculating aquaponics systems that promote the growth of specific microbial communities, which play a role in mineralising organic wastes (Sharrer et al. 2005). In aquaponics, both heterotrophic and autotrophic bacteria are present. Autotrophic bacteria may be chemolithotrophic, obtaining energy through the oxidation of iron, sulphur or inorganic nitrogen. Heterotrophic bacteria use undigested organic matter from fish faeces as a source of energy and carbon (Sharrer et al. 2005; Sugita et al. 2005) and are responsible for proteolysis and sulfate reduction (Sugita et al. 2005). Eutrophic bacterial biomass increases with the increase of suspended and dissolved organic matter (Leonard et al. 2002).

Most microbial studies in aquaponics systems are based on culture-dependent techniques, and the number of studies that utilise next-generation DNA sequencing techniques are limited (Munguia-Fragozo et al. 2015). In this study, the V4 16S rDNA diversity analysis together with culture techniques showed highly variable and diverse bacterial communities. We expected different communities in fish and plant compartments because various microniches exist within recirculating aquaponics systems that promote the growth of specific microbial communities, which play a role in mineralising organic wastes (Sharrer et al. 2005). However, the link between fish and plant compartments to specific bacterial communities was not strongly demonstrated due to high differences in the relative abundance of OTUs among replicates of independent aquaponics systems. While measured environmental parameters, like temperature, light, and metal elemental concentration were similar between replicated systems, other environmental factors could be responsible for the high variability in the relative abundance of OTUs.

In this study, Proteobacteria was the most dominant phylum in both fish and plant compartments. Alphaproteobacteria, Flavobacteria, Sphingobacteriia and Cytophagi were the most abundant classes observed. Similarly, in another study of bacterial communities by 16S rRNA sequencing of eight aquaponics and aquaculture systems it was demonstrated that Proteobacteria and Bacteroidetes were the most abundant phyla (Eck et al. 2019). Classes Alphaproteobacteria, Gammaproteobacteria, Actinobacteria, Bacteroidetes, Planctomycete, Bacilli, Nitrospirae, Betaproteobacteria, Nitrosomonas and Sphingobacteria have been reported as the most common ones found in freshwater aquaponics systems (Munguia-Fragozo et al. 2015). The differences between classes of bacteria in different studies could be explained by the type of aquatic organisms used to grow in the aquaponics system. For example, Sugita et al., (2005) reported that Alphaproteobacteria and Betaproteobacteria were the most abundant phyla in freshwater recirculating aquaponics when using common carp (*Cyprinus carpio*) species. When goldfish (*Carrassius auratus*) was used, the bacterial community was more diverse and included Planctomycetacia, Bacilli, Actinobacteria, Planctomycetacia, and Gammaproteobacteria bacterial groups.

In aquaponics, both heterotrophic and autotrophic bacteria are present. Autotrophic bacteria may be chemolithotrophic and obtain energy through the oxidation of iron, sulphur or inorganic nitrogen. Heterotrophic bacteria use undigested organic matter from fish faeces as a source of energy and carbon (Sharrer et al. 2005; Sugita et al. 2005) and are responsible for proteolysis and sulfate reduction (Sugita et al. 2005). Heterotrophic bacteria in aquaponics systems that can produce extracellular hydrolytic enzymes may possibly be of biotechnological interest for feed-enzyme supplement applications. In this study, 38 bacterial strains were isolated; most were identified to belong to the *Bacillus* genus. Even though the V4 16S rRNA diversity analysis (Figure 1) revealed that *Flavobacterium*, *Haliscomenobacter*, *Nitrospira*, *Thermomonas* and *Novosphingobium* were highly abundant genera, these were not found among our isolates. However, genera *Flavobacterium* and *Novosphingobium* were previously reported to carry phytases, and *Flavobacterium*, *Haliscomenobacter*, and *Cetobacterium* were reported to carry carbohydrases putative genes in PFAM database (<http://pfam.xfam.org/>). Nishioka et al. (2016) reported that the use of selective culture media is important for the effective isolation of *Flavobacterium*

spp. This might explain the failure in our study to isolate *Flavobacterium* in our study. Further work is needed, testing different minimal media and culture conditions for the isolation of targeted strains of interest that can be spotted through 16S rRNA diversity assays.

While screening for the isolates' capability of secreting fibre and starch degrading enzymes, four *Bacillus* isolates showed the ability for simultaneously producing xylanase, β -glucanase, cellulase and amylase (Table 4). These bacteria may be of high interest for enzyme production and biotechnological applications due to their high growth rate, their ability to harbour multi-enzyme complexes and their steadiness at extreme conditions (Ladeira et al. 2015). *Bacillus* strains are usually able to utilise different complex mixtures of organic material by producing numerous extracellular enzymes that hydrolyse polysaccharides (Ladeira et al. 2015). In this study, seven *Bacilli* strains were able to secrete xylanases, the least common carbohydrase detected. Different studies have cloned and characterised xylanases from *B. subtilis* (Marimuthu et al. 2019), and many other *Bacillus* are known to harbour xylanases (Subramaniyan and Prema 2002). In this study, five *B. subtilis* have been isolated, but only three (2Aq, 7Aq and 23Aq) were positive for xylanase activity, whereas two isolates 8Aq and 13Aq were negative for this activity. This shows that species identification may not be a good predictor of their ability to express any given enzyme; these traits may only be traceable for a given strain and not evenly distributed for all strains within a given species. Shakir et al. (2020) reported a *B. licheniformis* producing xylanase, and Singh et al. (2020) described xylanase production from *B. pumilus*. Among the isolates obtained in this study, the other *Bacilli* expressing extracellular xylanase activity in agar media were *B. subtilis* 2Aq, *B. velezensis* 6Aq, *B. licheniformis* 15Aq, *B. subtilis* 23Aq, and *B. tequilensis* 18Aq and 26Aq.

In our study, eight of the isolates obtained could produce extracellular β -glucanase activity. *Bacillus* strains are known β -1,3-1,4-glucanases sources, and previously their enzymes have been characterised from different donor species such as *B. subtilis*, *B. licheniformis*, *B. brevis*, *B. halodurans* and *B. circulans* (Pauly and Keegstra 2008). Furthermore, fourteen *Bacilli* isolates showed the ability to produce cellulase. The production of cellulases was previously detected from several *Bacilli* such as *B. subtilis*, *B. cereus* and *B. circulans* (Irfan et al. 2017). *Bacilli* commonly produce amylases, and *B. firmus* (Elayaraja et al. 2011) and *B. subtilis* (Quesada-Ganuza et al. 2019) are among the most common reported amylase producing bacteria. Among the *Bacilli* bacterial isolates in this study, eleven isolates were able to secrete amylase, although this trait is considered a widespread trait in aquatic environments and of overall lower biotechnological interest. Many enzymes used in the industrial sector are produced by *Bacilli*, especially by *B. amyloliquefaciens*, *B. subtilis* and *B. licheniformis*, because they are safe to handle, produce high enzyme yields, and have good fermentation properties (van Dijk and Hecker 2013). Isolation of similar species in our study suggested that bacteria in aquaponics systems could be a good source of novel industrial enzymes.

The detection of phytase genes (PCR amplification using degenerate primers from environmental DNA extracted from the two compartments was only positive when using BPP primers, suggesting that β -propeller phytase class is widespread and of high ecological importance in the studied aquaponics systems. Previous studies of bacteria in aquaponics systems mostly focused on the presence of pathogens and chemo-litho-autotrophic nitrifiers (Rurangwa and Verdegem 2015). This study, for the first time showed the presence of microbes with β -propeller phytase, which suggests that bacteria, such as *Bacilli*, can increase extracellular phosphorus concentration (Khan et al. (2009). Extracellular phosphorus is necessary for plant mineralization and growth in the aquaponics systems.

Cheng and Lim (2006) reported that among the four phytases classes, only β -propeller was identified in aquatic environments. Similarly, Lim et al. (2007) showed that HAP and PTP phytases are uncommon in aquatic bacteria, but β -propeller phytases play a central role in phytate-phosphorus cycling in aquatic habitats. β -propeller phytases are typically active in neutral-alkaline pHs, use calcium as a cofactor and are typically produced by a wide range of *Bacilli* (Jorquera et al. 2018; Menezes-Blackburn et al. 2013). Curiously,

although most of the isolates obtained were from the *Bacillus* genus, the only two isolates with positive phytase production belonged to *Staphylococcus* and *Rosellomorea* genus. Phytase activity is arguably more challenging to detect than carbohydrases, because: a) phytases are often exclusively intracellularly expressed in bacteria; b) their expression is often triggered by different environmental stresses (P deficiency, anaerobioses, etc.); and c) agar plate screening methods may produce false positives (Menezes-Blackburn et al. 2013). Furthermore, phytase from *Bacillus* have been proposed as feed additives for fish diets; The supplementation of 300 U Kg⁻¹ of *Bacillus* phytase was equivalent to the supplementation of 1000 U Kg⁻¹ acidic commercial phytase (Fu et al. 2008). β -proteins are considered good candidates for fish feed applications due to their optimal pH (6-7.5), while PAPs and HAPs often have optimal pH in the acidic range (2.5-5.5) (Cheng and Lim 2006; Menezes-Blackburn and Greiner 2015). Thus, phytase from microbes inhabiting aquaponics systems could be of high economical importance for fish feed formulations.

5. Conclusions

The studied aquaponics systems were highly diverse in their microbial community compositions, and the strong variations in the microbial communities within replicates prevented us from statistically demonstrating differences between communities present in fish and plant compartments. Evidence from DNA sequencing and biochemical assays performed on isolated strains showed that, among the fibre-degrading enzymes, cellulases are the most common enzymes expressed, followed by β -glucanase and xylanase. Phytases production was a far less common trait, with only two isolates showing a consistent increase in extracellular phosphate when grown in broth media supplemented with phytate. β -proteins appear to be the most ecologically relevant phytase class in our aquaponics systems. Further examination of isolates that showed different enzyme activities are needed (such as pH range of activity of the different enzymes detected) to assert their potential use either in aquaponics systems or as animal feed additives. Bacilli bacteria are here demonstrated to play a critical role in organic matter cycling in aquaponics systems, which can be a valuable source of niche microbes carrying carbohydrases and phytases enzymes with possible biotechnological applications.

Author Contributions: Daniel Menezes-Blackburn: Conceptualization, methodology, formal analysis, data curation, writing—draft preparation, supervision, project administration, funding acquisition; Nahad Al-Mahrouqi: laboratory analysis, writing—original draft preparation; Buthaina Al-Siyabi: laboratory analysis; Adhari Al-Kalbani: laboratory analysis, Ralf Greiner: writing—review and editing; Sergey Dobretsov: writing—review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Sultan Qaboos University through the project Screening of Omani desert soils for thermostable phytases (IG/AGR/SWAE/19/02) and by the The Research Council (TRC) of the Sultanate of Oman through the project Thermostable phytases and carbohydrases of Omani desert soils and their potential for biotechnological application as poultry feed supplements (RC/RG-AGR/SWAE/19/01).

Institutional Review Board Statement: Not applicable

Acknowledgments: In this section, you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments).

Conflicts of Interest: The authors declare no conflict of interest.

References

Al-Hafedh YS, Alam A, Beltagi MS (2008) Food production and water conservation in a recirculating aquaponic system in Saudi Arabia at different ratios of fish feed to plants Journal of the world aquaculture society 39:510-520

- Bandi MT-R, Zugravu AG, Turek A, Rahoveanu DNM Water Quality in Aquaponic Integrated Systems: An Overview of the Literature
- Buzby KM, Waterland NL, Semmens KJ, Lin L-S (2016) Evaluating aquaponic crops in a freshwater flow-through fish culture system *Aquaculture* 460:15-24
- Cheng C, Lim BL (2006) Beta-propeller phytases in the aquatic environment *Archives of microbiology* 185:1-13
- Council NR NRC.(2011). Nutrient Requirements of fish and shrimp Washington, DC
- da Silva Cerozi B, Fitzsimmons K (2017) Effect of dietary phytase on phosphorus use efficiency and dynamics in aquaponics *Aquaculture international* 25:1227-1238
- Delaide B, Goddek S, Gott J, Soyeurt H, Jijakli MH (2016) Lettuce (*Lactuca sativa* L. var. Sucrine) growth performance in complemented aquaponic solution outperforms hydroponics *Water* 8:467
- Eck M, Sare AR, Massart S, Schmutz Z, Junge R, Smits TH, Jijakli MH (2019) Exploring bacterial communities in aquaponic systems *Water* 11:260
- Fooladi J, Sajjadian A (2010) Screening the thermophilic and hyperthermophilic bacterial population of three Iranian hot-springs to detect the thermostable α -amylase producing strain Iranian journal of microbiology 2:46
- Francis G, Makkar HP, Becker K (2001) Antinutritional factors present in plant-derived alternate fish feed ingredients and their effects in fish *Aquaculture* 199:197-227
- Fu S, Sun J, Qian L, Li Z (2008) Bacillus phytases: present scenario and future perspectives *Applied Biochemistry and Biotechnology* 151:1-8
- Goddek S, Espinal CA, Delaide B, Jijakli MH, Schmutz Z, Wuertz S, Keesman KJ (2016) Navigating towards decoupled aquaponic systems: a system dynamics design approach *Water* 8:303
- Hammer Ø, Harper DA, Ryan PD (2001) PAST: Paleontological statistics software package for education and data analysis *Palaeontologia electronica* 4:9
- Heuer H, Krsek M, Baker P, Smalla K, Wellington E (1997) Analysis of actinomycete communities by specific amplification of genes encoding 16S rRNA and gel-electrophoretic separation in denaturing gradients *Applied environmental microbiology* 63:3233-3241
- Hien TTT, Be TT, Lee CM, Bengtson DA (2015) Development of formulated diets for snakehead (*Channa striata* and *Channa micropeltes*): Can phytase and taurine supplementation increase use of soybean meal to replace fish meal? *Aquaculture* 448:334-340
- Hill JE, Kysela D, Elimelech M (2007) Isolation and assessment of phytate-hydrolysing bacteria from the DelMarVa Peninsula *J Environmental Microbiology* 9:3100-3107
- Huang H, Shi P, Wang Y, Luo H, Shao N, Wang G, Yang P, Yao B (2009) Diversity of beta-propeller phytase genes in the intestinal contents of grass carp provides insight into the release of major phosphorus from phytate in nature *Applied and environmental microbiology* 75:1508-1516
- Huang H, Zhang R, Fu D, Luo J, Li Z, Luo H, Shi P, Yang P, Diao Q, Yao B (2011) Diversity, abundance and characterization of ruminal cysteine phytases suggest their important role in phytate degradation *Environmental microbiology* 13:747-757

- Irfan M, Mushtaq Q, Tabssum F, Shakir HA, Qazi JI (2017) Carboxymethyl cellulase production optimization from newly isolated thermophilic *Bacillus subtilis* K-18 for saccharification using response surface methodology *AMB Express* 7:1-9
- Joo H-S, Hirai M, Shoda M (2005) Characteristics of ammonium removal by heterotrophic nitrification-aerobic denitrification by *Alcaligenes faecalis* No. 4 *Journal of Bioscience and Bioengineering* 100:184-191
- Jorquera M, MARTÍNEZ O, Maruyama F, Marschner P, de la Luz Mora M (2008) Current and future biotechnological applications of bacterial phytases and phytase-producing bacteria *Microbes and environments* 23:182-191
- Jorquera MA, Gabler S, Inostroza NG, Acuña JJ, Campos MA, Menezes-Blackburn D, Greiner R (2018) Screening and characterization of phytases from bacteria isolated from Chilean hydrothermal environments *Microbial ecology* 75:387-399
- Kasozi N, Kaiser H, Wilhelmi B (2020) Metabarcoding analysis of bacterial communities associated with media grow bed zones in an aquaponic system *International journal of microbiology* 2020
- Khan AA, Jilani G, Akhtar MS, Naqvi SMS, Rasheed M (2009) Phosphorus solubilizing bacteria: occurrence, mechanisms and their role in crop production *J agric biol sci* 1:48-58
- Köhler J, Baumbach J, Taubert J, Specht M, Skusa A, Rüegg A, Rawlings C, Verrier P, Philippi S (2006) Graph-based analysis and visualization of experimental results with ONDEX *Bioinformatics* 22:1383-1390
- Krogdahl Å, Hemre GI, Mommsen T (2005) Carbohydrates in fish nutrition: digestion and absorption in postlarval stages *Aquaculture nutrition* 11:103-122
- Ladeira SA, Cruz E, Delatorre AB, Barbosa JB, Leal Martins ML (2015) Cellulase production by thermophilic *Bacillus* sp: SMIA-2 and its detergent compatibility *Electronic journal of biotechnology* 18:110-115
- Leonard N, Blancheton J, Guiraud J (2000) Populations of heterotrophic bacteria in an experimental recirculating aquaculture system *Aquacultural Engineering* 22:109-120
- Leonard N, Guiraud J, Gasset E, Cailleres J, Blancheton J (2002) Bacteria and nutrients—nitrogen and carbon—in a recirculating system for sea bass production *Aquacultural Engineering* 26:111-127
- Liebert F, Portz L (2005) Nutrient utilization of Nile tilapia *Oreochromis niloticus* fed plant based low phosphorus diets supplemented with graded levels of different sources of microbial phytase *Aquaculture* 248:111-119
- Maravi P, Kumar A (2020) Isolation, Screening and Identification of Cellulolytic Bacteria from Soil *Biotechnology Journal International*:1-8
- Marimuthu M, Sorimuthu A, Muruganantham S (2019) Production and Optimization of Xylanase Enzyme from *Bacillus subtilis* using Agricultural Wastes by Solid State Fermentation *International Journal of Pharmaceutical Investigation* 9:169-173
- Menezes-Blackburn D, Greiner R (2014) Enzymes used in animal feed: leading technologies and forthcoming developments. Wiley Online Library,
- Menezes-Blackburn D, Greiner R (2015) Enzymes used in animal feed: leading technologies and forthcoming developments. In: *Functional Polymers in Food Science*, Scrivener Publishing, vol 200. pp 47-73

- Menezes-Blackburn D, Jorquera MA, Greiner R, Gianfreda L, de la Luz Mora M (2013) Phytases and phytase-labile organic phosphorus in manures and soils *Crit Rev Environ Sci Technol* 43:916-954
- Munguia-Fragozo P, Alatorre-Jacome O, Rico-Garcia E, Torres-Pacheco I, Cruz-Hernandez A, Ocampo-Velazquez RV, Garcia-Trejo JF, Guevara-Gonzalez RG (2015) Perspective for aquaponic systems: “omic” technologies for microbial community analysis *BioMed research international* 2015
- Nishioka T, Elsharkawy MM, Suga H, Kageyama K, Hyakumachi M, Shimizu M (2016) Development of culture medium for the isolation of *Flavobacterium* and *Chryseobacterium* from rhizosphere soil *Microbes and environments:ME*15144
- Ohno T, Zibilske LM (1991) Determination of low concentrations of phosphorus in soil extracts using malachite green *Soil Science Society of America Journal* 55:892-895
- Ondov BD, Bergman NH, Phillippy AM (2011) Interactive metagenomic visualization in a Web browser *BMC bioinformatics* 12:1-10
- Pardee AB, Jacob F, Monod J (1959) The genetic control and cytoplasmic expression of “inducibility” in the synthesis of β -galactosidase by *E. coli* *Journal of Molecular Biology* 1:165-178
- Pauly M, Keegstra K (2008) Cell-wall carbohydrates and their modification as a resource for biofuels *The Plant Journal* 54:559-568
- Rakocy J, Masser MP, Losordo T (2016) Recirculating aquaculture tank production systems: aquaponics-integrating fish and plant culture
- Rurangwa E, Verdegem MC (2015) Microorganisms in recirculating aquaculture systems and their management *Reviews in aquaculture* 7:117-130
- Schmautz Z, Graber A, Jaenicke S, Goesmann A, Junge R, Smits TH (2017) Microbial diversity in different compartments of an aquaponics system *Archives of microbiology* 199:613-620
- Shakir HA, Anwar A, Irfan M, Khan M, Ali S, Qazi JI (2020) Statistical Optimization of Xylanase from *Bacillus licheniformis* Using Banana Peels in Submerged Fermentation *Iranian Journal of Science and Technology, Transactions A: Science*:1-11
- Sharrer MJ, Summerfelt ST, Bullock GL, Gleason LE, Taeuber J (2005) Inactivation of bacteria using ultraviolet irradiation in a recirculating salmonid culture system *Aquacultural Engineering* 33:135-149
- Singh A, Sharma D, Varghese LM, Mahajan R (2020) Fast flow rate processes for purification of alkaline xylanase isoforms from *Bacillus pumilus* AJK and their biochemical characterization for industrial application purposes *Biotechnology progress* 36:e2898
- Subajini M, Sandrasegarampillai B, Vasanthy A (2012) Screening and identification of a thermophilic and alkalophilic bacterium producing xylanase *Advances in Applied Science Research* 3:242-250
- Subramaniyan S, Prema P (2002) Biotechnology of microbial xylanases: enzymology, molecular biology, and application *Critical reviews in biotechnology* 22:33-64

Sugita H, Nakamura H, Shimada T (2005) Microbial communities associated with filter materials in recirculating aquaculture systems of freshwater fish *Aquaculture* 243:403-409

Turkmen G, Guner Y Aquaponic (integrating fish and plant culture) systems. In: International Symposium on Sustainable Development, Science Book, 2010. pp 657-666

Tye A, Siu F, Leung T, Lim B (2002) Molecular cloning and the biochemical characterization of two novel phytases from *B. subtilis* 168 and *B. licheniformis* *Applied Microbiology and Biotechnology* 59:190-197

Tyson RV, Treadwell DD, Simonne EH (2011) Opportunities and challenges to sustainability in aquaponic systems *HortTechnology* 21:6-13

van Dijk J, Hecker M (2013) *Bacillus subtilis*: from soil bacterium to super-secreting cell factory. *BioMed Central*,