

Article

Benzaldehyde attenuates the fifth stage larval excretory-secretory product of *Angiostrongylus cantonensis*-induced injury in mouse astrocytes via regulation of endoplasmic reticulum stress and oxidative stress

Kuang-Yao Chen ^{1†}, Yi-Ju Chen ¹, Chien-Ju Cheng ², Kai-Yuan Jhan ³, and Lian-Chen Wang ^{2,3,4†}

¹ Department of Parasitology, School of Medicine, China Medical University, Taichung, 404, Taiwan; kychen@mail.cmu.edu.tw (K.Y.C.); yiru830318@gmail.com (Y.J.C.)

² Department of Parasitology, College of Medicine, Chang Gung University, Taoyuan 333, Taiwan; d000014521@cgu.edu.tw (C.J.C.); wanglc@mail.cgu.edu.tw (L.C.W.)

³ Graduate Institute of Biomedical Sciences, College of Medicine, Chang Gung University, Taoyuan 333, Taiwan; ed790101@hotmail.com (K.Y.J.); wanglc@mail.cgu.edu.tw (L.C.W.)

⁴ Molecular Infectious Disease Research Center, Chang Gung Memorial Hospital, Taoyuan, Taiwan. wanglc@mail.cgu.edu.tw (L.C.W.)

† Correspondence: kychen@mail.cmu.edu.tw (K.Y.C.); wanglc@mail.cgu.edu.tw (L.C.W.)

Abstract: Excretory-secretory products (ESPs) are the main research targets for investigating the hosts and helminths interaction. Parasitic worms can migrate to parasitic sites and avoid the host immune response by secreting this product. *Angiostrongylus cantonensis* is an important food-borne zoonotic parasite that causes severe neuropathological damage and symptoms, including eosinophilic meningitis or meningoencephalitis in humans. Benzaldehydes are organic compounds composed of a benzene ring and formyl substituents. This compound has anti-inflammatory and anti-oxidation properties. Previous studies showed that 3-hydroxybenzaldehyde (3-HBA) and 4-hydroxybenzaldehyde (4-HBA) can reduce apoptosis in *A. cantonensis* ESPs treated astrocytes. These results on the protective effect underlying benzaldehyde have primarily focused on cell survival. The study was designed to investigate the molecular mechanisms of endoplasmic reticulum stress (ER stress) and oxidative stress in astrocytes in *A. cantonensis* ESPs treated astrocytes and to evaluate the therapeutic consequent of 3-HBA and 4-HBA. First, we initially established the RNA-seq dataset in each group, including Normal, ESPs, ESPs+3-HBA, and ESPs+4-HBA. We also found that benzaldehyde (3-HBA and 4-HBA) can stimulate astrocytes to express ER stress-related molecules after ESP treatment. The level of oxidative stress could also be decreased in astrocytes by elevating antioxidant activity and reducing ROS generation. These results suggested that benzaldehyde may be a potential therapeutic compound for human angiostrongyliasis to support brain cell survival by inducing the expression levels of ER stress- and oxidative stress-related pathway.

Keywords: Excretory-secretory products; *Angiostrongylus cantonensis*; Astrocytes; Benzaldehydes; endoplasmic reticulum stress; oxidative stress

1. Introduction

Angiostrongyliasis is a serious parasitic disease worldwide. It is caused by the parasitic helminth *Angiostrongylus cantonensis*. The life cycle of this nematode requires the rats and molluscan [1]. Adults live in the blood vessel of rat cardiopulmonary, and then eggs are become to the first-stage larvae (L1). Finally, L1 were released via rat faeces. After ingestion by the intermediate host, L1 can migrate to the muscle tissue and develop into infective third-stage larvae (L3). After rats eat the molluscan containing L3, the larvae can reach the host brain through blood circulation and develop into fifth-stage larvae (L5). Humans are the abnormal host for these worms through infection by ingesting infective

third-stage larvae (L3) in snails, terrestrial crabs, freshwater shrimp, frogs, and fish [2–5]. This parasite can induce eosinophilic meningitis/meningoencephalitis and cause some symptoms, including fever, headache, nausea, vomiting, neck stiffness, and paraesthesia [3]. Recently, human angiostrongyliasis has been known an emerging disease in many countries, such as China, America, Europe, Australia, Jamaica, and Brazil [6–11].

The treatment of angiostrongyliasis is limited to chemical anthelmintic or immunosuppressive reagents, which are often associated with severe side effects. At present, albendazole and corticosteroids are common therapeutic drugs for eosinophilic meningitis caused by *A. cantonensis* [12, 13]. Thus, the identification of more effective and safe drugs is highly necessary. *Gastrodia elata* (Tianma) is a Chinese herbal medicine that has antioxidative stress and anti-inflammation functions in humans. It has also been utilized for the treatment of headaches and dizziness [14].

In nature, Benzaldehyde has the anti-oxidation and –inflammatory function [15, 16]. 3-HBA and 4-HBA are the benzaldehyde from *G. elata*. Previous studies demonstrated that 3-HBA has the function of anti-oxidative stress, vasculoprotection, and inflammation suppression [16, 17]. On the other hand, 4-HBA is also the important bioactive constituent of *G. elata* and can promote antioxidant activation, wound repairment, and migration [18–20]. In protozoan parasite *Toxoplasma gondii* infection, 4-HBA inhibits *Toxoplasma gondii* growth in macrophages by inducing autophagy activation [21].

Our previous results found that the survival rate of astrocytes was increased after 3-HBA and 4-HBA treatment. Further results demonstrated that 3-HBA and 4-HBA can inhibit apoptosis-related molecule expression. Therefore, 3-HBA and 4-HBA may be the potential therapeutic molecules for human angiostrongyliasis treatment [22]. In this research, we want to evaluate the utility of 3-HBA and 4-HBA after ESPs treatment in astrocytes. The data demonstrated that the ESPs can cause endoplasmic reticulum stress (ER stress) and oxidative stress. Moreover, we found that 3-HBA and 4-HBA suppress ROS generation through the induction of antioxidant activation. These results suggest that benzaldehyde an effective compound for reducing oxidative stress caused by *A. cantonensis* ESPs treatment in astrocytes.

2. Materials and Methods

2.1. Ethics

The animal experiments were conducted in accordance with the standards of the Chang Gung University Institutional Animal Care and Use Committee (IACUC) in Taiwan (CGU109-198). Rats were kept in plastic cages and provided with food and water ad libitum.

2.2. Parasite and Experimental Infection

The lifecycle of *A. cantonensis* was maintained by *Biomphalaria glabrata* snails and Sprague–Dawley (SD) rats in laboratory [23]. Male SD rats were purchased from the National Laboratory Animal Center, Taipei. In *A. cantonensis* infection, the L3 were collected from infected snails by digestion with 0.6% (w/v) pepsin-HCl (pH 2–3) for 1 h after day 21 postinfection. SD rats were inoculated with L3 by stomach intubation.

2.3. *A. cantonensis* excretory/secretory product preparation

A. cantonensis L5 was collected from the host brains of infected rats. Next, the worms were incubated in fetal bovine serum (FBS)-free RPMI after washing with saline, phosphate buffered saline, ddH₂O and RPMI. *A. cantonensis* L5 ESPs were concentrated by centrifugal filter devices (Merck Millipore, Germany). The ESPs concentration was determined by a Bio–Rad Protein Assay Kit (Bio–Rad, Hercules, CA, USA) according to the instructions of the manufacturer. Finally, it was employed to treat astrocytes and then observe cell survival and gene and protein expression.

2.4. Astrocyte culture

The CRL2535 (mouse brain astrocytes) was cultured at 37 °C in Dulbecco's modified Eagle's medium/F-12 (DMEM/F-12) (Corning, USA) and with 10% fetal bovine serum (FBS) (Gibco, USA) in poly-L-lysine-coated culture flasks. The cells for assays were harvested from a 10 cm culture dish containing serum-free DMEM/F-12. Finally, the cells were pretreated with the drugs for 2 h and then stimulated to the ESPs of *A. cantonensis* L5.

2.5. Extraction of total RNA and cDNA synthesis

The total RNA of astrocytes cultured in medium containing ESPs and drugs was extracted as follows. TRIzol® Reagent was used to isolate total RNA from different treatments. Finally, the RNA was redissolved in DEPC-treated water. Next, the iScript™ Advanced cDNA Synthesis Kit (Bio-Rad, USA) was used to generate the cDNA.

2.6. RNA-sequencing (RNA-seq) library preparation and sequencing

The RNA was utilized to the preparation of the sequencing library by the TruSeq Stranded mRNA Library Prep Kit (Illumina, San Diego, CA, USA) following the manufacturer's recommendations. First, reverse transcriptase and random primers were used to generate First-strand cDNA, and the adaptors were ligated and purified with the AM-Pure XP system (Beckman Coulter, Beverly, USA). Agilent Bioanalyzer 2100 system was employed to evaluate the libraries quality. Finally, Illumina NovaSeq 6000 platform was employed to sequence the libraries (150 bp paired-end reads) (Genomics, Taiwan).

2.7. Bioinformatics Analysis

The filtered reads were aligned and quantified by Bowtie2 (version 2.3.4.1) and RSEM (version 1.2.28). The functional analysis, including GO and KEGG were carried out of GO terms and KEGG pathways [24–30].

2.8. Quantitative real-time PCR

The iQ™ SYBR® Green Supermix (Bio-Rad, USA) was employed to Quantitative real-time PCR on Real-Time PCR Detection System (Bio-Rad, USA). GAPDH is an internal control.

2.9. Protein extraction and Western blotting

The cells could be disrupted in lysis buffer (RIPA buffer) containing protease inhibitors (Sigma-Aldrich, USA). The proteins in 12% SDS-PAGE were transferred to NC membranes and incubated with antibodies against catalase (ABclonal, USA), glutathione reductase (ABclonal, USA), superoxide dismutase (ABclonal, USA), glutathione transferase kappa (ABclonal, USA), GRP78 (Proteintech, USA), IRE1 (Signalway Antibody, USA), ATF6 (Proteintech, USA), CHOP (Proteintech, USA), PERK (Proteintech, USA), eIF2α (Proteintech, USA), ATF4 (Proteintech, USA), and β-actin (Proteintech, USA). Next, the membrane was incubated with HRP-linked secondary antibody (Sigma-Aldrich, USA). ECL reagent (EMD Millipore, USA) and a ChemiDoc Imaging System (Bio-Rad, USA) were employed to detect and capture the immunoreactive bands. Finally, ImageJ was used to quantify the image densitometry.

2.10. Catalase activity assay

The activity of catalase in astrocytes was determined by using a catalase assay kit (Cayman, USA) according to the manufacturer's instructions. The activities were measured using a microplate reader by reading the absorbance at 540 nm.

2.11. Superoxide dismutase (SOD) activity assay

SOD activities in astrocytes were determined by using a SOD assay kit (Cayman, USA) according to the manufacturer's instructions. The activities were measured using a

microplate reader by reading the absorbance at 440-160 nm before incubation of the plate on a shake for 30 minutes.

2.12. Glutathione S-transferase (GST) activity assay

GST activities in astrocytes were determined by using a GST assay kit (Cayman, USA) according to the manufacturer’s instructions. The activities were measured at least 5 time points using a microplate reader by reading the absorbance at 340 nm.

2.13. ROS detection

ROS generation in astrocytes was detected by using the ROS Detection Cell-Based assay kit (Cayman, USA) according to the manufacturer’s instructions. For nuclear counterstaining, the cells were stained with DAPI. Finally, the samples were observed and photographed under a fluorescence microscope.

2.14. Antioxidant capacity assay

An antioxidant assay kit (Cayman, USA) was employed to detect antioxidant capacities in astrocytes. Controls and ESPs-, 3-HBA-, and 4-HBA-treated astrocytes (1 × 10⁷ cells/ml) were homogenized or sonicated in assay buffer. Cell lysate, Metmyoglobin, Chromogen, and Hydrogen Peroxide were added and incubated in the wells on the plate. Antioxidant capacity was detected using a spectrofluorometer (Bio–Rad, USA) by reading the absorbance at 405 or 750 nm.

2.15. Statistical Analysis

we used Student’s t-test to compare the expression levels by GraphPad Prism 5 software (GraphPad, USA). Data were showed as the mean ± standard deviation. The level of statistical significance was *P* < 0.05, < 0.01, and < 0.001.

3. Results

3.1. Transcriptome profile of mouse astrocytes after *A. cantonensis* L5 ESPs or benzaldehyde treatment.

First, we initially established the RNA-seq dataset in each treatment group, including normal, ESPs, ESPs+0.1 mM 3-HBA, ESPs+0.5 mM 3-HBA, 0.1 mM 4-HBA, and 0.5 mM 4-HBA. We obtained a total of 123,138,219 raw reads from polyA-tailed mRNA using the NovaSeq 6000 Sequencing System with 150 bp paired-end reads (Illumina, USA) (Table 1, Table S1). The transcriptome results showed that the expression of inflammation-related genes, such as serum amyloid A3, chemokines, interleukin 6, interleukin 12, and colony stimulating factor, was obviously elevated in astrocytes after *A. cantonensis* L5 ESPs treatment. In contrast, the expression of inflammation-related genes was reduced after 3-HBA or 4-HBA treatment (Table S2). These results suggested that benzaldehyde can reduce *A. cantonensis* L5 ESPs-induced inflammation in astrocytes.

Table 1. Characteristics of the transcriptome of mouse brain astrocytes after excretory/secretory product or benzaldehyde treatment.

Sample_Name	GC.	Length	Length_Mean	Phred	Q20_Ratio	Q30_Ratio	Qual_Mean	Read_Counts	Total_Bases
N_R1	48.65%	20-151	144.32	33	99.04%	96.29%	36.42	25,920,999	3,740,844,537
N_R2	48.78%	20-151	143.69	33	98.43%	94.35%	36.10	25,920,999	3,724,577,535
E_R1	48.00%	20-151	144.66	33	98.97%	96.05%	36.38	23,617,944	3,416,535,104
E_R2	48.22%	20-151	144.22	33	98.61%	94.87%	36.19	23,617,944	3,406,183,730
3H1_R1	48.48%	20-151	144.94	33	99.00%	96.18%	36.4	23,470,406	3,401,781,446
3H1_R2	48.68%	20-151	143.85	33	98.28%	94.04%	36.04	23,470,406	3,376,330,842

3H5_R1	49.22%	20-151	145.23	33	99.03%	96.28%	36.42	23,669,860	3,437,686,443
3H5_R2	49.38%	20-151	144.55	33	98.42%	94.38%	36.1	23,669,860	3,421,432,553
4H1_R1	48.82%	20-151	144.87	33	98.99%	96.18%	36.4	27,305,448	3,955,655,684
4H1_R2	48.97%	20-151	143.86	33	98.31%	94.10%	36.06	27,305,448	3,928,240,745
4H5_R1	49.20%	20-151	145.02	33	99.02%	96.25%	36.41	25,074,561	3,636,192,527
4H5_R2	49.37%	20-151	144.24	33	98.32%	94.09%	36.06	25,074,561	3,616,740,196

As shown in Fig. 1, the differentially expressed genes in each comparison were determined by RSEM and EBSeq (PPEE <= 0.05). In total, 4753 genes were shown in the union of comparisons, including E/N, 3H/E, and 4H/E (Figure 1A, B, C, and Table S2). Moreover, we used a heatmap to demonstrate the expression profiles of both union and intersect sets (Figure 1D). Next, we started with functional enrichment analysis of these genes. All unigene sequences were matched against Gene Ontology (GO) terms and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways (Figure 2 and Table S3).




Figure 1. Summary of the transcriptome profile of mouse brain astrocytes after excretory/secretory product or benzaldehyde treatment. (A) Boxplot of FPKM distribution among samples. N: Normal, E: 250 µg/ml ESPs treatment, 3H1: ESPs+0.1 mM 3-HBA, 3H5: ESPs+0.5 mM 3-HBA, 4H1: ESPs+0.1 mM 4-HBA, 4H5: ESPs+0.5 mM 4-HBA. (B) Number of DEGs in comparison sets E/N, 3H/E, and 4H/E. (C) Venn diagram was used to show the number of differentially expressed genes in each comparison determined by RSEM and EBSeq (PPEE <= 0.05). 4753 genes were shown in the union of comparisons. (D) A heatmap was employed to demonstrate the expression profiles of both union and intersect sets.




Figure 2. GO annotations of unigenes in transcriptome profile. (A) E compared to N. (B) 3H compared to E. (C) 4H compared to E.

3.2. The expression of ER stress-related molecules after 3-HBA and 4-HBA treatment.

To confirm whether 3-HBA and 4-HBA could induce ER stress-related gene expression. As shown in Figure 3, we used quantitative real-time PCR to measure the expression of genes after 3-HBA and 4-HBA treatment. The results showed that the mRNA expression levels of ER stress signalling molecules (GRP78, IRE1, ATF6, CHOP, PERK, eIF2, and ATF4) were significantly elevated after ESPs treatment. Moreover, the expression level of ER stress signalling was also significantly elevated after 0.5 mM 3-HBA and 4-HBA treatment compared with ESPs alone. On the other hand, we used western blotting to detect the protein expression levels of ER stress signalling in astrocytes (Figure 4). The results showed that the protein expression levels of ER stress signalling were also significantly elevated in ESPs alone or 3-HBA and 4-HBA (0.1 and 0.5 mM) treatment. Therefore, we suggested that HBA can stimulate the activation of the ER stress signalling pathway in astrocytes after *A. cantonensis* L5 ESPs treatment.




Figure 3. 3-HBA and 4-HBA induce the expression of ER stress-related genes in ESPs-treated astrocytes. Cells were pre-treated with 3-HBA and 4-HBA (0.1 and 0.5 mM) for 4 h and then incubated with 250 µg/ml ESPs for 12 h. mRNA expression were determined by real-time qPCR (n=3). [#]*P*< 0.05, ^{##}*P*< 0.001, compared to control. ^{**}*P*< 0.01, ^{***}*P*< 0.001, compared to cells exposed to ESPs.




Figure 4. 3-HBA and 4-HBA induce the expression of ER stress-related proteins in ESPs-treated astrocytes. Cells were pretreated with 3-HBA and 4-HBA (0.1 and 0.5 mM) for 4 h and then incubated with 250 µg/ml ESPs for 12 h. Protein expression levels were determined by western blotting (n=3). [#]*P*< 0.05, ^{##}*P*< 0.001, compared to control. ^{**}*P*< 0.01, ^{***}*P*< 0.001, compared to cells exposed to ESPs.

3.3. The expression of oxidative stress-related molecules after 3-HBA and 4-HBA treatment.

First, we wanted to determine whether benzaldehydes can protect astrocytes in ESPs treatment by activating antioxidants. Quantitative real-time PCR and western blotting were employed to detect the gene and protein expression of antioxidants after ESPs treatment alone or 3-HBA and 4-HBA treatment (Figures 5 and 6). The data showed that the expression of antioxidants (CAT, GSR, SOD, and GSTK) was significantly elevated in ESPs alone and increased in 3-HBA and 4-HBA (0.1 mM) treatment. The data demonstrated that HBA can induce the oxidative stress-related molecules expression after *A. cantonensis* L5 ESPs treatment.




Figure 5. 3-HBA and 4-HBA induce the gene expression of antioxidants in ESPs-treated astrocytes. Cells were pretreated with 3-HBA and 4-HBA (0.1 and 0.5 mM) for 4 h and then incubated with 250 µg/ml ESPs for 12 h. mRNA expression were determined by real-time qPCR (n=3). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, compared to the control. ** $P < 0.01$, *** $P < 0.001$, compared to cells exposed to ESPs.




Figure 6. 3-HBA and 4-HBA induce the protein expression of antioxidants in ESPs-treated astrocytes. Cells were pretreated with 3-HBA and 4-HBA (0.1 and 0.5 mM) for 4 h and then incubated with 250 µg/ml ESPs for 12 h. Protein expression levels of antioxidants were detected by western blotting (n=3). ** $P < 0.01$, *** $P < 0.001$, compared to control. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, compared to cells exposed to ESPs.

3.4. The activity of antioxidant after 3-HBA and 4-HBA treatment.

To confirm the effect of 3-HBA and 4-HBA on the activity of antioxidants in astrocytes, we employed ELISA to examine the activity of oxidative-related enzymes, such as CAT, SOD, and GST. As shown in Figure 7, the ESPs of *A. cantonensis* L5 induced antioxidant activation in astrocytes. Moreover, we found that the activity of antioxidants was elevated after 3-HBA and 4-HBA treatment compared with ESPs alone. Therefore, we propose that HBA reduce oxidative stress by inducing the activation of antioxidants in astrocytes.




Figure 7. 3-HBA and 4-HBA induce antioxidant activation in ESPs-treated astrocytes. Cells were pretreated with 3-HBA and 4-HBA (0.1 and 0.5 mM) for 4 h and then incubated with 250 µg/ml ESPs for 12 h. Activities of catalase, SOD, and GST were determined by activity assay (n=3). $^{##}P < 0.01$, $^{###}P < 0.001$, compared to control. $^{*}P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$, compared to cells exposed to ESPs.

3.5. 3-HBA and 4-HBA inhibit ROS generation after ESPs treatment.

To monitor the effect of 3-HBA and 4-HBA on ROS generation in astrocytes, we investigated whether 3-HBA and 4-HBA could reduce ESPs-induced ROS generation in astrocytes (Figure 8). Cells were incubated with *A. cantonensis* L5 ESPs in the absence or presence of 3-HBA and 4-HBA, and then DHE (dihydroethidium) was added to stain ROS. Fluorescence microscopy was employed to detect ROS expression in astrocytes. The data showed that ROS expression was significantly elevated after ESPs treatment. However, 3-HBA and 4-HBA reduced ESPs-induced ROS expression. The data demonstrated that HBA can inhibit oxidative stress generation through decreasing ESPs-induced ROS generation.




Figure 8. 3-HBA and 4-HBA reduce ROS generation in ESPs-treated astrocytes. Cells were pretreated with 3-HBA and 4-HBA (0.1 and 0.5 mM) for 4 h and then incubated with 250 µg/ml ESPs for 12 h. An ROS detection kit was used to detect ROS generation by fluorescence microscopy (n=3).

3.6. 3-HBA and 4-HBA elevate the antioxidant capacity after ESPs treatment.

Finally, we want to determine whether 3-HBA and 4-HBA could increase the antioxidant capacity in astrocytes (Figure 9). The results demonstrated that the level of antioxidant capacity was significantly higher in the cells pretreated with 3-HBA and 4-HBA than in the cells treated with ESPs alone. These data suggested that the antioxidant capacity in astrocytes can be elevated after HBA treatment.




Figure 9. Elevation of antioxidant capacity in astrocytes treated with 3-HBA and 4-HBA. Cells were pretreated with 3-HBA and 4-HBA (0.1 and 0.5 mM) for 4 h and then incubated with 250 µg/ml ESPs for 12 h. An antioxidant assay kit was employed to detect antioxidant capacities in astrocytes (n=3). ****P*< 0.001, compared to control. **P*< 0.05, ****P*< 0.001, compared to cells exposed to ESPs.

4. Discussion

In the research, we determined the protective efficiency of benzaldehydes in *A. cantonensis* L5 ESPs treatment. ESPs are a mixture of helminths secreted and contain a wide range of molecules, including DNA, mRNA, microRNA, proteins, and carbohydrate [31, 32]. This secretion can help parasites penetrate host barriers and reduce inflammatory responses. Moreover, ESPs also play a crucial role in reproduction and nourishment intake [33–35]. Because of their importance in nematode pathogenesis, ESPs are an important target in the investigation of host-parasite relationships and research in vaccine candidates and therapeutic targets. In previous investigation, we demonstrated that *A. cantonensis* ESPs can stimulate oxidative stress, autophagy, ER stress, and cell apoptosis in astrocytes [36–38]. However, we used proteomic analysis to profile, identify, and characterize the proteins in the *A. cantonensis* L5 ESPs. MALDI-TOF/TOF analysis showed that a total of 51 spots were identified, such as disulfide isomerases and calreticulin. On the other hand, approximately 8 immunoreactive proteins were identified, including protein disulfide-isomerase, a putative aspartic protease, and annexin [39].

In this study, we demonstrated that benzaldehydes (3-HBA and 4-HBA) can activate endoplasmic reticulum (ER) stress signalling by stimulating related molecules, including IRE1, ATF6, CHOP, PERK, eIF2, and ATF4. Moreover, our results on the expression of GRP78 reveal that benzaldehydes induce ER stress in astrocytes. The ER is the intracellular organelle for calcium (Ca²⁺) storage and has multiple functions, such as protein folding, modification, lipid biosynthesis, and membrane biogenesis [40, 41]. The ER also plays a major role in the regulation of Ca²⁺ signalling, such as Ca²⁺-induced gene expression, synaptic plasticity, embryonic development, and skeletal muscle excitation–contraction coupling [42, 43]. The unfolded proteins accumulation in the ER could stimulate unfolded protein response (UPR) activation, and this phenomenon is also called ER stress. Finally, this stress can induce the activation of related signalling pathways [44–46]. ER stress can inhibit protein synthesis and aggregation by activating chaperones, and it can also degrade abnormal proteins by the proteasome pathway. On the other hand, ER stress may increase the cell survival rate via activation of the antiapoptotic pathway [47].

ER stress activation can also lead to priming protein generation of glucose-regulated protein 78 (GRP78). GRP78 is a major marker for the evaluation of ER stress in target cells. When GRP78 is activated, GRP78 disassociates from and subsequently activates three signal transducers, IRE1, PERK and ATF6, and then induces the following pathway [48]. IRE1 can upregulate protein chaperones by activating the RNA of XBP1. PERK also activates chaperones by activating eIF2 α and ATF4. Moreover, ATF6 induces cell cycle arrest by initiating gene expression programs. Finally, these mechanisms could elevate cell survival through downregulation of cell death-related molecule expression [49].

In this study, we demonstrated that oxidative stress is induced after *A. cantonensis* L5 ESPs treatment in astrocytes. However, activation of antioxidants may decrease oxidative stress by inhibiting ROS generation in the treatment of 3-HBA and 4-HBA. Oxidative stress plays a significant role in the pathogenesis of parasite infection, such as *Plasmodium falciparum*, *Trypanosoma cruzi*, *Toxoplasma gondii*, *Giardia lamblia*, *Eurytrema coelomaticum*, and *Ornithodiplostomum ptychocheilus* [50–57]. After infection, oxidative stress is induced in the host and then causes cell apoptosis, inflammation, and DNA breakage. The generation of oxidative stress as a result of reactive oxygen species (ROS) (H₂O₂, O₂⁻ and OH⁻) induces the impairment of antioxidant defences [58, 59]. In *A. cantonensis* infection, the expression of oxidative stress and antioxidants, including glutathione reductase (GR), glutathione peroxidase (GPx) and glutathione S-transferase (GST), was increased in the *A. cantonensis*-infected mouse central nervous system (CNS) [60, 61].

In our previous study, the results showed that 3-HBA and 4-HBA can induce the protective function via regulating the generation of cell apoptosis after the treatment of ESPs [22]. At present, the therapeutic strategy for cerebral angiostrongyliasis remains controversial. Albendazole, the anthelmintics, may be effective drugs for angiostrongyliasis treatment by inhibiting nematode survival. Recent *A. cantonensis* studies have shown that the administration of anthelmintics or supportive therapy with corticosteroids is also the dispute. In our previous study, we found that administration of albendazole and dexamethasone has effective utility for cerebral angiostrongyliasis by reducing clinical symptoms and pathological changes, including meningitis, encephalitis, haemorrhage, and worm survival [13]. The results showed that the combination of albendazole with dexamethasone is the most effective therapy for cerebral angiostrongyliasis. Moreover, this therapeutic strategy could reduce inflammation in the CNS. For example, the cytokines expression, such as IL-1 β , IL-4, and IL-5, was decreased in the CNS after drug treatment [62].

5. Conclusions

In conclusion, we have demonstrated that *A. cantonensis* L5 ESPs induce ER stress and oxidative stress in mouse astrocytes. However, we found that 3-HBA and 4-HBA can reduce ROS generation by stimulating antioxidant activation, which results in decreased oxidative stress. On the basis of our findings, Benzaldehyde may be potential molecules for the therapy of angiostrongyliasis.

Author Contributions: KYC and LCW coordinated and designed this research. KYC wrote the manuscript. KYC, YJC, CJC and KYJ carrying out the experiments. All authors read and approved the final version of the manuscript

Funding: This work was supported in part by grants from the Ministry of Science and Technology (MOST) (grant MOST 109-2320-B-039-054 and 110-2320-B-039-055), the Chang Gung Memorial Hospital Research Grant (CMRPD1H0442), and the China Medical University Research Grant (CMU109-S-45, CMU109-MF-115, and CMU110-MF-101).

Institutional Review Board Statement: All procedures involving the experimental animals and their care were reviewed and approved by the Chang Gung University Institutional Animal Care and Use Committee (IACUC).

Data Availability Statement: All data generated or analyzed during this study are available from the corresponding author upon reasonable request.

Acknowledgments: The authors thank Pi-kai Chang for *A. cantonensis* L5 ESPs support.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Wang, L.C.; Chao, D.; Chen, E.R. Experimental infection routes of *Angiostrongylus cantonensis* in mice. *Journal of Helminthology* **1991**, *65*, 296-300, doi: 10.1017/s0022149x00010890.
2. Ash, L.R. The occurrence of *Angiostrongylus cantonensis* in frogs of New Caledonia with observations on paratenic hosts of metastrongyles. *J. Parasitol.* **1968**, *54*, 432-436.
3. Wang, Q.P.; Lai, D.H.; Zhu, X.Q.; Chen, X.G.; Lun, Z.R. Human angiostrongyliasis. *Lancet Infect. Dis.* **2008**, *8*, 621-630, doi: 10.1016/S1473-3099(08)70229-9.
4. Wallace, G.D.; and Rosen, L. Studies on eosinophilic meningitis. 2. Experimental infection of shrimp and crabs with *Angiostrongylus cantonensis*. *Am. J. Epidemiol.* **1966**, *84*, 20-31.
5. Wallace, G.D.; and Rosen, L. Studies on eosinophilic meningitis. IV. Experimental infection of fresh-water and marine fish with *Angiostrongylus cantonensis*. *Am. J. Epidemiol.* **1967**, *85*, 395-402.
6. Barbosa, T.A.; Thiengo, S.C.; Fernandez, M.A.; Graeff-Teixeira, C.; Morassutti, A.L.; Mourão, F.R.P.; Miranda, C.O.S.; Jorge, M.M.; Costa, L.F.; Gomes, S.R. Infection by *Angiostrongylus cantonensis* in both humans and the snail *Achatina* (Lissachatina) fulica in the city of Macapá, in the Amazon Region of Brazil. *Mem. Inst. Oswaldo. Cruz.* **2020**, *115*, e200115, doi: 10.1590/0074-02760200115.
7. Federspiel, F.; Skovmand, S.; Skarphedinsson, S. Eosinophilic meningitis due to *Angiostrongylus cantonensis* in Europe. *Int. J. Infect. Dis.* **2020**, *93*, 28-39, doi: 10.1016/j.ijid.2020.01.012.
8. Hochberg, N.S.; Park, S.Y.; Blackburn, B.G.; Sejvar, J.J.; Gaynor, K.; Chung, H.; Leniek, K.; Herwaldt, B.L.; Effler, P.V. Distribution of eosinophilic meningitis cases attributable to *Angiostrongylus cantonensis*, Hawaii. *Emerg. Infect. Dis.* **2007**, *13*, 1675-1680, doi:10.3201/eid1311.070367.
9. Lv, S.; Guo, Y.H.; Wei, F.R.; Zhang, Y.; Xiao, N.; Zhou, X.N. Control of eosinophilic meningitis caused by *Angiostrongylus cantonensis* in China. *Adv. Parasitol.* **2020**, *110*, 269-288, doi:10.1016/bs.apar.2020.02.002
10. Morton, N.J.; Britton, P.; Palasanthiran, P.; Bye, A.; Sugo, E.; Kesson, A.; Ardern-Holmes, S.; Snelling, T.L. Severe hemorrhagic meningoencephalitis due to *Angiostrongylus cantonensis* among young children in Sydney, Australia. *Clin. Infect. Dis.* **2013**, *57*, 1158-1161, doi:10.1093/cid/cit444.
11. Waugh, C.A.; Lindo, J.F.; Lorenzo-Morales, J.; Robinson, R.D. An epidemiological study of *A. cantonensis* in Jamaica subsequent to an outbreak of human cases of eosinophilic meningitis in 2000. *Parasitology* **2016**, *143*, 1211-1217, doi:10.1017/S0031182016000640.
12. Diao, Z.; Chen, X.; Yin, C.; Wang, J.; Qi, H.; Ji, A. *Angiostrongylus cantonensis*: effect of combination therapy with albendazole and dexamethasone on Th cytokine gene expression in PBMC from patients with eosinophilic meningitis. *Exp. Parasitol.* **2009**, *123*, 1-5, doi:10.1016/j.exppara.2009.04.016.
13. Jhan, K.Y.; Cheng, C.J.; Jung, S.M.; Lai, Y.J.; Chen, K.Y.; Wang, L.C. Co-Therapy of Albendazole and Dexamethasone Reduces Pathological Changes in the Cerebral Parenchyma of Th-1 and Th-2 Dominant Mice Heavily Infected with *Angiostrongylus cantonensis*: Histopathological and RNA-seq Analyses. *Biomolecules* **2021**, *11*, 536, doi:10.3390/biom11040536.
14. Huang, N.K.; Chern, Y.; Fang, J.M.; Lin, C.I.; Chen, W.P.; Lin, Y.L. Neuroprotective principles from *Gastrodia elata*. *J. Nat. Prod.* **2007**, *70*, 571-574, doi:10.1021/np0605182.
15. Moon, C.Y.; Ku, C.R.; Cho, Y.H.; Lee, E.J. Protocatechuic aldehyde inhibits migration and proliferation of vascular smooth muscle cells and intravascular thrombosis. *Biochem. Biophys. Res. Commun.* **2012**, *423*, 116-121, doi:10.1016/j.bbrc.2012.05.092.
16. Kong, B.S.; Im, S.J.; Lee, Y.J.; Cho, Y.H.; Do, Y.R.; Byun, J.W.; Ku, C.R.; Lee, E.J. Vasculoprotective Effects of 3-Hydroxybenzaldehyde against VSMCs Proliferation and ECs Inflammation. *PLoS One* **2016**, *11*, e0149394, doi:10.1371/journal.pone.0149394.
17. Wang, R.S.; Nakajima, T.; Kawamoto, T.; Honma, T. Effects of aldehyde dehydrogenase-2 genetic polymorphisms on metabolism of structurally different aldehydes in human liver. *Drug Metab. Dispos.* **2002**, *30*, 69-73, doi:10.1124/dmd.30.1.69.
18. Ha, J.H.; Lee, D.U.; Lee, J.T.; Kim, J.S.; Yong, C.S.; Kim, J.A.; Ha, J.S.; Huh, K. 4-Hydroxybenzaldehyde from *Gastrodia elata* B1. is active in the antioxidation and GABAergic neuromodulation of the rat brain. *J. Ethnopharmacol.* **2000**, *73*, 329-333, doi:10.1016/s0378-8741(00)00313-5.
19. Kang, C.W.; Han, Y.E.; Kim, J.; Oh, J.H.; Cho, Y.H.; Lee, E.J. 4-Hydroxybenzaldehyde accelerates acute wound healing through activation of focal adhesion signalling in keratinocytes. *Sci. Rep.* **2017**, *7*, 14192, doi:10.1038/s41598-017-14368-y.
20. Oh, S.H.; Ryu, B.; Ngo, D.H.; Kim, W.S.; Kim, D.G.; Kim, S.K. 4-hydroxybenzaldehyde-chitoooligomers suppresses H2O2-induced oxidative damage in microglia BV-2 cells. *Carbohydr. Res.* **2017**, *440-441*, 32-37, doi:10.1016/j.carres.2017.01.007.
21. Lee, J.; Choi, J. W.; Han, H.Y.; Kim, W.S.; Song, H.Y.; Byun, E.B.; Byun, E.H.; Lee, Y.H.; Yuk, J.M. 4-Hydroxybenzaldehyde Restricts the Intracellular Growth of *Toxoplasma gondii* by Inducing SIRT1-Mediated Autophagy in Macrophages. *Korean. J. Parasitol.* **2020**, *58*, 7-14, doi:10.3347/kjp.2020.58.1.7.
22. Chen, K.Y.; Chen, Y.J.; Cheng, C.J.; Jhan, K.Y.; Chiu, C.H.; Wang, L.C. 3-Hydroxybenzaldehyde and 4-Hydroxybenzaldehyde enhance survival of mouse astrocytes treated with *Angiostrongylus cantonensis* young adults excretory/secretory products. *Bio-med. J.* **2020**, doi.org/10.1016/j.bj.2020.11.008.
23. Wang, L.C.; Jung, S.M.; Chen, K.Y.; Wang, T.Y.; Li, C.H. Temporal-spatial pathological changes in the brains of permissive and non-permissive hosts experimentally infected with *Angiostrongylus cantonensis*. *Exp. Parasitol.* **2015**, *157*, 177-84, doi:10.1016/j.exppara.2015.08.006.

24. Bolger, A.M.; Lohse, M.; Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **2014**, *30*, 2114-2120, doi:10.1093/bioinformatics/btu170.
25. Langmead, B. and Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods.* **2012**, *9*, 357-359, doi:10.1038/nmeth.1923.
26. Li, B. and Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **2011**, *12*, 323, doi:10.1186/1471-2105-12-323.
27. Leng, N.; Dawson, J.A.; Thomson, J.A.; Ruotti, V.; Rissman, A.I.; Smits, B.M.; Haag, J.D.; Gould, M.N.; Stewart, R.M.; Kendzior-ski, C. EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq experiments. *Bioinformatics* **2013**, *29*, 1035-1043, doi:10.1093/bioinformatics/btt087.
28. Ashburner, M.; Ball, C.A.; Blake, J.A.; Botstein, D.; Butler, H.; Cherry, J.M.; Davis, A.P.; Dolinski, K.; Dwight, S.S.; Eppig, J.T.; Harris, M.A.; Hill, D.P.; Issel-Tarver, L.; Kasarskis, A.; Lewis, S.; Matese, J.C.; Richardson, J.E.; Ringwald, M.; Rubin, G.M.; Sherlock, G. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **2000**, *25*, 25-29, doi:10.1038/75556.
29. Kanehisa, M. and Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **2000**, *28*, 27-30, doi:10.1093/nar/28.1.27.
30. Yu, G.; Wang, L.G.; Han, Y.; He, Q.Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **2012**, *16*, 284-287, doi:10.1089/omi.2011.0118.
31. Guasconi, L.; Serradell, M.C.; Masih, D.T.; Chiapello, L.S. Immunomodulatory Effect of *Fasciola hepatica* Excretory-Secretory Products on Macrophages. *Methods. Mol. Biol.* **2020**, *2137*, 117-131, doi:10.1007/978-1-0716-0475-5_9.
32. Shepherd, C.; Navarro, S.; Wangchuk, P.; Wilson, D.; Daly, N.L.; Loukas, A. Identifying the immunomodulatory components of helminthes. *Parasite. Immunol.* **2015**, *37*, 293-303, doi:10.1111/pim.12192.
33. Crowe, J.; Lumb, F.E.; Harnett, M.M.; Harnett, W. Parasite excretory-secretory products and their effects on metabolic syn-drome. *Parasite. Immunol.* **2017**, *39*, 10.1111/pim.12410, doi:10.1111/pim.12410.
34. Mehrdana, F. and Buchmann, K. Excretory/secretory products of anisakid nematodes: biological and pathological roles. *Acta. Vet. Scand.* **2017**, *59*, 42, doi:10.1186/s13028-017-0310-3.
35. Wang, Z.; Hao, C.; Zhuang, Q.; Zhan, B.; Sun, X.; Huang, J.; Cheng, Y.; Zhu, X. Excretory/Secretory Products From *Trichinella spiralis* Adult Worms Attenuated DSS-Induced Colitis in Mice by Driving PD-1-Mediated M2 Macrophage Polarization. *Front. Immunol.* **2020**, *11*, 563784, doi:10.3389/fimmu.2020.563784.
36. Chen, K.Y.; Chiu, C.H.; Wang, L.C. Anti-apoptotic effects of Sonic hedgehog signalling through oxidative stress reduction in astrocytes co-cultured with excretory-secretory products of larval *Angiostrongylus cantonensis*. *Sci. Rep.* **2017**, *7*, 41574, doi:10.1038/srep41574.
37. Chen, K.Y.; Cheng, C.J.; Cheng, C.C.; Jhan, K.Y.; Chen, Y.J.; Wang, L.C. The excretory/secretory products of fifth-stage larval *Angiostrongylus cantonensis* induces autophagy via the Sonic hedgehog pathway in mouse brain astrocytes. *PLoS Negl. Trop. Dis.* **2020**, *14*, e0008290, doi:10.1371/journal.pntd.0008290.
38. Chen, K.Y.; Chen, Y.J.; Cheng, C.J.; Jhan, K.Y.; Wang, L.C. Excretory/secretory products of *Angiostrongylus cantonensis* fifth-stage larvae induce endoplasmic reticulum stress via the Sonic hedgehog pathway in mouse astrocytes. *Parasit. Vectors* **2020**, *13*, 317, doi:10.1186/s13071-020-04189-w.
39. Chen, K.Y.; Lu, P.J.; Cheng, C.J.; Jhan, K.Y.; Yeh, S.C.; Wang, L.C. Proteomic analysis of excretory-secretory products from young adults of *Angiostrongylus cantonensis*. *Mem. Inst. Oswaldo. Cruz.* **2019**, *114*, e180556, doi:10.1590/0074-02760180556.
40. Choi, H.H.; Shin, D.M.; Kang, G.; Kim, K.H.; Park, J.B.; Hur, G.M.; Lee, H.M.; Lim, Y.J.; Park, J.K.; Jo, E.K.; Song, C.H. Endoplas-mic reticulum stress response is involved in Mycobacterium tuberculosis protein ESAT-6-mediated apoptosis. *FEBS Lett.* **2010**, *584*, 2445-2454, doi:10.1016/j.febslet.2010.04.050.
41. Samali, A.; Fitzgerald, U.; Deegan, S.; Gupta, S. Methods for monitoring endoplasmic reticulum stress and the unfolded protein response. *Int. J. Cell. Biol.* **2010**, *2010*, 830307, doi:10.1155/2010/830307.
42. Calderón, J.C.; Bolaños, P.; Caputo, C. The excitation-contraction coupling mechanism in skeletal muscle. *Biophys. Rev.* **2014**, *6*, 133-160, doi:10.1007/s12551-013-0135-x.
43. Islam, M.S. Calcium Signaling: From Basic to Bedside. *Adv. Exp. Med. Biol.* **2020**, *1131*, 1-6, doi:10.1007/978-3-030-12457-1_1.
44. Chen, X. and Cubillos-Ruiz, J.R. Endoplasmic reticulum stress signals in the tumour and its microenvironment. *Nat. Rev. Cancer* **2021**, *21*, 71-88, doi:10.1038/s41568-020-00312-2.
45. Fribley, A.; Zhang, K.; Kaufman, R.J. Regulation of apoptosis by the unfolded protein response. *Methods Mol. Biol.* **2009**, *559*, 191-204, doi:10.1007/978-1-60327-017-5_14.
46. Oakes, S.A. Endoplasmic Reticulum Stress Signaling in Cancer Cells. *Am. J. Pathol.* **2020**, *90*, 934-946, doi:10.1016/j.aj-path.2020.01.010.
47. Castilla, J.; Hetz, C.; Soto, C. Molecular mechanisms of neurotoxicity of pathological prion protein. *Curr. Mol. Med.* **2004**, *4*, 397-403, doi:10.2174/1566524043360654.
48. Siwecka, N.; Rozpędek, W.; Pytel, D.; Wawrzynkiewicz, A.; Dżiki, A.; Dżiki, Ł.; Diehl, J.A.; Majsterek, I. Dual role of Endoplas-mic Reticulum Stress-Mediated Unfolded Protein Response Signaling Pathway in Carcinogenesis. *Int. J. Mol. Sci.* **2019**, *20*, 4354, doi:10.3390/ijms20184354.
49. Ghemrawi, R. and Khair, M. Endoplasmic Reticulum Stress and Unfolded Protein Response in Neurodegenerative Diseases. *Int. J. Mol. Sci.* **2020**, *21*, 6127, doi:10.3390/ijms21176127.

50. Becker, K.; Tilley, L.; Vennerstrom, J.L.; Roberts, D.; Rogerson, S.; Ginsburg, H. Oxidative stress in malaria parasite-infected erythrocytes: host-parasite interactions. *Int. J. Parasitol.* **2004**, *34*, 163-189, doi:10.1016/j.ijpara.2003.09.011.
51. Engin, A.B.; Dogruman-Al, F.; Ercin, U.; Celebi, B.; Babur, C.; Bukan, N. Oxidative stress and tryptophan degradation pattern of acute *Toxoplasma gondii* infection in mice. *Parasitol. Res.* **2012**, *111*, 1725-1730, doi:10.1007/s00436-012-3015-6.
52. Götz, A.; Ty, M.C.; Rodriguez, A. Oxidative Stress Enhances Dendritic Cell Responses to *Plasmodium falciparum*. *Immunohorizons* **2019**, *3*, 511-518, doi:10.4049/immunohorizons.1900076.
53. Machado-Silva, A.; Cerqueira, P.G.; Grazielle-Silva, V.; Gadelha, F.R.; Peloso, E.; Teixeira, S.M.; Machado, C.R. How *Trypanosoma cruzi* deals with oxidative stress: Antioxidant defence and DNA repair pathways. *Mutat. Res. Rev. Mutat. Res.* **2016**, *767*, 8-22, doi:10.1016/j.mrrev.2015.12.003.
54. Paiva, C.N.; Feijó, D.F.; Dutra, F.F.; Carneiro, V.C.; Freitas, G.B.; Alves, L.S.; Mesquita, J.; Fortes, G.B.; Figueiredo, R.T.; Souza, H.S.; Fantappiè, M.R.; Lannes-Vieira, J.; Bozza, M.T. Oxidative stress fuels *Trypanosoma cruzi* infection in mice. *J. Clin. Invest.* **2012**, *122*, 2531-2542, doi:10.1172/JCI58525.
55. Raj, D.; Ghosh, E.; Mukherjee, A.K.; Nozaki, T.; Ganguly, S. Differential gene expression in *Giardia lamblia* under oxidative stress: significance in eukaryotic evolution. *Gene* **2014**, *535*, 131-139, doi:10.1016/j.gene.2013.11.048.
56. Schwartz, C.I.; Gabriel, M.E.; Henker, L.C.; Bottari, N.B.; Carmo, G.d.; Guarda, N.; Moresco, R.N.; Machado, G.; Morsch, V.M.; Schetinger, M.R.; Stedille, F.A.; Baska, P.; Mattei, V.; da Silva, A.S.; Mendes, R.E. Oxidative stress associated with pathological changes in the pancreas of cattle naturally infected by *Eurytrema coelomaticum*. *Vet. Parasitol.* **2016**, *223*, 102-110, doi:10.1016/j.vetpar.2016.04.034.
57. Stumbo, A.D.; Goater, C.P.; Hontela, A. Parasite-induced oxidative stress in liver tissue of fathead minnows exposed to trematode cercariae. *Parasitology* **2012**, *139*, 1666-1671, doi:10.1017/S0031182012001023.
58. Betteridge, D.J. What is oxidative stress? *Metabolism* **2000**, *49*, 3-8, doi:10.1016/s0026-0495(00)80077-3.
59. Bhatia, S.; Sardana, S.; Senwar, K.R.; Dhillon, A.; Sharma, A.; Naved, T. In vitro antioxidant and antinociceptive properties of *Porphyra vietnamensis*. *Biomedicine (Taipei)* **2019**, *9*, 3, doi:10.1051/bmdcn/2019090103.
60. Chung, L.Y.; Chen, C.H.; Wang, L.C.; Chang, S.J.; Yen, C.M. Oxidative stress in mice infected with *Angiostrongylus cantonensis* coincides with enhanced glutathione-dependent enzymes activity. *Exp. Parasitol.* **2010**, *126*, 178-183, doi:10.1016/j.exppara.2010.04.017.
61. Chung, L.Y.; Wang, L.C.; Chen, C.H.; Lin, H.Y.; Yen, C.M. Kinetic change of oxidative stress in cerebrospinal fluid of mice infected with *Angiostrongylus cantonensis*. *Redox Rep.* **2010**, *15*, 43-48, doi:10.1179/174329210X12650506623203.
62. Diao, Z.; Wang, J.; Qi, H.; Li, X.; Zheng, X.; Yin, C. Treatment of angiostrongyliasis using a combination of albendazole and dexamethasone: the results of a retrospective and comparative study. *Ann. Trop. Med. Parasitol.* **2011**, *105*, 65-69, doi:10.1179/136485910X12851868780342.