

## Article

# Asian Hornet, *Vespa velutina* Lepeletier 1836 (Hym.: Vespidae), Venom Obtention Based on an Electric Stimulation Protocol

Xesús Feás<sup>1,2\*</sup>, Carmen Vidal<sup>3</sup>, M. Pilar Vázquez-Tato<sup>4</sup> and Julio A. Seijas<sup>4,\*</sup>

1 Academy of Veterinary Sciences of Galicia, Edificio EGAP, Rúa Madrid, No. 2-4, 15707 Santiago de Compostela, (A Coruña), Spain; [xesusfeas@gmail.com](mailto:xesusfeas@gmail.com)

2 Fundación del IDIS, Complejo Hospitalario Universitario de Santiago, Santiago de Compostela, (A Coruña), Spain.

3 Allergy Department, Complejo Hospitalario Universitario de Santiago, Faculty of Medicine, University of Santiago de Compostela, Spain; [carmen.vidal@sergas.es](mailto:carmen.vidal@sergas.es)

4 Departamento de Química Orgánica, Facultad de Ciencias, Universidad de Santiago de Compostela, Alfonso X el Sabio, 27002 Lugo, Spain; [pilar.vazquez.tato@usc.es](mailto:pilar.vazquez.tato@usc.es); [julioa.seijas@usc.es](mailto:julioa.seijas@usc.es)

\* Correspondence: [xesusfeas@gmail.com](mailto:xesusfeas@gmail.com) (X.F.); [julioa.seijas@usc.es](mailto:julioa.seijas@usc.es) (J.A.S)

**Abstract:** The yellow-legged Asian hornet (*Vespa velutina* Lepeletier 1836 (Hymenoptera: Vespidae)) is naturally distributed in China, Southeast Asia and India; however it has since detected outside of its native area, confirmed as being established in South Korea, Europe and Japan. Health risks and deaths caused by the invasive *Vespa velutina* stings have become a public health concern, being the most common cause of anaphylaxis due to hymenopterans in some European regions. This in turn has led to increased demand from medical practitioners and researchers for *Vespa velutina* venom for diagnostic and therapeutic purposes. In this study, a straightforward, quick and inexpensive method for obtaining *Vespa velutina* venom by electric stimulation is described. The venom extracts were analyzed by nuclear magnetic resonance spectroscopy (<sup>1</sup>H-NMR), confirming the composition of the obtained material. The availability of *Vespa velutina* venom will lead to improved diagnostic and therapeutic methods, mainly by venom immunotherapy (VIT), in patients allergic to this invasive species.

**Keywords:** Asian hornet; *Vespa velutina*; Venom; Electrical; Stimulation; Allergy; Stings; Invasive species

## 1. Introduction

Invasive alien species (IAS) are plants, animals, pathogens and other organisms that are introduced and/or spread outside of their natural geographic range and which may cause severe ecological, economic and social impacts on the invaded environments. The European Union experiences annual damages worth €12 billion as a result of IAS effects on human health, damaged infrastructure, and agricultural losses [1]. Recently it was estimated that IAS have cost North America \$2 billion per year in the early 1960s to over \$26 billion per year since 2010 [2] and that the economic cost of IAS has been \$1.288 trillion over the past 50 years [3]. Over 100 examples of IAS that affect human health, sometimes with devastating effects on our livelihood, have been described and documented around the world [4].

Several IAS insect species have migrated in the last decade to Galicia, located on the north western end of the Iberian Peninsula, and successfully colonized and spread, resulting in a broad range of consequences to recipient ecosystems and, thereby, human society [5]. Of these IAS, the yellow-legged Asian hornet (*Vespa velutina* Lepeletier 1836 (Hymenoptera: Vespidae)) was detected in Galicia in 2012.

*Vespa velutina* is naturally distributed in Southeast Asia, India, and China. It was first detected outside of its native habitat in South Korea in 2003 [6], in southwestern France in 2004 [7], on islands of Japan in 2012 [8] and on the Japanese western mainland in 2015 [9].

It was soon recognized as a pan-European threat after being detected in the province of Navarra, northern Spain (2010), in the north-western province of Minho in Portugal (2011), Belgium (2011), Italy (2012), Germany (2014), the Netherlands (2018), Majorca in the Balearic Islands (2015), England and the Channel Islands (2016) [5].

The species has become a major concern to apiculture and industries relying on pollination, given that the diet of these hornet colonies is predominantly based on honey bees and other insects. In Galicia, the annual cost of lost production is estimated at more than 4.5 million euros. The *Vespa velutina* could be responsible for the loss of 65% of the bee colonies in infested areas [10].

*Vespa velutina* is not only a problem for beekeepers and their industrious flying insects, since other agricultural sectors such as fruit producers and viticulturists are also impacted. Despite this, the medical-veterinary potential of *Vespa velutina* should also be outlined. In the Aculeates, the defining feature is that the egg-laying ovipositor is modified to form a sting. Hymenopteran insects are not predisposed to assault and sting humans, however social hornets, wasps and bees have a large defensive response to any risk to the colony. The evolution of the venom system shows to have further developed to cause pain and increase the immune response in humans and different vertebrate predators [11]. Due to its habits, abundance and wider distribution, the risk that the IAS *Vespa velutina* represents for human health is incomparable with other native species of hymenoptera [12].

Currently in invaded areas such as Spain and South Korea there is an increase in the number and severity of reactions in patients exposed to insect venom, mainly due to the invasive species *Vespa velutina*. The introduction of this new species has affected the number of stings of Hymenopterans seen in clinical toxicology units. *Vespa velutina* has become the commonest cause of Hymenoptera anaphylaxis and most cases report no previous *Vespa velutina* stings [13]. In the University Hospital in Santiago de Compostela, Spain, covering an area of 500,000 people, a total of 292 patients were receiving venom allergen immunotherapy in 2020 [14]. In South Korea, from 2010 to 2014, there were 483,233 calls requesting the removal of wasp nests and the stings of Hymenopterans caused 78,860 injuries and 49 deaths, with *Polistes rothneyi koreanus* Vecht and *Vespa velutina* being the most prevalent sources. The total medical costs associated with the stings of hornets and wasps over a five-year period were approximately 3.2 million dollars [15].

The medical community is requesting *Vespa velutina* venom extracts to aid diagnosis and treat allergy and/or anaphylaxis through immunotherapy [16,17]. Venom immunotherapy is the standard of care for people with severe reactions and has been shown to reduce the risk of future anaphylactic events and risk of death [18]. There is a moral imperative to ensure the availability of locally relevant venom extracts for the diagnosis and treatment of hymenopteran venom allergies by immunotherapy [19]. The key motivation behind the present research was to obtain extracts of venom from the invasive species *Vespa velutina*. It therefore has a very important practical application at the clinical level and could ultimately improve the lives of those highly reactive to the sting of the invasive species *Vespa velutina* and cannot “yet” receive immunotherapy due the lack of readily available and reliable sources of venom extracts.

## 2. Results and Discussion

Detailed information with all aspects required for finding, collecting and properly handling *Vespa velutina* specimens, as well as apparatus and methods used for venom extraction, are described in the present work.

*Vespa velutina* is almost ubiquitous in the invaded areas. The number of calls received by the emergency services “112 Galicia” related to “incidents” with the *Vespa velutina* totalled 42,901 in a period of 3 years (2015-2017) [5]. Two colonies of the *Vespa velutina* in

Galicia were detected in 2012, 17 in 2013, 769 in 2014, 5,022 in 2015 and 10,642 in 2016; in 2019, around 25,000 colonies were destroyed, settling widely in urban spaces [12].

At first glance, the sampling of the *Vespa velutina* specimens seems trivial. However, sampling insects requires knowledge of their biology, preferred habitats and activity patterns. Apiaries are hot spots for the collection of *Vespa velutina* as they are concentrated, abundant, and easy to capture [20]. *Vespa velutina* are notorious honey bee hawkers. They fly continuously and hover around the beehive entrance at a distance of 10-40 cm (**Figure 1**), and hunt in flight by intercepting arriving or departing honey bees to the beehive, grabbing the foragers with their outstretched legs.




**Figure 1.** A female *Vespa velutina* specimen in static flight in front of the beehive's entry at the apiary.

The target insect is not easily confused with any other hymenopteran species of hornet, bees and wasps normally present at the apiaries in different areas of Europe, such as the European hornet, *Vespa crabro* Linnaeus, 1758; *Vespa orientalis* Linnaeus, 1771; *Bombus* spp; *Vespula* spp; and/or *Polistes* spp. The *Vespa velutina* can be clearly differentiated because of its unique dark colour pattern, which is mostly black. Moreover, the *Vespa velutina* has the 4<sup>th</sup> abdominal segment almost entirely orange-yellow, is smaller than the native European hornet, *Vespa crabro* Linnaeus, 1758 and possesses yellow tipped legs (**Figure 1**).

The availability of arthropod venom still remains as the main barrier in arthropod toxinology. The difficulties in obtaining sufficient amounts of arthropod venom are either due to scarcity of the given venomous animal and/or the difficulties in the collection of its venom [21]. The distinctive *Vespa velutina* honey bee-capture behavior allows the pinpointing of apiaries as an ideal place, to catch enough live specimens for posterior venom extraction. The *Vespa velutina* colonies are typically at their maximal size in late summer and/or early autumn, with thousands of individuals in their nests, increasing their presence at the apiaries. *Vespa velutina* specimens were collected in an apiary in Viveiro, around 20 km away from the one of the first two original entry points of *Vespa velutina* detected in Galicia in October 2012 (GPS,UTM; X: 632451, Y: 4834800) [5].

Based on our field experiences and long term observations and research of the *Vespa velutina* we carried out an effective sampling and easy transferral of the *Vespa velutina* from the apiary with a net to the venom extraction box (**Figure 2**). Since *Vespa velutina* specimens are abundant in static flight in front of the beehives at the apiary, a fast horizontal swing of the net allows for the effective capture of this insect. Efficient use of a net to capture the *Vespa velutina* is gained only with experience. With a little practice, this becomes quite simple to perform. Because the hornets display positive phototropism,

they facilitated their transfer from the net to the venom extraction box by moving through the black tube towards the light.




**Figure 2.** A captured female *Vespa velutina* specimen by aril netting: (a) Walking up into the net bag; (b) Trapped in a fold of the tip net bag; (c) Inside the designed chamber for venom extraction.

The *Vespa velutina* venom extraction chamber is lightweight (1.3 kg) and mobile so it can be used out in the field. The electrical venom collection device is an all-in-one solid, smart compact, working with automatic tuning where a microprocessor monitors and adjusts pulses, based on humidity, number of insects, how long the device has been running, the battery level and the venom collector's overall condition. The access to on/off button allows users to operate the device safely. Every 50 seconds, the device pauses for 10 seconds. After 40 minutes of work time, the device turns off automatically. That was the maximum time for *Vespa velutina* venom collection in 1 session, for a total of 10 individuals.

A close visual inspection of the electrical venom collection device allows us to observe liquid globules on the over the glass plate (**Figure 3**), where globules crystallize or dehydrate quickly with exposure to the air. Once transported to the lab, in less than two hours, no liquid is observed on the plate, although an inspection under ultraviolet light does reveal small spots on the glass (**Figure 4a**). The dried venom was removed from the glass plates, carefully with a razor (**Figure 4b**).


At a lab-scale level, there are well tested and widely accepted techniques for obtaining venom from several hymenopterans, such as: (i) *Polybia paulista* Ihering, 1896 [22]; *Apis mellifera carnica* Linnaeus, 1758 [23] and *Vespa affinis* Linnaeus, 1764 [24]; and (ii) *Vespula maculifrons* Buysson, 1905; *Vespula germanica* Fabricius, 1793; *Vespula vulgaris* Linnaeus, 1758 [25]; *Polistes annularis* Linnaeus, 1763; *Polistes carolina* Linnaeus, 1767; *Polistes exclamans* Viereck, 1906; *Polistes fuscatus* Fabricius, 1793; *Polistes instabilis* Saussure, 1853; and *Vespula germanica* Fabricius, 1793 [26]; *Dolichovespula maculata* Linnaeus, 1763; *Polistes annularis* Linnaeus, 1763; and *Vespula vulgaris* Linnaeus, 1758 [27].



**Figure 3.** A female *Vespa velutina* specimens in the milking grid: (a) Resting, venom collection device off; (b) Bending the tip of their abdomen downward and stinging after receive an electrical shock, venom collection device on; (c) Deposited liquid globules of venom over the glass plate (red arrows).

They are based on dead, frozen insects, where the whole sting apparatus needs to be dissected, using microsurgical forceps, and the obtained venom sacs are basically: (i) manually extracted from the separated venom sacs, i.e. by gentle squeezing, or (ii) homogenized and/or just pooled to collect the liquid fraction by centrifugation.

The above existing venom extraction protocols are followed to obtain actual *Vespa velutina* venom [28-31]. The venom reservoir of the *Vespa velutina* is about 1 mm in length, white and transparent, [28,29] and the microdissection of the venom reservoir, usually by the precise manipulation of specialized needles and requires a high degree of operator skills. This makes the manual extraction of venom a tedious, laborious and time consuming task.



**Figure 4.** At laboratory, the removed glass plate from the venom collector device after *Vespa velutina* electric stimulation (a) under UV light, showing venom dried spots (red arrows); (b) scraped of the venom.

Li et al. demonstrate that honey bee venom extracted manually is different from venom extracted using electrical stimulation, and these differences may be important in their use as pharmacological agents [32]. The venom extracted manually is contaminated by non-toxin proteins that may leak from the gland tissue that is cut/disrupted during venom collection. Data showed that the toxin component in the venom of manual collection has significantly lower content than venom from electrical stimulation. The three newly identified phosphorylated venom proteins in the venom extracted through the use of electrical stimulation may elicit a different immune response through the specific recognition of antigenic determinants [32]. Comparison of venom from three vespid species (*Vespula maculifrons*, *Vespula maculata* and *Vespula arenaria*) collected by electrostimulation and by venom sac extraction were shown to be potent allergens, the last one having the

potential advantage of being free of contaminating tissue protein [33]. There is a scope for future research, using modern technology, to assess the potential differences between a vespid venom, in this case, venom from *Vespa velutina*, obtained by both methods: manual extraction and by electrical stimulation. Venom from gland extracts may possess components derived from the gland's epithelium, muscular layer, nerves, etc. Furthermore, it was suggested that foreign enzymes included in the gland extracts may affect the active components of the venom [21].

A variety of stimulation techniques have been evolved and are available for the collection of venom from individual Hymenoptera as well as simultaneously from large numbers of insects [34–38].

Since the first venom extraction carried out in 2017, we continued to develop the technique, adapting the venom extraction box, on account of a specific behaviour of a whole colony in their nest and were able to obtain venom extracts by electro stimulation of several hundreds of individual, obtained directly from a captive nest (**Figure 5**). The above allows the collection of more venom over a short time (**Supplementary video 1**). However to use a live *Vespa velutina* nest for milking the venom of their inhabitants is a high risk activity, needing skill which requires specialist training and, specific working places, more resources and infrastructure.




**Figure 5.** Scraped of the *Vespa velutina* venom obtained from a colony of 960 individuals from a captive nest last November 2020.

Targeted netting to capture the insect can provide enough samples of *Vespa velutina* specimens for venom collection by electrical stimulation. However, in the specific case of social wasps, there is a long tradition of harvesting wild nests to eat larvae and pupae, as well as the use of nests in medicine recipes. Moreover, collectors have also developed practices that can be understood to some extent as incipient vespiculture [39]. Although rigorous testing is required as current *Vespa* hornet rearing efforts are undeveloped, research indicates the prospect of a functional year-round *Vespa* hornet rearing process being developed. The current biological, ecological, medicinal, and culinary motivations justify the development of true hornet vespiculture [40]. We expect that the method developed here for venom extraction of the *Vespa velutina*, will prove satisfactory for collecting venom from other species of venomous arthropods and/or stinging insects, as well as reared colonies of bumblebees, wasps and hornets. Breeding of such target insects would reduce the necessity of wild harvesting.


A prerequisite to studying the nature of the venoms is the development of methods for their collection. The collection of venom should no longer be a limiting factor as this work details a successful method to allow for the easy collection of venom. Integrating transcriptomic and proteomic analyses should provide a better understanding of: the (i) venom composition of venomous hymenoptera in particular; and (ii) mammalian

immune system responses to those stinging insects. It is to be hoped that this will assist any future applications of venoms into diverse biomedicine, and the possible discovery and development of new pharmacological agents, and other related research areas [31, 41-45].


The collected venom was dissolved in methanol-d<sub>4</sub> (CD<sub>3</sub>OD) and analyzed by <sup>1</sup>H-NMR (500 MHz). In the proton spectrum (**Figure 6**) three regions can be observed in the enlargements (**Figures 7, 8 and 9**). Expanded region A ( $\delta = 6.5$ -8.7 ppm) shows the signals of aromatic hydrogens present in tryptophan, phenylalanine, tyrosine and histidine.




**Figure 6.** <sup>1</sup>H-NMR spectra (500 MHz, CD<sub>3</sub>OD) of *Vespa velutina* venom obtained by electrical stimulation.



**Figure 7.** Expanded region ( $\delta = 6.5$ -8.7 ppm) of a <sup>1</sup>H-NMR spectra (500 MHz, CD<sub>3</sub>OD) of *Vespa velutina* venom obtained by electrical stimulation.



**Figure 8.** Expanded region ( $\delta = 2.45\text{-}3.95$  ppm) of a  $^1\text{H-NMR}$  spectra (500 MHz,  $\text{CD}_3\text{OD}$ ) of *Vespa velutina* venom obtained by electrical stimulation.



**Figure 9.** Expanded region ( $\delta = 0.4\text{-}2.3$  ppm) of a  $^1\text{H-NMR}$  spectra (500 MHz,  $\text{CD}_3\text{OD}$ ) of *Vespa velutina* venom obtained by electrical stimulation.

### 3. Materials and Methods

The field study was conducted during September 2017 in one apiary ( $43^\circ 37' 15.2''\text{N}$   $7^\circ 35' 26.7''\text{W}$ ) located in San Pedro de Viveiro (municipality of Viveiro), in the Western Mariña at the province of Lugo (Galicia, Spain). At an altitude of 177 m above sea level, the climate in the area is characterized by mildness and rainfall, as corresponds to the oceanic climate. The average annual temperature exceeds  $14^\circ\text{C}$ , while the thermal oscillation is weak ( $10^\circ\text{C}$ ), as a result of a mild winter and moderate temperatures in summer.

Adult female *Vespa velutina* specimens were obtained while using an active collecting net method at an apiary. The apiary had: a total of (i) six beehives with a frontal protective module, a grid which prevents the entry of the *Vespa velutina* into the beehive;

and (ii) twelve traps situated at the beehives consisted of a 15 litre plastic box (width = 36.5 cm; depth = 28.5 cm; height = 18.5 cm) with four holes on the sides of each box. The bait used consisted of blueberry juice, brown beer and wax obtained from honey bee combs.

### 3.1. Materials


#### 3.1.1. Protective equipment

A full body protective apiary suit, consisting on one-piece pants and jacket, hat, veil and a pair of long sleeve beekeeping gloves was worn by the operators at the apiary when collecting of *Vespa velutina* specimens. For venom removal, at the lab, it is necessary to use protective equipment: glasses, gloves and mask to avoid contamination and potential direct contact with the skin and mucosas.

#### 3.1.2. The *Vespa velutina* venom extraction chamber

System set-up for *Vespa velutina* venom obtention by electric stimulation consists of a modified container where a venom collection device is located (**Figure 10**), composed of:

- A transparent plastic box (285 x 160 x 120 mm) with a lid and hermetically sealed, in which modifications were made for introduction of the captured insects (1) and to access the on/off switch of the electrical venom collection device with a stick (2). Two holes consisted of a circular, 30 mm diameter at a height of 70 mm from the base, for the introduction tube (1); and a rectangular aperture (10 mm x 20 mm) to give access to the on/off switch,(2). A 100 mm length plastic tubing, covered with grey adhesive tape, is inserted into hole number 1, of the plastic box with 45 mm protruding from the box and a screw cap at the end to prevent escape. Hole number two for the on/off switch mechanism is located at the back of the venom collection device, approx. 75-80 mm from the base of the box.
- An electrical venom collection device was supplied by I GK Electronics (Varna, Bulgaria), designed to harvest honey bee venom. The device consists of a solid wooden frame (250 x 158 x 38 mm) having an area 250 by 158 mm over which 39 wires are stretched at 3 mm intervals (**Figure 10**). A removable glass plate (201 x 140 x 4 mm), fits under the wires.



**Figure 10.** System set-up for *Vespa velutina* venom obtention chamber by electric stimulation.

### 3.2. Collecting insects and venom

#### 3.2.1. Identification of *Vespa velutina* individuals

Insects were identified using their external morphological characteristics. In brief, *Vespa velutina* average about 2 to 3 cm in length, head is black with face and mouthparts orange, the antennae are brown dorsally and orange ventrally. The thorax is dark brown, almost black. Metasomal terga brown, with thin yellow band on segment 1st and a thin orange band on 2nd and 3rd segments; metasomal segment 4th orange; metasomal segments 5th and 6th orange-brown. Legs are brown, with yellow tarsi and the wings are brownish hyaline.

### 3.2.2. Catching the insects

Collection of *Vespa velutina* specimens hovering in front of the beehives or the traps was performed by horizontally swinging the net quickly across the hornet, to capture the specimen and then follow through swipe to force the insect into the very bottom of the net bag tip. The above is performed with a fast twist of the wrist so that the bottom of the net bag hangs over the rim. If necessary, with the rim of the net in contact with the ground, hold the tip of the bag up with one hand. The *Vespa velutina* will fly or walk upward into the net at the tip of the bag, which can then be flipped over with the hand to entrap the specimen, keeping a sufficient amount of netting between the hand and insect.

### 3.2.3. Transfer the insects to the venom collection chamber

The trapped insect in a fold of the tip of the net, is then inserted into the *Vespa velutina* venom extraction chamber. The part of the open net is placed embedded at the entrance of tube number 1, until we reach the pocket that we are holding with one hand. At this time, the inlet of the tube number 1 is surrounded with the net, so the insect has free access to the tube entry. Captured insects will crawl voluntarily toward the illuminated side, in the venom collection box. Once the insect passes through the tube and enters the venom extraction box, which occurs in 3-5 seconds, the net is removed and the inlet tube is closed with the lid.

### 3.2.4. Electric stimulation

The glass panels should be sterilized previously with 70-90% ethyl alcohol. Turn on the venom collector device with the switch On/Off key with a stick through hole number 2. The device starts to work and you will see the green LED, flashing slowly: three times – pause - three times. Every minute the electrical device will pause for about 10 seconds. After about 40 minutes, the electrical venom collector turns itself off. Once the venom-harvesting is complete, the insects are removed. Then, the *Vespa velutina* venom extraction chamber with the fresh *Vespa velutina* venom on the electrical venom collection device is carefully packed into a container for transportation to the laboratory

### 3.2.5. *Vespa velutina* venom removal and processing in the laboratory

At the lab, the glass plate is pulled out from the electrical venom collection device, and inspected visually under UV light. The glass plate is then scraped, with the help of a razor and spatula, and transferred to a vial. The dehydrated venom, is then kept in dark bottles, and stored in a refrigerator at -15°C pending further analysis.

## 3.3. <sup>1</sup>H-NMR analysis

<sup>1</sup>H-NMR analyses were performed on a Varian VNMRS-500-WB spectrometer (500 MHz for <sup>1</sup>H) instrument (Agilent Technologies, Palo Alto, CA, USA), equipped with a 5 mm probe. The sample was dissolved in 500 μL of methanol-d<sub>4</sub> CD<sub>3</sub>OD, (Sigma-Aldrich, Madrid, Spain) shaken in a vortex mixer, and the resulting mixture was placed into a 5-mm diameter ultra-precision NMR sample tubes (Norell, Landisville, PA, USA). The temperature of the sample in the probe was 30 °C. The chemical shifts are reported in ppm, using the solvent proton signal as standard. The area of the signals was determined by using the equipment software, and the integrations were carried out three times to obtain average values. All figures of the <sup>1</sup>H-NMR spectra and of the expanded <sup>1</sup>H-NMR spectrum regions were plotted at a fixed value of absolute intensity to be valid for comparative purposes.

Values for the peaks detected are as follows: <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD) δ 8.45, 8.38, 8.09, 8.05, 8.00, 7.97, 7.93, 7.87, 7.76, 7.75, 7.62, 7.61, 7.55, 7.51, 7.50, 7.29, 7.28, 7.23, 7.14, 7.13, 7.12, 7.12, 7.02, 7.01, 6.99, 6.98, 6.92, 6.90, 6.87, 6.86, 6.86, 6.67, 6.66, 6.65, 6.65, 3.90, 3.83, 3.74, 3.73, 3.72, 3.68, 3.62, 3.62, 3.61, 3.60, 3.40, 3.40, 3.38, 3.16, 3.15, 3.14, 3.13, 3.13, 3.10, 3.09, 3.08, 3.07, 2.99, 2.98, 2.97, 2.80, 2.52, 2.50, 2.49, 2.47, 2.18, 2.16, 1.97, 1.97, 1.93, 1.92, 1.90, 1.88, 1.86, 1.72, 1.70, 1.67, 1.62, 1.52, 1.48, 1.47, 1.45, 1.43, 1.42, 1.36, 1.34, 1.33, 1.32, 1.30, 1.02, 0.89, 0.89, 0.88, 0.85, 0.81, 0.80, 0.80, 0.79, 0.77, 0.77, 0.75, 0.73, 0.71, 0.62, 0.44, 0.43.

#### 4. Conclusions

*Vespa velutina* is an invasive alien species with medical importance on the health of members of the public. We described a straightforward, quick and inexpensive method for obtaining *Vespa velutina* venom, based on an electric stimulation protocol; including all sequential steps for sampling, handling and milking the insect. The materials used are cheap and readily accessible in the market. The *Vespa velutina* venom extraction chamber can be used in field, since it is totally portable, by operators with minimal training. The method and detailed protocol developed here for venom extraction of the *Vespa velutina*, could be adapted satisfactorily for collecting venom from other species of venomous arthropods. Actually, the *Vespa velutina* venom extracts obtained by electro stimulation have allowed the initiation of new investigations in the Clinical Immunology Unit of the University Hospital of Santiago de Compostela, which we expect may ultimately promote substantial improvements in the diagnosis, prevention and medical treatment of severe allergies and anaphylaxis caused by *Vespa velutina* stings.

**Supplementary Materials:** The following are available online at [www.mdpi.com/xxx/s1](http://www.mdpi.com/xxx/s1), Video S1: Scraped of the *Vespa velutina* Lepeletier 1836 venom obtained from a captive nest.

**Author Contributions:** Conceptualization, X.F.; methodology, X.F., J.A.S. and P.V.; formal analysis, C.V.; investigation, C.V.; resources, C.V.; writing—original draft preparation, X.F.; writing—review and editing, X.F., C.V., P.V., and J.A.S; supervision, C.V.; project administration, C.V.; funding acquisition, C.V. All authors have read and agreed to the published version of the manuscript.

**Funding:** The authors thank the Ministerio de Ciencia y Tecnología (Project MAT2017-86109P) for the financial support. X.F. thanks Instituto de Salud Carlos III for a research contract (program number: RD16/0017/001).

**Institutional Review Board Statement:** Asian Hornet, *Vespa velutina* Lepeletier, 1836 (Hym.: Vespidae) is not a regulated invertebrate. Therefore, no ethical use approval is necessary

**Informed Consent Statement:** Not applicable.

**Acknowledgments:** We would like to thank to Robert Hogge (former President of the Jersey Beekeepers' Association and founder of Jersey Asian hornet Group) and Sam Day (former Kent Beekeeper's Asian hornet Team Coordinator) for critically reading the manuscript. Thanks also to Jose María Vázquez for support when sampling at the apiary and to Dani Slizt for obtaining the *Vespa velutina* nests. Thanks also to José Amoedo Montes and Carmen Cabadas Amoedo for providing the necessary facilities for this work, in San Amaro (Fornelos de Montes, Galicia). Assistance and comments provided by Arturo González-Quintela (Department of Medicine, Complejo Hospitalario Universitario de Santiago) was greatly appreciated.

**Conflicts of Interest:** The authors declare no conflict of interest.

**Sample Availability:** Samples of the *Vespa velutina* venom are available from the authors.

#### References

1. Genovesi, P.; Carnevali, L.; Scalera, R. The impact of invasive alien species on native threatened species in Europe. ISPRA - ISSG, Rome. Technical Report for the European Commission. 2005. Pp. 18.
2. Crystal-Ornelas, R.; Hudgins, E.J.; Cuthbert, R.N.; Haubrock, P.J.; Fantle-Lepczyk, J.; Angulo, E.; Kramer, A.M.; Ballesteros-Mejia, L.; Leroy, B.; Leung, B.; López-López, E.; Diagne, C.; Courchamp, F. Economic costs of biological invasions within North America. *NeoBiota* **2021**, *67*, 485–510.
3. Zenni, R.D.; Essl, F.; García-Berthou, E.; McDermott, S.M. The economic costs of biological invasions around the world. *NeoBiota* **2021**, *67*, 1–9.
4. Mazza, G.; Tricarico, E.; Genovesi, P.; Gherardi, F. Biological invaders are threats to human health: an overview. *Ethol. Ecol. Evol.* **2014**, *26*, 112–129.
5. Feás Sánchez, X.; Charles, R.J. Notes on the Nest Architecture and Colony Composition in Winter of the Yellow-Legged Asian Hornet, *Vespa velutina* Lepeletier 1836 (Hym.: Vespidae), in Its Introduced Habitat in Galicia (NW Spain). *Insects* **2019**, *10*, 237.

6. Kim, J.; Choi, M.; Moon, T. Occurrence of *Vespa velutina* Lepeletier from Korea, and a revised key for Korean *Vespa* species (Hymenoptera: Vespidae). *Entomol. Res.* **2006**, *36*, 112–115.
7. Haxaire, J.; Bouguet, J.-P.; Tamisier, J.-P. *Vespa velutina* Lepeletier, 1836, une redoutable nouveauté pour la faune de France (Hymenoptera, Vespidae). *Bull. Soc. Entomol. France* **2006**, *111*, 194.
8. Minoshima, Y.; Yamane, S.K.; Ueno, T. An invasive alien hornet, *Vespa velutina nigrithorax* du Buysson (Hymenoptera, Vespidae), found in Kitakyushu, Kyushu Island: A first record of the species from mainland Japan. *Jpn. J. Syst. Entomol.* **2015**, *21*, 259–261.
9. Takahashi, R.; Sakai, Y.; Yamamura, T.; Kiyoshi, T.; Takahashi, J. Analysis the nest of alien hornet, *Vespa velutina*, found for the first time in Tsushima Island, Japan. *Trans. Nagasaki Biol. Soc.* **2015**, *76*, 49–56.
10. Fedele, E.; Gervasini, E.; De Jesus Cardoso, A.; La Notte, A.; Vallecillo Rodriguez, S.; Tsiamis, K.; Maes, J., Invasive Alien Species impact on Ecosystem Services, EUR 29827 EN, Publications Office of the European Union, Luxembourg, 2019, ISBN 978-92-76-09510-1, doi:10.2760/646477, JRC111718.
11. Bauman, K. Evolution of the Venom System in Aculeate Hymenoptera. Ph.D. Thesis, The University of Queensland, Brisbane, Australia, 2018.
12. Feás, X. Human Fatalities Caused by Hornet, Wasp and Bee Stings in Spain: Epidemiology at State and Sub-State Level from 1999 to 2018. *Biology* **2021**, *10*, 73.
13. Vidal, C.; Armisen, M.; Monsalve, R.; González-Vidal, T.; Lojo, S.; López-Freire, S.; Méndez, P.; Rodríguez, V.; Romero, L.; Galán, A.; González-Quintela, A. Anaphylaxis to *Vespa velutina nigrithorax*: pattern of sensitization for an emerging problem in Western countries. *J. Investig. Allergol. Clin. Immunol.* **2021**, *31*, 228–235.
14. Martínez-Lourido, E.; Otero, A.; Armisen, M.; Vidal, C. Comment on: Bilò MB, Pravettoni V, Mauro M, Bonadonna P. Treating venom allergy during COVID-19 pandemic: Management of venom allergen immunotherapy during the COVID-19 outbreak in Spain. *Allergy* **2021**, *76*, 951–952.
15. Choi, M.B.; Kim, T.G.; Kwon, O. Recent Trends in Wasp Nest Removal and Hymenoptera Stings in South Korea. *J. Med. Entomol.* **2018**, *56*, 254–260.
16. Chugo, S.; Lizaso, M.T.; Alvarez, M.J.; Arroabaren, E.; Lizarza, S.; Tabar, A.I. *Vespa velutina nigrithorax*: A New Causative Agent in Anaphylaxis. *J. Investig. Allergol. Clin. Immunol.* **2015**, *25*, 231–232.
17. Tabar, A.I.; Chugo, S.; Joral, A.; Lizaso, M.T.; Lizarza, S.; Alvarez-Puebla, M.J.; Arroabarren, E.; Vela, C.; Lombardero, M. *Vespa Velutina Nigrithorax*: a new causative agent for anaphylaxis. *Clin. Transl. Allergy* **2015**, *5*, P43.
18. Sahiner, U.M.; Durham, S.R. Hymenoptera Venom Allergy: How Does Venom Immunotherapy Prevent Anaphylaxis From Bee and Wasp Stings?. *Front. Immunol.* **2019**, *10*, 1959.
19. Korošec, P.; Jakob, T.; Harb, H.; H eddle, R.; Karabus, S.; de Lima Zollner, R.; Selb, J.; Thong, B.Y.; Zaitoun, F.; Golden, D.; Levin, M. Worldwide perspectives on venom allergy. *World Allergy Organ. J.* **2019**, *12*, 100067.
20. Feás, X.; Vázquez-Tato, M.P.; Seijas, J.A.; Pratima G. Nikalje, A.; Fraga-López, F. Extraction and Physicochemical Characterization of Chitin Derived from the Asian Hornet, *Vespa velutina* Lepeletier 1836 (Hym.: Vespidae). *Molecules* **2020**, *25*, 384.
21. Breer, H.; Miller, T.A. Neurochemical Techniques in Insect Research. Springer Series in Experimental Entomology. 1985
22. Dias, N.B.; de Souza, B.M.; Gomes, P.C.; Palma, M.S. Peptide diversity in the venom of the social wasp *Polybia paulista* (Hymenoptera): a comparison of the intra- and inter-colony compositions. *Peptides* **2014**, *51*, 122–130.
23. Peiren, N.; Vanrobaeys, F.; de Graaf, D.C.; Devreese, B.; Van Beeumen, J.; Jacobs, F.J. The protein composition of honeybee venom reconsidered by a proteomic approach. *Biochim. Biophys. Acta* **2005**, *1752*, 1–5.
24. Sookrung, N.; Wong-din-Dam, S.; Tungtrongchitr, A.; Reamtong, O.; Indrawattana, N.; Sakolvaree Y.; Visitsunthorn, N.; Manuyakorn, W.; Chaicumpa, W. Proteome and allergenome of Asian wasp, *Vespa affinis*, venom and IgE reactivity of the venom components. *J. Proteome Res.* **2014**, *13*, 1336–1344.
25. King, T.P.; Alagon, A.C.; Kuan, J.; Sobotka, A.K.; Lichtenstein, L.M. Immunochemical studies of yellowjacket venom proteins. *Mol. Immunol.* **1983**, *20*, 297–308.
26. King, T.P.; Kochoumian, L.; Joslyn, A. Wasp venom proteins: phospholipase A1 and B. *Arch. Biochem. Biophys.* **1984**, *230*, 1–12.
27. King, T.P.; Lu, G.; Gonzalez, M.; Qian, N.; Soldatova, L. Yellow jacket venom allergens, hyaluronidase and phospholipase: sequence similarity and antigenic cross-reactivity with their hornet and wasp homologs and possible implications for clinical allergy. *J. Allergy Clin. Immunol.* **1996**, *98*, 588–600.

28. Le, T.N.; Da Silva, D.; Colas, C.; Darrouzet, E.; Baril, P.; Leseurre, L.; Maunit, B. Asian hornet *Vespa velutina nigrithorax* venom: Evaluation and identification of the bioactive compound responsible for human keratinocyte protection against oxidative stress. *Toxicon* **2020**, *176*, 1–9.
29. Le T,N.; da Silva, D.; Colas, C.; Darrouzet, E.; Baril, P.; Leseurre, L.; Maunit, B. Development of an LC-MS multivariate nontargeted methodology for differential analysis of the peptide profile of Asian hornet venom (*Vespa velutina nigrithorax*): application to the investigation of the impact of collection period variation. *Anal. Bioanal. Chem.* **2020**, *412*, 1419–1430.
30. Monsalve, R.I.; Gutiérrez, R.; Hoof, I.; Lombardero, M. Purification and molecular characterization of phospholipase, antigen 5 and hyaluronidases from the venom of the Asian hornet (*Vespa velutina*). *PLoS ONE* **2020**, *15*, e0225672.
31. Vidal, C.; Armisen, M.; Monsalve, R.; Gómez-Rial, J.; González-Fernández, T.; Carballada, F.; Lombardero, M.; González-Quintela, A. Vesp v 5 and glycosylated Vesp v 1 are relevant allergens in *Vespa velutina nigrithorax* anaphylaxis. *Clin. Exp. Allergy* **2020**, *50*, 1424–1427.
32. Li, R.; Zhang, L.; Fang, Y.; Han, B.; Lu, X.; Zhou, T.; Feng, M.; Li, J. Proteome and phosphoproteome analysis of honeybee (*Apis mellifera*) venom collected from electrical stimulation and manual extraction of the venom gland. *BMC Genomics* **2013**, *14*, 766.
33. Mueller, U.; Reisman, R.; Wypych, J.; Elliott, W.; Steger, R.; Walsh, S.; Arbesman, C. Comparison of vespoid venoms collected by electrostimulation and by venom sac extraction. *J. Allergy Clin. Immunol.* **1981**, *68*, 254–261.
34. Piek, T. (1986). 3 – Methods for the Collection of Venoms. Piek T. Venoms of the Hymenoptera: Biochemical, Pharmacological, and Behavioural Aspects. Academic Press; London, UK: 1986. p. 570.
35. Ali, M.A.A.S.M. Studies on Bee Venom and Its Medical Uses. *Int. J. Adv. Res. Technol.* **2012**, *1*, 69–83.
36. Eskridge, E.M.; Elliott W.B.; Elliott A.H.; Eskridge, P.B.; Doerr, J.C.; Schneller, N.; Reisman, R.E. Adaptation of the electrical stimulation procedure for the collection of vespoid venoms. *Toxicon* **1981**, *19*, 893–897.
37. Roger P, Simon.; Allen W, Benton. A Method for Mass Collection of Wasp Venoms. *Ann. Entomol. Soc. Am.* **1969**, *62*, 277–278.
38. de Graaf, D.C.; Brochetto Braga, M.R.; de Abreu, R.M.; Blank, S.; Bridts, C.H.; De Clerck, L.; Devreese, B.; Ebo, D.G.; Ferris, T.J.; Hagendorens, M.M.; Justo Jacomini, D.L.; Kanchev, I.; Kokot, Z.J.; Matysiak, J.; Mertens, C.M.; Sabato, V.; Van Gasse, A.L.; Van Vaerenbergh, M. Standard methods for *Apis mellifera* venom research. *J. Apic. Res.* **2020**, *60*, 1–31.
39. van Itterbeeck, J.; Feng, Y.; Zhao, M.; Wang, C.; Tan, K.; Saga, T.; Nonaka, K.; Jung, C. Rearing techniques for hornets with emphasis on *Vespa velutina* (Hymenoptera: Vespidae): A review. *J. Asia Pac. Entomol.* **2021**, *24*, 103–107.
40. Detoni, M.; Féas, X.; Jeanne, R.L.; Loope, K.J.; O'Donnell, S.; Santoro, D.; Sumner, S.; Jandt, J.M. Evolutionary and ecological pressures shaping social wasps collective defenses. *Ann. Entomol. Soc. Am.* **2020**, *113*, 407–424.
41. Lee, S.H.; Baek, J.H.; Yoon, K.A. Differential Properties of Venom Peptides and Proteins in Solitary vs. Social Hunting Wasps. *Toxins* **2016**, *8*, 32.
42. Abd El-Wahed, A.; Yosri, N.; Sakr, H.H.; Du, M.; Algethami, A.F.M.; Zhao, C.; Abdelazeem, A.H.; Tahir, H.E.; Masry, S.H.D.; Abdel-Daim, M.M.; Musharraf, S.G.; El-Garawani, I.; Kai, G.; Al Naggari, Y.; Khalifa, S.A.M.; El-Seedi, H.R. Wasp Venom Biochemical Components and Their Potential in Biological Applications and Nanotechnological Interventions. *Toxins* **2021**, *13*, :206.
43. VV, Achar R.R.; MUH, N A, T.Y.S.; Kameshwar, V.H.; Byrappa, K.; Ramadas, D. Venom peptides - A comprehensive translational perspective in pain management. *Curr. Res. Toxicol.* **2021**, *9*: 329–340.
44. Trim, C.M.; Byrne, L.J.; Trim, S.A. Utilisation of compounds from venoms in drug discovery. *Prog. Med. Chem.* **2021**, *60*, 1–66.
45. Pak, S.C. An Introduction to the Toxins Special Issue on “Bee and Wasp Venoms: Biological Characteristics and Therapeutic Application”. *Toxins* **2016**, *8*, 315.