

## Psychological distress in men during the COVID-19 pandemic in Brazil: The role of the sociodemographic variables, uncertainty and social support

Anderson Reis de Sousa<sup>1</sup>, Jules Ramon Brito Teixeira<sup>2</sup>, Wanderson Carneiro Moreira<sup>3,4</sup>, Milena Bitencourt Santos<sup>1</sup>, Herica Emilia Félix de Carvalho<sup>5</sup>, Éric Santos Almeida<sup>1</sup>, Raíssa Millena Silva Florencio<sup>6</sup>, Aline Macêdo de Queiroz<sup>7</sup>, Magno Conceição das Mercês<sup>8</sup>, Tilson Nunes Mota<sup>9</sup>, Isabella Félix Meira Araújo<sup>1</sup>, Josielson Costa da Silva<sup>1</sup>, Sélton Diniz dos Santos<sup>2</sup>, Emerson Lucas Silva Camargo<sup>10</sup>, Luciano Garcia Lourenção<sup>11</sup>, Richardson Augusto Rosendo da Silva<sup>12</sup>, Evanilda Souza de Santana Carvalho<sup>2</sup>, Iracema Lua<sup>13</sup>, Sônia Barros<sup>3</sup>, Tânia Maria de Araújo<sup>2</sup>, Márcia Aparecida Ferreira de Oliveira<sup>3,14</sup>, Álvaro Pereira<sup>1</sup>, Wilson Abreu<sup>15</sup> Carlos Alberto da Cruz Sequeira<sup>15\*</sup>

<sup>1</sup>Escola de Enfermagem, Universidade Federal da Bahia. 40231-300, Salvador, Bahia, Brasil.

<sup>2</sup>Universidade Estadual de Feira de Santana. 44001-970, Feira de Santana, Bahia, Brasil.

<sup>3</sup>Escola de Enfermagem, Universidade de São Paulo. 05403-000, São Paulo, São Paulo, Brasil.

<sup>4</sup>Coordenação de Saúde do Adolescente, Jovem e Homem, Diretoria Técnica, Secretaria de Saúde, Prefeitura Municipal de Ananindeua. 67130-600, Ananindeua, Pará, Brasil.

<sup>5</sup>Escola de Enfermagem de Ribeirão Preto, Universidade de São Paulo. 14040-902, Ribeirão Preto, São Paulo, Brasil.

<sup>6</sup>Faculdade Estácio de Alagoinhas. 48010-970, Alagoinhas, Bahia, Brasil.

<sup>7</sup>Faculdade de Enfermagem, Universidade Federal do Pará. 66075-110, Belém, Pará, Brasil.

<sup>8</sup>Universidade do Estado da Bahia. 1150-000, Salvador, Bahia, Brasil.

<sup>9</sup>Secretaria de Ciência, Tecnologia e Inovação. 41745-004, Salvador, Bahia, Brasil.

<sup>10</sup>Universidade de Ribeirão Preto. 11440-003, Ribeirão Preto, São Paulo, Brasil.

<sup>11</sup>Escola de Enfermagem, Universidade Federal do Rio Grande. 96.201-900, Rio Grande, Rio Grande do Sul, Brasil.

<sup>12</sup>Universidade Federal do Rio Grande do Norte. 59012-300, Natal, Rio Grande do Norte, Brasil.

<sup>13</sup>Instituto de Saúde Coletiva, Universidade Federal da Bahia. 40110-040, Salvador, Bahia, Brasil.

<sup>14</sup>Universidade Federal de Santa Maria. 97105-900, Santa Maria, Rio Grande do Sul, Brasil.

<sup>15</sup>Centro de Investigação em Tecnologias e Serviços de Saúde, Escola Superior de Enfermagem do Porto. 4200-072 Porto, Portugal.

\*Correspondence: [carlossequeira@esenf.pt](mailto:carlossequeira@esenf.pt), Escola Superior de Enfermagem do Porto. 4200-072 Porto, Portugal.

## ABSTRACT

**Objective:** to analyze the relationships between sociodemographic variables, intolerance to uncertainty (INT), social support and psychological distress (i.e., indicators of Common Mental Disorders [CMDs] and perceived stress [PS]) in Brazilian men during the COVID-19 pandemic.

**Methods:** a cross-sectional study with national coverage, of the web survey type, and conducted with 1,006 Brazilian men during the period of social circulation restriction imposed by the health authorities in Brazil, for suppression of the coronavirus and control of the pandemic. Structural equation modeling analysis was performed. **Results:** Statistically significant direct effects of race/skin color ( $\lambda=0.268$ ;  $p\text{-value}<0.001$ ), socioeconomic status (SES) ( $\lambda=0.306$ ;  $p\text{-value}<0.001$ ), household composition ( $\lambda=0.281$ ;  $p\text{-value}<0.001$ ), PS ( $\lambda=0.513$ ;  $p\text{-value}<0.001$ ) and INT ( $\lambda=0.421$ ;  $p\text{-value}<0.001$ ) were evidenced in the occurrence of CMDs. Black-skinned men, with higher SES, living alone and with higher PS and INT levels presented higher prevalence values of CMDs. **Conclusions:** high levels of PS and INT were the factors that presented the strongest associations with the occurrence of CMDs among the men. It is necessary to implement actions to reduce the stress-generating sources, as well as to promote an increase in resilience and the development of intrinsic reinforcements to deal with uncertain threats.

**Keywords:** Men's Health; COVID-19; Mental Disorders; Stress, Psychological; Social Support.

## Introduction

The health crisis caused by COVID-19 exerted a negative impact on the mental health of the world's population. The experience of unknown situations and the fear of contamination imposed by the burden of the disease and its repercussions produced new daily stressors and exacerbated previous ones, already structurally present in societies. Together, these factors generate continuous situations of suffering and contribute to the increase in psychological distress, as evidenced by indicators of Common Mental Disorders (CMDs) and perceived stress. While the former refer to a set of symptoms, such as fatigue, insomnia, irritability, problems

concentrating and somatizations [1-6], the latter involves feelings about the uncontrollable and unpredictable nature of everyday events and the individual assessment of the ability to face such questions [4-5].

It is estimated that nearly one billion people are affected by some mental disorder due to the pandemic, with an emphasis on symptoms of anxiety, anguish and depression, and disorders resulting from alcohol use and substance abuse, in addition to the increase in suicide rates [1-6]. The deleterious effects caused by the pandemic have been measured and show a reduction in life expectancy at birth, in the ability to work, and in absenteeism and presentism, in addition to growth in unemployment and informality. Jointly, these situations install contexts that culminate in massive mental ailments in the male population [7-11].

Consequently, the COVID-19 pandemic resulted in a psychologically chaotic and dismal setting. The radical changes in the individuals' daily routines and the general context of fear and insecurity converge to unfamiliar situations and significant uncertainties that, due to their persistence over time, also provoke reactions of intolerance to this uncertainty. Intolerance to uncertainty is described as the predisposition of an individual to consider the possibility of a negative event occurring, regardless of the possibility of its actual occurrence as unacceptable [1-2].

Thus, intolerance to uncertainty is considered as one of the main underlying components of anxiety disorders [13], Obsessive Compulsive Disorder (OCD) [14], post-traumatic stress disorder in depression [15], depression [16] and panic disorder [17]. Thus, inability to cope with uncertainty can be a negative predictor for well-being [18].

Epidemiological indicators, such as the number of new cases, the rates and length of hospital stay, the unfavorable outcomes of the disease and the number of deaths have shown that the male population has been more impacted, evidencing that being a man is a risk factor for COVID-19 [19-20]. However, these results are restricted to the physical dimension of the disease and hardly advance in the analysis of the mental health dimension [21-24].

In addition to that, population-based studies investigating the mental health situation in the context of the pandemic have included predominantly female samples [25], which limits a more comprehensive identification of the magnitude of the problem and, consequently, restricts the adoption of coping measures to a specific audience. In this context, this study contributes to overcoming this gap in scientific knowledge, based on the research and appreciation of aspects

related to the male population, which permeate the subjectivities and the social, symbolic, psychological and psycho-emotional constructions of the population and their repercussions on health [26-27].

Given the above, it is considered crucial to give visibility to the mental health of male populations and to the aspects associated with the occurrence of mental disorders, to strengthen the confrontation of the life dimensions affected by the pandemic, in addition to contributing to overcoming this moment of profound health and social crisis generated by the dissemination of COVID-19.

Thus, this study aimed at analyzing the relationships between sociodemographic variables, intolerance to uncertainty (INT), social support and psychological distress, that is, indicators of Common Mental Disorders (CMDs) and Perceived Stress (PS), in Brazilian men during the COVID-19 pandemic.

## **Methods**

### ***Type of study***

A cross-sectional study with national coverage, of the web survey type and carried out during the period of social circulation restriction imposed by the health authorities in Brazil, for suppression of the coronavirus and control of the pandemic.

### ***Sample and participants***

The sample was estimated at 923 participants, considering the population of 64,520,660 Brazilian men with Internet access [28], 50% prevalence estimate, 95% confidence level, 5% precision, 80% power, effect of study design of two and a 20% increase for losses.

The snowball technique [29] was used to recruit the participants through digital social networks (Facebook®, Instagram®, WhatsApp®, Grindr®). This is a non-probabilistic sampling technique, performed by means of reference chains, where the first eligible and recruited participants invite new participants from their network of contacts (family, friends and acquaintances) who, in turn, indicate new participants and, thus, successively, until the estimated sample is minimally reached.

Initially, 25 participants were recruited, five from each of the Brazilian regions, who were called seeds. These participants were encouraged to send the research link to other men from their contact networks. At the end, 27 seeds were recorded, one from each of the Brazilian states.

The inclusion criteria adopted were as follows: being digitally literate to access the Internet and being at least 18 years old. The individuals excluded were those non-residents, in Brazil, who were in the country at the time of data collection.

### ***Procedures, measurements, variables and outcome***

For data collection, a questionnaire that was structured in blocks was used, including the following:

a) Sociodemographic characteristics: sexual identity, age, schooling, self-reported race/skin color, house-sharing or not, work situation and use of health plans.

b) Common Mental Disorders (CMDs): the Self-Reporting Questionnaire (SRQ-20) was used to screen CMDs. SRQ-20 is validated for use in Brazil with satisfactory performance indicators [30-32]. It consists of 20 items with dichotomous answer categories (0=no; 1=yes). The cutoff point for men is at least five positive answers [31].

c) Perceived Stress level: the ten-item version of the Perceived Stress Scale (PSS-10) [33], cross-culturally adapted [34] and validated for use in the Brazilian population [35] was used. The items have five-point Likert-type answer options (0=never, 1=almost never, 2=sometimes, 3=almost always; 4=always) and are distinguished in questions with positive and negative connotations, having to invert the score of the positive questions. The perceived stress score is calculated by the sum of the scores obtained in the ten items [34,33]. The levels were categorized as low (from zero to 13 points), moderate (from 14 to 26 points) and high (27 points or more) [36].

d) Level of intolerance to uncertainty: the Intolerance of Uncertainty Scale (IUS) [37], 12-item version (IUS-12) [12], cross-culturally adapted [38] and validated for use in the Brazilian population [39] was applied. The items contain answer options arranged in a Likert scale, varying from one (not at all characteristics) to five (very characteristic) and assessing two dimensions: Prospective IU (seven items) and Inhibitory IU (five items). The validation study in Brazil evidenced a two-factor structure with high correlation between them (0.83) [39], which can indicate the existence of a higher order factor (intolerance to uncertainty) [16,40]. The literature


published to the present day indicates diverse evidence of the one-factor solution of intolerance to uncertainty, as well as the use of its overall score [41,42]. The higher the score, the greater the level of intolerance to uncertainty. The IUS-12 latent construct showed adequate internal consistency ( $\alpha=0.89$ ;  $\omega=0.89$ ).

e) Level of social support: the instrument used was the 2-Way Social Support Scale (2-WSSS) [43], 20-item version, transculturally adapted and validated in Brazil [44]. 2-WSSS assesses four social support dimensions: 1) receiving emotional support (seven items); 2) receiving instrumental support (four items); 3) offering emotional support (four items); and 4) offering instrumental support (five items). The items are answered in a Likert scale, varying from zero (it never applies) to five (it always applies). Higher scores indicate higher levels of social support. In this study, 2-WSSS presented satisfactory internal consistency ( $\alpha=0.85$ ;  $\omega=0.96$ ).

The self-reporting questionnaire was elaborated in a specific data collection platform and made available through the Internet. From then onwards, it was widely publicized on the Facebook®, Instagram®, WhatsApp® and Grindr® digital social networks, by five trained researchers, for autonomous and voluntary adherence, as well as by making a direct invitation to the individuals who met the eligibility criteria. The data were collected between May and September 2020. The participants had access to the questionnaire after reading and signing the Free and Informed Consent Form, agreeing to participate in the study. Before each block of questions, the necessary information to answer them were made available, as well as the option not to answer any question. The instructions about the recall period for each instrument were clarified.

### ***Conceptual framework and study hypotheses***

Considering the evidence found in the literature, the understanding of human behaviors related to coping with the coronavirus outbreak and its impacts on people's mental health, *Directed Acyclic Graphs* (DAGs) were constructed [45,46] to represent the conceptual structure of the common mental disorders in Brazilian men during the COVID-19 pandemic, with emphasis on the effect of race/skin color, house-sharing or not, social support, perceived stress and intolerance to uncertainty. The directly observed variables are represented by rectangles and the latent variables, by circles (Figure 1).



**Figure 1.** Conceptual structure of the determinants for the occurrence of CMDs in Brazilian men during the COVID-19 pandemic.

DAGs are diagrams that allow coding and explaining conceptual hypotheses [46], with growing recognition in the field of causal research in epidemiology [45,47,48]. In the DAGs, the relationships between events are represented by vertices connected by edges; the vertices represent the variables and the edges show the possible ways or paths of relationships between variables, explaining causal links [46]. These causal paths can indicate direct causes, if there is an arrow going from one variable to another; or indirect causes, if there is a sequence of arrows starting from one variable and reaching another, passing through one or more intermediate/mediating variables. Paths that do not follow the direction of the arrows linking exposure and outcome represent potential “confounding paths” [45].

### ***Statistical analysis***

For the statistical analysis, the data were transferred from the Google Forms platform to a Microsoft Excel spreadsheet and were later organized in the SPSS software, version 24.0. Initially, a descriptive analysis of the variables of interest was performed for an estimation of frequencies (absolute and relative) and of prevalence values for CMDs. Subsequently, the

database was exported to the Mplus software, version 8.0, for Structural Equation Modeling (SEM) analyses.

To perform the SEM, the latent constructs were measured (measuring models) [49]. To assess the factor structure of the observable items, Exploratory Factor Analysis (EFA) was performed, followed by Exploratory Structural Equation Models (ESEM) and Confirmatory Factor Analysis (CFA), to validate dimensionality of the construct elaborated [50], having as criteria a standardized factor load  $\geq 0.3$  and a residual variance  $\leq 0.7$  [51,52].

To evaluate the structural model, constituted by the observed and latent variables, unadjusted and standardized regression coefficients were estimated, with 95% confidence intervals (95%CI) and  $p$ -value  $< 0.05$ . The size of the direct, indirect and total effects were classified as follows: weak/small (around 0.10), moderate/medium (close to 0.30) and strong/big ( $> 0.50$ ) [53].

The Weighted Least Squares Means and Variance Adjusted (WLSMV) estimator was used as a function of the modeling with categorical data. To re-specify the model, the Modification Indices ( $MI \geq 10$ ) and the Expected Parameter Changes ( $EPC \geq 0.25$ ) were assessed [54]. To assess fit of the models, the Root Mean Square Error of Approximation ( $RMSEA > 0.06$  – exceptionally  $< 0.08$ , with a 90% confidence interval below 0.08) [54], the Comparative Fit Index ( $CFI \geq 0.95$ ) and the Tucker Lewis Index ( $TLI \geq 0.95$ ) were adopted [53].

### ***Ethical considerations***

Ethical approval regarding this study was obtained from the institutional ethics committee (decision: 4,087,611; CAAE: 32889420.9.0000.5531). All the participants in this study were only included after informed consent had been obtained from them. All procedures performed in this study were compatible with the ethical standards of the institutional research committee and with those of the Declaration of Helsinki and its comparable ethical standards.

### **Results**

The study participants were 1,006 men. The predominant profile included non-heterosexuals (54.1%), aged from 29 to 39 years old (45.1%), higher education (73.8%), black-skinned individuals (59.2%), without a partner (67.3%), monthly income up to two minimum



wages (41.6%), living with family members/friends (76.6%) and workers (75.0%). A high proportion of people depending exclusively on the SUS care and services was observed (41.9%).

The overall prevalence of CMDs was 54.3%. Higher prevalence of CMDs was observed among the youngest individuals (62.7%), with lower schooling (55.3%), black-skinned (55.1%), without a partner (58.2%), and among those who lived alone (58.3%), did not work (59.4%), had high levels of stress (92.4%) and of intolerance to uncertainty (82.3%) and received low social support (67.2%). The moderate stress levels reached 60.9%, while 36.0% reported high levels of intolerance to uncertainty. Receiving social support was mentioned by 36.6% and offering social support, by 39.5%.

Socioeconomic Status (SES), Intolerance to Uncertainty (INT) and Social Support (SS) were treated as latent constructs. In general, the factor loads of the measuring models were high and statistically significant. The factor loads of the SES latent construct were above 0.60. In this construct, the highest load was observed for the occupational situation (OCP) indicator ( $\lambda=0.690$ ) and the lowest was found in the use of health plan (HP) indicator ( $\lambda=0.609$ ) (Table 1).

As for the INT construct, the exploratory factor analysis revealed, based on the eigenvalues, a solution of one predominant factor (eigenvalues=5.907) with a marked reduction for two factors (eigenvalue=1283). In the ESEM, the correlation between prospective and inhibitory IU was 0.992 and only the one-factor solution presented satisfactory fit indices. For this one-factor model, the highest load was observed for the “uncertainty makes me vulnerable” item (I7) ( $\lambda=0.754$ ) and the lowest one for “it is necessary to think about the future to avoid surprises” (I4) ( $\lambda=0.393$ ) (Table 1).

The SS measuring model was initially evaluated by the indicators of the first-order factors: Receiving Emotional Support (RES), Receiving Instrumental Support (RIS), Offering Emotional Support (OES) and Offering Instrumental Support (OIS), all with high factor loads and statistically significant. For RES and RIS, the highest loads were observed for the items “when I'm feeling down, there's someone I can count on” (S3) ( $\lambda=0.907$ ) and “if I'm in trouble someone will help me” (S13) ( $\lambda=0.845$ ) and the lowest ones for “I feel I have a network of people who value me” (S7) ( $\lambda=0.609$ ) and “there's someone who can help me fulfill my responsibilities” (S16) ( $\lambda=0.594$ ), respectively. For GES and GIS, greater burdens were verified for “I helped someone with their responsibilities when they were not able to fulfill them” (S18) ( $\lambda=0.700$ ) and “I comfort other people in difficult times” (S11) ( $\lambda=0.867$ ) and lower

ones for “I'm a person to whom others ask for help with tasks” (S20) ( $\lambda=0.401$ ) and “I'm a person available to listen to the others' problems” (S8) ( $\lambda=0.697$ ), respectively. The high correlation between the constructs of receiving ( $r=0.832$ ) and offering ( $r=0.864$ ) support endorsed the existence of the respective second-order factors called Receiving Support (RS) and Offering Support (OS), without a significant residual correlation ( $r=0.430$ ) (Table 1).

**Table 1.** Standardized factor loads of the measuring models of socioeconomic status, intolerance to uncertainty and social support among Brazilian men in the COVID-19 pandemic context. Brazil, 2020.

Latent variables	Indicating variables (codes)	SFL <sup>a</sup>	p-value
<b>SES</b>			
	Schooling (SCH)	0.631	<0.001
	Occupational situation (OCP)	0.690	<0.001
	Monthly income (INC)	0.664	<0.001
	Use of health plan (HP)	0.609	<0.001
<b>INT</b>			
	Uncertainty prevents me from living a full life (I1)	0.747	<0.001
	I profoundly loathe unforeseen events (I2)	0.710	<0.001
	I feel frustrated when I don't have all the information I need (I3)	0.669	<0.001
	It is necessary to think about the future to avoid surprises (I4)	0.393	<0.001
	A small unforeseen event can ruin everything, even with the best planning (I5)	0.749	<0.001
	When it's time to act, uncertainty paralyzes me (I6)	0.620	<0.001
	Uncertainty makes me vulnerable (I7)	0.754	<0.001
	I always want to know what the future will bring me (I8)	0.611	<0.001
	I hate to be taken by surprise (I9)	0.610	<0.001
	The slightest sign of doubt dissuades me from acting (I10)	0.678	<0.001
	I should be able to organize everything beforehand (I11)	0.634	<0.001

Uncertainty does not allow me to sleep well (I12)	0.694	<0.001
<b>Receiving</b>		
I have someone I can talk to about the pressures in my life (S1)	0.838	<0.001
There is at least one person with whom I can share most of the things (S2)	0.838	<0.001
When I'm feeling down, there's someone I can count on (S3)	0.907	<0.001
I have someone in my life who offers me emotional support (S4)	0.899	<0.001
There is at least one person in whom I feel I can trust (S5)	0.820	<0.001
There is someone in my life who makes me feel that I'm worthy (S6)	0.689	<0.001
I feel that I have a network of people who value me (S7)	0.609	<0.001
<b>RIS</b>		
If I'm in trouble, someone will help me (S13)	0.845	<0.001
I have someone to help me when I'm ill (S14)	0.686	<0.001
If I need money, I know someone who can help me (S15)	0.649	<0.001
There is someone who can help me fulfill my responsibilities (S16)	0.594	<0.001
<b>OIS</b>		
I'm a person who is available to listen to others' problems (S8)	0.697	<0.001
I seek to encourage people when they're feeling down (S9)	0.723	<0.001
People close to me tell me their deepest concerns (S10)	0.710	<0.001
I comfort other people in difficult times (S11)	0.867	<0.001
People trust me when they have problems (S12)	0.737	<0.001
<b>OES</b>		
I help others when they are too busy (S17)	0.685	<0.001
I helped someone with their responsibilities when they were not able to fulfill them (S18)	0.700	<0.001
I provided help when someone who lived with me was ill (S19)	0.588	<0.001
I'm a person to whom others ask for help with tasks (S20)	0.401	<0.001
<b>RS<sup>a</sup></b>		

RES <sup>b</sup>	0.882	<0.001
RIS <sup>b</sup>	0.949	<0.001
<b>Offering</b>		
OIS <sup>b</sup>	0.857	<0.001
OES <sup>b</sup>	0.740	<0.001

SFL: Standardized Factor Loads; SES: Socioeconomic Status; INT: Intolerance to Uncertainty; RES: Receiving Emotional Support; RIS: Receiving Instrumental Support; OIS: Offering Instrumental Support; OES: Offering Emotional Support; RS: Receiving Support; OS: Offering Support.

<sup>a</sup> 2<sup>nd</sup> order factor;

<sup>b</sup> 1<sup>st</sup> order factor.

The SES, INT and SS measuring models obtained satisfactory fit indices. The evaluation of the correlations between these latent constructs, both between the second-order factors of the SS model and in the model considering the correlations between all the latent variables, evidenced adequate discriminant validity ( $r < 0.90$ ) (Table 2).


**Table 2.** Fit indicators of the measuring models, using CMDs as response variable Brazil, 2020.

Indices	SES	INT	SS (RS □ OS)	CMM
RMSEA				
Index	0.038	0.049	0.046	0.046
90%CI	0.000 – 0.078	0.040 – 0.059	0.041 – 0.050	0.043 – 0.048
p-value	0.671	0.518	0.939	0.998
CFI	0.997	0.992	0.959	0.976
TLI	0.991	0.986	0.950	0.974
$r^a$				
RS □ OS	-	-	0.430	-
SES □ INT	-	-	-	-0.098
SES □ RS	-	-	-	-0.019
SES □ OS	-	-	-	-0.033
INT □ RS	-	-	-	-0.325



increase in the level of perceived stress were the following: receiving low social support, living alone and offering high social support (Figure 2).

Living alone mediated the effect of SES on CMDs, evidencing that men with higher socioeconomic status had higher stress levels and more prevalence of CMDs. The level of intolerance to uncertainty was also an important effect mediator: men who lived with family members/friends ( $\lambda=0.170$ ;  $p\text{-value}=0.021$ ), notably, non-black-skinned and those with higher socioeconomic status, had a higher level of intolerance to uncertainty and higher prevalence of CMDs. Intolerance to uncertainty also mediated the effect of perceived stress on CMDs, indicating that men with higher stress levels, especially those with lower socioeconomic status, non-black-skinned, who lived alone and with low levels of receiving social support and high levels of offering social support, had higher levels of intolerance to uncertainty ( $\lambda=0.592$ ;  $p\text{-value}<0.001$ ), resulting in a higher prevalence of CMDs. The factor that most contributed to the increase in intolerance to uncertainty was the high level of perceived stress, with a strong and significant effect (Figure 2).



SES: Socioeconomic Status; INT: Intolerance to Uncertainty (I1-I12: indicating variables); RES: Receiving Emotional Support (S1-S7: indicating variables); RIS: Receiving Instrumental Support (S13-S16: indicating variables); OIS: Offering Instrumental Support (S8-S12: indicating variables); OES: Offering Emotional Support (S17-S20: indicating variables); RS: Receiving Support; OS: Offering Support; SCH: Schooling;

OCP: Occupational Status; INC: Monthly Income; HP: Use of Health Plan; COL: Race/Skin Color; HS: House-Sharing; PS: Perceived Stress; I1-I12: Items 1 to 12; S1-S20: Items 1 to 20.  
 RMSEA: Root Mean Square Error of Approximation; 90%CI: 90% Confidence interval; CFI: Comparative Fit Index; TLI: Tucker-Lewis Index.

\*Statistically significant loads (p-value<0.05)

**Figure 2.** Structural equation model with direct and indirect specific effects for CMDs among Brazilian men in the COVID-19 pandemic context. Brazil, 2020.

The greatest total effects in CMDs were observed for higher level of perceived stress ( $\lambda=0.763$ ; p-value<0.001), receiving low social support ( $\lambda= 0.716$ ; p-value< 0.001), living alone ( $\lambda=0.587$ ; p-value<0.001), high intolerance to uncertainty ( $\lambda=0.421$ ; p-value<0.001) and high social support ( $\lambda=0.409$ ; p-value<0.001), all strong and significant. There was a small overall effect of socioeconomic status ( $\lambda=0.183$ ; p-value<0.001), but the indirect effects, mediated by high levels of perceived stress and intolerance to uncertainty, were robust and significant (Table 3).

It is to be noted that there was no statistically significant direct effect for CMDs from receiving and offering social support. However, among the specific indirect paths, the greatest effects included receiving low social support ( $\lambda= 0.442$ ; p-value<0.001), with strong and significant magnitude, and offering high social support ( $\lambda=0.275$ ; p-value<0.001), of medium size and significant, both mediated by high stress levels. The medium effects of living alone, mediated by high stress level ( $\lambda=0.281$ ; p-value<0.001), of high stress level, mediated by high level of intolerance to uncertainty ( $\lambda=0.249$ ; p-value<0.001) and of low social support, mediated by high levels of stress and intolerance to uncertainty ( $\lambda= 0.214$ ; p-value<0.001) also stood out (Table 3).

**Table 3.** Standardized total and indirect effects of the structural equation model, using CMDs as response variable among Brazilian men in the COVID-19 pandemic context. Brazil, 2020.

Paths	SFL	SE	Est/SE <sup>a</sup>	p-value <sup>b</sup>
<b>Total effects</b>				
INT→CMDs	0.421	0.047	8.87	<0.001
RS→CMDs	-0.716	0.051	-14.12	<0.001
OS→CMDs	0.409	0.021	19.11	<0.001

SES→CMDs	0.183	0.043	4.21	<0.001
PS→CMDs	0.763	0.039	19.46	<0.001
COL→CMDs	-0.021	0.040	-0.53	0.595
HS→CMDs	0.587	0.035	16.75	<0.001
<b>Specific indirect effects</b>				
<b>RS</b>				
RS→PS→CMDs	-0.442	0.039	-11.46	<0.001
RS→INT→CMDs	-0.032	0.025	-1.31	0.190
RS→PS→INT→CMDs	-0.214	0.037	-5.84	<0.001
<b>OS</b>				
OS→PS→CMDs	0.275	0.012	23.48	<0.001
OS→PS→INT→CMDs	0.134	0.022	5.97	<0.001
<b>SES</b>				
SES→HS→CMDs	0.060	0.014	4.22	<0.001
SES→PS→CMDs	-0.179	0.036	-4.96	<0.001
SES→INT→CMDs	0.021	0.027	0.79	0.429
SES→RS→CMDs	0.001	0.002	0.38	0.701
SES→HS→PS→CMDs	0.060	0.014	4.22	<0.001
SES→RS→PS→CMDs	0.012	0.025	0.48	0.629
SES→OS→PS→CMDs	-0.015	0.015	-1.00	0.318
SES→HS→INT→CMDs	-0.015	0.008	-1.88	0.060
SES→PS→INT→CMDs	-0.087	0.022	-3.96	<0.001
SES→RS→INT→CMDs	0.001	0.002	0.49	0.625
SES→HS→RS→CMDs	0.001	0.001	-0.42	0.673
SES→HS→RS→PS→CMDs	-0.008	0.006	-1.37	0.170
SES→HS→OS→PS→CMDs	0.003	0.003	0.93	0.351
SES→HS→PS→INT→CMDs	0.029	0.009	3.31	0.001



SES→RS→PS→INT→CMDs	0.006	0.012	0.48	0.629
SES→OS→PS→INT→CMDs	-0.007	0.008	-0.99	0.325
SES→HS→RS→INT→CMDs	-0.001	0.001	-1.06	0.289
SES→HS→RS→PS→INT→CMDs	-0.004	0.003	-1.37	0.171
SES→HS→OS→PS→INT→CMDs	0.002	0.002	0.92	0.356
<b>PS</b>				
PS→INT→CMDs	0.249	0.040	6.22	<0.001
<b>COL</b>				
COL→PS→CMDs	0.135	0.006	20.80	<0.001
COL→INT→CMDs	-0.034	0.019	-1.78	0.075
COL→SES→CMDs	0.135	0.006	20.80	<0.001
COL→SES→HS→CMDs	0.026	0.006	4.21	<0.001
COL→SES→PS→CMDs	-0.079	0.016	-4.93	<0.001
COL→PS→INT→CMDs	0.066	0.011	5.92	<0.001
COL→SES→INT→CMDs	0.009	0.012	0.79	0.429
COL→SES→RS→CMDs	0.000	0.001	0.38	0.702
COL→SES→HS→PS→CMDs	0.026	0.006	4.21	<0.001
COL→SES→RS→PS→CMDs	0.005	0.011	0.48	0.629
COL→SES→OS→PS→CMDs	-0.007	0.007	-1.00	0.319
COL→SES→HS→INT→CMDs	-0.007	0.004	-1.88	0.060
COL→SES→PS→INT→CMDs	-0.038	0.010	-3.95	<0.001
COL→SES→RS→INT→CMDs	0.000	0.001	0.49	0.625
COL→SES→HS→RS→CMDs	0.000	0.001	-0.42	0.673
COL→SES→HS→RS→PS→CMDs	-0.004	0.003	-1.37	0.170
COL→SES→HS→OS→PS→CMDs	0.001	0.002	0.93	0.351
COL→SES→HS→PS→INT→CMDs	0.013	0.004	3.30	0.001
COL→SES→RS→PS→INT→CMDs	0.003	0.005	0.48	0.629

COL→SES→OS→PS→INT→CMDs	-0.003	0.003	-0.99	0.325
COL→SES→HS→RS→INT→CMDs	0.000	0.000	-1.06	0.289
COL→SES→HS→RS→PS→INT→CMDs	-0.002	0.001	-1.37	0.171
COL→SES→HS→OS→EP→INT→CMDs	0.001	0.001	0.92	0.356
<b>HS</b>				
HS→PS→CMDs	0.281	0.012	23.79	<0.001
HS→INT→CMDs	-0.071	0.033	-2.17	0.030
HS→RS→CMDs	-0.002	0.006	-0.43	0.668
HS→RS→PS→CMDs	-0.038	0.026	-1.45	0.146
HS→OS→PS→CMDs	0.015	0.016	0.97	0.334
HS→PS→INT→CMDs	0.136	0.023	5.93	<0.001
HS→RS→INT→CMDs	-0.003	0.003	-1.10	0.273
HS→RS→PS→INT→CMDs	-0.019	0.013	-1.46	0.144
HS→OS→PS→INT→CMDs	0.007	0.008	0.96	0.338

SFL: Standardized Factor Loads; SE: Standard Error; Est/SE: Ratio between SFL estimate and SE.

SES: Socioeconomic Status; INT: Intolerance to Uncertainty; RS: Receiving Support; OS: Offering Support. HS: House-Sharing; COL: Self-reported Race/Skin Color; PS: Perceived Stress; CMDs: Common Mental Disorders.

<sup>a</sup> Statistically significant if  $-1.96 > \text{Est/SE} > 1.96$ .

<sup>b</sup> Statistically significant if  $< 0.05$ .

## Discussion

The study evidenced a high prevalence of CMDs in men, which corroborates other findings on the high rates of mental illness in the Brazilian population during the COVID-19 pandemic. In the pandemic context, no studies analyzing CMDs were identified, hindering comparisons. However, high prevalence values of sadness/depression, anxiety/nervousness, worsening of sleep problems and panic syndrome were observed in Brazil during the pandemic [55-56].

The rapid spread of COVID-19, associated with the need for social distancing, forced quarantine and blockades across the world, has placed imposed high stress levels on people [57-58]; and sustaining these measures has increased the prevalence of CMDs in the populations [59-60], contributing to the increase in depressive and anxiety disorders, identified by the WHO as the most common psychiatric diseases in the world's population [61-62].

In this study, the high prevalence of CMDs is more pronounced among black-skinned men, higher socioeconomic status, living alone and higher levels of PS and INT, and these last two factors are the ones most strongly associated with CMDs. Although OS has not presented any direct effect, specific indirect effects, such as receiving low social support and offering high social support, both mediated by high stress levels, were associated with CMDs.

Although the association between CMDs and black skin color observed in our study can be related to the predominance of black-skinned individuals in the study population, being non-Hispanic and black-skinned is identified as a risk factor for suffering discrimination associated with COVID-19 [63]. Nevertheless, the direct effect of skin color on the prevalence of CMDs is justified by the literature, which evidences a direct association between race or skin color and mental health over time, either in pandemic situations or not [64-65], outlining the history of discrimination and its lasting consequences.

The socioeconomic differences seem to exert an effect on the development of CMDs, although the underlying mechanisms of that association are not yet well understood. The effect of higher socioeconomic status on the higher occurrence of CMDs is apparently controversial to what is documented in the literature [66-67], but some aspects can be evoked to understand this finding in this context of the COVID-19 pandemic: a) study carried out in a virtual environment, which contributed to the lower access of men with low socioeconomic status and possible underestimation of CMDs in this stratum; b) men with higher socioeconomic status may have a better perception of their mental health status due to a better schooling level [66], reflecting higher CMDs rates in a self-reporting questionnaire; c) due to the restrictions imposed by the COVID-19 pandemic, men with higher socioeconomic status may be experiencing, abruptly and suddenly, more financial deprivation related to business and work, in social relationships and in the life dynamics, situations that are already part of the daily life of those with lower socioeconomic status even before the pandemic and, due to the absence of intrinsic reinforcement, they end up in mental distress; and d) greater fear of poverty is associated with higher stress and anxiety levels among people with higher socioeconomic status during the pandemic, and maintenance of these levels reflects in mental illness [68]. Thus, this finding needs to be researched, longitudinally, in order to explore the possibilities of the causal link.

The “living alone” factor mediated the effect of SES on the CMDs, so that those with higher socioeconomic status presented higher stress levels and higher prevalence of CMDs.

These two factors, mediated by the high PS and INT levels, presented considerable indirect effects for CMDs. Living with family members/friends was related to a higher level of intolerance to uncertainty and to more prevalence of CMDs. In this context, although measures of social distancing and quarantine of the population are important for reducing morbidity and mortality due to COVID-19, their effects on the health of the population are undeniable. Regarding mental health, feelings and emotions such as boredom, loneliness, anger and sadness can emerge, not only because of the precautionary and control measures imposed, but also because of the perception of vulnerability to contagion and risk of illness, especially among those who they live alone [69].

The PS and INT factors are very inter-related, so that men with higher PS levels had higher INT levels, resulting in greater prevalence of CMDs. The PS level was influenced by several factors such as lower socioeconomic status, non-black-skinned people, living alone, receiving low social support and offering high social support. A study carried out with Brazilian parents also identified that PS was the variable most strongly related to the CMD symptoms, showing that the higher the level of perceived stress, the greater the suspicion of these disorders [70]. Several factors have been associated with the occurrence of stress during the pandemic, which include concern about physical health and precautionary and disease control measures [70], anger and confusion arising from quarantine and social distancing [69], income reduction [69,25], fear of infection [69] and being part of a risk group for COVID-19 [25]. Consequently, the public policies must target their efforts towards reducing the stress sources with the aim of preventing occurrence of CMDs.

Regarding the socioeconomic situation of the men, the excess of demands, the fear of income loss and the holding of the capital power may have contributed to increased stress levels and, consequently, to a higher prevalence of CMDs. This association is an important warning sign to be included in the global public health agenda, in a commitment to promote actions to reduce stress, increase literacy in male mental health for the perception of stress and self-management of mental health care [71].

INT is a vulnerability factor associated with the development and maintenance of mental disorders [72]. Uncertainty about the future is a potentially stressful condition [73], hence the strong association with the occurrence of CMDs. The study evidenced that INT is negatively associated with mental well-being, especially when mediated by loneliness and fear of COVID-

19, which are feelings intensely experienced during the pandemic [74]. Thus, it is necessary to implement actions to promote increased resilience and development and/or strengthening of intrinsic reinforcements to deal with the impacts of regular exposure to uncertain threats.

These results reinforce that mental health goes beyond the biological issues, being the result of multiple social, economic, psychological and cultural interactions[75]. Therefore, it is not only related to the absence of manifestations of mental distress, with the need to assess the frequency and intensity of these events and how this affects people's quality of life [76-78]. Predominance of negative feelings reported by the male population is noticed, them being expected and compatible given the continuity and even worsening of the pandemic scenario. Therefore, it is essential to investigate and understand the factors that precede mental illness during the COVID-19 pandemic in different populations, so that policy makers can implement effective countermeasures [59].

It is also highlighted that, although there are groups at higher risk, COVID-19 ends up affecting different social segments and men tend to adhere less to the care measures and to neglect their health [79]. The occurrence and extension of this scenario can be even more impactful when considering social support, whether from the perspective of receiving or offering emotional and instrumental support. Thus, it is indispensable to formulate and strengthen already existing actions to promote psychosocial and emotional support for men, in an exercise of citizenship and health care promotion that strengthens socio-affective networks, enabling men to have someone to count on in critical moments like a pandemic.

As shown by the results, men presented moderate stress levels and high indices of intolerance to uncertainty and social support. Feeling such as anxiety, insecurity, stress, irritation, agony, boredom, shock and fear had already been detected in this population during the pandemic [27]. However, they can be generators of greater impacts on mental health, as men start to perceive themselves as more vulnerable in the face of uncertainties, which directly compromises tolerance to uncertain threats and increases the prevalence of CMDs, mainly due to the fact that, during the research period, Brazil continued with the disease lacking sanitary control and without collective vaccination or consistent therapeutic treatment measures.

In addition to the categories of psychological disorders such as stress, other structural categories need to be analyzed in the scenario of impacts on male mental health, such as race/skin color, territory (geographic location and area of residence), since our findings identified that

black-skinned men, those with higher socioeconomic status, who lived alone, and with higher levels of stress and intolerance to uncertainty had a higher prevalence of CMDs. Thus, an intersectional perspective needs to be employed, as a way of explaining the vulnerabilities, inequalities, inequities and necropolitics that promote overlapping impacts on the health, quality of life and well-being of men who belong to marginalized groups, an aspect confirmed in our study when we identified the disparity in economic status among non-black-skinned men.

Cultural factors can partially explain men's adherence to the health protection measures and consequent increase in the incidence of CMDs during the pandemic period. Generally, men are encouraged to adopt patriarchal values where self-care and health practices go against their masculinity and reinforce the stereotypes of independence, virility, courage and strength which, in turn, can prevent them from adopting protective health [78-82].

Although knowledge about the disease is advanced, there are still many gaps. Commonly, guided behaviors are based on scientific evidence, protocols and guidelines. However, the recommendations themselves present certain degree of uncertainty, which generates doubts and discomfort in the face of the unknown on the part of authorities, health professionals and the population [82]. Given this scenario, it is necessary to accept the existence of these uncertainties, limitations and incompleteness in current knowledge. Transparency regarding these issues is a strategy to deal with the unknown in a rational and ethical way. In addition, the relevance of providing information and guidelines as recommended by the health authorities is emphasized, as a way to avoid excessive simplifications and the dissemination of easy answers if they do not exist. Thus, gaining trust becomes one of the greatest challenges and ends up fighting mistrust and fake news [83], which intensify the emergence of conspiracy theories, misinformation and infodemic [84-88].

Our study adopted a cross-sectional design and a sampling technique with recruitment by non-probabilistic methods, which limit the causal inference and external validity of the results. In addition to that, participation of men with no access to the interview via the Internet was excluded, which may have over- or under-estimated the prevalence of CMDs in the strata evaluated.

## **Conclusion**

The results of this study point out the high prevalence of CMDs in men in the pandemic context and reinforce the need to assess the factors that precede this disease and to replicate the study with women. In addition to that, they endorse the importance of constructing latent variables to measure subjective aspects, since three important latent constructs were revealed to be better studied, namely: Socioeconomic Status, Intolerance to Uncertainty, and Social Support.

In addition to that, the study allowed identifying a higher prevalence of CMDs among younger men, with lower schooling, black-skinned, without a partner, who lived alone, did not work, with high levels of stress and intolerance to uncertainty and who received low social support. It was also evidenced that the high levels of stress and intolerance to uncertainty were the factors that presented the strongest direct associations with the occurrence of CMDs.

Thus, the importance of developing mental health promotion policies and actions for male populations is highlighted, considering that intervening in the aspects associated with the occurrence of CMDs is crucial to facing the impacts of the COVID-19 pandemic in the different life dimensions.

**Author Contributions:** Conceptualisation, Sousa AR and Pereira A; methodology, validation, formal analysis, investigation, data curation, writing-original draft preparation, writing-review and editing and visualisation, all authors; supervision, Sousa AR. All authors have read and agreed to the published version of the manuscript

**Funding:**

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki, and the study was approved by the Ethical Review Committee.

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The dataset generated during the current study are not publicly available but are available from the corresponding author on reasonable request.

**Acknowledgments:** To the men who participated in the study.

**Conflicts of Interest:** The authors declare no conflicts of interest

**References**

1. Dal'Bosco EB, Floriano LSM, Rangel AGSS, Ribas MC, Cavalheiro APG, Silva CLD, Cabral LPA. Coping in mental health during social isolation: analysis in light of Hildegard Peplau. *Rev Bras Enferm.* **2021** Oct 1;75(2):e20201207. English, Portuguese. doi: 10.1590/0034-7167-2020-1207.
2. Sousa AR, Teixeira JRB, Mota TN, Santana TDS, Santos SDD, Merces MCD, Carvalho ESS, Sousa ÁFL. Coping strategies, concerns, and habits of Brazilian men in the COVID-19 context. *Rev Bras Enferm.* **2021** Jul 12;74Suppl 1(Suppl 1):e20210040. English, Portuguese. doi: 10.1590/0034-7167-2021-0040.
3. Faro, A.; Bahiano, M. A.; Nakano, T. C.; Reis, C.; Silva, B. F. P.; Vitti, L. S. COVID-19 e saúde mental: a emergência do cuidado. *Estudos de Psicologia (Campinas)* **2020**, 37, 1-14, doi:10.1590/1982-0275202037e200074.
4. Mari, J. J.; Oquendo, M. A. Mental health consequences of COVID-19: the next global pandemic. *Trends Psychiatry Psychother* **2020**, 42, 219-220, doi:10.1590/2237-6089-2020-0081.
5. Moreira, W.C.; Sousa, A.R.; Nóbrega, M.D.P.S.S. Mental illness in the general population and health professionals during COVID-19: A scoping review. *Texto Contexto Enferm.* **2020**, 29, e20200215, English, Portuguese. doi:10.1590/1980-265X-TCE-2020-0215
6. Paiano M, Jaques AE, Nacamura PAB, Salci MA, Radovanovic CAT, Carreira L. Mental health of healthcare professionals in China during the new coronavirus pandemic: an integrative review. *Rev Bras Enferm.* **2020** Sep 18;73(suppl 2):e20200338. English, Portuguese. doi: 10.1590/0034-7167-2020-0338.
7. Mamun, M. A.; Griffiths, M. D. First COVID-19 suicide case in Bangladesh due to fear of COVID-19 and xenophobia: possible suicide prevention strategies. *Asian J Psychiatr.* **2020**, 51, 102073, doi: 10.1016/j.ajp.2020.102073
8. Goyal, K.; Chauhan, P.; Chhikara, K.; Gupta, P; Singh MP. Fear of COVID 2019: first suicidal case in India. *Asian J Psychiatr.* **2020**, 49, 101989, doi: 10.1016/j.ajp.2020.101989
9. Gonzalez-Diaz, J. M.; Cano, J. F.; Pereira-Sanchez, V. Psychosocial impact of COVID-19-related quarantine: reflections after the first case of suicide in Colombia. *Cad Saúde Pública* **2020**, 36, e00117420, doi:10.1590/0102-311X00117420



10. Gonçalves Júnior, J.; de Sales, J. P.; Moreno, M. M.; Rolim-Neto, M. L. The Impacts of SARS-CoV-2 Pandemic on Suicide: A Lexical Analysis. *Front Psychiatry* **2021**, *10*, 593918, doi: 10.3389/fpsyt.2021.593918.
11. Sousa, A. R.; Moreira, W. C.; Queiroz, A. M.; Rezende, M. F.; Teixeira, J. R. B.; Mercês, M. C.; Silva, A. V.; Camargo, E. L. S. COVID-19 pandemic decrease men's mental health: background and consequence analysis. *J. bras. Psiquiatr.* **2021**, *70*, 141-148, doi: 10.1590/0047-2085000000327.
12. Carleton, R. N.; Norton, M. A.; Asmundson, G. J. Fearing the unknown: a short version of the Intolerance of Uncertainty Scale. *J Anxiety Disord.* **2007**, *21*, 105-117. doi: 10.1016/j.janxdis.2006.03.014.
13. Morriss, J.; Christakou, A.; van Reekum, C. M. Nothing is safe: Intolerance of uncertainty is associated with compromised fear extinction learning. *Biological Psychology.* **2016**, *121*, 187–193. doi: 10.1016/j.biopsycho.2016.05.001.
14. Holaway, R. M.; Heimberg, R. G.; Coles, M. E. A comparison of intolerance of uncertainty in analogue obsessive-compulsive disorder and generalized anxiety disorder. *J. Anxiety Disord.* **2006**, *20*, 158-174, doi: 10.1016/j.janxdis.2005.01.002.
15. Hollingsworth, D. W.; Gauthier, J. M.; McGuire, A. P.; Peck, K. R.; Hahn, K. S.; Connolly, K. M. Intolerance of uncertainty mediates symptoms of PTSD and depression in African American veterans with comorbid PTSD and substance use disorders. *J. Black Psychol.* **2018**; *44*, 667–688. doi: 10.1177/0095798418809201.
16. McEvoy, P. M.; Mahoney, A. E. J. Achieving certainty about the structure of intolerance of uncertainty in a treatment-seeking sample with anxiety and depression. *J. Anxiety Disord.* **2011**, *25*, 112-122, doi:10.1016/j.janxdis.2010.08.010.
17. Tolin, D. F.; Abramowitz, J. S.; Brigidi, B. D.; Foa, E. B. Intolerance of Uncertainty in obsessive-compulsive disorder. *J. Anxiety Disord* **2003**, *17*, 233-242, doi: 10.1016/s0887-6185(02)00182-2
18. Satici, B.; Saricali, M.; Satici, S. A.; Griffiths, M. D. Intolerance of Uncertainty and Mental Wellbeing: Serial Mediation by Rumination and Fear of COVID-19. *Int J Ment Health Addict.* **2020**, *15*:1-12. doi: 10.1007/s11469-020-00305-0
19. World Health Organization. WHO Coronavirus (COVID-19) Dashboard. 2021. Available from: <https://covid19.who.int/>

20. Brasil. Ministério da Saúde (BR). Boletim epidemiológico especial. Doença pelo Coronavírus COVID-19. Brasília, DF (BR): MS, 2021. Available online: [https://www.gov.br/saude/pt-br/media/pdf/2021/outubro/01/boletim\\_epidemiologico\\_covid\\_82.pdf](https://www.gov.br/saude/pt-br/media/pdf/2021/outubro/01/boletim_epidemiologico_covid_82.pdf)
21. Brasil. Ministério da Saúde (BR). Boletim epidemiológico especial. Doença pelo Coronavírus COVID-19. Brasília, DF (BR): MS, 2020. Available online: <http://saude.gov.br/images/pdf/2020/July/15/Boletim-epidemiologico-COVID-22.pdf> (accessed on 30 April 2021).
22. Peckham, H.; de Gruijter, N. M.; Raine, C.; Radziszewska, A.; Ciurtin, C.; Wedderburn LR et al. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission. *Nat Commun.* **2020**,11, 1-10. doi: 10.1038/s41467-020-19741-6
23. Gorbalenya, A. E.; Baker, S. C.; Baric, R. S.; Groot, R. J.; Drosten, C.; Gulyaeva, A. A.; Ziebuhr, J. et al. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat. Microbiol.* **2020**, 5, 536–544, doi: 10.1038/s41564-020-0695-z
24. Scully, E. P.; Haverfield, J.; Ursin, R. L.; Tannenbaum, C.; Klein, S. L. Considering how biological sex impacts immune responses and COVID-19 outcomes. *Nat. Rev. Immunol.* **2020**, 20, 442–447, doi: 10.1038/s41577-020-0348-8
25. Duarte, M. Q.; Santo, M. A. S.; Lima, C. P.; Giordani, J. P.; Trentini, C. M. COVID-19 e os impactos na saúde mental: uma amostra do Rio Grande do Sul, Brasil. *Cien Saude Colet* **2020**, 25, 3401–3412. doi:10.1590/1413-81232020259.16472020
26. Windsor, L. C.; Reinhardt, G. Y.; Windsor, A. J, OstergardI, R.; Allen, S.; Burns, C.; Giger, J.; Wood, R. Gender in the time of COVID-19: Evaluating national leadership and COVID-19 fatalities. *PLOS ONE* **2020**,15, e0244531, doi.org/10.1371/journal.pone.0244531
27. Sousa, A. R.; Santana, T. S.; Moreira, W. C.; Sousa, Á. F. L.; Carvalho, E. S. S.; Craveiro, I. Emoções e estratégias de coping de homens à pandemia da covid-19 no Brasil. *Texto e Contexto Enferm*, **2020**, 29, e20200248. doi:10.1590/1980-265x-tce-2020-0248.
28. Núcleo de Informação e Coordenação do Ponto BR (NIC.br). Pesquisa sobre o uso das tecnologias de informação e comunicação: pesquisa TIC Domicílios, ano 2019: Tabelas. São Paulo: 2020. Available online:

- <http://cetic.br/pt/arquivos/domicilios/2019/individuos/#tabelas>. Acesso em: 18 mai. 2021 (accessed on 30 April 2021).
29. Biernacki, P.; Waldorf, D. Snowball sampling: Problems and techniques of chain referral sampling. *Sociol. Methods Res.* **1981**, 10, 141-163, doi:10.1177/004912418101000205
  30. Mari, J. J.; Williams, P. A validity study of a psychiatric screening questionnaire (SRQ-20) in primary care in the city of Sao Paulo. *Br J Psychiatry* **1986**, 148, 23-6. doi: 10.1192/bjp.148.1.23.
  31. Santos, K. O. B.; Araújo, T. M.; Sousa Pinho, P.; Silva, A. C. C. Avaliação de um instrumento de mensuração de morbidade psíquica: estudo de validação do Self-Reporting Questionnaire (SRQ-20). *Rev. Baiana de Saúde Pública* **2010**, 34, 544-544. doi: 10.22278/2318-2660.
  32. Santos, K. O. B.; Carvalho, F. M.; Araújo, T. M. Consistência interna do self-reporting questionnaire-20 em grupos ocupacionais. *Rev Saude Publica* 2016, 50, 6. doi:10.1590/S1518 8787.2016050006100.
  33. Cohen, S.; Kamarck, T.; Mermelstein, R. A global measure of perceived stress. *J Health Soc Behav*, **1983**, 24, 385–396, doi:10.2307/2136404.
  34. Luft, C. D.; Sanches, S. O.; Mazo, G. Z.; Andrade, A. Versão brasileira da Escala de Estresse Percebido: tradução e validação para idosos. *Rev Saúde Pública* **2007**, 41, 606-15, doi: 10.1590/S0034-89102007000400015.
  35. Machado, W. L.; Damásio, B. F.; Borsa, F. C.; Silva, J. P. Dimensionalidade da Escala de Estresse Percebido (Perceived Stress Scale, PSS10) em uma amostra de professores. *Psicol Reflex Crit* **2014**, 27,38-43, doi: 10.1590/S0102-79722014000100005.
  36. Ma. Y.; Rosenheck, R.; He, H. Psychological stress among health care professionals during the 2019 novel coronavirus disease Outbreak: Cases from online consulting customers. *Intensive Crit Care Nurs.* **2020**,61, 102905. doi: 10.1016/j.iccn.2020.102905.
  37. Freeston, M. H.; Rheaume, J.; Letarte. H.; Dugas, M. J.; Ladouceur, R. Why Do People Worry? *Personality and Individual Differences* **1994**,17, 791-802, doi: 10.1016/0191-8869(94)90048-5
  38. Kretzmann, R. P. Intolerance of Uncertainty Scale (IUS-12) para uso no Brasil: adaptação transcultural e propriedades psicométricas [dissertation]. Porto Alegre: Universidade Federal do Rio Grande do Sul; 2018.

39. Kretzmann, R. P., Gauer, G. Psychometric properties of the Brazilian Intolerance of Uncertainty Scale – Short Version (IUS-12). *Trends Psychiatry Psychother*, **2020**, 42, 129-137, doi:10.1590/2237-6089-2018-0087.
40. Norton, P. J. A psychometric analysis of the Intolerance of Uncertainty Scale among four racial groups. *J. Anxiety Disord*, **2005**, 6, 699–707, doi: 10.1016/j.janxdis.2004.08.002
41. Yao, N.; Qian, M.; Jiang, Y.; Elhai, J. D. The Influence of Intolerance of Uncertainty on Anxiety and Depression Symptoms in Chinese-speaking Samples: Structure and Validity of The Chinese Translation of The Intolerance of Uncertainty Scale. *J Pers Assess*. **2021**, 103, 406-415, doi: 10.1080/00223891.2020.1739058
42. Bottesi, G.; Noventa, S.; Freeston, M. H.; Ghisi, M. Seeking certainty about Intolerance of Uncertainty: Addressing old and new issues through the Intolerance of Uncertainty Scale-Revised. *PloS one* **2019**, 14, e0211929, doi: 10.1371/journal.pone.0211929.
43. Shakespeare-Finch, J.; Obst, P. L. The development of the 2-Way Social Support Scale: a measure of giving and receiving emotional and instrumental support. *J Pers Assess*. **2011**, 93, 483-90, doi: 10.1080/00223891.2011.594124.
44. Bastianello, M. R. Implicações do Otimismo, Autoestima e Suporte Social Percebido na Qualidade de Vida de Mulheres com Câncer de Mama [Implications of optimism, self esteem and perceived social support on quality of life of women with breast cancer]. PhD Thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil, **2015**.
45. Glymour, M. M.; Greenland, S.; Causal diagrams. Ch. 12. In: Rothman KJ, Greenland S, Lash TL (eds) *Modern epidemiology*, 3rd edn. Lippincott, Philadelphia. **2008**.
46. Hernán, M. A.; Robins, J. M. Causal Inference: What If. Boca Raton: Chapman & Hall/CRC. **2020**.
47. Cortes, T. R.; Faerstein, E. S.; Claudio, J. Utilização de diagramas causais em epidemiologia: um exemplo de aplicação em situação de confusão. *Cad Saúde Pública* **2016**, 32, e00103115, doi: 10.1590/0102-311X00103115.
48. Werneck, G. L. Diagramas causais: a epidemiologia brasileira de volta para o futuro. *Cadernos de Saúde Pública* **2016**, 32, e00120416. doi:10.1590/0102-311X00120416
49. Marôco, J. Análise de Equações Estruturais - Fundamentos teóricos, Software e Aplicações. 2. ed. Pêro Pinheiro: Report Number, 2014.

50. Marsh, H. W.; Muthén, B.; Asparouhov, T.; Lüdtke, O.; Robitzsch, A.; Morin, A. J. S.; Trautwein, U. Exploratory structural equation modeling, integrating CFA and EFA: Application to students' evaluations of university teaching, *Structural Equation Modeling. Structural equation modeling: A multidisciplinary journal* **2009**.  
doi:10.1080/10705510903008220
51. Hair, J. F. J.; Black, W. C.; Babin, B. J.; Anderson R.E.; Tatham, R. L. *Análise multivariada de dados*. 6.ed.; Bookman: Porto Alegre, Brazil, 2009.
52. Reichenheim, M. E.; Hökerberg, Y. H. M.; Moraes, C. L. Assessing construct structural validity of epidemiological measurement tools: a seven-step roadmap. *Cad Saúde Pública* **2014**, 30, 927-939, doi: 10.1590/0102-311X00143613.
53. Kline, R. B. *Principles and Practice of Structural Equation Modeling*. New York, NY. 2015.
54. Brown, T. A. *Methodology in the social sciences. Confirmatory factor analysis for applied research*, 2nd ed. New York, NY, US: 2015.
55. Barros, A. J. D.; Victora, C. G.; Menezes, A. M. B.; Horta, B. L.; Hartwig, F.; Victora, G. et al. Social distancing patterns in nine municipalities of Rio Grande do Sul, Brazil: the Epicovid19/RS study. *Rev Saude Pública* **2020**, 54: 75, doi: 10.11606/s1518-8787.2020054002810.
56. Lipp, M. E. N.; Lipp, L. M. N. Stress e transtornos mentais durante a pandemia da COVID-19 no Brasil. *Revista da Academia Paulista de Psicologia* **2020**, 40, 180-19.
57. Bao, Y.; Sun, Y.; Meng, S.; Shi, J.; Lu, L. 2019-nCoV epidemic: address mental health care to empower society. *The Lancet*, 395, e37-e38, doi:10.1016/S0140-6736(20)30309-3.
58. Dubey, S.; Biswas, P.; Ghosh, R.; Chatterjee, S.; Dubey, M. J.; Chatterjee, S.; Lahiri, D.; Lavie, C. J. Psychosocial impact of COVID-19. *Diabetes Metab. Syndr* **2020**, 14, 779–788, doi: /10.1016/j.dsx.2020.05.035.
59. Wu, T.; Jia, X.; Shi, H.; Niu, J.; Yin, X.; Xie, J.; Wang, X. Prevalence of mental health problems during the COVID-19 pandemic: A systematic review and meta-analysis. *J Affect Disord.* **2021**, 15, 281:91-98. doi: 10.1016/j.jad.2020.11.117.
60. Cui, L.; Wang, X.; Wang, H. Challenges facing coronavirus disease 2019: Psychiatric services for patients with mental disorders. *Psychiatry and clinical neurosciences.* **2020**, 74, 371–372, doi: 10.1111/pcn.13003.

61. World Health Organization (WHO). Depression and other common mental disorders: Global Health Estimates. 2017 Available online: [www.who.int/mental\\_health/management/depression/prevalence\\_global\\_health\\_estimates/en/](http://www.who.int/mental_health/management/depression/prevalence_global_health_estimates/en/) (accessed on 30 April 2021).
62. Souza, I. M.; Sousa, J. P. M. Brazil: world leader in anxiety and depression rates. *Revista Brasileira de Psiquiatria* **2017**, *39*, 384. doi: 10.1590/1516-4446-2017-2300.
63. Liu, Y.; Finch, B. K.; Brenneke, S. G.; Thomas, K.; Le, P. D. Perceived Discrimination and Mental Distress Amid the COVID-19 Pandemic: Evidence From the Understanding America Study. *Am J Prev Med* **2020**, *59*, 481-492. doi:10.1016/j.amepre.2020.06.007.
64. Smolen, J. R.; Araujo, E. M. Raça/cor da pele e transtornos mentais no Brasil: uma revisão sistemática. *Cienc Saude Coletiva* **2017**, *22*, 4021-30, doi:10.1590/1413-812320172212.19782016
65. Souza, A. S. R.; Souza, G. F.A.; Souza, G. A.; Cordeiro, A. L. N.; Praciano, G. A. F.; Alves, A. C. S, et al. Factors associated with stress, anxiety, and depression during social distancing in Brazil. *Rev Saúde Publica* 2021; *55*: 5. doi:10.11606/s1518-8787.2021055003152.
66. Dorner, T. E.; Mittendorfer-Rutz, E. Socioeconomic inequalities in treatment of individuals with common mental disorders regarding subsequent development of mental illness. *Soc Psychiatry Psychiatr Epidemiol.* **2017**, *52*, 1015-1022. doi:10.1007/s00127-017-1389-6
67. Jokela, M.; Batty, G.; Vahtera, J.; Elovainio, M.; Kivimäki, M. Socioeconomic inequalities in common mental disorders and psychotherapy treatment in the UK between 1991 and 2009. *British Journal of Psychiatry*, **2013**, *202*(2), 115-120. doi:10.1192/bjp.bp.111.098863
68. Mental Health Outcomes of the COVID-19 Pandemic and a Collapsing Economy: Perspectives from a Developing Country. *Psychiatry Res* **2020**, *294*:113520. doi: 10.1016/j.psychres.2020.113520.
69. Brooks, S. K.; Webster, R. K.; Smith, L. E.; Woodland, L.; Wessely, S.; Greenberg, N. et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* **2020**, *395*, 912-20. doi: 10.1016/S0140-6736(20)30460-8
70. Riter, H.S; Almeida, M.L; Vescovi, G.; Marques, F.M.; Pedrotti, B.G.; Mallmann, M.Y.; et al. Symptoms of Common Mental Disorders in Brazilian Parents During the COVID-19 Pandemic: Associated Factors. *Psychol Stud.* **2021**, 1-10, doi: 10.1007/s12646-021-00609-8.

71. Sousa, A.R.; Alves, G.V.; Queiroz, A.M.; Florêncio, R.M.S.; Moreira, W.C.; Nóbrega, M.D.P.S.S.; Teixeira, E.; Rezende, M.F. Men's mental health in the COVID-19 pandemic: Is there a mobilization of masculinities? *Rev. Bras. Enferm.* **2021**, *74* (Suppl. S1), e20200915, doi: 10.1590/0034-7167-2020-0915
72. Angehrn, A.; Krakauer, R.L.; Carleton, R.N.; The Impact of Intolerance of Uncertainty and Anxiety Sensitivity on Mental Health Among Public Safety Personnel: When the Uncertain is Unavoidable *Int. J. Cogn. Ther* **2020**; *44*, 919-930, doi: 10.1007/s10608-020-10107-2.
73. Giusti, L.; Salza, A.; Mammarella, S.; Bianco, D.; Ussorio, D.; Casacchia, M.; Roncone, R. #Everything Will Be Fine. Duration of Home Confinement and "All-or-Nothing" Cognitive Thinking Style as Predictors of Traumatic Distress in Young University Students on a Digital Platform During the COVID-19 Italian Lockdown. *Front. Psychiatry* **2020**, *11*, 574812, doi: 10.3389/fpsyt.2020.574812
74. Rehman, S.U.; Ullah, R.; Imran, M.; Iqbal, K.; Saleem, Z.; Khattak, S.R. Intolerance of Uncertainty and Mental Well-being: Important Role of Loneliness and Fear of COVID-19: Evidence from Higher Educational Institutions. *Multicultural Education* **2021**, *7*, doi: 10.5281/zenodo.4750645
75. Moreira, W.C.; Nóbrega, M.P.S.S.; Lima, F.P.S.; Lago, E.C.; Lima, M.O. Effects of the association between spirituality, religiosity and physical activity on health/mental health: a systematic review. *Rev Esc Enferm USP.* **2020**, *54*:e03631. doi: <https://doi.org/10.1590/S1980-220X2019012903631>
76. Queiroz, A.M.; Sousa, A.R.; Moreira, W.C.; Nóbrega, M.P.S.S.; Santos, M.B.; Barbosa, L.J.; et al. The novel COVID-19: impacts on nursing professionals' mental health?. *Acta Paul Enferm.* **2021**, *34*:eAPE02523. Doi: <https://doi.org/10.37689/acta-ape/2021AO02523>.
77. Nasi, C.; Marcheti, P.M.; Oliveira, E.; Rezio, L.A.; Zerbetto, S.R.; Queiroz, A.M.; et al. Meanings of nursing professionals' experiences in the context of the pandemic of COVID-19. *Rev Rene.* **2021**, *22*:e67933. DOI: <https://doi.org/10.15253/2175-6783.20212267933>
78. Nóbrega, M.P.S.S.; et al. Nurses' attitudes facing the family involvement in caring for people with mental disorder. *Rev Bras Enferm.* **2020**, *73*, Suppl 1, e20200041. Doi: <https://doi.org/10.1590/0034-7167-2020-0041>.

79. Alves, A.A.M.; Rodrigues, N.F.R. Determinantes sociais e econômicos da Saúde Mental. *Rev Port Saúde Pública* **2010**, 28, 127-31. doi: [https://doi.org/10.1016/S0870-9025\(10\)70003-1](https://doi.org/10.1016/S0870-9025(10)70003-1)
80. Sousa, A.R.; Queiroz, A.M.; Florentino, R.M.S.; Portela, P.P.; Fernandes, J.D.; Pereira, A. Homens nos serviços de atenção básica à saúde: repercussões da construção social das masculinidades. *Rev. baiana enferm* **2016**, 30, 1-10, doi: 10.18471/rbe.v30i3.16054
81. Sousa, A.R.; Santana, T.S.; Carvalho, E.S.S.; Mendes, I.A.C.; Santos, M.B.; Reis, J.L.; et al. Vulnerabilities perceived by men in the context of the Covid-19 pandemic. *Rev Rene*. **2021**, 22:e60296. DOI: <https://doi.org/10.15253/2175-6783.20212260296>
82. Sousa, A.R.; Araújo, I.F.M.; Borges, C.C.L.; Oliveira, J.A.; Almeida, M.S.; Caribé, W.; et al. Men's health in the covid-19 pandemic: brazilian panorama. *Rev baiana enferm*. **2021**;35:e38683. DOI 10.18471/rbe.v35.38683
83. Botton, A., Cúnico, SD, Strey, MN. Diferenças de gênero no acesso aos serviços de saúde: Problematizações obrigatórias. *Mudanças - Psicologia da Saúde*, **2017**, 25, 67-72. doi: 10.15603/2176-1019/mud.v25n1p67-72
84. Sousa, A.R.; et al. Sentidos e significados atribuídos por homens ao vivido na pandemia da Covid-19. *Rev Esc Enferm USP* **2021**, 55, e03763. Doi: <https://doi.org/10.1590/S1980-220X2020046403763>.
85. Moreira, W. C.; Sousa, K. H. J. F.; Sousa, A. R.; Santana, T. S.; Zeitoune, R. C. G.; Nóbrega, M. P. S. S. Mental health interventions implemented in the COVID-19 pandemic: what is the evidence?. *Rev. Bras. Enferm* **2021**,74, e20200635, English, Portuguese. doi: 10.1590/0034-7167-2020-0635.
86. Romiti, G.F.; Talerico, G. Embracing the Uncertainty: an Important Lesson from COVID-19. *J Gen Intern Med*. **2021**,1-3. doi: 10.1007/s11606-021-06809-2
87. Vasconcellos-Silva, P. R.; Castiel, L. D. COVID-19, as fake news e o sono da razão comunicativa gerando monstros: a narrativa dos riscos e os riscos das narrativas. *Cad Saúde Pública* **2020**, 36, e00101920, doi: 10.1590/0102-311X00101920>.
88. Sousa ÁFL, Teixeira JRB, Lua I, Oliveira Souza F, Ferreira AJF, Schneider G, et al. Determinants of COVID-19 Vaccine Hesitancy in Portuguese-Speaking Countries: A Structural Equations Modeling Approach. *Vaccines* (Basel). **2021** Oct 12;9(10):1167. doi: 10.3390/vaccines9101167. PMID: 34696275; PMCID: PMC8541102.