

Article

Bioaccessibility of antioxidants in blackcurrant juice after treatment using supercritical carbon dioxide

Urszula Trych^{1,*}, Magdalena Buniowska², Sylwia Skapska¹, Ireneusz Kapusta³ and Krystian Marszałek^{1,3}

¹ Prof. Wacław Dąbrowski Institute of Agricultural and Food Biotechnology – State Research Institute, Department of Fruit and Vegetable Product Technology, 36 Rakowiecka St., 02532, Warsaw, Poland; urszula.trych@ibprs.pl, sylwia.skapska@ibprs.pl, krystian.marszalek@ibprs.pl

² University of Rzeszów, Department of Dairy Technology, Institute of Food Technology and Nutrition, Ćwiklińskiej 2D St., 35601 Rzeszów, Poland; mbuniowska@ur.edu.pl

³ University of Rzeszów, Institute of Food Technology and Nutrition, College of Natural Science, Department of Food Technology and Human Nutrition, 2D Zelwerowicza St., 35601, Rzeszów, Poland; ikapusta@ur.edu.pl

* Correspondence: urszula.trych@ibprs.pl

Abstract: Blackcurrant juice (*Ribes nigrum* L.) was subjected to supercritical carbon dioxide (SCCD) at 10, 30 and 60 MPa for 10 min at 45°C as well as thermally treated at 45°C and 85°C for 10 min to determine the stability, antioxidant capacity (AC) and bioaccessibility (BAc) of vitamin C, total anthocyanins and their individual monomers. An *in vitro* gastrointestinal digestion model completed with dialysis was used to assess BAc. The use of SCCD at each of the pressures applied improved the stability of vitamin C, total anthocyanins, and AC before *in vitro* digestion. As a result of digestion, L-ascorbic acid was oxidized to L-dehydroascorbic acid, and finally, the total content of vitamin C, anthocyanins, and AC decreased. SCCD did not significantly improve the BAc of vitamin C and total anthocyanins. The highest BAc of vitamin C was noted in fresh juice (FJ) (40%) and after mild heat treatment at 45°C (T45) (46%). The highest BAc of total anthocyanins was also noted in the FJ (4.4%). The positive effect of the application of SCCD on the BAc of the delphinidin-3-O-glycosides was observed compared to T45 and thermal pasteurization at 85°C (T85). Moreover, cyanidins were generally more bioaccessible than delphinidins in all samples. AC after digestion was higher in SCCD samples compared to thermally treated measured using ABTS+• and DPPH• assays, whereas in dialysate similar trends were observed only for AC measured using the ABTS+• assay. This phenomenon was justified by the formation of individual metabolites detected by UPLC-PDA-MS / MS in the model experiment with delphinidin-3-O-rutinoside. The protocatechuic acid which is well known as a strong antioxidant was detected in the model experiment after digestion. Further research is needed to better understand the metabolic pathway of anthocyanins and the possible uses of SCCD to improve the health properties of fruit products.

Keywords: bioaccessibility, supercritical carbon dioxide, blackcurrant, vitamin C, anthocyanins, antioxidant activity, *in vitro* digestion model

1. Introduction

Growing awareness of the impact of proper nutrition on the health and quality of human life is one of the most important topics for food technologists, especially with regard to the application of emerging techniques for food preservation. Modern food preservation techniques, such as treatment with supercritical carbon dioxide (SCCD) might be a good alternative to thermal heating since it allows maintaining a high antioxidant content while ensuring food safety and high sensorial properties [1–4]. Supercritical CO₂ is a non-toxic, inexpensive, readily available, non-flammable medium that is approved as safe. It is characterized by a low viscosity (3–7×10⁻⁵ Pa s) and zero surface tension, thanks to which it is able to quickly and effectively penetrate the sample [4]. SCCD carried out at a pressure ranging from 7.28 MPa up to 60 MPa allowed for the effective reduction of

microorganisms, even by more than 3 logarithmic cycles. The use of the parameters of 60 MPa, 30 minutes and 45°C made it possible to obtain strawberry juice while maintaining good microbiological quality even for 12 weeks of storage at 6°C [2]. SCCD also enables the inactivation of tissue enzymes such as polyphenol oxidase or peroxidase, thereby inhibiting the browning processes [2,3,5,6]. Penetration of a food sample with supercritical carbon dioxide causes a decrease in the pH inside the cell, physical disruption of the tissues, modification of cell membranes, and the extraction of intracellular compounds. In fruit and vegetable processing SCCD is used primarily in liquid foods such as juices, where processing fruit caused significant tissue damage before preservation [7].

Vitamin C and anthocyanins are well known for their anti-inflammatory, antibacterial and neuroprotective properties. The high content of these antioxidants in blackcurrants makes this fruit a recommended dietary component in the prevention of cancer, cardiovascular diseases, and vision defects [8–12]. Studies in mouse models confirmed the possibility of anthocyanin extracts, obtained from Kenyan purple tea, crossing the blood-brain barrier. Due to their neuroprotective properties, they are promising dietary supplements for inhibiting oxidative stress in the brain and preventing neurodegenerative diseases such as Alzheimer's, Parkinson's, and multiple sclerosis [13].

There are indications that anthocyanins may have a positive effect on the body even though their absorption by the intestinal epithelium is very limited. The metabolism and absorption of these substances are complicated and not fully understood. However, to take full advantage of the health benefits that can be associated with the consumption of antioxidant-rich fruits, it is important to ensure the highest possible bioavailability and bioaccessibility of these compounds so that they can reach the target tissues and organs in the human body [12,14]. Bioaccessibility is defined as the amount of an active compound that is released from the matrix and is available for absorption into the bloodstream [15,16]. The bioaccessibility study was performed using an *in vitro* model that simulated digestion in the mouth, stomach and small intestine along with dialysis through a cellulose membrane simulating the passive transport of components through the intestinal epithelium into the bloodstream.

Technological processing of food is one of the factors influencing the bioaccessibility of nutrients and bioactive compounds. The aim of the present research was to investigate whether SCCD also affects the stability and bioaccessibility of bioactive ingredients such as vitamin C and anthocyanins in blackcurrant juice after digestion in an *in vitro* gastrointestinal model. Moreover the metabolites of selected anthocyanins after *in vitro* digestion and dialysis were detected using UPLC-PDA-MS/MS.

2. Results and Discussion

2.1. Stability and bioaccessibility of vitamin C after SCCD treatment

According to the data presented in Table 1, the vitamin C content in the control samples of blackcurrant juice ranged from 150 mg/L in untreated juice to 172.54 mg/L in juice treated using SCCD at 60 MPa. SCCD improved the stability and extractability of the vitamin C from the tissue to the juice. Thermal treatment did not significantly affect the vitamin C content. The vitamin C content in blackcurrant juice was similar to the values obtained in other studies, which ranged from 60 to 280 mg/L depending on the country of origin of the fruit [17,18]. Marszałek et al. (2018) reported no effect of SCCD treatment under pressure of 10–60 MPa on the vitamin C content in apple juice [3]. The higher sensitivity of vitamin C was observed in strawberry juice, where about 30% of the vitamin C was degraded after SCCD treatment under a pressure of 30 and 60 MPa [1]. In the studies conducted on orange juice, SCCD caused a 12% reduction in vitamin C compared to the control samples, but a 43% decrease in this vitamin after pasteurization was also noted. Vitamin C is susceptible to degradation in the presence of oxygen and active oxidative enzymes. In turn, supercritical CO₂ is a poor solvent for the extraction of this vitamin and does not contribute to its degradation. The replacement of the oxygen dissolved in the liquid juice matrix with supercritical CO₂ could be beneficial for the better stability of

vitamin C in products [19,20]. It transpires that the vitamin C contained in blackcurrant juice is resistant to the application of temperature. A low pH of the environment could have a protective effect on the chemical structure of vitamin C [21,22].

98% of vitamin C in the control samples was present in the L-ascorbic acid (AA) form. Due to digestion, it underwent reversible oxidation to L-dehydroascorbic acid (DHAA). During the simulated digestion in the small intestine, DHAA became the dominant form of vitamin C, reaching 76–93%. In the dialysate, the DHAA content again fell slightly, ranging from 29% in the FJ samples to 54% in the T45 samples. In the conditions prevailing in the digestive system, DHAA could also undergo further decomposition due to diketogulonic acid, 2-hydroxyfurfural, and other 5-carbon compounds [22].

The total vitamin C content in all samples was higher than the control as a result of digestion at the mouth stage and then decreased after gastric digestion. Due to intestinal digestion and dialyse, the total vitamin C content decreased by 45–66% compared to the stomach stage. In our previous studies, we observed a decrease in the vitamin C content at the stage of oral digestion, and then, an increase in vitamin C concentration in the acidic environment of the stomach, which could have been caused by using a different matrix (puree). After intestinal digestion and dialysis, the vitamin C was significantly degraded (~98%). Studies conducted on blended fruit juice made from orange, pineapple, and kiwi showed high vitamin C stability after gastric digestion (recovery >75%) and a significant decrease in vitamin C stability in an alkaline pH which occurred in the intestinal step of digestion [21]. During the study of the bioaccessibility (BAc) of bioactive ingredients, including vitamin C from broccoli inflorescences, only a 7% loss of vitamin C after digestion in gastric conditions was noted, but as much as a 91% degradation in intestinal conditions [23]. Similarly, in the studies of pomegranate juice, a 95% loss of vitamin C after intestinal digestion was reported [22]. Vitamin C is commonly known to be an unstable bioactive compound sensitive to changes in pH (pH>4), high temperature, enzyme activity, oxygen, light, and some metal ions [24].

It can also be observed that most of the vitamin C remaining after the simulation of intestinal digestion was passively transported across a cellulose membrane simulating the epithelial barrier of the small intestine. The most effective diffusion took place in samples treated using SCCD60 (87%), followed by the T45 sample (81%). Other processing conditions resulted in an absorption level of vitamin C ranging from 76 to 78%. The absorption of vitamin C in the human organism differs depending on the type of metabolite. Riboflavin and ascorbic acid are absorbed via a sodium-dependent active transporter (SVCT1) in the upper part of the small intestine. Dehydroascorbic acid can be absorbed thanks to glucose transporters and facilitated diffusion in the duodenum and jejunum. It is also worth mentioning that age, habits, medications taken, and physiological characteristics affect the high variability of the actual BAc of vitamin C ingested by the human alimentary tract [22,25].

The highest BAc of total vitamin C was noted for FJ (40%) and T45 (46%) (Figure 1). BAc after T85 and SCCD treatment was about 25–28%. SCCD did not improve the BAc of vitamin C. The effect of a lower pressure level in the SCCD treatment was insignificant compared to other pressure parameters and pasteurization. To the best of our knowledge, there are no studies on the BAc of vitamin C in samples treated using SCCD. In our previous research, the influence of High Hydrostatic Pressure on the BAc of antioxidants in blackcurrant puree was indicated. We reported that the BAc of vitamin C in dialysate was unfortunately very low and reached only 1% in samples treated using HHP in 400 and 600 MPa. Due to the possibility of the absorption of vitamin C just after gastric digestion, BAc after that stage was considered and ranged between 60% in pasteurized samples and 90% in HHP 400 and 600 MPa [24]. The differences in the bioaccessibility obtained in our studies could largely be due to using a different food matrix, which was also noted by other researchers. For instance, polyphenols contained in wine were more readily bioaccessible than those derived from grapes. In the case of solid food matrices, digestion at the mouth and stomach stages plays a very important role. The process of mastication in an

appropriate pH and the presence of digestive enzymes, provides additional extraction of bioactive compounds and is essential for their release and availability for absorption [26].

In other studies, the BAc of vitamin C varied depending on the type and composition of the food matrix and the method of assessing the absorption of substances into the bloodstream (centrifugation or dialysis). Rodríguez-Roque et al. [21] who used a dialysis membrane in their studies, reported a BAc level of vitamin C from blended fruit juice of 15%. In other studies, also using a cellulose membrane, the BAc of vitamin C was obtained at a lower level of approx. 2.5% (in pomegranate juice) and 3.2% (in broccoli inflorescences) [22,23]. Rodríguez-Roque et al., (2015) reported a positive effect of HHP (400 MPa, 5 min, 40°C) on the BAc of vitamin C in juices with the addition of water, cow's milk or soy milk compared to thermal treatment (90°C, 60 s). BAc of the vitamin C in control samples of beverages and after different processing such as: high intensity pulsed electric fields, high pressure processing, thermal pasteurization, were determined from 11 to 23% [27]. Among the studies using the centrifugation technique to assess BAc, Cilla et al., (2012) noted large differences in the BAc of vitamin C depending on the addition of a protein component to the mixed fruit juice and the processing technique. HHP treatment (400 MPa, 40°C, 5 min) did not improve the BAc of vitamin C compared to short thermal treatment (90°C, 30 s). The BAc of vitamin C reached about 70% when whole cow's milk was added to the control samples, which was significantly higher compared to the samples with the addition of soy milk (BAc=12.6%). It has been assumed that HHP can promote the formation of whey protein aggregates better than thermal treatment and, therefore, can reduce the BAc of vitamin C to a greater extent [28]. *In vivo* studies on humans have shown that the consumption of vitamin C-rich foods, such as vegetable 'gazpacho' soup treated with non-thermal processing techniques, increased the concentration of this vitamin in the plasma and reduced oxidative stress [29].

Table 1. L-ascorbic acid (AA), L-dehydroascorbic acid (DHAA) and total vitamin C content at each stage of the digestion of black-currant juice. FJ- fresh juice; T45- thermal treated juice at 45°C; T85- thermal treated juice at 85°C; SCCD10- juice after supercritical carbon dioxide treatment at 10 MPa; SCCD30- juice after supercritical carbon dioxide treatment at 30 MPa; SCCD60- juice after supercritical carbon dioxide treatment at 60 MPa

	Sample	AA (mg/100 ml)	DHAA (mg/100 ml)	Total vitamin C (AA+DHAA) (mg/100 ml)
control	FJ	147.71 ^c ± 11.04	2.29 ^c ± 0.71	150.00 ^c ± 11.08
	T45	155.09 ^{bc} ± 3.56	3.17 ^{ab} ± 0.91	158.25 ^{bc} ± 3.14
	T85	155.81 ^{bc} ± 2.82	1.95 ^c ± 0.65	157.76 ^{bc} ± 2.60
	SCCD10	162.68 ^{ab} ± 8.80	3.23 ^{ab} ± 1.21	165.91 ^{ab} ± 9.36
	SCCD30	156.93 ^{bc} ± 5.60	3.84 ^a ± 0.38	160.77 ^{bc} ± 5.82
	SCCD60	169.39 ^a ± 2.66	3.15 ^{ab} ± 0.92	172.54 ^a ± 2.25
oral	FJ	188.72 ^{bc} ± 15.05	9.10 ^a ± 1.64	197.82 ^{ab} ± 16.59
	T45	201.55 ^{ab} ± 11.03	1.83 ^c ± 0.49	203.38 ^{ab} ± 11.44
	T85	180.99 ^c ± 12.66	6.96 ^b ± 1.56	187.95 ^b ± 13.71
	SCCD10	210.91 ^a ± 4.06	2.35 ^c ± 0.83	213.26 ^a ± 3.66
	SCCD30	201.71 ^{ab} ± 4.87	8.69 ^{ab} ± 1.26	210.39 ^a ± 4.95
	SCCD60	196.97 ^{abc} ± 3.59	7.93 ^{ab} ± 0.48	204.90 ^{ab} ± 3.12
gastric	FJ	170.03 ^a ± 14.71	2.73 ^c ± 0.51	172.77 ^a ± 14.42
	T45	161.06 ^{ab} ± 3.57	2.40 ^c ± 0.77	163.46 ^{ab} ± 3.73
	T85	151.76 ^b ± 9.45	8.19 ^{ab} ± 5.27	159.96 ^{ab} ± 9.56
	SCCD10	151.88 ^b ± 11.08	11.65 ^a ± 2.31	163.53 ^{ab} ± 9.43
	SCCD30	131.45 ^c ± 12.82	4.08 ^{bc} ± 1.09	135.52 ^c ± 12.79
	SCCD60	152.94 ^{ab} ± 5.37	1.87 ^c ± 0.62	154.81 ^b ± 5.49
intestina I	FJ	4.25 ^a ± 0.93	13.83 ^{ab} ± 2.18	18.08 ^a ± 2.93
	T45	2.19 ^{bc} ± 0.70	14.81 ^a ± 3.15	16.99 ^{ab} ± 3.67
	T85	1.50 ^{cd} ± 0.17	12.20 ^{ab} ± 2.45	13.70 ^{bc} ± 2.61

	SCCD10	2.53 ^b ± 0.36	10.79 ^b ± 1.72	13.32 ^{bc} ± 1.36
	SCCD30	1.06 ^{de} ± 0.36	10.94 ^b ± 1.12	12.00 ^c ± 0.72
	SCCD60	0.46 ^e ± 0.06	6.28 ^c ± 0.67	6.74 ^d ± 0.67
dialysate	FJ	42.61 ^a ± 4.33	17.26 ^b ± 1.69	59.87 ^b ± 5.87
	T45	33.61 ^b ± 2.50	38.75 ^a ± 5.30	72.37 ^a ± 3.09
	T85	23.04 ^c ± 1.21	20.17 ^c ± 3.36	43.21 ^c ± 4.14
	SCCD10	30.22 ^b ± 2.89	16.04 ^c ± 1.21	46.26 ^c ± 3.79
	SCCD30	21.94 ^c ± 3.47	18.69 ^c ± 2.21	40.63 ^c ± 1.84
	SCCD60	21.43 ^c ± 1.79	23.86 ^c ± 3.59	45.28 ^c ± 2.52

The same letter indicate no significant difference between mean ($p \leq 0.05$) in the columns within each digestion stage.




Figure 1. Bioaccessibility of total vitamin C in the dialysate after different processing: FJ- fresh juice; T45- thermal treated juice at 45°C; T85- thermal treated juice at 85°C; SCCD10- juice after supercritical carbon dioxide treatment at 10 MPa; SCCD30- juice after supercritical carbon dioxide treatment at 30 MPa; SCCD60- juice after supercritical carbon dioxide treatment at 60 MPa (The same letter indicates no significant difference between the mean ($p \leq 0.05$) in the columns within each digestion stage.)

2.2. Stability and bioaccessibility of anthocyanins and determination of delphinidin metabolites after *in vitro* gastro-intestinal digestion

The total anthocyanin content in blackcurrant juice in the control samples ranged from 1 173 mg/L in the SCCD30 samples to 498 mg/L in the T85 samples (Table 2). Other authors indicated the total anthocyanin content in blackcurrants to be between 160–411 mg/100 g of fresh fruit [30,31]. In our previous study, the total anthocyanin content in blackcurrant was determined as 737 mg/kg of raw material [24]. The lower anthocyanin content in the raw material compared to juice may be related to the higher concentration of pectin in the fruits, which are able to bind anthocyanins in plant tissue. To obtain the highest possible anthocyanins concentration the mash was enzymatically treated with pectinolytic enzymes before pressing.

The SCCD treatment of the juice positively influenced the retention and extractability of anthocyanins, which, regardless of the pressure applied, were higher than in the fresh juice. The best effect was obtained using SCCD treatment at 30 and 60 MPa. On the other hand, pasteurization at 85°C definitely contributed to the degradation of anthocyanins. Thermal treatment at 45°C did not have a significant effect on the total anthocyanin content. Four major anthocyanins were determined in the blackcurrant juice, in descending

order of concentration: delphinidin-3-*O*-rutinoside (df-3-*O*-rut), cyanidin-3-*O*-rutinoside (cy-3-*O*-rut), delphinidin-3-*O*-glucoside (df-3-*O*-glu), and cyanidin-3-*O*-glucoside (cy-3-*O*-glu). Other studies confirmed that SCCD treatment allows maintaining the high stability of anthocyanins derived from plants [7], thanks to the inactivation of the polyphenol oxidases and reduction in the activity of peroxidases [32]. Processing using the SCCD method may result in the occurrence of reversible changes in the structure of proteins, such as enzymes, and inactivate them [32]. As a result of the SCCD treatment (30 and 60 MPa, 45°C, 30 min) on strawberry juice, anthocyanin degradation following first-order kinetics was observed. The processing temperature and time were the most important for preservation conditions followed by pressure [1]. In the other studies, SCCD treatment at 23 MPa decreased the anthocyanin content compared to the control samples of blood orange juice by about 8% which did not differ significantly from the values obtained for the thermal treated samples (88–91°C, 30 s). On the other hand, the application of 13 MPa showed a slight loss of anthocyanins compared to the control samples, and the stability of the anthocyanins was higher than in the pasteurized samples [20].

Table 2. Total anthocyanin content at each stage of digestion of blackcurrant juice. FJ- fresh juice; T45- thermal treated juice at 45°C; T85- thermal treated juice at 85°C; SCCD10- juice after supercritical carbon dioxide treatment at 10 MPa; SCCD30- juice after supercritical carbon dioxide treatment at 30 MPa; SCCD60- juice after supercritical carbon dioxide treatment at 60 MPa

	Sample	Df-3- <i>O</i> -glu (mg/L)	Df-3- <i>O</i> -rut (mg/L)	Cy-3- <i>O</i> -glu (mg/L)	Cy-3- <i>O</i> -rut (mg/L)	Total anthocyanins (mg/L)
control	FJ	111.13 ^c ± 5.85	371.91 ^d ± 18.76	45.75 ^c ± 2.18	259.66 ^c ± 12.28	788.45 ^c ± 39.07
	T45	101.36 ^c ± 9.04	342.26 ^d ± 30.38	41.45 ^c ± 3.54	236.86 ^c ± 20.05	721.93 ^c ± 62.99
	T85	73.62 ^d ± 4.14	236.99 ^e ± 2.42	28.53 ^d ± 1.94	159.28 ^d ± 2.95	498.43 ^d ± 11.27
	SCCD10	138.80 ^b ± 19.38	460.60 ^c ± 31.44	56.08 ^b ± 7.35	306.72 ^b ± 27.17	962.20 ^b ± 25.31
	SCCD30	159.95 ^a ± 8.23	579.44 ^a ± 34.45	71.55 ^a ± 6.19	383.67 ^a ± 28.74	1173.02 ^a ± 64.69
	SCCD60	159.90 ^a ± 4.85	521.73 ^b ± 11.37	62.77 ^b ± 1.90	334.94 ^b ± 7.45	1079.34 ^a ± 25.30
oral	FJ	96.14 ^a ± 11.09	314.95 ^a ± 19.52	34.95 ^a ± 6.09	186.23 ^a ± 10.10	632.28 ^a ± 45.36
	T45	85.24 ^a ± 21.99	289.64 ^a ± 27.35	31.98 ^a ± 2.91	162.13 ^{ab} ± 18.40	568.99 ^a ± 59.44
	T85	44.97 ^b ± 13.15	153.49 ^b ± 16.45	17.25 ^b ± 5.37	105.52 ^c ± 22.08	321.23 ^b ± 27.05
	SCCD10	48.74 ^b ± 2.42	165.39 ^b ± 6.50	18.55 ^b ± 0.65	111.90 ^c ± 4.34	344.58 ^b ± 13.21
	SCCD30	49.97 ^b ± 8.19	150.17 ^b ± 2.04	19.05 ^b ± 3.26	108.77 ^c ± 10.14	327.95 ^b ± 20.68
	SCCD60	56.08 ^b ± 7.80	208.56 ^b ± 13.46	21.12 ^b ± 2.95	132.90 ^{bc} ± 9.04	418.66 ^b ± 21.29
gastric	FJ	66.37 ^a ± 6.28	237.17 ^a ± 15.89	27.14 ^a ± 2.56	165.86 ^a ± 6.94	496.55 ^a ± 26.55
	T45	54.24 ^c ± 6.95	238.86 ^a ± 13.54	22.11 ^b ± 2.31	126.41 ^c ± 16.47	441.62 ^b ± 26.36
	T85	60.33 ^{ab} ± 6.08	212.41 ^b ± 10.14	21.86 ^b ± 3.69	165.46 ^a ± 11.30	460.05 ^b ± 19.41
	SCCD10	66.57 ^a ± 1.54	228.10 ^{ab} ± 4.48	25.84 ^a ± 0.56	154.12 ^{ab} ± 3.29	474.62 ^{ab} ± 9.26
	SCCD30	62.22 ^{ab} ± 1.45	213.37 ^b ± 4.27	24.00 ^{ab} ± 0.38	143.22 ^b ± 2.57	442.82 ^b ± 8.53
	SCCD60	62.56 ^{ab} ± 2.98	212.83 ^b ± 7.91	23.97 ^{ab} ± 0.82	142.92 ^b ± 5.06	442.27 ^b ± 16.64
intestinal	FJ	1.88 ^d ± 0.73	11.11 ^c ± 1.88	1.03 ^b ± 0.31	11.53 ^b ± 1.62	25.55 ^c ± 2.30
	T45	4.13 ^a ± 0.49	18.59 ^a ± 1.83	2.27 ^a ± 0.69	17.40 ^a ± 2.55	42.39 ^a ± 5.28
	T85	3.17 ^{bc} ± 0.31	16.11 ^{ab} ± 2.67	1.91 ^a ± 0.29	15.86 ^a ± 2.33	37.04 ^{ab} ± 3.52
	SCCD10	3.40 ^{abc} ± 0.33	16.34 ^{ab} ± 1.16	2.05 ^a ± 0.10	16.91 ^a ± 0.62	38.69 ^{ab} ± 2.04
	SCCD30	3.77 ^{ab} ± 0.32	16.26 ^{ab} ± 3.04	2.03 ^a ± 0.26	16.78 ^a ± 1.80	38.83 ^{ab} ± 4.06
	SCCD60	2.80 ^c ± 0.64	12.88 ^{bc} ± 2.64	1.77 ^a ± 0.19	14.61 ^{ab} ± 1.27	32.05 ^{bc} ± 4.73
dialysate	FJ	3.63 ^a ± 0.79	15.72 ^a ± 0.95	1.93 ^a ± 0.29	13.29 ^a ± 2.11	34.56 ^a ± 3.94
	T45	1.67 ^b ± 0.14	6.53 ^c ± 0.91	1.28 ^b ± 0.14	9.98 ^b ± 0.90	19.46 ^b ± 1.63
	T85	0.96 ^b ± 0.16	4.03 ^d ± 0.93	1.08 ^b ± 0.08	8.21 ^{bc} ± 1.15	14.28 ^c ± 1.21

SCCD10	3.96 ^a ± 0.83	12.51 ^b ± 2.27	1.76 ^a ± 0.22	13.96 ^a ± 2.59	32.19 ^a ± 5.41
SCCD30	0.94 ^b ± 0.06	4.59 ^{cd} ± 0.25	1.04 ^b ± 0.08	8.17 ^{bc} ± 0.40	14.75 ^{bc} ± 0.73
SCCD60	0.96 ^b ± 0.39	4.26 ^d ± 0.18	1.19 ^b ± 0.26	6.40 ^c ± 0.27	12.81 ^c ± 1.04

The same letter indicates no significant difference between the mean ($p \leq 0.05$) in the columns within each digestion stage.

As a result of digestion at the oral stage, the total anthocyanin content decreased in all samples. In the FJ and T45 samples, there was a decrease of 20–21%, in the T85 of 36%, and in the samples treated with SCCD by as much as 61–72%. After digestion in gastric conditions, a further decrease in the total anthocyanins in the FJ and T45 samples of 22% was recorded. In the remaining samples, the anthocyanin content increased in the range of from 6% in SCCD60 to 43% in T85. The total anthocyanin content after simulated gastric digestion was the highest in the FJ samples. Insignificant differences were observed in the remaining samples. The digestion at the small intestine and after dialysis stage decreased the amount of total anthocyanins in all samples by 85–90%. Less anthocyanins were absorbed by a membrane into the dialysate compared to vitamin C. The amount of anthocyanins in the dialysate was higher only in the FJ samples than detected on the other side of the membrane. Finally, insignificant differences were observed between SCCD10 and the FJ sample after dialysis.

Other studies have confirmed a decrease in anthocyanins as a result of simulated digestion. The conditions reflecting those in the stomach were the most favourable for anthocyanin stability. On the other hand, at the small intestine digestion stage, there was a significant decrease in the anthocyanin content [24,26,33]. This phenomenon was explained mainly by changes in pH, which are one of the factors that significantly influence the structural changes of anthocyanins. An alkaline pH contributes to the aglycone ring breakage and formation of insoluble polymeric brown pigments [34]. In the studies conducted on blackcurrant puree, a more pronounced increase in the anthocyanin content was noticed during simulated gastric digestion. In a more complex matrix, containing more plant tissue and pectin than the juice, pepsin digestion and the effect of an acidic pH played a greater role in the extraction of bioactive compounds [24]. Anthocyanins are located in the vacuoles of cells, therefore, after consuming less processed fruit, the release of these compounds occurs during digestion [35].

The mechanisms of anthocyanin absorption in the gastrointestinal tract have not been thoroughly investigated. *In vivo* studies on humans and animals, based on the analysis of anthocyanins concentration in blood and tissues, provide evidence that, apart from the small intestine, they can be partially absorbed in the oral cavity [36], and also in a glycosidic form through the active transporters in the gastric mucosa [37–40]. The microbiota of the large intestine is also of great importance for the metabolism and absorption of anthocyanins [34,40,41].

The BAc of total anthocyanins was the highest in fresh juice (4.4%), then in the T45, T80 and SCCD10 samples (2.7–3.4%), the lowest in SCCD30 and 60 (1.3% and 1.2%) (Figure 2). SCCD did not improve the BAc of total anthocyanins compared to the thermally treated and untreated samples. The most accessible glycoside was cy-3-O-rut, in all types of samples. Cyanidins turned out to be more bioaccessible than delphinidins. Among the SCCD variants, the highest BAc of total anthocyanins was obtained after the application of 10 MPa. The SCCD treatment at 10 MPa improved the BAc of df-3-O-glu and df-3-O-rut compared to the samples subjected to thermal treatment at 45°C and 85°C. Peixoto et al., (2018) reported in a study of anthocyanins derived from blueberries that delphinidin derivatives were degraded by simulation of digestion and were the least bioaccessible glycoside. The greater the number of -OCH₃ groups in anthocyanin, the less susceptible the molecule was to convert or decay to undetectable forms. Consequently, these compounds were more available for absorption [42]. Delphinidin and malvidin derivatives in acetylated form maintained a high resistance to pH changes and lower degradation under digestive conditions. These can be one of the metabolites that are stable in the gastrointestinal tract and therefore enhance the antioxidant status of the blood plasma [34].

Irrespective of the structure of different anthocyanins, the metabolites released after enzymatic and/ or acidic hydrolysis may play important role in human health.

To better understand this phenomenon, UPLC-PDA-MS/MS analysis were used to determine anthocyanin metabolites during digestion in model conditions (Table 3.). The results obtained showed that the major metabolite of df-3-O-rut, the most abundant anthocyanin in blackcurrants, was protocatechuic acid. This compound was present in an amount possible to quantify after the first step of digestion with the use of salivary enzymes, and after simulation of gastric digestion. It was not detected in the intestinal and dialyse stage, possibly due to the low concentration caused by sequential dilutions (Table 4.). Depending on the material tested, different anthocyanin metabolites have been found in the available publications. Our results have been confirmed in some *in vivo* studies on humans [43,44]. Mallery et al. (2011) conducted clinical studies on the effect of oral digestive conditions on metabolism and bioactivation of black raspberry anthocyanins using the LC/MS-MS technique. An analysis of the salivary samples after rinsing the mouth with preparations based on lyophilized black raspberry showed the presence of the parent anthocyanins as well as protocatechuic acid as a functional, stable metabolite, up to 4 hours after administration [43]. Other studies dealt with the *in vivo* metabolism of cyanidin glycosides in blood orange juice (HPLC/MS/MS). 44% of the metabolites detected in the blood plasma collected within 24 hours after juice consumption were protocatechuic acid, a significant amount was also detected in the fecal samples, but not in the urine. It was shown that after digestion and absorption into the body, 73% of the ingested cyanidin glycosides turned into protocatechuic acid. This metabolite may have contributed to the short-term increased antioxidant capacity of blood plasma after the consumption of cyanidin-rich foods [44]. Goszcz et al. (2017) found that delphinidin was a very unstable anthocyanin ($t_{1/2}$ 2–30 min) and the main product of its degradation was gallic acid. The degradation of delphinidin was investigated by LC-MS / MS analysis of solutions incubated for 30 min in phenol red-free tissue culture medium (pH 7.4, temperature 37°C). It was shown that at a concentration of 100 μ M both delphinidins and gallic acid showed cytotoxicity to a cultured cell model of the human umbilical vein endothelium, and had a protective effect on the cells at lower concentrations (100 nM – 1 μ M) [45]. In the *in vivo* studies of the metabolism of berry anthocyanins in humans, it was found that during 48 hours of observation after consuming bilberry-lingonberry purée with and without oat cereals, the phenolic acid content in urine increased. Homovanillic and vanillic acids have been detected as the main metabolites of anthocyanins. As in this study, gallic acid was not found to be a breakdown product of delphinidin glycosides [46]. Other authors reported that hippuric acid was the main metabolite of anthocyanins detected in urine and the tissues of rats on a wild blueberry-enriched diet [47]. Phenolic acids are compounds that present very good antioxidant properties. The metabolites and degradation products catalyzed by the microbiota may be responsible for the health benefits attributed to anthocyanins [40].

Table 3. UPLC-PDA-MS/MS properties of delphinidin 3-O-rutinoside and protocatechuic acid as its major detectable metabolite (condition as in p. 3.4.5.2.).

No	Compound	RT (min.)	[M-H] (m/z)	Fragment ions (m/z)	Absorbance maxima (nm)
1	Delphinidin 3-O-rutinoside	2.71	611 ⁺	303	277, 525
2	3,4-Dihydroxybenzoic acid (protocatechuic acid)	2.14	153 ⁻	109	260, 294

Table 4. The content of Df-3-O-rut and protocatechuic acid at individual stages of the digestion simulation

Stage of digestion	Df-3-O-rut (μ g/ml)	Protocatechuic acid (μ g/ml)
control	49.74 \pm 0.89	0.0

oral	16.25 ± 1.34	0.63 ± 0.02
gastric	52.66 ± 3.25	0.44 ± 0.02
intestinal	3.65 ± 0.02	0.0
Dialysate	0.0	0.0

The bioavailability of anthocyanins in *in vivo* studies is reported to be very low, ranging from 1–2%, as they are often found only in slight amounts in urine, blood and target organs. However, this value may be underestimated as the calculations do not consider all metabolites, activity of the microbiota in the anthocyanin’s degradation, and conjugates that may have an additional effect on the bioavailability of anthocyanins. The proven beneficial effects of consuming anthocyanin-rich foods on many health aspects further indicates the shortcomings of these data [12].


In our previous studies, the BAc of anthocyanins calculated as the ratio of the substance in the dialysate to that in the control samples was very low (maximum 1.6% in fruit crushed in a mortar). No improvement in the BAc of anthocyanins was observed using HHP treatment, irrespective of pressure conditions. However, due to the possibility of absorption of the glycoside form of anthocyanins from the stomach, BAc after this digestive step was also considered. The BAc of total anthocyanins after simulation of gastric digestion was significantly higher in the samples treated with HHP than with thermal pasteurization, reaching the highest value after the application of 600 MPa (68%) [24].

There are studies showing the beneficial effect of the SCCD technique on bioaccessibility when used in the pharmaceutical industry to complex the active substance of a drug onto polymeric carriers such as silica or cyclodextrin. SCCD activation and SCCD-assisted impregnation methods have proved to be promising strategies to obtain better quality medical preparations [48–51]. However, with regard to food-derived substances, only sparse research was found and all of it concerned the effect of SCCD on the BAc of lipophilic compounds such as carotenoids. Zhao et al. [52] investigated the effect of SCCD at a pressure of 10, 20 and 30 MPa, at 55°C for 10, 20, 40 and 60 minutes on the physicochemical properties, isomerization and *in vitro* BAc of lycopene in tomato juice. The effect of the SCCD parameters on phenolics was different, depending on the type of compound detected. The application of 10 MPa did not show a positive effect on the total phenolic compound content compared to the control samples, but it positively influenced the caffeic and chlorogenic acid content compared to the heat-treated samples. Conditions of 20 and 30 MPa had a positive effect and in particular the use of 30 MPa for 40 and 60 minutes had a more positive effect on the phenolic compound content (myricinic acid, ferulic acid, naringin and chlorogenic acid) compared to the control. The BAc of lycopene increased significantly after treatment at 20 MPa for 20, 40, 60 min compared to the control and heat-treated samples. The SCCD samples at 20 MPa for 10 min, and 10 MPa at all pressure-holding times also showed a higher BAc of lycopene than the control and thermally treated samples. Damage to the cell barriers and sample homogenization may be the reason for the improvement in BAc. SCCD supports the release of bioactive ingredients from the food matrix [52]. It was found that SCCD increased the content of *cis*-lycopene isomers which are more accessible than the *trans* isomers. Other researchers noticed that SCCD caused an increase in the volume of the proportion of small particles while there was a decrease in the volume of the proportion of large particles in the settling pulp of orange juice compared to untreated juice, thus having a homogenizing effect on the food matrix. The cause of this process was the induction of high internal stresses causing the breakdown of particles during CO₂ removal from the vessel [53,54]. In the research of Ubeyitogullari (2018), SCCD was used to produce low-crystallinity phytosterol nanoparticles by impregnating phytosterols into nonporous starch aerogels. The use of this technique allowed for a twenty-fold increase in the BAc of phytosterols [55]. In a study of the BAc of lycopene extracted using SCCD from tomato pomace, it was shown that it was about 2.4 times higher compared to hexane extraction. Extracts rich in *cis*-lycopene, obtained using SCCD treatment can be used to design novel functional foods while increasing the use of by-products of the tomato processing industry [56]. SCCD processing may affect the BAc

of polyphenols by increasing the extractability of these compounds under pressure. Moreover, penetration of the sample by CO₂ changes the pH of the environment to more acidic, which may contribute to the hydrolysis of polyphenols to simpler phenolic compounds [3].

Figure 2. Bioaccessibility of blackcurrant juice anthocyanins in the dialysate after different processing: FJ- fresh juice; T45- thermal treated juice at 45°C; T85- thermal treated juice at 85°C; SCCD10- juice after super critical carbon dioxide at 10 MPa; SCCD30- juice after super critical carbon dioxide at 30 MPa; SCCD60- juice after super critical carbon dioxide at 60 MPa. (The same letter indicates no significant difference between the mean ($p \leq 0.05$) in the columns within each digestion stage.)

2.3. Effect of processing on the antioxidant capacity of blackcurrant juice in a simulated digestive system



The antioxidant capacity (AC) in control samples measured using the method with DPPH• radicals ranged from 13.90 to 15.35 mg/L TEAC and was statistically significantly higher in FJ and SCCD treated samples compared to T85 (Figure 3). AC increased after digestion at the oral and stomach stage, and then significantly decreased after digestion in intestinal conditions and in dialysate. Following simulated oral digestion, the AC of FJ, SCCD at 10 and 60 MPa increased significantly, while SCCD at 30 MPa, T45, and T85 statistically had a significantly lower AC. After simulated gastric digestion, the AC of the FJ and T45 did not change significantly, but in the other samples the AC decreased in the range of from 11% in SCCD treated samples at 60 MPa to 27% in SCCD treated at 30 MPa. Due to simulated digestion in intestinal conditions and dialysate, a significant decrease in AC was noticed in all the samples. In the intestinal stage, samples treated using SCCD in all the pressure variants had similar level of AC as FJ and T45 samples and were significantly higher than T85 samples. However, no statistically significant differences were noted between the AC after dialysis.




Figure 3. Antioxidant capacity (AC) of blackcurrant juice determined using the DPPH• assay at each step of digestion after different processing: FJ- fresh juice; T45- thermal treated juice at 45°C; T85- thermal treated juice at 85°C; SCCD10- juice after super critical carbon dioxide at 10 MPa; SCCD30- juice after super critical carbon dioxide at 30 MPa; SCCD60- juice after super critical carbon dioxide at 60 MPa. (The same letter indicates no significant difference between the mean ($p \leq 0.05$) in the columns within each digestion stage.)

Similar trends in the AC changes were confirmed by the results obtained using the ABTS+• assay but the AC was significantly higher in all the SCCD treated samples than in those subjected to thermal treatment and FJ samples (Figure 4). After digestion at the oral stage, there was a noticeable increase of AC in FJ, T45, T85 and SCCD60. The AC changes in SCCD10 and SCCD30 were insignificant. The highest AC after oral digestion was recorded in FJ samples, followed by SCCD, regardless of the type of pressure applied. T85 samples had the lowest AC. The increase in AC in FJ (70%) may have occurred due to the action of salivary enzymes, which contributed to the release of bioactive compounds from plant cells into the juice. Likewise, in other studies, at the earlier stages of digestion, a greater release of bioactive ingredients was noticeable in the non-processed samples. It was only at the intestinal stage that the advantage of the treated samples over the control ones in terms of antioxidant content could be observed [58][59][59]. Simulation of gastric digestion resulted in the unification of the AC to similar values in all the samples (25.4–30.3 mg/L), with no statistically significant differences between them. Similarly to the DPPH• assay, there was a decrease in AC after the simulation of intestinal digestion, however, it was smaller than that observed using ABTS+• (ABTS+•: decrease of 39%–74%; DPPH•: decrease of 83–94%). SCCD samples, in all pressure variants were characterized by significantly higher AC than the samples subjected to thermal treatment and did not differ significantly from the FJ. This trend was maintained in the dialysate, with the exception of the SCCD10 sample, where the lowest AC was observed. The AC in the dialysate ranged from 8.8 to 15.3 mg/L TEAC, and these values were similar to those at the intestinal stage. In contrast, thermal treatment at 45°C did not cause an increase in AC, therefore the positive effect of SCCD treatment can only be attributed only to the application of supercritical carbon dioxide.




Figure 4. Antioxidant capacity (AC) using the ABTS+• assay at each step of digestion after different processing: FJ- fresh juice; T45- thermal treated juice at 45°C; T85- thermal treated juice at 85°C; SCCD10- juice after super critical carbon dioxide at 10 MPa; SCCD30- juice after super critical carbon dioxide at 30 MPa; SCCD60- juice after super critical carbon dioxide at 60 MPa. (The same letter indicates no significant difference between the mean ($p \leq 0.05$) in the columns within each digestion stage.)

The decrease in AC at each steps of digestive tract was significant but much lower in comparison to the anthocyanins degradation. This phenomenon can be justified by the high AC of anthocyanins metabolites, which concentration increased after digestion. The calculated coefficient expressed as antioxidant capacity per 1 mg/L of anthocyanins in blackcurrant juice (Figure 5) indicate that the AC increase significantly in relation to 1 mg of anthocyanins per liter. Considering the calculated coefficient in the dialysate step of digestion, the AC of the SCCD60 sample was three-fold higher, and the SCCD30 sample was two-fold higher than the control's. This phenomenon may indicate the influence of SCCD on improvement the accessibility of antioxidants for digestion, thanks to which more metabolites were released and they could influence the AC of the fruit juice.




Figure 5. Coefficient of the ratio of increase the antioxidant capacity (AC) in relation to the total anthocyanins content in blackcurrant juice after different processing: FJ- fresh juice; T45- thermal treated juice at 45°C; T85- thermal treated juice at 85°C; SCCD10- juice after super critical carbon dioxide at 10 MPa; SCCD30- juice after super critical carbon dioxide at 30 MPa; SCCD60- juice after super critical carbon dioxide at 60 MPa.

Other author studies indicate the effectiveness of the SCCD method in maintaining the high antioxidant capacity of food products [58–61]. In the study of red grapefruit juice after SCCD treatment, at several pressure variants and times, no significant differences were detected in the total phenol content, ascorbic acid content and AC compared to untreated juice [58]. Similarly, an examination of a hibiscus infusion treated using SCCD (34.5 MPa, 8% CO₂, 6.5 min and 40°C) indicated a slight loss of anthocyanins (9%) and no significant changes in phenolic compounds or AC during 14 days of storage [59]. The AC of apple juices subjected to SCCD at 35°C, 15 min and pressure of 15 or 25 MPa turned out to be significantly higher with the use of higher pressure. Moreover, both pressures applied improved the AC compared to the fresh samples. As mentioned before, SCCD can promote conformational changes in the secondary structure of proteins and the inactivation of tissue enzymes [60]. The lychee juice treated with SCCD (8 MPa, 36°C, 120 s) also showed higher AC measured using FRAP and ABTS+• assays, as well as more total phenols and flavonoids compared to thermal treated juice (100°C, 60 s), and UHT (134°C, 4 s). Treatment with SCCD supported preservation of the bioactive compounds due to the exclusion of oxygen and a mild temperature [61].

In our previous studies AC measured using ABTS+• radicals also increased after oral digestion and additionally the DPPH• and ABTS+• assays were in line with the improvement in AC after gastric digestion and the lowering of AC in intestinal conditions [62]. The effect of simulated gastrointestinal digestion on the content of phenolic compounds and their antioxidant capacity in wild Chilean currants was investigated. As a result of digestion, there was an approx. 50% decrease in the content of total phenols and flavonoids, and approx. 80% in anthocyanins and hydroxycinnamic acids, as well as a decrease in antioxidant activity at each stage of digestion, correlated with the loss of bioactive compounds [63]. In studies conducted on strawberry juices enriched with inulin and oligofructose, treated with ultrasound, simulated gastrointestinal digestion resulted in a decrease

in the content of phenolic compounds, flavonoids and AC at each subsequent stage of the process. However, at the stage of the small intestine, higher AC (TEAC) and the content of the aforementioned bioactive compounds were noticed in the ultrasonically treated samples compared to the unprocessed ones [58].

The metabolites formed during the digestion of vitamin C and anthocyanins contribute remarkably to the antioxidant properties presented by the food products. For instance, protocatechuic acid detected in the samples in model conditions was the main metabolite of delphinidin-3-*O*-rutinoside. After the consumption of blood orange juice, as a source of anthocyanins, by six healthy subjects, high plasma concentrations of protocatechuic acid were noted, which contributed to an increase in short-term antioxidant activity [44]. Protocatechuic acid is one of the many metabolites that can be formed as a result of anthocyanin digestion in the gastrointestinal tract. The pathway of its formation from delphinidin-3-*O*-rutinoside is shown in Figure 6, according to available publications [7,41,64].




Figure 6. Proposed pathway for the formation of protocatechuic acid from delphinidin-3-*O*-rutinoside in the gastrointestinal tract.

Bouayed et al., (2011) showed that the AC (FRAP and ABTS+•) of antioxidants in apples after digestion with dialysis was about 50% lower than the values originally contained in the pre-digested samples [33]. According to available studies, AC is correlated with the antioxidant content at a given stage of digestion. In addition, AC is influenced by environmental conditions such as pH, the interactions of antioxidants with other matrix components (iron and other mineral ions, dietary fibre, proteins), and the chemical structure of the compounds. The free radical scavenging activity of polyphenols depends on the number and position of the hydrogen donating hydroxyl groups on the aromatic rings of the molecules. Therefore, aglycones indicate a higher AC than their glycosides [33].

It has also been found that due to its low stability, delphinidin does not have a significant influence on AC, but its metabolites may be responsible for strong antioxidant properties. The parent compound is not necessary to obtain high AC. This explains the paradox of the low bioavailability of polyphenols, including anthocyanins, while exerting strong antioxidant effects and health benefits [66].

3. Materials and Methods

3.1. Reagents and solvents

Sigma-Aldrich (St. Louis, MO, USA) was the supplier of the dialysis tubing cellulose membrane (avg. flat width 25 mm) and most of the enzymes and reagents, as follows: mucin from the porcine stomach – type II, α -amylase, heat-stable, (TDF-100A, 24975 U/ml), pepsin from the porcine gastric mucosa (250 U/mg solid), pancreatin from the porcine pancreas (8 x USP specifications), porcine bile extract, sodium dodecyl sulfate – ACS reagent, sodium bicarbonate $\geq 99.5\%$, 2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic) acid, diammonium salt (ABTS+• radical), 2,2-diphenyl-1-picrylhydrazyl (DPPH• radical), (\pm)-6-hydroxy-2,5,7,8-tetramethyl-chromane-2-carboxylic acid (Trolox), DL-dithiothreitol (HPLC) (DTT), phosphoric acid 85%, acetonitrile (HPLC), formic acid $\geq 95.0\%$ and sodium hydroxide pellets $\geq 98.0\%$, (NaOH).

Other reagents were obtained from Chempur (Piekary Śląskie, Poland), such as: di-sodium hydrogen phosphate anhydrous pure p.a. $\geq 99.0\%$ (Na_2HPO_4), di-potassium hydrogen phosphate (K_2HPO_4), sodium chloride pure p.a. $\geq 99.9\%$ (NaCl) and di-sodium edetate standard solution 0,01 mol/L (EDTA). Hydrochloric acid pure p.a. ACS reagent 37% (HCl) and potassium peroxodisulfate $\geq 99.0\%$ were purchased from Honeywell Fluka (Seelze, Germany). Ethanol – 96% CZDA and methanol (HPLC grade) came from Avantor (Gliwice, Poland).

3.2. Testing material

3.2.1. Preparation of blackcurrant juice

The research material was juice made from frozen blackcurrants (*Ribes nigrum* L.) of the *Tisel* cultivar which is of Polish origin and bought from the local wholesaler. To produce blackcurrant juice, the fruit was enzymatically treated with a pectinolytic preparation (Klerzyme 150, DSM, France) for 1.5 h at 45°C , and then squeezed in a hydraulic layer press (Tako, Poland). The juice was transferred into 250 mL glass bottles and divided into six portions. The first was left without further processing (FJ). The second was heated to 45°C for 10 min (T45), the third was pasteurized at 85°C for 10 min (T85), both thermal processes were carried out in a laboratory pasteurizer (Labo Play, Bytom, Poland). The next portions were treated with supercritical carbon dioxide in the batch system using a Spe-ed SFE 4 (Applied Separations, USA). The method consists of placing the sample in a thermostatic pressure chamber, to pump CO_2 at the applicable pressure, and then leave it for a specified period of time to ensure penetration. Three pressure variants were applied: 10, 30 and 60 MPa for 10 min, at 45°C (SCCD10, SCCD30, SCCD60). The samples were frozen before carrying out further research. Samples not subjected to digestion were considered to be control samples.

3.3. In vitro digestion model with dialysis and calculation of bioaccessibility

In vitro gastrointestinal digestion was conducted according to a slightly modified method presented by Buniowska et al. (2017) [66] and international consensus on the *in vitro* digestion method suitable for food [15]. Three replications of the juice samples, and distilled water as a blank sample, were poured into dark glass bottles and mixed with 5 mL of salivary enzyme solution (2.38 g of Na_2HPO_4 , 0.19 g of K_2HPO_4 , 8 g of NaCl, 100 mg of mucin and α - amylase with enzymatic activity 200 U/L, solution per 1 L of distilled water). The pH of the solution was then adjusted (HI 211 meter, Hanna Instruments, Woonsocket, RI, USA) to 6.75 ± 0.20 by adding HCl (12 mol/L) or NaOH (2 mol/L). The solution was incubated in a shaking water bath (Labo Play, SWB 8N, Bytom, Poland) at 37°C and 90 rpm for 10 minutes. Gastric digestion was then performed by adding pepsin solution and adjusting the pH to 2.0 with HCl (12 mol/L) and again incubating for 2 h. 20 mL of each gastric digestion sample was placed in a clean bottle, titrated with NaOH (2 mol/L) to pH 5.00 ± 0.20 and 5 mL of pancreatin (1 g/L) and bile solution (25 g/L) were added to carry out the intestinal phase. The previously prepared dialysis membranes (cellulose, width 25 mm, length 30 cm) filled with 25 mL of NaHCO_3 solution (0.5 M, pH 7.5) were immersed in the digested samples. The samples were incubated once more for 2 h under the same conditions. In order to finish the digestion simulation, the samples were cooled in an ice

bath for 10 min. Samples were taken from each phase of the simulated digestion and frozen and stored until analysis. The dialysate solution inside the membrane is part of the sample potentially accessible to get into the bloodstream. Bioaccessibility (BAc) was determined using Equation 1 and is expressed as a percentage.

$$\text{BAc [\%]} = 100 \times (\text{BC}_{\text{digested}} / \text{BC}_{\text{non-digested}}) \quad (1)$$

Equation 1. Calculation of bioaccessibility (BAc – bioaccessibility of bioactive compound; $\text{BC}_{\text{digested}}$ – the concentration of bioactive compound in the digested sample; $\text{BC}_{\text{non-digested}}$ – the concentration of bioactive compound in the non-digested sample).

3.4. Chemical analysis

3.4.1. Determination of vitamin C

The vitamin C content was determined as a sum of L-ascorbic acid (AA) and L-dehydroascorbic acid (DHAA) according to the method introduced by Odriozola-Serrano et al. (2007) [67]. The sample was diluted using 0.01% phosphoric acid and filtered on a disposable syringe filter (0.45 μm , Macherey-Nagel, Düren, Germany). A solution of dithiothreitol (DTT) (1 g/L in 0.01% phosphoric acid) was used as the reducing agent to indicate the sum of AA and DHAA. Samples mixed with DTT in a 1:1 proportion were kept for 1 h in a dark place and at 4°C before further analysis. The devices used for the analysis were: Waters chromatographic system (Milford, MA, USA), 2695 Separations Module, 2995 Photodiode Array Detector, Sunfire C 28 column, 5 μm , 4.6 mm x 250 mm with reversed phase and Sunfire C18 Sentry guard insert, 5 μm , 4.6 mm x 20 mm (both Waters, Milford, MA, USA). Samples were eluted isocratically using 0.01% phosphoric acid at a flow rate of 1 ml/min. Compounds were quantified using UV absorption at 245 nm. The amount of DHAA was calculated based on the difference between the sum of both acids and AA.

3.4.2. Determination of anthocyanins

The total anthocyanin content and their individual monomers were determined according to the method presented by Oszmiański (2002) [68]. The sample injection was 10 μl , and the analysis time was 26 minutes, the column was heated to 25°C as before. A 4.5% aqueous formic acid solution (A) and an 80% acetonitrile solution in the previous formic acid solution (B) at a flow rate of 1.0 ml / min were the eluent. Anthocyanins were quantified using vis absorption at 520 nm. The amount of anthocyanins was calculated as cyanidin-3-glucoside.

3.4.3. Antioxidant activity according to the ABTS+• radical assay

The method described by Re et al. (1999) [69] was the principle used to determine the antioxidant activity with the ABTS+• radical. 18 hours before starting the analyses, a cationic radical solution was prepared by combining 7 mM ABTS+• and 2.45 mM potassium persulfate and was kept in a dark place. The radical solution was diluted with ethanol to an absorbance of 0.740–0.750 at a wavelength of 734 nm. The 1 mg/mL solution of Trolox in ethanol was the reference material, which was also used to prepare the standard curve. During the analysis, 0.025 mL of the sample, or distilled water as a blank, was dispensed into cuvettes, 2.5 mL of ABTS+• radical solution was added, mixed thoroughly and incubated at 30°C for 6 min. Absorbance was measured at 734 nm in ethanol. The antioxidant activity was expressed as Trolox Equivalent Antioxidant Capacity (TEAC). The absorbance was measured with the use of a Pharmacia Biotech UV / Vis spectrophotometer, model Ultrospec 2000 (England).

3.4.4. Antioxidative activity according to DPPH• radical assay

The method presented by Yen and Chen (1995) [70] was the guideline used to determine the antioxidant activity with the DPPH• radical. A solution of the DPPH• radical (1 mM) in methanol was prepared 3 h before starting the analyses and left to incubate in a dark place. After this time, it was diluted with 80% methanol to obtain a concentration of 0.1 mM and an absorbance in the range of 0.700–0.800. A standard curve was prepared using Trolox dissolved in methanol (1 mg/ml). 2 mL of DPPH• solution and 0.1 mL of juice, or a blank sample, were mixed in a cuvette. After incubation at room temperature in the dark for 30 min, the absorbance of the samples was measured at 517 nm. A Pharmacia Biotech spectrophotometer (Ultrospec 2000; England) was also used this time. The antioxidant activity is expressed as Trolox Equivalent Antioxidant Capacity (TEAC).

3.4.5. Determination of anthocyanin metabolites in a model system

3.4.5.1. Preparation of the delphinidin-3-*O*-rutinoside solution

In order to recognize the metabolites that are formed after anthocyanin digestion, an aqueous solution of 0.05 mg/ml of delphinidin-3-*O*-rutinoside (Df 3-*O*-rut) was prepared and subjected to simulated digestion in the same conditions as the blackcurrant juice. The solution was subjected to LC-MS/MS analysis before digestion and after each digestion step.

3.4.5.1. LC-MS/MS analysis

The analysis was carried out according to the methodology developed by Kapusta et al., (2018) [71]. Anthocyanin metabolites were investigated using a UPLC-PDA-MS/MS Waters ACQUITY system (Waters, Milford, MA, USA), with binary pump manager, sample manager, column manager, PDA detector and tandem quadrupole mass spectrometer (TQD) with electrospray ionization (ESI). The separation was performed using a BEH C18 column (100 mm × 2.1 mm i.d., 1.7 µm, Waters) kept at 50°C. The applied solvents were: 2% formic acid in water v/v (mobile phase A) and 2% formic acid in 40% acetonitrile in water v/v (mobile phase B). The gradient program was set as follows: 0 min 5% B, 0 – 8 min linear to 100% B, 8 – 9.5 min for washing and return to initial conditions. Before injection, samples were filtered through a membrane filter (0.45 µm, Merck Millipore) and injected directly into a chromatographic column. The injection volume was 5 µL (partial loop with needle overfill) and the flow rate was 0.35 mL/min. The parameters applied for TQD were: capillary voltage 3.5 kV; cone voltage 30 V in positive and negative mode; the source was maintained at 250°C and the desolvation temperature was 350°C; cone gas flow 100 l/h; and desolvation gas flow 800 l/h. Argon was used as the collision gas at a flow rate of 0.3 ml/min. The detection and identification of chemical compounds were based on characteristic PDA spectra, mass to charge ratio and ion fragments formed after collision-induced dissociation (CID). The quantitative analysis was possible only for one compound and was based on specific MS transitions in Multiple Reaction Monitoring (MRM) mode. Quantification was performed by injecting a standard solution of certain concentration ranging from 0.05 to 5 mg/ml ($R^2 \leq 0.9998$). Waters MassLynx software v.4.1 was used for data acquisition and processing.

3.5. Statistical analysis

A statistical analysis of the results obtained was carried out using the Statistica 7.1 program (StatSoft, Tulsa, OK, USA). A one-way analysis of variance with the ANOVA test and a significance analysis of the differences in mean values were performed using Tukey's test with a confidence level of $\alpha = 0.05$. Each sample was digested in three independent trials and analysed in duplicate.

4. Conclusions

Supercritical carbon dioxide (SCCD) processing at 10, 30 and 60 MPa for 10 min and 45°C contributed to an improvement in the stability of vitamin C, total anthocyanins and antioxidant capacity (AC) measured using the ABTS+• assay in blackcurrant juices. Thermal treatment at 85°C was insignificant for the stability of vitamin C, but caused a significant

degradation of total anthocyanins and a decrease in AC. As a result of simulated gastrointestinal digestion in the *in vitro* model, ascorbic acid was gradually oxidized to dehydroascorbic acid, and the total vitamin C and anthocyanin content decreased. No improvement in the BAC of vitamin C and total anthocyanins was reported after SCCD treatment. The positive effect of SCCD at 10 MPa was noted for the BAC of glycosides: delphinidin-3-*O*-glucoside and delphinidin-3-*O*-rutinoside compared to thermal treated samples. Overall, cyanidins were more bioaccessible than delphinidins, led by cyanidin-3-*O*-rutinoside in all sample types. An inverse correlation between AC and BAC of antioxidants was noted. This may indicate the formation of metabolites that were impossible to detect but showed a high antioxidant value. The calculated BAC concerns dialysate compounds in the parent form while they may have degraded to undetectable forms which are still of high antioxidant value. The UPLC-PDA-MS / MS analysis of the digested delphinidin-3-*O*-rutinoside extract showed that the main metabolite of this glycoside is protocatechuic acid, distinguished by its high antioxidant activity. SCCD is a non-thermal processing technique that increases the extractivity of bioactive compounds and reduces their degradation by penetrating the sample using CO₂ at supercritical state, while displacing oxygen, and inactivating tissue enzymes.

The significantly higher antioxidant activity in the SCCD samples, especially at 30 and 60 MPa, than in the thermal samples, suggests that this technique may be an effective method of improving the health value of antioxidant-rich products. It also confirms the paradox of the low BAC of antioxidants such as anthocyanins, while maintaining antioxidant activity and their beneficial effects on the body. Further research is needed to understand antioxidant metabolites and their bioaccessibility to better explore the possible uses of SCCD to improve the health properties of food.

Author Contributions: Conceptualization, U. Trych, K. Marszałek and M. Buniowska; methodology, K. Marszałek and M. Buniowska; formal analysis, U. Trych, I. Kapusta; investigation, U. Trych; data curation, U. Trych; writing—original draft preparation, U. Trych; writing—review and editing, K. Marszałek, S. Skąpska; supervision, K. Marszałek, M. Buniowska and S. Skąpska. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Marszałek, K.; Skąpska, S.; Woźniak, Ł.; Sokołowska, B. Application of Supercritical Carbon Dioxide for the Preservation of Strawberry Juice: Microbial and Physicochemical Quality, Enzymatic Activity and the Degradation Kinetics of Anthocyanins during Storage. *Innovative Food Science and Emerging Technologies* **2015**, *32*, 101–109, doi:10.1016/j.ifset.2015.10.005.
2. Marszałek, K.; Woźniak, Ł.; Skąpska, S. Effect of Supercritical Carbon Dioxide on Selected Quality Parameters of Preserved Strawberry Juice. *Zywnosc.Nauka.Technologia.Jakosc/Food.Science.Technology.Quality* **2015**, *21*, 114–123, doi:10.15193/zntj/2015/99/026.
3. Marszałek, K.; Woźniak, Ł.; Barba, F.J.; Skąpska, S.; Lorenzo, J.M.; Zambon, A.; Spilimbergo, S. Enzymatic, Physicochemical, Nutritional and Phytochemical Profile Changes of Apple (Golden Delicious L.) Juice under Supercritical Carbon Dioxide and Long-Term Cold Storage. *Food Chemistry* **2018**, *268*, 279–286, doi:10.1016/j.foodchem.2018.06.109.
4. Xue, Z.; Li, J.; Yu, W.; Lu, X.; Kou, X. Effects of Nonthermal Preservation Technologies on Antioxidant Activity of Fruits and Vegetables: A Review. *Food Science and Technology International* **2016**, *22*, 440–458, doi:10.1177/1082013215606835.
5. Zhou, L.; Wang, Y.; Hu, X.; Wu, J.; Liao, X. Effect of High Pressure Carbon Dioxide on the Quality of Carrot Juice. *Innovative Food Science and Emerging Technologies* **2009**, *10*, 321–327, doi:10.1016/j.ifset.2009.01.002.

6. Marszałek, K.; Doesburg, P.; Starzonek, S.; Szczepańska, J.; Woźniak, Ł.; Lorenzo, J.M.; Skaopska, S.; Rzoska, S.; Barba, F.J. Comparative Effect of Supercritical Carbon Dioxide and High Pressure Processing on Structural Changes and Activity Loss of Oxidoreductive Enzymes. *Journal of CO2 Utilization* **2019**, *29*, 46–56, doi:10.1016/j.jcou.2018.11.007.
7. Marszałek, K.; Woźniak, Ł.; Kruszewski, B.; Skapska, S. The Effect of High Pressure Techniques on the Stability of Anthocyanins in Fruit and Vegetables. *International Journal of Molecular Sciences* **2017**, *18*.
8. Lyall, K.A.; Hurst, S.M.; Cooney, J.; Jensen, D.; Lo, K.; Hurst, R.D.; Stevenson, L.M. Short-Term Blackcurrant Extract Consumption Modulates Exercise-Induced Oxidative Stress and Lipopolysaccharide-Stimulated Inflammatory Responses. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology* **2009**, *297*, 70–81, doi:10.1152/ajpregu.90740.2008.
9. Li, L.; Li, S.; Hu, C.; Zhou, L.; Zhang, Y.; Wang, M.; Qi, Z. BKCa Channel Is a Molecular Target of Vitamin C to Protect against Ischemic Brain Stroke. *Molecular Membrane Biology* **2019**, *35*, 9–20, doi:10.1080/09687688.2019.1608378.
10. Bakowska-Barczak, A.M.; Kolodziejczyk, P.P. Black Currant Polyphenols: Their Storage Stability and Microencapsulation. *Industrial Crops and Products* **2011**, *34*, 1301–1309, doi:10.1016/j.indcrop.2010.10.002.
11. Diaconeasa, Z.; Leopold, L.; Rugină, D.; Ayvaz, H.; Socaciu, C. Antiproliferative and Antioxidant Properties of Anthocyanin Rich Extracts from Blueberry and Blackcurrant Juice. *International Journal of Molecular Sciences* **2015**, *16*, 2352–2365, doi:10.3390/ijms16022352.
12. Lila, M.A.; Burton-Freeman, B.; Grace, M.; Kalt, W. Unraveling Anthocyanin Bioavailability for Human Health. *Annual Review of Food Science and Technology* **2016**, *7*, 375–393, doi:10.1146/annurev-food-041715-033346.
13. Rashid, K.; Wachira, F.N.; Nyabuga, J.N.; Wanyonyi, B.; Murilla, G.; Isaac, A.O. Kenyan Purple Tea Anthocyanins Ability to Cross the Blood Brain Barrier and Reinforce Brain Antioxidant Capacity in Mice. *Nutritional Neuroscience* **2014**, *17*, 178–185, doi:10.1179/1476830513Y.0000000081.
14. Stahl, W.; Van Den Berg, H.; Arthur, J.; Bast, A.; Dainty, J.; Faulks, R.M.; Gärtner, C.; Haenen, G.; Hollman, P.; Holst, B.; et al. Bioavailability and Metabolism. In *Molecular Aspects of Medicine*; Azzi, A., Ed.; PERGAMON, 2002; Vol. 23, pp. 39–100.
15. Minekus, M.; Alming, M.; Alvito, P.; Ballance, S.; Bohn, T.; Bourlieu, C.; Carrière, F.; Boutrou, R.; Corredig, M.; Dupont, D.; et al. A Standardised Static in Vitro Digestion Method Suitable for Food-an International Consensus. *Food and Function* **2014**, *5*, 1113–1124, doi:10.1039/c3fo60702j.
16. Fernández-García, E.; Carvajal-Lérida, I.; Pérez-Gálvez, A. In Vitro Bioaccessibility Assessment as a Prediction Tool of Nutritional Efficiency. *Nutrition Research* **2009**, *29*, 751–760.
17. Mattila, P.H.; Hellström, J.; McDougall, G.; Dobson, G.; Pihlava, J.M.; Tiirikka, T.; Stewart, D.; Karjalainen, R. Polyphenol and Vitamin C Contents in European Commercial Blackcurrant Juice Products. *Food Chemistry* **2011**, *127*, 1216–1223, doi:10.1016/j.foodchem.2011.01.129.
18. Nour, V.; Trandafir, I.; Ionica, M.E. Ascorbic Acid, Anthocyanins, Organic Acids and Mineral Content of Some Black and Red Currant Cultivars. *Fruits* **2011**, *66*, 353–362, doi:10.1051/fruits/2011049.
19. Oulé, K.M.; Dickman, M.; Arul, J. Properties of Orange Juice with Supercritical Carbon Dioxide Treatment. *International Journal of Food Properties* **2013**, *16*, 1693–1710, doi:10.1080/10942912.2011.604893.
20. Fabroni, S.; Amenta, M.; Timpanaro, N.; Rapisarda, P. Supercritical Carbon Dioxide-Treated Blood Orange Juice as a New Product in the Fresh Fruit Juice Market. *Innovative Food Science and Emerging Technologies* **2010**, *11*, 477–484, doi:10.1016/j.ifset.2010.02.004.
21. Rodríguez-Roque, M.J.; Rojas-Graü, M.A.; Elez-Martínez, P.; Martín-Belloso, O. Changes in Vitamin C, Phenolic, and Carotenoid Profiles throughout in Vitro Gastrointestinal Digestion of a Blended Fruit Juice. *Journal of Agricultural and Food Chemistry* **2013**, *61*, 1859–1867.
22. Pérez-Vicente, A.; Gil-Izquierdo, A.; García-Viguera, C. In Vitro Gastrointestinal Digestion Study of Pomegranate Juice Phenolic Compounds, Anthocyanins, and Vitamin C. *Journal of Agricultural and Food Chemistry* **2002**, *50*, 2308–2312, doi:10.1021/jf0113833.

23. Vallejo, F.; Gil-Izquierdo, A.; Pérez-Vicente, A.; García-Viguera, C. In Vitro Gastrointestinal Digestion Study of Broccoli Inflorescence Phenolic Compounds, Glucosinolates, and Vitamin C. *Journal of Agricultural and Food Chemistry* **2004**, *52*, 135–138, doi:10.1021/jf040108o.
24. Trych, U.; Buniowska, M.; Skapska, S.; Starzonek, S.; Marszałek, K. The Bioaccessibility of Antioxidants in Black Currant Puree after High Hydrostatic Pressure Treatment. *Molecules* **2020**, *25*, doi:10.3390/molecules25153544.
25. Rodríguez-Roque, M.J. In Vitro Bioaccessibility of Health-Related Compounds from Beverages Based on Fruit Juice, Milk or Soymilk: Influence of Food Matrix and Processing, Universitat de Lleida, 2014.
26. Lingua, M.S.; Wunderlin, D.A.; Baroni, M. V. Effect of Simulated Digestion on the Phenolic Components of Red Grapes and Their Corresponding Wines. *Journal of Functional Foods* **2018**, *44*, 86–94, doi:10.1016/j.jff.2018.02.034.
27. Rodríguez-Roque, M.J.; de Ancos, B.; Sánchez-Moreno, C.; Cano, M.P.; Elez-Martínez, P.; Martín-Belloso, O. Impact of Food Matrix and Processing on the in Vitro Bioaccessibility of Vitamin C, Phenolic Compounds, and Hydrophilic Antioxidant Activity from Fruit Juice-Based Beverages. *Journal of Functional Foods* **2015**, *14*, 33–43.
28. Cilla, A.; Alegría, A.; De Ancos, B.; Sánchez-Moreno, C.; Cano, M.P.; Plaza, L.; Clemente, G.; Lagarda, M.J.; Barberá, R. Bioaccessibility of Tocopherols, Carotenoids, and Ascorbic Acid from Milk- and Soy-Based Fruit Beverages: Influence of Food Matrix and Processing. *Journal of Agricultural and Food Chemistry* **2012**, *60*, 7282–7290.
29. Sánchez-Moreno, C.; Cano, M.P.; De Ancos, B.; Plaza, L.; Olmedilla, B.; Granado, F.; Martín, A. Consumption of High-Pressurized Vegetable Soup Increases Plasma Vitamin C and Decreases Oxidative Stress and Inflammatory Biomarkers in Healthy Humans. *Journal of Nutrition* **2004**, *134*, 3021–3025.
30. Moyer, R.A.; Hummer, K.E.; Finn, C.E.; Frei, B.; Wrolstad, R.E. Anthocyanins, Phenolics, and Antioxidant Capacity in Diverse Small Fruits: *Vaccinium*, *Rubus*, and *Ribes*. *Journal of Agricultural and Food Chemistry* **2002**, *50*, 519–525.
31. Paredes-López, O.; Cervantes-Ceja, M.L.; Vigna-Pérez, M.; Hernández-Pérez, T. Berries: Improving Human Health and Healthy Aging, and Promoting Quality Life-A Review. *Plant Foods for Human Nutrition* **2010**, *65*, 299–308, doi:10.1007/s11130-010-0177-1.
32. Pozo-Insfran, D. del; Balaban, M.O.; Talcott, S.T. Inactivation of Polyphenol Oxidase in Muscadine Grape Juice by Dense Phase-CO₂ Processing. *Food Research International* **2007**, *40*, 894–899, doi:10.1016/j.foodres.2007.03.002.
33. Bouayed, J.; Hoffmann, L.; Bohn, T. Total Phenolics, Flavonoids, Anthocyanins and Antioxidant Activity Following Simulated Gastro-Intestinal Digestion and Dialysis of Apple Varieties: Bioaccessibility and Potential Uptake. *Food Chemistry* **2011**, *128*, 14–21, doi:10.1016/j.foodchem.2011.02.052.
34. Correa-Betanzo, J.; Allen-Vercos, E.; McDonald, J.; Schroeter, K.; Corredig, M.; Paliyath, G. Stability and Biological Activity of Wild Blueberry (*Vaccinium Angustifolium*) Polyphenols during Simulated in Vitro Gastrointestinal Digestion. *Food Chemistry* **2014**, *165*, 522–531, doi:10.1016/j.foodchem.2014.05.135.
35. Fernandes, I.; Faria, A.; de Freitas, V.; Calhau, C.; Mateus, N. Multiple-Approach Studies to Assess Anthocyanin Bioavailability. *Phytochemistry Reviews* **2015**, *14*, 899–919, doi:10.1007/s11101-015-9415-3.
36. Sigurdson, G. T., Giusti, M.M. The Stability and Absorption of Anthocyanins in the Mouth. In *Anthocyanins from Natural Sources*; Brooks, M. S., Celli, G.B., Ed.; The Royal Society of Chemistry: London (UK), 2019; pp. 186–215.
37. Milbury, P.E.; Cao, G.; Prior, R.L.; Blumberg, J. Bioavailability of Elderberry Anthocyanins. *Mechanisms of Ageing and Development* **2002**, *123*, 997–1006, doi:10.1016/S0047-6374(01)00383-9.
38. Cao, G.; Muccitelli, H.U.; Sánchez-Moreno, C.; Prior, R.L. Anthocyanins Are Absorbed in Glycated Forms in Elderly Women: A Pharmacokinetic Study. *American Journal of Clinical Nutrition* **2001**, *73*, 920–926, doi:10.1093/ajcn/73.5.920.
39. Fernandes, I.; Faria, A.; Calhau, C.; de Freitas, V.; Mateus, N. Bioavailability of Anthocyanins and Derivatives. *Journal of Functional Foods* **2014**, *7*, 54–66, doi:10.1016/j.jff.2013.05.010.
40. Fang, J. Bioavailability of Anthocyanins. *Drug Metabolism Reviews* **2014**, *46*, 508–520, doi:10.3109/03602532.2014.978080.

41. Mueller, D.; Jung, K.; Winter, M.; Rogoll, D.; Melcher, R.; Richling, E. Human Intervention Study to Investigate the Intestinal Accessibility and Bioavailability of Anthocyanins from Bilberries. *Food Chemistry* **2017**, *231*, 275–286, doi:10.1016/j.foodchem.2017.03.130.
42. Peixoto, F. M.; Senna Gouvêa, A.; de Araújo Santiago, M. C.; de Sá Veloso Martins, Z. E.; Galhardo Borguini, R.; & de Oliveira Godoy, R.L. Characterization and Bioaccessibility of Anthocyanins from Blueberry (*Vaccinium Corymbosum* L.) after Simulated Gastro-Intestinal Digestion: A Positive Effect on Malvidin Derivatives. *Fruits* **2018**, *73*, 101–109.
43. Mallery, S.R.; Budendorf, D.E.; Larsen, M.P.; Pei, P.; Tong, M.; Holpuch, A.S.; Larsen, P.E.; Stoner, G.D.; Fields, H.W.; Chan, K.K.; et al. Effects of Human Oral Mucosal Tissue, Saliva, and Oral Microflora on Intraoral Metabolism and Bioactivation of Black Raspberry Anthocyanins. *Cancer Prevention Research* **2011**, *4*, 1209–1221, doi:10.1158/1940-6207.CAPR-11-0040.
44. Vitaglione, P.; Donnarumma, G.; Napolitano, A.; Galvano, F.; Gallo, A.; Scalfi, L.; Fogliano, V. Protocatechuic Acid Is the Major Human Metabolite of Cyanidin-Glucosides. *Journal of Nutrition* **2007**, *137*, 2043–2048, doi:10.1093/jn/137.9.2043.
45. Goszcz, K.; Deakin, S.J.; Duthie, G.G.; Stewart, D.; Megson, I.L. Bioavailable Concentrations of Delphinidin and Its Metabolite, Gallic Acid, Induce Antioxidant Protection Associated with Increased Intracellular Glutathione in Cultured Endothelial Cells. *Oxidative Medicine and Cellular Longevity* **2017**, *2017*, doi:10.1155/2017/9260701.
46. Nurmi, T.; Mursu, J.; Heinonen, M.; Nurmi, A.; Hiltunen, R.; Voutilainen, S. Metabolism of Berry Anthocyanins to Phenolic Acids in Humans. *Journal of Agricultural and Food Chemistry* **2009**, *57*, 2274–2281, doi:10.1021/jf8035116.
47. Del Bò, C.; Ciappellano, S.; Klimis-Zacas, D.; Daniela, M.; Claudio, G.; Riso, P.; Porrini, M. Anthocyanin Absorption, Metabolism, and Distribution from a Wild Blueberry-Enriched Diet (*Vaccinium Angustifolium*) Is Affected by Diet Duration in the Sprague-Dawley Rat. *Journal of Agricultural and Food Chemistry* **2010**, *58*, 2491–2497, doi:10.1021/jf903472x.
48. He, Y.; Hou, X.; Guo, J.; He, Z.; Guo, T.; Liu, Y.; Zhang, Y.; Zhang, J.; Feng, N. Activation of a Gamma-Cyclodextrin-Based Metal–Organic Framework Using Supercritical Carbon Dioxide for High-Efficient Delivery of Honokiol. *Carbohydrate Polymers* **2020**, *235*, doi:10.1016/j.carbpol.2020.115935.
49. Zhang, Z.; Quan, G.; Wu, Q.; Zhou, C.; Li, F.; Bai, X.; Li, G.; Pan, X.; Wu, C. Loading Amorphous Asarone in Mesoporous Silica SBA-15 through Supercritical Carbon Dioxide Technology to Enhance Dissolution and Bioavailability. *European Journal of Pharmaceutics and Biopharmaceutics* **2015**, *92*, 28–31, doi:10.1016/j.ejpb.2015.02.018.
50. Hassan, H.A.; Al-Marzouqi, A.H.; Jobe, B.; Hamza, A.A.; Ramadan, G.A. Enhancement of Dissolution Amount and in Vivo Bioavailability of Itraconazole by Complexation with β -Cyclodextrin Using Supercritical Carbon Dioxide. *Journal of Pharmaceutical and Biomedical Analysis* **2007**, *45*, 243–250, doi:10.1016/j.jpba.2007.06.011.
51. Kankala, R.K.; Zhang, Y.S.; Wang, S. bin; Lee, C.H.; Chen, A.Z. Supercritical Fluid Technology: An Emphasis on Drug Delivery and Related Biomedical Applications. *Advanced Healthcare Materials* **2017**, *6*, doi:10.1002/adhm.201700433.
52. Zhao, W.; Sun, Y.; Ma, Y.; Zhao, X. Dense Phase Carbon Dioxide Treatment of Tomato Juice: Effect on Physico-Chemical Properties, Phenolic Composition, Lycopene Isomerisation and in Vitro Bioaccessibility. *International Journal of Food Science and Technology* **2019**, *54*, 1658–1669, doi:10.1111/ijfs.14042.
53. Briongos, H.; Illera, A.E.; Sanz, M.T.; Melgosa, R.; Beltrán, S.; Solaesa, A.G. Effect of High Pressure Carbon Dioxide Processing on Pectin Methylesterase Activity and Other Orange Juice Properties. *LWT - Food Science and Technology* **2016**, *74*, 411–419, doi:10.1016/j.lwt.2016.07.069.
54. Niu, L.; Hu, X.; Wu, J.; Liao, X.; Chen, F.; Zhao, G.; Wang, Z. Effect of Dense Phase Carbon Dioxide Process on Physicochemical Properties and Flavor Compounds of Orange Juice. *Journal of Food Processing and Preservation* **2010**, *34*, 530–548, doi:10.1111/j.1745-4549.2009.00369.x.
55. Ubeyitogullari, A. *ENHANCING BIOACCESSIBILITY OF PHYTOSTEROLS USING NANOPOROUS STARCH AEROGELS AND SUPERCRITICAL CARBON DIOXIDE*; 2018;
56. Vallecilla Yopez, L.; Yopez, V. *Increasing Cis-Lycopene Content of the Oleoresin from Tomato Processing Byproducts Using Supercritical Carbon Dioxide and Assessment of Its Bioaccessibility*; 2017; Vol. 89;.

57. Cassani, L.; Gerbino, E.; Moreira, M. del R.; Gómez-Zavaglia, A. Influence of Non-Thermal Processing and Storage Conditions on the Release of Health-Related Compounds after in Vitro Gastrointestinal Digestion of Fiber-Enriched Strawberry Juices. *Journal of Functional Foods* **2018**, *40*, 128–136, doi:10.1016/j.jff.2017.11.005.
58. Ferrentino, G.; Plaza, M.L.; Ramirez-Rodrigues, M.; Ferrari, G.; Balaban, M.O. Effects of Dense Phase Carbon Dioxide Pasteurization on the Physical and Quality Attributes of a Red Grapefruit Juice. *Journal of Food Science* **2009**, *74*, doi:10.1111/j.1750-3841.2009.01250.x.
59. Ramírez-Rodrigues, M.M.; Plaza, M.L.; Azeredo, A.; Balaban, M.O.; Marshall, M.R. Phytochemical, Sensory Attributes and Aroma Stability of Dense Phase Carbon Dioxide Processed Hibiscus Sabdariffa Beverage during Storage. *Food Chemistry* **2012**, *134*, 1425–1431, doi:10.1016/j.foodchem.2012.03.042.
60. Porto, C. Da; Decorti, D.; Tubaro, F. Effects of Continuous Dense-Phase CO₂ System on Antioxidant Capacity and Volatile Compounds of Apple Juice. *International Journal of Food Science and Technology* **2010**, *45*, 1821–1827, doi:10.1111/j.1365-2621.2010.02339.x.
61. Liu, L.; Zeng, Q.; Zhang, R.; Wei, Z.; Deng, Y.; Zhang, Y.; Tang, X.; Zhang, M. Comparative Study on Phenolic Profiles and Antioxidant Activity of Litchi Juice Treated by High Pressure Carbon Dioxide and Thermal Processing. *Food Science and Technology Research* **2015**, *21*, 41–49, doi:10.3136/fstr.21.41.
62. Trych, U.; Buniowska, M.; Skapska, S.; Starzonek, S.; Marszałek, K. The Bioaccessibility of Antioxidants in Black Currant Puree after High Hydrostatic Pressure Treatment. *Molecules* **2020**, *25*, 1–15, doi:10.3390/molecules25153544.
63. Burgos-Edwards, A.; Jiménez-Aspee, F.; Thomas-Valdés, S.; Schmeda-Hirschmann, G.; Theoduloz, C. Qualitative and Quantitative Changes in Polyphenol Composition and Bioactivity of Ribes Magellanicum and R. Punctatum after in Vitro Gastrointestinal Digestion. *Food Chemistry* **2017**, *237*, 1073–1082, doi:10.1016/j.foodchem.2017.06.060.
64. Prior, R.L. *Fruits and Vegetables in the Prevention of Cellular Oxidative Damage 1-5*; 2003; Vol. 78;.
65. Goszcz, K.; Deakin, S.J.; Duthie, G.G.; Stewart, D.; Megson, I.L. Bioavailable Concentrations of Delphinidin and Its Metabolite, Gallic Acid, Induce Antioxidant Protection Associated with Increased Intracellular Glutathione in Cultured Endothelial Cells. *Oxidative Medicine and Cellular Longevity* **2017**, *2017*, doi:10.1155/2017/9260701.
66. Buniowska, M.; Carbonell-Capella, J.M.; Frigola, A.; Esteve, M.J. Bioaccessibility of Bioactive Compounds after Non-Thermal Processing of an Exotic Fruit Juice Blend Sweetened with Stevia Rebaudiana. *Food Chemistry* **2017**, *221*, 1834–1842.
67. Odriozola-Serrano, I.; Hernández-Jover, T.; Martín-Belloso, O. Comparative Evaluation of UV-HPLC Methods and Reducing Agents to Determine Vitamin C in Fruits. *Food Chemistry* **2007**, *105*, 1151–1158, doi:10.1016/j.foodchem.2007.02.037.
68. Oszmianski, J. Stabilizacja i Zastosowanie Barwnika Antocyjanowego Aronii Do Barwienia Napoi. *Acta Scientiarum Polonorum. Technologia Alimentaria* **2002**, *01*, 37–45.
69. Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans, C. Antioxidant Activity Applying an Improved ABTS Radical Cation Decolorization Assay. *Free Radical Biology and Medicine* **1999**, *26*, 1231–1237.
70. Yen, G. C., Chen, H.Y. Antioxidant Activity of Various Tea Extracts in Relation to Their Antimutagenicity. *Journal of Agricultural and Food Chemistry* **1995**, *43*, 27–32.
71. Kapusta, I.; Cebulak, T.; Oszmiański, J. Characterization of Polish Wines Produced from the Interspecific Hybrid Grapes Grown in South-East Poland. *European Food Research and Technology* **2018**, doi:10.1007/s00217-017-2972-5.