

Article

Oral hygiene practices and knowledge among adolescents aged between 15 and 17 years old during fixed orthodontic treatment: multicentre study conducted in France.

Camille Inquimbert ^{1,2}, Celine Clement ^{2,3,4}, Antoine Couatarmanach ⁵, Paul Tramini ¹, Denis Bourgeois ², Florence Carrouel ^{2*}

¹ Department of Public Health, Faculty of Dental Medicine, University of Montpellier, France; camille.inquimbert@umontpellier.fr (C.I.); paul.tramini@umontpellier.fr (P.T.)

² Laboratory "Systemic Health Care", EA4129, University Claude Bernard Lyon 1, University of Lyon, 69008 Lyon, France; denis.bourgeois@univ-lyon1.fr (D.B.); florence.carrouel@univ-lyon1.fr (F.C.)

³ "Interpsy" Laboratory, University of Lorraine, EA 4432, 54015 Nancy, France

⁴ Department of Public Health, Faculty of Dental Medicine, University of Nancy, France; celine.clement@univ-lorraine.fr (C.C.)

⁵ Faculty of Dentistry, University of Rennes, UMR 6051, CHU Rennes, Rennes, France; antoine.couatarmanach@univ-rennes1.fr (A.C.)

* Correspondance: camille.inquimbert@umontpellier.fr; Tel.: +33666957674

Abstract: The aims of this study were to assess oral health knowledge, attitudes, and practices among orthodontic patients between the ages of 15 and 17 years old compared to adolescents without orthodontic treatment. This cross-sectional study included 392 adolescents drawn from various French teaching hospitals. A closed-ended questionnaire was used to collect data. Adolescents undergoing orthodontic treatment had a higher knowledge of oral health than adolescents without orthodontic treatment. The majority of adolescents for both groups (69%) claimed to brush their teeth twice a day. Regarding complimentary dental material, 81.9% of adolescents without orthodontic treatment never used an interdental brush and 78.8% never used dental floss. For those undergoing orthodontic treatment, 48.5% never used interdental brush. Only 4% of adolescents without and 3% of adolescents with orthodontic treatment never consumed fizzy drinks, 4.9% and 3% never consumed sweets, 4% and 8.4% never ate fast-food. Adolescents without treatment consumed more sodas ($p=0.04$) and more fast food ($p=0.03$). Adolescents had insufficient knowledge of oral health. Health education programmes should be implemented to improve adolescents' knowledge and individual oral prophylaxis with interdental brushes.

Keywords: Oral Health; Adolescent; Knowledge, Practice, Behaviour

1. Introduction

Orthodontic treatment is often performed on teenagers and its prevalence may range from 10% to 35% in developed countries (1). It usually involves fixed or removable appliances for the correction of mild to severe malocclusions. A systematic review found that orthodontic treatment usually lasts an average of 20 months, with a mean number of required visits at 17.8 (2). During this period, orthodontic appliances may increase biofilm accumulation and plaque retention, and inadequate oral hygiene can lead to permanent damage to dental tissue, caries, or periodontal lesions. Moreover, orthodontic patients demonstrate higher proportions of gram-negative species (3) resulting in more inflammation and bleeding (4). Previous studies (5,6) have shown that orthodontic care is often associated with oral diseases due to excessive plaque retention adjacent to

brackets and attachments. Therefore, adherence to prophylactic practices is essential to maintaining good oral health during orthodontic treatment.

Adolescence may increase the risk of dental diseases, as it is a period during which oral care habits are being established, with lower motivation regarding good oral hygiene maintenance (7–10). Moreover, adolescents tend to consume more snacks and beverages between meals. For all these reasons, oral care plays a key role (11–13) in adolescents becoming at increasing risk for caries [14–16] and early periodontal disease [17–19].

Currently, we know that maintaining oral hygiene in interproximal spaces calls for special devices, the use of interdental brushes being the most effective method for interproximal plaque removal (15,16). The daily use of interdental brushes can reduce inflammation and lead to the reestablishment of symbiotic interdental microbiota (17,18).

Changing behaviour is a very long and complex process that takes place over time and at different stages of life. Health literacy has been found to be a strong predictor of individual health, health behaviour and health outcomes (19,20). Health literacy is a known mediator between socio-economic factors, health behaviour and oral health outcomes in various populations, explaining gradients in oral health status and outcomes (21). An increase in oral health knowledge is often associated with an increased awareness of oral health and better oral hygiene practices (22).

Although many studies on oral health knowledge and the practice of preventive measures have been conducted among children, few studies are available among adolescents (23–27), and epidemiological studies on oral health status and dental knowledge are scarce among French adolescents (28,29).

Since adolescents undergoing orthodontic treatment receive more specific information and adequate hygiene recommendations at each regular orthodontic visit than adolescents without orthodontic treatment, the hypothesis of this study was that orthodontically treated adolescents would have better dental knowledge and even more adequate oral hygiene practices than non-treated adolescents. Therefore, the objective of this study was to describe the oral health knowledge, attitudes, and practices amongst a sample of 15-17-year-old French adolescents with and without orthodontic treatment.

2. Materials and Methods

2.1. Study site and population

A cross-sectional questionnaire-based survey was conducted from September 2018 to December 2018. A convenience sample was recruited from four teaching dental hospitals (public health and orthodontic departments) located from the north, east, west, and south of France. To be included in this study, (i) the patients had to be at least 15 years old and at most 17 on the day of the inclusion, (ii) they had to have a good understanding of the French language, (iii) the study's terms had to have been accepted by one of their parents and by the patients, (iv) and the written informed consent had to have been signed by one of their parents and also by the patients.

The exclusion criteria were: (i) patients who were mentally or physically challenged, and (ii) patients in need for urgent care, because they were not fit for answering the questions in the survey.

Two groups were set up, one for adolescents undergoing orthodontic treatment (OT) and another for adolescents without orthodontic treatment (NT).

2.2. Ethics

The protocol and study design were approved by ethics and regulatory agencies and were implemented in accordance with provisions of the Declaration of Helsinki. The appropriate Committee (Local Research Ethics Committee, Montpellier, France) approved the protocol on January 10, 2018 (2017_CLER_12-01). Informed consent had to be signed by all adolescents and one of the parents or legal representatives. The informed consent form contained the name and affiliation of the investigator, a plain language description of the study (intervention), the approximate duration of the interview and the ethics committee approval.

2.3. Measurement Tool

A questionnaire was established, comprising 36 questions (see Appendix 1). It was drawn from the ESCARCEL survey where a validated questionnaire was previously developed in different languages [42]. All questions were closed-ended questions. The patient's knowledge, attitude, and practices (KAP) were assessed by using a questionnaire which included four parts. The first part referred to general information: year of birth, gender, the father's and mother's occupations and social security coverage. The second part concerned the characteristics of the dental consultation. The third part described oral health, oral hygiene practices, dietary habits, and tobacco, drugs, and alcohol consumption. The last section of the questionnaire dealt with knowledge, in particular questions about oral health knowledge on hygiene, eating habits and the role of fluoride. Responses on self-perception of general and oral health and on dietary and hygiene habits were made on a four-point scale: "Often" (more than 3 times a week), "Sometimes" (one or two times a week), "Rarely" (less than once a week), and "never".

Adolescents were interviewed face-to-face by the dental practitioners who took part in the survey (C.I., X.X., Y.Y., Z.Z., ...). To avoid potential information bias, interviewers explained the purposes and confidentiality of the survey and explained that the study had no impact on participants' examinations. The data was collected on a sheet of paper (printed questionnaire) with an anonymity number (allocation of a random number).

2.4. Statistical method

Sample size: With a confidence level of 95%, a confidence interval of 5% and an average of 50% for any unknown percentage of the questionnaire, and considering the total number of adolescents in France (approximately 2,000,000 according to INSEE data in 2017), it was necessary to include at least 380 subjects in this study, with approximately 50% in each group (orthodontic treatment/no orthodontic treatment).

Statistical tests: Categorical variables were described by frequencies or percentages and quantitative variables using means and standard deviations. The relationships between oral health knowledge and other qualitative variables were analysed using chi-square tests. Student tests or Wilcoxon tests were used for quantitative variables, depending on the normality of these variables. Multivariate analyses, such as factor analysis, were used to analyse the relationships between different items of the questionnaire. Statistical significance was assumed when $p < 0.05$. All analyses were performed with the statistical software Stata 16.1 (StataCorp).

3. Results

3.1. Patient characteristics

The total sample comprised 392 adolescents, with 226 (57.7%) in the NT group versus 166 (42.3%) in the OT group. The gender distribution was 44.6% (175) males and 55.4% (217) females. The mean age of the participants was 16.43 (± 0.51), without gender difference: 16.41 and 16.45 for males and females respectively ($p = 0.73$). Sociodemographic characteristics of the sample are displayed in Table 1. The fathers' occupations were mostly administration, or employees, and a quarter were executives. The mothers' occupations were statistically different between the two groups ($p = 0.06$) with mostly employees and executives among the OT group. The percentage of adolescents benefiting from medical assistance was significantly higher in the OT group (81.3%) than in the NT group (70.3%), ($p = 0.01$).

Table 1. Sociodemographic characteristics of the sample according to orthodontic treatment.

Variable	No orthodontic treatment (n = 266) %	Orthodontic treatment (n=166) %	p-value (chi-square)
Mean Age (sd)	16.48 (0.53)	16.38 (0.49)	

Gender	Males	56.6	53.6	0.55
	Females	43.4	46.4	
Father's occupation	Farm worker/Shopkeeper	11.1	10.3	0.13
	Executive/Company manager	24.8	30.7	
	Administration	17.7	9.6	
	Employee/Manual worker	29.6	37.4	
	Unemployed/Disabled/Deceased	7.5	7.2	
	Unknown	9.3	4.8	
Mother's occupation	Farm worker/Shopkeeper	5.7	3.6	0.06
	Executive/Company manager	15.9	22.3	
	Administration	29.7	23.5	
	Employee/Manual worker	21.7	32.5	
	Unemployed/Disabled/Deceased	19.5	15.7	
	Unknown	7.5	2.4	
Any medical aid	No	29.7	18.7	0.01
	Yes	70.3	81.3	

3.2. Assessment of adolescents' knowledge regarding prevention of oral diseases (Table 2)

The proportion of adolescents who knew that fluorides prevent decay was higher in the OT group (65.1%) than in the NT group (46.0%), $p=0.0001$. There was no other significant difference in knowledge between the two groups. Regarding the duration of toothbrushing, 36.1% of adolescents from the OT group thought that the correct answer was between 2 and 3 minutes, while this percentage was 28.3% in the NT group ($p=0.06$).

Almost half of the sample in both groups thought that the role of fluorides was to prevent calculus (48.2% and 37.6%).

Table 2. Knowledge of adolescents regarding prevention of oral diseases according to orthodontic treatment.

Variable		No orthodontic treatment (n = 266) %	Orthodontic treatment (n=166) %	p-value (*) (chi-square)
Frequency of daily toothbrushing	Less than twice	3.6%	4.9%	0.91
	Twice	43.5%	43.3%	
	More than twice	52.9%	51.8%	
Duration of toothbrushing	1 minute or less	0.4%	0.0%	0.06
	2 minutes	6.6%	6.0%	
	3 minutes	21.7%	30.1%	
	> 3 minutes	59.7%	55.4%	
	I don't know	11.1%	7.8%	
Snacking reduces caries	No	80.5%	76.5%	0.33
	Yes	9.7%	12.1%	
Sticky food protects from decay	No	92.9%	96.4%	0.14
	Yes	1.8%	1.2%	
	I don't know	5.3%	2.4%	

F toothpaste prevents calculus	No	12.4%	13.3%	0.80
	Yes	37.6%	48.2%	
	I don't know	50.0%	38.6%	
F toothpaste makes teeth white	No	14.6%	18.1%	0.36
	Yes	44.3%	52.4%	
	I don't know	41.2%	29.5%	
F toothpaste provides fresh breath	No	22.6%	24.1%	0.72
	Yes	32.7%	39.8%	
	I don't know	44.7%	36.1%	
F toothpaste prevents decay	No	12.0%	7.2%	0.0001
	Yes	46.0%	65.1%	
	I don't know	42.0%	27.7%	

(*) *p-values were calculated without considering the answer 'I don't know'*

3.3. Assessment of adolescents' attitudes and practices towards oral health (Table 3)

3.3.1. Oral hygiene habits

The majority of participants from both groups (NT: 88.0%, OT: 91.0%) claimed to brush their teeth at least twice a day. Regarding oral hygiene practice, there was no significant difference between the two groups, except for the use of interdental brushes: adolescents from the OT group used them more often ($p=0.0001$). Concerning at what time the participants brushed their teeth, there were also some differences between the two groups: in the OT group, adolescents were more likely to brush their teeth after breakfast ($p=0.01$), and after lunch ($p=0.01$), while in the NT group, they were more likely to brush before breakfast ($p=0.02$). For brushing after dinner, there was no difference between the groups ($p=0.89$).

3.3.2. Nutrition and harmful habits

Fizzy drinks were more often consumed by adolescents without orthodontic treatment ($p=0.04$). In the same way, fast foods and cigarettes showed significantly higher percentages in the NT group. Sweets were more often consumed by adolescents without orthodontic treatment, but the difference was not significant ($p=0.17$). High frequency of snacking (5 times a day or more) was significantly increased in the NT group (63.3% vs 52.4%, $p=0.03$). Participants claimed that they rarely consumed alcohol and drugs (77.0%), and 96.4% never used alcohol and drugs.

Table 3. Oral hygiene, nutrition, and harmful habits among adolescents aged between 15 and 17 years old according to orthodontic treatment.

Variable	No orthodontic treatment (n = 266) %	Orthodontic treatment (n=166) %	p-value (*) (chi-square)
Dental floss use			0.63
	Often	0.9%	1.2%
	Sometimes	7.5%	6.0%
	Rarely	11.5%	15.7%
	Never	78.8%	76.5%
	I don't know	1.3%	0.6%
Interdental brush use			0.0001

Frequency of snacking	Often	0.4%	12.1%	0.24
	Sometimes	4.0%	23.0%	
	Rarely	12.0%	13.9%	
	Never	81.9%	48.5%	
	I don't know	1.8%	2.4%	
Fizzy drinks	Less than 3 times	2.2%	3.0%	0.04
	3 times	11.1%	11.5%	
	4 times	23.5%	33.1%	
	5 times	23.9%	18.7%	
	More than 5 times	39.4%	33.7%	
Sweets	Often	48.7%	39.8%	0.17
	Sometimes	38.5%	39.2%	
	Rarely	8.9%	18.1%	
	Never	4.0%	3.0%	
	I don't know	0.0%	0.0%	
Fast foods	Often	33.6%	24.7%	0.03
	Sometimes	38.9%	45.2%	
	Rarely	22.6%	27.1%	
	Never	4.9%	3.0%	
	I don't know	0.0%	0.0%	
Cigarettes	Often	25.8%	15.1%	0.002
	Sometimes	40.9%	42.2%	
	Rarely	29.3%	34.3%	
	Never	4.0%	8.4%	
	I don't know	0.0%	0.0%	
Drugs	Often	11.1%	4.2%	0.42
	Sometimes	6.7%	2.4%	
	Rarely	9.8%	4.8%	
	Never	72.4%	88.0%	
	I don't know	0.0%	0.6%	
Alcohol	Often	0.4%	1.2%	0.23
	Sometimes	3.5%	1.8%	
	Rarely	3.5%	1.8%	
	Never	92.5%	94.6%	
	I don't know	0.0%	0.6%	

Never	52.2%	62.1%
I don't know	0.0%	0.6%

(*) *p-values were calculated without considering the answer 'I don't know'*

3.4. Self-perception of oral and general health (Table 4)

Several situations and behaviours were significantly different between the two groups. Adolescents from the OT group were more likely to have difficulty chewing ($p=0.002$). When grouping 'rarely' and 'sometimes' together, adolescents from the OT group still had more difficulty eating ($p=0.02$) and suffered from toothache ($p=0.04$). The presence of orthodontic appliances was related to feeling embarrassed about his/her teeth ($p=0.01$). The perception of general, oral, or gingival health was comparable between the two groups.

Table 4. Self-perception of oral and general health according to orthodontic treatment.

Variable	No orthodontic treatment (n = 266) %	Orthodontic treatment (n=166) %	p-value (*) (chi-square)
Difficulty eating			0.06
Often	4.4	3.0	
Sometimes	11.1	13.3	
Rarely	23.0	33.7	
Never	61.5	49.4	
I don't know	0.0	0.6	
Difficulty chewing			0.002
Often	3.5	3.0	
Sometimes	9.4	14.5	
Rarely	21.0	34.9	
Never	66.1	47.0	
I don't know	0.0	0.6	
Toothache			0.11
Often	6.6	3.0	
Sometimes	17.3	24.7	
Rarely	34.5	36.8	
Never	41.2	35.5	
I don't know	0.4	0.0	
Embarrassed about their teeth			0.01
Often	8.1	10.3	
Sometimes	8.5	18.8	
Rarely	13.8	15.1	
Never	67.4	55.8	
I don't know	2.2	0.0	

(*) *p-values were calculated without considering the answer 'I don't know'*

Perceived oral health was linked to some functional difficulties such as toothache or difficulties chewing or talking. The proportion of participants who had difficulty talking increased

proportionally with their self-perceived bad oral health in both the NT group ($p=0.04$) and the OT group ($p=0.03$). The same relationship was found with difficulty chewing in the NT group ($p=0.001$) but not in the OT group ($p=0.10$). Adolescents who felt embarrassed with the appearance of their teeth were those reporting worse self-perceived oral health, in both the NT group ($p=0.0001$) and in the OT group ($p=0.04$). Talking difficulties were also related to self-perceived oral health in the NT group ($p=0.004$) but not in the OT group ($p=0.87$). Adolescents having had toothache were those reporting bad self-perceived oral health in the NT group ($p=0.0001$), but not in the OT group ($p=0.09$). The relationship between bleeding gums and self-perceived oral health was significant in the NT group ($p=0.0001$) but not in the OT group ($p=0.24$).

4. Discussion

Studies on knowledge, attitudes and practices have been conducted at international level in the field of public health for many years. This study, to our knowledge, represents the first study in France of this type which compares two groups of adolescents with or without orthodontic treatment.

We know today that adolescence is a critical period of transition where risky behaviours appear and that adolescent optimism is associated with positive health outcomes [38].

Orthodontists give oral health hygiene and dietary instructions before starting a treatment. We hypothesised that adolescents undergoing orthodontic treatment have better oral health knowledge, attitudes, and practices.

In this study, the knowledge of the adolescents was insufficient in both groups; except for the question about the role of fluoride in the prevention of caries, as adolescents undergoing treatment had a significantly higher percentage of correct answers ($p=0.0001$).

96.5% of adolescents without orthodontic treatment and 95.1% of adolescents with treatment knew that it is necessary to brush their teeth at least twice a day. In a study by Wahengbam et al. de 2016, adolescents' knowledge about brushing twice a day was 86.4% [34].

Most participants reported having correct knowledge about the impact of snacking (78.5%) and sticky foods (94.7%) on the risk of caries. There was no significant difference between the two groups. Studies had similar results; 91.9% of adolescents knew about the role of diet in tooth decay in India (35–37)[34], and this percentage was 86.3% in a recent study among adolescents from Portugal, Romania and Sweden [35].

Adolescents' knowledge of fluoride was incorrect in both of the groups. However, adolescents with treatment had better results, mainly regarding the role of fluoride in the prevention of caries with 65.1% of correct answers, as opposed to adolescents without treatment, who had 46% correct answers ($p=0.0001$). Indeed, some studies had similar results in adolescents [33,35,36,39–41]. Adolescents undergoing orthodontic treatment had a better understanding of the role of fluoride.

Most adolescents in both groups (69%) claimed to brush their teeth twice a day. Adolescents undergoing orthodontic treatment brushed more. In the literature, the frequency of brushing in adolescents showed different results, from 40% [33] to 77.4% [35].

Adolescents undergoing orthodontic treatment use interdental brushes significantly more than other adolescents (35.1% vs 4.4%) ($p=0.0001$). 20% of orthodontic patients in the study of R. Aljohani et al. used an interdental brush at least once daily [42]. Previous studies reported an infrequent use of interdental brushes (7–23%) among orthodontic patients [43–45]. A recent study carried out among adolescents aged 15 to 17 years old highlighted an average interdental inflammation of 96.5%. We now know that interdental spaces harbour pathogenic bacteria and that daily brushing reduces bleeding by 47% after 1 week of use and that symbiosis can be restored.

Our results show that very few adolescents did not consume fizzy drinks (3.5%). Many adolescents consumed fizzy drinks (48.7% and 39.8%). But adolescents undergoing orthodontic treatment consumed significantly less ($p=0.04$). In R. Graça et al.'s study, this concerned 35.5% of adolescents [35] and 56% in the study of R. Aljohani et al. [43]. 25.8% and 15.1% of adolescents - with and without orthodontic treatment, respectively - consumed fast-food ($p=0.03$). For sweets, there were no significant differences between the two groups (33.6% and 24.7%), even though

adolescents without orthodontic treatment tended to consume sweets more often. Results of between 40 % and 60% can be found in the literature [35,36,42].

With regards to the use of drugs and alcohol, the results are similar in both groups. On the contrary, cigarette consumption is significantly higher among adolescents who do not have orthodontic treatment ($p=0.002$). 11.1% of adolescents without treatment and 4.2% of adolescents with treatment smoke on a regular basis. Smoking daily or occasionally was reported in 11.7% of adolescents in the study of T.Graça et al. [35].

5. Conclusions

Adolescents lack oral health knowledge, and their attitudes and practices are not optimal. We know adolescence is an important period for the development of oral pathologies, and that this period is associated with hormonal changes, behavioural shifts, and changing eating habits. Adolescents start becoming more autonomous by making their own decisions. Orthodontists should be more aware of the need to teach their patients how to maintain good oral hygiene and have good dietary habits during orthodontic treatment to prevent cavities and periodontal disease. Oral health education is essential for all adolescents and the use of interdental brushes should be implemented.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, S1: questionnaire.

Author Contributions: C.I., D.B., F.C., C.C., A.C. participated in the study design conceptualization, supervision, D.B., P.T. involved in the methodology, F.C. validation, C.I., P.T., F.C. contributed the figures drawing, and F.C. analysis, editing, and proofreading. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Petersen, P.E. The World Oral Health Report 2003: Continuous Improvement of Oral Health in the 21st Century--the Approach of the WHO Global Oral Health Programme. *Community Dent. Oral Epidemiol.* **2003**, *31*, 3–23, doi:10.1046/j.2003.com122.x.
2. Jackson, S.L.; Vann, W.F.; Kotch, J.B.; Pahel, B.T.; Lee, J.Y. Impact of Poor Oral Health on Children's School Attendance and Performance. *Am. J. Public Health* **2011**, *101*, 1900–1906, doi:10.2105/AJPH.2010.200915.
3. Kassebaum, N.J.; Bernabé, E.; Dahiya, M.; Bhandari, B.; Murray, C.J.L.; Marcenes, W. Global Burden of Untreated Caries: A Systematic Review and Metaregression. *J. Dent. Res.* **2015**, *94*, 650–658, doi:10.1177/0022034515573272.
4. Califano, J.V.; AAP; AAPD Periodontal Diseases of Children and Adolescents. *Pediatr. Dent.* **2005**, *27*, 189–196.
5. Coolidge, T.; Heima, M.; Johnson, E.K.; Weinstein, P. The Dental Neglect Scale in Adolescents. *BMC Oral Health* **2009**, *9*, 2, doi:10.1186/1472-6831-9-2.
6. Al-Ghutaimel, H.; Riba, H.; Al-Kahtani, S.; Al-Duhaimi, S. Common Periodontal Diseases of Children and Adolescents. *Int. J. Dent.* **2014**, doi:10.1155/2014/850674.
7. Agustsdottir, H.; Gudmundsdottir, H.; Eggertsson, H.; Jonsson, S.H.; Gudlaugsson, J.O.; Saemundsson, S.R.; Eliasson, S.T.; Arnadottir, I.B.; Holbrook, W.P. Caries Prevalence of Permanent Teeth: A National Survey of Children in Iceland Using ICDA. *Community Dent. Oral Epidemiol.* **2010**, *38*, 299–309, doi:10.1111/j.1600-0528.2010.00538.x.
8. Fleming, E.; Afful, J. Prevalence of Total and Untreated Dental Caries Among Youth: United States, 2015-2016. *NCHS Data Brief* **2018**, 1–8.

9. Olczak-Kowalczyk, D.; Gozdowski, D.; Kaczmarek, U. Oral Health in Polish Fifteen-Year-Old Adolescents. *Oral Health Prev. Dent.* **2019**, *17*, 139–146, doi:10.3290/j.ohpd.a42373.
10. Jordão, L.M.R.; Vasconcelos, D.N.; Moreira, R. da S.; Freire, M. do C.M. Individual and Contextual Determinants of Periodontal Health in 12-Year-Old Schoolchildren in a Brazilian Capital City. *Int. J. Dent.* **2012**, *2012*, 325475, doi:10.1155/2012/325475.
11. Das, U.M.; Vadakkettical, R.J.; Kanakkath, H.; Shankunni, S.P. Dental Health Awareness, Attitude, and Dental Health-Care Seeking Practices as Risk Indicators for the Prevalence of Periodontal Disease among 15–17-Year-Old School Children in Kozhikode District, Kerala, India. *J. Indian Soc. Periodontol.* **2017**, *21*, 144–151, doi:10.4103/jisp.jisp_160_17.
12. Albandar, J.M.; Rams, T.E. Global Epidemiology of Periodontal Diseases: An Overview. *Periodontol.* **2000**, *29*, 7–10.
13. Brukiene, V.; Aleksejuniene, J. Theory-Based Oral Health Education in Adolescents. *Stomatologija* **2010**, *12*, 3–9.
14. Silk, H.; Kwok, A. Addressing Adolescent Oral Health: A Review. *Pediatr. Rev.* **2017**, *38*, 61–68, doi:10.1542/pir.2016-0134.
15. Graves, D.T.; Corrêa, J.D.; Silva, T.A. The Oral Microbiota Is Modified by Systemic Diseases. *J. Dent. Res.* **2019**, *98*, 148–156, doi:10.1177/0022034518805739.
16. Hajishengallis, G.; Lamont, R.J. Beyond the Red Complex and into More Complexity: The Polymicrobial Synergy and Dysbiosis (PSD) Model of Periodontal Disease Etiology. *Mol. Oral Microbiol.* **2012**, *27*, 409–419, doi:10.1111/j.2041-1014.2012.00663.x.
17. Lamont, R.J.; Hajishengallis, G. Polymicrobial Synergy and Dysbiosis in Inflammatory Disease. *Trends Mol. Med.* **2015**, *21*, 172–183, doi:10.1016/j.molmed.2014.11.004.
18. Tanner, A.C.R.; Kressirer, C.A.; Rothmiller, S.; Johansson, I.; Chalmers, N.I. The Caries Microbiome: Implications for Reversing Dysbiosis. *Adv. Dent. Res.* **2018**, *29*, 78–85, doi:10.1177/0022034517736496.
19. Hajishengallis, G. Immunomicrobial Pathogenesis of Periodontitis: Keystones, Pathobionts, and Host Response. *Trends Immunol.* **2014**, *35*, 3–11, doi:10.1016/j.it.2013.09.001.
20. Inquimbert, C.; Bourgeois, D.; Bravo, M.; Viennot, S.; Tramini, P.; Llodra, J.C.; Molinari, N.; Dussart, C.; Giraudeau, N.; Carrouel, F. The Oral Bacterial Microbiome of Interdental Surfaces in Adolescents According to Carious Risk. *Microorganisms* **2019**, *7*, doi:10.3390/microorganisms7090319.
21. Tarannum, F.; Faizuddin, M.; Swamy, S.; Hemalata, M. Efficacy of a New Interdental Cleaning Aid. *J. Indian Soc. Periodontol.* **2012**, *16*, 375–380, doi:10.4103/0972-124X.100914.
22. Ng, E.; Lim, L.P. An Overview of Different Interdental Cleaning Aids and Their Effectiveness. *Dent. J.* **2019**, *7*, doi:10.3390/dj7020056.
23. Bourgeois, D.; Saliassi, I.; Llodra, J.C.; Bravo, M.; Viennot, S.; Carrouel, F. Efficacy of Interdental Calibrated Brushes on Bleeding Reduction in Adults: A 3-Month Randomized Controlled Clinical Trial. *Eur. J. Oral Sci.* **2016**, *124*, 566–571, doi:10.1111/eos.12302.
24. Bourgeois, D.; Bravo, M.; Llodra, J.-C.; Inquimbert, C.; Viennot, S.; Dussart, C.; Carrouel, F. Calibrated Interdental Brushing for the Prevention of Periodontal Pathogens Infection in Young Adults - a Randomized Controlled Clinical Trial. *Sci. Rep.* **2019**, *9*, 15127, doi:10.1038/s41598-019-51938-8.
25. Baskaradoss, J.K. Relationship between Oral Health Literacy and Oral Health Status. *BMC Oral Health* **2018**, *18*, doi:10.1186/s12903-018-0640-1.
26. Berkman, N.D.; Davis, T.C.; McCormack, L. Health Literacy: What Is It? *J. Health Commun.* **2010**, *15 Suppl 2*, 9–19, doi:10.1080/10810730.2010.499985.
27. Berkman, N.D.; Sheridan, S.L.; Donahue, K.E.; Halpern, D.J.; Crotty, K. Low Health Literacy and Health Outcomes: An Updated Systematic Review. *Ann. Intern. Med.* **2011**, *155*, 97–107, doi:10.7326/0003-4819-155-2-201107190-00005.
28. Paulander, J.; Axelsson, P.; Lindhe, J. Association between Level of Education and Oral Health Status in 35-, 50-, 65- and

75-Year-Olds. *J. Clin. Periodontol.* **2003**, *30*, 697–704.

29. Batista, M.J.; Lawrence, H.P.; Sousa, M. da L.R. de Oral Health Literacy and Oral Health Outcomes in an Adult Population in Brazil. *BMC Public Health* **2017**, *18*, 60, doi:10.1186/s12889-017-4443-0.
30. Das, D.; Menon, I.; Gupta, R.; Arora, V.; Ashraf, A.; Ahsan, I. Oral Health Literacy: A Practical Strategy towards Better Oral Health Status among Adult Population of Ghaziabad District. *J. Fam. Med. Prim. Care* **2020**, *9*, 764, doi:10.4103/jfmpejfmpej_1049_19.
31. El Osta, N.; Pichot, H.; Soulier-Peigue, D.; Hennequin, M.; Tubert-Jeannin, S. Validation of the Child Oral Health Impact Profile (COHIP) French Questionnaire among 12 Years-Old Children in New Caledonia. *Health Qual. Life Outcomes* **2015**, *13*, 176, doi:10.1186/s12955-015-0371-9.
32. Himida, T.; Promise, U. School-Based Dental Sealant Programmes May Be Effective in Caries Prevention. *Evid. Based Dent.* **2017**, *18*, 13–14, doi:10.1038/sj.ebd.6401218.
33. Blaggana, A.; Grover, V.; Anjali; Kapoor, A.; Blaggana, V.; Tanwar, R.; Kaur, H.; Haneet, R.K. Oral Health Knowledge, Attitudes and Practice Behaviour among Secondary School Children in Chandigarh. *J. Clin. Diagn. Res. JCDR* **2016**, *10*, ZC01–ZC06, doi:10.7860/JCDR/2016/23640.8633.
34. Wahengbam, P.P.; Kshetrimayum, N.; Wahengbam, B.S.; Nandkeoliar, T.; Lyngdoh, D. Assessment of Oral Health Knowledge, Attitude and Self-Care Practice Among Adolescents - A Statewide Cross- Sectional Study in Manipur, Northeastern India. *J. Clin. Diagn. Res. JCDR* **2016**, *10*, ZC65–ZC70, doi:10.7860/JCDR/2016/20693.8002.
35. Graça, S.R.; Albuquerque, T.S.; Luis, H.S.; Assunção, V.A.; Malmqvist, S.; Cuculescu, M.; Slusanschi, O.; Johannsen, G.; Galuscan, A.; Podariu, A.C.; et al. Oral Health Knowledge, Perceptions, and Habits of Adolescents from Portugal, Romania, and Sweden: A Comparative Study. *J. Int. Soc. Prev. Community Dent.* **2019**, *9*, 470–480, doi:10.4103/jispcd.JISPCD_194_19.
36. Al-Qahtani, S.M.; Razak, P.A.; Khan, S.D. Knowledge and Practice of Preventive Measures for Oral Health Care among Male Intermediate Schoolchildren in Abha, Saudi Arabia. *Int. J. Environ. Res. Public Health* **2020**, *17*, doi:10.3390/ijerph17030703.
37. Nihtila, A.; West, N.; Lussi, A.; Bouchard, P.; Ottolenghi, L.; Senekola, E.; Llodra, J.C.; Viennot, S.; Bourgeois, D. Oral Health Behavior and Lifestyle Factors among Overweight and Non-Overweight Young Adults in Europe: A Cross-Sectional Questionnaire Study. *Healthcare* **2016**, *4*, doi:10.3390/healthcare4020021.
38. Malinowska-Cieślak, M.; Mazur, J.; Nałęcz, H.; Małkowska-Szkućnik, A. Social and Behavioral Predictors of Adolescents' Positive Attitude towards Life and Self. *Int. J. Environ. Res. Public Health* **2019**, *16*, doi:10.3390/ijerph16224404.
39. Prasai Dixit, L.; Shakya, A.; Shrestha, M.; Shrestha, A. Dental Caries Prevalence, Oral Health Knowledge and Practice among Indigenous Chepang School Children of Nepal. *BMC Oral Health* **2013**, *13*, 20, doi:10.1186/1472-6831-13-20.
40. Jensen, O.; Gabre, P.; Sköld, U.M.; Birkhed, D. Fluoride Toothpaste and Toothbrushing: Knowledge, Attitudes and Behaviour among Swedish Adolescents and Adults. *Swed. Dent. J.* **2011**, *35*, 203–213.
41. Al-Darwish, M.S. Oral Health Knowledge, Behaviour and Practices among School Children in Qatar. *Dent. Res. J.* **2016**, *13*, 342–353.
42. Aljohani, S.R.; Alsaggaf, D.H. Adherence to Dietary Advice and Oral Hygiene Practices Among Orthodontic Patients. *Patient Prefer. Adherence* **2020**, *14*, 1991–2000, doi:10.2147/PPA.S277034.
43. Pandey, V.; Chandra, S.; Dilip Kumar, H.P.; Gupta, A.; Bhandari, P.P.; Rathod, P. Impact of Dental Neglect Score on Oral Health among Patients Receiving Fixed Orthodontic Treatment: A Cross-Sectional Study. *J. Int. Soc. Prev. Community Dent.* **2016**, *6*, 120–124, doi:10.4103/2231-0762.178752.
44. Anuwongnukroh, N.; Dechkunakorn, S.; R, K. Oral Hygiene Behavior during Fixed Orthodontic Treatment. *Dentistry* **2017**, *7*, doi:10.4172/2161-1122.1000457.
45. Choi, Y.-J.; Choi, J.-O. The Effects of the Oral Care Behaviors of Adolescents on Self-Perceived Oral Health. *Curr. Pediatr. Res.* **2017**, *21*, 553–558.