

## Article

# Geochemical Association Rules of Elements Mined using Clustered Events of Spatial Auto-correlation: A Case Study in the Chahanwusu River Area, Qinghai Province, China

Baoyi Zhang <sup>1,2</sup>, Zhengwen Jiang <sup>2</sup>, Yiru Chen <sup>2</sup>, Nanwei Cheng <sup>2</sup>, Umair Khan <sup>2</sup> and Jiqui Deng <sup>1,2,\*</sup>

<sup>1</sup> Key Laboratory of Metallogenic Prediction of Nonferrous Metals and Geological Environment Monitoring (Ministry of Education), Central South University, Changsha 410083, China; zhangbaoyi@csu.edu.cn (B.Z.)

<sup>2</sup> School of Geosciences and Info-Physics, Central South University, Changsha 410083, China; jiang-zhengwen@csu.edu.cn (Z.J.); 195011096@csu.edu.cn (Y.C.); 215011050@csu.edu.cn (N.C.); Umair77@csu.edu.cn (U.K.)

\* Correspondence: csugis@csu.edu.cn; Tel.: +86-731-88877676 (J.D.)

**Abstract:** The spatial distribution of elements can be regarded as a numerical field of concentration values with a continuous spatial coverage. An active area of research is to discover geologically meaningful relationships among elements from their spatial distribution. To solve this problem, we propose an association rule mining method based on clustered events of spatial auto-correlation and applied it to the polymetallic deposits of the Chahanwusu River area, Qinghai Province, China. The elemental data for stream sediments were first clustered into HH (high-high), LL (low-low), HL (high-low), and LH (low-high) groups by using local Moran's I clustering map (LMIC). Then the Apriori algorithm was used to mine the association rules among different elements in these clusters. More than 86% of the mined rule points are located within 1000 m of faults and near known ore occurrences, and occur in the upper reaches of the stream and catchment areas. In addition, we found that the Indosinian granodiorite is enriched in sulfophile elements, e.g., Zn, Ag and Cd, and the Variscan granite quartz diorite (P<sub>1</sub>γδo) coexists with Cu and associated elements. Therefore, the proposed algorithm is an effective method for mining co-existence patterns of elements and provides an insight into their enrichment mechanisms.

**Keywords:** Concentration field; Spatial auto-correlation; Association rules; Apriori algorithm; Element co-occurrence

## 1. Introduction

Spatial auto-correlation analysis focuses on the similarity of attributes as well as spatial similarity between one geological entity and adjacent entities. The spatial distribution of concentrations of elements can be regarded as a numerical field with a continuous spatial coverage, which can be characterized by using the spatial auto-correlation among different elements. Korobova and Romanov (2009) stressed that the nonrandom characteristics and spatial structure of geochemical data depend on the concentration field [1]. The analysis of the concentration field includes comparing samples to recognize anomalies and using the spatial correlation among elements to explain geochemical processes. Geological interactions between elements result in mutual influence and restriction. Therefore, it is necessary to consider spatial auto- and cross-correlation in geochemical studies. The concentrations and spatial association of different elements are usually related to parent lithostrata. Therefore, it is of great significance to study the distribution, enrichment, and relationships among different elements to understand regional magmatism and ore-forming process [2].

Tobler (1970) proposed the first law of geography: everything is related to everything else, but near things are more related to distant things [3]. The measurement of spatial

auto-correlation includes global and local indicators. The global indicators reveal the spatial pattern of the whole region and reflect the global characteristics. In contrast, local indicators measure the relationship between each location and its neighbors to reveal more detailed local spatial patterns. Global metrics include Moran's I [4] and Geary's C [5]. Improvements in spatial theory and statistical tests have made Moran's I and Geary's C become the most widely used global indicators [6-8]. Based on Moran's I, Cliff and Ord (1981) also proposed a simple spatio-temporal auto-correlation indicator form  $I_{s-t}$  [8]. Getis and Ord (1992, 1995) proposed global G statistic and local G\* statistic [9,10]. Anselin (1995) developed local indicators of spatial association (LISA), including local Moran's I and local Geary's C [11]. Boots and Okabe (2007) proposed the concept of local spatial statistical analysis (LoSSA) both as an integrative structure for existing methods and as a framework that facilitates the development of new local and global statistics [12]. Anselin (2019) extended the application of the local Geary's C statistic to a multivariate context. According to the characteristics of experimental data, each local auto-correlation indicator has its advantages and disadvantages [13]. Spatial auto-correlation indicators have been used in the fields of environmental science, regional economy, identification of diseases and mortality, and detection of geochemical anomalies [14-18].

The spatial pattern of the concentration field is caused by different geological processes [19]. The concentration field reflects the migration of elements and the spatio-temporal distribution of various elements. Therefore, both the spatial characteristics of a single element and the spatial relationship among multiple elements need to be considered.

For a long time, the identification and evaluation of geochemical anomalies has been a key issue in the field of geochemical exploration [20-22]. A geochemical anomaly is the enrichment or dilution of elements. The enriched area often has high mineral resource potential [23,24]. Geologists seek to use the spatial pattern to distinguish an anomaly from the background. For many years, various statistical methods, such as mean  $\pm 2 \times$  standard deviations [25], probability graphs [26], univariate analysis [27], multivariate analysis [28,29], logistic regression [30,31], weights of evidence [32-34], fractal/multifractal models [35-37], and geostatistics [38,39] have been used to identify geochemical anomalies. In recent years, machine learning methods have been used in geological prospecting. These methods include support vector machines [40,41], random forests [42,43], Bayesian networks [44-46], and deep autoencoder networks [47].

Some small ore deposits or occurrences are overlooked in actual mineral prospecting if the association rules among elements are not considered [19]. How to efficiently delineate the metallogenic target area has become one of the main objectives of geochemical exploration. Nguyen et al. (2014) found that local Moran's I could better detect the spatial clustering of elements in stream sediments on a small spatial scale than classical statistics, and local G\* is suitable for detecting high clustered on a large scale [48]. Wang et al. (2015) used geostatistics, fractal and spatial auto-correlation methods to study the spatial characteristics of geochemical data for stream sediments in southwest Fujian, and concluded that the spatial auto-correlation method delineates the geochemical anomaly [49]. Ji et al. (2017) used local Moran's I to analyze the spatial clustering and outliers of elemental concentrations and extracted geochemical anomalies [50]. Yu et al. (2021) proposed a local correlation coefficient based on spatial neighborhoods to characterize the global distribution of elements [16].

The mutual influence and interaction among different elements produce a spatial pattern [51,52]. The effects of regional geological and geochemical processes can be inferred from the spatial patterns in the concentration field. Therefore, exploring the association rules among different elements is of great significance for understanding geological processes. Association rule mining is one of the branch fields of data mining. The Apriori algorithm can uncover Boolean association rules between itemsets and has been widely used in spatial data mining [53-56]. The Apriori algorithm was proposed by Agrawal et al. (1993), who used it to mine association rules of sales data obtained from a large retailing company [57]. Liu and Zhou (2019) used the Apriori algorithm to derive the anomalies of elements for metallogenic prediction [58].

In this paper, we proposed an association rule mining method to study the cross-correlation of concentration fields based on clustered events of spatial auto-correlation. This method can be used to comprehensively understand the spatial distribution of geochemical concentrations and co-existing of elements. Moreover, we compared the advantages and limitations of bivariate spatial auto-correlation and association rule mining results, and finally explored the relationship of specific geological features with the results of association rule mining.

2. Study area and data

2.1. Geological background

The Chahanwusu River area (98°15'E-98°45'E, 35°50'N-36°00'N) covers approximately 893 km<sup>2</sup> in the eastern part of the East Kunlun tectonic belt in Dulan County in central Qinghai Province. The area is a polymetallic belt where one gold deposit, three copper deposits, one lead-zinc deposit two magnetite deposits, and one gem-stone deposit have been found [59]. Figure 1 shows the sketch geological map of the study area [43].

The main faults in the study area are EW, NW, and NE-trending, which constitute the structural framework of the area. NW-trending faults are the most developed, and control the distribution of strata and magmatic rocks. The sedimentary strata in the study area are undeveloped and dispersed. The outcropping strata from old to new are the Paleoproterozoic Baishahe rock Formation (Pt**b**), the late Triassic Elashan Formation (T**3e**), the Neogene Guide Group (NG) and Quaternary (Q). Outcrops of intrusive rocks are widespread in the study area, and are dominated by Variscan and Indosinian intrusives.




Figure 1. Geological map of the study area [43].


2.2. Geochemical data

The datasets used in this study were geochemical analyses of 4959 stream sediment samples taken at a density of 5.55 points per 1 km<sup>2</sup> by the Geological Survey Institute of



Qinghai Province (Figure 2). The concentrations of 15 elements (Au, Sn, Ag, As, Sb, Bi, Co, Cu, La, Pb, Zn, W, Mo, Nb, and Cd) were measured in each sample. The methods used to analyze the concentration of heavy metals include atomic emission spectrometry (AES) for Au, Ag, and Sn, atomic fluorescence spectrometry (AFS) for As, Sb, and Bi, atomic absorption spectrometry (AAS) for Cu, Pb, Zn, Co, and Ni, and polarography (POL) for W and Mo.

The elemental concentrations are summarized in Table 1. The coefficient of variation (CV) is expressed as the ratio of the standard deviation to the mean and is an important parameter that reflects the homogenization of element distribution. The elements with the  $CV > 1$  from largest to smallest are Bi, W, Sb, As, Ag, Sn, Au, Cu, Pb, Mo. Larger CV represents more inhomogeneous elemental concentrations. The higher the coefficient of variation, the greater the level of dispersion around the mean.



**Figure 2.** Map of stream sediment geochemical sampling points.

**Table 1.** Main statistical results of the stream sediment elements in the study area.

Element	Mean	Median	Standard Deviation	Skewness	Kurtosis	Coefficient of Variation
Au	1.42	1.20	1.81	25.07	793.15	1.27
Sn	2.31	1.70	3.21	10.35	134.68	1.39
Ag	76.31	41.00	117.35	6.77	67.02	1.54
As	13.07	8.10	21.89	13.21	303.48	1.68
Sb	0.82	0.47	1.45	14.56	372.59	1.88
Bi	0.37	0.17	1.00	13.68	256.65	2.50
Co	7.07	6.40	3.44	3.07	24.09	0.48
Cu	16.05	12.10	20.44	15.15	330.62	1.27
La	13.73	12.00	8.30	8.99	192.65	0.61
Pb	17.00	12.80	21.29	13.35	314.69	1.25
Zn	48.29	40.40	32.06	4.27	27.61	0.66
W	2.74	1.70	5.99	18.99	528.95	2.22
Mo	1.20	0.96	1.36	9.31	117.22	1.17
Nb	3.93	3.30	2.26	4.53	37.90	0.59
Cd	0.15	0.10	0.18	5.28	49.30	0.85

Au、Ag:10<sup>-9</sup>, others:10<sup>-6</sup>.

### 3. Methods

#### 3.1. Spatial auto-correlation

##### 3.1.1. Univariate spatial auto-correlation

Spatial auto-correlation indicates the extent to which one attribute of a feature is influenced by nearby features [60]. Spatial auto-correlation indicators are the sum of the cross product of a similarity matrix  $c_{ij}$  and spatial similarity matrix  $w_{ij}$ , and include global (Eq. 1) and local (Eq. 2) metric indicators [11]. In general form, they are written as

$$\Gamma_g = \sum_{i=1}^n \sum_{j=1}^n c_{ij} w_{ij} \tag{1}$$

$$\Gamma(i) = \sum_{j=1}^n c_{ij} w_{ij} \tag{2}$$

where  $n$  is the total numbers of observations,  $c_{ij}$  is the self-similarity matrix, and  $w_{ij}$  is the spatially weighted matrix.

Global indicators give the degree of spatial association for a single value, and local indicators assess the extent to which observations of similar and dissimilar values are clustered for each location [11]. Different measures of similarity yield different indices for spatial association [11]. For example, using  $c_{ij} = (x_i - \bar{x})(x_j - \bar{x})$  yields a Moran-like indicator, setting  $c_{ij} = (x_i - x_j)^2$  yields a Geary-like indicator, and setting  $c_{ij} = x_i x_j$  yields a Getis–Ord-like indicator. The corresponding spatial auto-correlation indicators are global Moran’s I [4,6-8], Geary’s C [5,6,8], and Getis–Ord’s G [9], respectively. A global spatial auto-correlation indicator can only reflect the overall spatial trend and auto-correlation of the geographical entity or phenomenon. However, local spatial auto-correlation indicators measure the correlation among various locations and their neighbors to reveal more detailed local spatial patterns. These indicators include local Moran’s I [11], Geary’s C [11], and Getis–Ord’s G [9]. The calculation method of univariate global and local spatial auto-correlation statistics is shown in Table 2.

Table 2. Spatial auto-correlation statistics.

Spatial auto-correlation statistics	Calculation formula	Remarks	References
global Moran’s I	$I = \frac{n \sum_i^n \sum_j^n w_{ij} (x_i - \bar{x})(x_j - \bar{x})}{(\sum_i^n \sum_j^n w_{ij}) \sum_i^n (x_i - \bar{x})^2}$	The range of $I$ is [-1,1], $I < 0$ indicates negative spatial auto-correlation, $I > 0$ indicates positive spatial auto-correlation, $I$ tends to 0 indicates spatial random distribution.	[4,6-8]
global Geary’s C	$C = \frac{(n-1) \sum_i^n \sum_j^n w_{ij} (x_i - x_j)^2}{(2 \sum_i^n \sum_j^n w_{ij}) \sum_i^n (x_i - \bar{x})^2}$	The range of $C$ is [0,2], $C > 1$ indicates negative spatial auto-correlation, $C < 1$ indicates positive spatial auto-correlation, $C$ tends to 1 indicates spatial random distribution.	[5,6,8]
global Getis–Ord’s G	$G = \frac{\sum_i^n \sum_j^n w_{ij} x_i x_j}{\sum_i^n \sum_j^n x_i x_j}$	$G <$ mathematical expectation (ME) indicates low value clustered, $G >$ ME indicates high value clustered, $G$ tends to ME indicates spatial random distribution.	[9]
local Moran’s I	$I(i) = \frac{(x_i - \bar{x})}{S^2} \sum_j^n w_{ij} (x_j - \bar{x})$ $S^2 = \frac{\sum_i^n (x_i - \bar{x})^2}{n}$	$Z(I(i)) < 0$ indicates negative spatial auto-correlation, $Z(I(i)) > 0$ indicates positive spatial auto-correlation,	[9]

		$Z(I(i))$ tends to 0 indicates spatial random distribution.
local Geary's C	$C(i) = \frac{1}{S^2} \sum_j^n w_{ij} (x_i - x_j)^2$ $S^2 = \frac{\sum_i^n (x_i - \bar{x})^2}{n}$	$Z(C(i)) < 0$ indicates negative spatial auto-correlation, $Z(C(i)) > 0$ indicates positive spatial auto-correlation,
		$Z(C(i))$ tends to 0 indicates spatial random distribution. [11]
local Getis-Ord's G	$G(i) = (\sum_j^n w_{ij} x_j) / \sum_j^n x_j$	$Z(G(i)) < 0$ indicates negative spatial auto-correlation, $Z(G(i)) > 0$ indicates positive spatial auto-correlation,
		$Z(G(i))$ tends to 0 indicates spatial random distribution. [9]

$x_i$  and  $x_j$  is the observed value at positions  $i$  and  $j$ ,  $x_j$  is  $x_i$  position's neighbor point in a certain distance,  $n$  is the total numbers of observations,  $\bar{x}$  is the mean value of the observations,  $w_{ij}$  is the spatial weight matrix.  $Z(\Gamma) = (\Gamma - E(\Gamma)) / \sqrt{VAR(\Gamma)}$ ,  $\Gamma$  is a spatial statistic,  $E(\Gamma)$  is the mathematical expectation of  $\Gamma$ ,  $VAR(\Gamma)$  is the variance of  $\Gamma$ .

To take an example, given the spatial distribution of points in Figure 3a, we obtain a distance matrix (Figure 3b) between point pairs, then we get a spatially weighted matrix (Figure 3c) between point pairs according to whether their distance is less than 4 units. According to Figure 3, we find that the global Moran's I value is -0.052, the p-value is 0.057, and the local Moran's I indicators are as listed in Table 3.




Figure 3. (a) Spatial distribution of points, (b) distance matrix between point pairs, and (c) spatially weighted matrix.

Table 3. Local Moran's I calculation results of the example in Figure 3.

ID	X	Y	Value	Local Moran's I	Clustered Type	P-value
P1	1	4	0.2	-0.1758203	Low-High	0.001
P2	1	3	0.4	-0.0028961	Low-High	0.001
P3	1	1	0.8	-0.323655	Insignificant	0.382
P4	2	1	0.5	-0.0335362	Insignificant	0.134
P5	3	2	0.1	-0.1007592	Low-High	0.001
P6	4	3	0.6	0.0541114	High-High	0.001
P7	3	4	0.4	-0.0111241	Low-High	0.001
P8	4	5	0.3	0.2280927	Low-Low	0.001

### 3.1.2. Multivariate spatial cross-correlation

Spatial cross-correlation indicates the extent to which the multiple attributes of a feature are cross-influenced by nearby features. The exploration of multivariate spatial cross-correlation is a core functionality of current exploratory data analysis (EDA), knowledge discovery and data mining tools [61]. Anselin et al. (2002) proposed bivariate global (Eq. 3) and local (Eq. 4) Moran's I to quantify bivariate spatial cross-correlation [62]. They are calculated from

$$I_{ab} = \frac{n \sum_i^n \sum_j^n w_{ij} (a_i - \bar{a})(b_j - \bar{b})}{(\sum_i^n \sum_j^n w_{ij}) \sum_i^n (a_i - \bar{a})^2} \quad (3)$$

$$I_{ab}(i) = \frac{n(a_i - \bar{a})}{\sum_i^n (a_i - \bar{a})^2} \sum_j^n w_{ij} (b_j - \bar{b}) \quad (4)$$

where  $a_i$  and  $b_j$  are the observed values of variables  $a$  and  $b$  at positions  $i$  and  $j$ ,  $n$  is the total number of observations,  $\bar{a}$  and  $\bar{b}$  are the mean values of the observations of variables  $a$  and  $b$ , and  $w_{ij}$  is the spatial weighted matrix.

Anselin (2019) proposed using the univariate local Geary's C to measure the squared distance in attribute space (i.e., along a line for the univariate case) between the values at a geographic location and its neighboring locations, which is summarized in the form of a weighted sum [13]. This indicator can be readily extended to a multivariate context. For example, consider two variables,  $p$  and  $q$ . The squared distance  $d_{ij}^2$  in two-dimensional attribute space between the values at observation  $i$  and its geographic neighbor  $j$  is:

$$d_{ij}^2 = (p_i - p_j)^2 + (q_i - q_j)^2 \quad (5)$$

The bivariate local Geary's C can be defined as:

$$c_{ab}(i) = \frac{1}{2} \sum_j w_{ij} d_{ij}^2 = \frac{1}{2} \sum_j w_{ij} [(p_i - p_j)^2 + (q_i - q_j)^2] = \frac{1}{2} [\sum_j w_{ij} (p_i - p_j)^2 + \sum_j w_{ij} (q_i - q_j)^2] = \frac{1}{2} (c_a(i) + c_b(i)) \quad (6)$$

where  $p_i$  and  $p_j$  are the observed values of variable  $a$  at positions  $i$  and  $j$ ,  $q_i$  and  $q_j$  are the observed values of variable  $b$  at positions  $i$  and  $j$ , and  $w_{ij}$  is the spatial weighted matrix.

Following standard practice in multivariate clustering analysis, these variables have been standardized such that the mean of the transformed variable is zero and its variance is one. Moreover, the concept of a local Geary's C is additive in the attribute dimension. Therefore, a multivariate local Geary's C can be defined as:

$$c_{total}(i) = \sum_{v=1}^k c_v(i) / k \quad (7)$$

where  $k$  represents  $k$ -dimensional attribute space, and  $c_v(i)$  represents the univariate local Geary's C of variable  $v$ .

### 3.2. Association rule mining and Apriori algorithm

Association rule mining is used to reveal the association among items in a dataset. We assume that  $D = \{t_1, t_2, \dots, t_N\}$  is the event dataset,  $t_k = \{i_1, i_2, \dots, i_K\}$  represents an event corresponding to a geochemical sample, and  $i_k$  represents an item belonging to an aggregated event  $t_k$ . Itemset  $I = \{i_1, i_2, \dots, i_M\}$  is a specific item combination, which contains  $M$  different items. For a subset  $X$  in  $I$ , if  $X \subseteq t_k$ , then the event  $t_k$  contains  $X$ . The goal of association rule mining is to find an implicit form of  $X \Rightarrow Y$ , where  $X \subseteq I$ ,  $Y \subseteq I$ , and  $X \cap Y = \emptyset$ . If the rule  $X \Rightarrow Y$  exists, there are two key coefficients, the support degree  $S$  and confidence  $C$ . The support degree  $S(X \Rightarrow Y) = P(X \cup Y)$  represents the probability of co-occurrence of itemsets  $X$  and  $Y$ . The confidence  $C(X \Rightarrow Y) = P(Y|X) = P(X \cup Y)/P(X)$  represents conditional probability of occurrence of itemset  $Y$  given that itemset  $X$  has occurred. The itemsets which satisfy the minimum threshold ( $S_{min}$ ) of support degree are so-called frequent itemsets, and those which satisfy both  $S_{min}$  and a minimum threshold of confidence ( $C_{min}$ ) are strong association rules.

The Apriori algorithm [57] can be decomposed into two main steps. The basic intuition is that any subset of a frequent itemset must be frequent. The first step is to generate frequent itemsets, as shown in Figure 4. The second step is to extract strong association rules based on frequent itemsets, as shown in Figure 5. The Apriori algorithm generates the candidate itemsets to be counted in a pass by using only the frequent itemsets in the previous pass. To improve the efficiency of frequent itemset extraction, the method utilizes a pruning strategy in order to compress the search space, that is, all non-empty subsets of frequent item-sets must also be frequent, and all parent sets of nonfrequent item-sets are nonfrequent.




Figure 4. Algorithm for generating frequent itemsets.




Figure 5. Algorithm for extracting strong association rules based on frequent itemsets.

## 4. Results and discussion

### 4.1. Spatial auto-correlation of elements

#### 4.1.1. Univariate spatial auto-correlation of individual elements

We calculated the spatial auto-correlation and cross-correlation indicators of each element using open-source software packages Geoda (<http://geodacenter.github.io>) and spdep (<https://github.com/r-spatial/spdep>). Then, we applied the Z-score to test the significance of spatial auto-correlation and cross-correlation statistics. Because the global Moran's I can be tested by normal or permutation tests [8], and the Z-score was calculated by Monte Carlo simulation by randomly sampling 999 permutations. The global Moran's I, global Geary's C, and global Getis-Ord's G for 15 elements passed the statistical significance test and are consistent with each other (Table 4). The global Moran's I and Geary's C are both suitable for characterizing the overall spatial pattern of an element, however, the global Getis-Ord's G only indicates whether an element's concentration exhibits a positive correlation (LL clustered or HH clustered) or is randomly distributed. It can't be used to ascertain a negative correlation or compare the correlation between elements.

Table 4. Univariate global spatial auto-correlation indicators for 15 elements in the study area.

Variable	Global Moran's I	Global Geary's C	Global Getis-Ord's G
----------	------------------	------------------	----------------------



	I	P-value	C	P-value	G	P-value	E(G)
log10(Au)	0.059	<2.2e-16	0.949	<2.2e-16	0.009	<2.2e-16	5.1e-09
log10(Sn)	0.439	<2.2e-16	0.563	<2.2e-16	0.014	<2.2e-16	6.7e-09
log10(Ag)	0.430	<2.2e-16	0.570	<2.2e-16	0.013	<2.2e-16	9.0e-09
log10(As)	0.451	<2.2e-16	0.551	<2.2e-16	0.012	<2.2e-16	1.1e-08
log10(Sb)	0.608	<2.2e-16	0.394	<2.2e-16	0.014	<2.2e-16	1.3e-08
log10(Bi)	0.438	<2.2e-16	0.564	<2.2e-16	0.019	<2.2e-16	5.2e-08
log10(Co)	0.423	<2.2e-16	0.578	<2.2e-16	0.009	<2.2e-16	5.8e-10
log10(Cu)	0.468	<2.2e-16	0.535	<2.2e-16	0.012	<2.2e-16	5.3e-09
log10(La)	0.259	<2.2e-16	0.740	<2.2e-16	0.009	<2.2e-16	9.1e-10
log10(Pb)	0.500	<2.2e-16	0.500	<2.2e-16	0.011	<2.2e-16	5.1e-09
log10(Zn)	0.530	<2.2e-16	0.468	<2.2e-16	0.010	<2.2e-16	1.1e-09
log10(W)	0.365	<2.2e-16	0.638	<2.2e-16	0.015	<2.2e-16	2.4e-08
log10(Mo)	0.430	<2.2e-16	0.572	<2.2e-16	0.012	<2.2e-16	4.0e-09
log10(Nb)	0.181	<2.2e-16	0.817	<2.2e-16	0.009	<2.2e-16	8.2e-10
log10(Cd)	0.459	<2.2e-16	0.539	<2.2e-16	0.012	<2.2e-16	4.0e-09


P-value < 0.05 means that the indicator passes the statistical significance test.

The global Getis–Ord’s G shows that all the 15 elements have a positive correlation in the study area. The global Moran’s I and Geary’s C show that Au is randomly distributed, and the other 14 elements are positively correlated. The elements ordered from high to low correlation are Sb, Zn, Pb, Cu, Cd, As, Sn, Bi, Ag, Mo, Co, W, La and Nb (Table 4). Except for Au, the global Moran’s I and Geary’s C are consistent with Getis–Ord’s G. According to the geological survey report, an Au deposit was found in the study area [59]. However, because of the low concentrations of Au in most sampling points of the study area, it would be easy to overlook the local clustering in the global spatial auto-correlation analysis.

We calculated the local Moran’s I of major elements in the study area and visualized the results via a Voronoi diagram (Figure 6). Anselin (1995) proposed a local indicator of spatial association (LISA) statistic that satisfies the following two requirements: a. the LISA for each observation gives an indication of the extent of significant spatial clustering of similar values around that observation; b. the sum of LISAs for all observations is proportional to a global indicator of spatial association [11]. By calculating the local Moran’s I  $I(i)$  in each quadrant, this divides the concentrations of elements into 5 categories: insignificant or high-high (HH), low-low (LL), low-high (LH), or high-low (HL) clustering [11]. A local Moran’s I clustering map (LMIC) represents different types of association between the value at a given location and its spatial lag, i.e., the weighted average of the values in the surrounding locations. The LISA significance map is shown in Figure 7, in which we set  $P=0.05$ . The local Moran’s I clustering map is shown in Figure 8. These results are consistent with the Moran’s I clustering results, which show that the HH and LL clustering in LMIC can reflect the spatial pattern of elements’ concentrations with a certain statistical significance. In addition, maps of local Moran’s I have natural transitions from strong to weak, which capture the local details and are consistent with the distributions of elements in nature.


Moreover, we also calculated indicators of the local Geary’s C and the local Getis–Ord’s G of all the elements in the study area. The HH and LL clustered values of the local Moran’s I and the local Geary’s C are similar; however, the local Getis–Ord’s G covers a broader space, especially Sb, As, Cu and Co. Compared with local Geary’s C and local Getis–Ord’s G, we can identify points with HH, LL, LH, and HL clustering with a precise

meaning for each category from the local Moran’s I. Therefore, we chose the LMIC results to mine the association rules of various elements.




**Figure 6.** Voronoi diagrams of local Moran’s I indicators for major elements: (a) Ag, (b) As, (c) Sb, (d) Bi, (e) Co, (f) Cu, (g) La, (h) Zn, and (i) Cd.





**Figure 7.** The LISA significance map of local Moran's I indicators for major elements: (a) Ag, (b) As, (c) Sb, (d) Bi, (e) Co, (f) Cu, (g) La, (h) Zn, and (i)Cd.





**Figure 8.** Clustering map of local Moran's I indicators for major elements: (a) Ag, (b) As, (c) Sb, (d) Bi, (e) Co, (f) Cu, (g) La, (h) Zn, and (i) Cd.

4.1.2. Bivariate spatial cross-correlation between two elements



The bivariate global Moran’s I for 15 elements in the study area are shown in Table 5, and all the calculated results passed the statistical significance test. The diagonal elements in Table 5 are consistent with the univariate global Moran’s I. The elements with strong positive correlations include Pb and Cd, Pb and Zn, Cu and Bi, and Zn and Cd, and those with negative correlations include La and Co, and La and Cu (Tables 6 and 7).

Table 5. Bivariate global Moran’s I for 15 elements in the study area.

Element/ log10()	Au	Sn	Ag	As	Sb	Bi	Co	Cu	La	Pb	Zn	W	Mo	Nb	Cd
Au	0.06	0.03	0.03	0.03	0.03	0.01	-0.03	0.00	0.01	0.02	-0.02	0.01	-0.02	-0.01	0.02
Sn	0.03	0.44	0.28	0.27	0.20	0.30	0.15	0.29	-0.03	0.29	0.15	0.19	0.08	-0.03	0.30
Ag	0.03	0.28	0.43	0.31	0.26	0.25	0.15	0.26	0.01	0.36	0.30	0.22	0.14	0.02	0.34
As	0.03	0.27	0.31	0.45	0.35	0.22	0.19	0.22	-0.02	0.34	0.26	0.16	0.09	0.02	0.33
Sb	0.03	0.20	0.26	0.35	0.61	0.15	0.11	0.10	0.04	0.32	0.26	0.12	-0.01	0.00	0.30
Bi	0.01	0.30	0.25	0.22	0.15	0.44	0.18	0.36	-0.06	0.28	0.21	0.27	0.23	0.03	0.24
Co	-0.03	0.15	0.15	0.19	0.11	0.18	0.42	0.31	-0.15	0.20	0.22	0.15	0.15	0.01	0.16
Cu	0.00	0.29	0.26	0.22	0.10	0.36	0.31	0.47	-0.13	0.26	0.22	0.29	0.27	0.04	0.23
La	0.01	-0.03	0.01	-0.02	0.04	-0.06	-0.15	-0.13	0.26	0.06	0.06	-0.02	-0.01	0.07	0.04
Pb	0.02	0.29	0.36	0.34	0.32	0.28	0.20	0.26	0.06	0.50	0.42	0.23	0.16	0.05	0.43
Zn	-0.02	0.15	0.30	0.26	0.26	0.21	0.22	0.22	0.06	0.42	0.53	0.22	0.21	0.22	0.36
W	0.01	0.19	0.22	0.16	0.12	0.27	0.15	0.29	-0.02	0.23	0.22	0.37	0.32	0.07	0.19
Mo	-0.02	0.08	0.14	0.09	-0.01	0.23	0.15	0.27	-0.01	0.16	0.21	0.32	0.43	0.10	0.11
Nb	-0.01	-0.03	0.02	0.02	0.00	0.03	0.01	0.04	0.07	0.05	0.22	0.07	0.10	0.18	0.03
Cd	0.02	0.30	0.34	0.33	0.30	0.24	0.16	0.23	0.04	0.43	0.36	0.19	0.11	0.03	0.46


Table 6. Bivariate global Moran’s I for elements with positive correlations.

$I_{ab}$	Positive correlation
$I_{ab} = 0.43$	Pb-Cd
$I_{ab} = 0.42$	Pb-Zn
$I_{ab} = 0.36$	Ag-Pb、Cu-Bi、Zn-Cd
$I_{ab} = 0.35$	As-Sb
$I_{ab} = 0.34$	Ag-Cd、Pb-As
$I_{ab} = 0.33$	As-Cd
$I_{ab} = 0.32$	Pb-Sb、Mo-W
$I_{ab} = 0.31$	Ag-As、Cu-Co
$I_{ab} = 0.30$	Sn-Bi、Sn-Cd、Ag-Zn、Sb-Cd

Table 7. Bivariate global Moran’s I for elements with negative correlations.


$I_{ab}$	Negative correlation
$I_{ab} = -0.13$	La-Co
$I_{ab} = -0.15$	La-Cu

The clustering map of bivariate local Moran's I divides the sampling points into 5 categories, i.e., insignificant, high-high (HH), low-low (LL), low-high (LH), and high-low (HL) clustered. However, their meanings are different from categories in a univariate clustering map. In the clustering map of bivariate local Moran's I,  $I_{ab}(i)$  indicates the spatial pattern of the related element  $b$  around the main element  $a$ . From this, we plotted  $I_{CuCo}$ ,  $I_{CoCu}$ ,  $I_{CuBi}$  and  $I_{AsSb}$ , as shown in Figure 9. In the  $I_{CuCo}$  and  $I_{CoCu}$ , the sampling points with high-high (HH) and low-low (LL) clustering are consistent with the univariate  $I_{Cu}$  and  $I_{Co}$ . Therefore,  $I_{CuCo}$  and  $I_{CoCu}$  show that the spatial distributions of Cu and Co in the study area are similar and positively cross-correlated. Due to the differences in the spatial distribution of Cu and Co, there are some differences in  $I_{CuCo}$  and  $I_{CoCu}$  after exchanging the main variable and related variable. The high-high (HH) clustering in  $I_{CuBi}$  and  $I_{AsSb}$  also has obvious regionality. The map for  $I_{LaCu}$  and  $I_{LaCo}$  is shown in Figure 10. There are apparent areas of low-high (LH) and high-low (HL) clustering in Figure 10, which indicate a negative cross-correlation of the two elements. Therefore, the bivariate local Moran's I effectively reveals whether two elements have a spatial cross-correlation.



**Figure 9.** Clustering map of bivariate local Moran's I with mainly positive cross-correlation: (a)  $I_{CuCo}$ , (b)  $I_{CoCu}$ , (c)  $I_{CuBi}$ , and (d)  $I_{AsSb}$ .





**Figure 10.** Clustering map of bivariate local Moran's I with mainly negative cross-correlation: (a)  $I_{LaCu}$  and (b)  $I_{LaCo}$ .

4.2. Association rules among multiple elements

4.2.1. Association rule mining

The 4959 geochemical sampling points were each taken as an event in the Apriori algorithm. Then we reorganized the geochemical concentration data into the original dataset D for association rule mining according to clustering by local univariate Moran's I, as shown in Table 8. Table 9 shows the statistics for the LMIC analysis. Some items frequently appear in events, while some items are very sparse. If the support threshold in the Apriori algorithm is set too small, the efficiency of the mining algorithm is low, and a large number of meaningless rules may be extracted. If the support threshold is set too large, the efficiency of the mining algorithm is high, but it may filter out some sparse items. For this study, we took the support threshold  $S_{min} = 0.05$  and confidence threshold  $C_{min} = 0.7$ .

**Table 8.** Original example dataset for association rule mining.

$t_k$ (Point)	$i_1$ (Au)	$i_2$ (Sn)	$i_3$ (Ag)	$i_4$ (As)	$i_5$ (Sb)	$i_6$ (Bi)	$i_7$ (Co)	$i_8$ (Cu)	$i_9$ (La)	$i_{10}$ (Pb)	$i_{11}$ (Zn)	$i_{12}$ (W)	$i_{13}$ (Mo)	$i_{14}$ (Nb)	$i_{15}$ (Cd)
1	HL			HH	HH				HH	HH	HH	HH	HH	HH	HH
2			LL		HH		HH		HL						
3		HH			LH	HH	LH					HH			
4	HH	HH			HH			HH		LH					HH
5		LL	LL	LL		LL	LL	LL	HH	LL					LL
6		LL	LL	LL	LL	LL		LL		LL	LL	LL	LL		LL
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
4959		HH	HH	HH	HH	HH	HH	HH	LH	HH	HH	HH	HH		HH



**Table 9.** Counts of items in local Moran’s I clustering (LMIC) of elements.

Element	Insignificant	High-high	Low-low	Low-high	High-low
Au	4023	113	466	163	194
Sn	2584	300	1786	190	99
Ag	1706	560	2286	336	71
As	1931	521	2137	259	111
Sb	1192	867	2509	305	86
Bi	2352	351	1973	194	89
Co	1810	981	1613	327	228
Cu	2036	570	2044	169	140
La	2025	659	1529	417	329
Pb	3740	65	796	315	43
Zn	1662	793	2049	200	255
W	2442	332	1874	193	118
Mo	2289	414	1952	128	176
Nb	2749	478	1132	329	271
Cd	2054	556	1927	303	119

We used the Apriori algorithm to mine out dozens of association rules, of which 15 rules were selected for interpretation (Table 10). The supports for Au and Pb in the Apriori algorithm are smaller than the threshold, so no relevant association rules were mined. Meanwhile, the relevance of these association rules was judged according to the coexistence of elements and the geological environment in the study area.

**Table 10.** Mined association rules among elements.

ID	Association rules	Support degree	Confidence
a	{As (HH)} ⇒ {Sb (HH)}	0.076	0.73
b	{Cd (HH)} ⇒ {Zn (HH)}	0.090	0.81
c	{W (HH)} ⇒ {Cu (HH)}	0.051	0.76
d	{Cu (HH)} ⇒ {Co (HH)}	0.089	0.77
e	{Bi (HH)} ⇒ {Cu (HH)}	0.059	0.83
f	{Mo (HH)} ⇒ {Sb (LL)}	0.065	0.77
g	{Zn (HH), Ag (HH)} ⇒ {Cd (HH)}	0.058	0.81
h	{Cd (HH), Ag (HH)} ⇒ {Zn (HH)}	0.058	0.92
i	{Cd (HH), As (HH)} ⇒ {Zn (HH)}	0.053	0.93
j	{As (HH), Zn (HH)} ⇒ {Cd (HH)}	0.053	0.82
k	{Zn (HH), Sb (HH)} ⇒ {Cd (HH)}	0.055	0.71
l	{Cd (HH), Sb (HH)} ⇒ {Zn (HH)}	0.055	0.93
m	{Zn (HH), As (HH)} ⇒ {Sb (HH)}	0.052	0.82
n	{Cu (HH), La (LL)} ⇒ {Co (HH)}	0.052	0.76
o	{Co (HH), Sb (LL)} ⇒ {La (LL)}	0.056	0.73


HH (high-high clustered), LL (low-low clustered).

4.2.2. Comparison with bivariate spatial cross-correlation


The affinity of elements is the ability of elements to preferentially coexist with each other. The most abundant anions in the crustal system are oxygen (O) and sulfur (S), and the most abundant element that can exist as a natural metal is iron (Fe). Therefore, according to the geochemical affinities, the 15 elements are divided into the following three categories: (1) siderophile elements, i.e., Au; (2) sulfophile elements, i.e., Sn, Ag, As, Sb, Bi, Cu, Co, Pb, Zn, and Cd; and (3) oxyphile elements, i.e., Mo, Nb, W, and La.

The mining of association rules shows that there are positive correlations among all sulfophile elements with HH clustering, that is,  $\{As(HH)\} \Rightarrow \{Sb(HH)\}$ ,  $\{Cd(HH)\} \Rightarrow \{Zn(HH)\}$ ,  $\{Cu(HH)\} \Rightarrow \{Co(HH)\}$ ,  $\{Bi(HH)\} \Rightarrow \{Cu(HH)\}$ ,  $\{Zn(HH), Ag(HH)\} \Rightarrow \{Cd(HH)\}$ ,  $\{Cd(HH), Ag(HH)\} \Rightarrow \{Zn(HH)\}$ ,  $\{Cd(HH), As(HH)\} \Rightarrow \{Zn(HH)\}$ ,  $\{As(HH), Zn(HH)\} \Rightarrow \{Cd(HH)\}$ ,  $\{Zn(HH), Sb(HH)\} \Rightarrow \{Cd(HH)\}$ ,  $\{Cd(HH), Sb(HH)\} \Rightarrow \{Zn(HH)\}$ , and  $\{Zn(HH), As(HH)\} \Rightarrow \{Sb(HH)\}$ . In rules  $\{Mo(HH)\} \Rightarrow \{Sb(LL)\}$ ,  $\{Cu(HH), La(LL)\} \Rightarrow \{Co(HH)\}$ , and  $\{Co(HH), Sb(LL)\} \Rightarrow \{La(LL)\}$ , there are positive correlations between sulfophile elements with HH clustering and oxyphile elements with LL clustering.

We next compared the bivariate spatial cross-correlations and association rules for Cu and Co (Figure 11), and As and Sb (Figure 12). The distributions of  $I_{CuCo}$  HH clustering and  $I_{AsSb}$  HH clustering are spatially similar to the association rules  $\{Cu(HH)\} \Rightarrow \{Co(HH)\}$  and  $\{As(HH)\} \Rightarrow \{Sb(HH)\}$ , respectively; however,  $I_{CuCo}$  HH clustering and  $I_{AsSb}$  HH clustering cover wider areas. In addition,  $I_{CuCo}$  and  $I_{AsSb}$  reveal not only high HH clustering but also LL, LH, and HL clustering, which shows the simultaneous relationship between two elements, but does not scale efficiently to massive data sets. In contrast, association rule mining is suitable for revealing the association among items in a large geochemical dataset.



**Figure 11.** (a) Association rule mining result  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  and (b) bivariate spatial cross-correlation indicator  $I_{CuCo}$  of Cu and Co.



**Figure 12.** (a) Association rule mining result  $\{As (HH)\} \Rightarrow \{Sb (HH)\}$  and (b) bivariate spatial cross-correlation indicator  $I_{AsSb}$  of As and Sb.

#### 4.2.3. Controls of geological features

Due to the influence of multiple stages of tectonic and magmatic activities, the fault structures in the study area are relatively well developed. We calculated the Euclidean distance field for the faults in the study area (see Figure 13). Then, the 15 mined association rules were overlaid with the fault distance field (Figure 14). We found that more than 86% of the mined rule points are located within 1000 m distance of the fault, especially  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  (Figure 15) and  $\{Zn (HH), Sb (HH)\} \Rightarrow \{Cd (HH)\}$  (Figure 16). The rule  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  is most predominant near the faults in the northwest and southeast parts of the study area, and three known copper ore occurrences are also near the faults. The rule  $\{Zn (HH), Sb (HH)\} \Rightarrow \{Cd (HH)\}$  is most strongly associated with the faults in the southeast part of the study area, and a known lead-zinc ore occurrence is near the faults. That is, the fault structure has an obvious control effect on clustering of the elements. In addition, we extracted streams and catchments area to analyze whether element co-occurrence is related to stream transport. As shown in Figures 17 and 18, most  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  and  $\{Zn (HH), Sb (HH)\} \Rightarrow \{Cd (HH)\}$  events are distributed in the upper reaches of the streams and catchment areas, so the impact of stream transport on element association rule mining is weak in the study area.






Figure 13. Euclidean distance field of faults.




Figure 14. Numbers of mined rule points that are close to faults.




Figure 15. Rule  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  points within 500 m and 1000 m of faults.






Figure 16. Rule {Zn (HH), Sb (HH)} ⇒ {Cd (HH)} points within 500 m and 1000 m of faults.




Figure 17. Rule {Cu (HH)} ⇒ {Co (HH)} points overlaid with streams and catchments area.




Figure 18. Rule {Zn (HH), Sb (HH)} ⇒ {Cd (HH)} points overlaid with streams and catchments area.

Figure 10 displays 18 subplots (a-o) showing the spatial distribution of point density (point/km²) and point density (point/km²) for various locations. Each plot has two y-axes: the left axis represents 'point' (0 to 180) and the right axis represents 'density(point/km²)' (0.0 to 4.0). The x-axis lists locations: Pt.b, T.yb, T.e, P.ny, J.fy, T.ny, P.yb, O.ybo, T.mny, P.bo, J.fy. The legend indicates 'density(point/km²)' (grey bar) and 'point' (purple line with circles). Data values are labeled on the plots.

(a) Pt.b: 141, T.yb: 15, T.e: 36, P.ny: 63, J.fy: 9, T.ny: 21, P.yb: 72, O.ybo: 12, T.mny: 21, P.bo: 2, J.fy: 0.

(b) P.ybo: 22, T.yb: 13, Pt.b: 79, P.bo: 19, T.ny: 24, P.ny: 40, T.mny: 34, P.yb: 60, O.ybo: 4, T.e: 0, J.fy: 0.

(c) P.ybo: 28, P.bo: 37, T.yb: 12, Pt.b: 83, P.ny: 68, T.ny: 24, T.mny: 40, P.yb: 51, O.ybo: 4, T.e: 0, J.fy: 0.

(d) T.yb: 18, P.ybo: 19, Pt.b: 95, P.bo: 15, T.mny: 44, P.ny: 32, T.ny: 13, P.yb: 72, O.ybo: 0, T.e: 0, J.fy: 0.

(e) P.ybo: 92, T.e: 20, T.yb: 4, P.ny: 34, J.fy: 6, P.yb: 44, O.ybo: 2, T.mny: 2, T.ny: 2, P.bo: 0, P.ybo: 0.

(f) Pt.b: 95, T.yb: 9, T.e: 22, P.ny: 43, T.mny: 20, P.yb: 42, O.ybo: 2, T.ny: 2, J.fy: 0, P.bo: 0, P.ybo: 0.

(g) Pt.b: 134, T.e: 32, J.fy: 7, T.yb: 4, P.ny: 18, P.yb: 25, P.bo: 3, O.ybo: 5, T.mny: 0, P.ybo: 0.

(h) Pt.b: 115, T.e: 15, T.yb: 4, J.fy: 5, P.ny: 26, T.ny: 35, P.yb: 11, O.ybo: 2, T.mny: 2, T.ny: 2, P.bo: 0, P.ybo: 0.

(i) T.yb: 49, P.ybo: 19, Pt.b: 107, P.bo: 23, T.mny: 34, P.ny: 15, P.yb: 26, O.ybo: 52, T.e: 4, J.fy: 0.

(j) T.yb: 25, P.ybo: 18, P.bo: 32, Pt.b: 73, P.yb: 35, P.ny: 26, T.mny: 5, O.ybo: 2, T.e: 2, T.ny: 0, J.fy: 0.

(k) T.yb: 49, P.ybo: 19, Pt.b: 107, P.bo: 23, T.mny: 34, P.ny: 15, P.yb: 26, O.ybo: 52, T.e: 4, J.fy: 0.

(l) T.yb: 49, P.ybo: 19, Pt.b: 107, P.bo: 23, T.mny: 34, P.ny: 15, P.yb: 26, O.ybo: 52, T.e: 4, J.fy: 0.

(m) T.yb: 49, P.ybo: 19, Pt.b: 107, P.bo: 23, T.mny: 34, P.ny: 15, P.yb: 26, O.ybo: 52, T.e: 4, J.fy: 0.


(n) T.yb: 49, P.ybo: 19, Pt.b: 107, P.bo: 23, T.mny: 34, P.ny: 15, P.yb: 26, O.ybo: 52, T.e: 4, J.fy: 0.

(o) T.yb: 49, P.ybo: 19, Pt.b: 107, P.bo: 23, T.mny: 34, P.ny: 15, P.yb: 26, O.ybo: 52, T.e: 4, J.fy: 0.

**Figure 19.** Total number of points and density of each mined rule, e.g., (a) {As (HH)}  $\Rightarrow$  {Sb (HH)}, (b) {Cd (HH)}  $\Rightarrow$  {Zn (HH)}, (c) {W (HH)}  $\Rightarrow$  {Cu (HH)}, (d) {Cu (HH)}  $\Rightarrow$  {Co (HH)}, (e) {Bi (HH)}  $\Rightarrow$  {Cu (HH)}, (f) {Mo (HH)}  $\Rightarrow$  {Sb (LL)}, (g) {Zn (HH), Ag (HH)}




$\Rightarrow \{Cd (HH)\}$ , (h)  $\{Cd (HH), Ag (HH)\} \Rightarrow \{Zn (HH)\}$ , (i)  $\{Cd (HH), As (HH)\} \Rightarrow \{Zn (HH)\}$ , (j)  $\{As (HH), Zn (HH)\} \Rightarrow \{Cd (HH)\}$ , (k)  $\{Zn (HH), Sb (HH)\} \Rightarrow \{Cd (HH)\}$ , (l)  $\{Cd (HH), Sb (HH)\} \Rightarrow \{Zn (HH)\}$ , (m)  $\{Zn (HH), As (HH)\} \Rightarrow \{Sb (HH)\}$ , (n)  $\{Cu (HH), La (LL)\} \Rightarrow \{Co (HH)\}$ , and (o)  $\{Co (HH), Sb (LL)\} \Rightarrow \{La (LL)\}$ , in the main litho-strata.




**Figure 20.** Rule  $\{As (HH)\} \Rightarrow \{Sb (HH)\}$  overlaid with the main litho-strata.

According to the geological survey data, the enrichment of Cu, Pb, Zn, Ag, Bi, and other elements in the lower Proterozoic Baishahe Formation (Pt<sub>1b</sub>) provides the main ore-forming materials in the study area. The locations of rules  $\{As (HH)\} \Rightarrow \{Sb (HH)\}$ ,  $\{Zn (HH), Ag (HH)\} \Rightarrow \{Cd (HH)\}$  and  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  are related to Pt<sub>1b</sub>, as shown in Figure 20, 21 and 22. The association rules of sulfophile elements, e.g.,  $\{As (HH)\} \Rightarrow \{Sb (HH)\}$ ,  $\{Cd (HH)\} \Rightarrow \{Zn (HH)\}$ ,  $\{Zn (HH), Ag (HH)\} \Rightarrow \{Cd (HH)\}$ ,  $\{Cd (HH), Ag (HH)\} \Rightarrow \{Zn (HH)\}$ ,  $\{Cd (HH), As (HH)\} \Rightarrow \{Zn (HH)\}$ ,  $\{Cd (HH), As (HH)\} \Rightarrow \{Zn (HH)\}$ ,  $\{Zn (HH), Sb (HH)\} \Rightarrow \{Cd (HH)\}$ ,  $\{Cd (HH), Sb (HH)\} \Rightarrow \{Zn (HH)\}$ , and  $\{Zn (HH), As (HH)\} \Rightarrow \{Sb (HH)\}$ , are mainly distributed in the lower Proterozoic Baishahe Formation (Pt<sub>1b</sub>), the upper Triassic Elashan Formation (T<sub>3e</sub>), and the Indosinian granodiorite (T<sub>2γδ</sub>). Indosinian magmatism resulted in the intrusion of the middle Triassic Kekesai Sequence granite and the late Triassic Zamari Sequence granite, which provided conditions for enrichment of many sulfophile elements in the study area, especially represented by the rule  $\{Zn (HH), Ag (HH)\} \Rightarrow \{Cd (HH)\}$  (Figure 21). Therefore, the Indosinian magmatism provided a heat and material source to enrich elements, and is an important geological unit for aggregating sulfophile elements. Cu mineralization often occurs in the contact between the Variscan magmatic rocks and surrounding rocks such as the Baishahe Formation (Pt<sub>1b</sub>), forming the Keregou East copper occurrence and the Hariza copper deposit. The rules of  $\{W (HH)\} \Rightarrow \{Cu (HH)\}$ ,  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$ ,  $\{Bi (HH)\} \Rightarrow \{Cu (HH)\}$ ,  $\{Cu (HH), La (LL)\} \Rightarrow \{Co (HH)\}$ ,  $\{Co (HH), Sb (LL)\} \Rightarrow \{La (LL)\}$  related to Cu HH and Co HH clustering also have high density in the Variscan granite quartz diorite (P<sub>1γδo</sub>), especially  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  (Figure 22). Therefore, we may infer that a co-existing relationship between Cu and other elements developed in the Variscan granite quartz diorite.



**Figure 21.** Rule  $\{Zn (HH), Ag (HH)\} \Rightarrow \{Cd (HH)\}$  overlaid with the main litho-strata.



**Figure 22.** Rule  $\{Cu (HH)\} \Rightarrow \{Co (HH)\}$  overlaid with the main litho-strata.

## 5. Conclusions

The case study of association rule mining in the Chahanwusu River area yields the following conclusions.

(1) According to the global auto-correlation indicators, Au shows a random distribution in the study area, and other 14 elements have positive correlations ranked from large to small are Sb, Zn, Pb, Cu, Cd, As, Sn, Bi, Ag, Mo, Co, W, La, and Nb. Compared with local Geary's C and local Getis-Ord's G, local Moran's I can identify points of HH, LL, LH and HL clustering with a precise meaning for each category, which makes it a better local auto-correlation indicator for association rule mining.

(2) Based on the univariate LMIC results, the proposed method successfully mined 15 association rules among various elements in the study area. Bivariate spatial cross-correlation can also detect distribution pattern details of pair elements co-occurrence compared with association rule mining method. However, it cannot be used to efficiently explore massive geochemical datasets. In contrast, association rule mining can reveal the association among items in a large geochemical dataset.

(3) Overlaying the mining results of association rules on the faults, ore occurrences and catchment areas, we found that more than 86% of the mined rule points are located within 1000 m of faults and near known ore occurrences, and the impact of stream transport on element co-occurrences is weak. Greater densities and numbers of points of association rules were found in the lower Proterozoic Baishahe Formation (Pt,b) and the



Variscan granodiorite (P<sub>1</sub>γδ). Therefore, the association rules are closely related to specific geological features.

The association rules mined in this paper are mainly high-value element co-occurrence. Where these combinations appear, higher concentration of the element is more likely, which can improve the prediction of unknown ore deposits or occurrences. However, the mining efficiency of low-value element co-occurrence is low and the local dilution of elements in the study area cannot be effectively detected. In the future, we will build an element association rule database to find combinations of anomalies for known metallogenic elements and to map the probability of unknown mineralization in the study area.

**Author Contributions:** Conceptualization, B.Z. and J.D.; methodology, Y.C.; software, Z.J. and N.C.; validation, B.Z., Z.J., and Y.C.; data curation, Y.C.; writing—original draft preparation, B.Z. and Z.J.; writing—review and editing, B.Z. and U.K.; funding acquisition, B.Z. All authors have read and agreed to the published version of the manuscript.

**Funding:** This study was supported by grants from the National Natural Science Foundation of China (Grant Nos. 42072326 and 41772348), China Geological Survey Project (Grant No. DD20190156), and the National Key Research and Development Program of China (Grant No. 2019YFC1805905).

**Acknowledgments:** The authors would like to thank the Co-Construction MapGIS Library by Engineering Research Center for Geographic Information System of China and Central South University for providing MapGIS® software. We also thank senior engineer Professor ZHANG Shao-ning (The 8th Team of Qinghai Provincial Bureau of Nonferrous Metals and Geological Exploration), Professor LAI Jian-qing (Central South University) for their kind assistance with data collection and Professor Jeffrey Dick (Central South University) for revising scientific-English writing of the article.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Korobova, E.M.; Romanov, S.L. A Chernobyl 137Cs contamination study as an example for the spatial structure of geochemical fields and modeling of the geochemical field structure. *Chemometrics and Intelligent Laboratory Systems* **2009**, *99*, 1-8.
2. Zhang, B.; Chen, Y.; Huang, A.; Lu, H.; Cheng, Q. Geochemical field and its roles on the 3D prediction of concealed ore-bodies. *Acta Petrologica Sinica* **2018**, *34*, 352-362.
3. Tobler, W.R. A computer movie simulating urban growth in the Detroit region. *Economic geography* **1970**, *46*, 234-240.
4. Moran, P.A. Notes on continuous stochastic phenomena. *Biometrika* **1950**, *37*, 17-23.
5. Geary, R.C. The Contiguity Ratio and Statistical Mapping. *Incorporated Statian* **1954**, *5*, 115-146.
6. Cliff, A.D.; Ord, J.K. The Problem of Spatial Autocorrelation. *Regional Science* **1969**, *1*, 26-55.
7. Cliff, A.D.; Ord, J.K. Evaluating the percentage points of a spatial autocorrelation coefficient. *Geographical Analysis* **1971**, *3*, 51-62.
8. Cliff, A.D.; Ord, J.K. *Spatial processes: models & applications*; Taylor & Francis: 1981.
9. Getis, A.; Ord, J.K. *The Analysis of Spatial Association by Use of Distance Statistics*; Springer: 2010; pp. 127-145.
10. Ord, J.K.; Getis, A. Local spatial autocorrelation statistics: distributional issues and an application. *Geographical analysis* **1995**, *27*, 286-306.
11. Anselin, L. Local indicators of spatial association—LISA. *Geographical analysis* **1995**, *27*, 93-115.
12. Boots, B.; Okabe, A. Local statistical spatial analysis: Inventory and prospect. *International Journal of Geographical Information Science* **2007**, *21*, 355-375.
13. Anselin, L. A local indicator of multivariate spatial association: extending Geary's C. *Geographical Analysis* **2019**, *51*, 133-150.

14. Goovaerts, P.; Jacquez, G.M. Accounting for regional background and population size in the detection of spatial clusters and outliers using geostatistical filtering and spatial neutral models: the case of lung cancer in Long Island, New York. *International Journal of Health Geographics* **2004**, *3*, 14.
15. McLaughlin, C.C.; Boscoe, F.P. Effects of randomization methods on statistical inference in disease cluster detection. *Health & place* **2007**, *13*, 152-163.
16. Yu, X.; Wang, S.; Wang, H.; Liang, Y.; Chen, S.; Wu, K.; Yang, Z.; Li, C.; Chang, Y.; Zhan, Y. Detection of Geochemical Element Assemblage Anomalies Using a Local Correlation Approach. *Journal of Earth Science* **2021**, *32*, 408-414.
17. Xiao, G.; Hu, Y.; Li, N.; Yang, D. Spatial autocorrelation analysis of monitoring data of heavy metals in rice in China. *Food Control* **2018**, *89*, 32-37.
18. Bivand, R.S.; Wong, D.W. Comparing implementations of global and local indicators of spatial association. *Test* **2018**, *27*, 716-748.
19. Cheng, Q. Singularity theory and methods for mapping geochemical anomalies caused by buried sources and for predicting undiscovered mineral deposits in covered areas. *Journal of Geochemical Exploration* **2012**, *122*, 55-70.
20. Carranza, E.J.M. *Geochemical anomaly and mineral prospectivity mapping in GIS*; Elsevier: 2008.
21. Zuo, R.; Xiong, Y. Geodata science and geochemical mapping. *Journal of Geochemical Exploration* **2019**, *209*, 106431.
22. Wang, J.; Zhou, Y.; Xiao, F. Identification of multi-element geochemical anomalies using unsupervised machine learning algorithms: A case study from Ag–Pb–Zn deposits in north-western Zhejiang, China. *Applied Geochemistry* **2020**, *120*, 104679.
23. Taylor, R.; Steven, T. Definition of mineral resource potential. *Economic Geology* **1983**, *78*, 1268-1270.
24. Wang, L.; Wu, X.; Zhang, B.; Li, X.; Huang, A.; Meng, F.; Dai, P. Recognition of Significant Surface Soil Geochemical Anomalies Via Weighted 3D Shortest-Distance Field of Subsurface Orebodies: A Case Study in the Hongtoushan Copper Mine, NE China. *Natural Resources Research* **2019**, *28*, 587-607.
25. Hawkes, H.E.; Webb, J.S. Geochemistry in mineral exploration. *Soil Science* **1963**, *95*, 283.
26. Sinclair, A. Selection of threshold values in geochemical data using probability graphs. *Journal of Geochemical Exploration* **1974**, *3*, 129-149.
27. Govett, G.; Goodfellow, W.; Chapman, R.; Chork, C. Exploration geochemistry—distribution of elements and recognition of anomalies. *Journal of the International Association for Mathematical Geology* **1975**, *7*, 415-446.
28. El-Makky, A.M. Statistical analyses of La, Ce, Nd, Y, Nb, Ti, P, and Zr in bedrocks and their significance in geochemical exploration at the Um Garayat Gold mine area, Eastern Desert, Egypt. *Natural resources research* **2011**, *20*, 157.
29. Ravani, P.; Barrett, B.J.; Parfrey, P.S. Longitudinal Studies 2: Modeling Data Using Multivariate Analysis. *Methods in molecular biology (Clifton, NJ)* **2021**, *2249*, 103-124.
30. Cox, D.R.; Snell, E.J. *Analysis of binary data*; Routledge: 2018.
31. Cioci, A.C.; Cioci, A.L.; Mantero, A.M.; Parreco, J.P.; Yeh, D.D.; Rattan, R. Advanced statistics: Multiple logistic regression, Cox proportional hazards, and propensity scores. *Surgical Infections* **2021**, *22*, 604-610.
32. Agterberg, F.P. Computer programs for mineral exploration. *Science* **1989**, *245*, 76-81.
33. Cheng, Q.; Agterberg, F. Fuzzy weights of evidence method and its application in mineral potential mapping. *Natural resources research* **1999**, *8*, 27-35.
34. Goyes-Penafiel, P.; Hernandez-Rojas, A. Double landslide susceptibility assessment based on artificial neural networks and weights of evidence. *Boletín De Geología* **2021**, *43*, 173-191.
35. Cheng, Q.; Agterberg, F.; Ballantyne, S. The separation of geochemical anomalies from background by fractal methods. *Journal of Geochemical Exploration* **1994**, *51*, 109-130.
36. Cheng, Q.; Xu, Y.; Grunsky, E. Integrated spatial and spectrum method for geochemical anomaly separation. *Natural Resources Research* **2000**, *9*, 43-52.



37. Cheng, Q. Mapping singularities with stream sediment geochemical data for prediction of undiscovered mineral deposits in Gejiu, Yunnan Province, China. *Ore Geology Reviews* **2007**, 32, 314-324.
38. Goovaerts, P. Geostatistical modelling of spatial uncertainty using p-field simulation with conditional probability fields. *International Journal of Geographical Information Science* **2002**, 16, 167-178.
39. Naik, M.R.; Barik, M.; Prasad, K.; Kumar, A.; Verma, A.K.; Sahoo, S.K.; Jha, V.; Sahoo, N.K. Hydro-geochemical analysis based on entropy and geostatistics model for delineation of anthropogenic ground water pollution for health risks assessment of Dhenkanal district, India. *Ecotoxicology* **2021**, 2, 1-16.
40. Zuo, R.; Carranza, E.J.M. Support vector machine: a tool for mapping mineral prospectivity. *Computers & Geosciences* **2011**, 37, 1967-1975.
41. Xiong, J.; Li, J.; Cheng, W.; Wang, N.; Guo, L. A GIS-based support vector machine model for flash flood vulnerability assessment and mapping in China. *ISPRS International Journal of Geo-Information* **2019**, 8, 297.
42. Rodriguez-Galiano, V.; Chica-Olmo, M.; Chica-Rivas, M. Predictive modelling of gold potential with the integration of multisource information based on random forest: a case study on the Rodalquilar area, Southern Spain. *International Journal of Geographical Information Science* **2014**, 28, 1336-1354.
43. Zhang, B.; Li, M.; Li, W.; Jiang, Z.; Khan, U.; Wang, L.; Wang, F. Machine learning strategies for lithostratigraphic classification based on geochemical sampling data: a case study in the area of the Chahanwusu River, Qinghai Province, China. *Journal of Central South University* **2021**, 28, 1422-1447.
44. Porwal, A.; Carranza, E.J.M. *Classifiers for modeling of mineral potential*; Wiley-Blackwell: 2008.
45. Porwal, A.; Carranza, E.J.M.; Hale, M. Bayesian network classifiers for mineral potential mapping. *Computers & Geosciences* **2006**, 32, 1-16.
46. Klüppelberg, C.; Krali, M. Estimating an extreme Bayesian network via scalings. *Journal of Multivariate Analysis* **2021**, 181, 104672.
47. Xiong, Y.; Zuo, R. Recognition of geochemical anomalies using a deep autoencoder network. *Computers & Geosciences* **2016**, 86, 75-82.
48. Nguyen, T.T.; Liu, X.; Ren, Z. A study of geochemical exploration spational cluster identification based on local spatial autocorrelation. *Geographical & Geochemical Exploration* **2014**, 38, 370-376.
49. Wang, H.; Cheng, Q.; Zuo, R. Spatial characteristics of geochemical patterns related to Fe mineralization in the southwestern Fujian province (China). *Journal of Geochemical Exploration* **2015**, 148, 259-269.
50. Ji, b.; Zhou, T.; Yuan, F.; Zhang, D.; Liu, L.; Liu, G. A method for identifying geochemical anomalies based on spatial autocorrelation. *Science of Surveying and Mapping* **2017**, 42, 24-27+123.
51. Sadeghi, M.; Morris, G.A.; Carranza, E.J.M.; Ladenberger, A.; Andersson, M. Rare earth element distribution and mineralization in Sweden: an application of principal component analysis to FOREGS soil geochemistry. *Journal of Geochemical Exploration* **2013**, 133, 160-175.
52. Wang, J.; Zuo, R. Quantifying the Distribution Characteristics of Geochemical Elements and Identifying Their Associations in Southwestern Fujian Province, China. *Minerals* **2020**, 10, 183.
53. Zhang, C.-S.; Li, Y. Extension of local association rules mining algorithm based on apriori algorithm. In Proceedings of the 2014 5th IEEE International Conference on Software Engineering and Service Science (ICSESS), 2014; pp. 340-343.
54. Zhang, X. Study of an improved Apriori algorithm for data mining of association rules. In Proceedings of the International Conference on Applied Science & Engineering Innovation, 2015.
55. Xu, T.; Dong, X. Mining frequent patterns with multiple minimum supports using basic Apriori. In Proceedings of the 2013 Ninth International Conference on Natural Computation (ICNC), 2013; pp. 957-961.
56. Wu, X.; Kumar, V.; Quinlan, J.R.; Ghosh, J.; Yang, Q.; Motoda, H.; McLachlan, G.J.; Ng, A.; Liu, B.; Philip, S.Y. Top 10 algorithms in data mining. *Knowledge and information systems* **2008**, 14, 1-37.

- 
57. Agrawal, R.; Imieliński, T.; Swami, A. Mining association rules between sets of items in large databases. In Proceedings of the Proceedings of the 1993 ACM SIGMOD international conference on Management of data, 1993; pp. 207-216.
  58. Liu, X.; Zhou, Y. Application of association rule algorithm in studying abnormal elemental associations in the Pangxidong area in western Guangdong Province, China. *Earth Science Frontiers* **2019**, *26*, 57-71.
  59. Tang, J.; Ji, B.; Yan, Y. Comprehensive survey report of 1:50000 regional mineral geology, stream sediment geochemistry and high-precision magnetic survey in the Chahanwusu River area, Dulan County, Qinghai Province. **2008**, 254-273.
  60. Chou, Y.H. Spatial pattern and spatial autocorrelation. In Proceedings of the International Conference on Spatial Information Theory, 1995; pp. 365-376.
  61. Buja, A.; Cook, D.; Swayne, D.F. Interactive high-dimensional data visualization. *Journal of computational and graphical statistics* **1996**, *5*, 78-99.
  62. Anselin, L.; Syabri, I.; Smirnov, O. Visualizing multivariate spatial correlation with dynamically linked windows. In Proceedings of the Proceedings, CSISS Workshop on New Tools for Spatial Data Analysis, Santa Barbara, CA, 2002.