

## Review

# Review and Comparison of Antimicrobial Resistance Gene Databases

Márton Papp<sup>1</sup> and Norbert Solymosi<sup>1,\*</sup>

<sup>1</sup> Centre for Bioinformatics, University of Veterinary Medicine Budapest, 1078 Budapest, Hungary; [papp.marton@univet.hu](mailto:papp.marton@univet.hu) (M. P.); [solymosi.norbert@gmail.com](mailto:solymosi.norbert@gmail.com) (N. S.)

\* Correspondence: [solymosi.norbert@gmail.com](mailto:solymosi.norbert@gmail.com)

**Abstract:** As the prevalence of antimicrobial resistance genes is increasing in microbes, we are facing the return of the preantibiotic era. Consecutively, the number of studies concerning antibiotic resistance and its spread in the environment is rapidly growing. Next generation sequencing technologies are widespread used in many areas of biological research and antibiotic resistance is no exception. For the rapid annotation of whole genome sequencing and metagenomic results considering antibiotic resistance, several tools and data resources were developed. These databases, however can differ fundamentally in the number and type of genes and resistance determinants they comprise. Furthermore, the annotation structure and metadata stored in these resources can also contribute to their differences. Several previous reviews were published on the tools and databases of resistance gene annotation, however, to our knowledge, no previous review focused solely and in depth on the differences in the databases. In this review, we compare the most well-known and widely used antibiotic resistance gene databases based on their structure and content. We believe that this knowledge is fundamental for selecting the most appropriate database for a research question and for the development of new tools and resources of resistance gene annotation.

**Keywords:** antibiotic resistance genes, antibiotic resistance gene database, annotation of antibiotic resistance genes

## 1. Introduction

Antimicrobial resistance (AMR) means an emerging threat on humanity. Based on a 2017 report it is estimated that ~ 700.000 deaths can be attributed to AMR worldwide [1]. As stated by a CDC study, approximately 35000 people dies in the United States yearly due to antibiotic resistance [2], and a similar number was estimated in the European Union as well [3]. However, it is expected that the impact of AMR will further increase and claim approximately 10 million lives yearly by 2050 [1]. The emergence of resistant microbes will not only cause untreatable primer infections, but the safe performance of routine medical procedures (such as surgeries or chemotherapeutic treatment of oncological patients) will become impossible due to the inability of a successful antibiotic prophylaxis [1,2,4]. Even though one usually associates AMR with hospitals and the misuse/overuse of antibiotics by medical professionals, the influence of agriculture and the environment is no less important [1,2,4–6]. Therefore to tackle this global challenge the investigation of the spread of AMR between different environments is required.

Phenotypic AMR in microbes is necessarily lies in genetic fundamentals which can be categorized to two main mechanisms. On one hand, AMR can arise through genetic mutations (e.g. modification of the antibiotic target site, overexpression of efflux pumps or the antibiotic target molecule etc.), under the selective pressure of antibiotics, or by the acquisition of specific genes conferring resistance (e.g. genes coding enzymes that degrade the antibiotic compounds or open alternative metabolic pathways for evading the effects of the antibiotic) through horizontal gene transfer (HGT) [7,8]. It is believed that the

majority of antimicrobial resistance genes (ARG) transmitted between bacteria is not the novel product of widespread antibiotic usage by humans, but evolved previously for a variety of functions and enriched with the extensive usage of antibiotics since the mid 20th century [9–11]. As environmental microbes have a significant role in the spread of resistance genes, the global surveillance of ARGs in various environments is critical for understanding and combating AMR [6,11]. As bacterial AMR is currently the most important form of resistance in microbes, we will refer it to when we mention AMR throughout this review.

With Next-Generation genome Sequencing (NGS) technologies become widespread in recent years, they are commonly used in AMR surveillance studies either in clinical settings [12,13], or in the agriculture and food industry [14–17] and the environment [12,18,19]. In line with the importance of genomic surveillance of AMR, several annotation tools and databases has been developed for the analysis of ARG content of bacterial genomes or NGS metagenomic samples [20–35]. Table 1 presents some information on the most well known AMR databases.

**Table 1.** Well-known ARG databases. The table contains the most well known general ARG databases with additional information (the year of the last update of the database (Last modified), link address, where the database can be accessed (URL) and the publications associated with the database (References)).

| Database               | Last modified               | URL   | References |
|------------------------|-----------------------------|---|------------|
| ARDB                   | Archived, last update 2009  | <a href="https://ardb.cbcb.umd.edu/">https://ardb.cbcb.umd.edu/</a>   | [20]       |
| ARG-ANNOT              | Archived, last update: 2018 | not available   | [22]       |
| ARGminer*              | 2019                        | <a href="https://bench.cs.vt.edu/argminer/#/home">https://bench.cs.vt.edu/argminer/#/home</a>                               | [32]       |
| CARD*                  | 2021                        | <a href="https://card.mcmaster.ca/">https://card.mcmaster.ca/</a>   | [21,36,37] |
| FARME                  | 2019                        | <a href="http://staff.washington.edu/jwal-lace/farme/index.html">http://staff.washington.edu/jwal-lace/farme/index.html</a> | [29]       |
| MEGAres*               | 2019                        | <a href="https://megares.meglab.org/">https://megares.meglab.org/</a>   | [27,38]    |
| Mustard                | 2018                        | <a href="http://mgps.eu/Mustard/index.php?id=accueil">http://mgps.eu/Mustard/index.php?id=accueil</a>                       | [31]       |
| NDARO*                 | 2021                        | <a href="https://www.ncbi.nlm.nih.gov/pathogens/refgene/">https://www.ncbi.nlm.nih.gov/pathogens/refgene/</a>               | [34,39]    |
| PATRIC                 | 2017                        | <a href="https://patricbrc.org/">https://patricbrc.org/</a>   | [35,40,41] |
| ResFams                | 2015                        | <a href="http://www.dantaslab.org/resfams">http://www.dantaslab.org/resfams</a>   | [24]       |
| ResFinder/PointFinder* | 2021                        | <a href="https://cge.cbs.dtu.dk/services/ResFinder/">https://cge.cbs.dtu.dk/services/ResFinder/</a>                         | [30,33,42] |
| SARG*                  | 2019                        | <a href="https://smile.hku.hk/SARGs#">https://smile.hku.hk/SARGs#</a>   | [26,43]    |

\* Considered in this review

ARG databases can be divided to two major types [44], some of them contain species specific information (e.g. the MUBI database containing mutations conferring resistance in *Mycobacterium tuberculosis* [45]), while others focusing on ARGs from every sources (e.g. the CARD database [21]). However, ARG databases can differ not only in the covered species but on the type of AMR mechanism as well. Some database specialize only on acquired resistance genes, while others contain only mutations (e.g. the ResFinder [42] database focuses on acquired resistance genes, while the PointFinder [30] database from the same research group covers only AMR associated mutations). Unsurprisingly, however, there are databases with information on both AMR mechanisms (e.g. the CARD [21] or NDARO [34] databases).

The number of tools and databases focusing on AMR has rapidly grown in recent years, and many review articles were published trying to summarize the information on these resources. However, they put more emphasis on the different tools designed for ARG annotation rather than the databases supplying the information for these tasks [18,44,46–48].

As the performance of each tool heavily relies on the underlying database [36], it is important to understand the advantages and limitations of all databases available for the

research community. By understanding it, researchers can select the best database for their purpose. Furthermore, this knowledge can be important for choosing the best resource for developing new annotation tools as well. In this review we compare the most important ARG resources available today. Firstly, we review the structure of each database and then we directly compare them by their content. We present this comparison from the acquired and mutation based resistance mechanisms separately as databases can differ in these regards significantly. Researchers might prefer one mechanism of AMR more in their study, for example in a study investigating environmental ARGs with potential mobilization properties, acquired resistance genes might be the primary focus, whereas mutations can be more important in a clinical context [49].

## 2. Comparison of the structure of databases

### 2.1. Databases reviewed in this article

From the databases summarized in Table 1, ARDB, ARG-ANNOT and ResFams is not covered by this review as they are not actively updated (they haven't been update since 2008, 2018 and 2015, respectively). Furthermore, Mustard is also not reviewed here as it was constructed for a study of the gut resistome profiling of humans and wasn't dedicated as a comprehensive resource of ARGs [31]. FARME and PATRIC database are not covered here as well. FARME is based on several metagenomic studies, which were characterized based on their predicted ARG content and AMR phenotype, however those genes were not extensively validated and might contain false positives [29,48]. PATRIC is constructed for collecting genome sequence data and associated metadata of pathogen microorganisms [35], and necessarily relies on a specialized annotation system for the curation of the data. The ARG annotation pipeline employed by PATRIC is based on the NDARO and CARD databases as well as data from scientific literature, which was reannotated by experts [41], however this is not available on their FTP site. Therefore the following 6 databases are covered in detail only in this review: ARGminer, CARD, MEGARes, NDARO, ResFinder and SARG.

### 2.2. ARGminer

ARGminer is an ensemble database assembled from several independent ARG resources. It is based on the CARD [21], ARDB [20], DeepARG [50], MEGARes [27], ResFinder [42], and SARG [26] databses [32]. Only the acquired resistance genes were collected from these resources. After the acquisition of the sequences from these databases, they have clustered them to remove duplicates and annotated them by the best match from each of the above data resources. After the assignment of UniProt and GeneOntology metadata to the sequences, they guessed the best nomenclature of each gene names by a machine learning model. However, as several differences can be found between databases, they also utilize a crowdsourcing model to refine annotations (with a Trust-validation Filter to prevent misuse).

Furthermore, they have collected mobility and pathogen predictions by fitting the sequences to the ACLAME [51] and PATRIC [40] databases, respectively.

The database is periodically updated with the method described above and published after the verification of ARGminer evaluators. The date of the latest update of the database at the writing of this review is April 2019.

### 2.3. CARD

The Comprehensive Antibiotic Resistance Database (CARD) is a hand curated resource that is developed to cover the entire spectrum of ARGs [21]. Every ARG is included in the database based on three criteria. All ARG sequences must be available in the GenBank repository and increase the Minimal Inhibitory Concentration (MIC) in an

experimental validation setting which needs to be published in a peer-reviewed journal. Only a handful of historical  $\beta$ -lactam antibiotics are an exception from the above as they don't have an associated peer-reviewed publication [37]. The CARD database is built around an ontology-driven framework, where the resistance determinants and their associated metadata is recorded in the Antibiotic Resistance Ontology (ARO) network and even the sequences and the threshold used for their detection is stored in a specialized ontology (Model Ontology, MO) [36]. CARD contains resistance genes and resistance mutations as well, which are organized in a species specific manner. Furthermore, as CARD uses a strict curation procedure for incorporating genes, to increase sensitivity, they have developed a special database (the CARD Resistomes & Variants module) that contains *in silico* validated ARGs based on the genes stored in CARD [37]. The database is regularly updated based on reviewing the scientific literature by expert curators, whose work is augmented by a machine learning algorithm (CARD\*Shark) that sorts scientific publications based on reference for the process. The current version of CARD was updated in October 2021. It is important to note that CARD is freely accessible for academic researchers only and every use by commercial parties are only permitted with a written license.

#### 2.4. MEGARes

MEGARes is also an assembly of multiple resources in a way that is designed specifically for annotating metagenomic data [27]. The first version of the database was based on ResFinder [42], ARG-ANNOT [22], CARD [21] and the Lahey Clinic  $\beta$ -lactamase database curated by NCBI. During the update of the database to MEGARes 2.0 [38], further sequences were collected from the newer versions of CARD [36] and ResFinder [42] and the NCBI Bacterial Antimicrobial Resistance Reference Database [39]. Furthermore, MEGARes 2.0 also incorporates biocid and metal resistance genes derived from the BacMet database [52]. After they have removed the duplicates from the sequences collected from these resources, the genes were reannotated which revealed several overlapping genes between the ARG databases and BacMet. As the purpose of the database is to form a basis of the ARG annotation of metagenomic reads that can be used for read abundance based analysis, the annotations are stored in the form of an acyclic graph which avoids that one read or contig is assigned to multiple nodes [27]. The database contains antibiotic resistance genes and mutations as well, however the mutations are not ordered to microbial species due to the nature of the annotation graph. The current version of the database at the time of the writing of this review was updated at October 2019.

#### 2.5. NDARO

The National Database of Antibiotic Resistant Organisms (NDARO) is a comprehensive database dedicated to antibiotic resistance in the curation of NCBI [34]. The resistance genes are stored in The Reference Gene Catalog which predecessor was the Bacterial Antimicrobial Resistance Reference Gene Database, with the RefSeq PRJNA313047 BioProject (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA313047>) for storing the reference sequences [39]. This database was constructed from the ResFinder [42], CARD [36], RAC [53] and INTEGRALL [54] databases with extensive curation of the associated scientific literature. Since the expansion of the database in 2021, AMR mutations, general stress response genes and virulence genes are also curated within NDARO for the clinically important pathogens [34]. NDARO is updated regularly, the latest database version was released in December 2021.

#### 2.6. ResFinder/PointFinder

ResFinder [42] and PointFinder [30] are dedicated tools for acquired resistance genes and resistance mutations, respectively. These were separate AMR data resources, however since ResFinder 4.0, they are developed under the same project [33]. ResFinder was

originally developed on the basis of the Lahey Clinic  $\beta$ -lactamase database, ARDB [20], and an extensive literature review. To develop a more comprehensive resource of AMR determinants, the developers of ResFinder constructed a database dedicated to mutations conferring resistance only, named PointFinder. During the concatenation of the two databases under the ResFinder 4.0 project, not only an extensive expert curation was applied to the data, but phenotype prediction tables were also constructed to help researchers connecting genotype information with potential phenotypic traits. With regular updates, the latest version of ResFinder and PointFinder was released in September and February 2021, respectively.

### 2.7. SARG


The Structured ARG reference database (SARG) is a hierarchically constructed database [26] based on the CARD [21] and ARDB [20] data resources. They only retained the acquired resistance genes from these databases, and after duplicate removal, they have ordered the genes to a two-level hierarchical architecture. The highest level of this hierarchy is the type of the resistance indicating the antibiotic that the genes confer resistance to, while the lower level is the class of the genes. In 2018 the developers of SARG expanded the database by ARG homologs found by aligning the NCBI nt database to SARG [43]. They are regularly updating the database in a similar manner, with the latest aired in January 2022, however it did not introduced any new ARGs since the 2019 version. SARG, similarly to CARD only accessible freely for academic purposes, and for commercial use a written permit is necessary.

## 3. Comparison of the database contents

### 3.1. Number of sequences and ARGs in the databases

To compare the ARG content of the different databases we first matched the number of sequences stored in them and the associated count of unique genes (Figure 1). Figure 1 only shows resistance genes and biocid resistance genes (to maintain comparability between databases), and virulence or metal resistance genes were omitted. In the case of ARGminer we have found several different nomenclature forms of the same ARGs, which is not surprising as one of the main goal of the database was to collect and standardize this information with the aid of crowdsourcing. However as we did not intended to make such standardization through this review it might be possible that the same gene was counted multiple times in the case of ARGminer in Figure 1. We tried to reduce the risk of this bias by converting gene names to lowercase when comparing them, as usually the ARG name nomenclature differences concerning only the casing of the letters. Furthermore, we have found 13 and 3 duplicate sequences in the NDARO and MEGARes databases, respectively (the number of sequences in Figure 1 is corrected for the presence of duplicates). The presence of duplicate genes and corresponding sequences in the database might cause overestimation of those genes if the user does not pay enough attention on thoroughly reviewing the results. In Figure 1 a clear difference can be observed between SARG, ARGminer and MEGARes databases in contrast with NDARO, ResFinder and CARD in the relationship between the number of unique sequences and corresponding genes. The major differences between the two groups is that CARD, NDARO and ResFinder are primer databases mainly based on the extensive review of the corresponding scientific literature [37], while the others are assembled from other data resources. One might expect that with keeping one reference sequence for each genes, these databases are prone to producing false negatives in homology searches, however, this is overcome in CARD with the use of individual detection threshold for genes stored in the Model Ontology [36] and the use of HMMs (Hidden Markov Models) in the detection software of NDARO when no high confidence match can be retrieved [34].






**Figure 1.** ARG and sequence content of the databases. Only antibiotic and biocid resistance genes were considered for the plot. For each databases on the x axis, the number of unique sequences and the corresponding number of unique genes were determined. The y axis represents the number of genes and sequences. Red bars show the gene number while blue bars represent the number of sequences stored in each database.

### 3.2. Gene count of antibiotic classes in each database

Figure 2 shows the differences in the number of antibiotic resistance genes (without those conferring resistance through mutations) associated with the antibiotic classes stored in the respective database for CARD and ResFinder. We have selected these two due to the extensive differences in the depth of the antibiotic classification. For the rest of the databases (MEGARes, NDARO and SARG) the same figures can be found in the Supplementary File 1. In the case of ARGminer we couldn't construct such figure as notable differences were found in some cases between the antibiotic classification of different records for the same genes in. In either of the above figures the respective classification scheme of each database is used. As one would expect, aminoglycoside and  $\beta$ -lactam antibiotics are the most popular categories in either of the databases. However there is a significant difference in the classification depth of  $\beta$ -lactams between the CARD and other resources. In CARD, separate  $\beta$ -lactam groups have their respective categories (such as penems, penams, carbapenems, cephalosporins etc.), while others label them only as  $\beta$ -lactams. Furthermore, the presence of several collective category in the MEGARes database is notable (e.g. Multi-drug resistance or Drug and biocide resistance, etc.). The reason for the presence of such categories is the acyclic form of the MEGARes annotation graph, which not allows linking the same gene to multiple groups. These figures clearly show that the most comprehensive antibiotic classification of the genes can be expected in the case of the CARD database, however the differences also emphasize that expert knowledge is important for understanding the results of ARG annotation and one can not expect to rely entirely on the output of a database.




**Figure 2.** Number of unique genes for each antibiotic class stored in CARD and ResFinder. Bars represents the number of gene in each unique antibiotic or biocid categories, where colors are associated with the specific antibiotics themselves. As one gene can confer resistance to multiple antibiotics, it is possible that the same gene is counted for two or more antibiotics. The plots show the data for CARD (A) and ResFinder (B), respectively.

### 3.3. Microbial genus with corresponding AMR mutations in the databases

Next we have compared the number of genes conferring resistance through mutations for microbial species in each database (Figure 3). Among the databases covered in depth in this review, only CARD, MEGARes, NDARO and PointFinder (element of ResFinder 4.0) comprise such information. Although MEGARes also has information on mutations causing resistance, connecting them to species is not applicable in that case due to the nature of the annotation architecture. For comparing the microbes for which data is stored in each database we had to find a taxonomical level that can achieve a standardized comparison between all databases. We have decided to count genes in the genus level. We had to diverge from this principle only in one case, where the arbitrary group

Propionibacteria had to be used instead of the corresponding genus. For simplicity, despite of this exception, we further refer to the groups of microbes used for the classification in Figure 3 as genus.

It is upfront in Figure 3 that CARD contains mutations for the highest number genus (37 genus) between the three databases, and it even has 59 non species classified genes as well. In contrast, NDARO store genes for 11 genus while PointFinder only for 10. Not every genus considered by the databases belong to bacteria. CARD stores 10 genes for *Chlamydomonas* algae and 5 for the archeal genus *Halobacterium*, while PointFinder has 6 genes for *Plasmodium* protozoas. Those genus considered by the NDARO and PointFinder databases are primarily human pathogens. Especially those among the critically important bacteria for human health, determined by the WHO in 2017 [55–57] (ESKAPE pathogens: *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* species). Furthermore, NDARO has 11 genes enrolled to the *Salmonella* genus, one of the most important food born pathogen and often associated with resistance conferring mutations [58]. PointFinder has a significant collection of *Mycobacterium* genes (36 genes), which is even more notable in CARD with 1168 genes associated with this genus. Pathogens from the *Mycobacterium* genus, especially *M. tuberculosis* is one of the most important among disease causing bacteria that develops resistance through mutations. Furthermore, as it needs long incubation times for culturing, whole-genome sequencing based approaches are accountable alternatives [59].



**Figure 3.** Number of genes conferring resistance through mutations for each microbial genus in CARD, NDARO and MEGARes databases. Genes conferring resistance through mutations was calculated at the genus level for each microbes stored in each database. Only one group could not be summarized at the genus level (Propionibacteria). Microbial genus is on the x axis and the number of genes associated with each group in the database is represented by the y axis. Rows show the data separately for each database. Columns are colored by the microbial genus.



CARD database extremely differs from the other data resources reviewed here, due to the high number of microbial genus it collects data for and the number of genes it stores considering AMR mutations. These properties makes it especially suitable for AMR mutation screening in a wide variety study settings. Even in the case of environmental AMR surveillance as it stores mutations for typical environmental genus such as *Thermus* or *Halobacterium*. An especially high number of genes is stored in the database for the *Mycobacterium* genus (37 genes), which is in the forefront as a potential bacterium for degrading plastic pollutants [60].

#### 4. Conclusions

Previously several ARG database were constructed to form the bases of ARG annotation of whole-genome sequencing and metagenome samples. With the advent of NGS, their significance is even more profound, and they became an important augmentation of previous phenotypic screening based studies. We compared the accessible and regularly updated ARG databases in this review, that have new versions released lately. The main focus in this review was on the architecture and content of the different databases, in contrast with previous studies mainly focusing on the tools used for annotation. However, understanding how databases are constructed and the differences between them is crucial for every researcher in the field of AMR, so they can use the most powerful tool for their research question. Based on the differences outlined in this review it seems that CARD and NDARO is prominent among the databases. NDARO contains the most acquired resistance genes, however CARD comprises of a similarly high number of genes, making both of them a suitable tool for ARG annotation. In the case of mutations conferring resistance, however CARD dominates other tools. We advise that in cases where mutations or both type of resistance is considered, CARD should be the number one data resource. Furthermore, the differences in antibiotic classification of the databases emphasize the importance of expert knowledge for interpreting the results. Moreover, as some databases are accessible for non academic parties only with a written permit, it is important for one to be familiar with the terms of using these resources.

#### 5. Future Perspective

We believe that during the evaluation of the performance of different ARG annotation tools, differences in the underlying database should also be considered. Moreover, as major differences can be observed in ARG nomenclature between databases, a standardization procedure would be advantageous for enabling direct comparisons between results generated from different resources.

**Supplementary Materials:** File S1: Gene counts for antibiotic classes in MEGARes, NDARO and SARG databases .

**Author Contributions:** Conceptualization, M.P. and N.S.; writing—original draft preparation, M.P.; writing—review and editing, N.S.; visualization, M.P.; supervision, N.S.. All authors have read and agreed to the published version of the manuscript.

**Funding:** The project is supported by the European Union and co-financed by the European Social Fund (No. EFOP-3.6.3-VEKOP-16-2017-00005). It has also received funding from the European Union's Horizon 2020 research and innovation program under Grant Agreement No. 874735 (VEO).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. O'Neill, J. Tackling Drug-Resistant Infections Globally: Final Report and Recommendations; Review on Antimicrobial Resistance. London, UK, **2016**; pp. 1–84.
2. Centers for Disease Control and Prevention (U.S.) Antibiotic Resistance Threats in the United States, 2019. Centers for Disease Control and Prevention (U.S.), **2019**.
3. Cassini, A.; Högberg, L.D.; Plachouras, D.; Quattrocchi, A.; Hoxha, A.; Simonsen, G.S.; Colomb-Cotinat, M.; Kretzschmar, M.E.; Devleeschauwer, B.; Cecchini, M.; et al. Attributable Deaths and Disability-Adjusted Life-Years Caused by Infections with Antibiotic-Resistant Bacteria in the EU and the European Economic Area in 2015: A Population-Level Modelling Analysis. *The Lancet Infectious Diseases* **2019**, *19*, 56–66, doi:10.1016/S1473-3099(18)30605-4.
4. Dadgostar, P. Antimicrobial Resistance: Implications and Costs. *IDR* **2019**, Volume 12, 3903–3910, doi:10.2147/IDR.S234610.
5. Baquero, F.; Coque, T.M.; Martínez, J.-L.; Aracil-Gisbert, S.; Lanza, V.F. Gene Transmission in the One Health Microbiome and the Channels of Antimicrobial Resistance. *Front. Microbiol.* **2019**, *10*, 2892, doi:10.3389/fmicb.2019.02892.
6. Singh, K.S.; Anand, S.; Dholpuria, S.; Sharma, J.K.; Blankenfeldt, W.; Shouche, Y. Antimicrobial Resistance Dynamics and the One-Health Strategy: A Review. *Environ Chem Lett* **2021**, *19*, 2995–3007, doi:10.1007/s10311-021-01238-3.
7. Blair, J.M.A.; Webber, M.A.; Baylay, A.J.; Ogbolu, D.O.; Piddock, L.J.V. Molecular Mechanisms of Antibiotic Resistance. *Nat Rev Microbiol* **2015**, *13*, 42–51, doi:10.1038/nrmicro3380.
8. Munita, J.M.; Arias, C.A. Mechanisms of Antibiotic Resistance. *Microbiol Spectr* **2016**, *4*, doi:10.1128/microbiolspec.VMBF-0016-2015.
9. Brown, E.D.; Wright, G.D. Antibacterial Drug Discovery in the Resistance Era. *Nature* **2016**, *529*, 336–343, doi:10.1038/nature17042.
10. Perry, J.; Waglechner, N.; Wright, G. The Prehistory of Antibiotic Resistance. *Cold Spring Harb Perspect Med* **2016**, *6*, a025197, doi:10.1101/cshperspect.a025197.
11. Larsson, D.G.J.; Flach, C.-F. Antibiotic Resistance in the Environment. *Nat Rev Microbiol* **2021**, doi:10.1038/s41579-021-00649-x.
12. Crofts, T.S.; Gasparrini, A.J.; Dantas, G. Next-Generation Approaches to Understand and Combat the Antibiotic Resistance. *Nat Rev Microbiol* **2017**, *15*, 422–434, doi:10.1038/nrmicro.2017.28.
13. Saracino, I.M.; Pavoni, M.; Zullo, A.; Fiorini, G.; Lazzarotto, T.; Borghi, C.; Vaira, D. Next Generation Sequencing for the Prediction of the Antibiotic Resistance in *Helicobacter pylori*: A Literature Review. *Antibiotics* **2021**, *10*, 437, doi:10.3390/antibiotics10040437.
14. Caniça, M.; Manageiro, V.; Abriouel, H.; Moran-Gilad, J.; Franz, C.M.A.P. Antibiotic Resistance in Foodborne Bacteria. *Trends in Food Science & Technology* **2019**, *84*, 41–44, doi:10.1016/j.tifs.2018.08.001.
15. Thanner, S.; Drissner, D.; Walsh, F. Antimicrobial Resistance in Agriculture. *mBio* **2016**, *7*, doi:10.1128/mBio.02227-15.
16. Tóth, A.G.; Csabai, I.; Krikó, E.; Tózsér, D.; Maróti, G.; Patai, Á.V.; Makrai, L.; Szita, G.; Solymosi, N. Antimicrobial Resistance Genes in Raw Milk for Human Consumption. *Sci Rep* **2020**, *10*, 7464, doi:10.1038/s41598-020-63675-4.
17. Tóth, A.G.; Csabai, I.; Maróti, G.; Jerzsele, Á.; Dubecz, A.; Patai, Á.V.; Judge, M.F.; Nagy, S.Á.; Makrai, L.; Bányai, K.; et al. A Glimpse of Antimicrobial Resistance Gene Diversity in Kefir and Yoghurt. *Sci Rep* **2020**, *10*, 22458, doi:10.1038/s41598-020-80444-5.
18. The Global Sewage Surveillance project consortium; Hendriksen, R.S.; Munk, P.; Njage, P.; van Bunnik, B.; McNally, L.; Lukjancenko, O.; Röder, T.; Nieuwenhuijse, D.; Pedersen, S.K.; et al. Global Monitoring of Antimicrobial Resistance Based on Metagenomics Analyses of Urban Sewage. *Nat Commun* **2019**, *10*, 1124, doi:10.1038/s41467-019-08853-3.
19. Garner, E.; Davis, B.C.; Milligan, E.; Blair, M.F.; Keenum, I.; Maile-Moskowitz, A.; Pan, J.; Gnegy, M.; Liguori, K.; Gupta, S.; et al. Next Generation Sequencing Approaches to Evaluate Water and Wastewater Quality. *Water Research* **2021**, *194*, 116907, doi:10.1016/j.watres.2021.116907.
20. Liu, B.; Pop, M. ARDB--Antibiotic Resistance Genes Database. *Nucleic Acids Research* **2009**, *37*, D443–D447, doi:10.1093/nar/gkn656.
21. McArthur, A.G.; Waglechner, N.; Nizam, F.; Yan, A.; Azad, M.A.; Baylay, A.J.; Bhullar, K.; Canova, M.J.; De Pascale, G.; Ejim, L.; et al. The Comprehensive Antibiotic Resistance Database. *Antimicrob. Agents Chemother.* **2013**, *57*, 3348–3357, doi:10.1128/AAC.00419-13.
22. Gupta, S.K.; Padmanabhan, B.R.; Diene, S.M.; Lopez-Rojas, R.; Kempf, M.; Landraud, L.; Rolain, J.-M. ARG-ANNOT, a New Bioinformatic Tool To Discover Antibiotic Resistance Genes in Bacterial Genomes. *Antimicrob. Agents Chemother.* **2014**, *58*, 212–220, doi:10.1128/AAC.01310-13.
23. Kaminski, J.; Gibson, M.K.; Franzosa, E.A.; Segata, N.; Dantas, G.; Huttenhower, C. High-Specificity Targeted Functional Profiling in Microbial Communities with ShortBRED. *PLoS Comput Biol* **2015**, *11*, e1004557, doi:10.1371/journal.pcbi.1004557.
24. Gibson, M.K.; Forsberg, K.J.; Dantas, G. Improved Annotation of Antibiotic Resistance Determinants Reveals Microbial Resistomes Cluster by Ecology. *ISME J* **2015**, *9*, 207–216, doi:10.1038/ismej.2014.106.
25. de Man, T.J.B.; Limbago, B.M. SSTAR, a Stand-Alone Easy-To-Use Antimicrobial Resistance Gene Predictor. *mSphere* **2016**, *1*, doi:10.1128/mSphere.00050-15.

26. Yang, Y.; Jiang, X.; Chai, B.; Ma, L.; Li, B.; Zhang, A.; Cole, J.R.; Tiedje, J.M.; Zhang, T. ARGs-OAP: Online Analysis Pipeline for Antibiotic Resistance Genes Detection from Metagenomic Data Using an Integrated Structured ARG-Database. *Bioinformatics* **2016**, *32*, 2346–2351, doi:10.1093/bioinformatics/btw136.
27. Lakin, S.M.; Dean, C.; Noyes, N.R.; Dettenwanger, A.; Ross, A.S.; Doster, E.; Rovira, P.; Abdo, Z.; Jones, K.L.; Ruiz, J.; et al. MEGARes: An Antimicrobial Resistance Database for High Throughput Sequencing. *Nucleic Acids Res* **2017**, *45*, D574–D580, doi:10.1093/nar/gkw1009.
28. Hunt, M.; Mather, A.E.; Sánchez-Busó, L.; Page, A.J.; Parkhill, J.; Keane, J.A.; Harris, S.R. ARIBA: Rapid Antimicrobial Resistance Genotyping Directly from Sequencing Reads. *Microbial Genomics* **2017**, *3*, doi:10.1099/mgen.0.000131.
29. Wallace, J.C.; Port, J.A.; Smith, M.N.; Faustman, E.M. FARME DB: A Functional Antibiotic Resistance Element Database. *Database (Oxford)* **2017**, *2017*, baw165, doi:10.1093/database/baw165.
30. Zankari, E.; Allesøe, R.; Joensen, K.G.; Cavaco, L.M.; Lund, O.; Aarestrup, F.M. PointFinder: A Novel Web Tool for WGS-Based Detection of Antimicrobial Resistance Associated with Chromosomal Point Mutations in Bacterial Pathogens. *J Antimicrob Chemother* **2017**, *72*, 2764–2768, doi:10.1093/jac/dkx217.
31. Ruppé, E.; Ghoulane, A.; Tap, J.; Pons, N.; Alvarez, A.-S.; Maziers, N.; Cuesta, T.; Hernando-Amado, S.; Clares, I.; Martínez, J.L.; et al. Prediction of the Intestinal Resistome by a Three-Dimensional Structure-Based Method. *Nat Microbiol* **2019**, *4*, 112–123, doi:10.1038/s41564-018-0292-6.
32. Arango-Argoty, G.A.; Guron, G.K.P.; Garner, E.; Riquelme, M.V.; Heath, L.S.; Pruden, A.; Vikesland, P.J.; Zhang, L. ARGminer: A Web Platform for the Crowdsourcing-Based Curation of Antibiotic Resistance Genes. *Bioinformatics* **2020**, *36*, 2966–2973, doi:10.1093/bioinformatics/btaa095.
33. Bortolaia, V.; Kaas, R.S.; Ruppe, E.; Roberts, M.C.; Schwarz, S.; Cattoir, V.; Philippon, A.; Allesøe, R.L.; Rebelo, A.R.; Florensa, A.F.; et al. ResFinder 4.0 for Predictions of Phenotypes from Genotypes. *Journal of Antimicrobial Chemotherapy* **2020**, *75*, 3491–3500, doi:10.1093/jac/dkaa345.
34. Feldgarden, M.; Brover, V.; Gonzalez-Escalona, N.; Frye, J.G.; Haendiges, J.; Haft, D.H.; Hoffmann, M.; Pettengill, J.B.; Prasad, A.B.; Tillman, G.E.; et al. AMRFinderPlus and the Reference Gene Catalog Facilitate Examination of the Genomic Links among Antimicrobial Resistance, Stress Response, and Virulence. *Sci Rep* **2021**, *11*, 12728, doi:10.1038/s41598-021-91456-0.
35. Davis, J.J.; Wattam, A.R.; Aziz, R.K.; Brettin, T.; Butler, R.; Butler, R.M.; Chlenski, P.; Conrad, N.; Dickerman, A.; Dietrich, E.M.; et al. The PATRIC Bioinformatics Resource Center: Expanding Data and Analysis Capabilities. *Nucleic Acids Research* **2019**, *47*, gkz943, doi:10.1093/nar/gkz943.
36. Jia, B.; Raphenya, A.R.; Alcock, B.; Waglechner, N.; Guo, P.; Tsang, K.K.; Lago, B.A.; Dave, B.M.; Pereira, S.; Sharma, A.N.; et al. CARD 2017: Expansion and Model-Centric Curation of the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Res* **2017**, *45*, D566–D573, doi:10.1093/nar/gkw1004.
37. Alcock, B.P.; Raphenya, A.R.; Lau, T.T.Y.; Tsang, K.K.; Bouchard, M.; Edalatmand, A.; Huynh, W.; Nguyen, A.-L.V.; Cheng, A.A.; Liu, S.; et al. CARD 2020: Antibiotic Resistome Surveillance with the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Research* **2019**, *47*, gkz935, doi:10.1093/nar/gkz935.
38. Doster, E.; Lakin, S.M.; Dean, C.J.; Wolfe, C.; Young, J.G.; Boucher, C.; Belk, K.E.; Noyes, N.R.; Morley, P.S. MEGARes 2.0: A Database for Classification of Antimicrobial Drug, Biocide and Metal Resistance Determinants in Metagenomic Sequence Data. *Nucleic Acids Research* **2020**, *48*, D561–D569, doi:10.1093/nar/gkz1010.
39. Feldgarden, M.; Brover, V.; Haft, D.H.; Prasad, A.B.; Slotta, D.J.; Tolstoy, I.; Tyson, G.H.; Zhao, S.; Hsu, C.-H.; McDermott, P.F.; et al. Validating the AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates. *Antimicrob Agents Chemother* **2019**, *63*, doi:10.1128/AAC.00483-19.
40. Wattam, A.R.; Abraham, D.; Dalay, O.; Disz, T.L.; Driscoll, T.; Gabbard, J.L.; Gillespie, J.J.; Gough, R.; Hix, D.; Kenyon, R.; et al. PATRIC, the Bacterial Bioinformatics Database and Analysis Resource. *Nucl. Acids Res.* **2014**, *42*, D581–D591, doi:10.1093/nar/gkt1099.
41. Antonopoulos, D.A.; Assaf, R.; Aziz, R.K.; Brettin, T.; Bun, C.; Conrad, N.; Davis, J.J.; Dietrich, E.M.; Disz, T.; Gerdes, S.; et al. PATRIC as a Unique Resource for Studying Antimicrobial Resistance. *Briefings in Bioinformatics* **2019**, *20*, 1094–1102, doi:10.1093/bib/bbx083.
42. Zankari, E.; Hasman, H.; Cosentino, S.; Vestergaard, M.; Rasmussen, S.; Lund, O.; Aarestrup, F.M.; Larsen, M.V. Identification of Acquired Antimicrobial Resistance Genes. *Journal of Antimicrobial Chemotherapy* **2012**, *67*, 2640–2644, doi:10.1093/jac/dks261.
43. Yin, X.; Jiang, X.-T.; Chai, B.; Li, L.; Yang, Y.; Cole, J.R.; Tiedje, J.M.; Zhang, T. ARGs-OAP v2.0 with an Expanded SARG Database and Hidden Markov Models for Enhancement Characterization and Quantification of Antibiotic Resistance Genes in Environmental Metagenomes. *Bioinformatics* **2018**, *34*, 2263–2270, doi:10.1093/bioinformatics/bty053.
44. Boolchandani, M.; D'Souza, A.W.; Dantas, G. Sequencing-Based Methods and Resources to Study Antimicrobial Resistance. *Nat Rev Genet* **2019**, doi:10.1038/s41576-019-0108-4.
45. Flandrois, J.-P.; Lina, G.; Dumitrescu, O. MUBII-TB-DB: A Database of Mutations Associated with Antibiotic Resistance in *Mycobacterium tuberculosis*. *BMC Bioinformatics* **2014**, *15*, 107, doi:10.1186/1471-2105-15-107.
46. McArthur, A.G.; Tsang, K.K. Antimicrobial Resistance Surveillance in the Genomic Age: AMR Surveillance in the Genomic Age. *Ann. N.Y. Acad. Sci.* **2017**, *1388*, 78–91, doi:10.1111/nyas.13289.

47. Van Camp, P.-J.; Haslam, D.B.; Porollo, A. Bioinformatics Approaches to the Understanding of Molecular Mechanisms in Antimicrobial Resistance. *IJMS* **2020**, *21*, 1363, doi:10.3390/ijms21041363.
48. Lal Gupta, C.; Kumar Tiwari, R.; Cytryn, E. Platforms for Elucidating Antibiotic Resistance in Single Genomes and Complex Metagenomes. *Environment International* **2020**, *138*, 105667, doi:10.1016/j.envint.2020.105667.
49. Woodford, N.; Ellington, M.J. The Emergence of Antibiotic Resistance by Mutation. *Clinical Microbiology and Infection* **2007**, *13*, 5–18, doi:10.1111/j.1469-0691.2006.01492.x.
50. Arango-Argoty, G.; Garner, E.; Pruden, A.; Heath, L.S.; Vikesland, P.; Zhang, L. DeepARG: A Deep Learning Approach for Predicting Antibiotic Resistance Genes from Metagenomic Data. *Microbiome* **2018**, *6*, 23, doi:10.1186/s40168-018-0401-z.
51. Leplae, R. ACLAME: A CLAssification of Mobile Genetic Elements. *Nucleic Acids Research* **2004**, *32*, 45D – 49, doi:10.1093/nar/gkh084.
52. Pal, C.; Bengtsson-Palme, J.; Rensing, C.; Kristiansson, E.; Larsson, D.G.J. BacMet: Antibacterial Biocide and Metal Resistance Genes Database. *Nucl. Acids Res.* **2014**, *42*, D737–D743, doi:10.1093/nar/gkt1252.
53. Tsafnat, G.; Coptly, J.; Partridge, S.R. RAC: Repository of Antibiotic Resistance Cassettes. *Database* **2011**, *2011*, bar054–bar054, doi:10.1093/database/bar054.
54. Moura, A.; Soares, M.; Pereira, C.; Leitao, N.; Henriques, I.; Correia, A. INTEGRALL: A Database and Search Engine for Integrations, Integrases and Gene Cassettes. *Bioinformatics* **2009**, *25*, 1096–1098, doi:10.1093/bioinformatics/btp105.
55. Tacconelli, E. GLOBAL PRIORITY LIST OF ANTIBIOTIC-RESISTANT BACTERIA TO GUIDE RESEARCH, DISCOVERY, AND DEVELOPMENT OF NEW ANTIBIOTICS. 7.
56. De Oliveira, D.M.P.; Forde, B.M.; Kidd, T.J.; Harris, P.N.A.; Schembri, M.A.; Beatson, S.A.; Paterson, D.L.; Walker, M.J. Antimicrobial Resistance in ESKAPE Pathogens. *Clin Microbiol Rev* **2020**, *33*, doi:10.1128/CMR.00181-19.
57. Mancuso, G.; Midiri, A.; Gerace, E.; Biondo, C. Bacterial Antibiotic Resistance: The Most Critical Pathogens. *Pathogens* **2021**, *10*, 1310, doi:10.3390/pathogens10101310.
58. Hur, J.; Jawale, C.; Lee, J.H. Antimicrobial Resistance of Salmonella Isolated from Food Animals: A Review. *Food Research International* **2012**, *45*, 819–830, doi:10.1016/j.foodres.2011.05.014.
59. Dookie, N.; Rambaran, S.; Padayatchi, N.; Mahomed, S.; Naidoo, K. Evolution of Drug Resistance in *Mycobacterium tuberculosis*: A Review on the Molecular Determinants of Resistance and Implications for Personalized Care. *Journal of Antimicrobial Chemotherapy* **2018**, *73*, 1138–1151, doi:10.1093/jac/dkx506.
60. Cortés-Albayay, C.; Sangal, V.; Klenk, H.-P.; Nouioui, I. Comparative Genomic Study of Vinyl Chloride Cluster and Description of Novel Species, *Mycolicibacterium vinylchloridicum* Sp. Nov. *Front. Microbiol.* **2021**, *12*, 767895, doi:10.3389/fmicb.2021.767895.