

Article

Gravity-Assist might be a Solution to Save Earth from Global Warming

Sohrab Rahvar^{1,†}  0000-0002-7084-5725

¹ Department of Physics, Sharif University of Technology, P. O. Box 11365-9161, Tehran, Iran rahvar@sharif.edu

Abstract: Global warming is one of the problems of human civilization and decarbonization policy is the main solution to this problem. In this work, we propose an alternative method of using the gravity-assist by the asteroids to increase the orbital distance of the Earth from the Sun. We can manipulate the orbit of asteroids in the asteroid belt by solar sailing and propulsion engines to guide them towards the Mars orbit and a gravitational scattering can put asteroids in a favorable direction to provide an energy loss scattering from the Earth. The result would be increasing the orbital distance of the earth and consequently cooling down the Earth's temperature. We calculate the increase in the orbital distance of the earth for each scattering and investigate the feasibility of performing this project.

Keywords: Dynamics; Solar System; Global Warming

1. Introduction

The monitoring of the Earth's climate shows a significant change in temperature and climate pattern due to the emission of greenhouse gases by humans [1]. The two gases of carbon dioxide and methane as a result of fuel burning are the main components of the greenhouse gases causing the climate change. In recent years the temperature rise is accelerated as a result of feedback from the melting of ices and decreasing of the light reflection from sun and increasing of the water vapor in the atmosphere [2]. Figure (1) shows the trend of temperature increase in terms of the year which compares the measured temperature of earth with the simulations [3]. In recent years the increase in extreme events of weather also shows a direct correlation with the increase of the temperature of the atmosphere and oceans [4].

The present policy to prevent the global temperature is mainly based on decreasing fossil fuel consumption and investment in green sources of energy. The policy of decarbonization is based on the rapid development of renewable energies [5]. The Paris agreement as an international implementation plans that the mean global temperature doesn't rise beyond 2° C above pre-industrial levels.

In this work, we propose an alternative method for decreasing the global temperature by changing the orbital distance of Earth from the Sun. This operation can be done by the gravity-assist mechanism with the flyby motion of the asteroids close to the earth where the energy loss of asteroids can increase the orbital distance of the Earth from the Sun and results in a decrease in the Earth temperature. While this plan may look like an out-of-practical idea, we will show the feasibility of this project at least for future technology.

In Section (2), we introduce the gravity-assist and provide the physical mechanism to use the asteroids to change the mechanical energy of the Earth. In section (3) we discuss the dependence of mechanical energy of the Earth to the global temperature and calculate the temperature change on earth for an asteroid scattering with a given mass. In Section (4), we provide a practical plan for the asteroid scattering from the earth. The Discussion is given in (5).

2. Gravity-assist and energy loss process

The gravity-assist is suggested by Yuri Kondratyuk in 1938 for accelerating a spacecraft traveling between the planets of the solar system using the gravity of planets as the slingshot



Citation: Gravity-Assist might be a Solution to Save Earth from Global Warming. *Preprints* 2022, 1, 0. <https://doi.org/>

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 1. Change in global surface temperature (annual average) as observed and simulated using human & natural and only natural factors (both 1850-2020), adapted from [3]

[6]. In gravity assist the relative speed of the spacecraft before and after the gravitational scattering with respect to a planet is the same, however since the planet is moving with respect to the sun, the spacecraft can accelerate or decelerate with respect to the solar frame. The gravity assist maneuver was first used in 1959 by the Soviet probe Luna 3 for monitoring the Moon. The other interplanetary probes such as the Pioneer program, Mariner 10, Voyager program, Galileo, Ulysses, and many other programs used gravity-assist to reach the other planets at the edge of the solar system. The gravity assist not only can change the kinetic energy of the spacecraft also can change the energy of photons or the frequency of light during the gravitational scattering so-called gravitational lensing [7].

Here we want to use the gravity-assist mechanism for the scattering of asteroids from the Earth and instead of energy-gain by the asteroid, we manage the energy loss of asteroid where in this case the energy of asteroid transfers to the mechanical energy of Earth, orbiting around the Sun. This idea also discussed in [8,9]. The result of this scattering would be increasing the orbital distance from the sun and consequently cooling down the temperature of the planet.

In the following, we investigate the scattering process of an asteroid from Earth. Let us take m_a as the mass of an asteroid and M_e as the mass of the earth. \mathbf{v}_i and \mathbf{v}_f are the initial and final velocity vectors of the asteroid (as a result of scattering) with respect to the Earth and \mathbf{v}'_i and \mathbf{v}'_f are the corresponding velocities with respect to the Sun. Assuming that the mass of the asteroid is much smaller than earth (i.e. $m_a \ll M_e$), the speed of the asteroid before and after the scattering with respect to the Earth is equal (i.e. $|\mathbf{v}_i| = |\mathbf{v}_f|$). However the velocity of asteroid with respect to the sun before the scattering is $\mathbf{v}'_i = \mathbf{v}_i + \mathbf{v}_e$ and after the scattering is $\mathbf{v}'_f = \mathbf{v}_f + \mathbf{v}_e$. Here \mathbf{v}_e is the velocity of the earth with respect to the sun. We note that the incoming and outgoing speed of asteroid with respect to the solar coordinate are not equal.

The energy difference of asteroid after the scattering compare to that before the scattering in the frame of sun is given by $\Delta E_a = m_a \mathbf{v}_e \cdot (\mathbf{v}_f - \mathbf{v}_i)$. According to the conservation of energy, the energy gain of earth would be $\Delta E_e = -\mathbf{v}_e \cdot (\mathbf{p}_f - \mathbf{p}_i)$ where $\mathbf{p}_f = m_a \mathbf{v}_f$ and $\mathbf{p}_i = m_a \mathbf{v}_i$ are the momentum of asteroid with respect to the Earth frame where $|\mathbf{p}_i| = |\mathbf{p}_f|$. Also the momentum of Earth would change by the amount of




Figure 2. Schematic figure for the scattering where in this case $\Delta E_e > 0$ and earth can gain energy from this scattering. \mathbf{v}_e (with black arrow) is the velocity of earth. \mathbf{v}_i and \mathbf{v}_f (in blue arrows) are the relative velocity of asteroid with respect to the earth before and after the scattering. The red arrow is the difference of $\mathbf{v}_f - \mathbf{v}_i$ and the grey axis are the symmetric coordinate system with respect to \mathbf{v}_i and \mathbf{v}_f .

$\Delta \mathbf{P}_e = -(\mathbf{p}_f - \mathbf{p}_i)$. In order to calculate the energy and momentum change of earth, we project the velocities on x and y axis in Fig (2) as follows:

$$\mathbf{v}_e = v_e(\hat{i} \sin \phi + \hat{j} \cos \phi), \quad (1)$$

$$\mathbf{v}_i = v_i(-\hat{i} \sin \theta + \hat{j} \cos \theta), \quad (2)$$

$$\mathbf{v}_f = v_i(-\hat{i} \sin \theta - \hat{j} \cos \theta), \quad (3)$$

where the y axis of this coordinate system is designed to be symmetric with respect to the incoming and outgoing velocities of the asteroid with respect to the earth. The θ angle is the direction of incoming velocity with the y -axis and the ϕ angle is earth's velocity with the y -axis. Substituting the velocities in the energy and momentum gain of earth results in

$$\Delta E_e = 2m_a v_i v_e \cos \theta \cos \phi, \quad (4)$$

$$\Delta \mathbf{p}_e = 2m_a v_i \cos \theta \hat{j}. \quad (5)$$

Using the total mechanical energy of earth around the sun $E_e = -GM_\odot M_e / 2R_e$, then the relative energy change of Earth orbiting around the sun would be

$$\frac{\Delta E_e}{|E_e|} = 4\left(\frac{m_a}{M_e}\right)\left(\frac{v_i}{v_e}\right) \cos \theta \cos \phi. \quad (6)$$

3. Temperature change of earth as a result of mechanical energy change

The change in the total energy of earth results in the change in the orbital distance as $\Delta E/|E| = \Delta R/R$. Since the power of light received by the earth depends to distance as $S \propto 1/R^2$, we can conclude that $\Delta S/S_0 = -2\Delta E_e/|E_e|$. Then from equation (6), the relative change in the power of sun received by the earth would be

$$\frac{\Delta S}{S_0} = -0.5 \times 10^{-9} \left(\frac{m_a}{10^{15} \text{kg}}\right) \left(\frac{v_i}{11 \text{km/s}}\right) \cos \theta \cos \phi. \quad (7)$$

The change in the flux of light received by the earth can change the average global temperature. While this is a complicated process, however, we can have a rough estimation on the temperature decrease.

The spectrum of the sun's light that is absorbed by the earth is reemitted in the longer wavelengths as the blackbody radiation. The total rate of energy that is absorbed by the earth is $\dot{\mathcal{E}}_{in} = S_0 \pi r_e^2 (1 - A_e)$ where S_0 is the energy flux of the sun at the position of the earth, πr_e^2 is the effective area of earth and A_e is the Albedo of earth. On the other hand earth radiates as a blackbody in the longer wavelength with the rate of $\dot{\mathcal{E}}_{out} = 4\pi r_e^2 \sigma T_e^4$. For the equilibrium condition of energy between the inward and outward energy flux, the effective temperature of the earth would be

$$T_e = \left(\frac{S_0(1 - A_e)}{4\sigma} \right)^{1/4}, \quad (8)$$

where for the earth albedo of $A \simeq 0.3$ [10], the effective temperature of earth would be $T_e = 252\text{K}$. This temperature is smaller than the average ground temperature of the earth. Using the greenhouse effect where the atmosphere can absorb part of energy radiating from the surface of earth, the temperature of the ground modified to

$$T_g = \left(\frac{S_0(1 - A_e)}{2\sigma(2 - \epsilon)} \right)^{1/4}, \quad (9)$$

where ϵ is the effective emissivity of ground. We adapt $\epsilon \simeq 0.9$. Then the temperature on ground of earth enhances to $T_g \sim 300\text{K}$. Substituting equation (7) in the derivative of equation (9) results in the temperature change on the ground of earth due to orbital change of earth as

$$\Delta T_g = -0.35 \times 10^{-7} \text{ K} \left(\frac{T_g}{300\text{K}} \right) \left(\frac{m_a}{10^{15}\text{kg}} \right) \left(\frac{v_i}{11\text{km/s}} \right) \cos \theta \cos \phi. \quad (10)$$

We note that the deflection angle of the asteroid scattering by the earth (i.e. θ) depends on the impact parameter (as the minimum distance of asteroid from the earth) as well as the initial velocity of the asteroid. For an asteroid with positive energy of E with respect to the earth, the shape of orbit in polar coordinate is a hyperbola with the equation of $a/r = 1 + \epsilon \cos \gamma$. Here, ϵ is the eccentricity and depends on the physical parameters of asteroid as [11]

$$\epsilon = \left(1 + \frac{2EL^2}{\mu \kappa^2} \right)^{1/2}, \quad (11)$$

where L is the angular momentum of the asteroid with respect to the earth, $\kappa = GM_e m_a$ and μ is the reduced mass of the asteroid and since $\mu \ll M_e$ then $\mu \simeq m_a$. The scattering angle is determined from ϵ (for $r \rightarrow \infty$) as $\cos \gamma = -1/\epsilon$ where γ depends in θ in Fig. (2) by $\gamma = \pi - \theta$, then $\cos \theta = 1/\epsilon$.

Using $L = m_a v_i b$ where b is the impact factor, the eccentricity can be written as $\epsilon = (1 + m_a^2 v_i^2 b^2 / \kappa^2)^{1/2}$. We replace $\cos \theta = 1/\epsilon$ in equation (10) in terms of the eccentricity parameter and in terms of the escape velocity from the earth and the impact parameter, where equation (10) simplifies to

$$\Delta T_g = -0.35 \times 10^{-7} \text{ K} \left(\frac{T_e}{300\text{K}} \right) \left(\frac{m_a}{10^{15}\text{kg}} \right) \left(\frac{v_i}{11\text{km/s}} \right) \left(1 + \left(\frac{v_i}{11\text{km/s}} \right)^2 \left(\frac{b}{6400\text{ km}} \right)^2 \right)^{-1/2} \cos \phi. \quad (12)$$

Here we normalized the initial velocity of the asteroid to the escape velocity from the earth and the impact parameter to the radius of the earth.

The distribution of the asteroids in terms of their size in the main belt of asteroids within the range of $2.82 \text{ a.u.} < a < 2.96 \text{ a.u.}$ is a power law function as $N(D) \sim D^q$ where q is -2.5 for the range of $100 \text{ km} < D < 1000 \text{ km}$ and a shallower slope for $D < 100 \text{ km}$,




Figure 3. Schematic path of an asteroid with the equation of motion provided by $r = r_0 \exp(\beta\varphi)$ with $\beta < 0$.

the exponent is $q \sim -1.8$ down to $D \sim 10$ km [12]. Let us assume an average density of $\bar{\rho} = 2 \text{ gr/cm}^3$ for the asteroids [13]. As a result, we have more low mass asteroids compare to the large ones. Using the mass of asteroid in terms of the size

$$m_a(R) = \left(\frac{L}{10\text{km}}\right)^3 \times 10^{15} \text{ kg}, \quad (13)$$

then we rewrite equation (12) as

$$\Delta T_g = -0.35 \times 10^{-7} K \left(\frac{T_e}{300\text{K}}\right) \left(\frac{L}{10\text{km}}\right)^3 \left(\frac{v_i}{11\text{km/s}}\right) \left(1 + \left(\frac{v_i}{11\text{km/s}}\right)^2 \left(\frac{b}{6400 \text{ km}}\right)^2\right)^{-1/2} \cos \phi. \quad (14)$$

In the rest of this paper, we will discuss how to manage the asteroids to have a gravitational scattered from the earth.

4. Asteroid scattering from the Earth

Most of the asteroids in the plane of solar system are moving in the same direction as the planets move. Our aim is to lower the orbit of the asteroids and encounter them gravitationally with the earth. However, we note that in order to have an energy gain to the earth, asteroids should have a head-on gravitational scattering with the earth.

The closest-large number of asteroids to earth are located in the asteroid belt and in order to have gravitational interaction of asteroids with the earth, we manage to reduce their orbital distance with respect to the sun. The least costing method might be using solar sailing. Also, we can consider a hybrid braking system using solar sailing and install propulsion engines on the asteroids. For solar sailing, the momentum transfer from the photons of the sun to the asteroid plays the role of braking along with the orbital velocity and the result would be a spiral motion of asteroids towards the lower orbits.

Let us denote r as the distance of the sun to the asteroid and φ as the polar coordinate. The general solution for an object with the solar pressure results in the solution of $r = r_0 \exp(\beta\varphi)$ where r_0 is the initial orbital distance and β is a constant where depending on the sign of it orbital distance changes as shown in Figure (3).

The dynamics of an asteroid with a solar sail (in the polar coordinate) is given by the following equations

$$\ddot{r} - r\dot{\varphi}^2 = -\frac{k}{r^2} + \frac{D_r^{(0)}}{mr^2} \cos \psi, \quad (15)$$

$$r\ddot{\varphi} + 2\dot{r}\dot{\varphi} = \frac{D_r^{(0)}}{mr^2} \sin \psi, \quad (16)$$

where (r, φ) are the polar coordinates, $k = GM_{\odot}$ and $-k/r^2$ is the gravitational acceleration due to Sun and $D_r^{(0)}/(mr^2)$ is the extra drag term as the result of the pressure of solar radiation on the sail installed on the asteroid, ψ angle is the angle between the r vector and a normal vector to the sail (e.g. $\psi = \pi/2$ means the sail is aligned perpendicular to the r direction). Assuming P_0 as the radiation pressure at the position of the earth, then $D_r^{(0)} = P_0 r_0^2 A \cos \psi$ where r_0 is the earth-sun distance and A is the area of the sail. $D_r^{(0)}/(mr^2) \times \cos \psi$ is the extra acceleration in the radial direction and $D_r^{(0)}/(mr^2) \times \sin \psi$ is the acceleration along the tangent direction. For $0 < \psi < \pi/2$ the solar pressure is accelerating and results increasing the angular momentum of asteroid and causes the increase of the orbital distance and for $-\pi/2 < \psi < 0$ the effect would be decreasing the angular momentum of the asteroid and hence decreasing the orbital distance to the sun.

Substituting the suggested solution of $r = r_0 \exp(\beta\varphi)$ in the equations (15) and (16), we will have a modified Kepler law as $r^3 \dot{\varphi}^2 = B(\psi)$ where

$$B(\psi) = \frac{k}{1 + \beta^2} - \frac{D_r^{(0)}(\cos \psi - \beta \sin \psi)}{m(1 + \beta^2)}. \quad (17)$$

Also the dynamics of asteroid as a function of time obtain as

$$r(t) = \left(r_0^{3/2} + \frac{3\beta}{2} \sqrt{B(\psi)} t \right)^{2/3}, \quad (18)$$

where depending on the sign of β , the orbit of asteroid increase or decrease. We substitute $r = r_0 \exp(\beta\varphi)$ in the dynamical equations (15 and 16) and use the equation (17) to obtain β in terms of the gravitational force and radiation pressure as

$$\beta = \frac{P_0 r_0^2 A \sin 2\psi}{km - P_0 r_0^2 A \cos^2 \psi} \simeq \frac{P_0 r_0^2 A}{km} \sin 2\psi. \quad (19)$$

For $-\pi/2 < \psi < 0$ the braking effect results in a spiral motion of the asteroid towards the lower orbit and $\psi = -\pi/4$ provides the maximum braking. Figure (3) represents the schematic trajectory of the asteroid towards the lower orbit. We note that our result for the spiral dynamics of the object under solar pressure is different than that is reported in [14].

The time scale for changing the orbital distance of asteroid can be obtained from $t_c = r/\dot{r}$ where substituting the dynamics of spiral motion of asteroid we obtain

$$t_c = 74 \text{yr} \left(\frac{m_a}{10^{10} \text{kg}} \right) \left(\frac{v_a}{10 \text{km/s}} \right) \left(\frac{A}{100 \text{km}^2} \right)^{-1}, \quad (20)$$

where m_a is the mass of the asteroid, v_a is the orbital velocity of the asteroid and A is the area of solar sail installed on the asteroid. In this calculation, we set the solar pressure at the position of sun, $P_0 = S_0/c \simeq 4.5 \mu\text{pa}$. Installing the jet propulsion engines on the asteroids can boost brakes and provides a shorter time scale for changing the orbit of the asteroid. If we lower the orbit of the asteroids to the position of Mars, then a controlled scattering of the asteroid from Mars can guide it toward the earth through an elliptical path. We note that the asteroid and Mars are moving in the same direction and the asteroid can have an energy gain scattering from Mars to an elliptical Hohmann orbit. Now the direction of interception of earth and asteroid is not parallel the asteroid can have a proper configuration of scattering as Fig (2). The consecutive scattering of asteroids from the earth can enhance the mechanical energy of the earth, or in other words the orbital distance of the earth from the sun.

5. Discussion

Concluding this work, we have introduced the hazards of global warming and besides decarbonization policy, we proposed a mechanical method to change the orbit of the earth

by an energy-loss scattering of asteroids from the earth. We proposed the solar-sail as the braking tool to decrease the orbit of the asteroids from the asteroid belt orbit to the Mars orbit. The time scale to lower the orbit is about 70 yrs for a 10^{10} kg mass asteroid. Using the installed propulsion jet engines on the asteroids will decrease this time scale and enable us to do the asteroid maneuvering for a larger number of asteroids. This project can enable us to change the earth's orbit and cool down its temperature by decreasing the energy flux of the sun received by the earth. This project could be feasible for future technology on earth. Also, habitable planets with intelligent life might use it to save their planets while global warming or when their parent star started to inflate to a giant star.

Acknowledgements

This research was supported by Sharif University of Technology's Office of Vice President for Research under Grant No. G950214

1. Jenkins, S.; Millar, R.J.; Leach, N.; Allen, M.R. Framing Climate Goals in Terms of Cumulative CO₂-Forcing-Equivalent Emissions. *Geophysical Research Letters* **2018**, *45*, 2795–2804. doi:10.1002/2017GL076173.
2. Held, I.M.; et al. Structure and Performance of GFDL's CM4.0 Climate Model. *Journal of Advances in Modeling Earth Systems* **2019**, *11*, 3691–3727. doi:10.1029/2019MS001829.
3. Ades, M.; et al. Global Climate. *Bulletin of the American Meteorological Society* **2020**, *101*, S9–S128. doi:10.1175/BAMS-D-20-0104.1.
4. Velden, C.; Olander, T.; Herndon, D.; Kossin, J.P. Reprocessing the Most Intense Historical Tropical Cyclones in the Satellite Era Using the Advanced Dvorak Technique. *Monthly Weather Review* **2017**, *145*, 971–983. doi:10.1175/MWR-D-16-0312.1.
5. Barton, J.P.; Infield, D.G. Energy Storage and Its Use With Intermittent Renewable Energy. *IEEE Transactions on Energy Conversion* **2004**, *19*, 441–448. doi:10.1109/TEC.2003.822305.
6. Jenkins, S.; Millar, R.J.; Leach, N.; Allen, M.R. Framing Climate Goals in Terms of Cumulative CO₂-Forcing-Equivalent Emissions. *Geophysical Research Letters* **2018**, *45*, 2795–2804. doi:10.1002/2017GL076173.
7. Rahvar, S. Frequency-shift in the gravitational microlensing. *Phys Rev D* **2020**, *101*, 024015, [arXiv:astro-ph.GA/1908.01361]. doi:10.1103/PhysRevD.101.024015.
8. Korycansky, D.G.; Laughlin, G.; Adams, F.C. Astronomical Engineering: A Strategy For Modifying Planetary Orbits. *Astrophysics and Space Science* **2001**, *275*, 349–366, [arXiv:astro-ph/0102126].
9. McInnes, C.R. Astronomical Engineering Revisited: Planetary Orbit Modification Using Solar Radiation Pressure. *Astrophysics and Space Science* **2002**, *282*, 765–772. doi:10.1023/A:1021178603836.
10. Susskind, J.; Schmidt, G.A.; Lee, J.N.; Iredell, L. Recent global warming as confirmed by AIRS. *Environmental Research Letters* **2019**, *14*, 044030. doi:10.1088/1748-9326/aafd4e.
11. Thornton, S.T.; Marion, J.B. *Classical dynamics of particles and systems*; Fort Worth: Saunders College Pub., 1995.
12. Tsirvoulis, G.; Morbidelli, A.; Delbo, M.; Tsiganis, K. Reconstructing the size distribution of the primordial Main Belt. *Icarus* **2018**, *304*, 14–23, [arXiv:astro-ph.EP/1706.02091]. doi:10.1016/j.icarus.2017.05.026.
13. Krasinsky, G.A.; Pitjeva, E.V.; Vasilyev, M.V.; Yagudina, E.I. Hidden Mass in the Asteroid Belt. *Icarus* **2002**, *158*, 98–105. doi:10.1006/icar.2002.6837.
14. Bacon, R.H. Logarithmic Spiral: An Ideal Trajectory for the Interplanetary Vehicle with Engines of Low Sustained Thrust. *American Journal of Physics* **1959**, *27*, 164–165. doi:10.1119/1.1934788.