

Novel recurrence relations for volumes and surfaces of n -balls, regular n -simplices, and n -orthoplices in real dimensions

Szymon Łukaszyk*

Łukaszyk Patent Attorneys, ul. Głowackiego 8, 40-052 Katowice, Poland

The study examines n -balls, n -simplices, and n -orthoplices in real dimensions using novel recurrence relations that removed indefiniteness present in known formulas. They show that in the negative, integer dimensions the volumes of n -balls are zero if n is even, positive if $n = -4k - 1$, and negative if $n = -4k - 3$, for natural k . The volumes and surfaces of n -cubes inscribed in n -balls in negative dimensions are complex, wherein for negative, integer dimensions they are associated with integral powers of the imaginary unit. The relations are continuous for $n \in \mathbb{R}$ and show that the constant of π is absent for $0 \leq n < 2$. For $n < -1$ self-dual n -simplices are undefined in the negative, integer dimensions and their volumes and surfaces are imaginary in the negative, fractional ones and divergent with decreasing n . In the negative, integer dimensions n -orthoplices reduce to the empty set, and their real volumes and imaginary surfaces are divergent in negative, fractional ones with decreasing n . Out of three regular, convex polytopes present in all natural dimensions, only n -orthoplices, n -cubes (and n -balls) are defined in the negative, integer dimensions.

Keywords: regular convex polytopes; negative dimensions; fractal dimensions

1. Introduction

The notion of dimension n of a set has various definitions [1,2]. Natural dimensions define a minimum number of independent parameters (coordinates) needed to specify a point within Euclidean space $\mathbb{R}^n$, where $n = -1$ is the dimension of the empty set, the void, having zero volume and undefined surface. Negatively dimensional spaces can be defined by analytic continuations from positive dimensions [3]. A spectrum, topological generalization of the notion of space allows for negative dimensions [2,4,5,6] that refer to densities, rather than to sizes, as the natural ones.

Fractional (or fractal) dimensions extend the notion of dimension to real, including negative [7], numbers. Negative dimensions are considered in probabilistic fractal measures [8]. Fractal dimension and lacunarity [9,10] allow for an investigation of the fractal nature of prime sequences [11]. Fractal dimensions are verified to be consistent with the experimental observations and allow for the analysis of the transport properties, such as permeability, thermal dispersion and conductivities (both thermal and electrical) in multiphase fractal media [12]. The probability models for pore distribution and for permeability of porous media can also be expressed as a function of fractal dimension [13]. Interestingly the dimension of the boundary of the Mandelbrot set equals 2 [14], and the generalized Mandelbrot set in higher-dimensional hypercomplex number spaces, when the power α of the iterated complex variable z tends to infinity, is convergent to the unit $(\alpha-1)$ -sphere [15].

Complex dimensions can also be considered [2]. Furthermore, geometric concepts (such as lengths, volumes, surfaces) can be related to negative, fractional, and complex numbers. Complex geodesic paths emerge in the presence of black hole singularities [16] and when studying entropic dynamics on curved statistical manifolds [17]. Fractional derivatives of complex functions could be able to describe different physical phenomena [18].

In $\mathbb{R}^2$ there is a countably infinite number of regular, convex polygons; in $\mathbb{R}^3$ there are five regular, convex

Platonic solids; in $\mathbb{R}^4$ there are six regular, convex polytopes. For $n > 4$, there are only three: self-dual n -simplex, and n -cube dual to n -orthoplex [19]. Furthermore, $\mathbb{R}^n$ is also equipped with a perfectly regular, convex, n -ball. Properties of these three regular, convex polytopes in natural dimensions are well known [20,21,22]. Fractal dimensions of hyperfractals based on these polytopes in natural dimensions were disclosed in [23].

The study examines n -balls, regular n -simplices, and n -orthoplices in real dimensions using novel recurrence relations that remove indefiniteness present in known formulas.

The paper is structured as follows. Section 2 presents known formulas for volumes and surfaces of n -balls, regular n -simplices and n -orthoplices in natural dimensions. Section 3 defines novel recurrence relations for these geometric objects in real dimensions. Section 4 refers to n -balls circumscribed about and inscribed in n -cubes in real dimensions. Finally, Section 5 summarizes the finding of this paper, whereas their possible applications are discussed in Section 6.

2. Known formulas

The volume of an n -ball (B) is known to be

$$V_n(R)_B = \frac{\pi^{n/2}}{\Gamma(n/2+1)} R^n, \quad (1)$$

where Γ is the Euler's gamma function and R is the n -ball radius. This becomes

$$V_{2k}(R)_B = \frac{\pi^k R^{2k}}{k!}, \quad (2)$$

if n is even ($n = 2k$, $k \in \mathbb{N}_0$) and

$$V_{2k-1}(R)_B = \frac{2^{2k} \pi^{k-1} k!}{(2k)!} R^{2k-1}, \quad (3)$$

if n is odd ($n = 2k - 1$, $k \in \mathbb{N}$). Expressed in terms of n -ball diameter (1) is the rescaling factor between the n -

* szymon@patent.pl

dimensional Lebesgue measure and Hausdorff measure for $n \in \mathbb{R}^+$ [24,2].

Another known [21] recurrence relation expresses the volume of an n -ball in terms of the volume of an $(n-2)$ -ball of the same radius

$$V_n(R)_B = \frac{2\pi R^2}{n} V_{n-2}(R)_B, \quad (4)$$

where $V_0(R)_B = 1$ and $V_1(R)_B = 2R$. It is also known [21] that the $(n-1)$ -dimensional surface of an n -ball can be expressed as

$$S_n(R)_B = \frac{n}{R} V_n(R)_B. \quad (5)$$

Furthermore, it is known [25] that the sequence

$$f_n = \frac{2\pi}{n} f_{n-2} \quad (6)$$

satisfies the same recursion formula as (4) for unit radius.

Volume of a regular n -simplex (S) is known [20,26] to be

$$V_n(A)_S = \frac{\sqrt{n+1}}{n! \sqrt{2^n}} A^n, \quad (7)$$

where A is the edge length. A regular n -simplex has $n+1$ $(n-1)$ -facets [21] so its surface is

$$S_n(A)_S = (n+1) V_{n-1}(A)_S. \quad (8)$$

Volume of n -orthoplex (O) is known [22] to be

$$V_n(A)_O = \frac{\sqrt{2^n}}{n!} A^n. \quad (9)$$

As n -orthoplex has 2^n facets [21] being regular $(n-1)$ -simplices, its surface is

$$S_n(A)_O = 2^n V_{n-1}(A)_S. \quad (10)$$

Formulas (1)-(3) and (7)-(10) are undefined in negative dimensions since factorial is defined only for non-negative integers, while gamma function is undefined for non-positive integers. Relations (4), (5) are undefined if $n = 0$.

3. Novel recurrence relations

A radius recurrence relation

$$f_n \doteq \frac{2}{n} f_{n-2}, \quad (11)$$

for $n \in \mathbb{N}_0$, where $f_0 := 1$ and $f_1 := 2$, allows to express the volumes and, using (5), surfaces of n -balls as

$$V_n(R)_B \doteq f_n \pi^{\lfloor n/2 \rfloor} R^n, \quad (12)$$

$$S_n(R)_B \doteq n f_n \pi^{\lfloor n/2 \rfloor} R^{n-1} = \frac{d}{dR} V_n(R)_B, \quad (13)$$

where " $\lfloor x \rfloor$ " denotes the floor function giving the greatest integer less than or equal to its argument x .

Proof:

If $n = 2k$ for $k \in \mathbb{N}_0$, then by equating (2) with (12)

$$\frac{\pi^k R^{2k}}{k!} = f_{2k} \pi^k R^{2k} \Leftrightarrow f_{2k} = \frac{1}{k!} = \frac{1}{(n/2)!}. \quad (14)$$

Then, with (11), e.g. for $k = 3$

$$f_6 = \frac{2}{6} f_4, f_4 = \frac{2}{4} f_2, f_2 = \frac{2}{2} f_0$$

$$f_6 = \frac{2}{6} \frac{2}{4} \frac{2}{2} 1 = \frac{2^3}{6!!} \Leftrightarrow f_{2k} = \frac{2^k}{(2k)!!} = \frac{2^k}{2^k k!}.$$

For even $n \geq 0$, $n!! = 2^k k!$, which completes the proof.

If $n = 2k-1$, $k \in \mathbb{N}$, then by equating (3) with (12), we have

$$\frac{2^{2k} \pi^{k-1} k!}{(2k)!} R^{2k-1} = f_{2k-1} \pi^{\lfloor (2k-1)/2 \rfloor} R^{2k-1}$$

$$\frac{2^{2k} \pi^{k-1} k!}{(2k)!} = f_{2k-1} \pi^{k-1} \Leftrightarrow \quad (15)$$

$$f_{2k-1} = \frac{2^{2k} k!}{(2k)!} = \frac{2^{2k-1} (k-1)!}{(2k-1)!} = \frac{2^n \left(\frac{n-1}{2} \right)!}{n!}$$

Then, with (11), e.g. for $k = 4$

$$f_7 = \frac{2}{7} f_5, f_5 = \frac{2}{5} f_3, f_3 = \frac{2}{3} f_1, f_1 = \frac{2}{1} f_{-1}$$

$$f_7 = \frac{2}{7} \frac{2}{5} \frac{2}{3} \frac{2}{1} 1 = \frac{2^4}{7!!} \Leftrightarrow f_{2k-1} = \frac{2^k}{(2k-1)!!}.$$

For odd $n \geq 1$, $n!! = (2k-1)!(2^{k-1}(k-1)!)$, which completes the proof.

The sequence (11) allows for presenting n -balls volume and surface recurrence relations (12), (13) as a product of the rational factor f_n or $n f_n$, the irrational factor $\pi^{\lfloor n/2 \rfloor}$, and the metric (radius) factor R^n or R^{n-1} . The relation (11) can be extended into negative dimensions as

$$f_n = \frac{n+2}{2} f_{n+2}, \quad (16)$$

solving for f_{n-2} and assigning new $n \in \mathbb{Z}$ as old $n-2$. It is sufficient to define $f_{-1} = f_0 := 1$ (for the empty set and point dimension) to initiate (11) and (16).

The same assignment of new $n \in \mathbb{Z}$ as old $n-2$ can be made in (4) solved for $V_{n-2}(R)_B$, yielding

$$V_n(R)_B = \frac{n+2}{2\pi R^2} V_{n+2}(R)_B, \quad (17)$$

which enables to avoid the indefiniteness of factorial and gamma function in negative dimensions present in formulas (1)-(3) and removes singularity present in relation (4).




Fig. 1: n -ball radius recurrence relation f_n for $n \in \mathbb{Z}$ (blue), its spline interpolation (red), recurrence relation f_n initialized by spline interpolation for $[-9, -7]$ (cyan), even algebraic form of f_n (yellow), odd algebraic form of f_n (black), and the $\pi^{\lfloor n/2 \rfloor}$ factor (green); for $-7 \leq n \leq 7$.

If $n \leq -3$ and odd

$$f_n = i^{n+1} \frac{2^{n+2} (-n-2)!}{\left(\frac{-n-3}{2}\right)!}. \quad (18)$$

Proof:

Set $n = -2k - 1$, $k \in \mathbb{N}$. Then, with (16), e.g. for $k = 3$

$$f_{-7} = -\frac{5}{2} f_{-5}, \quad f_{-5} = -\frac{3}{2} f_{-3}, \quad f_{-3} = -\frac{1}{2} f_{-1}$$

$$f_{-7} = (-1)^3 \frac{5}{2} \frac{3}{2} \frac{1}{2} 1 = (-1)^3 \frac{5!!}{2^3} \Leftrightarrow$$

$$f_{-2k-1} = \frac{(-1)^k (2k-1)!!}{2^k} = \frac{(-1)^k (2k-1)!}{2^{2k-1} (k-1)!}$$

Also

$$\begin{aligned} (-1)^k &= (-1)^{(-n-1)/2} = \left[(-1)^{\frac{1}{2}} \right]^{-n-1} = i^{-(n+1)} = \\ &= -i^{n+1} = (-1)^{n+1} i^{n+1} = i^{n+1} \end{aligned}$$

since n is odd.

The factorial can be expressed by the gamma function. Thus, for $n = 2k$, $k \in \mathbb{N}$, (14) becomes

$$f_{2k} = \frac{2}{n\Gamma(n/2)}, \quad (19)$$

while for $n = 2k - 1$, $k \in \mathbb{N}$, (15) becomes

$$f_{2k-1} = \frac{2\sqrt{\pi}}{n\Gamma(n/2)}. \quad (20)$$

Radius recurrence relation f_n (16) is shown in Fig. 1 along with its spline interpolation, recurrence relation f_n

initialized by spline interpolation for $[-9, -7]$, even algebraic form of f_n (19), odd algebraic form of f_n (20), and the $\pi^{\lfloor n/2 \rfloor}$ factor for $n \in \mathbb{R}$, and listed in Table 1 for $n \in \mathbb{Z}$. Volumes and surfaces of n -balls calculated with relations (12) and (13) are shown in Fig. 2.




Fig. 2: Graphs of volumes (V) and surface areas (S) of n -balls of unit radius for $n = -25, -24, \dots, 15$.




Fig. 3: Graphs of volumes (V) and surface areas (S) of n -balls of unit diameter for $n = -10, -9, \dots, 8$.

Table 1: Volumes and surfaces of n -balls for $-11 \leq n \leq 9$.

n	f_n	g_n	$V_n(R=1)_B$	$S_n(R=1)_B$	$V_n(D=1)_B$	$S_n(D=1)_B$
-11	-945/32	-60480	-0.031	0.338	-62.909	1383.997
-9	105/16	3360	0.021	-0.193	10.980	-197.634
-7	-15/8	-240	-0.019	0.135	-2.464	34.494
-5	3/4	24	0.024	-0.121	0.774	-7.7404
-3	-1/2	-4	-0.051	0.152	-0.405	2.432
-1	1	2	0.318	-0.318	0.637	-1.273
0	1	1	1	0	1	0
1	2/1	1	2	2	1	2
2	1/1	1/4	3.142	6.283	0.785	3.142
3	4/3	1/6	4.189	12.566	0.524	3.142
4	1/2	1/32	4.935	19.739	0.308	2.467
5	8/15	1/60	5.264	26.319	0.164	1.645
6	1/6	1/384	5.168	31.006	0.081	0.969
7	16/105	1/840	4.725	33.073	0.037	0.517
8	1/24	1/6144	4.059	32.470	0.016	0.254
9	32/945	1/15120	3.299	29.687	0.006	0.116

Furthermore, for $n \in \mathbb{Z}$

$$f_n f_{-n-2} = \operatorname{Re}(i^{n+1}) = \cos\left(\frac{\pi}{2}(n+1)\right), \quad (21)$$

where

$$i^{n+1} = e^{i\pi(n+1)/2} = \cos\left(\frac{\pi}{2}(n+1)\right) + i \sin\left(\frac{\pi}{2}(n+1)\right). \quad (22)$$

Proof:

If $n = 2k$, then

$$f_n f_{-n-2} = f_{2k} f_{-2k-2} = 0 = \operatorname{Re}(i^{n+1}),$$

since $f_n = 0$ for negative, even n .

If $n = 2k - 1$ then, using (15) and (18)

$$\begin{aligned} f_n f_{-n-2} &= f_{2k-1} f_{-2k-1} = \\ &= \frac{2^{2k-1}(k-1)!}{(2k-1)!} (-1)^k \frac{(2k-1)!}{2^{2k-1}(k-1)!} =, \\ &= (-1)^k = (-1)^{(n+1)/2} = i^{n+1} = \operatorname{Re}(i^{n+1}) \end{aligned}$$

since n is odd, which completes the proof.

Furthermore, for $n \in \mathbb{R}$, $k \in \mathbb{Z}$

$$\pi^{\lfloor n/2 \rfloor} \pi^{\lfloor (-n-2)/2 \rfloor} = \begin{cases} \pi^{-1} & n = 2k \\ \pi^{-2} & n \neq 2k \end{cases}. \quad (23)$$

Proof:

If $n = 2k$, then $\pi^k \pi^{-k-1} = \pi^{-1}$. Otherwise, set $n = 2k \pm \varepsilon$, where $0 < \varepsilon \leq 1$, $\varepsilon \in \mathbb{R}$. For $n = 2k + \varepsilon$

$$\pi^{\lfloor k+\varepsilon/2 \rfloor} \pi^{\lfloor -k-\varepsilon/2-1 \rfloor} = \pi^k \pi^{-k-2} = \pi^{-2},$$

while for $n = 2k - \varepsilon$

$$\pi^{\lfloor k-\varepsilon/2 \rfloor} \pi^{\lfloor -k+\varepsilon/2-1 \rfloor} = \pi^{k-1} \pi^{-k-1} = \pi^{-2},$$

which completes the proof.

Also the following holds for n -balls surfaces (13)

$$S_{nB} S_{(2-n)B} = n(2-n) f_n f_{2-n} = 4 \operatorname{Re}(i^{n-1}), \quad (24)$$

for $n \in \mathbb{Z}$, where

$$\begin{aligned} i^{n-1} &= e^{i\pi(n-1)/2} = \\ &= \cos\left(\frac{\pi}{2}(n-1)\right) + i \sin\left(\frac{\pi}{2}(n-1)\right) = -i^{n+1}. \end{aligned} \quad (25)$$

Proof:

If $n = 2k$ then

$$\begin{aligned} S_{(2k)B} S_{(2-2k)B} &= \\ &= 2k f_{2k} \pi^k R^{2k-1} (2-2k) f_{2-2k} \pi^{1-k} R^{1-2k} =, \\ &= 4k(1-k) f_{2k} f_{2-2k} \pi = 0 = 4 \operatorname{Re}(i^{n-1}) \end{aligned}$$

for $k = \{0, 1\}$ and for the remaining k 's, as $f_{-2k} = 0$. Also $\operatorname{Re}(i^{n-1}) = 0$ and $\operatorname{Im}(i^{n-1}) = \pm 1$, as n is even.

If $n = 2k - 1$ then

$$\begin{aligned} S_{(2k-1)B} S_{(3-2k)B} &= \\ &= (2k-1)(3-2k) f_{2k-1} f_{3-2k} \pi^{\lfloor (2k-1)/2 \rfloor} \pi^{\lfloor (3-2k)/2 \rfloor} = \\ &= (2k-1)(3-2k) f_{2k-1} f_{3-2k} \pi^{k-1} \pi^{-k+1} = \\ &= (2k-1)(3-2k) f_{2k-1} f_{3-2k} \end{aligned}$$

For $k = 1$, using (15)

$$S_{(1)B} S_{(1)B} = f_1^2 = 4 = 4 \operatorname{Re}(i^0),$$

wherein for the remaining k 's, we shall use both (15) and (18) (and $f_{-1} := 1$). E.g. for $k = \{0, 2\}$

$$S_{(3)B} S_{(-1)B} = 3(-1) f_3 f_{-1} = -3 \frac{4}{3} 1 = -4 = 4 \operatorname{Re}(i^2),$$

and further, for $k \leq -1$ or $k \geq 3$

$$\begin{aligned} S_{(2k-1)B} S_{(3-2k)B} &= \\ &= (2k-1)(3-2k) \frac{2^{2k-1}(k-1)!}{(2k-1)!} \frac{(-1)^{k-2} (2k-5)!}{2^{2k-5}(k-3)!} = \\ &= (2k-1)(3-2k) (-1)^{k-2} 2^4 \frac{(k-1)!(2k-5)!}{(2k-1)!(k-3)!} = \\ &= 4(-1)^{k-1} = 4i^{n-1} = 4 \operatorname{Re}(i^{n-1}) \end{aligned}$$

since n is odd and, thus $n-1$ is even.

Furthermore, for $n \in \mathbb{R}$, $k \in \mathbb{Z}$

$$\pi^{\lfloor n/2 \rfloor} \pi^{\lfloor (2-n)/2 \rfloor} = \begin{cases} \pi & n = 2k \\ 1 & n \neq 2k \end{cases}. \quad (26)$$

Proof:

If $n = 2k$, then $\pi^k \pi^{1-k} = \pi$. Otherwise, set $n = 2k \pm \varepsilon$, where $0 < \varepsilon \leq 1$, $\varepsilon \in \mathbb{R}$. For $n = 2k + \varepsilon$

$$\pi^{\lfloor k+\varepsilon/2 \rfloor} \pi^{\lfloor 1-k-\varepsilon/2 \rfloor} = \pi^k \pi^{-k} = \pi^0 = 1,$$

while for $n = 2k - \varepsilon$

$$\pi^{\lfloor k-\varepsilon/2 \rfloor} \pi^{\lfloor 1-k+\varepsilon/2 \rfloor} = \pi^{k-1} \pi^{-k+1} = \pi^0 = 1,$$

which completes the proof.

Also the following holds for n -balls volumes (12)

$$\frac{n\pi}{2} V_{nB} V_{(-n)B} = \operatorname{Re}(i^{n-1}), \quad (27)$$

for $n \in \mathbb{Z}$, $n \neq 0$.

Proof:

If $n = 2k$, $k \in \mathbb{N}$, then

$$\begin{aligned} V_{(2k)B} V_{(-2k)B} &= f_{2k} \pi^k R^{2k} f_{-2k} \pi^{-k} R^{-2k} = \\ &= f_{2k} f_{-2k} = 0 = \frac{2}{2k\pi} \operatorname{Re}(i^{2k-1}) \end{aligned}$$

If $n = 2k - 1$, $k \in \mathbb{N}$, then

$$V_{(2k-1)B} V_{(1-2k)B} = f_{2k-1} \pi^{k-1} f_{1-2k} \pi^{-k} = f_{2k-1} f_{1-2k} \pi^{-1}.$$

For $k = 1$, using (15) and $f_{-1} := 1$

$$V_{(1)B} V_{(-1)B} = f_1 f_{-1} \pi^{-1} = \frac{2}{1\pi} = \frac{2}{1\pi} \operatorname{Re}(i^0).$$

For the remaining k 's, we shall use both (15) and (18)

$$\begin{aligned} V_{(2k-1)B} V_{(-2k+1)B} &= f_{2k-1} f_{1-2k} \pi^{-1} = \\ &= \frac{2^{2k-1} (k-1)! (-1)^{k-1} (2k-3)!}{(2k-1)! 2^{2k-3} (k-2)!} \pi^{-1} = , \\ &= (-1)^{k-1} \frac{2}{n\pi} = \frac{2}{n\pi} i^{n-1} = \frac{2}{n\pi} \operatorname{Re}(i^{n-1}) \end{aligned}$$

since n is odd and, thus $n-1$ is even.

Furthermore, for $n \in \mathbb{R}$, $k \in \mathbb{Z}$

$$\pi^{\lfloor n/2 \rfloor} \pi^{\lfloor -n/2 \rfloor} = \begin{cases} 1 & n = 2k \\ \pi^{-1} & n \neq 2k \end{cases}. \quad (28)$$

Proof:

If $n = 2k$, then $\pi^k \pi^{-k} = 1$. Otherwise, set $n = 2k \pm \varepsilon$, where $0 < \varepsilon \leq 1$, $\varepsilon \in \mathbb{R}$. For $n = 2k + \varepsilon$

$$\pi^{\lfloor k+\varepsilon/2 \rfloor} \pi^{\lfloor -k-\varepsilon/2 \rfloor} = \pi^k \pi^{-k-1} = \pi^{-1},$$

while for $n = 2k - \varepsilon$

$$\pi^{\lfloor k-\varepsilon/2 \rfloor} \pi^{\lfloor -k+\varepsilon/2 \rfloor} = \pi^{k-1} \pi^{-k} = \pi^{-1},$$

which completes the proof.

One can also express the volumes and, using (5), surfaces of n -balls in terms of their diameters D as

$$V_n(D)_B \doteq g_n \pi^{\lfloor n/2 \rfloor} D^n, \quad (29)$$

$$S_n(D)_B \doteq 2n g_n \pi^{\lfloor n/2 \rfloor} D^{n-1} = 2 \frac{d}{dD} V_n(D)_B, \quad (30)$$

defining diameter recurrence relation

$$g_n \doteq \frac{1}{2n} g_{n-2} \quad (31)$$

having inverse

$$g_n = 2(n+2) g_{n+2}, \quad (32)$$

for $n \in \mathbb{Z}$, where $g_{-1} := 2$ and $g_0 := 1$. Diameter recurrence relation (31) is related to radius recurrence relation (11) by

$$f_n = 2^n g_n. \quad (33)$$

Proof:

By equating (12) with (29), we have

$$f_n \pi^{\lfloor n/2 \rfloor} R^n = g_n \pi^{\lfloor n/2 \rfloor} 2^n R^n,$$

which completes the proof.

Furthermore (proof follows from (21) and (33))

$$g_n g_{-n-2} = 4 \operatorname{Re}(i^{n+1}). \quad (34)$$

Diameter recurrence relation g_n (32) is shown in Fig. 4 along with its spline interpolation, recurrence relation g_n initialized by spline interpolation for $[-9, -7]$, even algebraic form of g_n , odd algebraic form of g_n , and the $\pi^{\lfloor n/2 \rfloor}$ factor for $n \in \mathbb{R}$, and listed in Table 1 for $n \in \mathbb{Z}$. Volumes and surfaces of n -balls calculated with relations (29) and (30) are shown in Fig. 3.




Fig. 4: n -ball diameter recurrence relation g_n for $n \in \mathbb{Z}$ (blue), its spline interpolation (red), recurrence relation g_n initialized by spline interpolation for $[-9, -7]$ (cyan), even algebraic form of g_n (yellow), odd algebraic form of g_n (black), and the $\pi^{\lfloor n/2 \rfloor}$ factor (green) for $-4 \leq n \leq 4$.

In the case of regular n -simplices, equation (7) can be written as a recurrence relation, with $V_0(A)_S := 1$

$$V_n(A)_S \doteq AV_{n-1}(A)_S \sqrt{\frac{n+1}{2n^3}}. \quad (35)$$

Proof:

By equating (7) with (35), we have

$$\begin{aligned} \frac{\sqrt{n+1}}{n! 2^{n/2}} A^n &= AV_{n-1}(A)_S \frac{\sqrt{n+1}}{2^{1/2} n^{3/2}} \\ V_{n-1}(A)_S &= \frac{2^{(1-n)/2} n^{3/2}}{n!} A^{n-1}, \\ V_n(A)_S &= \frac{(n+1)^{3/2}}{2^{n/2} (n+1)!} A^n = \frac{\sqrt{n+1}}{n! \sqrt{2}^n} A^n \end{aligned}$$

which recovers (7) and completes the proof.

The relation (35) removes the indefiniteness of factorial for $n < 0$ and singularity for $n = -1$ present in (7). Solving (35) for V_{n-1} and assigning new $n \in \mathbb{Z}$ as old $n-1$, yields

$$V_n(A)_S = \frac{V_{n+1}(A)}{A} \sqrt{\frac{2(n+1)^3}{n+2}}, \quad (36)$$

which shows that n -simplices are indefinite only for integer $n < -1$, as shown in Fig. 5. The volume of an empty or

void (-1) -simplex is $V_{-1}(A)_S = 0$, while its surface $S_{-1}(A)_S$ (8) is undefined, as the void itself.




Fig. 5: Graphs of volumes (V) and surface areas (S) of regular n -simplices of unit edge length for $n = -1, \dots, 7$.




Fig. 6: Graphs of volumes (V) and surface areas (S) of n -orthoplices of unit edge length for $n = -1, \dots, 7$.

In the case of n -orthoplices, equation (9) can be written as a recurrence relation

$$V_n(A)_O \doteq AV_{n-1}(A)_O \frac{\sqrt{2}}{n}, \quad (37)$$

with $V_0(A)_O := 1$.

Proof:

By equating (9) with (37), we have

$$\begin{aligned} \frac{\sqrt{2^n}}{n!} A^n &= AV_{n-1}(A)_O \frac{\sqrt{2}}{n} \\ V_{n-1}(A)_O &= \frac{n}{n!} A^{n-1} 2^{(n-1)/2} \\ V_n(A)_O &= \frac{n+1}{(n+1)!} A^n 2^{n/2} = \frac{\sqrt{2^n}}{n!} A^n \end{aligned}$$

which recovers (9) and completes the proof.

The relation (37) removes the indefiniteness of factorial for $n < 0$ present in (9). Solving (37) for V_{n-1} and assigning new $n \in \mathbb{Z}$ as old $n - 1$, yields

$$V_n(A)_O = V_{n+1}(A)_O \frac{n+1}{A\sqrt{2}}, \quad (38)$$

which removes singularity from (37) and is zero for integer $n \leq -1$ showing that for negative, integer dimensions volumes of n -orthoplices are zero, while their surfaces (10) are undefined, as shown in Fig. 6.

4. n -balls circumscribed about and inscribed in n -cubes




Fig. 7: Graphs of volumes (a, pink) and surface areas (b, blue) of n -balls of radius 1, along with volumes and surface areas of n -cubes circumscribed about (yellow) and inscribed in (green) these n -balls. In negative dimensions the latter are complex.

The edge length A_{CC} of n -cube circumscribed (CC) about n -ball corresponds to the diameter D of this n -ball. Thus, the volume of this cube is $V_n(D)_{CC} = D^n$ and the surface is $S_n(D)_{CC} = 2nD^{n-1}$. The edge length A_{CI} of n -cube inscribed (CI) inside the n -ball of diameter D is $A_{CI} = D/\sqrt{n}$, which is singular for $n = 0$ and complex for $n < 0$. Thus, the volume of n -cube inscribed in n -ball is

$$V_n(D)_{CI} = A_{CI}^n = D^n n^{-n/2}, \quad (39)$$

and the surface is

$$\begin{aligned} S_n(D)_{CI} &= 2nA_{CI}^{n-1} = 2D^{n-1} n^{(3-n)/2} = \\ &= \frac{2V_n(D)_{CI} n\sqrt{n}}{D} \end{aligned} \quad (40)$$

The volume (39) is real if n is negative and even and imaginary if n is negative and odd. The surface (40) is real if n is negative and odd and imaginary if n is negative and even. In negative, integer dimensions volumes (39) are associated with a coefficient i^{-n} , while surfaces (40) with a coefficient i^{-n-1} . By convention $0^0 := 1$. Volumes and surfaces of n -cubes given by formulas (39) and (40) are shown in Fig. 7 and listed in Table 2. This peculiar mixture of integer, rational, and irrational coefficients requires further research.

The ratio of volume or surface of n -ball to volume or surface of n -cube circumscribing this n -ball can be expressed using diameter recurrence relations (29), (30) as

$$\frac{V_{nB}}{V_{nCC}} = \frac{S_{nB}}{S_{nCC}} = g_n \pi^{\lfloor n/2 \rfloor}, \quad (41)$$

and similarly, the ratio of volume and surface of n -ball to volume (39) and surface (40) of n -cube inscribed in this n -ball can be expressed as

$$\frac{V_{nB}}{V_{nCI}} = g_n \pi^{\lfloor n/2 \rfloor} n^{n/2}, \quad (42)$$

$$\frac{S_{nB}}{S_{nCI}} = g_n \pi^{\lfloor n/2 \rfloor} n^{(n-1)/2} = \frac{V_{nB}}{V_{nCI} \sqrt{n}}. \quad (43)$$

Table 2: Volumes and surfaces of n -cubes inscribed in n -balls of unit radius and diameter for $-8 \leq n \leq 3$ (rational fraction approximation using Matlab rats function).

n	$V_n(R=1)_{CI}$	$S_n(R=1)_{CI}$	$V_n(D=1)_{CI}$	$S_n(D=1)_{CI}$
-8	16	-362.0387i	4096	-185363.8i
-7	-7.0898i	-16807/128	-907.4927i	-33614
-6	-27/8	49.6022i	-216	6349.077i
-5	1.7469i	625/32	55.9017i	1250
-4	1	-8i	16	-256i
-3	-0.6495i	-27/8	-5.1961i	-54
-2	-1/2	$i\sqrt{2}$	-2	$8i\sqrt{2}$
-1	$i/2$	$1/2$	i	2
0	1	0	1	0
1	2	2	1	2
2	2	$4\sqrt{2}$	$1/2$	$2\sqrt{2}$
3	$8 \cdot 3^{-3/2}$	8	$3^{-3/2}$	2

5. Summary

Novel radius recurrence relation f_n (11) enables to express known recurrence relation (4) for n -ball volume and known relation (5) for n -ball surface as a function of $\pi^{\lfloor n/2 \rfloor}$ showing that the value of π as n -ball volume and surface irrational factor appears only for $n < 0$ and $n \geq 2$ ($\pi^{\lfloor n/2 \rfloor} = 1$ for $0 \leq n < 2$).

Sequence (16), inverse to sequence (11), enables for examination of n -ball volumes and surfaces in the negative dimensions. Since $f_{-2} = 0$, in negative, even dimensions n -balls have zero (void-like) volumes and zero (point-like) surfaces and become divergent with decreasing n . Curiously, the double factorial $n!!$ can be extended to negative, odd integers by inverting its recurrence relation and is not defined for negative even integers.

For positive dimensions $n = 5$ (the largest unit radius n -ball volume) is the largest odd n where $f_n > f_{n-1}$, while $n = 7$ (the largest unit radius n -ball surface) is the smallest odd n where $f_n < f_{n-1}$. Diameter recurrence relation g_n (32) is related with (16) by $f_n = 2^n g_n$.

Algebraic forms (14), (15), (18)-(20) of sequence (16) were presented for even and odd dimensions. Constant (21) of products of pairs of this sequence values for integer n and $-n-2$ reveal symmetry that is the additive inverse of the symmetry $\{n, n-2\}$ or equivalence of an ordinary $(n-2)$ -dimensional space to the n -dimensional superspace [3]. Furthermore sequence (16) reveal symmetry $\{n, 2-n\}$ (24) and $\{n, -n\}$ (27), respectively between n -balls surfaces and volumes in integer dimensions.

Sequence (16) comprises rational numbers, while all $\pi^{\lfloor n/2 \rfloor}$ (for $n < 0$ and $n \geq 2$) are most likely transcendental numbers.

It was shown that the known formula (7) for the volume of a regular n -simplex can be expressed as a recurrence relation (35) to remove indefiniteness of factorial and further expressed as (36) to remove singularity for $n = 0$. Thus, n -simplices are undefined in the negative, integer dimensions if $n < -1$. This is congruent with the fact that every simplicial n -manifold inherits a natural topology from Euclidean space $\mathbb{R}^n$ [27] and by researching Euclidean space $\mathbb{R}^n$ as a simplicial n -manifold topological (metric-independent) and geometrical (metric-dependent) content of the modeled quantities are disentangled [27]. Therefore, the lack of n -simplices in the negative, integer dimensions excludes the notion of negatively dimensional Euclidean space $\mathbb{R}^n$ for $n < -1$. Volumes and surfaces of regular n -simplices are imaginary in negative, fractional dimensions for $n < -1$ (surfaces also for $n < 0$) and are divergent with decreasing n .

It was shown that the known formula (9) for the volume of n -orthoplex can be expressed as a recurrence relation (37) to remove indefiniteness of factorial and further expressed as (38) to remove singularity for $n = 0$. Thus, the volumes of n -orthoplices are zero in the negative, integer dimensions and divergent in the negative, fractional ones with decreasing n . Moreover, the surfaces of n -orthoplices are undefined for integer $n < -1$ (n -orthoplex has facets being regular simplices of the previous dimension (10), and these are undefined for integer $n \leq -1$), imaginary for fractional $n < 0$, and also divergent with decreasing n . Peculiarly, in 1 dimension the volume $V_1(A)_O := A\sqrt{2}$ not A , as in the case of 1-simplex and 1-cube.

Relations (4), (5), (8), (10), (12), (13), (17), (19)-(22), (24), (25), (27), (29), (30), (33)-(43) are continuous on their domains of definitions for $n \in \mathbb{R}$. The starting points for fractional dimensions can be provided e.g. using spline interpolation between two (or three in the case of n -balls) subsequent integer dimensions.

In the negative dimensions n -simplices, n -orthoplices, and n -balls have different properties than their positively dimensional counterparts. n -cube is an exception. A volume $V_n(A)_C = A^n$ and surface $S_n(A)_C = 2nA^{n-1} = 2dV_n(A)_C/dA$ of n -cube are defined for any $n \in \mathbb{R}$ and are real if $A \in \mathbb{R}$. Interestingly, in $\mathbb{R}^3$, the fractal dimension of the Sierpiński 3-simplex is 2, of the Sierpiński 3-orthoplex is 2.585, and only the Sierpiński 3-cube retains its regular dimension [28].

Out of three regular, convex polytopes (and n -balls) present in all non-negative dimensions [19] only n -cubes, n -orthoplices, and n -balls are defined in the negative, integer dimensions with n -cubes being dual to the void.

This should not be surprising. There are no 0-dimensional points in negative dimensions.

6. Discussion

Once upon a time, there was a (-1) -dimensional void of volume zero and undefined surface. Then a 0-dimensional point of unit volume and null surface somehow appeared in this void. This first point is now called primordial Big Bang singularity. The existence of the first point implied a countably infinite number of other labeled points forming various relations among each other. And thus, the void expanded into real and imaginary dimensionalities.

Presented recurrence relations remove indefiniteness and singularities present in known formulas revealing the properties of the relevant geometric objects in negative and real dimensions.

The results of this study could perhaps be applied in linguistic statistics, where the dimension in the distribution for frequency dictionaries is chosen to be negative [4], and in fog computing, where n -simplex is related to a full mesh pattern, n -orthoplex is linked to a quasi-full mesh structure and n -cube is referred to as a certain type of partial mesh layout [29].

Another possible application of the results of this study could be molecular physics and crystallography. There are countably infinitely many spherical harmonics, but nature uses only the first four as subshells of s, p, d, and f electron shells that can hold 2, 6, 10, and 14 electrons, respectively. Further subshells are not populated in the ground states of all the observed elements. The first element that would require a g subshell (18 electrons) would have an atomic number of 121, while the heaviest element synthesized is Oganesson, with an atomic number of 118 and a half-life of about 1/1000 of a second. Perhaps this is linked with properties of the unit radius n -balls in negative dimensions, as illustrated in Fig. 2. The “flattening” occurring between dimensions -14 and -2 is intriguing. Dimensions -2 , -6 , -10 , and -14 are bounded from both sides, with -14 , which would represent the f subshell, already at the onset of divergence. In nature, the f subshell occurs essentially only in lanthanides and actinides. A simple and approximate formula for a spherical nuclear radius that generates very precise results in quantum and nuclear techniques is $R = r_0 A^{1/3}$, where A is the atomic number and $r_0 = 1.25 \pm 0.2$ fm.

Acknowledgments

I truly thank my wife and Mirek for their support. I thank Wawrzyniec for hinting derivation of the preliminary forms of formulas (18), (15).

Data Availability Statement

https://github.com/szluk/balls_simplices_orthoplices

References

1. D. Schleicher, “Hausdorff Dimension, Its Properties, and Its Surprises”. The American Mathematical Monthly Vol. 114, No. 6 (Jun-Jul 2007), <https://www.jstor.org/stable/27642249>.
2. Y. I. Manin, “The notion of dimension in geometry and algebra”. Bulletin of the American Mathematical Society. 43 (2): 139–161 (8 Feb 2006).
3. G. Parisi, et al., “Random Magnetic Fields, Supersymmetry, and Negative Dimensions”. Phys. Rev. Lett. 43, 744 (10 Sep 1979), <https://doi.org/10.1103/PhysRevLett.43.744>.
4. V. P. Maslov, “Negative dimension in general and asymptotic topology”. arXiv:math/0612543 (19 Dec 2006).
5. V. P. Maslov, “General notion of a topological space of negative dimension and quantization of its density”. Mathematical Notes 81 (1–2): 140–144 (6 Dec 2006), <https://doi.org/10.1134/S0001434607010166>.
6. Office chair philosophy, Generalised definition for negative dimensional geometry. Available online: <https://tgrad.blogspot.com/2017/08/reframing-geometry-to-include-negative.html> (accessed on 20 May 2022).
7. B. B. Mandelbrot, “Negative Fractal Dimensions And Multifractals”. Physica A 163 306-315 (1990), [https://doi.org/10.1016/0378-4371\(90\)90339-T](https://doi.org/10.1016/0378-4371(90)90339-T).
8. A. B. Chhabra et al, “Negative dimensions: Theory, computation, and experiment”, Phys. Rev. A 43, 114(R) (1991), <https://doi.org/10.1103/PhysRevA.43.1114>.
9. C. Allain, et al., “Characterizing the lacunarity of random and deterministic fractal sets”. Phys. Rev. A, 44, 3552–3558 (1991), <https://doi.org/10.1103/physreva.44.3552>.
10. J. E. Hutchinson, “Fractals and self similarity”. Indiana Univ. Math. J., 30, 713–747 (1981), <http://dx.doi.org/10.1512/iumj.1981.30.30055>.
11. E. Guariglia, “Primality, Fractality, and Image Analysis”. Entropy, 21(3), 304 (2019); <https://doi.org/10.3390/e21030304>.
12. B. Yu, “Fractal Dimensions for Multiphase Fractal Media”. Fractals Vol. 14, No. 02, pp. 111-118 (2006) <https://doi.org/10.1142/S0218348X06003155>.
13. B. Yu, et al., “Permeability of fractal porous media by Monte Carlo simulations”. International Journal of Heat and Mass Transfer, Vol 48, Issue 13, (Jun 2005), <https://doi.org/10.1016/j.ijheatmasstransfer.2005.02.008>.
14. M. Shishikura, “The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets”. Annals of Mathematics. Second Series. 147 (2): (1998), <https://doi.org/10.2307/121009>.
15. A. Katunin, et al., “On a Visualization of the Convergence of the Boundary of Generalized Mandelbrot Set to $(n-1)$ -Sphere”. Journal of Applied Mathematics and Computational Mechanics (2015), <https://doi.org/10.17512/jamcm.2015.1.06>.
16. L. Fidkowski et al., “The black hole singularity in AdS/CFT”, Journal of High Energy Physics 02, 014 (2004), <https://doi.org/10.1088/1126-6708/2004/02/014>.
17. S. Gassner et al., “Information geometric complexity of entropic motion on curved statistical manifolds under different metrizations of probability spaces”, Int. J. Geometric Methods in Modern Physics 16,

- 1950082, (2019),
<https://doi.org/10.1142/s0219887819500828>.
18. E. Guariglia, et al., "Fractional-Wavelet Analysis of Positive definite Distributions and Wavelets on $D'(\mathbb{C})$ ". Engineering Mathematics II. Springer Proceedings in Mathematics & Statistics, vol 179. Springer, Cham., (2016)
https://doi.org/10.1007/978-3-319-42105-6_16.
 19. J. Baez, "Platonic Solids in All Dimensions", Available online:
<https://math.ucr.edu/home/baez/platonic.html> (accessed on 20 May 2022).
 20. D. M. Y. Sommerville, "An Introduction to The Geometry of N Dimensions". Methuen & Co., London, (1929) ISBN-10: 1781830312.
 21. H. S. M. Coxeter, "Regular Polytopes (3rd ed.)" (1973). ISBN-13: 978-0486614809.
 22. M. Henk, et al., "Basic Properties of Convex Polytopes". Handbook of Discrete and Computational Geometry (1997), ISBN9781315119601.
 23. A. Katunin, "Fractals based on regular convex polytopes". Scientific Research of the Institute of Mathematics and Computer Science, Vol 11, Issue 2, p. 53-62 (2012).
 24. The Chinese University of Hong Kong, "Chapter 3, Lebesgue and Hausdorff Measures", Math 5011.
 25. J. Gipple, "The Volume of n -balls". Rose-Hulman Undergraduate Mathematics Journal, Vol. 15, Issue 1 Art. 14. (2014).
 26. R. H. Buchholz, "Perfect Pyramids". Bulletin of the Australian Mathematical Society, 45(3), (1991),
<https://doi.org/10.1017/S0004972700030252>.
 27. M. Desbrun, et al., "Discrete Differential Forms for Computational Modeling". Discrete Differential Geometry. Oberwolfach Seminars, vol 38. Birkhäuser Basel (2008),
https://doi.org/10.1007/978-3-7643-8621-4_16.
 28. A. Kunnen, et al., "Regular Sierpinski polyhedral". Pi Mu Epsilon J., 10, 607-619 (1998).
 29. P. J. Roig, et al., "Applying Multidimensional Geometry to Basic Data Centre Designs". Journal of Electrical and Computer Engineering Research, Vol. 1, No. 1, (2021),
<https://doi.org/10.53375/ijecer.2021.17>.