

Novel recurrence relations for volumes and surfaces of n -balls, regular n -simplices, and n -orthoplices in real dimensions

Szymon Łukaszyk*

Łukaszyk Patent Attorneys, ul. Głowackiego 8, 40-052 Katowice, Poland

The aim of this study was to examine n -balls, n -simplices and n -orthoplices in real dimensions using novel recurrence relations that removed indefiniteness present in known formulas. They show that in negative, integer dimensions volumes of n -balls are zero if n is even, positive if $n = -4k - 1$, and negative if $n = -4k - 3$, for natural k . Volumes and surfaces of n -cubes inscribed in n -balls in negative dimensions are complex, wherein for negative, integer dimensions they are associated with integral powers of the imaginary unit. The relations are continuous for $n \in \mathbb{R}$ and show that the constant of π is absent for $0 \leq n < 2$. For $n < -1$ self-dual n -simplices are undefined in negative, integer dimensions and their volumes and surfaces are imaginary in negative, fractional ones, and divergent with decreasing n . In negative, integer dimensions n -orthoplices reduce to the empty set, and their real volumes and imaginary surfaces are divergent in negative, fractional ones with decreasing n . Out of three regular, convex polytopes present in all natural dimensions, only n -orthoplices, n -cubes (and n -balls) are defined in negative, integer dimensions.

Keywords: regular convex polytopes; negative dimensional spectra; fractal dimensions

1. Introduction

The notion of dimension n of a set has various definitions [1]. Natural dimensions define a minimum number of independent parameters (coordinates) needed to specify a point within Euclidean space \mathbb{R}^n , where $n = -1$ is the dimension of the empty set, the void, having zero volume and undefined surface. Negatively dimensional spaces can be defined by analytic continuations from positive dimensions [2]. A spectrum, topological generalization of the notion of space allows for negative dimensions [3, 4, 5, 6] that refer to densities, rather than to sizes, as the natural ones. Fractional (or fractal) dimensions extend the notion of dimension to real, including negative [7], numbers. Interestingly the dimension of the boundary of the Mandelbrot set equals 2 [8] and the generalized Mandelbrot set in higher-dimensional hypercomplex number spaces, when the power α of the iterated complex variable z tends to infinity, is convergent to the unit $(\alpha-1)$ -sphere [9]. By no means this limits the meaning of dimension. Complex dimensions [3] are also considered, for example.

In \mathbb{R}^2 there are countably infinite number of regular, convex polygons, in \mathbb{R}^3 there are five regular, convex Platonic solids, in \mathbb{R}^4 there are six regular, convex polytopes. For $n > 4$, there are only three: self-dual n -simplex, and n -cube dual to n -orthoplex [10]. Furthermore \mathbb{R}^n is also equipped with perfectly regular, and obviously also convex, n -ball.

Properties of these three regular, convex polytopes in natural dimensions are well known [11, 12, 13], while fractal dimensions of hyperfractals based on these polytopes in natural dimensions were disclosed in [14].

The aim of this study was to examine n -balls, n -simplices and n -orthoplices in real dimensions using novel recurrence relations that remove indefiniteness present in known formulas.

The paper is structured as follows. Section 2 presents known formulas for volumes and surfaces of n -balls, n -simplices and n -orthoplices in natural dimensions. Section 3 defines novel recurrence relations for these geometric objects in real dimensions. Section 4 refers to n -balls

circumscribed about and inscribed in n -cubes in real dimensions. Section 5 summarizes the finding of this paper, whereas their possible applications are discussed in Section 6.

2. Known formulas

Volume of an n -ball (B) is known to be

$$V_n(R)_B = \frac{\pi^{n/2}}{\Gamma(n/2 + 1)} R^n, \quad (1)$$

where Γ is the Euler's gamma function and R is the n -ball radius. This becomes

$$V_{2k}(R)_B = \frac{\pi^k R^{2k}}{k!}, \quad (2)$$

if n is even and

$$V_{2k+1}(R)_B = \frac{2(k!)(4\pi)^k}{(2k+1)!} R^{2k+1}, \quad (3)$$

if n is odd. Expressed in terms of n -ball diameter (1) is the rescaling factor between n -dimensional Lebesgue measure and Hausdorff measure for $n \in \mathbb{R}^+$ [15, 3].

Another known [12] recurrence relation expresses the volume of an n -ball in terms of the volume of an $(n-2)$ -ball of the same radius

$$V_n(R)_B = \frac{2\pi R^2}{n} V_{n-2}(R)_B, \quad (4)$$

where $V_0(R)_B = 1$ and $V_1(R)_B = 2R$. It is also known [12] that $n-1$ dimensional surface of an n -ball can be expressed as

$$S_n(R)_B = \frac{n}{R} V_n(R)_B. \quad (5)$$

Furthermore, it is known [16] that the sequence

* szymon@patent.pl

$$f_n = \frac{2\pi}{n} f_{n-2} \quad (6)$$

satisfies the same recursion formula as (4) for unit radius.

Volume of a regular n -simplex (S) is known [11, 17] to be

$$V_n(A)_S = \frac{\sqrt{n+1}}{n! \sqrt{2^n}} A^n, \quad (7)$$

where A is the edge length. A regular n -simplex has $n+1$ ($n-1$)-facets [12] so its surface is

$$S_n(A)_S = (n+1)V_{n-1}(A)_S. \quad (8)$$

Volume of n -orthoplex (O) is known [13] to be

$$V_n(A)_O = \frac{\sqrt{2^n}}{n!} A^n. \quad (9)$$

As n -orthoplex has 2^n facets [12] being ($n-1$)-simplices, its surface is

$$S_n(A)_O = 2^n V_{n-1}(A)_S. \quad (10)$$

Relations (1)-(3) and (7)-(10) are undefined in negative dimensions, since factorial is defined only for non-negative integers, while gamma function is undefined for non-positive integers. Relations (4), (5) are undefined if $n = 0$.

3. Novel recurrence relations

A radius recurrence relation

$$f_n \doteq \frac{2}{n} f_{n-2}, \quad (11)$$

for $n \in \mathbb{N}_0$, where $f_0 := 1$ and $f_1 := 2$, allows to express the volumes (4) and surfaces (5) of n -balls as

$$V_n(R)_B \doteq f_n \pi^{\lfloor n/2 \rfloor} R^n, \quad (12)$$

$$S_n(R)_B \doteq n f_n \pi^{\lfloor n/2 \rfloor} R^{n-1}. \quad (13)$$




Fig. 1: Radius n -ball recurrence relation (blue) with the $\pi^{\lfloor n/2 \rfloor}$ factor (green) for $n = -7, -6, \dots, 7$.

The sequence (11) allows to present n -balls volume and surface recurrence relations (12), (13) as a product of a rational factor f_n or $n f_n$, an irrational factor $\pi^{\lfloor n/2 \rfloor}$ (for $n \neq 0$ and $n \neq 1$), and a metric (radius) factor R^n or R^{n-1} . The relation (11) can be then extended into negative dimensions as

$$f_n \doteq \frac{n+2}{2} f_{n+2}, \quad (14)$$

solving for f_{n-2} and assigning new $n \in \mathbb{Z}$ as old $n-2$.

The same assignment of new $n \in \mathbb{Z}$ as old $n-2$ can be made in (4) solved for $V_{n-2}(R)_B$ yielding

$$V_n(R)_B = \frac{n+2}{2\pi R^2} V_{n+2}(R)_B. \quad (15)$$

This enables to avoid the indefiniteness of factorial and gamma function in negative dimensions present in relations (1)-(3) and removes singularity present in (4).

Radius recurrence relations (11), (14) are shown in Fig. 1 along with the $\pi^{\lfloor n/2 \rfloor}$ factor and listed in Table 1. Volumes and surfaces of n -balls calculated with relations (12) and (13) are shown in Fig. 2 and Fig. 3.

For positive dimensions $n = 5$ (the largest unit radius n -ball volume) is the last odd n where $f_n > f_{n-1}$, while $n = 7$ (the largest unit radius n -ball surface) is the first odd n where $f_n < f_{n-1}$. It is sufficient to define $f_{-1} = f_0 := 1$ (for the empty set and point dimension) to initiate (11) and (14).




Fig. 2: Graphs of volumes (V) and surface areas (S) of n -balls of unit radius for $n = -25, -6, \dots, 15$.




Fig. 3: Graphs of volumes (V) and surface areas (S) of n -balls of unit diameter for $n = -10, -9, \dots, 8$.

Furthermore, if $n < -1$ and odd

$$f_n = -(-1)^{\lfloor n/2 \rfloor} \frac{\prod_{k=0}^{\lfloor n/2 \rfloor - 1} (2k+1)}{2^{\lfloor n/2 \rfloor}}, \quad (16)$$

if $n > 0$ and odd

$$f_n = \frac{2^{(n+1)/2}}{\prod_{k=0}^{\lfloor n/2 \rfloor} (2k+1)}, \quad (17)$$

and if $n > 0$ and even

$$f_n = \left(\prod_{k=1}^{n/2} k \right)^{-1}. \quad (18)$$

Also, if n is odd

$$-(-1)^{\lfloor n/2 \rfloor} f_{-(n+2)} f_n = 1. \quad (19)$$

One can also express the volumes and, using (5), surfaces of n -balls in terms of their diameters D as

$$V_n(D)_B = g_n \pi^{\lfloor n/2 \rfloor} D^n, \quad (20)$$

$$S_n(D)_B = 2n g_n \pi^{\lfloor n/2 \rfloor} D^{n-1}, \quad (21)$$

defining diameter recurrence relation

$$g_n \doteq \frac{1}{2n} g_{n-2} \quad (22)$$

having inverse

$$g_n \doteq 2(n+2) g_{n+2}, \quad (23)$$

for $n \in \mathbb{Z}$, where $g_{-1} := 2$ and $g_0 := 1$.




Fig. 4: Diameter n -ball recurrence relation (blue) with the $\pi^{\lfloor n/2 \rfloor}$ factor (green) for $n = -4, -3, \dots, 4$.

Diameter recurrence relation (22), (23) is shown in Fig. 4 along with the $\pi^{\lfloor n/2 \rfloor}$ factor and listed in Table 1. Furthermore, if $n \geq 0$

$$g_n = \frac{\pi^{n/2}}{2^n \Gamma(n/2 + 1) \pi^{\lfloor n/2 \rfloor}}, \quad (24)$$

is the reciprocal of OEIS A087299 sequence, and if $n < 0$ and odd

$$g_n = -(-1)^{\lfloor n/2 \rfloor} \frac{2(2\lfloor -n/2 \rfloor)!}{(\lfloor -n/2 \rfloor)!}, \quad (25)$$

corresponds to OEIS A151817 (for $k = \lfloor -n/2 \rfloor$ and excluding the sign factor) and OEIS A052718 sequence (for $n < -2$ and excluding the sign factor).

Also, if n is odd

$$-(-1)^{\lfloor n/2 \rfloor} g_{-(n+2)} g_n = 4. \quad (26)$$

Since $f_{-2} = 0$ (14) and $g_{-2} = 0$ (23), in negative, even dimensions volumes and surfaces of n -balls vanish.

Table 1: Volumes and surfaces of n -balls for $-11 \leq n \leq 9$.

n	f_n	g_n	$V_n(R=1)_B$	$S_n(R=1)_B$	$V_n(D=1)_B$	$S_n(D=1)_B$
-11	-945/32	-60480	-0.031	0.338	-62.909	1383.997
-9	105/16	3360	0.021	-0.193	10.980	-197.634
-7	-15/8	-240	-0.019	0.135	-2.464	34.494
-5	3/4	24	0.024	-0.121	0.774	-7.7404
-3	-1/2	-4	-0.051	0.152	-0.405	2.432
-1	1	2	0.318	-0.318	0.637	-1.273
0	1	1	1	0	1	0
1	2/1	1	2	2	1	2
2	1/1	1/4	3.142	6.283	0.785	3.142
3	4/3	1/6	4.189	12.566	0.524	3.142
4	1/2	1/32	4.935	19.739	0.308	2.467
5	8/15	1/60	5.264	26.319	0.164	1.645
6	1/6	1/384	5.168	31.006	0.081	0.969
7	16/105	1/840	4.725	33.073	0.037	0.517
8	1/24	1/6144	4.059	32.470	0.016	0.254
9	32/945	1/15120	3.299	29.687	0.006	0.116




Fig. 5: Graphs of volumes (V) and surface areas (S) of regular n -simplices of unit edge length for $n = -1, \dots, 7$.

In the case of regular n -simplices, equation (7) can be written as a recurrence relation

$$V_n(A)_S = A V_{n-1}(A)_S \sqrt{\frac{n+1}{2n^3}}, \quad (27)$$

with $V_0(A)_S := 1$. This removes indefiniteness of factorial for $n = -1$ present in (7). Solving (27) for V_{n-1} , and assigning new $n := n - 1 \in \mathbb{Z}$ yields

$$V_n(A)_S = \frac{V_{n+1}(A)}{A} \sqrt{\frac{2(n+1)^3}{n+2}}, \quad (28)$$

which shows that n -simplices are indefinite only for integer $n < -1$, as shown in Fig. 5. The volume of an empty or void (-1) -simplex is $V_{-1}(A)_S = 0$, while its surface $S_{-1}(A)_S$ (8) is undefined, as the void itself.

In the case of n -orthoplices, equation (9) can be written as a recurrence relation

$$V_n(A)_O = AV_{n-1}(A)_O \frac{\sqrt{2}}{n}, \quad (29)$$

with $V_0(A)_O := 1$ and reversed solving for $n-1$ as

$$V_n(A)_O = V_{n+1}(A)_O \frac{n+1}{A\sqrt{2}}, \quad (30)$$

which removes singularity from (29) and is zero for integer $n \leq -1$ showing that for negative, integer dimensions volumes of n -orthoplices are zero, while their surfaces (10) are undefined, as shown in Fig. 6. It is not surprising: n -orthoplex has facets being simplexes of the previous dimension, and these are undefined for integer $n \leq -1$. Peculiarly, $V_1(A)_O := A\sqrt{2}$ not A , as in the case of 1-simplex and 1-cube.




Fig. 6: Graphs of volumes (V) and surface areas (S) of n -orthoplices of unit edge length for $n = -1, \dots, 7$.

4. n -balls circumscribed about and inscribed in n -cubes

The edge length A_C of n -cube circumscribed about n -ball corresponds to the diameter D of this n -ball. The ratio of volume (D^n) or surface ($2nD^{n-1}$) of n -ball to volume or surface of n -cube circumscribing this n -ball can be expressed using diameter recurrence relation (20), (21) as

$$\frac{V_{nB}}{V_{nC_C}} = \frac{S_{nB}}{S_{nC_C}} = g_n \pi^{\lfloor n/2 \rfloor}. \quad (31)$$

The edge length A_I of n -cube inscribed inside the n -ball of diameter D is

$$A_I = \frac{D}{\sqrt{n}}, \quad (32)$$

which is singular for $n = 0$ and complex for $n < 0$. The volume of n -cube inscribed in n -ball is

$$V_n(D)_{C_I} = A_I^n = D^n n^{-n/2}, \quad (33)$$

and the surface is

$$S_n(D)_{C_I} = 2nA_I^{n-1} = 2D^{n-1} n^{(3-n)/2}. \quad (34)$$

The volume (33) is real if n is negative and even, and imaginary if n is negative and odd. The opposite holds true for the surface (34). In negative, integer dimensions volumes (33) are associated with a coefficient i^n , while surfaces (34) with a coefficient i^{n+1} . By convention $0^0 := 1$.

Volumes and surfaces of n -cubes given by formulas (33) and (34) are shown in Fig. 7 and listed in Table 2.




Fig. 7: Graphs of volumes (a, pink) and surface areas (b, blue) of n -balls of radius 1, along with volumes and surface areas of n -cubes circumscribed about (yellow) and inscribed in (green) these n -balls. In negative dimensions the latter are complex.

Table 2: Volumes and surfaces of n -cubes inscribed in n -balls of unit radius and diameter for $-8 \leq n \leq 2$.

n	$V_n(R=1)_{C_I}$	$S_n(R=1)_{C_I}$	$V_n(D=1)_{C_I}$	$S_n(D=1)_{C_I}$
-8	16	362.039i	4096	185363.8i
-7	7.090i	-16807/128	907.493i	-33614
-6	-216/64	-49.602i	-216	-6349.077i
-5	-1.747i	19.531	-55.902i	1250
-4	1	8i	16	256i

n	$V_n(R=1)_{Ci}$	$S_n(R=1)_{Ci}$	$V_n(D=1)_{Ci}$	$S_n(D=1)_{Ci}$
-3	0.650i	-216/64	5.196i	-54
-2	-1/2	-i√2	-2	-8i√2
-1	-i/2	1/2	-i	2/1
0	1	0	1	0
1	2	2	1	2
2	2	4√2	1/2	2√2

5. Summary

Novel radius (11) and diameter (22) recurrence relations enable to express known recurrence relation (4) for n -ball volume and known relation (5) for n -ball surface as a function of $\pi^{[n/2]}$ showing that the value of π as n -ball volume and surface irrational factor appears only for $n < 0$ and $n \geq 2$ ($\pi^{[n/2]} = 1$ for $0 \leq n < 2$). Inverse sequences (14) and (23) enable to examine n -ball volumes and surfaces in negative dimensions showing that in negative, even dimensions n -balls have zero (void-like) volumes and zero (point-like) surfaces and become divergent with decreasing n . Novel forms (16)-(18) and (24), (25) of sequences (11), (14), (23), and (22) were presented for even and odd dimensions, as well as particular constants (19), (26) of their products in odd dimensions. Sequences (11), (14), (23), and (22) are rational numbers, while all $\pi^{[n/2]}$ (for $n < 0$ and $n \geq 2$) are most likely transcendental numbers. Doubled maxima for unit diameter n -balls (volume for $n = 0, 1$ and surface for $n = 2, 3$) are also interesting.

It was shown that known formula (7) for the volume of a regular n -simplex can be expressed as a recurrence relation (27) to remove indefiniteness of factorial and further expressed as (28) to remove singularity for $n = 0$. Thus, n -simplices are undefined in negative, integer dimensions if $n < -1$. This is congruent with the fact that every simplicial n -manifold inherits a natural topology from Euclidean space \mathbb{R}^n [18] and by researching Euclidean space \mathbb{R}^n as a simplicial n -manifold topological (metric-independent) and geometrical (metric-dependent) content of the modeled quantities are disentangled [18]. Therefore, lack of n -simplices in negative, integer dimensions excludes the notion of negatively dimensional Euclidean space \mathbb{R}^n for $n < -1$. Volumes and surfaces and surfaces of regular n -simplices are imaginary in negative, fractional dimensions for $n < -1$ (surfaces also for $n < 0$) and are divergent with decreasing n .

It was shown that known formula (9) for the volume of n -orthoplex can be expressed as a recurrence relation (29) to remove indefiniteness of factorial and further expressed as (30) to remove singularity for $n = 0$. Thus, volumes of n -orthoplices are zero in negative, integer dimensions, and divergent in negative, fractional ones with decreasing n . Surfaces of n -orthoplices are undefined for integer $n < -1$, imaginary for fractional $n < 0$, and also divergent with decreasing n .

Relations (4), (11)-(15), (20)-(23), (27)-(30) are continuous for $n \in \mathbb{R}$. The starting points for fractional dimensions can be provided e.g. using spline interpolation between two (or three in the case of n -balls) subsequent integer dimensions.

In negative dimensions n -simplices, n -orthoplices, and n -balls have different properties than their positively dimensional counterparts. n -cube is an exception. A volume $V_n(A)_C = A^n$ and surface $S_n(A)_C = 2nA^{n-1}$ of n -cube are defined for any $n \in \mathbb{R}$ and are real if $A \in \mathbb{R}$. Interestingly

in \mathbb{R}^3 , fractal dimension of the Sierpiński 3-simplex is 2, of the Sierpiński 3-orthoplex is 2.585, and only the Sierpiński 3-cube retains its regular dimension [19].

Out of three regular, convex polytopes (and n -balls) present in all non-negative dimensions [10] only n -cubes, n -orthoplices, and n -balls are defined in negative, integer dimensions with n -cubes being dual to the void.

6. Discussion

Once upon a time there was a (-1)-dimensional void of volume zero and undefined surface. A 0-dimensional point of unit volume and zero surface somehow appeared in this void. This first point is now called primordial Big Bang singularity. An existence of the first point implied countably infinite number of other labelled points forming various relations among each other. And thus the void expanded into real and imaginary dimensionalities.

Presented recurrence relations remove indefiniteness and singularities present in known formulas revealing the properties of the relevant geometric objects in negative and real dimensions.

The results of this study could perhaps be applied in linguistic statistics, where the dimension in the distribution for frequency dictionaries is chosen to be negative [4] and fog computing, where n -simplex is related to a full mesh pattern, n -orthoplex is linked to a quasi-full mesh structure and n -hypercube is referred to as a certain type of partial mesh layout [20].

Another possible application of the results of this study could be molecular physics and crystallography. There are countably infinitely many spherical harmonics but nature uses only the first four as subshells of s, p, d, and f electron shells that can hold 2, 6, 10, and 14 electrons respectively. Further subshells are not populated in ground states of all the observed elements. The first element that would require a g subshell (18 electrons) would have an atomic number of 121, while the heaviest element synthesized is Oganesson, with an atomic number of 118 and a half-life of about 1/1000 of a second. Perhaps this is linked with properties of the unit radius n -balls in negative dimensions as illustrated in Fig. 2. The “flattening” occurring between dimensions -14 and -2 is intriguing. Dimensions -2, -6, -10, and -14 are bounded from both sides, with -14, that would represent the f subshell, already at the onset of divergence. In nature, the f subshell occurs essentially only in lanthanides and actinides. A simple and approximate formula for a spherical nuclear radius that generates very precise results in quantum and nuclear techniques is $R = r_0 A^{1/3}$, where A is the atomic number and $r_0 = 1.25 \pm 0.2$ fm.

Acknowledgments

I thank Wawrzyniec for hinting the derivation of the formulas (16)-(18).

References

- [1] Dierk Schleicher, “Hausdorff Dimension, Its Properties, and Its Surprises”. The American Mathematical Monthly Vol. 114, No. 6 (Jun. - Jul., 2007).
- [2] G. Parisi, N. Sourlas, “Random Magnetic Fields, Supersymmetry, and Negative Dimensions”. Sep 1979. <https://doi.org/10.1103/PhysRevLett.43.744>

- [3] Yuri I. Manin, "The notion of dimension in geometry and algebra". Bulletin of the American Mathematical Society. 43 (2): 139–161. Retrieved 17 Dec 2021.
- [4] V. P. Maslov, "Negative dimension in general and asymptotic topology". arXiv:math/0612543, 19 Dec 2006.
- [5] V. P. Maslov, "General notion of a topological space of negative dimension and quantization of its density". Mathematical Notes 81 (1–2): 140–144. doi:10.1134/S0001434607010166, 6 Dec 2006.
- [6] <https://tglas.blogspot.com/2017/08/reframing-geometry-to-include-negative.html>.
- [7] Benoit B. Mandelbrot, "Negative Fractal Dimensions And Multifractals". Physica A 163 (1990) 306-315.
- [8] Mitsuhiro Shishikura, "The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets". Annals of Mathematics. Second Series. 147 (2): 225–267. arXiv:math.DS/9201282. doi:10.2307/121009. JSTOR 121009. MR 1626737. S2CID 14847943, 1998.
- [9] Andrzej Katunin, Kamil Fedio, "On a Visualization of the Convergence of the Boundary of Generalized Mandelbrot Set to $(n-1)$ -Sphere". Journal of Applied Mathematics and Computational Mechanics, doi:10.17512/jamcm.2015.1.06. 2015.
- [10] John Baez, "Platonic Solids in All Dimensions", 11 Jun 2020, <https://math.ucr.edu/home/baez/platonic.html>
- [11] D.M.Y. Sommerville, "An Introduction to The Geometry of N Dimensions". Methuen & Co., London, ISBN-10: 1781830312, 1929.
- [12] H.S.M. Coxeter, "Regular Polytopes (3rd ed.)". ISBN-13: 978-0486614809, 1973.
- [13] Martin Henk, Jürgen Richter-Gebert, Günter M. Ziegler, "Basic Properties of Convex Polytopes". Handbook of Discrete and Computational Geometry, ISBN9781315119601, 1997.
- [14] Andrzej Katunin, "Fractals based on regular convex polytopes". Scientific Research of the Institute of Mathematics and Computer Science, 2012, Volume 11, Issue 2, pages 53-62.
- [15] "Chapter 3, Lebesgue and Hausdorff Measures" The Chinese University of Hong Kong, Math 5011.
- [16] Jake Gipple, "The Volume of n -balls". Rose-Hulman Undergraduate Mathematics Journal, Vol. 15, Issue 1 Art. 14. 2014.
- [17] Ralph Heiner Buchholz, "Perfect Pyramids". Bulletin of the Australian Mathematical Society, 45(3), 353-368. doi:10.1017/S0004972700030252. 1991.
- [18] Mathieu Desbrun, Eva Kanso, Yiyang Tong, "Discrete Differential Forms for Computational Modeling". Discrete Differential Geometry. Oberwolfach Seminars, vol 38. Birkhäuser Basel (2008).
- [19] Aimee Kunnen, Steven Schlicker, "Regular Sierpinski polyhedral". Pi Mu Epsilon J. 1998, 10, 607-619.
- [20] Pedro Juan Roig, Salvador Alcaraz, Katja Gilly, Carlos Juiz, "Applying Multidimensional Geometry to Basic Data Centre Designs". Journal of Electrical and Computer Engineering Research, Vol. 1, No. 1, 2021.