
Review

Shiftless, A Critical Piece of the Innate Immune Response to Viral Infection

William Rodriguez¹ and Mandy Muller^{1*}

¹ Department of Microbiology, University of Massachusetts, Amherst, USA

* Correspondence: mandymuller@umass.edu

Abstract: Since its initial characterization in 2016, the interferon stimulated gene Shiftless (SHFL) has proven to be a critical piece of the innate immune response to viral infection. SHFL expression stringently restricts the replication of multiple DNA, RNA, and retroviruses with an extraordinary diversity of mechanisms that differ from one virus to the next. These inhibitory strategies include the negative regulation of viral RNA stability, translation, and even the manipulation of RNA granule formation during viral infection. Even more surprisingly, SHFL is the first human protein found to directly inhibit the activity of the -1 programmed ribosomal frameshift, a translation recoding strategy utilized across nearly all domains of life and a several human viruses. Recent literature has shown that SHFL expression also significantly impacts viral pathogenesis in mouse models, highlighting its *in-vivo* efficacy. To help reconcile the many mechanisms by which SHFL restricts viral replication, we provide here a comprehensive review of this complex ISG, its influence over viral RNA fate, and the implications of its functions on the virus-host arms race for control of the cell.

Keywords: C19ORF66; FLJ11286; shiftless; SVA-1; RyDEN; IRAV; ISG; innate immune response; RNA stability; translation; RNA granules; ribosomal frameshift

1. Introduction

To overcome viral infection and ensure survival, cells have evolved a robust and efficient array of antiviral defenses. These defenses manifest as complex signaling cascades tied to equally elaborate shifts in cellular gene expression to regain control of cell fate. While the sensing of viral products - nucleic acids and proteins - is an essential first step in this process, the true sword of the intrinsic immune response are the thousands of interferon stimulated genes (ISG) that work to establish an anti-viral state[1-4]. Each of these ISG target distinct steps of viral infection from viral entry and viral gene expression to virion assembly and egress (reviewed here [5]). Given the stunning diversity among viruses, ISGs are correspondingly adaptable at targeting multiple steps simultaneously to efficiently stamp out infection. One such ISG, *C19orf66* (herein referred to as Shiftless (SHFL)), has proven to be among the most versatile and yet still enigmatic anti-viral host factors identified.

An increasing number of large-scale screens have independently identified SHFL as both an anti-viral factor and downstream effector of the Type I, II, and III interferon response to viral infection [6-13]. However, some studies have gone even further to characterize SHFL as a potent, broad-spectrum virus restriction factor for a growing list of DNA, RNA, and Retroviruses (Summarized in Table 1) [14-25]. Many ISGs have evolved to target viral gene expression as it is often the primary battleground that nearly all viral replication steps depend upon. However, while a few multifunctional ISGs have been described, they rarely demonstrate the same breadth of restriction mechanisms as SHFL. So far, for each virus SHFL has been shown to restrict, a novel mechanism has been described. With each study of SHFL, numerous avenues of research have emerged into RNA stability, ribosomal frameshifts, and even the regulation of RNA granule formation both outside of and within the context of viral infection. The study of SHFL has improved our

understanding of the pathways that govern the balance between host and viral gene expression and continues to provide insights into weaknesses in the viral life cycle that can be exploited for therapeutic development. Here, we provide a comprehensive review of the many mechanisms by which SHFL restricts viral infection, the consequences of its many functions on the regulation viral and host RNA fate, and the implications of SHFL role as a critical piece of the innate immune response to viral infection.

2. Shiftless: An ISG of Many Names.

Located on chromosome 19 (NC_000019.10), *C19orf66* is an eight exon long gene that encodes 4 isoforms, with the full-length transcript (C19ORF66-201) (1928 bp) encoding a 291 amino acid (aa) protein of 33 kD molecular weight, C19ORF66. In nearly every context, this protein was identified as a broad-spectrum anti-viral factor whose expression is up-regulated in response to Type I interferon (IFN) induction and generally viral infection. Six of these studies have attributed several unique identifiers to the *C19orf66* gene product including FLJ11286 (genomic location), Repressor of Yield of Dengue Virus (RyDEN), Suppressor of Viral Activity-1 (SVA-1), Interferon Regulated Anti-Viral gene (IRAV), and more recently Shiftless (SFL/SHFL) with SHFL finally approved for its official gene symbol [14-25]. SHFL expression can often vary dramatically between cell types as demonstrated by Balinsky and colleagues [16]. This is in-line with previous observations of differences in both IFN induction and ISG expression among diverse cell lines [26-28].

The predicted protein structure of SHFL shares little sequence homology with other human proteins. From Jarred structural prediction by Suzuki and colleagues and our own Alphafold2 structural prediction, SHFL consists of eight α -helices, seven β -strands, a zinc-ribbon motif (aa 112-135), a coiled-coil motif (aa 261-285), a nuclear localization signal (NLS) (aa 121-173), a nuclear export signal (NES) (aa 261-269), and lastly a Glutamic acid (E) rich motif in the C-terminus [14] (Figure 1A). Apart from secondary structure predictions, SHFL also has two validated functional domains which we define here as the PABPC-binding domain (PABPC-BD) (aa102-150) and -1PRF domain (aa164-199), both of which will be discussed in more detail in later sections [14; 18].

Several groups have hypothesized, although not experimentally validated, that SHFL has the capacity to dimerize with itself *via* its putative C-terminal coiled-coil motif. These higher order SHFL complexes would then be more readily able to bind its mRNA and ribosomal subunit targets to facilitate its regulatory functions relative to translation. Interestingly, none of the top 5 ranked SHFL Alphafold2 predicted structures reflect a coiled-coil motif, rather, a single alpha helix was predicted to form across the far C-terminal end of SHFL (Figure 1B). This structural prediction would in-fact still support the idea of a SHFL dimerization, as the two C-terminal coils could in-fact interact with one another to bridge the interaction in place of a coiled-coil motif. Importantly, this domain of SHFL also contains the critical NES, and therefore several layers of SHFL function may be simultaneously tied to this region of the protein. Our current understanding of SHFL structure relative to its multitude of functions is undoubtedly complex. The overlap between NLS and NES signals with known functional domains, the capacity to dimerize, and the myriad of different functions all culminate into an exciting challenge for understanding SHFL as a protein moving forward. However, incredibly important milestones such as distinct residues involved in RNA binding, a growing list of validated interactors, and reports of multiple SHFL mechanisms acting simultaneously continue to set a critical foundation for future SHFL structural studies.




Figure 1. Shiftless Protein Structure. Shiftless is a 291aa long protein of 33 kDa molecular weight. (A) Highlighted in Red is the PABPC-binding domain (PABPC-BD) which also encompasses the Zinc-Ribbon Domain (112-135) and the Nuclear Localization Signal (121-173). Highlighted in Blue is the -1 Programmed Ribosomal Frameshift (PRF) (169-199). Highlighted in Yellow is the C-terminal domain containing the Glutamic Acid (E)-Rich Domain (270-286) and the Nuclear Export Signal (261-269). (B) Alphafold2 predicted protein structure of Shiftless. The PABPC-BD, -1PRF Domain, and the C-terminal Domain are Red, Blue, and Yellow respectively. Also highlighted in orange is the Nuclear Localization Signal and in pink is the Nuclear Export Signal [29; 30].

In terms of biological function, SHFL itself remains rather enigmatic. To date, no enzymatic functions have been attributed to SHFL itself. As such, SHFL functionality has often been tied to its interactions with a diverse set of cellular proteins (Table 1). The SHFL interactome across various viral infection backgrounds is most often enriched in RNA-binding proteins, most notably Poly-A Binding Protein Cytoplasmic 1 (PABPC) and the Moloney Leukemia Virus 10 (MOV10) proteins which will be discussed in more detail in Section 3. In-line with these observations, SHFL itself has been experimentally validated as a bona fide RNA-binding protein with a strong, yet non-specific affinity for ssRNA [16; 18; 22]. While specific RNA binding domains have yet to be fully mapped out, Recent *in vitro* work by Napthine and colleagues demonstrated that three arginine residues (R131, R133, and R136, Figure 1) within the predicted zinc-finger motif are critical for SHFL RNA binding [22]. This observation is further reinforced by the findings of three other SHFL studies that also mutagenize the SHFL NLS at the same three arginine residues albeit with additional mutated residues in select cases [14; 19; 24]. Furthermore, it appears that the binding interaction between SHFL and RNA-binding proteins (RBP) such as PABPC substantially enhances the binding affinity between SHFL and its target RNA [14]. These are rather surprising observations in consideration of findings by Wang and colleagues which identified an entirely separate domain of SHFL, here referred to as the -1PRF domain, that appeared to be necessary for its interaction with target RNA [18]. Thus, while the zing-

finger motif/NLS of SHFL is demonstrably critical for SHFL RNA binding, it will be important moving forward to investigate potential synergistic functions across these seemingly disparate SHFL domains that could explain these complex observations of SHFL functionality.

3. Shiftless and Viral RNA Fate.

The life of an RNA from the moment of transcription, execution of its functional output (i.e., translation), and unto its eventual decay, is collectively referred to as RNA fate [31-33]. While transcription and translation are considered the cornerstones of gene expression, RNA fate is undoubtedly the linchpin that inexorably ties these two processes together. Studies across years of RNA biology have uncovered hundreds of positive and negative feedback loops that allow the cell to finely tune gene expression in response to environmental challenges, especially that of viral infection [34-38]. This interconnectedness presents a significant challenge when investigating the mechanisms of RNA-binding proteins such as SHFL. Particularly in the context of virus replication, which almost always coincides with dramatic alterations to the global RNP landscape. In this section, we will review the relationship between SHFL and its capacity to influence RNA fate during viral infection.

3.1. SHFL and Viral RNA Translation

In 2016, Suzuki and colleagues set out to identify novel anti-viral factors that are up-regulated in response type I IFN treatment and are refractory to Dengue Virus (DENV) infection [14]. DENV is a positive sense ssRNA virus belonging to Flaviviridae, a large family of arthropod borne RNA viruses. Like most flaviviruses, following receptor mediated endocytosis, DENV replication takes place exclusively in the cytoplasm. DENV proteins work in concert to execute viral replication while simultaneously rewiring host cell organization and restricting the activation of the innate immune response [39]. Striking back against this manipulation, a plethora of ISG including IFIT2, IFIT1, ISG15 and TRIM69 have been shown to restrict flavivirus replication [40-42]. From their screen, Suzuki and colleagues identified yet another ISG, *C19orf66*, (referred to as RyDEN in the study) as capable of conferring greater survivability to DENV infected cells and is upregulated in response type I, II, and III interferon treatment. Upon further investigation, they next demonstrated using a knockdown approach that steady-state levels of SHFL are sufficient in restricting the replication of all four DENV serotypes. Further investigation revealed that a dramatic downregulation of both viral RNA and viral protein levels when SHFL was overexpressed in DENV infected cells. Compounding this, Suzuki and colleagues would be the first to demonstrate that loss of SHFL expression *via* knockdown (KD) results in catastrophic loss of IFN-mediated protection of the host cell upon DENV infection. Thus, highlighting that SHFL is a vital piece of the innate immune response to DENV infection in human cells. To facilitate this restriction of DENV, SHFL was shown to interact with La-related protein 1 (LARP1) and PABPC, two RBP that are both known to regulate mRNA stability and the cellular translation apparatus (Both extensively reviewed here [43]). Interestingly, in this same study, Suzuki and colleagues showed that both PABPC and LARP1 act as pro-viral factors during DENV infection as previously reported [44]. This could suggest that SHFL repurposes both proteins toward antiviral functions during flavivirus infection. Exploring the structure further, the authors next mapped SHFL anti-viral capacity to its NLS located within the PABPC-BD. Loss of this domain leads to an inability of SHFL to impact DENV replication and binding to PABPC as per its namesake. Recent work by Napthine and colleagues has shown that mutation of three arginine residues to alanines (R131A, R133A, and R136A, Figure 1) within the NLS results in the loss of SHFL RNA binding capacity [22]. One key difference to this study, however, is that Suzuki and colleagues also mutagenized several other arginine residues throughout the PABPC binding domain. Since Suzuki and colleagues showed that SHFL itself binds to the 3'UTR of DENV genomic RNA (gRNA), this NLS mutant thus proved that

SHFL can bind viral RNA independently of PABPC, albeit with less affinity. Given SHFL ability to bind DENV gRNA, the authors next explored SHFL possible influence over DENV translation. Interestingly, SHFL stringently restricted the translation of DENV but did not significantly impact global cellular translation of the host. Thus, in their final model, Suzuki and colleagues concluded that through its binding to DENV 3'UTR, SHFL can restricts its translation while possibly also triggering viral RNA degradation.

Excitingly, a recent study by Hanners and colleagues recently studied the *in-vivo* efficacy of SHFL and broadly addressed its capacity to restrict a wide range of positive and negative sense RNA viruses [21]. To identify ISG with strong anti-viral activity against the flavivirus West Nile Virus (WNV), Hanners and colleagues utilized a large lentiviral library of ISGs and measured WNV infectivity for each ISG tested. Among roughly 387 candidate genes, they found SHFL exhibited a potent anti-viral effect against WNV. When testing the specificity of the SHFL phenotype, they found that SHFL was able to restrict all Flaviviruses tested including Yellow Fever Virus (YFV), WNV, DENV, and Zika Virus (ZIKV) and several other positive sense RNA viruses including Equine arteritis virus (EAV), Coxsackievirus B3 virus (CVB3), Venezuelan Equine Encephalitis virus (VEEV), Sindbis virus (SINV), and O'nyong'nyong virus (ONNV). Surprisingly, they did not observe an effect of SHFL on the infectivity of the human coronaviruses OC43 and SARS-CoV-2 or five negative sense RNA viruses including Vesicular stomatitis virus (VSV), parainfluenza virus 3 (PIV3), respiratory syncytial virus (RSV), measles (MV), or IAV. Given the breadth of SHFL among positive sense RNA viruses, this lack of an impact on negative sense RNA viruses most likely represents a staunch evolutionary divergence in the SHFL mechanism relative to differences in viral replication strategies. As has been posited previously by Suzuki and colleagues, this could boil down to difference between the RNP structures assembled on the viral genome and/or viral mRNA at the earliest stages of primary infection. Perhaps positive sense RNA viruses, readily translated, are more easily recognized by SHFL or rather SHFL is recruited to viral translation complexes that are recognized by specific SHFL co-factors and therein triggers viral RNA decay/translational arrest.

Hanners and colleagues further explored the mechanism by which SHFL restricts HCV and YFV replication and found that similarly to DENV, SHFL binds to viral RNA, does not affect primary translation, but was still capable of restricting later stages of viral infection. They also observed a distinct loss of HCV replication organelles which could explain the dramatic impact of SHFL on overall viral gene expression. Correspondingly, there was also a distinct restriction of dsRNA intermediates produced as byproducts on viral genome amplification. Lastly, Hanners and colleagues studied the impact of the loss of SHFL on infection by the flavivirus ZIKV and alphavirus SINV in mice lacking SHFL expression. SHFL knockout (KO) mice displayed a more rapid onset of symptoms and increased severity of clinical disease compared to control mice, highlighting the significant role SHFL plays in viral pathogenesis in an *in-vivo* model. Interestingly, SHFL KO mice also exhibited enhanced inflammation and ZIKV titers in the brain and spinal cord but no observable dysregulation of cytokine levels. This suggests that the cause of lethality is likely due to increased viral replication in the central nervous system and not an effect on the overall immune response to ZIKV infection. Thus, SHFL may have a significant impact on the neuroinvasiveness of ZIKV and perhaps other neurotropic flaviviruses such as WNV and Japanese Encephalitis virus (JEV) [45; 46]. Given the myriad of mechanisms described thus far, several possibilities present themselves. First, SHFL targeting of viral RNA may be governed by the expression/functionality of a diverse set of RBP that bind to and facilitate subsequent rounds of viral translation. Second, SHFL could either, as suggested by Balinsky and colleagues, re-localize viral RNA to sites of RNA decay (namely P-bodies) or recruit RBPs involved in RNA decay directly to the viral genome. Lastly, SHFL could be acting on pathways completely independent of viral products as has been suggested recently by Kinast and colleagues regarding the biophysical formation of HCV replication compartments, known as the HCV membranous web, following infection. This observation was also observed by Hanners and colleagues with YFV by electron

microscopy [19]. Much work remains to parse out each of these different possibilities, a challenge that mirrors the interconnectedness of RNA stability and translation during viral infection.

3.2. *Shiftless and Viral RNA Stability*

While each of the studies discussed thus far identified SHFL as a downstream effector of the IFN response, not all studies of SHFL study discovered it within these confines. In 2019, our group set out to investigate the back-and-forth struggle between herpesviruses and their hosts for control of cellular mRNA stability during viral infection [17]. During lytic replication, herpesviruses rapidly seize control of the cellular gene expression apparatus by triggering a massive cellular RNA decay event termed “host-shutoff” [47-49]. This global RNA decay is orchestrated within the cytoplasm by a single viral endoribonuclease. This strategy is surprisingly well conserved among related alpha- and gammaherpesviruses, including Kaposi’s sarcoma-associated herpesvirus (KSHV) ORF37 (SOX), Herpes Simplex virus-1 (HSV-1) vhs, Epstein-Barr Virus (EBV) BGLF5, and the Murine Herpesvirus 68 (MHV68) muSOX [50-52]. There are also several RNA viruses that have evolved the same host-shutoff strategy, including several human coronaviruses and influenza A virus (IAV) [53-55]. Each of these viral endonucleases, though diverse in mechanism, ultimately trigger the internal cleavage of messenger RNA (mRNA) targets, rendering these transcripts susceptible to both 3' to 5' as well as 5' to 3' directional RNA decay machinery such as by the major cellular exonuclease XRN1 [48]. During KSHV infection, the viral endonuclease SOX is responsible for the degradation of about 70% of the host transcriptome, profoundly altering the host gene expression landscape [17]. Despite the global efficiency of host-shutoff, there remains a pool of cellular mRNAs that we and others have shown evade cleavage by SOX [52; 56; 57]. Among these “escapees” are those that actively evade SOX cleavage *via* an RNA element located within their 3' UTRs that we refer to as the “SOX Resistant Element” or SRE [57]. Among these SRE-bearing transcripts, we identified the SHFL mRNA and showed that it can evade cleavage by multiple herpesviral endonucleases as well as the Influenza A virus PA-X endonuclease [17]. We next showed that SHFL expression climbs over the course of KSHV lytic replication and remains diffusely cytoplasmic. Given SHFL previously demonstrated anti-viral capacity for DENV, we next found that SHFL also restricts KSHV infection, impacting nearly every step of the viral life cycle following lytic reactivation. The presence of an SRE on the SHFL mRNA coupled with its essential place in the innate immune response point starkly to a strong evolutionary imperative for its expression during viral infection. However, whether this impact on viral RNA stability is direct or indirect remains an important open question for our group.

3.3. *Shiftless and RNA Granules during RNA virus infection*

One pattern that emerges among studies of the SHFL interactome is that it is consistently dominated by constituents of phase-separated RNA granules [14; 16; 19; 25]. RNA-granules are membrane-free, phase-separated ribonucleoprotein (RNP) complexes that function in the storage, translational arrest, and/or degradation of RNA throughout the cell [58-61]. These granules vary in size, shape, and overall function between nucleus and the cytoplasm [62; 63]. In the cytoplasm, two distinct granule types have gained increasing attention in the past 15 years, these include Processing bodies (P-bodies) and Stress Granules (SG). In a broader biological context, SGs and P-bodies are conserved across all eukaryotes and have been implicated in multiple processes ranging from stem cell differentiation to cancer development [64-67]. An important distinction between these two granule types is that SGs are exclusively stress induced, while P-Bodies are constitutive in some cell lines but increase in size and number in response to stress. Generally, it is accepted that SGs are sites of translational arrest while P-bodies are sites of RNA-storage and decay. However, these functions relative to the phenotypes that regulate their formation remain an active area of research and is often contested [68-72]. Interestingly, the

formation of both SGs and P-bodies are also induced during viral infection by a diverse array of viruses [73-77]. A growing number of studies have begun to find a range of anti-viral roles for RNA granules during viral infection, many of which often link back to the translational arrest or degradation of viral RNA at these sites. However, the extent to which these cytoplasmic RNP granules facilitate anti-viral function or in some cases are coopted by viruses to facilitate replication remain to be explored [78-81]. It is important to note that while the function of RNA granules has been actively contested in the literature, it is generally agreed that their formation is still reflective of broader changes in gene expression. P-bodies for example are constitutively formed in most human cell types and are often a consequence of the activity of RNA-induced silencing complexes (RISC) which mediate the translational silencing of cellular transcripts [82-84]. SG on the other hand are formed often as a direct consequence of cellular translational arrest, a phenotype that can be both naturally and artificially induced with chemical/environmental stimuli such as Sodium Arsenate and cold shock [85-87]. In either case, RNP granules allow the cell a substantial level of control over gene expression, acting as “membraneless organelles” that allow for the fine-tuning of RNA stability and protein translation in response to challenges to cell health.

The next SHFL study by Balinsky and colleagues (here referred to as IRAV) would go on to verify that SHFL expression is upregulated in response to DENV infection [16]. They first demonstrated that SHFL itself is an ISG whose expression is coupled to the canonical IFN-induced ISG3 pathway in an IFN- β dependent manner. It was then shown that SHFL restricts the replication of both DENV and Encephalomyocarditis virus (EMCV), drastically decreasing viral titers and viral RNA levels. To dig further into the mechanism of DENV/EMCV restriction, Balinsky and colleagues next showed that SHFL, whose steady-state localization is often diffusely cytoplasmic, is re-localized to the DENV replication complex, co-localizing with the viral proteins NS3 and NS4A, both of which are essential to the capping and amplification of DENV gRNA. Mirroring findings by Suzuki and colleagues, SHFL was also shown to interact with several known RBP that play critical roles in RNA stability including the RNA helicases MOV10, UPF1, and AU-rich element (ARE) mRNA binding protein HuR [88-90]. Given that SHFL interactors are predominately RNP-granule constituents, Balinsky and colleagues proceeded to investigate whether SHFL localizes to P-bodies in response to IFN treatment. Interestingly, they found that SHFL does in-fact re-localize to P-bodies in IFN-treated cells. This a particularly interesting observation considering this experiment was performed in the absence of DENV infection. SHFL granularization in response to IFN may suggest SHFL functions over gene expression extends beyond viral infection. Exploring the SHFL interactome further, Balinsky and colleagues lastly confirmed the co-localization of SHFL with MOV10 in DENV infected cells, both of which re-localize to the DENV replication complex to restrict DENV replication. Importantly, several studies have previously shown that MOV10, a known P-body constituent, is itself an ISG with several anti-viral functions (Reviewed Here [91]). This represents one of the first studies to demonstrate a direct relationship between the anti-viral function of SHFL and phase-separated RNA granules, providing a novel link between the innate immune response to viral infection and RNP granule dynamics.

Further reinforcing the ties between SHFL and RNA granules, recent work by Kinast and colleagues also demonstrated that SHFL can re-localize to SG upon HCV infection [19]. Interestingly, Kinast and colleagues also found that a Zinc-finger mutant, was unable to restrict HCV infection. Given this RBP-enriched interactome, one outstanding question remains: What is the direct functional link between SHFL and RNA granules? More specifically, what benefit would RNA phase-separation have toward the capacity of SHFL to bind viral RNA or regulate viral gene translation? These are questions that unfortunately remain difficult to answer with the current lack of studies tracking the localization of SHFL relative to target RNA transcripts. In a similar vein, the size and shape of the granules reported to date also suggest that there may be human transcripts also caught by

SHFL in its attempt to establish an anti-viral state. What these transcripts are, and whether they serve pro-viral roles remains entirely underexplored.

3.4. *Shiftless and RNA Granules during DNA Virus Infection*

In a recent preprint released by our group, we set out to dive deeper into the relationship between SHFL and KSHV lytic replication, seeking to understand the mechanism by which SHFL restricts lytic viral gene expression [25]. We mapped SHFL interactome across KSHV infection and, in-line with previous SHFL literature discussed here, we showed that SHFL interactome is dominated by protein constituents of both P-bodies and SG. As such, we further explored whether SHFL localizes to RNA granules and demonstrated for the first time that exogenous expression of SHFL leads to a striking disassembly of P-bodies and a simultaneous induction of SG-like densities. It is important to note that in all our cell lines, there was no observed reduction in expression levels for either of our P-body markers suggesting that this is a true disassembly of these granules. Our observations raise several questions: first, how does SHFL trigger P-body disassembly? Given the lack of change in marker expression, one hypothesis is that SHFL could promote the diffusion of these markers across the cytoplasm to facilitate its anti-viral function. Second, what does this suggest about P-bodies relative to SHFL being an ISG? Most studies thus far, suggest that P-body formation is likely refractory to herpesviral infection [92; 93]. However, their disassembly by SHFL could also suggest that there may be a pro-viral, dual functionality of P-bodies that is yet to be described that SHFL restricts. Lastly, what are these SG-like densities that form upon SHFL induction? Several works throughout this review highlight SHFL capacity to influence and interface with various components of the translational machinery and as such its ability to induce translational arrest of viral/host transcripts could be reflected by these densities. Among these transcripts which comprise these granules could be those that encode components of the RISC complex, which could explain why we see a concomitant disassembly of P-bodies [84]. Needless to say, these observations warrant extensive further investigation but are undoubtedly an exciting frontier in understanding the place of SHFL in regulating so many different facets of the broader axis of gene expression between RNA stability and translation.

To date, the interplay between these two arms of the anti-viral response, RNA granules and ISGs, remains an active area of research. A growing number of studies have demonstrated that IFN signaling proteins often re-localize to SG upon viral infection. Furthermore, several anti-viral proteins, some of which also ISGs, are known constituents of RNP granules including FUS, MOV10, and more recently TDRD3 [94; 95]. These observations all raise critical questions regarding SHFL: Does SHFL restrict the expression of select mRNA by re-localizing them to P-bodies as a part of the IFN response? If so, how does this relate to SHFL disassembly of P-bodies when exogenously expressed? Do these target mRNA otherwise serve pro-viral roles? And if so, what are these transcripts and what dictates their targeting by SHFL from cell type to cell type? And lastly, could these differences in targeting explain the expansive breadth SHFL modes of function? Complicating this further, work by Suzuki and colleagues also reports that SHFL had no impact on global gene translation, which is contradictory to what would be assumed of its impact on translation and SG localization. This could suggest that SHFL-containing granules, though reminiscent of SG in composition, may not reflect a global impact on translation but rather a localized accumulation of specific mRNAs. Nonetheless, these questions will be critical to assess moving forward, as RNA granules (and phase-separation itself) could be a critical missing link that may bridge several SHFL mechanisms.

4. Shiftless and the -1 Programmed Ribosomal Frameshift.

While we still cannot fully mechanistically define the SHFL anti-viral strategies discussed thus far, most are similar to strategies employed by other ISGs expressed in response to viral infection. However, in this section we will discuss a mechanism that is unique to SHFL and is especially effective against a subset of human viruses including

retroviruses such as Human Immunodeficiency virus-1 (HIV-1), select flaviviruses such as WNV and JEV and of recent notoriety coronaviruses such as SARS-CoV-2 [96-99].

In 2019, a study by Wang, Xinlu and colleagues leveraged the anti-viral response to HIV-1 to identify novel host factors that modulate a viral translation strategy known as the -1 programmed ribosomal frameshift (-1PRF) [18]. The -1PRF is a translation-recoding strategy to expand genetic coding capacity, where an actively translating ribosomes slips back one nucleotide on select viral open reading frames, resulting in translation of a new reading frame (Reviewed Here [100]). The -1PRF signal itself is a *cis*-acting RNA element consisting of two RNA motifs: one is a heptameric slippery sequence X XXY YYZ, wherein X is any nucleotide; Y is A or U; and Z is A, U, or C. Prior to the shift in the -1 direction, the original frame (0 frame), are ordered as codons XXY and YYZ, but in the -1 frame, the reading frame is back shifted one nucleotide resulting in the codons XXX and YYY. The second RNA motif is an RNA secondary structure that serves as stimulatory signal to the ribosome known as an RNA pseudoknot downstream of the slippery sequence. During lytic replication, HIV-1 uses a -1PRF cue to control the ratio between the Gag and Gag-Pol polyproteins [101]. Several groups have shown that mutagenesis of this region can severely inhibit this frameshift which results in a stringent restriction of HIV-1 productive infection [102; 103]. Using a screen for IFN-linked effectors modulating the -1-frameshift efficiency of Gag-Pol, Wang, Xinlu and colleagues identified SHFL, which displayed the highest inhibitory activity against the -1PRF reporter construct assayed. This is how SHFL became "SHFL", which stands for Shiftless. Upon further investigation, SHFL was shown to restrict HIV-1 replication by directly disrupting the balance between Gag and Gag-Pol expression in favor of a short-form premature-translation termination (PMT) product. Furthermore, when testing the inhibitory capacity of SHFL against other -1PRFs, Wang, Xinlu and colleagues found that SHFL appears to be a broad inhibitor of the -1PRF for multiple retroviruses including Rous sarcoma virus (RSV), Mouse mammary tumor virus (MMTV), Human T-lymphotropic virus (HTLV), HIV-2, and SIV. Surprisingly, two notable human mRNAs, CCR5 and PEG10, also contain -1PRF signals that are also restricted by SHFL suggesting that SHFL may be the only known trans-acting human protein capable of restricting -1PRF signals [104; 105]. To better understand the mechanisms by which SHFL restricts the -1PRF, Wang, Xinlu and colleagues next investigated SHFL interactome and found that SHFL directly interfaces with active ribosomes and target -1PRF bearing mRNAs. This binding of SHFL to -1PRF transcripts triggers a PMT event that leads to the enrichment of shortform products from the shorter open reading frame. The authors then posited that this inhibition may be due to a restriction of a non-canonical ribosome rotation by SHFL through recruitment of factors that facilitate PMT. In line with this, SHFL was found to bind to the ribosomal binding protein Eukaryotic polypeptide chain release factor 3 (eRF3) in complex with eRF1 which are both known to help facilitate ribosome rotation [106]. Thus, it was concluded that SHFL binds to -1PRF bearing mRNAs in response to ribosome stalling at the -1PRF element and therein triggers a PMT event that leads to the formation of the PMT product. This study marks SHFL as a unique human gene capable of regulating the -1 frameshifts, further emphasizing the extraordinary versatility of this ISG.

Recent studies have also followed up on this unique feature of SHFL, testing its ability to modulate -1PRFs outside of retroviruses. A recent study by Sun and colleagues found that SHFL could restrict the -1PRF signal within SARS-CoV-2 genome *in-vitro*. While this would imply that SHFL has the capacity to restrict SARS-CoV-2 replication, the recent report by Hanners and colleagues showed that SHFL was unable to restrict SARS-CoV-2 infection [21]. This could suggest that inhibition of -1PRF is not necessarily sufficient to restrict SARS-CoV-2 or that other confounding factors could prevent SHFL function during coronavirus infection *in-vivo*. Fascinatingly, another recent study has also showed that -1PRF mechanisms can work in tandem with other SHFL anti-viral strategies to restrict flavivirus infection. In a study conducted by Yu and colleagues, SHFL was identified as a potent anti-viral ISG capable of restricting JEV, yet another neurogenic flavivirus [24]. However, surprisingly, this group reported that SHFL not only inhibited the

-1PRF-mediated expression of NS1' but also triggered the lysosomal degradation of JEV NS3 in a manner parallel to findings by Wu and colleagues for ZIKV NS3 [20]. Of note, the capacity of SHFL to modulate these -1PRF elements was mapped by Wang, Xinlu and colleagues to a C-terminal domain of SHFL (aa164-199, labeled in **Figure 1** as “-1PRF Domain”). SHFL lacking the -1PRF domain mirrors a true isoform of SHFL (C19orf66-209) which is expressed in select cell types but to date has displayed no attributable anti-viral function [18]. Interestingly, work by Napthine and colleagues, who demonstrated that bacterially expressed, purified SHFL can inhibit the -1PRF, also showed that this domain of SHFL is dispensable for its ability to restrict the -1PRF *in-vitro* [22]. They also showed that SHFL binds indiscriminately to single stranded RNA substrates regardless of the presence or absence of the -1PRF. These findings directly contradict those made previously by Wang, Xinlu and Yu and colleagues more recently and therefore warrant further investigation. Taken together, with SHFL influence over viral RNA and viral protein stability, SHFL has emerged a critical regulator of viral gene expression. But observations of SHFL influence over human -1PRF signals, such as in CCR5 and PEG10, also suggest that SHFL phenotype may have a much broader impact on cellular gene expression than previously hypothesized.

Table 1. Summary of Shiftless Studies to-date. Table listing each of study of Shiftless and its capacity to restrict viral infection through diverse mechanisms, human protein interactions, and targets. Studies marked with * also examined the impact of Shiftless on multiple viruses in larger panels.

Authors	Viruses Studied	Restriction Strategy	Target	Human Protein Interactors	RNA Granule Association	Reference
Suzuki (2016)	DENV (Serotypes 1-4)*	Viral Translation, Viral RNA Stability	Genomic RNA	PABPC, LARP1	-	[14]
Xiong (2016)	HIV-1	Viral Gene Expression	p24	-	-	[15]
Balinsky (2017)	DENV and EMCV	Viral RNA Stability	Genomic RNA	UPF1, HuR, MOV10	Localization to P-bodies	[16]
Rodriguez (2019)	KSHV	Viral Gene Expression	Early/Delayed Early Genes	-	-	[17]
Wang, Xinlu (2019)	HIV-1	(-1) Frameshift	Ratio of Gag-Pol	Ribosomal Subunits, eRF1, eRF3	-	[18]
Kinast (2020)	HCV	Viral Replication Compartment	Membranous Web, PI(4)P	-	Localization to Stress Granules	[19]
Wu (2020)	ZIKV	Lysosomal Degradation	NS3	-	-	[20]
Hanners (2021)	YFV and HCV*	Viral Gene Expression	Genomic RNA	-	-	[21]
Wang, Hua (2021)	PEDV	Ubiquitinylation-based Degradation	Nucleocapsid (N) Protein	MARCH8	-	[23]
Yu (2021)	JEV	(-1) Frameshift, Lysosomal Degradation	NS1'-NS1 ratio, NS3	-	-	[24]
Rodriguez (2022)	KSHV	Viral Gene Expression	ORF57	PABPC	Restricts P-Body Formation, Stress Granule-like Densities	[25]

6. Conclusions

Of the human ISGs studied to date, few have demonstrated such extraordinary versatility in restricting viral infection as SHFL. While our present review has covered much of our current understanding of SHFL and viral infection, new studies continue to unveil even more complex mechanisms-of-action. Recent work by Wu and colleagues as well as Wang, Hua and colleagues have both begun to tie SHFL function directly to the anti-viral capacity of autophagy and the ubiquitination pathways respectively [20; 23]. For certain viruses, as suggested by work by Kinast and colleagues, these mechanisms may also be entirely distinct from those directly targeting viral replication products [19; 21]. A more exciting possibility and a significant challenge to overcome moving forward will be to determine which of the many SHFL mechanisms are applicable from one virus to the next and within these systems are their multiple strategies working simultaneously as seen with JEV [24]. While not often consistent from one infection context to the next, further investigation of how SHFL regulates both viral and host mRNA along the axis of RNA stability and translation will likely prove to be essential for one or more of the mechanisms discussed here. In line with this, understanding the relationship between SHFL and RNA granules may lead to novel insights into a broader influence of SHFL over global gene expression and how this relates to the establishment of an anti-viral state. Taken together, SHFL has proven to be a critical piece of the innate immune response to viral infection that absolutely warrants further study. In doing so, we will undoubtedly uncover further weaknesses within viral replication and better understand the complex web that ties the innate immune response to the fate of RNA during viral infection.

Funding: Our work is funded by the National Institute of General Medical Sciences R35GM138043

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Christgen, S., Place, D. E., Zheng, M., Briard, B., Yamamoto, M., & Kanneganti, T. D. (2021). The IFN-inducible GTPase IRGB10 regulates viral replication and inflammasome activation during influenza A virus infection in mice. *European Journal of Immunology*.
2. Fernbach, S., Spieler, E. E., Busnadio, I., Karakus, U., Lkharrazi, A., Stertz, S., & Hale, B. G. (2022). Restriction factor screening identifies RABGAP1L-mediated disruption of endocytosis as a host antiviral defense. *Cell Reports*, 38(12), 110549.
3. Guo, X., Zeng, S., Ji, X., Meng, X., Lei, N., Yang, H., & Mu, X. (2022). Type I Interferon-Induced TMEM106A Blocks Attachment of EV-A71 Virus by Interacting With the Membrane Protein SCARB2. *Frontiers in Immunology*, 13.
4. Yang, X., Xiang, Z., Sun, Z., Ji, F., Ren, K., & Pan, D. (2022). Host MOV10 is induced to restrict herpes simplex virus 1 lytic infection by promoting type I interferon response. *PLOS Pathogens*, 18(2), e1010301.
5. Schoggins, J. W. (2019). Interferon-Stimulated Genes: What Do They All Do? *Annu Rev Virol*, 6(1), 567-584.
6. Bull, T. M., Meadows, C. A., Coldren, C. D., Moore, M., Sotto-Santiago, S. M., Nana-Sinkam, S. P., Campbell, T. B., & Geraci, M. W. (2008). Human herpesvirus-8 infection of primary pulmonary microvascular endothelial cells. *American journal of respiratory cell and molecular biology*, 39(6), 706-716.
7. Gaucher, D., Therrien, R., Kettaf, N., Angermann, B. R., Boucher, G., Filali-Mouhim, A., Moser, J. M., Mehta, R. S., Drake III, D. R., & Castro, E. (2008). Yellow fever vaccine induces integrated multilineage and polyfunctional immune responses. *The Journal of experimental medicine*, 205(13), 3119-3131.
8. Harvey, S. A., Romanowski, E. G., Yates, K. A., & Gordon, Y. J. (2005). Adenovirus-directed ocular innate immunity: the role of conjunctival defensin-like chemokines (IP-10, I-TAC) and phagocytic human defensin- α . *Investigative ophthalmology & visual science*, 46(10), 3657-3665.
9. Kash, J. C., Muñoz-Blanchard, E., Carter, V., Grosch, M., Perwitasari, O., Proll, S. C., Thomas, M. J., Weber, F., Klenk, H.-D., & Katze, M. G. (2006). Global suppression of the host antiviral response by Ebola-and Marburgviruses: increased antagonism of the type I interferon response is associated with enhanced virulence. *JOURNAL OF VIROLOGY*, 80(6), 3009-3020.
10. Miyazaki, D., Haruki, T., Takeda, S., Sasaki, S.-i., Yakura, K., Terasaka, Y., Komatsu, N., Yamagami, S., Touge, H., & Touge, C. (2011). Herpes simplex virus type 1-induced transcriptional networks of corneal endothelial cells indicate antigen presentation function. *Investigative ophthalmology & visual science*, 52(7), 4282-4293.
11. Wang, J., Nikrad, M. P., Phang, T., Gao, B., Alford, T., Ito, Y., Edeen, K., Travanty, E. A., Kosmider, B., & Hartshorn, K. (2011). Innate immune response to influenza A virus in differentiated human alveolar type II cells. *American journal of respiratory cell and molecular biology*, 45(3), 582-591.
12. Xiong, W., Contreras, D., Irudayam, J. I., Ali, A., Yang, O. O., & Arumugaswami, V. (2016). C19ORF66 is an interferon-stimulated gene (ISG) which Inhibits human immunodeficiency virus-1. *bioRxiv*, 050310.
13. Zapata, J. C., Carrion Jr, R., Patterson, J. L., Crasta, O., Zhang, Y., Mani, S., Jett, M., Poonia, B., Djavani, M., & White, D. M. (2013). Transcriptome analysis of human peripheral blood mononuclear cells exposed to Lassa virus and to the attenuated Mopeia/Lassa reassortant 29 (ML29), a vaccine candidate. *PLOS Neglected Tropical Diseases*, 7(9), e2406.
14. Suzuki, Y., Chin, W. X., Han, Q. E., Ichiyama, K., Lee, C. H., Eyo, Z. W., Ebina, H., Takahashi, H., Takahashi, C., Tan, B. H., Hishiki, T., Ohba, K., Matsuyama, T., Koyanagi, Y., Tan, Y. J., Sawasaki, T., Chu, J. J. H., Vasudevan, S. G., Sano, K., & Yamamoto, N. (2016). Characterization of RyDEN (C19orf66) as an Interferon-Stimulated Cellular Inhibitor against Dengue Virus Replication. *PLOS Pathogens*, 12(1), e1005357-e1005357.
15. Xiong, W., Contreras, D., Irudayam, J. I., Ali, A., Yang, O. O., & Arumugaswami, V. (2016). C19ORF66 is an Interferon-Stimulated Gene (ISG) which Inhibits Human Immunodeficiency Virus-1. *bioRxiv*, 050310-050310.

16. Balinsky, C. A., Schmeisser, H., Wells, A. I., Ganesan, S., Jin, T., Singh, K., & Zoon, K. C. (2017). IRAV (FLJ11286), an Interferon-Stimulated Gene with Antiviral Activity against Dengue Virus, Interacts with MOV10. *J Virol*, 91(5).

17. Rodriguez, W., Srivastav, K., & Muller, M. (2019). C19ORF66 Broadly Escapes Virus-Induced Endonuclease Cleavage and Restricts Kaposi's Sarcoma-Associated Herpesvirus. *J Virol*, 93(12).

18. Wang, X., Xuan, Y., Han, Y., Ding, X., Ye, K., Yang, F., Gao, P., Goff, S. P., & Gao, G. (2019). Regulation of HIV-1 Gag-Pol Expression by Shiftless, an Inhibitor of Programmed -1 Ribosomal Frameshifting. *Cell*, 176(3), 625-635.e614.

19. Kinast, V., Plociennikowska, A., Anggakusuma, Bracht, T., Todt, D., Brown, R. J. P., Boldanova, T., Zhang, Y., Bruggemann, Y., Friesland, M., Engelmann, M., Vieyres, G., Broering, R., Vondran, F. W. R., Heim, M. H., Sitek, B., Bartenschlager, R., Pietschmann, T., & Steinmann, E. (2020). C19orf66 is an interferon-induced inhibitor of HCV replication that restricts formation of the viral replication organelle. *J Hepatol*, 73(3), 549-558.

20. Wu, Y., Yang, X., Yao, Z., Dong, X., Zhang, D., Hu, Y., Zhang, S., Lin, J., Chen, J., An, S., Ye, H., Zhang, S., Qiu, Z., He, Z., Huang, M., Wei, G., & Zhu, X. (2020). C19orf66 interrupts Zika virus replication by inducing lysosomal degradation of viral NS3. *PLOS Neglected Tropical Diseases*, 14(3), e0008083-e0008083.

21. Hanners, N. W., Mar, K. B., Boys, I. N., Eitson, J. L., De la Cruz-Rivera, P. C., Richardson, R. B., Fan, W. C., Wight-Carter, M., & Schoggins, J. W. (2021). Shiftless inhibits flavivirus replication in vitro and is neuroprotective in a mouse model of Zika virus pathogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, 118(49), e211126611810.1073/pnas.2111266118

22. Napthine, S., Hill, C. H., Nugent, H. C. M., & Brierley, I. (2021). Modulation of Viral Programmed Ribosomal Frameshifting and Stop Codon Readthrough by the Host Restriction Factor Shiftless. *Viruses*, 13(7), 1230.

23. Wang, H., Kong, N., Jiao, Y., Dong, S., Sun, D., Chen, X., Zheng, H., Tong, W., Yu, H., Yu, L., Zhang, W., Tong, G., & Shan, T. (2021). EGR1 Suppresses Porcine Epidemic Diarrhea Virus Replication by Regulating IAV To Degrade Viral Nucleocapsid Protein. *J Virol*, 95(19), e0064521.

24. Yu, D., Zhao, Y., Pan, J., Yang, X., Liang, Z., Xie, S., & Cao, R. (2021). C19orf66 Inhibits Japanese Encephalitis Virus Replication by Targeting -1 PRF and the NS3 Protein. *Virologica Sinica* 2021, 1-13.

25. Rodriguez, W., Mehrmann, T., & Muller, M. (2022). Shiftless Restricts Viral Gene Expression and Influences RNA Granule Formation during KSHV lytic replication. *bioRxiv*.

26. Wu, X., Thi, V. L. D., Huang, Y., Billerbeck, E., Saha, D., Hoffmann, H.-H., Wang, Y., Silva, L. A. V., Sarbanes, S., & Sun, T. (2018). Intrinsic immunity shapes viral resistance of stem cells. *Cell*, 172(3), 423-438. e425.

27. Green, J. L., Osterhout, R. E., Klova, A. L., Merkwirth, C., McDonnell, S. R., Zavareh, R. B., Fuchs, B. C., Kamal, A., & Jakobsen, J. S. (2021). Molecular characterization of type I IFN-induced cytotoxicity in bladder cancer cells reveals biomarkers of resistance. *Molecular Therapy-Oncolytics*, 23, 547-559.

28. Aso, H., Ito, J., Koyanagi, Y., & Sato, K. (2019). Comparative description of the expression profile of interferon-stimulated genes in multiple cell lineages targeted by HIV-1 infection. *Frontiers in Microbiology*, 10, 429.

29. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., & Potapenko, A. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, 596(7873), 583-589.

30. Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D., Stroe, O., Wood, G., & Laydon, A. (2022). AlphaFold Protein Structure Database: Massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Research*, 50(D1), D439-D444.

31. Boo, S. H., & Kim, Y. K. (2020). The emerging role of RNA modifications in the regulation of mRNA stability. *Experimental & molecular medicine*, 52(3), 400-408.

32. Gillen, S. L., Giacomelli, C., Hodge, K., Zanivan, S., Bushell, M., & Wilczynska, A. (2021). Differential regulation of mRNA fate by the human Ccr4-Not complex is driven by coding sequence composition and mRNA localization. *Genome biology*, 22(1), 1-31.

33. Turner, M., & Diaz-Munoz, M. D. (2018). RNA-binding proteins control gene expression and cell fate in the immune system. *Nature Immunology*, 19(2), 120-129.

34. Dremel, S. E., Sivrich, F. L., Tucker, J. M., Glaunsinger, B. A., & DeLuca, N. A. (2022). Manipulation of RNA polymerase III by herpes simplex virus-1. *Nature Communications*, 13(1), 1-14.

35. Gilbertson, S., Federspiel, J. D., Hartenian, E., Cristea, I. M., & Glaunsinger, B. (2018). Changes in mRNA abundance drive shuttling of RNA binding proteins, linking cytoplasmic RNA degradation to transcription. *elife*, 7, e37663.

36. Liu, Z., Qin, Q., Wu, C., Li, H., Shou, J. n., Yang, Y., Gu, M., Ma, C., Lin, W., & Zou, Y. (2018). Downregulated NDR1 protein kinase inhibits innate immune response by initiating an miR146a-STAT1 feedback loop. *Nature Communications*, 9(1), 1-16.

37. Rigby, R. E., Wise, H. M., Smith, N., Digard, P., & Rehwinkel, J. (2019). PA-X antagonises MAVS-dependent accumulation of early type I interferon messenger RNAs during influenza A virus infection. *Scientific Reports*, 9(1), 1-13.

38. Zhao, Y., Ye, X., Shehata, M., Dunker, W., Xie, Z., & Karijolich, J. (2020). The RNA quality control pathway nonsense-mediated mRNA decay targets cellular and viral RNAs to restrict KSHV. *Nature Communications*, 11(1), 1-15.

39. Uno, N., & Ross, T. M. (2018). Dengue virus and the host innate immune response. *Emerging microbes & infections*, 7(1), 1-11.

40. Wang, K., Zou, C., Wang, X., Huang, C., Feng, T., Pan, W., Wu, Q., Wang, P., & Dai, J. (2018). Interferon-stimulated TRIM69 interrupts dengue virus replication by ubiquitinating viral nonstructural protein 3. *PLOS Pathogens*, 14(8), e1007287.

41. Singh, P. K., Singh, S., Farr, D., & Kumar, A. (2019). Interferon-stimulated gene 15 (ISG15) restricts Zika virus replication in primary human corneal epithelial cells. *The ocular surface*, 17(3), 551-559.

42. Desole, G., Sinigaglia, A., Riccetti, S., Masi, G., Pacenti, M., Trevisan, M., & Barzon, L. (2019). Modelling neurotropic flavivirus infection in human induced pluripotent stem cell-derived systems. *International Journal of Molecular Sciences*, 20(21), 5404.

43. Mattijssen, S., Kozlov, G., Fonseca, B. D., Gehring, K., & Maraia, R. J. (2021). LARP1 and LARP4: up close with PABP for mRNA 3'poly (A) protection and stabilization. *RNA Biology*, 18(2), 259-274.

44. Polacek, C., Friebe, P., & Harris, E. (2009). Poly (A)-binding protein binds to the non-polyadenylated 3' untranslated region of dengue virus and modulates translation efficiency. *Journal of General Virology*, 90(3), 687-692.

45. Zidovec-Lepej, S., Vilibic-Cavlek, T., Barbic, L., Ilic, M., Savic, V., Tabain, I., Ferenc, T., Grgic, I., Gorenec, L., & Bogdanic, M. (2021). Antiviral cytokine response in neuroinvasive and non-neuroinvasive west nile virus infection. *Viruses*, 13(2), 342.

46. Khou, C., Díaz-Salinas, M. A., da Costa, A., Préhaud, C., Jeannin, P., Afonso, P. V., Vignuzzi, M., Lafon, M., & Pardigon, N. (2021). Comparative analysis of neuroinvasion by Japanese encephalitis virulent and vaccine viral strains in an in vitro model of human blood-brain barrier. *PLoS ONE*, 16(6), e0252595.

47. Covarrubias, S., Richner, J. M., Clyde, K., Lee, Y. J., & Glaunsinger, B. A. (2009). Host Shutoff Is a Conserved Phenotype of Gammaherpesvirus Infection and Is Orchestrated Exclusively from the Cytoplasm. *JOURNAL OF VIROLOGY*, 83(18), 9554-9566.

48. Covarrubias, S., Gaglia, M. M., Kumar, G. R., Wong, W., Jackson, A. O., & Glaunsinger, B. A. (2011). Coordinated destruction of cellular messages in translation complexes by the gammaherpesvirus host shutoff factor and the Mammalian Exonuclease Xrn1. *PLOS Pathogens*, 7(10).

49. Richner, J. M., Clyde, K., Pezda, A. C., Cheng, B. Y. H., Wang, T., Kumar, G. R., Covarrubias, S., Coscoy, L., & Glaunsinger, B. (2011). Global mrna degradation during lytic gammaherpesvirus infection contributes to establishment of viral latency. *PLOS Pathogens*, 7(7).

50. He, T., Wang, M., Cheng, A., Yang, Q., Wu, Y., Jia, R., Liu, M., Zhu, D., Chen, S., Zhang, S., Zhao, X.-X., Huang, J., Sun, D., Mao, S., Ou, X., Wang, Y., Xu, Z., Chen, Z., Zhu, L., . . . Chen, X. (2020). Host shutoff activity of VHS and SOX-like proteins: role in viral survival and immune evasion. *Virology Journal*, 17(1), 68-68.

51. Daly, R., Khaperskyy, D. A., & Gaglia, M. M. (2020). Fine-tuning a blunt tool: Regulation of viral host shutoff RNases. *PLOS Pathogens*, 16(4), e1008385-e1008385.

52. Clyde, K., & Glaunsinger, B. A. (2011). Deep sequencing reveals direct targets of gammaherpesvirus-induced mRNA decay and suggests that multiple mechanisms govern cellular transcript escape. *PLoS ONE*, 6(5).

53. Schubert, K., Karousis, E. D., Jomaa, A., Scaiola, A., Echeverria, B., Gurzeler, L.-A., Leibundgut, M., Thiel, V., Mühlmann, O., & Ban, N. (2020). SARS-CoV-2 Nsp1 binds the ribosomal mRNA channel to inhibit translation. *Nature structural & molecular biology*, 27(10), 959-966.

54. Gaucherand, L., Porter, B. K., Levene, R. E., Price, E. L., Schmaling, S. K., Rycroft, C. H., Kevorkian, Y., McCormick, C., Khaperskyy, D. A., & Gaglia, M. M. (2019). The influenza A virus endoribonuclease PA-X usurps host mRNA processing machinery to limit host gene expression. *Cell Reports*, 27(3), 776-792. e777.

55. Bermudez, Y., Miles, J., & Muller, M. (2022). Nonstructural protein 1 (nsp1) widespread RNA decay phenotype varies among Coronaviruses. *bioRxiv*.

56. Muller, M., Hutin, S., Marigold, O., Li, K. H., Burlingame, A., & Glaunsinger, B. A. (2015). A Ribonucleoprotein Complex Protects the Interleukin-6 mRNA from Degradation by Distinct Herpesviral Endonucleases. *PLOS Pathogens*, 11(5), 1-23.

57. Muller, M., & Glaunsinger, B. A. (2017). Nuclease escape elements protect messenger RNA against cleavage by multiple viral endonucleases. *PLOS Pathogens*, 13(8).

58. Tian, S., Curnutt, H. A., & Trcek, T. (2020). RNA granules: a view from the RNA perspective. *Molecules*, 25(14), 3130.

59. Tauber, D., Tauber, G., & Parker, R. (2020). Mechanisms and regulation of RNA condensation in RNP granule formation. *Trends in Biochemical Sciences*, 45(9), 764-778.

60. Riggs, C. L., Kedersha, N., Ivanov, P., & Anderson, P. (2020). Mammalian stress granules and P bodies at a glance. *Journal of Cell Science*, 133(16), jcs242487.

61. Corbet, G. A., & Parker, R. (2019). RNP granule formation: lessons from P-Bodies and stress granules. *Cold Spring Harbor symposia on quantitative biology*,

62. Zhou, H., Mangelsdorf, M., Liu, J., Zhu, L., & Wu, J. Y. (2014). RNA-binding proteins in neurological diseases. *Science China Life Sciences*, 57(4), 432-444.

63. Kedersha, N., & Anderson, P. (2007). Mammalian stress granules and processing bodies. *Methods in enzymology*, 431, 61-81.

64. Min, H., Xu, L., Parrott, R., Overall, C. C., Lillich, M., Rabjohns, E. M., Rampersad, R. R., Tarrant, T. K., Meadows, N., & Fernandez-Castaneda, A. (2021). Mesenchymal stromal cells reprogram monocytes and macrophages with processing bodies. *Stem Cells*, 39(1), 115-128.

65. Hutchins, E. J., Piacentino, M. L., & Bronner, M. E. (2020). P-bodies are sites of rapid RNA decay during the neural crest epithelial-mesenchymal transition.

66. Fan, A. C., & Leung, A. K. (2016). RNA granules and diseases: a case study of stress granules in ALS and FTLD. In *RNA Processing* (pp. 263-296). Springer.

67. Anderson, P., Kedersha, N., & Ivanov, P. (2015). Stress granules, P-bodies and cancer. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, 1849(7), 861-870.

68. Turakhiya, A., Meyer, S. R., Marincola, G., Böhm, S., Vanselow, J. T., Schlosser, A., Hofmann, K., & Buchberger, A. (2018). ZFAND1 recruits p97 and the 26S proteasome to promote the clearance of arsenite-induced stress granules. *Molecular Cell*, 70(5), 906-919. e907.

69. Mateju, D., Eichenberger, B., Voigt, F., Eglinger, J., Roth, G., & Chao, J. A. (2020). Single-molecule imaging reveals translation of mRNAs localized to stress granules. *Cell*, 183(7), 1801-1812. e1813.

70. Mateju, D., & Chao, J. A. (2022). Stress granules: regulators or by-products? *The FEBS Journal*, 289(2), 363-373.

71. Buddika, K., Huang, Y.-T., Ariyapala, I. S., Butrum-Griffith, A., Norrell, S. A., O'Connor, A. M., Patel, V. K., Rector, S. A., Slovan, M., & Sokolowski, M. (2022). Coordinated repression of pro-differentiation genes via P-bodies and transcription maintains Drosophila intestinal stem cell identity. *Current Biology*, 32(2), 386-397. e386.

72. Ford, L., Ling, E., Kandel, E. R., & Fioriti, L. (2019). CPEB3 inhibits translation of mRNA targets by localizing them to P bodies. *Proceedings of the National Academy of Sciences*, 116(36), 18078-18087.

73. Robles-Luna, G., Furman, N., Barbarich, M. F., Carlotto, N., Attorresi, A., Garcia, M. L., & Kobayashi, K. (2020). Interplay between potato virus X and RNA granules in Nicotiana benthamiana. *Virus Res*, 276, 197823.

74. Moosa, M. M., & Banerjee, P. R. (2020). Subversion of host stress granules by coronaviruses: Potential roles of π -rich disordered domains of viral nucleocapsids. In *Journal of Medical Virology* (Vol. 92, pp. 2891-2893): John Wiley and Sons Inc.

75. Liu, D., Ndongwe, T. P., Puray-Chavez, M., Casey, M. C., Izumi, T., Pathak, V. K., Tedbury, P. R., & Sarafianos, S. G. (2020). Effect of P-body component Mov10 on HCV virus production and infectivity. *FASEB Journal*, 34(7), 9433-9449.

76. Liu, L., Weiss, E., Panas, M. D., Gotte, B., Sellberg, S., Thaa, B., & McInerney, G. M. (2019). RNA processing bodies are disassembled during Old World alphavirus infection. *J Gen Virol*, 100(10), 1375-1389.

77. Sharma, N. R., Majerciak, V., Kruhlak, M. J., & Zheng, Z. M. (2017). KSHV inhibits stress granule formation by viral ORF57 blocking PKR activation. *PLOS Pathogens*, 13(10), e1006677-e1006677.

78. Fan, S., Xu, Z., Liu, P., Qin, Y., & Chen, M. (2021). Enterovirus 71 2A Protease Inhibits P-Body Formation To Promote Viral RNA Synthesis. *JOURNAL OF VIROLOGY*, 95(19), e00922-00921.

79. Bonenfant, G., Williams, N., Netzband, R., Schwarz, M. C., Evans, M. J., & Pager, C. T. (2019). Zika virus subverts stress granules to promote and restrict viral gene expression. *JOURNAL OF VIROLOGY*, 93(12), e00520-00519.

80. Blázquez, A.-B., Martín-Acebes, M. A., Poderoso, T., & Saiz, J.-C. (2021). Relevance of oxidative stress in inhibition of eIF2 alpha phosphorylation and stress granules formation during Usutu virus infection. *PLOS Neglected Tropical Diseases*, 15(1), e0009072.

81. Arakawa, M., Tabata, K., Ishida, K., Kobayashi, M., Arai, A., Ishikawa, T., Suzuki, R., Takeuchi, H., Tripathi, L. P., & Mizuguchi, K. (2022). Flavivirus recruits the valosin-containing protein (VCP)/NPL4 complex to induce stress granule disassembly for efficient viral genome replication. *Journal of Biological Chemistry*, 101597.

82. Youn, J.-Y., Dyakov, B. J., Zhang, J., Knight, J. D., Vernon, R. M., Forman-Kay, J. D., & Gingras, A.-C. (2019). Properties of stress granule and P-body proteomes. *Molecular Cell*, 76(2), 286-294.

83. Eulalio, A., Behm-Ansmant, I., Schweizer, D., & Izaurralde, E. (2007). P-body formation is a consequence, not the cause, of RNA-mediated gene silencing. *Molecular and Cellular Biology*, 27(11), 3970-3981.

84. Campbell, A. M., De La Cruz Herrera, C. F., Marcon, E., Greenblatt, J., & Frappier, L. (2022). Epstein-Barr Virus BGLF2 commandeers RISC to interfere with cellular miRNA function. *PLOS Pathogens*, 18(1), e1010235.

85. Van Treeck, B., Protter, D. S., Matheny, T., Khong, A., Link, C. D., & Parker, R. (2018). RNA self-assembly contributes to stress granule formation and defining the stress granule transcriptome. *Proceedings of the National Academy of Sciences*, 115(11), 2734-2739.

86. Fu, Y., & Zhuang, X. (2020). m6A-binding YTHDF proteins promote stress granule formation. *Nature chemical biology*, 16(9), 955-963.

87. Chen, Y., & Cohen, T. J. (2019). Aggregation of the nucleic acid-binding protein TDP-43 occurs via distinct routes that are coordinated with stress granule formation. *Journal of Biological Chemistry*, 294(10), 3696-3706.

88. Park, Y., Park, J., Hwang, H. J., Kim, B., Jeong, K., Chang, J., Lee, J.-B., & Kim, Y. K. (2020). Nonsense-mediated mRNA decay factor UPF1 promotes aggresome formation. *Nature Communications*, 11(1), 1-15.

89. Pabis, M., Popowicz, G. M., Stehle, R., Fernández-Ramos, D., Asami, S., Warner, L., García-Mauriño, S. M., Schlundt, A., Martínez-Chantar, M. L., & Díaz-Moreno, I. (2019). HuR biological function involves RRM3-mediated dimerization and RNA binding by all three RRMs. *Nucleic Acids Research*, 47(2), 1011-1029.

90. Kenny, P. J., Kim, M., Skariah, G., Nielsen, J., Lannom, M. C., & Ceman, S. (2020). The FMRP-MOV10 complex: a translational regulatory switch modulated by G-Quadruplexes. *Nucleic Acids Research*, 48(2), 862-878.

91. Su, F., Liu, X., & Jiang, Y. (2020). Roles of MOV10 in Animal RNA Virus Infection. In *Frontiers in Veterinary Science* (Vol. 7, pp. 569737-569737): Frontiers Media S.A.

92. Sharma, N. R., Majerciak, V., Kruhlak, M. J., Yu, L., Kang, J. G., Yang, A., Gu, S., Fritzler, M. J., & Zheng, Z. M. (2019). KSHV RNA-binding protein ORF57 inhibits P-body formation to promote viral multiplication by interaction with Ago2 and GW182. *Nucleic Acids Research*, 47(17), 9368-9385.

93. Castle, E. L., Robinson, C.-A., Douglas, P., Rinker, K. D., & Corcoran, J. A. (2021). Viral manipulation of mechanoresponsive signaling disassembles processing bodies. *bioRxiv*, 2020.2005.2015.091876-092020.091805.091815.091876.

94. Deater, M., Tamhankar, M., & Lloyd, R. E. (2022). TDRD3 is an antiviral restriction factor that promotes IFN signaling with G3BP1. *PLOS Pathogens*, 18(1), e1010249.

95. An, H., Litscher, G., Watanabe, N., Wei, W., Hashimoto, T., Iwatsubo, T., Buchman, V. L., & Shelkovnikova, T. A. (2022). ALS-linked cytoplasmic FUS assemblies are compositionally different from physiological stress granules and sequester hnRNPA3, a novel modifier of FUS toxicity. *Neurobiology of disease*, 162, 105585.

96. Zimmer, M. M., Kibe, A., Rand, U., Pekarek, L., Ye, L., Buck, S., Smyth, R. P., Cicin-Sain, L., & Caliskan, N. (2021). The short isoform of the host antiviral protein ZAP acts as an inhibitor of SARS-CoV-2 programmed ribosomal frameshifting. *Nature Communications*, 12(1), 1-15.

97. Zhou, D., Jia, F., Li, Q., Zhang, L., Chen, Z., Zhao, Z., Cui, M., Song, Y., Chen, H., & Cao, S. (2018). Japanese encephalitis virus NS1' protein antagonizes interferon beta production. *Virologica Sinica*, 33(6), 515-523.

98. Halma, M. T., Ritchie, D. B., Cappellano, T. R., Neupane, K., & Woodside, M. T. (2019). Complex dynamics under tension in a high-efficiency frameshift stimulatory structure. *Proceedings of the National Academy of Sciences*, 116(39), 19500-19505.

99. Benner, B. E., Bruce, J. W., Kentala, J. R., Murray, M., Becker, J. T., Garcia-Miranda, P., Ahlquist, P., Butcher, S. E., & Sherer, N. M. (2021). Perturbing HIV-1 Ribosomal Frameshifting Frequency Reveals a cis Preference for Gag-Pol Incorporation into Assembling Virions. *JOURNAL OF VIROLOGY*, 96(1), e01349-01321.

100. Anokhina, V. S., & Miller, B. L. (2021). Targeting Ribosomal Frameshifting as an Antiviral Strategy: From HIV-1 to SARS-CoV-2. *Accounts of Chemical Research*, 54(17), 3349-3361.

101. Ritchie, D. B., Cappellano, T. R., Tittle, C., Rezajooei, N., Rouleau, L., Sikkema, W. K., & Woodside, M. T. (2017). Conformational dynamics of the frameshift stimulatory structure in HIV-1. *RNA*, 23(9), 1376-1384.

102. Dulude, D., Berchiche, Y. A., Gendron, K., Brakier-Gingras, L., & Heveker, N. (2006). Decreasing the frameshift efficiency translates into an equivalent reduction of the replication of the human immunodeficiency virus type 1. *Virology*, 345(1), 127-136.

103. Brakier-Gingras, L., Charbonneau, J., & Butcher, S. E. (2012). Targeting frameshifting in the human immunodeficiency virus. *Expert opinion on therapeutic targets*, 16(3), 249-258.

104. Clark, M. B., Janicke, M., Gottesbuhren, U., Kleffmann, T., Legge, M., Poole, E. S., & Tate, W. P. (2007). Mammalian gene PEG10 expresses two reading frames by high efficiency-1 frameshifting in embryonic-associated tissues. *Journal of Biological Chemistry*, 282(52), 37359-37369.

105. Belew, A. T., Meskauskas, A., Musalgaonkar, S., Advani, V. M., Sulima, S. O., Kasprzak, W. K., Shapiro, B. A., & Dinman, J. D. (2014). Ribosomal frameshifting in the CCR5 mRNA is regulated by miRNAs and the NMD pathway. *Nature*, 512(7514), 265-269.

106. Chiabudini, M., Tais, A., Zhang, Y., Hayashi, S., Wölfle, T., Fitzke, E., & Rospert, S. (2014). Release factor eRF3 mediates premature translation termination on polylysine-stalled ribosomes in *Saccharomyces cerevisiae*. *Molecular and Cellular Biology*, 34(21), 4062-4076.