

Article

DNA Identification and Diversity of The Vector Mosquitoes *Culex Pipiens* S.S. and *Culex Torrentium* in Belgium (Diptera: Culicidae)

Ann Vanderheyden ^{1*}, Nathalie Smitz ^{2*}, Katrien De Wolf ^{3,4}, Isra Deblauwe ³, Wouter Dekoninck ⁵, Kenny Meganck ², Sophie Gombeer ¹, Adwine Vanslembrouck ^{3,5}, Jacobus De Witte ³, Anna Schneider ³, Ingrid Verlé ³, Marc De Meyer ², Thierry Backeljau ^{1,6}, Ruth Müller ³ and Wim Van Bortel ^{3,7}

¹ Royal Belgian Institute of Natural Sciences (Barcode Facility for Organisms and Tissues of Policy Concern), Vautierstraat 29, 1000 Brussels, Belgium; avanderheyden@naturalsciences.be; sgombeer@naturalsciences.be

² Royal Museum for Central Africa (Barcode Facility for Organisms and Tissues of Policy Concern), Leuvensesteenweg 17, 3080 Tervuren, Belgium; nathalie.smitz@africamuseum.be; kenny.meganck@africamuseum.be; marc.de.meyer@africamuseum.be

³ The Unit of Entomology, Dept. Biomedical Sciences, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerp, Belgium; ideblauwe@itg.be; avanslembrouck@itg.be; jdewitte@itg.be; aschneider@itg.be; ingridverle@hotmail.com; rmuller@itg.be

⁴ Terrestrial Ecology Unit, Dept. of Biology, Ghent University, K.L. Ledeganckstraat 35, 9000 Ghent, Belgium; katrien.dewolf@ugent.be

⁵ Royal Belgian Institute of Natural Sciences (Scientific Heritage Service), Vautierstraat 29, 1000 Brussels, Belgium; wdekoninck@naturalsciences.be

⁶ Evolutionary Ecology Group, University of Antwerp, Universiteitsplein 1, 2610 Antwerp, Belgium; tbackeljau@naturalsciences.be

⁷ Outbreak Research team, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerp, Belgium; wvanbortel@itg.be

* Correspondence: AV: avanderheyden@naturalsciences.be; Tel.: +32 (0)2 627 41 23; NS: nathalie.smitz@africamuseum.be; Tel.: +32 (0)2 769 58 55

Abstract: This survey reports on the DNA identification and occurrence of *Culex torrentium* and *Cx. pipiens* s.s. in Belgium. These native disease vector mosquito species are morphologically difficult to separate and the biotypes of *Cx. pipiens* s.s. are morphologically indistinguishable. *Culex torrentium* and *Cx. pipiens* s.s. were identified using the COI and ACE2 loci. We recorded 1,248 *Cx. pipiens* s.s. and 401 *Cx. torrentium* specimens from 24 locations in Belgium (collected between 2017 and 2019). *Culex pipiens* biotypes *pipiens* and *molestus*, and their hybrids, were differentiated by fragment size analysis of the CQ11 locus (956 *pipiens* and 227 *molestus* biotype specimens, 29 hybrids). Hybrids were observed at 13 out of 16 sympatric sites. These results confirm that both species are widespread in Belgium, but while *Cx. torrentium* revealed many COI haplotypes, *Cx. pipiens* s.s. showed only one abundant haplotype. This latter observation may either reflect a recent population-wide demographic or range expansion, or a recent bottleneck, possibly linked to a *Wolbachia* infection. Finally, new evidence is provided for the asymmetric but limited introgression of the *molestus* biotype into the *pipiens* biotype.

Keywords: *Culex pipiens* biotypes *pipiens* / *molestus*; hybrids; disease vectors; DNA-based identification; cytochrome c oxidase I (COI); fragment size analyses (ACE2, CQ11)

1. Introduction

Culex pipiens s.l. is a complex of three species, viz. *Cx. australicus* Dobrotworsky & Drummond, 1953, *Cx. pipiens* s.s. Linnaeus, 1758 and *Cx. quinquefasciatus* Say, 1823. The latter species is common in (sub)tropical regions with no known established populations

in Europe [1], but has been introduced in the Netherlands with airplane traffic [2]. In contrast, *Culex australicus* is endemic to Australia. In Europe, hybrids between *Cx. quinquefasciatus* and *Cx. pipiens* s.s. were characterized in Southern coastal regions and the Mediterranean region by applying DNA methods [3]. Despite morphological similarities with *Cx. pipiens* s.s. [4], *Culex torrentium* Martini, 1925 is no longer considered as belonging to the *Cx. pipiens* species complex [5,6]. Yet, both species occur throughout Europe, with *Cx. torrentium* being more common in northern Europe and at high elevations further south, whereas *Cx. pipiens* s.s. is more common in the south, but the exact species distribution limits are still unclear [7]. The two species occur in sympatry and are native in Belgium, where *Cx. pipiens* s.s. appears to be more abundant and widespread [8–13]. Within *Cx. pipiens* s.s., two biotypes are recognized, viz. *Cx. pipiens* biotype *pipiens* Linnaeus, 1758 and *Cx. pipiens* biotype *molestus* Forskal, 1775 [14].

The identification of *Cx. torrentium* and *Cx. pipiens* s.s. is difficult as the two species differ by a few subtle morphological characters only [4,15]. The biotypes of *Cx. pipiens* s.s. are morphologically indistinguishable [14], but show four key behavioural differences [16–19]. Females of *Cx. pipiens* biotype *pipiens* need a bloodmeal to produce their first batch of viable eggs, prefer feeding on birds, breed in open spaces, and overwinter in a state of diapause. In contrast, females of *Cx. pipiens* biotype *molestus* can produce a first batch of viable eggs without a bloodmeal, prefer feeding on mammals, can breed in confined mating spaces, and do not overwinter in a state of diapause. In temperate regions of Europe, including Belgium, both biotypes co-occur in open aboveground spaces, but *Cx. p. biotype molestus* has a preference for confined spaces like cellars, cesspits, man-made basements or subways, where these mosquitoes mate and remain active throughout the year [20,21]. Hybrids between biotypes have been reported [13,19,22], displaying a combination of the behavioural traits of both biotypes [20,23]. However, since hybrids are less frequent than expected under random mating, the biotypes may show some degree of reproductive isolation [24,25]. Across the Mediterranean basin, populations in open spaces are genetically more homogenous with individuals displaying mixed biotype ancestry and a mix of the four key behavioural traits [20]. Thus, the genetic differentiation between biotypes decreases gradually from north to south across the western Palearctic. This may be linked to winters being less severe, allowing the non-diapausing *molestus* biotype to survive in open space environments and admix [20].

In Europe, *Culex pipiens* s.s. is the principal vector for West Nile Virus (WNV), and several other arboviruses [26]. *Culex pipiens* s.s. biotype hybrids with an opportunistic feeding behaviour seem to transmit WNV between birds and humans easier compared to non-hybrid *Cx. pipiens* biotypes [23,27–29]. *Culex torrentium* in turn, is an important vector for Sindbis virus (SINV) in Sweden [30]. However, *Cx. torrentium* also has a high potential to transmit WNV [31,32]. In view of the recent outbreaks of WNV infections in Germany and the Netherlands [33], it is important to closely monitor competent *Culex* vectors.

The distinction between *Cx. torrentium*, *Cx. pipiens* s.s., and *Cx. pipiens* biotypes, has not been investigated systematically. Hence, the distribution and identity of these taxa is still poorly known in most European countries [34]. Yet, this information is essential to establish reliable risk projections and control programmes, particularly for the early detection of WNV vectors and their potential spreading in Europe [35,36]. Therefore, the present paper reports on the DNA-based identification, occurrence, and diversity of *Cx. torrentium* and *Cx. pipiens* s.s., as well as *Cx. pipiens* biotypes *molestus* and *pipiens* and their hybrids, in Belgium.

2. Materials and Methods

2.1. Sampling

Adult and larval mosquitoes were collected in 2017 (August – November), 2018 and 2019 (both April – November) in the framework of the MEMO project (Monitoring of Exotic MOsquito species in Belgium [11]). Thirty-one potential points of entry (PoEs) for exotic mosquitoes in Belgium were surveyed using different sampling and trapping methods [11]. The PoEs included ports and airports, used tire and lucky bamboo import companies, parking lots along highways, whole sale markets, a flower auction, an allotment garden, an industrial area and cemeteries along the border with Germany. Specimens were morphologically identified as *Cx. pipiens* s.l. / *Cx. torrentium* using the keys of [4,37]. A random subset of 1,689 *Cx. pipiens* s.l. / *Cx. torrentium* specimens were selected for DNA-based identifications, using the sample_frac function of the dplyr package in R v4.03 [38].

2.2. DNA extraction and COI amplification

DNA was extracted from legs (adults) or abdomen (larvae) using the NucleoSpin® Tissue DNA extraction kit (Macherey-Nagel), following the manufacturer's protocol, but with an elution volume of 70 µl. Remaining parts of the specimens and dried DNA extracts are stored at the Royal Belgian Institute of Natural Sciences (Collection Identifier: IG34179). The universal primers LCO1490 and HCO2198 [39] were used to amplify the mitochondrial cytochrome c oxidase subunit I (COI) barcode region (658 bp) [40]. If this was unsuccessful, the C1N-2191 and C1J-1718 primer combination [41] was used to amplify a 472 bp fragment of the COI barcode region. All PCR mixtures, cycling conditions, purification and sequencing details are as described by [42] (Table S1). Raw sequences were trimmed, corrected, translated into amino acids and assembled using Geneious® v.10.0.4 (Biomatters Ltd.). A consensus sequence was generated for each specimen.

2.3. Fragment size analyses

To distinguish between *Cx. pipiens* s.s. and *Cx. torrentium*, a fragment of the Acetylcholinesterase-2 locus (ACE2) was amplified in a 10 µl PCR reaction volume as described by [43] (Table S1). This method also allows to detect the eventual presence of introduced exotic *Cx. quinquefasciatus*. Using the forward primer B1246s, and the reverse primers ACEpip, ACEquin and ACEtorr species-specific fragment sizes were produced, viz. 610 bp for *Cx. pipiens* s.s., 416 bp for *Cx. torrentium*, and 274 bp for *Cx. quinquefasciatus* [43]. PCR products were checked on a 2.5 % agarose gel (45 min; 90 V).

To identify the two *Cx. pipiens* s.s. biotypes and their hybrids, the CQ11 microsatellite locus was amplified using the forward primer CQ11F2 and the reverse primers pipCQ11R and molCQ11R, following [44] (Table S1). PCR products were checked on a 2.5 % agarose gel (45 min; 90V), with a band at 200 bp for *Cx. p.* biotype *pipiens* and at 250 bp for *Cx. p.* biotype *molestus*. Hybrids show both bands. Such hybrids were subsequently re-extracted and re-amplified for the CQ11 locus to exclude possible DNA contamination and confirm their status by visualisation of the two bands.

2.4. COI data analyses

The species identification engine of BOLD was used (www.boldsystems.org) with the species level barcode records option to find the closest matching reference sequence. A Neighbour-Joining (NJ) tree was constructed to examine the clustering support of each *Culex* species occurring in Belgium [12,45] (Geneious® v10.0.4., Tamura-Nei distance model, 1,000 bootstrap replicates). To do so, all publicly available COI sequences (<http://www.boldsystems.org/index.php/databases>, March 2020) for these *Culex* species were aligned using ClustalW in Geneious® v10.0.4 (Biomatters Ltd.) with the newly generated COI sequences in this study. COI sequences of four species of the genus *Coquillettidia* Dyar, 1905 were included as outgroup (GenBank accession numbers: GQ165785, GQ165801, GQ165802 and GQ165803). The alignment was checked for stop codons and

trimmed to retain 658 bp. Sequences of less than 300 bp and conspecific identical sequences were discarded.

Pairwise differences in COI nucleotide frequencies between species, biotypes and biotype hybrids, were evaluated using Wright's F -statistics in Arlequin v3.5 (1,000 random permutations for significance, with subsequent standard Bonferroni correction) [46]. Haplotype frequencies, mean number of pairwise nucleotide differences (k) and nucleotide diversity (Pi) were also estimated with Arlequin v3.5, excluding sequences with ambiguous sites.

2.5. Habitat characterization: Land Cover Classes

The percentage of Corine Land Cover (CLC) classes (© European Union, Copernicus Land Monitoring Service 2021, European Environment Agency (EEA)) was calculated in a 2.5 km buffer zone around each sampling locations. The latest raster file (CLC 2018) was used and calculation was done in Q-GIS and R v4.03. The levels were grouped into five main CLC classes *i.e.*, artificial or urban areas, agricultural areas, forest and seminatural areas, wetlands, and water bodies.

3. Results

In total, 34,401 specimens from 27 out of 31 PoEs were morphologically identified as *Cx. pipiens* s.l. / *Cx. torrentium* of which 1,689 specimens from 24 sites were selected for DNA-based verification. The four PoEs where these species were not collected were only surveyed using oviposition traps. Based on the BOLD similarity percentages, the COI NJ-tree (Figure 1), and the ACE2 fragment sizes (Figure S1), 401 specimens were identified as *Cx. torrentium*, and 1,248 as *Cx. pipiens* s.s.. Thirty seven specimens did not provide ACE2 results and were therefore considered as *Cx. pipiens* s.l. / *Cx. torrentium*. Three sequences were of too low quality for identification. The ACE2 fragment size analysis provided no evidence of *Cx. quinquefasciatus*. In the NJ-tree, *Cx. torrentium* forms a cluster with 74.9 % bootstrap support inside the *Cx. pipiens* s.s. / *Cx. torrentium* group (Figure 1). The identified *Cx. torrentium* cluster includes all generated and downloaded (BOLD) COI sequences.




Figure 1. Neighbour-Joining tree based on the COI sequences available on BOLD / GenBank for *Culex* species present in Belgium [12,45] including the sequences generated in this study, and identified based on the ACE2 fragment size analysis. Bootstrap values are indicated at the branches. Sequences were collapsed in species-clusters. N = total number of unique sequences included, following Geneious® v10.0.4.

Based on the CQ11 fragment size analysis 956 specimens were assigned to *Cx. p.* biotype *pipiens* and 227 specimens to *Cx. p.* biotype *modestus* (Figure S2). Additionally, 29 specimens were identified as hybrids between both biotypes, while the biotypes of 36 *Cx. pipiens* s.s. sequences were not determined due to missing CQ11 results. These sequences, together with those identified as *Cx. pipiens* s.l. / *Cx. torrentium* (N = 37), were excluded from further analyses. The abundance of each taxon at each sampling site is shown in Figure 2. In most sites where both biotypes co-occur (N = 16), crossbreeding was identified, with hybrids detected at 13 sampling locations (Figure 2, Table S2). Sites where biotypes co-occurred included environments dominated by urban (Kallo, Charleroi, Zeebrugge and Zaventem), agricultural (Villers-Le-Bouillet, Vrasene, Frameries, Aische-en-Refail, Büllingen and Natoye), and forest and seminatural (Eupen, Dilsen-Stokken, Houyet and Maasmechelen) areas (Figure 3). At these sites, larval stages of both biotypes were collected in the same type of larval habitats on the same days, viz. used tires, drainage holes, plastic containers and cemetery flower vases, on multiple occasions. They were also found once together in a large artificial pond. *Culex pipiens* s.s. and *Cx. torrentium* were collected on the same days in the following same type of larval habitats: used tires, drainage holes, cemetery flower vases, plastic sheets, metal and plastic containers. The new COI sequences were deposited in GenBank (accession numbers: *Cx. torrentium*: OM749168-OM749568; *Cx. p.* biotype *pipiens*: OM748139-OM749094; *Cx. p.* biotype *modestus*: OM747912-OM748138; *Cx. p.* biotype *pipiens* X *Cx. p.* biotype *modestus*: OM747883-OM747911; *Cx. pipiens* s.s.: OM749132-OM749167; *Cx. pipiens* s.l. / *Cx. torrentium*: OM749095-OM749131).




Figure 2. Distribution of the specimens identified using DNA-based techniques and collected during the MEMO survey 2017-2019 [11]. Hybrids = *Cx. p. biotype pipiens* X *Cx. p. biotype molestus*. The numbers indicated on the pie charts are the collection site numbers used in Table S2.




Figure 3. The percentage of main Corine Land Cover classes in a 2.5 km buffer zone around the sampling locations (group levels based on five classes).

Pairwise F_{ST} values between *Cx. p. biotype pipiens* and *Cx. p. biotype molestus* were significantly different from zero, which was also the case between the latter and the hybrids (Table 1). Hybrids and *Cx. p. biotype pipiens* showed the smallest average pairwise nucleotide differences and lowest average nucleotide diversities (Table 2), despite *Cx. p. biotype pipiens* was the most widespread taxon in this survey (Figure 2; Table S4). *Culex torrentium* showed higher average pairwise nucleotide differences and nucleotide diversities than *Cx. p. biotype pipiens* (Table 2).




Table 1. Pairwise F_{ST} estimates between biotypes and biotype hybrids of *Culex pipiens* s.s. based on COI sequences, calculated using Arlequin v3.5. Hybrids = *Cx. p. biotype molestus* X *Cx. p. biotype pipiens*. Significant values after standard Bonferroni correction marked by an asterisk ($p < 0.0005$).

<i>F_{ST}</i>	Hybrids	<i>Cx. p. biotype molestus</i>	<i>Cx. p. biotype pipiens</i>
Hybrids	0	-	-
<i>Cx. p. biotype molestus</i>	0.2172*	0	-
<i>Cx. p. biotype pipiens</i>	0.0164	0.4737*	0

Table 2. COI sequence diversity of *Culex pipiens* s.s. biotypes and *Cx. torrentium*, calculated using Arlequin v3.5. Hybrids = *Cx. p. biotype molestus* × *Cx. p. biotype pipiens*, N_{COI} = number of new COI sequences, N_{Hap} = number of haplotypes, N_{UnHap} = number of unique haplotypes, S = number of polymorphic sites, k = average pairwise nucleotide differences, Pi = nucleotide diversity.

	Hybrids	<i>Cx. p. biotype molestus</i>	<i>Cx. p. biotype pipiens</i>	<i>Cx. torrentium</i>
N_{COI}	29	227	956	401
N_{Hap}	3	6	24	50
N_{UnHap}	1	3	20	50
S	3	8	19	31
k	0.2069 ± 0.2606	0.5458 ± 0.4522	0.0703 ± 0.1418	0.8990 ± 0.6275
Pi	0.0006 ± 0.0008	0.0014 ± 0.0012	0.0002 ± 0.0004	0.0024 ± 0.0018

The most common COI haplotypes in *Cx. pipiens* s.s. were H1 (698 out of 1,248 sequences, including 19 hybrids, 91 *Cx. p. biotype molestus* and 588 *Cx. p. biotype pipiens*) and H2 (84 out of 1,248 sequences, including 82 *Cx. p. biotype molestus* and two *Cx. p. biotype pipiens*) (Figure 4). The most common COI haplotype in *Cx. torrentium* was H3 (125 out of 401 sequences) (Figure 4). The haplotype of 510 sequences could not be identified because of ambiguous sites or short fragment lengths ($N_{Cx. torrentium} = 123$; $N_{Cx. p. biotype pipiens} = 330$; $N_{Cx. p. biotype molestus} = 49$; $N_{hybrids} = 8$).

*Culex pipiens* biotype *pipiens**Culex pipiens* biotype *molestus*

Hybrids


*Culex torrentium*

Figure 4. COI haplotype frequencies calculated with Arlequin v.3.5. Hybrids = *Culex pipiens* biotype *pipiens* X *Cx. p.* biotype *molestus*, blue = *Cx. pipiens* s.s. haplotype H1, orange = *Cx. pipiens* s.s. haplotype H2, green = *Cx. torrentium* haplotype H3, yellow = sequences to which no haplotypes could be assigned, grey = haplotypes with a frequency < 6 %.

4. Discussion

4.1. Species occurrence in Belgium

In line with previous studies [8,9,13], *Cx. pipiens* s.s. appears to be more common in Belgium than *Cx. torrentium*. Likewise, *Culex p.* biotype *pipiens* is more common and widespread in Belgium than *Cx. p.* biotype *molestus* as the latter comprises only 13 % of the specimens (Table S4). Yet, since industrial areas were overrepresented in this survey for

the early detection of exotic *Aedes* species [11], the sampling may have been biased. As such, the *molestus* biotype with its preference for hypogean habitats [21] and highly eutrophic waters in confined mating spaces [20,21], may be underrepresented as these environments were not surveyed during the MEMO project [11]. Thus, more targeted surveys are needed to determine the actual prevalence and distribution of *Cx. p. biotype molestus* in Belgium. Nevertheless, this study confirms the co-occurrence of both biotypes in urban, agricultural, and forest and seminatural habitats [13,47–50] (Figure 3). Despite the sympatric occurrence of both biotypes, only few hybrid specimens were found (1.7 %; Table S2-S4), *i.e.* less than in Germany (4.2 %) [19], Portugal (8–10 %) [47], and Italy (14.4 %) [49]. These low hybridisation rates suggest at least partial reproductive / ecological isolation between biotypes [20,47], with some rare haplotypes identified as specific to one biotype (Table 2, see next section). The sympatric co-occurrence of both biotypes and their hybrids with their opportunistic feeding behaviour [47,51] hints at the potential danger of viral transmissions from birds to humans (*i.e.* to act as bridge disease vectors). However, the low frequency of hybrids likely limits their potential epidemiological role in WNV outbreaks.

While *Cx. pipiens* s.s. and *Cx. torrentium* are sympatric in some areas, the latter species was not collected in the north of Belgium (Figure 2). Again, this may be a sampling artefact, since the species was collected all over Belgium during the nationwide MODIRISK mosquito survey (2007–2010) [9,48]. Both species were equally observed in different habitats and have adapted to a life in human neighbourhoods [52], with *Cx. torrentium* and *Cx. pipiens* s.s. larvae often found in small artificial and nutrient-rich bodies of water [7,52]. Thus, both species are widespread in Belgium, but their exact distribution limits in Europe remain to be determined.

4.2. COI haplotype composition and genetic variability

Belgian *Cx. torrentium* showed a higher COI variability than *Cx. pipiens* s.s., which is in line with [52,53]. Yet, we found no evidence of a further sub-structuring or taxonomic differentiation within *Cx. torrentium*. The limited intraspecific variation within *Cx. pipiens* s.s. is consistent with [54,55]. Hence, COI haplotype H1 had a prevalence of 61.51 % in *Cx. p. biotype pipiens* and of 40.09 % in *Cx. p. biotype molestus*. Similar prevalences were reported by [52] (H1 = haplotype 1), [3] (H1 = haplotype A/C) and [56] (H1 = haplotype H). As such, haplotype H1 is widespread and most frequent in northern temperate climates (Germany, Japan, North America and Russia) [3,52,56]. Situations in which populations show limited genetic variation and consist of a highly frequent haplotype jointly with a few rare haplotypes, can be explained by either a recent population-wide demographic or range expansion, or a recent bottleneck, possibly in combination with a *Wolbachia* infection [52]. Such a *Wolbachia* infection can severely reduce mitochondrial diversity [52,57,58]. This might, in part, explain the limited COI diversity in *Cx. pipiens* s.s., which shows *Wolbachia* infection rates of > 90 % [52,59–61], whereas COI diversity might have been retained in *Cx. torrentium*, within which *Wolbachia* infections appear to be very rare [59,62].

In Belgium haplotype H2 was almost exclusively found in *Cx. p. biotype molestus*. This is somewhat unexpected as this haplotype is rarely found in temperate climates, but associated with (sub)tropical climates ([3] H2 = haplotype E/E1; [56] H2 = haplotype C). Elsewhere, haplotype H2 is prevalent in *Cx. quinquefasciatus* (42 %) and its hybrids with *Cx. pipiens* s.s. (32 %) [3]. Hence, COI haplotypes in *Cx. pipiens* s.l. are not species-specific [56].

Currently, the biotypes *pipiens* and *molestus* of *Cx. pipiens* s.s. are regarded as different monophyletic evolutionary units undergoing incipient ecological speciation, so that they

may be distinct phylogenetic entities [22,27,47,63–65]. This was supported by the significant F_{ST} estimates found in the present study. The different mating behaviour of both biotypes was considered as an initial factor of a sympatric speciation process [47]. The limited level of hybridisation is not bidirectional, with a mainly male-mediated introgression from *molestus* to the *pipiens* biotype [47], which explains the prevalence of H1 (typical of *pipiens*) and absence of H2 (typical of *molestus*) in hybrids. This asymmetric introgression may reflect a mating strategy, wherein stenogamous *molestus* males mate with both *molestus* and *pipiens* females in above-ground habitats, while *pipiens* males mate (specialised swarming behaviour) in open spaces and therefore have a higher disposition to mate with *pipiens* females [47]. An experimental study revealed at least one reproductive isolating mechanism, with females actively avoiding copulation with males of the other biotype, and *pipiens* females being unsuccessful in receiving *molestus* male's sperm [66].

Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1, Figure S1: Example of ACE2 size fragment analysis on a 2.5 % agarose gel. PCR multiplex including the primers B1246s (Forward), ACEpip (Reverse), ACEtorr (Reverse) and ACEquin (Reverse). Cycling conditions are provided in Table S1. P = *Cx. pipiens* s.s. (610 bp); T = *Cx. torrentium* (416 bp); Figure S2: Example of CQ11 size fragment analysis on a 2.5 % agarose gel. PCR multiplex including the primers CQ11F2 (Forward), molCQ11R (Reverse) and pipCQ11R (Reverse). Cycling conditions are provided in Table S1. PP = *Culex p. biotype pipiens* (200 bp); PM = *Culex p. biotype molestus* (250 bp), H = *Cx. p. biotype pipiens* X *Cx. p. biotype molestus* (200 bp and 250 bp); Table S1: Summary of PCR cycling conditions for the amplification of the COI, ACE2 and CQ11 loci; Table S2: Map codes, municipalities and coordinates of sampling localities with taxon occurrence; Table S3: Overview of the COI sequencing success per year. N = number of specimens.

Author Contributions: Conceptualization: N.S., I.D., K.D.W., W.D., R.M., W.V.B.; Methodology: N.S., K.D.W., I.D., A.V., J.D.W., A.S., I.V., W.V.B.; Formal Analysis & Software: N.S., A.V., K.M., S.G.; Data Curation: W.D., N.S., I.D.; Writing – Original Draft Preparation: A.V., N.S.; Writing – Review & Editing: I.D., K.D.W., W.D., K.M., S.G., A.V., J.D.W., A.S., I.V., M.D.M., T.B., R.M., W.V.B.; Visualization: A.V., N.S.; Supervision: I.D., W.V.B., R.M., T.B., M.D.M.; Project Administration and Funding Acquisition: I.D., W.V.B., R.M., T.B., M.D.M. All authors have read and agreed to the published version of the manuscript.

Funding: This work is part of the MEMO project, funded by the Flemish, Walloon and Brussels regional governments and the Federal Public Service (FPS) Public Health, Food Chain Safety and Environment in the context of the National Environment and Health Action Plan (NEHAP) (Belgium). The Barcoding Facility for Organisms and Tissues of Policy Concern (BopCo - <http://bopco.myspecies.info/>) is financed by the Belgian Science Policy Office (Belspo). The Outbreak Research Team of the Institute of Tropical Medicine is funded by the Department of Economy, Science and Innovation of the Flemish government.

Data Availability Statement: The data presented in this study are openly available in GenBank at <https://www.ncbi.nlm.nih.gov/genbank/>, reference number: OM747883-OM749568.

Acknowledgments: We would like to thank all cooperating companies for giving access to their private property during mosquito sampling surveys. We would like to thank the laboratory and technical staff at the Institute of Tropical Medicine Antwerp (ITM), at the Royal Belgian Institute of Natural Sciences (RBINS) and at the Royal Museum for Central Africa (RMCA).

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. ECDC. *Culex pipiens* factsheet. 2020. Available online: <https://www.ecdc.europa.eu/en/all-topics-z/disease-vectors/facts/mosquito-factsheets/culex-pipiens-factsheet-experts> (accessed on 16 Feb 2022)
2. Scholte, E., Ibáñez-Justicia, A., Stroo, A., Zeeuw, J.D., Hartog, W.D., Reusken, C. Mosquito collections on incoming intercontinental flights at Schiphol International airport, the Netherlands, 2010–2011. *J Eur Mosq Control Assoc* **2014**, *32*, 17–21.
3. Shaikevich, E.V., Vinogradova, E.B., Bouattour, A., Gouveia de Almeida, A.P. Genetic diversity of *Culex pipiens* mosquitoes in distinct populations from Europe: contribution of *Cx. quinquefasciatus* in Mediterranean populations. *Parasit Vectors* **2016**, *9*, 47.

4. Becker, N., Petric, D., Zgomba, M., Boase, C., Madon, M., Dahl, C., et al. *Mosquitoes and their control*; Springer: Heidelberg; New York, 2010.
5. Miller, B.R., Crabtree, M.B., Savage, H.M. Phylogeny of fourteen *Culex* mosquito species, including the *Culex pipiens* complex, inferred from the internal transcribed spacers of ribosomal DNA. *Insect Mol Biol* **1996**, *5*, 93–107.
6. Harbach, R.E. *Culex pipiens*: species versus species complex taxonomic history and perspective. *J Am Mosq Control Assoc* **2012**, *28*, 10–23.
7. Hesson, J.C., Rettich, F., Merdić, E., Vignjević, G., Östman, Ö., Schäfer, M., et al. The arbovirus vector *Culex torrentium* is more prevalent than *Culex pipiens* in northern and central Europe. *Med Vet Entomol* **2014**, *28*, 179–86.
8. Dekoninck, W., Hendrickx, F., Versteirt, V., Coosemans, M., Clercq, E.M.D., Hendrickx, G., et al. Changes in species richness and spatial distribution of mosquitoes (Diptera: Culicidae) inferred from museum specimen records and a recent inventory: a case study from Belgium suggests recent expanded distribution of arbovirus and malaria vectors. *J Med Entomol* **2013**, *50*, 237–43.
9. Versteirt, V., Boyer, S., Damiens, D., De Clercq, E., Dekoninck, W., Ducheyne, E., et al. Nationwide inventory of mosquito biodiversity (Diptera: Culicidae) in Belgium, Europe. *Bull Entomol Res* **2013**, *103*, 193–203.
10. Deblauwe, I., Sohier, C., Schaffner, F., Rakotoarivony, L.M., Coosemans, M. Implementation of surveillance of invasive mosquitoes in Belgium according to the ECDC guidelines. *Parasites and Vectors* **2014**, *7*, 1–11.
11. Deblauwe, I., De Wolf, K., Smitz, N., Vanslembrouck, A., Schneider, A., De Witte, J., et al. Monitoring of exotic mosquitoes in Belgium (MEMO): Final Report Phase 7 Part 1: MEMO results. Antwerp, Belgium, 2020.
12. Boukraa, S., Dekoninck, W., Versteirt, V., Schaffner, F., Coosemans, M., Haubrige, E., et al. Updated checklist of the mosquitoes (Diptera: Culicidae) of Belgium. *J Vector Ecol* **2015**, *40*, 398–407.
13. Boukraa, S., de La Grandiere, M.A., Bawin, T., Raharimalala, F.N., Zimmer, J.Y., Haubrige, E., et al. Diversity and ecology survey of mosquitoes potential vectors in Belgian equestrian farms: A threat prevention of mosquito-borne equine arboviruses. *Prev Vet Med* **2016**, *124*, 58–68.
14. Harbach, R.E., Harrison, B.A., Gad, A.M. *Culex (Culex) molestus* Forskal (Diptera: Culicidae): Neotype designation, description, variation, and taxonomic status. *Proc Entomol Soc Wash* **1984**, *86*, 521–42.
15. Razygraev, A.V., Sulesco, T.M. The Use of the Bayes Factor for identification of *Culex pipiens* and *C. torrentium* (Diptera: Culicidae) based on morphometric wing characters. *Entomol Rev* **2020**, *100*, 220–7.
16. Byrne, K., Nichols, R.A. *Culex pipiens* in London Underground tunnels: differentiation between surface and subterranean populations. *Heredity* **1999**, *82*, 7–15.
17. Lundström, J. Mosquito-borne viruses in western Europe: a review. *J Vector Ecol* **1999**, *24*, 1–39.
18. Vinogradova, E.B. *Culex pipiens pipiens* mosquitoes: taxonomy, distribution, ecology, physiology, genetics, applied importance and control. Pensoft. Sofia-Moscow, 2000.
19. Rudolf, M., Czajka, C., Börstler, J., Melaun, C., Jöst, H., von Thien, H., et al. First nationwide surveillance of *Culex pipiens* Complex and *Culex torrentium* mosquitoes demonstrated the presence of *Culex pipiens* biotype *pipiens/molestus* hybrids in Germany. *PLoS One* **2013**, *8*:e71832.
20. Haba, Y., McBride, L. Origin and status of *Culex pipiens* mosquito ecotypes. *Curr Biol* **2022**, *32*, R237–46.
21. Becker, N., Jöst, A., Weitzel, T. The *Culex pipiens* complex in Europe. *J Am Mosq Control Assoc* **2012**, *28*, 53–67.
22. Yurchenko, A.A., Masri, R.A., Khrabrova, N.V., Sibataev, A.K., Fritz, M.L., Sharakhova, M.V. Genomic differentiation and intercontinental population structure of mosquito vectors *Culex pipiens pipiens* and *Culex pipiens molestus*. *Sci Rep* **2020**, *10*, 7504.
23. Fritz, M.L., Walker, E.D., Miller, J.R., Severson, D.W., Dworkin, I. Divergent host preferences of above- and below-ground *Culex pipiens* mosquitoes and their hybrid offspring. *Med Vet Entomol* **2015**, *29*, 115–23.
24. Osório, H.C., Zé-Zé, L., Amaro, F., Nunes, A., Alves, M.J. Sympatric occurrence of *Culex pipiens* (Diptera, Culicidae) biotypes *pipiens*, *molestus* and their hybrids in Portugal, Western Europe: feeding patterns and habitat determinants. *Med Vet Entomol* **2014**, *28*, 103–9.
25. Gomes, B., Sousa, C.A., Vicente, J.L., Pinho, L., Calderón, I., Arez, E., et al. Feeding patterns of *molestus* and *pipiens* forms of *Culex pipiens* (Diptera: Culicidae) in a region of high hybridization. *Parasit Vectors* **2013**, *6*, 93.
26. Brugman, V.A., Hernández-Triana, L.M., Medlock, J.M., Fooks, A.R., Carpenter, S., Johnson, N. The role of *Culex pipiens* L. (Diptera: Culicidae) in virus transmission in Europe. *Int J Environ Res Public Health* **2018**, *15*, 389.
27. Fonseca, D.M., Keyghobadi, N., Malcolm, C.A., Mehmet, C., Schaffner, F., Mogi, M., et al. Emerging Vectors in the *Culex pipiens* Complex. *Science* **2004**, *303*, 1535–8.
28. Gomes, B., Kioulos, E., Papa, A., Almeida, A.P.G., Vontas, J., Pinto, J. Distribution and hybridization of *Culex pipiens* forms in Greece during the West Nile virus outbreak of 2010. *Infect Genet Evol* **2013**, *16*, 218–25.
29. Ciota, A.T., Chin, P.A., Kramer, L.D. The effect of hybridization of *Culex pipiens* complex mosquitoes on transmission of West Nile virus. *Parasites and Vectors*. *BioMed Central* **2013**, *6*, 1–4.
30. Hesson, J.C., Verner-Carlsson, J., Larsson, A., Ahmed, R., Lundkvist, Å., Lundström, J.O. *Culex torrentium* mosquito role as major enzootic vector defined by rate of Sindbis virus infection, Sweden, 2009. *Emerg Infect Dis* **2015**, *21*, 875.
31. Leggewie, M., Badusche, M., Rudolf, M., Jansen, S., Börstler, J., Krumkamp, R., et al. *Culex pipiens* and *Culex torrentium* populations from Central Europe are susceptible to West Nile virus infection. *One Heal* **2016**, *2*, 88–94.

32. Jansen, S., Heitmann, A., Lühken, R., Leggewie, M., Helms, M., Badusche, M., et al. *Culex torrentium*: A potent vector for the transmission of west nile virus in central Europe. *Viruses* **2019**, *11*, 1–11.
33. ECDC. Surveillance and disease data for West Nile virus infections. 2020. Available online: <https://www.ecdc.europa.eu/en/west-nile-fever/surveillance-and-disease-data> (accessed on 21 Feb 2022)
34. Weitzel, T., Braun, K., Collado, A., Jöst, A., Becker, N. Distribution and frequency of *Culex pipiens* and *Culex torrentium* (Culicidae) in Europe and diagnostic allozyme markers. *Eur Mosq Bull* **2011**, *29*, 22–37.
35. Calistri, P., Giovannini, A., Hubalek, Z., Ionescu, A., Monaco, F., Savini, G., et al. Epidemiology of West Nile in Europe and in the Mediterranean Basin. *Open Virol J* **2010**, *4*, 29.
36. García-Carrasco, J.M., Muñoz, A.R., Olivero, J., Segura, M., Real, R. Predicting the spatio-temporal spread of West Nile virus in Europe. *PLoS Negl Trop Dis* **2021**, *15*, e0009022.
37. Gunay, F., Picard, M., Robert, V. MosKeyTool, an interactive identification key for mosquitoes of Euro-Mediterranean. 2018. Available online: <http://medilabsecure.com/moskeytool> (accessed on 5 Aug 2020)
38. Wickham, H., François, R., Henry, L., Müller, K. A Grammar of Data Manipulation • dplyr 1.0.8. 2020. Available online: <https://dplyr.tidyverse.org/index.html> (accessed on 1 Apr 2022)
39. Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* **1994**, *3*, 294–9.
40. Hebert, P., Cywinska, A., Ball, S., DeWaard, J. Biological identifications through DNA barcodes. *Proc R Soc - Biol Sci* **2003**, *270*, 313–21.
41. Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H., Flook, P. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Entomol Soc Am* **1994**, *87*, 653–701.
42. Ibáñez-Justicia, A., Smitz, N., den Hartog, W., van de Vossenberg, B., De Wolf, K., Deblauwe, I., et al. Detection of exotic mosquito species (Diptera: Culicidae) at international airports in Europe. *Int J Environ Res Public Health* **2020**, *17*, 3450.
43. Smith, J.L., Fonseca, D.M. Rapid assays for identification of members of the *Culex (Culex) pipiens* complex, their hybrids, and other sibling species (Diptera: Culicidae). *Am J Trop Med Hyg* **2004**, *70*, 339–45.
44. Bahnck, C.M., Fonseca, D.M. Rapid assay to identify the two genetic forms of *Culex (Culex) pipiens* L. (Diptera : Culicidae) and hybrid populations. *Am J Trop Med Hyg* **2006**, *75*, 251–5.
45. De Wolf, K., Vanderheyden, A., Deblauwe, I., Smitz, N., Gombeer, S., Vanslembrouck, A., et al. First record of the West Nile virus bridge vector *Culex modestus* Ficalbi (Diptera: Culicidae) in Belgium, validated by DNA barcoding. *Zootaxa* **2021**, *4920*, 131–9.
46. Excoffier, L., Lischer, H.E.L. Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* **2010**, *10*, 564–7.
47. Gomes, B., Sousa, C.A., Novo, M.T., Freitas, F.B., Alves, R., Côrte-Real, A.R., et al. Asymmetric introgression between sympatric molestus and pipiens forms of *Culex pipiens* (Diptera: Culicidae) in the Comporta region, Portugal. *BMC Evol Biol* **2009**, *9*, 1–15.
48. Versteirt, V., De Clercq, E., Dekoninck, W., Damiens, D., Ayrinhac, A., Jacobs, F., et al. MODIRISK: Mosquito vectors of disease: spatial biodiversity, drivers of change, and risk. Antwerp, Belgium; 2011. Available online: [https://www.belspo.be/belspo/SSD/science/Reports/MODIRISK_Phase 1 Summary.pdf](https://www.belspo.be/belspo/SSD/science/Reports/MODIRISK_Phase%201%20Summary.pdf)
49. Di Luca, M., Toma, L., Boccolini, D., Severini, F., La Rosa, G., Minelli, G., et al. Ecological Distribution and CQ11 Genetic Structure of *Culex pipiens* Complex (Diptera: Culicidae) in Italy. *PLoS One* **2016**, *11*, e0146476.
50. Zittra, C., Flechl, E., Kothmayer, M., Vitecek, S., Rossiter, H., Zechmeister, T., et al. Ecological characterization and molecular differentiation of *Culex pipiens* complex taxa and *Culex torrentium* in eastern Austria. *Parasit Vectors* **2016**, *9*, 197.
51. Huang, S., Hamer, G.L., Molaei, G., Walker, E.D., Goldberg, T.L., Kitron, U.D., et al. Genetic variation associated with mammalian feeding in *Culex pipiens* from a West Nile virus epidemic region in Chicago, Illinois. *Vector Borne Zoonotic Dis* **2009**, *9*, 637–42.
52. Werblow, A., Klimpel, S., Bolius, S., Dorresteijn, A.W.C., Sauer, J., Melaun, C. Population structure and distribution patterns of the sibling mosquito species *Culex pipiens* and *Culex torrentium* (Diptera: Culicidae) reveal different evolutionary paths. *PLoS One* **2014**, *9*, e102158.
53. Werblow, A., Bolius, S., Dorresteijn, A.W.C., Melaun, C., Klimpel, S. Diversity of *Culex torrentium* Martini, 1925 - A potential vector of arboviruses and filaria in Europe. *Parasitol Res* **2013**, *112*, 2495–501.
54. Guillemaud, T., Pasteur, N., Rousset, F. Contrasting levels of variability between cytoplasmic genomes and incompatibility types in the mosquito *Culex pipiens*. *Proc R Soc London Ser B Biol Sci* **1997**, *264*, 245–51.
55. Fedorova, M.V., Shaikevich, E.V. Morphological and molecular-genetic distinctions between adult mosquitoes *Culex torrentium* Martini and *C. pipiens* Linnaeus (Diptera, Culicidae) from Moscow Province. *Entomol Rev* **2007**, *87*, 127–35.
56. Koosha, M., Oshaghi, M.A., Sedaghat, M.M., Vatandoost, H., Azari-Hamidian, S., Abai, M.R., et al. Sequence analysis of mtDNA COI barcode region revealed three haplotypes within *Culex pipiens* assemblage. *Exp Parasitol* **2017**, *181*, 102–10.
57. Cariou, M., Duret, L., Charlat, S. The global impact of *Wolbachia* on mitochondrial diversity and evolution. *J Evol Biol* **2017**, *30*, 2204–10.
58. Hurst, G.D.D., Jiggins, F.M. Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. *Proc R Soc B* **2005**, *272*, 1525–34.
59. Bergman, A., Hesson, J.C. *Wolbachia* prevalence in the vector species *Culex pipiens* and *Culex torrentium* in a Sindbis virus-endemic

region of Sweden. *Parasites and Vectors* **2021**, *14*, 1–7.

60. Duron, O., Lagnel, J., Raymond, M., Bourtzis, K., Fort, P., Weill, M. Transposable element polymorphism of *Wolbachia* in the mosquito *Culex pipiens*: Evidence of genetic diversity, superinfection and recombination. *Mol Ecol* **2005**, *14*, 1561–73.
61. Vinogradova, E.B., Fedorova, M.V., Shaikevich, E.V. Endosymbiotic bacterium *Wolbachia pipiensis* in synanthropic populations of the mosquito *Culex pipiens pipiens* L. (Diptera, Culicidae). *Dokl Biol Sci* **2003**, *389*, 172–5.
62. Leggewie, M., Krumkamp, R., Badusche, M., Heitmann, A., Jansen, S., Schmidt-Chanasit, J., et al. *Culex torrentium* mosquitoes from Germany are negative for *Wolbachia*. *Med Vet Entomol* **2018**, *32*, 115–20.
63. Shaikevich, E. PCR-RFLP of the COI gene reliably differentiates *Cx. pipiens*, *Cx. pipiens f. molestus* and *Cx. torrentium* of the *Pipiens* Complex. *Eur Mosq Bull* **2007**, *23*, 25–30.
64. Gomes, B., Wilding, C.S., Weetman, D., Sousa, C.A., Novo, M.T., Savage, H.M., et al. Limited genomic divergence between intraspecific forms of *Culex pipiens* under different ecological pressures. *BMC Evol Biol* **2015**, *15*, 197.
65. Kothera, L., Godsey, M., Mutebi, J.P., Savage, H.M. A comparison of aboveground and belowground populations of *Culex pipiens* (Diptera: Culicidae) mosquitoes in Chicago, Illinois, and New York City, New York, using microsatellites. *J Med Entomol* **2010**, *47*, 805–13.
66. Kim, S., Trocke, S., Sim, C. Comparative studies of stenogamous behaviour in the mosquito *Culex pipiens* complex. *Med Vet Entomol* **2018**, *32*, 427–35.