

Article

Fusarium Culmorum Produces NX-2 Toxin Simultaneously With Deoxynivalenol and 3-Acetyl-Deoxynivalenol or Nivalenol (For Submission to Toxins)

Simon Schiwek^{1*}, Mohammad Alhussein², Charlotte Rodemann³, Tuvshinjargal Budragchaa⁴, Lukas Beule⁵, Andreas von Tiedemann³ and Petr Karlovsky^{2*}

¹ Julius Kuehn-Institute, Institute for Plant Protection in Field Crops and Grassland, D-38104 Braunschweig, Germany

² Molecular Phytopathology and Mycotoxin Research, University of Goettingen, D-37077 Goettingen, Germany

³ Plant Phytopathology and Crop Protection, University of Goettingen, D-37077 Goettingen, Germany

⁴ Leibniz Institute for Plant Biochemistry, Department of Bioorganic Chemistry, D-06120 Halle (Saale), Germany

⁵ Julius Kuehn-Institute, Institute for Ecological Chemistry, Plant Analysis and Stored Product Protection, D-14195 Berlin, Germany

* Correspondence: simon.schiwek@julius-kuehn.de; pkarlov@gwdg.de

Abstract: *Fusarium culmorum* is a major pathogen of grain crops. Infected plants accumulate deoxynivalenol (DON), 3-acetyl-deoxynivalenol (3-ADON), or nivalenol (NIV), which are mycotoxins of the trichothecene B group. These toxins are also produced by *F. graminearum* species complex. New trichothecenes structurally similar to trichothecenes B but lacking the carbonyl group on C-8, designated NX toxins, were recently discovered in atypical isolates of *F. graminearum* from North America. Only these isolates and a few strains of a yet to be characterized *Fusarium* species from South Africa are known to produce NX-2 and other NX toxins. Here we report that among 20 *F. culmorum* strains isolated from maize, wheat, and oat in Europe and Asia over a period of 70 years, 18 strains produced NX-2 simultaneously with 3-ADON and DON or NIV. Rice cultures of strains producing 3-ADON accumulated NX-2 in amounts corresponding to 2-8% of 3-ADON (1.2 - 36 mg/kg). A strain producing NIV accumulated NX-2 and NIV at comparable amounts (13.6 and 10.3 mg/kg, respectively). In *F. graminearum*, producers of NX-2 possess a special variant of cytochrome P450 monooxygenase encoded by *TRI1* that is unable to oxidize C-8. In *F. culmorum*, producers and nonproducers of NX-2 possess identical *TRI1*; the reason for the production of NX-2 is unknown. Our results indicate that production of NX-2 simultaneously with trichothecenes B is a common feature of *F. culmorum*. **Key result:** Isolates of *Fusarium culmorum* obtained from different hosts at different locations over several decades produced NX-2, which is a type A trichothecene structurally related to 3-ADON.

Keywords: NX-2; NX toxins; *Fusarium culmorum*; 3-acetyl-deoxynivalenol; nivalenol; trichothecenes; chemotype

1. Introduction

Fusarium head blight (FHB) is a cosmopolitan disease of small-grain cereals with a high economic impact [1] caused by several *Fusarium* species. *F. graminearum* and *F. culmorum* belong to major causal agents of FHB [2]. Infection of grain crops with *Fusarium* spp. causes yield losses and contamination of grains with toxic metabolites (mycotoxins), impairing food safety [3-4].

F. culmorum belongs to the *F. sambucinum* species complex [5]. It infects a wide range of grain crops such as wheat [2,6], maize [7-9], barley [10,11], oat [10], triticale [11], and rye [11]. *F. culmorum* typically co-occur with other *Fusarium* species. In most reports, the

incidence of *F. culmorum* in crops was second to *F. graminearum*, but in some crops, growing regions, and years, *F. culmorum* dominated [2,6,10]. Infected plant material often contains trichothecene mycotoxins deoxynivalenol (DON), nivalenol (NIV), their acetylated derivatives 3-acetyl-deoxynivalenol (3-ADON) and 15-acetyl-deoxynivalenol (15-ADON), fusarenon X (4-acetyl-nivalenol), and zearalenone (ZEN) [4,12].

Biosynthesis of trichothecenes in *Fusarium* spp. is encoded by *TRI* genes, comprising the *TRI1-TRI16* cluster and the *TRI101* gene located outside of the cluster [13]. According to the dominant trichothecene produced, *Fusarium* strains producing trichothecenes B are partitioned into the 3-ADON chemotype, the 15-ADON chemotype, and the NIV chemotype [13].

The distribution of chemotypes among wheat-producing regions exhibits a strong geographic pattern with most areas dominated by either 3-ADON or 15-ADON chemotype (reviewed in [15]; a succinct overview can also be found in the Introduction of [16]). The 3-ADON chemotype in the USA was assumed to be introduced from Europe [17]. Some studies have not found any relationship between chemotype and aggressiveness of *F. graminearum* [18,19], but other studies found the 3-ADON chemotype more aggressive than the 15-ADON chemotype [20-23]. As several authors suggested [20,21], the discrepancy between earlier and later studies may be accounted for by the use of different inoculation methods because DON is not required for initial infection but it facilitates spread of the pathogen along the spike. Greater aggressiveness of the 3-ADON chemotype as compared to the 15-ADON chemotype is in line with a shift of *F. graminearum* populations from 15-DON to 3-ADON producers in North America over the last two decades [16,21]. In several studies of the population structure of *F. graminearum*, fitness advantage of 3-ADON producers due to their higher aggressiveness was postulated and correlations with fitness parameters were determined. The reason for a greater aggressiveness of 3-ADON producers as compared to 15-ADON producers has rarely been addressed. A plausible hypothesis was coined by Poppenberger et al. [24] based on their finding that 3-ADON was protected against glucosylation by UDP-glucosyltransferase from *Arabidopsis thaliana* (see also Section 3.5.).

Trichothecene chemotype has been monitored extensively in *F. graminearum* (e.g., [9,16,18-23,25]; reviewed in [15]). Fewer studies monitored the chemotype in *F. culmorum*, which only comprises 3-ADON and NIV types [9,26-28]. In the past, chemotypes were often assigned according polymorphisms in *TRI* genes [25], but frequently reported discrepancies between the chemotype prediction by PCR-RFLP and results of chemical analysis have questioned this approach [14,16,29].

During an investigation of *F. graminearum* strains colonizing wheat heads in North America, a new type A trichothecene mycotoxin, the NX-2, was discovered [30]. The structure of NX-2 is similar to 3-ADON but it lacks a carbonyl group at C-8. Deacetylated derivative of NX-2, called NX-3, has also been reported. Very recently, these metabolites have also been found in a yet to be characterized species of the *Fusarium sambucinum* species complex from South Africa [31]. The first strains of *F. graminearum* from North America producing NX toxins did not produce any trichothecene type B under laboratory conditions. The lack of known trichothecenes actually motivated the investigation that led to the discovery of NX toxins [30]. Strains producing NX-2 were originally found at very low frequency but a recent study by Lofgren et al. [32] estimated that 20% of the *F. graminearum* population in the USA produce NX-2. An even more recent study of *F. graminearum* strains in Canada [16] identified producers of NX-2 (which they designated 3ANX for consistency with the nomenclature of ADONs) among strains of *F. graminearum* assigned to the 15-ADON chemotype. Contrary to the reports that NX-2-producing strains from the USA have not produced DON, NIV, or their acetylated derivatives [30,33], most strains in their work [16] produced NX-2 simultaneously with 15-ADON.

Here we report that most isolates of *F. culmorum* collected in Europe and Asia produced NX-2 toxin simultaneously with DON, 3-ADON, or NIV.

2. Results

2.1. Mycotoxin production in rice cultures

Our set of 20 *F. culmorum* strains consisted of 14 strains isolated from maize, wheat, and oat in Germany and 6 strains obtained from other laboratories and culture collections, which were isolated in different countries in a time span of 70 years. The analysis of rice culture extracts revealed accumulation of NX-2 at concentrations larger than 1 mg/kg in 18 cultures (**Table 1**). All strains except one were primarily 3-ADON producers; one isolate was assigned to the NIV chemotype. The amounts of NX-2 and 3-ADON in cultures of strains of the 3-ADON chemotype were tightly correlated (**Figure 1**). NX-2 in these cultures accumulated to levels corresponding to 2-8% of 3-ADON. In the culture of the only isolate of the NIV chemotype (isolate 59.6st), the concentrations of NIV and NX-2 were comparable (**Table 1**).

Table 1. Accumulation of trichothecenes in rice cultures of *F. culmorum* strains.

Isolate	Mycotoxin content (mg/kg)				
	NIV	DON	15-ADON	3-ADON	NX-2
31.6st	0.04	25.6	<LOD	156	3.2
215.1st	<LOD	4.2	<LOD	90.4	2.4
227.2cst	<LOD	0.5	<LOD	2.1	<LOD
240.2sp	0.38	355	<LOD	552	35.6
IPP0211	0.31	566	<LOD	412	12.2
IPP0212	0.10	162	<LOD	463	16.4
IPP0213	0.33	639	<LOD	388	9.4
IPP0618	<LOD	20.3	<LOD	175	3.7
IPP0619	<LOD	15.9	<LOD	272	14.4
IPP0999	<LOQ	<LOD	<LOD	<LOD	<LOD
IPP1000	<LOD	12.6	<LOD	132	2.7
DSM62184	0.05	108	<LOD	173	13.5
966	<LOD	25.7	<LOD	118	3.2
969	<LOQ	14.3	<LOD	147	3.2
3.37	<LOD	9.0	<LOD	68.6	1.4
DSM62188	<LOQ	46.5	<LOD	294	13.6
55.6st	<LOQ	31.3	<LOD	85.9	1.2
59.6st	13.6	0.4	0.9	0.63	10.3
K11.2	<LOD	1.0	<LOD	46.3	1.2
J31.2	0.08	127	<LOD	447	18.6

LOD = limit of determination; LOQ = limit of quantification.

2.2. Confirmation of the structure of NX-2

Putative NX-2 accumulating in rice cultures of *F. culmorum* was originally identified by comparing its retention time in HPLC and MS/MS fragmentation with data obtained with purified NX-2 from Prof. Franz Berthiller (BOKU Vienna, Austria). Because production of NX-2 by *F. culmorum* has not been reported before and an isomer of NX-2 could possess the same retention time and generate product ions with the same m/z values, we purified putative NX-2 from rice cultures of *F. culmorum* 240.2sp (**Figure 2**) to verify its structure. Approx. 5 mg of pure metabolite were obtained from 480 g of dry rice culture (see Section 5.4 for details).

One- (¹H and ¹³C) and two-dimensional (¹H,¹³C-HSQC, ¹H,¹³C-HMBC, ¹H,¹H-COSY) NMR spectroscopic analysis was performed on the purified metabolite (**Supplementary Figure S1-S6**). The spectroscopic data were in accordance with published data for NX-2 (**Supplementary Table S1**). The structure of NX-2 is shown in **Figure 3**.




Figure 1. Relationship (Pearson correlation) between NX-2 and 3-ADON concentrations in rice cultures of *F. culmorum*. Each culture (**Table 1**) is represented by a single data point (black dots). Isolate 59.6st, which was assigned to the NIV chemotype, and isolates 227.2 and IPP0999, in which the concentrations of NX-2 were below the limit of detection, were excluded.

2.3. Species assignment and investigation of polymorphisms in *TRI1*

Examination of spores assigned all isolates analyzed for trichothecene production (**Table 1**) to *Fusarium culmorum*. Morphological characterization was complemented by the analysis of melting curves of amplicons of taxonomically informative genes encoding translation elongation factor 1 α (TEF-1 α) and the second largest component of the RNA polymerases II (RPB2) [34]. Species identification was further strengthened by the analysis of full-length sequence (1,753 nt) of the *TRI1* gene, which encodes cytochrome P450 monooxygenase catalyzing oxygenation of calonecetrin on C-7 and C-8. The sequences of *TRI1* obtained from isolates of *F. culmorum* used in this study (accession numbers OM144918 to OM144937) were aligned with a set of reference sequences (**Supplementary Table S3**) and used for phylogenetic analysis by the maximum-likelihood method. Separation of *F. culmorum* from other *Fusarium* species was highly supported, indicating that *TRI1* is taxonomically informative in *Fusarium* species producing trichothecenes.

The reason for selecting the *TRI1* gene for the analysis was that the product of *TRI1* catalyzes biosynthetic steps distinguishing NX toxins from trichothecenes B, and that polymorphisms in *TRI1* differentiating *F. graminearum* strains producing NX toxins from nonproducers were identified [30]. As expected, all *F. graminearum* strains not producing NX toxins were separated from all strains producing NX toxins. In contrast, no polymorphisms separating *F. culmorum* strains producing NX toxins from nonproducers were found in the *TRI1* gene (**Figure 4**).




Figure 2. Purification of NX-2 toxin from rice culture of *F. culmorum* 240.2sp. Three weeks-old rice cultures were extracted with methanol/water/acetic acid and putative NX-2 was enriched by chromatography on C18 cartridge (a), polar-modified C18 column (b), and Sephadex LH-20 (c). Purity of the metabolite was established by HPLC-MS in a full-scan mode (d).




Figure 3. Structure of NX-2 toxin.

Amino acid sequences of translation products of *TRI1* genes, designated Tri1, were identical for all isolates of *F. culmorum* in this study. Isolates of *F. graminearum* producing NX toxins differed from nonproducers in 14 amino acid residues within the heme-binding motif [35] (Table 2). In all *F. culmorum* strains used in this study, comprising 18 producers

and 2 nonproducers of NX-2, these amino acid residues were identical, and they matched the corresponded residues in strains of *F. graminearum* that did not produce NX toxins. Thus, the reason for the production of NX-2 by *F. culmorum* is not an NX-specific form of *TRI1* found in NX-2 producers of *F. graminearum*.



Figure 4. Maximum likelihood estimate of phylogenetic relationships among *TRI1* genes in *Fusarium* spp. Complete sequences of *TRI1* genes of the investigated isolates of *F. culmorum* (see **Table 3**) and reference sequences were subjected to maximum likelihood analysis [36] assuming Tamura-Nei model [37]. Bootstrap values (1,000 replications) are shown next to the nodes. Production of NX-2 was determined by HPLC-MS/MS; n.d. stands for no data available. Nucleotide sequences were deposited at NCBI with accession nos. listed in **Supplementary Table S3**.

3. Discussion

3.1. Production of NX-2 is a characteristic feature of *F. culmorum*

So far, production of NX-2 has exclusively be reported in *F. graminearum*, which shares a high level of genomic similarity with *F. culmorum*, produces the same mycotoxins, and has the same host range [6,38]. The production of NX-2 was discovered in an atypical population of *F. graminearum* from the Midwest of the U.S.A., named Northland popula- tion, which did not produce any known trichothecene [33]. Producers of NX-2 were later found in this population [30] but only a small fraction of *F. graminearum* strains collected in the area produced NX toxins. A later study on *F. graminearum* in wheat in Ontario (Can- ada), however, reported that 80% of investigated strains produced NX toxins [16]. A key finding of the current study is that NX-2 production is not limited to *F. graminearum*, and, contrary to the previous assumption [35], it is not endemic to Northern USA and Southern Canada.

Table 2. Amino acid residues specific for the production of NX-2 in the translation product of *TRI1* of *F. graminearum* and corresponding residues in *F. culmorum*. Species-specific positions are highlighted in purple for *F. culmorum* and blue for *F. graminearum*. Positions reported to distinguish NX-2-producing strains of *F. graminearum* [35] are marked red.

Location in <i>TRI1</i>	Position of amino acid residue																			
	5	5	8	33	35	100	115	134	210	252	254	256	346	361	373	418	430	450		
<i>F. culmorum</i> NX+ (n = 18) ^a	I	S	S	A	K	S	F	G	S	R	L	T	F	I	D	Q	T	A		
<i>F. culmorum</i> NX- (n = 2) ^b	I	S	S	A	K	S	F	G	S	R	L	T	F	I	D	Q	T	A		
<i>F. graminearum</i> NX+ (n = 3) ^c	L	T	Q	T	Q	N	L	A	T	S	M	N	I	F	E	K	P	V		
<i>F. graminearum</i> NX- (n = 3) ^d	L	T	Q	A	Q	S	F	A	S	R	L	T	F	I	D	Q	T	A		
<i>F. cerealis</i> 25805	I	S	S	A	K	S	F	G	S	R	L	T	F	I	D	Q	T	A		
<i>F. pseudograminearum</i> 28062	I	S	S	A	K	S	F	G	S	R	L	T	F	I	D	Q	T	A		
<i>F. poae</i> FRC T-0962	I	D	P	T	K	R	F	A	S	N	L	N	F	V	D	Q	T	A		
<i>F. incarnatum</i> NRRL31160	M	N	N	A	K	K	F	A	S	H	L	D	K	V	D	D	T	A		
<i>F. sambucinum</i> FRC R-07843	L	D	S	R	K	N	F	A	T	R	L	N	K	V	D	Q	T	A		
<i>F. langsethiae</i> NRRL53410	F	D	E	R	K	N	F	A	A	R	L	N	K	V	D	L	T	A		
<i>F. sporotrichioides</i> NRRL29977	F	D	E	R	K	N	F	A	A	R	L	N	K	V	D	Q	T	A		

^a All isolates of *F. culmorum* that produced NX-2 (**Table 1**); ^b Isolates 227.2cst and IPP0999. ^c Isolates 06-204, 02-264, and 03-348. ^d Isolates PH-1, 06-627, and 38383. The accession numbers are listed in **Supplementary Table S3**.

The ability to produce NX-2 appears to be a common feature of *F. culmorum*. Among 20 strains of *F. culmorum* isolated in Europe and Asia over a period of 70 years, 18 strains produced NX-2 (**Table 1**). We suppose that the production of NX-2 toxins in cultures of *F. culmorum* remained unnoticed because NX toxins were not monitored in routine surveys. No commercial standards for NX-2 are available at the time of writing. Because infection of grain crops with *F. culmorum* is widespread, we assume that contamination of grains and grain products with NX-2 might be common.

3.2. Both chemotypes of *F. culmorum* produce NX-2

Isolates of *F. culmorum* belong to the 3-ADON and NIV chemotypes; 15-ADON chemotype is absent [26-28,39,40]. Most *F. culmorum* isolates studied in this work belonged to the 3-ADON chemotype and produced NX-2 toxin. The only strain of the NIV chemotype produced NX-2 toxin, too (**Table 1**). Investigation of further isolates has yet to confirm that this generally holds for strains of the NIV chemotype. In *F. graminearum*, all producers of NX toxins identified in the first report from the USA belonged to the 3-ADON chemotype [30]. Another group studying *F. graminearum* strains from Canada confirmed this finding: NX-2 producers were rare, and all of them (including 5 isolates from Ontario, see below) were assigned to the 3-ADON chemotype [41]. The most recent study on *F. graminearum* from Ontario reported contradictory results, assigning most NX-2 producers to the 15-ADON chemotype [16]. Production of NX-2 was verified by chemical analysis in both studies. In the study of Kelly et al. [41], the 3-ADON/15-ADON chemotype was assessed by PCR-RFLP, while Crippin et al. [16] used both PCR-RFLP and chemical analysis. Incongruencies between the chemotype prediction by PCR-RFLP and the results of chemical analysis, reported by several studies [14,16,29,42], cannot explain the contradictory chemotype assignments mentioned above because both studies used the same PCR assay (*TRI3/TRI12*), yet Crippin et al. [16] reported that all their NX-2 producers belonged to the 15-ADON chemotype while Kelly et al. [41] assigned all their NX-2 producers to the 3-ADON chemotype. Crippin et al. [16] also analyzed trichothecenes in cultures on three growth media by HPLC-MS/MS, using a special elution gradient for the separation of 3-

ADON and 15-ADON. (These mycotoxins often co-elute, and they cannot be reliably distinguished by MS fragmentation). They reported the concentrations of 15-ADON, but unfortunately not the concentrations of 3-ADON and/or DON.

The acetylation of C-3-OH of NX-2 (**Figure 3**) is reminiscent of 3-ADON. Varga et al. [30] showed that when part of the translation product of *TRI1* in a strain of the 15-ADON chemotype was replaced with a *TRI1* segment specific for the production of NX-2, the recombinant strain accumulated NX-4 (a derivative acetylated on C-15-OH, reminiscent of 15-ADON) rather than NX-2. The replacement of NX-specific polypeptide in an NX-2 producer with a protein segment from a 15-ADON-producer converted the NX-2-producing strain into a 3-ADON producer. Both results strongly support the hypothesis that NX-2 producers in *F. graminearum* developed from strains of the 3-ADON chemotype. The controversy between the assignment of chemotypes to NX-2 producers from Ontario in [41] and [16] remains unresolved.

3.3. *F. culmorum* produces NX-2 simultaneously with DON and 3-ADON or NIV

A search for new trichothecenes in *F. graminearum* that facilitated the discovery of NX toxins was motivated by a failure to detect any trichothecene in a group of atypical strains of *F. graminearum* from the USA [30,33]. These strains produced NX toxins but no trichothecenes B. The recent study from Ontario cited above [16] reported that cultures of *F. graminearum* accumulating NX-2 simultaneously accumulated 15-ADON.

Cultures of NX-2-producing *F. culmorum* strains in our study accumulated 3-ADON (most strains) or NIV (a single strain). In cultures of the 3-ADON chemotype, NX-2 accumulated at amounts corresponding to 2-8% of 3-ADON, and the content of NX-2 and trichothecenes B tightly correlated (**Figure 1**). In contrast, the concentrations of NX-2 to 15-ADON in *F. graminearum* cultures in the study of Crippin et al. [16] were not correlated; the ratio NX-2/15-ADON varied from 0.003 to 33.

The only strain in our work producing NX-2 at an amount comparable to trichothecenes B was a strain of the NIV chemotype (**Table 1**). In rice cultures of this strain, NX-2 reached 75% of the concentration of NIV and 66% of total trichothecenes B. Future studies have to clarify whether relatively high production of NX-2 is a typical feature of strains of *F. culmorum* with the NIV chemotype. In *F. graminearum*, no strain of the NIV chemotype have so far been reported to produce NX-2.

3.4. NX-2 production by *F. culmorum* is not caused by a variant of *TRI1*

The reason for the production of NX toxins by certain strains of *F. graminearum* is that these strains harbor a special variant of the *TRI1* gene [30]. The gene, which is only present in genomes of trichothecene-producing *Fusarium* species [43], encodes cytochrome P450 monooxygenase (calonecetrin C7/C8 hydroxylase) Tri1 that catalyzes oxidation of C-7 and C-8 [44,45]. The difference between *TRI1* of NX-2 producers and nonproducers allowed a PCR-RFLP assay for NX-2 producers [46]. We have not found such a polymorphism in the *TRI1* gene of *F. culmorum*.

In *F. graminearum*, changes in Tri1 specific for NX toxins occurred in the heme binding motif close to the C-terminus. Furthermore, Ramdass et al. [47] found an new potential glycosylation site in the enzyme of NX-2 producers. It was located at a large distance from the heme binding site but the authors suggested that glycosylation may modulate enzyme activity via protein folding. None of these changes occurred in the *TRI1* gene of NX-2-producing *F. culmorum*. Some residues in the heme-binding segment differed from corresponding residues in *F. graminearum* (purple in **Table 2**), but these residues occurred in NX-2 producers as well as nonproducers.

Factors other than the amino acid sequence of Tri1 must suppress its activity towards C-8 of calonecetrin in *F. culmorum* strains producing NX-2. The enzyme is located inside endoplasmic reticulum with two hydrophobic segments close to the ends of the protein crossing the membrane [47]. Other proteins and especially other cytochromes P450, which compete with Tri1 for the same NADPH-cytochrome P450 reductase, share this location. Cytochromes P450 are known to interact with each other and form heterooligomers,

which modifies their activity [48]. Interaction of Tri1 with other proteins anchored in the membrane of endoplasmic reticulum may suppresses the oxidation of C-8 in NX-2-producing strains of *F. culmorum*.

3.5. NX-2 and the aggressiveness of *F. graminearum* and *F. culmorum*

In *F. graminearum*, a shift from 15-ADON to 3-ADON chemotype observed in the last decades supports the hypothesis that the 3-ADON chemotype is more aggressive. For instance, an increase of the 3-ADON chemotype at locations where 15-ADON producers used to be predominant was reported in a study from North America [49]. Many studies of the population structure in *F. graminearum* elucidated the relationship between genotype, chemotype and aggressiveness, aiming to explain how selection and gene flow shaped the populations (e.g., [17,20,35,41,46,49,51,53]). Field studies documented the success of the 3-ADON chemotype of *F. graminearum* but they could not address its cause, which requires a biochemical approach. According to a hypothesis from the lab of Gerhard Adam [24], acetylation of C-3-OH prevents detoxification of DON, which is a virulence factor, by plant UDP-glycosyltransferases. In NX-2, the hydroxyl on C-3 is also protected by acetylation, and Varga et al. [30] suggested that the production of NX-2 may benefit *F. graminearum* during the infection in the same way. Glucosylation of DON takes place in the cytoplasm while the target of DON is protein synthesis in rough endoplasmic reticulum, where the acetyl group would have to be removed. This is plausible because endoplasmic reticulum is rich in hydrolases [50]. Varga et al. [15] also speculated that the lack of carbonyl on C-8 circumvents detoxification by glutathionylation [15]. This hypothesis holds for *F. culmorum*, too.

Can the effect of NX-2 on the aggressiveness of *F. graminearum* or *F. culmorum* be proved? Field trials with natural isolates are unlikely to generate a conclusive proof. The same situation exists for the claim that the 3-ADON chemotype is more aggressive than the 15-ADON chemotype, which was supported by some studies [20-23] yet rejected by others [18,19,53]. Field isolates differ in many properties modulating aggressiveness. Some of them are likely linked to the chemotype. Isogenic strains differing only in the gene in question are required. Regarding relative aggressiveness of the chemotypes 3-ADON and 15-ADON, strains with swapped *TRI8* genes or *TRI8* chimeras, constructed by Alexander et al. [57], could be used. Similarly, infection experiments with *F. graminearum* strains harboring *TRI1* with swapped domains controlling the NX-2 production, constructed by Varga et al. [30], would reveal the effect of NX-2 production on aggressiveness.

3.6. Do further *Fusarium* species produce NX toxins?

Comparison of the amino acid residues in Tri1 sequence distinguishing NX-2-producing isolates from nonproducers in *F. graminearum* (Table 2) showed that certain residues specific for the production of NX-2 are also present in Tri1 of other *Fusarium* species. For instance, Tri1 in a particular *F. sambucinum* strain shared 3 residues with NX-2 producing strains of *F. graminearum*, and it differed from Tri1 of strains not producing NX-2 in other 3 positions. We suggest that this species be examined for NX-2 production. *F. cerealis* and *F. pseudograminearum* harbor *TRI1* genes very similar to *TRI1* of *F. culmorum* (Figure 4), and the amino acid sequence in the heme-binding segment of their Tri1 protein is identical with the corresponding sequence in *F. culmorum* (Table 2). Examination of these species for the production of NX toxins, too, appears worthwhile.

3.7. Can a single strain of *F. culmorum* produce both DON and NIV?

It is generally assumed that *F. culmorum* and *F. graminearum* produces either NIV or DON but not both. We believe that this view is biased by low sensitivity of analytical methods in the past, which only detected trichothecenes that were present at high concentrations (e.g., [27]). Accounts of the production of both trichothecenes by a single strain have been largely overlooked. For instance, Foroud et al. [3] write in their recent review "NIV chemotypes do not produce DON", though many studies, such as [58,20,59] (the first

two of which are even cited in [3]), reported simultaneous production of DON and NIV by single strains of *F. culmorum* or *F. graminearum*.

NIV and DON have a common precursor. In NIV producers, the precursor is hydroxylated at C-4 by Tri13 [57]. We speculate that before the entire pool is hydroxylated, some enters the path leading to DON. Therefore, we assume that all cultures producing NIV contain small amounts of DON. The presence of NIV in cultures of DON producers can be explained by a residual activity of Tri13 or by the presence of hydroxylases with a relaxed substrate specificity. In line with this reasoning, cultures of NIV-producing strain 59.6st in our work accumulated relatively large amounts of DON while cultures of most DON producers contained small amounts of NIV (Table 1). We expect that with the widespread use of sensitive analytical methods, small amounts of NIV will be often found in cultures of *Fusarium* strains producing DON, and substantial amounts of DON will be found in cultures of all strains producing NIV.

4. Conclusions

Most strains of *Fusarium culmorum* produce NX-2 toxin simultaneously with deoxynivalenol, 3-acetyldeoxynivalenol, or nivalenol. The strains producing NX-2 do not possess a specific variant of the *TRI1* gene known from NX-2-producing strains of *F. graminearum*.

5. Materials and Methods

5.1. Fungal strains

Strains of *F. culmorum* are listed in **Table 3**. Strains isolated in the course of this study were obtained from maize grains, rachis, stalks, and from oat grains according to Leslie and Summerell [54]. Briefly, samples were surface sterilized for 10 min with 0.1% silver nitrate or 3% sodium hypochlorite, rinsed, and placed on potato dextrose agar (PDA). Isolates were purified via single-spore cultures and grown on PDA for colony characteristics and on low-nutrient agar (SNA, [55]) under long-wave UV light for morphological characterization of spores. For long-term storage, fungal cultures were freeze-dried.

5.2. DNA methods

Fungal DNA was extracted using a CTAB-based protocol [56] from 10 mg of lyophilized mycelium and dissolved in 50 μ l TE (10 mM Tris, 1 mM EDTA, pH 8.0). Segments of marker genes *TEF-1 α* and *RPB2* were amplified and the PCR products subjected to high-resolution melting curve (HRM) analysis as described previously [34]. The *TRI1* gene was amplified for sequencing as four overlapping fragments (**Table 4**), which were sequenced by the Sanger method and assembled to full-length gene sequences. PCR was carried out in a peqSTAR 96 thermocycler (PEQLAB, Erlangen, Germany) in a total reaction volume of 25 μ l. Reaction mixtures were composed of a buffer (10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl₂, pH 8.3 at 25°C; New England Biolabs, Beverly, MA, USA) adjusted to 2 mM final MgCl₂ concentration containing 100 μ M of each deoxyribonucleoside triphosphate (Bioline, Luckenwalde, Germany), 0.3 μ M of each primer, 0.62 U HotStart-polymerase (New England Biolabs, Beverly, Massachusetts, USA), and 1 μ L template DNA solution diluted 100-times. The thermocycling conditions are specified in **Supplementary Table S2**. PCR products were precipitated with isopropanol, washed with 80% ethanol and sequenced at the facilities of MacroGen Europe (MacroGen Europe, Amsterdam, The Netherlands). The sequences were quality-trimmed with Chromas version 2.6.6 (Technelysium Pty. Ltd., South Brisbane, Australia) and assembled to full-length gene sequences; the accession numbers are listed in **Supplementary Table S3**. Multiple sequence alignments were performed with ClustalW [60] in MEGAX version 10.1.8 [36]. Phylogenetic relationships among *TRI1* genes in *Fusarium* spp. were investigated by maximum-likelihood analysis using MEGA X under the assumption of Tamura's and Nei's substitution model [37].

5.3. Mycotoxin extraction and HPLC-MS/MS

Rice cultures were prepared in 50-ml Falcon tubes (Sarstedt, Nümbrecht, Germany) by autoclaving 3 g dry polished rice with 5 ml tap water. The tubes were inoculated with plugs of PDA (0.5 cm diameter) overgrown with 5-days-old mycelium. Rice medium incubated with agar plugs without mycelium served as a control. The cultures were incubated for 21 days at 25 °C in the dark. Fungal metabolites were extracted by shaking with 30 ml acetonitrile/water/acetic acid (84:15:1 (v/v/v)) overnight. Extracts were dried in a vacuum concentrator (Martin Christ, Osterode am Harz, Germany) and residues re-dissolved in methanol/water (20:80 (v/v)) as described previously [61]. Mycotoxin analysis was carried out using a 1290 Infinity II HPLC system (Agilent Technologies, Waldbronn, Germany) coupled with a 6460 triple quadrupole detector (Agilent Technologies, Waldbronn, Germany). The separation was performed on a Zorbax Eclipse Plus C18 column, 50 x 2.1 mm with 1.8 µm particle size (Agilent Technologies, Waldbronn, Germany). The column oven temperature was 40 °C. Mobile phase A was water with 0.1% formic acid (v/v), and phase B was methanol with 0.1% formic acid (v/v). The gradient was as follows: 0 to 0.2 min, 5% B; 0.2 to 8 min, 5% to 35% B; 8 to 8.5 min, 35% to 98% B; 8.5 to 12 min, 98% B; 12 to 12.5 min, 98% to 5% B; 12.5 to 16 min, 5% B. The calibration curve included 11 concentrations from 0.48 to 500 µg/L. A blank was analyzed after every 7th sample and a quality control standard after every 15th sample. The metabolites were detected in a multiple reaction monitoring (MRM) mode. The acquisition parameters and the limits of detection (LOD) and quantification (LOQ) are listed in **Table 5**.

Table 3. Strains of *F. culmorum* used in this study.

Species	Origin	Year	Isolate	Host
<i>F. culmorum</i>	Germany	2017	31.6st	Maize stalk
<i>F. culmorum</i>	Germany	2018	215.1st	Maize stalk
<i>F. culmorum</i>	Germany	2018	227.2cst	Maize stalk
<i>F. culmorum</i>	Germany	2018	240.2sp	Maize rachis
<i>F. culmorum</i>	-	< 1984 ^a	IPP0211 ^b	Wheat
<i>F. culmorum</i>	Italy	< 1993 ^a	IPP0212 ^b	Wheat
<i>F. culmorum</i>	-	1993	IPP0213 ^b	Barley
<i>F. culmorum</i>	Hungary	1991	IPP0618 ^b	Wheat
<i>F. culmorum</i>	Germany	1991	IPP0619 ^b	Wheat
<i>F. culmorum</i>	Germany	2010	IPP0999 ^b	Wheat
<i>F. culmorum</i>	Germany	2010	IPP1000 ^b	Wheat
<i>F. culmorum</i>	Germany	1952	DSM62184 ^c	Maize grain
<i>F. culmorum</i>	Syria	2009-2010	966 ^d	Wheat
<i>F. culmorum</i>	Syria	2009-2010	969 ^d	Wheat
<i>F. culmorum</i>	Germany	2004	3.37 ^b	Wheat
<i>F. culmorum</i>	Germany	1990	DSM62188 ^c	Maize stalk
<i>F. culmorum</i>	Germany	2017	55.6st	Maize stalk
<i>F. culmorum</i>	Germany	2017	59.6st	Maize stalk
<i>F. culmorum</i>	Germany	2021	K11.2	Oat
<i>F. culmorum</i>	Germany	2021	J31.2	Oat

^a Isolates were collected before the specified year. ^b Isolates of the Plant Phytopathology and Crop Protection Section at the University of Göttingen, Göttingen, Germany. ^c Strains were obtained from the German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany.

^d The isolates were described by Alkadri et al. [40].

5.4. Purification of NX-2

F. culmorum isolate 240.2sp, which produced the largest amounts of NX-2, was cultivated on rice media. The cultures were prepared in 4 l flat penicillin flasks with 120 g

organic polished rice and 270 ml tap water, which were autoclaved and inoculated with 12 agar-plugs (0.6 cm diameter) from 5-day old fungal cultures. Rice cultures were incubated at 25 °C for three weeks in the dark. NX-2 was extracted with a 5-time excess of methanol/water/acetic acid (90:9:1) by shaking overnight. The extract was concentrated to approximately 15% of its original volume using a rotary evaporator R-100 (Buchi, Flawil, Switzerland). The concentrated extract was partitioned with the same volume of ethyl acetate (EtOAc) three times. Combined EtOAc fractions were dried in a rotary evaporator. The residue was dissolved in methanol/water (30:70) and cleared by centrifugation. The supernatant was subjected to flash chromatography (Sepacore Flash system X10/X50, Buchi, Flawil, Switzerland) equipped with a binary pump (modules C-601/C-605) and an UV detector (module C-635). Metabolites were separated on a reverse-phase column (Chromoband Flash cartridge RP C₁₈ ec 40–63 µm, 262 x 37 mm, Machery-Nagel, Düren, Germany). Mobile phase A was water with 0.2% acetic acid; phase B was methanol with 0.2% acetic acid. The gradient elution program was as follows: 0-2 min, 5% B; 2-42 min, from 5% to 50% B; 42-45 min, from 50% to 98% B; 45-55 min, 98% B; 55-58 min, from 98% to 5% B; 58-66 min, 5% B. The flow rate was 50 ml min⁻¹. The separation was monitored at 254 nm. NX-2 fraction was collected at 33 min, the solvent was evaporated, fractions containing NX-2 were re-dissolved in methanol/water (30:70) and subjected to a preparative-HPLC (PU-2086 plus, Jasco, Gross-Umstadt, Germany) equipped with a preparative reverse-phase column (Nucleodur C18 pyramid, 5 µm, 250 x 21mm, Macherey-Nagel, Düren, Germany) and coupled to a UV/VIS detector (UV-970, Jasco, Gross-Umstadt, Germany). NX-2 was eluted by a gradient of water with 0.2% acetic acid (v/v) (solvent A) and methanol with 0.2% acetic acid (solvent B) as follows: 0-5 min, 5% B; 5-50 min, from 5% to 50% B; 50-55 min, from 50% to 98% B; 55-70 min, 98% B; 70-75 min, from 98% to 5% B; 75-90 min, 5% B at a flow rate of 14 ml min⁻¹. The separation was monitored at 254 nm. NX-2 was collected at 43 min and the solvent evaporated in a vacuum concentrator (Christ, Osterode am Harz, Germany). NX-2 was further polished on Sephadex LH-20 (Sigma-Aldrich, Darmstadt, Germany) in a 460 x 15 mm column eluted isocratically with methanol at a flow rate of 1 ml min⁻¹. The eluent was monitored at 254 nm.

Table 4. Oligonucleotides used for the sequence analysis of the gene *TRI1*.

Primer name	Primer sequence (5'-3')	Location ^a	Source
<i>TRI16IF1</i>	GCCTSATAGCGACGATCTTGC	0 - 485	[30]
<i>Tri1_5prime_RV</i>	GACCGTGAATAACCTCCTTGATCAGT		This study
<i>Tri1_5prime_FW</i>	ACTGATCAAGGAGGTTATTCACGGTC	460 - 1225	This study
<i>Tri1_SH_R</i>	CGCTGTCGAGAAGGAACATCTTG		This study
<i>Tri1_SH_F</i>	GGCTATGTACAAGATGTTCTTCTCG	1194 - 1753	This study
<i>FgTRI1-R1</i>	AACAAGTGGCGAGATCAAACC		[30]
<i>FcTri1F</i>	ATGGCTATCATCAGCAG	1 - 1745	This study
<i>Tri1R</i>	CAATTCCAATCGCAGACAA		[46]

^a Location of the PCR product within the nucleotide sequence of the gene *TRI1* (1753 bp).

Table 5. Parameters for HPLC-MS/MS analysis of trichothecenes.

Toxin	Ioniz. mode	Parent ion [m/z]	Fragm. voltage [V]	Collision energy [V]	Product ion [m/z] ^c	LOD ^b [mg/kg]	LOQ ^b [mg/kg]
NIV	Pos	313.1	93	7	205.0 <u>175.0</u>	0.01	0.03
DON	Pos	297.1	100	4	<u>249.1</u> 91.2	0.04	0.13
3-ADON	Pos	339.2	100	8	<u>231.1</u> 203.0	0.02	0.08
15-ADON	Pos	339.2	90	10	321.1	0.07	0.23

				10	<u>261.0</u>		
				10	137.0		
NX-2 ^a	Pos	325.2	90	10	247.2	0.04	0.12
				10	229.2		
				10	199.1		
				25	121.2		
				35	<u>105.1</u>		

^a Analytical reference for NX-2 was kindly provided by Dr. Franz Berthiller (BOKU, Vienna, Austria). ^b Limit of detection (LOD) and limit of quantification (LOQ) were calculated based on the standard deviation of the blank [62]. ^c m/z of product ions used for quantification are underlined.

5.5. Nuclear magnetic resonance (NMR) spectroscopy

1D (¹H and ¹³C) NMR spectra were obtained from Bruker Avance 500 Hz and 2D (¹H,¹³C-HSQC, ¹H,¹³C-HMBC, ¹H,¹H-COSY) NMR spectra were obtained from an Agilent DD2 400 MHz system. The spectra were recorded at 500/400 MHz (¹H) and 126/101 MHz (¹³C). Chemical shifts were referenced to internal TMS (δ_H 0, ¹H) or MeOH-*d*₄ (δ_C 49.0, ¹³C). The data are provided as **Supplementary Figures S1 to S5**.

Supplementary Data: Figure S1: ¹H NMR spectrum of NX-2; Figure S2: ¹³C NMR spectrum of NX-2; Figure S3: ¹H, ¹³C HSQC NMR spectrum of NX-2; Figure S4: ¹H, ¹³C HMBC spectrum of NX-2; Figure S5: ¹H, ¹H COSY spectrum of NX-2. Table S1: Comparison of NMR spectra of NX-2 with metabolite purified from *F. culmorum* 240.2sp. Table S2: Conditions used for the amplification of *TRI1* gene. Table S3: Acc. nos. of sequences of the *TRI1* gene.

References

1. Parry, D.W.; Jenkinson, P.; McLeod, L. *Fusarium* ear blight (scab) in small grain cereals—a review. *Plant Pathol.* **1995**, *44*, 207–238, doi:10.1111/j.1365-3059.1995.tb02773.x.

2. Birr, T.; Hasler, M.; Verreet, J.-A.; Klink, H. Composition and predominance of *Fusarium* species causing fusarium head blight in winter wheat grain depending on cultivar susceptibility and meteorological factors. *Microorganisms* **2020**, *8*, 617, doi:10.3390/microorganisms8040617.

3. Foroud, N.A.; Baines, D.; Gagkaeva, T.Y.; Thakor, N.; Badea, A.; Steiner, B.; Bürstmayr, M.; Bürstmayr, H. Trichothecenes in cereal grains - an update. *Toxins* **2019**, *11*, 634, doi:10.3390/toxins11110634.

4. Polak-Śliwińska, M.; Paszczyk, B. trichothecenes in food and feed, relevance to human and animal health and methods of detection: a systematic review. *Molecules* **2021**, *26*, 454, doi:10.3390/molecules26020454.

5. O'Donnell, K.; Rooney, A.P.; Proctor, R.H.; Brown, D.W.; McCormick, S.P.; Ward, T.J.; Frandsen, R.J.N.; Lysøe, E.; Rehner, S.A.; Aoki, T.; et al. Phylogenetic analyses of *RPB1* and *RPB2* support a middle Cretaceous origin for a clade comprising all agriculturally and medically important Fusaria. *Fungal Genet. Biol.* **2013**, *52*, 20–31, doi:10.1016/j.fgb.2012.12.004.

6. Scherm, B.; Balmas, V.; Spanu, F.; Pani, G.; Delogu, G.; Pasquali, M.; Migheli, Q. *Fusarium culmorum*: causal agent of foot and root rot and head blight on wheat. *Molecular Plant Pathology* **2013**, *14*, 323–341, doi:10.1111/mpp.12011.

7. Logrieco, A.; Mulè, G.; Moretti, A.; Bottalico, A. Toxigenic *Fusarium* species and mycotoxins associated with maize ear rot in Europe. *Eur. J. Plant Pathol.* **2002**, *108*, 597–609, doi:10.1023/A:1020679029993.

8. Pfordt, A.; Ramos Romero, L.; Schiwek, S.; Karlovsky, P.; von Tiedemann, A. Impact of environmental conditions and agronomic practices on the prevalence of *Fusarium* species associated with ear- and stalk rot in maize. *Pathogens* **2020**, *9*, 236, doi:10.3390/pathogens9030236.

9. Basler, R. Diversity of *Fusarium* species isolated from UK forage maize and the population structure of *F. graminearum* from maize and wheat. *PeerJ* **2016**, *4*, 2143, doi:10.7717/peerj.2143.

10. Sundheim, L.; Brodal, G.; Hofgaard, I.S.; Rafoss. Temporal variation of mycotoxin producing fungi in Norwegian cereals. *Microorganisms* **2013**, *1*, 188–198, doi:10.3390/microorganisms1010188.

11. Nielsen, L.K.; Jensen, J.D.; Nielsen, G.C.; Jensen, J.E.; Spliid, N.H.; Thomsen, I.K.; Justesen, A.F.; Collinge, D.B.; Jørgensen, L.N. *Fusarium* head blight of cereals in Denmark: species complex and related mycotoxins. *Phytopathology* **2011**, *101*, 960–969, doi:10.1094/PHYTO-07-10-0188.

12. Ropejko, K.; Twarużek, M. Zearalenone and its metabolites - general overview, occurrence, and toxicity. *Toxins* **2021**, *13*, 35, doi:10.3390/toxins13010035.

13. McCormick, S.P.; Stanley, A.M.; Stover, N.A.; Alexander, N.J. Trichothecenes: from simple to complex mycotoxins. *Toxins*. **2011**, *3*, 802–814, doi:10.3390/toxins3070802.

14. Villafana, R.T.; Ramdass, A.C.; Rampersad, S.N. *TRI* genotyping and chemotyping: a balance of power. *Toxins* **2020**, *12*, 64, doi:10.3390/toxins12020064.
15. van der Lee, T.; Zhang, H.; van Diepeningen, A.; Waalwijk, C. Biogeography of *Fusarium graminearum* species complex and chemotypes: a review. *Food Addit. Contam.* **2015**, *32*, 453–460, doi:10.1080/19440049.2014.984244.
16. Crippin, T.; Renaud, J.B.; Sumarah, M.W.; Miller, J.D. Comparing genotype and chemotype of *Fusarium graminearum* from cereals in Ontario, Canada. *PLoS One* **2019**, *14*, e0216735, doi:10.1371/journal.pone.0216735.
17. Gale, L.R.; Ward, T.J.; Balmas, V.; Kistler, H.C. Population subdivision of *Fusarium graminearum sensu stricto* in the upper midwestern United States. *Phytopathology* **2007**, *97*, 1434–1439, doi:10.1094/PHYTO-97-11-1434.
18. Gilbert, J.; Clear, R.M.; Ward, T.J.; Gaba, D.; Tekauz, A.; Turkington, T.K.; Woods, S.M.; Nowicki, T.; O'Donnell, K. Relative aggressiveness and production of 3- or 15-acetyl deoxynivalenol and deoxynivalenol by *Fusarium graminearum* in spring wheat. *Can. J. Plant Pathol.* **2010**, *32*, 146–152, doi:10.1080/07060661003740231.
19. von der Ohe, C.; Gauthier, V.; Tamburic-Ilincic, L.; Brule-Babel, A.; Fernando, W.G.D.; Clear, R.; Ward, T.J.; Miedaner, T. A comparison of aggressiveness and deoxynivalenol production between Canadian *Fusarium graminearum* isolates with 3-acetyl and 15-acetyldeoxynivalenol chemotypes in field-grown spring wheat. *Eur. J. Plant Pathol.* **2010**, *127*, 407–417, doi:10.1007/s10658-010-9607-z
20. Puri, K.D.; Zhong, S. The 3ADON population of *Fusarium graminearum* found in North Dakota is more aggressive and produces a higher level of DON than the prevalent 15ADON population in spring wheat. *Phytopathology* **2010**, *100*, 1007–1014, doi:10.1094/PHYTO-12-09-0332.
21. Malhipour, A.; Gilbert, J.; Piercey-Normore, M.; Cloutier, S. Molecular phylogenetic analysis, trichothecene chemotype patterns, and variation in aggressiveness of *Fusarium* isolates causing head blight in wheat. *Plant Dis.* **2012**, *96*, 1016–1025, doi:10.1094/PDIS-10-11-0866-RE.
22. Liu, Y.Y.; Sun, H.Y.; Li, W.; Xia, Y.L.; Deng, Y.Y.; Zhang, A.X.; Chen, H.G. Fitness of three chemotypes of *Fusarium graminearum* species complex in major winter wheat-producing areas of China. *PLOS One* **2017**, *12*, e0174040, doi:10.1371/journal.pone.0174040.
23. Amarasinghe, C.; Sharanowski, B.; Dilantha Fernando, W.G. Molecular phylogenetic relationships, trichothecene chemotype diversity and aggressiveness of strains in a global collection of *Fusarium graminearum* species. *Toxins* **2019**, *11*, 263, doi:10.3390/toxins11050263.
24. Poppenberger, B.; Berthiller, F.; Lucyshyn, D.; Sieberer, T.; Schuhmacher, R.; Krska, R.; Kuchler, K.; Glössl, J.; Luschnig, C.; Adam, G. Detoxification of the *Fusarium* mycotoxin deoxynivalenol by a UDP-glucosyl-transferase from *Arabidopsis thaliana*. *Journal of Biological Chemistry* **2003**, *278*, 47905–47914, doi:10.1074/jbc.M307552200.
25. Pasquali, M.; Beyer, M.; Logrieco, A.; Audenaert, K.; Balmas, V.; Basler, R.; Boutigny, A.L.; Chrpová, J.; Czembor, E.; Gagkaeva, T.; et al. A European database of *Fusarium graminearum* and *F. culmorum* trichothecene genotypes. *Front. Microbiol.* **2016**, *7*, 406, doi:10.3389/fmicb.2016.00406.
26. Jennings, P.; Coates, M.E.; Turner, J.A.; Chandler, E.A.; Nicholson, P. Determination of deoxynivalenol and nivalenol chemotypes of *Fusarium culmorum* isolates from England and Wales by PCR assay. *Plant Pathol.* **2004**, *53*, 182–190, doi:10.1111/j.1365-3059.2004.00985.x.
27. Quarta, A.; Mita, G.; Haidukowski, M.; Santino, A.; Mulè, G.; Visconti, A. Assessment of trichothecene chemotypes of *Fusarium culmorum* occurring in Europe. *Food Addit. Contam.* **2005**, *22*, 309–315, doi:10.1080/02652030500058361.
28. Matny, O.N.; Bates, S.T.; Song, Z. Geographic distribution of *Fusarium culmorum* chemotypes associated with wheat crown rot in Iraq. *J. Plant. Prot. Res.* **2017**, *57*, 43–49, DOI: <https://doi.org/10.1515/jppr-2017-0006>.
29. Kulik, T.; Buško, M.; Bilska, K.; Ostrowska-Kołodziejczak, A.; Van Diepeningen, A.D.; Perkowski, J.; Stenglein, S. Depicting the discrepancy between *TRI* genotype and chemotype on the basis of strain CBS 139514 from a field population of *F. graminearum sensu stricto* from Argentina. *Toxins* **2016**, *8*, 330, doi:10.3390/toxins8110330.
30. Varga, E.; Wiesenberger, G.; Hametner, C.; Ward, T.J.; Dong, Y.; Schöffbeck, D.; McCormick, S.; Broz, K.; Stückler, R.; Schuhmacher, R.; et al. New tricks of an old enemy: Isolates of *Fusarium graminearum* produce a type A trichothecene mycotoxin. *Environ. Microbiol.* **2015**, *17*, 2588–2600, doi:10.1111/1462-2920.12718.
31. Laraba, I.; McCormick, S.P.; Vaughan, M.M.; Geiser, D.M.; O'Donnell, K. Phylogenetic diversity, trichothecene potential, and pathogenicity within *Fusarium sambucinum* species complex. *PLoS One* **2021**, *16*, e0250812, doi:10.1371/journal.pone.0245037.
32. Lofgren, L.; Riddle, J.; Dong, Y.; Kuhnem, P.R.; Cummings, J.A.; Del Ponte, E.M.; Bergstrom, G.C.; Kistler, H.C. A high proportion of NX-2 genotype strains are found among *Fusarium graminearum* isolates from northeastern New York State. *Eur J Plant Pathol* **2018**, *150*, 791–796, doi:10.1007/s10658-017-1314-6.
33. Gale, L.R.; Ward, T.J.; Kistler H.C. A subset of the newly discovered Northland population of *Fusarium graminearum* from the U.S. does not produce the B-type trichothecenes DON, 15ADON, 3ADON or NIV.

- In Proceedings of the National Fusarium Head Blight Forum, 2010, Canty, S., Clark, A., Anderson-Scully, A., Ellis, D., and Van Sanford, D. (eds). Milwaukee, WI, USA, pp. 48–49.
34. Schiwek, S.; Beule, L.; Vinas, M.; Tiedemann, A. Von; Karlovsky, P. High-resolution melting (HRM) curve assay for the identification of eight *Fusarium* species causing ear rot in maize. *Pathogens*. **2020**, *9*, 270, doi:10.3390/pathogens9040270.
 35. Kelly, A.; Proctor, R.H.; Belzile, F.; Chulze, S.N.; Clear, R.M.; Cowger, C.; Elmer, W.; Lee, T.; Obanor, F.; Waalwijk, C.; et al. The geographic distribution and complex evolutionary history of the NX-2 trichothecene chemotype from *Fusarium graminearum*. *Fungal Genet. Biol.* **2016**, *95*, 39–48, doi:10.1016/j.fgb.2016.08.003.
 36. Kumar, S.; Stecher, G.; Li, M.; Knyaz, C.; Tamura, K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* **2018**, *35*, 1547–1549, doi:10.1093/molbev/msy096.
 37. Tamura, K.; Nei, M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* **1993**, *10*, 512–526, doi:10.1093/oxfordjournals.molbev.a040023.
 38. Goswami, R.S.; Kistler, H.C. Heading for Disaster: *Fusarium graminearum* on cereal crops. *Mol Plant Pathol* **2004**, *5*, 515–525, doi:10.1111/j.1364-3703.2004.00252.x.
 39. Oufensou, S.; Balmas, V.; Scherm, B.; Rau, D.; Camilio, S.; Prota, V.A.; Ben-Attia, M.; Gargouri, S.; Pasquali, M.; El-Bok, S. Genetic variability, chemotype distribution, and aggressiveness of *Fusarium culmorum* on durum wheat in Tunisia. *Phytopathol. Mediterr.* **2019**, *58*, 103–113, doi:10.14601/Phytopathol_Mediterr-23907.
 40. Alkadri, D.; Nipoti, P.; Döll, K.; Karlovsky, P.; Prodi, A.; Pisi, A. Study of fungal colonization of wheat kernels in Syria with a focus on *Fusarium* species. *International Journal of Molecular Sciences* **2013**, *14*, 5938–5951, doi:10.3390/ijms14035938.
 41. Kelly, A.C.; Clear, R.M.; O'Donnell, K.; McCormick, S.; Turkington, T.K.; Tekauz, A.; Gilbert, J.; Kistler, H.C.; Busman, M.; Ward, T.J. Diversity of *Fusarium* head blight populations and trichothecene toxin types reveals regional differences in pathogen composition and temporal dynamics. *Fungal Genet. Biol.* **2015**, *82*, 22–31, doi:10.1016/j.fgb.2015.05.016.
 42. Pasquali, M.; Beyer, M.; Bohn, T.; Hoffmann, L. Comparative analysis of genetic chemotyping methods for *Fusarium*: Tri13 polymorphism does not discriminate between 3- and 15-acetylated deoxynivalenol chemotypes in *Fusarium graminearum*. *Journal of Phytopathology* **2011**, *159*, 700–704, doi:10.1111/j.1439-0434.2011.01824.x.
 43. Proctor, R.H.; McCormick, S.P.; Kim, H.-S.; Cardoza, R.E.; Stanley, A.M.; Lindo, L.; Kelly, A.; Brown, D.W.; Lee, T.; Vaughan, M.M.; et al. Evolution of structural diversity of trichothecenes, a family of toxins produced by plant pathogenic and entomopathogenic fungi. *PLOS Pathogens* **2018**, *14*, e1006946, doi:10.1371/journal.ppat.1006946.
 44. McCormick, S.P.; Harris, L.J.; Alexander, N.J.; Ouellet, T.; Saparno, A.; Allard, S.; Desjardins, A.E. Tri1 in *Fusarium graminearum* encodes a P450 oxygenase. *Appl. Environ. Microbiol.* **2004**, *70*, 2044–2051, doi:10.1128/AEM.70.4.2044–2051.2004.
 45. Kimura, M.; Tokai, T.; Takahashi-Ando, N.; Ohsato, S.; Fujimura, M. Molecular and genetic studies of *Fusarium* trichothecene biosynthesis: pathways, genes, and evolution. *Bioscience, Biotechnology, and Biochemistry* **2007**, *71*, 2105–2123, doi:10.1271/bbb.70183.
 46. Liang, J.M.; Xayamongkhon, H.; Broz, K.; Dong, Y.; McCormick, S.P.; Abramova, S.; Ward, T.J.; Ma, Z.H.; Kistler, H.C. Temporal dynamics and population genetic structure of *Fusarium graminearum* in the upper Midwestern United States. *Fungal Genet. Biol.* **2014**, *73*, 83–92, doi:10.1016/j.fgb.2014.10.002.
 47. Ramdass, A.C.; Villafana, R.T.; Rampersad, S.N. Comparative Sequence Analysis of TRI1 of *Fusarium*. *Toxins* **2019**, *11*, 689, doi:10.3390/toxins11120689.
 48. Davydov, D.R.; Davydova, N.Y.; Sineva, E.V.; Halpert, J.R. Interactions among cytochromes P450 in mitochondrial membranes. *J Biol Chem* **2015**, *290*, 3850–3864, doi:10.1074/jbc.M114.615443.
 49. Ward, T.J.; Clear, R.M.; Rooney, A.P.; O'Donnell, K.; Gaba, D.; Patrick, S.; Starkey, D.E.; Gilbert, J.; Geiser, D.M.; Nowicki, T.W. An adaptive evolutionary shift in *Fusarium* head blight pathogen populations is driving the rapid spread of more toxigenic *Fusarium graminearum* in North America. *Fungal Genet. Biol.* **2008**, *45*, 473–484, doi:10.1016/j.fgb.2007.10.003.
 50. Hara-Nishimura, I.; Matsushima, R.; Shimada, T.; Nishimura, M. Diversity and formation of endoplasmic reticulum-derived compartments in plants. are these compartments specific to plant cells? *Plant Physiol* **2004**, *136*, 3435–3439, doi:10.1104/pp.104.053876.
 51. Castiblanco, V.; Castillo, H.E.; Miedaner, T. Candidate genes for aggressiveness in a natural *Fusarium culmorum* population greatly differ between wheat and rye head blight. *Journal of Fungi* **2018**, *4*, 14, doi:10.3390/jof4010014.
 52. Miedaner, T.; Vasquez, A.; Castiblanco, V.; Castillo, H.E.; Foroud, N.; Würschum, T.; Leiser, W. genome-wide association study for deoxynivalenol production and aggressiveness in wheat and rye head blight by resequencing 92 isolates of *Fusarium culmorum*. *BMC Genomics* **2021**, *22*, 1–16, doi:10.1186/s12864-021-07931-5.

53. Mueller, B.; Groves, C.L.; Smith, D.L. chemotype and aggressiveness evaluation of *Fusarium graminearum* and *Fusarium culmorum* isolates from wheat fields in Wisconsin. *Plant Disease* **2021**, *105*, 3686–3693, doi:10.1094/PDIS-06-20-1376-RE.
54. Leslie, John F. Summerell, B.A. The *Fusarium* Laboratory Manual; Kansas State University, Manhattan, USA; ISBN: 9780813819198; <https://doi.org/10.1002/9780470278376>, **2006**.
55. Nirenberg, H. Untersuchungen über die morphologische und biologische Differenzierung in der *Fusarium*-Sektion Liseola. *Mitt. Biol. Bundesanst. Land und Forstwirtschaft. Berlin-Dahlem* **1976**, *169*: 1-117.
56. Brandfass, C.; Karlovsky, P. Upscaled CTAB-based DNA extraction and real-time PCR assays for *Fusarium culmorum* and *F. graminearum* DNA in plant material with reduced sampling error. *Int. J. Mol. Sci.* **2008**, *9*, 2306–2321, doi: <https://doi.org/10.3390/ijms9112306>.
57. Alexander, N.J.; McCormick, S.P.; Waalwijk, C.; van der Lee, T.; Proctor, R.H. The genetic basis for 3-ADON and 15-ADON trichothecene chemotypes in *Fusarium*. *Fungal Genet Biol* **2011**, *48*, 485–495, doi:10.1016/j.fgb.2011.01.003.
58. Gang, G.; Miedaner, T.; Schuhmacher, U.; Schollenberger, M.; Geiger, H.H. Deoxynivalenol and nivalenol production by *Fusarium culmorum* isolates differing in aggressiveness toward winter rye. *Phytopathology* **1998**, *88*, 879–884, doi:[10.1094/PHYTO.1998.88.9.879](https://doi.org/10.1094/PHYTO.1998.88.9.879).
59. Mateo, J.J.; Mateo, R.; Hinojo, M.J.; Llorens, A.; Jiménez, M. Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains. *J. Chromatogr. A* **2002**, *955*, 245–256, doi:10.1016/S0021-9673(02)00214-5.
60. Thompson, J.D.; Higgins, D.G.; Gibson, T.J. CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **1994**, *22*, 4673–4680, doi:10.1093/nar/22.22.4673.
61. Beule, L.; Lehtsaar, E.; Rathgeb, A.; Karlovsky, P. Crop diseases and mycotoxin accumulation in temperate agroforestry systems. *Sustainability* **2019**, *11*, 1–21, doi:10.3390/su11102925.
62. Wenzl, T.; Haedrich, J.; Schaechtele, A.; Robouch, P.; Stroka, J. Guidance document on the estimation of LOD and LOQ for measurements in the field of contaminants in feed and food; EUR 28099, Publications Office of the European Union, Luxembourg, **2016**, ISBN 978-92-79-61768-3; doi:10.2787/8931.