

Review

Building Consensus around the Assessment and Interpretation of Symbiodiniaceae Diversity

Davies SW^{1,*}, Gamache MH², Howe-Kerr LI³, Kriefall NG¹, Baker AC⁴, Banaszak AT⁵, Bay LK⁶, Bellantuono AJ⁷, Bhattacharya D⁸, Chan CX⁹, Claar DC¹⁰, Coffroth MA¹¹, Cunnning R¹², Davy SK¹³, del Campo J¹⁴, Díaz-Almeyda EM¹⁵, Frommlet JC¹⁶, Fuess LE¹⁷, González-Pech RA¹⁸, Goulet TL¹⁹, Hoadley KD²⁰, Howells EJ²¹, Hume BCC²², Kemp DW²³, Kenkel CD²⁴, Kitchen SA²⁵, LaJeunesse TC¹⁸, Lin S²⁶, McIlroy SE²⁷, McMinds R²⁸, Nitschke MR⁶, Oakley CA¹³, Peixoto RS²⁹, Prada C³⁰, Putnam HM³⁰, Quigley KM³¹, Reich HG³⁰, Reimer JD³², Rodriguez-Lanetty M⁷, Rosales SM³³, Saad OS³⁴, Sampayo EM³⁵, Santos SR³⁶, Shoguchi E³⁷, Smith EG³⁸, Stat M³⁹, Stephens TG⁸, Strader ME⁴⁰, Suggett DJ⁴¹, Swain TD⁴², Tran C⁴³, Traylor-Knowles N⁴, Voolstra CR²², Warner ME⁴⁴, Weis VM⁴⁵, Wright RM⁴⁶, Xiang T³⁸, Yamashita H⁴⁷, Ziegler M⁴⁸, Correa AMS^{3*} and Parkinson JE^{2,*}

¹Department of Biology, Boston University, Boston, MA, USA

²Department of Integrative Biology, University of South Florida, Tampa, FL, USA

³Department of BioSciences, Rice University, Houston, TX, USA

⁴Department of Marine Biology and Ecology, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL, USA

⁵Unidad Academica de Sistemas Arrecifales, Universidad Nacional Autonoma de Mexico, Puerto Morelos, Mexico

⁶Reef Recovery, Adaptation, and Restoration, Australian Institute of Marine Science, Townsville, QLD, Australia

⁷Department of Biological Sciences, Florida International University, Miami, FL, USA

⁸Department of Biochemistry and Microbiology, Rutgers University, New Brunswick, NJ, USA

⁹Australian Centre for Ecogenomics, School of Chemistry and Molecular Biosciences, University of Queensland, Brisbane, Australia

¹⁰Hakai Institute, Campbell River, Canada

¹¹Department of Geology, University at Buffalo, Buffalo, NY, USA

¹²Daniel P. Haerther Center for Conservation and Research, John G. Shedd Aquarium, Chicago, IL, USA

¹³School of Biological Sciences, Victoria University of Wellington, Wellington, New Zealand

¹⁴Biodiversity Program, Institut de Biologia Evolutiva, Universitat Pompeu Fabra, Barcelona, Spain

¹⁵Department of Natural Sciences, New College of Florida, Sarasota, FL, USA

¹⁶Centre for Environmental and Marine Studies, Department of Biology, University of Aveiro, Aveiro, Portugal

¹⁷Department of Biology, Texas State University, San Marcos, TX, USA

¹⁸Department of Biology, Pennsylvania State University, University Park, PA, USA

¹⁹Department of Biology, University of Mississippi, Oxford, MS, USA

²⁰Department of Biological Sciences, University of Alabama, Tuscaloosa, AL, USA

²¹National Marine Science Centre, Faculty of Science and Engineering, Southern Cross University, Coffs Harbour, NSW, Australia

²²Department of Biology, University of Konstanz, Konstanz, Germany

²³Department of Biology, University of Alabama at Birmingham, Birmingham, AL, USA

²⁴Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA

²⁵Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA, USA

²⁶Department of Marine Sciences, University of Connecticut, Storrs, CT, USA

²⁷Swire Institute of Marine Science, School of Biological Sciences, The University of Hong Kong, Hong Kong

²⁸Center for Global Health and Infectious Disease Research, University of South Florida, Tampa, FL, USA

²⁹Red Sea Research Center, Division of Biological and Environmental Science and Engineering, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

³⁰Department of Biological Sciences, University of Rhode Island, Kingston, RI, USA

³¹Minderoo Foundation, Perth, WA, Australia

³²MISE, Department of Biology, Chemistry and Marine Sciences, Faculty of Science, University of the Ryukyus, Nishihara, Okinawa, Japan

³³The Cooperative Institute For Marine And Atmospheric Studies, University of Miami, Miami, FL, USA

³⁴Department of Biological Oceanography, Red Sea University, Port-Sudan, Sudan

³⁵School of Biological Sciences, The University of Queensland, St. Lucia, QLD, Australia

³⁶Department of Biological Sciences, University at Buffalo, Buffalo, NY, USA

³⁷Marine Genomics Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan

³⁸Department of Biological Sciences, University of North Carolina at Charlotte, Charlotte, NC, USA

³⁹School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW, Australia

40Department of Biological Sciences, Auburn University, Auburn, AL USA

41Climate Change Cluster, University of Technology Sydney, Ultimo, NSW, Australia

42Department of Marine and Environmental Science, Nova Southeastern University, Dania Beach, FL, USA

43Department of Biology, University of San Diego, San Diego, CA, USA

44School of Marine Science and Policy, University of Delaware, Lewes, DE, USA

45Department of Integrative Biology, Oregon State University, Corvallis, OR, USA

46Department of Biological Sciences, Smith College, Northampton, MA, USA

47Fisheries Technology Institute, Japan Fisheries Research and Education Agency, Ishigaki, Okinawa, Japan

48Department of Animal Ecology & Systematics, Justus Liebig University Giessen, Giessen, Germany

Abstract

Within microeukaryotes, genetic and functional variation sometimes accumulate more quickly than morphological differences. To understand the evolutionary history and ecology of such lineages, it is key to examine diversity at multiple levels of organization. In the dinoflagellate family Symbiodiniaceae, which can form endosymbioses with cnidarians (*e.g.*, corals, octocorals, sea anemones, jellies), other marine invertebrates (*e.g.*, sponges, molluscs, flatworms), and protists (*e.g.*, foraminifera), molecular data have been used extensively over the past three decades to describe phenotypes and to make evolutionary and ecological inferences. Despite advances in Symbiodiniaceae genomics, a lack of consensus among researchers with respect to interpreting genetic data has slowed progress in the field and acted as a barrier to reconciling observations. Here, we identify key challenges regarding the assessment and interpretation of Symbiodiniaceae genetic diversity across three levels: species, populations, and communities. We summarize areas of agreement and highlight techniques and approaches that are broadly accepted. In areas where debate remains, we identify unresolved issues and discuss technologies and approaches that can help to fill knowledge gaps related to genetic and phenotypic diversity. We also discuss ways to stimulate progress, in particular by fostering a more inclusive and collaborative research community. We hope that this perspective will inspire and accelerate coral reef science by serving as a resource to those designing experiments, publishing research, and applying for funding related to Symbiodiniaceae and their symbiotic partnerships.

Keywords: Symbiodiniaceae; genetic diversity; coral symbiosis

Introduction

Dinoflagellates in the family Symbiodiniaceae occupy multiple ecological niches on tropical, subtropical, and temperate reefs, ranging from species that are exclusively free-living to those that form symbioses with marine invertebrates (LaJeunesse et al., 2018). The biology of symbiotic Symbiodiniaceae has been a major research focus due to the integral role these mutualists play in the health of scleractinian corals and other marine invertebrates (Glynn, 1996; Hughes et al., 2017). Although many scleractinian coral species exhibit specificity for particular Symbiodiniaceae (reviewed by Baker, 2003; Hume et al., 2020; Thornhill et al., 2014), some coral species and even individual coral colonies can associate with a diversity of algal symbionts (Baker and Romanski, 2007; Silverstein et al., 2012). Moreover, not all host-symbiont pairings are equally resistant or resilient to stress (Abrego et al., 2008; Berkelmans and Van Oppen, 2006; Hoadley et al., 2019; Howells et al., 2013a; Sampayo et al., 2008), and a change in symbiont community may enhance tolerance to future stress. Thus, efforts to characterize the genetic and functional diversity within Symbiodiniaceae not only advance our fundamental knowledge of the evolution and ecology of microeukaryotes, but also provide insights into

the potential for cnidarian-Symbiodiniaceae partnerships, and ultimately for coral reefs, to respond to rapidly changing environments.


The first Symbiodiniaceae species was formally described by Hugo Freudenthal in 1962 (Freudenthal, 1962). As more associations with these endosymbiotic dinoflagellates were cataloged, the utility of allozymes (Schoenberg and Trench, 1980) and later ribosomal markers (LaJeunesse, 2001; Rowan and Powers, 1991) to distinguish different lineages became apparent. Continued exploration of Symbiodiniaceae diversity through molecular genetics ultimately resulted in a recent systematic revision, delineating at least eleven genera and many species (LaJeunesse et al., 2021, 2018; Nitschke et al., 2020; Pochon and LaJeunesse, 2021). However, despite numerous advances in our ability to resolve Symbiodiniaceae genera, species, populations, and strains (Thornhill et al., 2017), these dinoflagellates continue to pose substantial challenges to diversity assessments (Figure 1). A single square centimeter of host tissue typically contains 1-2 million Symbiodiniaceae cells (Fitt et al., 2000) and these cells may be individuals of a single species or a mixture of multiple species and/or genera (Baker and Romanski, 2007; Coffroth et al., 2010; Kemp et al., 2008; Rowan and Knowlton, 1995; Thornhill et al., 2017, 2006; van Oppen et al., 2005). In addition, each symbiont possesses an expansive genome, including multi-copy genes (Lin, 2011), and some Symbiodiniaceae genomes feature extensive gene duplication (González-Pech et al., 2021). Many approaches to resolve Symbiodiniaceae taxonomy rely on multi-copy gene markers and data from such markers may need to be treated differently than data derived from single-copy genes. Since the 2000s, these factors have contributed to an active debate regarding the interpretation of molecular and functional diversity (Apprill and Gates, 2007; Correa and Baker, 2009; E. J. Howells et al., 2016; Pochon et al., 2012; Rodriguez-Lanetty, 2003; Stat et al., 2012, 2008; Thornhill et al., 2007; Wham and LaJeunesse, 2016; Wirshing and Baker, 2016). Due to its accessibility, many in the community have defaulted to using the internal transcribed spacer 2 (ITS2) region of Symbiodiniaceae rDNA to differentiate lineages despite growing recognition that ITS2 data straddle intergenomic and intragenomic variation (the latter of which is abbreviated as IGV), limiting its utility for some applications (Smith et al., 2017). Debate regarding the interpretation of molecular data has also contributed to underuse of other molecular markers, even though they may be more appropriate in some contexts (LaJeunesse and Thornhill, 2011; Takishita et al., 2003).

Low Complexity *Ex Hospite*

Example:
Isoclonal Culture

1 Genus
1 Species
1 Strain

No other
strains present




Low Complexity *In Hospite*

Example:
Acropora cervicornis

1 Dominant Genus
1 Dominant Species
1 Dominant Strain

Background strains
present




High Complexity *Ex Hospite*

Example:
Water Column

2+ Genera
2+ Species
2+ Strains

Free-living &
symbiotic
strains
present




High Complexity *In Hospite*

Example:
Orbicella faveolata

2+ Codominant Genera
2+ Codominant Species
2+ Codominant Strains

Background strains
present



Genus Species MLG Symbol Strain

X	1	α	●	X1α
X	1	β	⬡	X1β
X	2	γ	⬢	X2γ
X	2	δ	⬣	X2δ
Y	3	ε	■	Y3ε
Y	3	ζ	◆	Y3ζ
Y	4	η	▲	Y4η
Y	4	θ	▼	Y4θ

Free-Living Strains: ☆ ☆ ☆ ☆

Notes

MLG = Multi-Locus Genotype. Typically assessed using microsatellite markers.

Strain = Unique MLG nested within a species. Equivalent to an "individual" (isoclonal lineage)

Background Strains = Low-abundance members of the Symbiodiniaceae community. Typically <1% or <10%.

(Co)Dominant Strains = High-abundance member(s) of the Symbiodiniaceae community. Typically 10-50% for codominance, and >50% for dominance.

For visual simplicity, only one Symbiodiniaceae cell per host cell is depicted above. However, different cnidarians vary in the average number of Symbiodiniaceae cells per host cell.

Figure 1 A representation of the various degrees of complexity in Symbiodiniaceae genetic diversity among different habitats (e.g., cultures, corals, and the water column). Communities of Symbiodiniaceae within a given sample can encompass multiple strains, populations, species, and genera.

The increasing popularity of amplicon-sequencing methods (Arif et al., 2014; Green et al., 2014; Howe-Kerr et al., 2020; Hume et al., 2019; Quigley et al., 2014), exploration of additional molecular markers (Pochon et al., 2019, 2012; Smith et al., 2020; Takabayashi et al., 2004), and incorporation of whole-genome datasets (Dougan et al., 2022; González-Pech et al., 2021; Liu et al., 2018) have led to novel insights into

Symbiodiniaceae diversity. Novel challenges in assessing this diversity have also been identified. For example, most non-ITS2 genetic loci exhibit differential utility across Symbiodiniaceae genera (Pochon et al., 2014). Furthermore, different analytical pipelines and thresholds applied to the same marker(s) among studies have led to different estimates of genetic variation and interpretation of their functional importance (Cunning et al., 2017; Howells et al., 2016; Wham and LaJeunesse, 2016; Wirshing and Baker, 2016). These issues have further fueled scientific debate around which markers to use and how to interpret the resulting data.

Recognizing that debate has complicated the process of scientific inquiry, we sought to identify areas of consensus regarding the assessment and interpretation of Symbiodiniaceae genetic diversity. Sixty-one scientists from 12 countries, spanning expertise in the taxonomy, physiology, genomics, and ecology of Symbiodiniaceae and other marine microbes, participated in a workshop funded by the National Science Foundation titled “Building consensus around the quantification and interpretation of Symbiodiniaceae diversity,” held virtually in July 2021. The overall aim was to reduce barriers to those designing experiments, publishing research, and applying for funding related to Symbiodiniaceae and their partnerships. The major workshop outcomes are summarized herein, though not exhaustively. We highlight techniques that are broadly accepted by many experts in the field and point out caveats and considerations for these approaches (Box 1). Where agreement was not reached, we identify the key issues that remain unresolved and point to technologies that might help fill knowledge gaps so that consensus can be achieved in the future. We conclude with suggestions for how to make the Symbiodiniaceae research community a more inclusive and welcoming space that promotes innovation as we navigate the coral reef crisis. Above all, we wish to stress that the choice of genetic marker(s) and analytical framework(s) for interpreting Symbiodiniaceae diversity will always depend on the research question at hand, along with the availability of resources (*e.g.*, for sample preservation, processing, and computation), and that these options will inevitably evolve as our understanding of the system continues to develop.

Box 1: Major Workshop Outcomes and Consensus Highlights

General

- a. Different research questions require different levels of resolution of Symbiodiniaceae diversity (*e.g.* species, populations, communities).
- b. Molecular markers evolve at different rates and vary in their ability to resolve different Symbiodiniaceae taxonomic levels and lineages, requiring careful selection of the appropriate marker(s) for a given question.
- c. Many markers and analytical approaches are available, each with their own strengths and weaknesses. As genomic resources for Symbiodiniaceae continue to be developed and technologies advance, so will options for analyzing and interpreting diversity.
- d. Collaborations among research groups can ameliorate methodological and analytical disconnect within the Symbiodiniaceae community, while also reducing costs associated with answering complex and integrative research questions.

1. Species-Level Assessment of Symbiodiniaceae Diversity

- a. Resolving Symbiodiniaceae to the species level is important. Species identification forms the basis of comparative physiological, ecological, and evolutionary investigations within Symbiodiniaceae.
- b. A robust Symbiodiniaceae taxonomy is required to facilitate scientific communication, link past and future research, and establish legal frameworks for conservation. Funding to develop and maintain up-to-date public taxonomic tools and databases should be increased.
- c. There are distinctions between *describing* a new species, *recognizing* a new species, and *identifying* a known species. Reef researchers benefit from incorporation and consideration of the current taxonomy whenever possible.
- d. Expanding publicly accessible Symbiodiniaceae culture collections and their formal genetic, morphological, and physiological description will drive taxonomic, ecological, physiological, and genomic research. Supporting these resources for use by the scientific community should be a priority for long-term federal funding.

2. Population-Level Assessment of Symbiodiniaceae

- a. Population-level studies evaluate the distribution of genetic variation within Symbiodiniaceae species, often across spatiotemporal gradients or among host taxa, to understand the influence of evolutionary processes such as gene flow, genetic drift, and natural selection.
- b. When multiple Symbiodiniaceae lineages are present within host colonies, population-level questions are more challenging to address.
- c. Pre-screening to determine which lineages are present within samples is necessary to determine the marker(s) needed to address population-level questions in Symbiodiniaceae.
- d. Microsatellite loci can be effective at addressing population-level questions in Symbiodiniaceae if used appropriately.
- e. The ITS2 region of Symbiodiniaceae rDNA can be an effective marker for distinguishing between different populations, but requires thorough validation with multilocus analysis to distinguish intra- and inter- genomic variation.

3. Community-Level Assessment of Symbiodiniaceae

- a. Symbiodiniaceae communities can be conceptualized at different scales. The presence of two or more Symbiodiniaceae species within a host individual constitutes a “local community.” Total Symbiodiniaceae diversity at larger scales (*e.g.*, among conspecific host colonies, multiple host species, or across environments including free-living Symbiodiniaceae) constitutes a “macroscale community.” The total diversity of both local and macroscale communities is likely underestimated.
- b. Local Symbiodiniaceae communities are often composed of representatives of different genera, rather than multiple species/lineages within the same genus.
- c. Marker genes that exhibit inter- and intra-genomic variation (as well as variation in copy number across lineages) make it challenging to characterize Symbiodiniaceae community composition. Quantifying this molecular variation for Symbiodiniaceae genera and species is a priority.
- d. The Symbiodiniaceae ITS2 marker can be useful for describing Symbiodiniaceae communities but there are circumstances where multiple markers or other approaches may be more appropriate. A majority of researchers at the workshop reported greatest familiarity and comfort with the ITS2 marker, which may have contributed to its popularity in characterizing Symbiodiniaceae communities.
- e. There is a lack of consensus regarding best practices for interpreting Symbiodiniaceae gene amplicon data to identify species, and for applying and interpreting community diversity metrics. Authors are encouraged to clearly highlight assumptions associated with their data interpretation, acknowledge that other interpretations exist, and discuss whether or not alternative interpretations change the biological or ecological findings of their study.

4. Beyond Genotype: Phenotyping Symbiodiniaceae

- a. Phenotypic diversity varies greatly within and between Symbiodiniaceae species, thus it is critical to avoid overestimating the functional significance of a given symbiont based on taxonomic assignment alone (*e.g.*, assuming that all *Durisdinium* spp. are heat-tolerant).
- b. There is a need to develop technologies to functionally assay Symbiodiniaceae in culture, *in hospite*, and in the environment—and to better contextualize the resulting phenotypes—with the understanding that functional diversity will vary depending on the metrics used.
- c. When attempting to understand phenotypic variability among strains and species, using cultures of Symbiodiniaceae can help control confounding variables. However, because cultures are artificial environments, performance *in vitro* may differ from performance *in hospite*.

5. Integrating Multiomic Technologies to Study Symbiodiniaceae

- a. Various 'omics techniques have been used to address Symbiodiniaceae taxonomic, functional, and physiological research questions. Because each technique has unique considerations, leveraging these novel tools requires stringent ground-truthing and the development of quality standards.
- b. Genome projects have improved tremendously over the past decade, but there are unique biological obstacles that have restricted Symbiodiniaceae genome assembly quality. Examples include large genome sizes, high repeat content, and

difficulty annotating gene functions. Integrating multiple techniques, such as transcriptomics and proteomics, and coupling these with phenotyping methods, can help answer outstanding questions regarding Symbiodiniaceae-host interactions. Efficient experimentation will require combining expertise across laboratories.

6. Ensuring an Inclusive Symbiodiniaceae Research Community

- a. Critical examination is at the heart of scientific inquiry. A diversity of perspectives has always been and will continue to be needed to move the Symbiodiniaceae field forward.
- b. The publication process should be equitable. Recommendations include increasing diversity on relevant journal editorial boards, scaling publication costs for researchers employed in countries with lower income economies, implementing double-blind review, and citing articles led by diverse colleagues.
- c. Parachute science should be avoided. Recommendations include fostering long-term international collaborations and exchange programs to involve local scientists in Symbiodiniaceae research, improving sensitivity to the challenges facing colleagues in funding-limited partner institutions, and extending full collaborative benefits including authorship and grant writing opportunities to these colleagues.
- d. Accessibility and collaboration should be fostered. Recommendations include establishing a voluntary database of all Symbiodiniaceae researchers and their research products, maintaining hybrid format options for conferences, and expanding long-term federal funding to include more international collaborations.
- e. It is critical to improve recruitment, retention, and promotion of scholars of diverse backgrounds. Recommendations include working actively to increase diversity at all levels of academia and science, promoting the work of minority scientists, and providing strong multidimensional mentorship to support and retain these scientists throughout each career stage.

1. Guidance for Species-Level Assessment of Symbiodiniaceae Diversity


1.1 Why is species-level resolution important for Symbiodiniaceae?

Species are evolutionarily independent lineages and therefore represent a fundamental level of biological organization. Species-level resolution provides insight into the ecological and evolutionary mechanisms that create diversity, and forms the basis of comparative physiological investigations (Kareiva and Levin, 2015). The delineation of species can affect nearly all scales of inquiry, from biochemical pathways to ecosystem processes. Species-level diversity in Symbiodiniaceae has been discussed in the literature since the description of *Symbiodinium microadriaticum* in 1962 by Hugo Freudenthal (Freudenthal, 1962). As more diversity was uncovered and more species were recognized (LaJeunesse, 2001; LaJeunesse and Trench, 2000; Rowan and Powers, 1992, 1991; Schoenberg and Trench, 1976; Trench and Blank, 1987), controversy arose as to where to draw species boundaries (Apprill and Gates, 2007; Correa and Baker, 2009; Cuning et al., 2013; LaJeunesse et al., 2014, 2012; Stat et al., 2012; Thornhill et al., 2007; Wham and LaJeunesse, 2016). At present, there is general consensus among Symbiodiniaceae specialists about the need for species-level resolution, as well as support for current taxonomic methodologies that are underpinned by genetic, ecological, and morphological data (LaJeunesse et al., 2018; Voolstra et al., 2021b). Such taxonomic descriptions facilitate scientific communication and are necessary for establishing legal frameworks for conservation (IUCN, 2021). The recent elevation of Symbiodiniaceae “Clades” to genera has provided some clarity (LaJeunesse et al., 2021, 2018; Nitschke et al., 2020; Pochon and LaJeunesse, 2021), but the paucity of formal species-level descriptions for the large genetic diversity found within most Symbiodiniaceae genera constitutes a formidable barrier to progress.

Without robust species delineation, functional differences can inadvertently be ascribed to incorrect taxonomic levels or non-existent biological entities. For example, the genus level may be too coarse and lead to over-generalizations regarding the physiology or function of Symbiodiniaceae variants (see Section 4). A statement such as “*the genus Cladocopium consists of heat-sensitive species*” overlooks the superior stress tolerance of some *Cladocopium* species (Abrego et al., 2008), including the dominance of *Cladocopium thermophilum* in corals on some of the world’s hottest reefs in the Persian/Arabian Gulf (Hume et al., 2015). However, diversity assessments based on gene sequence variants may recover both interspecific variation (resolving distinct species) and intraspecific variation (sequence diversity within a single genome). This is a major issue for the multi-copy rDNA internal transcribed spacer 2 (ITS2) gene, one of the most commonly used diversity markers for Symbiodiniaceae. Consequently, a statement such as “*Symbiodiniaceae harboring the ITS2 D13 sequence variant are adapted to temperate environments*” overlooks the fact that ITS2 sequence variants D8, D8-12, D12-13, and D13 are all characteristic of the same species, *Durusdinium eurythalpos*, which is found in mutualisms with Oulastreidae corals across temperate and tropical regions (LaJeunesse et al., 2014). Such issues are problematic because they obfuscate ecological interpretations of sequence data.

1.2 What types of data can identify Symbiodiniaceae species?

The Biological Species Concept dictates that if two organisms cannot reproduce and create viable offspring, they should be considered different species (Mayr, 1942). Unfortunately, this criterion cannot be applied to Symbiodiniaceae, as no direct observation of sexual reproduction has been made to date (but see Figueroa et al., 2021; Shah et al., 2020). Fortunately, many other species concepts exist, each placing emphasis on different criteria (De Queiroz, 2007; Leliaert et al., 2014). Robust species descriptions satisfy multiple species concepts using independent lines of evidence. For Symbiodiniaceae, the field has largely applied three key types of data: morphological (cell size and cell wall features), ecological (host specificity and biogeographic distribution), and phylogenetic (divergence across multiple DNA markers), along with the assignment of type material and strains (Figure 2). The taxonomic framework for describing species has matured since the earliest effort by Freudenthal (Freudenthal, 1962). For example, in line with the Morphological Species Concept, Trench and Blank (Trench and Blank, 1987) proposed three new species based on Symbiodiniaceae cell ultrastructure. They used transmission electron microscopy (TEM) to reveal features such as the nucleus, chromosomes, pyrenoid, chloroplast thylakoid membranes, and cell size; additionally, they used scanning electron microscopy (SEM) to observe thecal plates and the arrangement of the two flagella. Technological advancements in SEM resolution now enable complete morphological characterization of amphiesmal vesicles in the cell wall (Jeong et al., 2014; LaJeunesse et al., 2015; Lee et al., 2020, 2015; Nitschke et al., 2020), though such plate tabulations tend to be variable within species (LaJeunesse et al., 2018).



*Biological evidence in the Gray Zone may arise with different timing and order than depicted.

Figure 2 A simplified representation of Symbiodiniaceae speciation, species concepts (SC), and associated biological evidence. In this example, one ancestral species splits and diverges to become two descendant lineages after barriers to gene flow are established. Through selection and drift, these lineages evolve different properties, which satisfy the criteria of different species concepts (represented by horizontal lines). Because these properties may arise at different times and in different orders, there is a “gray zone” where conflict among species concepts may result in controversy about whether one or two species exist. Modified from de Queiroz (2007) and Leliart *et al.* (2014).

As an increasing number of host species are sampled, it has become clear that the Ecological Species Concept can also be used to support Symbiodiniaceae species descriptions. Although not diagnostic in all cases (Cunning *et al.*, 2013), symbiosis ecology can be particularly useful for Symbiodiniaceae species that exhibit host-specificity or coadaptation with their hosts (Davies *et al.*, 2020; Finney *et al.*, 2010; Howells *et al.*, 2020; Santos *et al.*, 2004; Smith *et al.*, 2017; Thornhill *et al.*, 2014). For example, *Cladocopium pacificum* and *Cladocopium latusorum* are found exclusively within corals of the genus *Pocillopora* (Turnham *et al.*, 2021). Ultimately, because Symbiodiniaceae do not always have distinct morphological characteristics, nor do they always exhibit host specificity, the collection of genetic data to satisfy a Phylogenetic Species Concept has

also become a necessity in the description of species (see Section 1.3).

Although taxonomic descriptions are fundamental, *describing* a new species is not the same as *recognizing* a new species or *identifying* a known species. The former should be based on multiple lines of evidence, while the latter may require generating and interpreting data from only one or two diagnostic methods. Incorporating existing taxonomy (*i.e.*, species names) into current research whenever possible is encouraged. Due to a general lack of funding for taxonomic descriptions, formal species names are not always available for a given entity, and therefore accommodating sequence variant terminology in the literature will continue to be important. Providing synonyms (*e.g.*, the ITS2 sequence variant and its species name) when a species is first mentioned will improve clarity. Resources that consolidate current and past taxonomic assignments are critical for connecting historical and future research (Voolstra et al., 2021b).

Species description projects can be extremely fruitful for researchers who are new to the field and who are seeking to increase their knowledge of the system. There are six major components to creating a valid species description: 1) collecting information on at least two congruent genes (see our recommendations below in Section 1.3), 2) comparing genetic data against that from other Symbiodiniaceae, 3) comparing cell size measurements against those from other Symbiodiniaceae, 4) providing cell images (simple light microscopy is sufficient), 5) depositing type specimens in a permanent archive (*e.g.*, herbarium or museum), and 6) proposing a valid name (according to the International Code of Nomenclature for Algae, Fungi, and Plants; Turland et al., 2018). Those that do not endeavor to describe Symbiodiniaceae species can encourage and incentivize those that do by accurately treating taxa names as hypotheses and citing the work of taxonomists at the first mention of previously described taxa within manuscripts.

1.3 How can we resolve Symbiodiniaceae species with genetic markers?

No single marker is likely able to distinguish species across all Symbiodiniaceae genera reliably (Table 1). Instead, ecological and physiological studies will benefit from adopting a multi-gene approach (see Section 3). Congruence among sequence data from different cellular compartments (nuclear, chloroplast, and mitochondrial; Table 1) indicates that classifying Symbiodiniaceae using a lineage-based species concept is achievable (De Queiroz, 2007; LaJeunesse and Thornhill, 2011; Sampayo et al., 2009). This multi-gene approach, supported with ecological, morphological, and sometimes physiological data, has led to the formal description (or re-validation) of 39 Symbiodiniaceae species in 11 genera thus far (Table 1; Hume et al., 2015; Jeong et al., 2014; LaJeunesse, 2017; LaJeunesse et al., 2021, 2018, 2015, 2014, 2012; Lee et al., 2020; Lewis et al., 2019; Nitschke et al., 2020; Parkinson et al., 2015; Pochon and LaJeunesse, 2021; Ramsby et al., 2017;

Turnham et al., 2021; Wham et al., 2017; Xiang et al., 2013). This taxonomic list will continue to grow, and recent whole-genome data already point toward the need for further revision of some genera and species (Dougan et al., 2022; González-Pech et al., 2021).

Table 1 Current list of formally described Symbiodiniaceae species and associated diagnostic information. Mito = Mitochondrial; Chloro = Chloroplast; R = Resolves all species within the genus; D = Diagnostic (uniquely differentiates a particular species of the genus); ND = Not diagnostic (sequence/trait identical in two or more species); M = Measured but lacking congeners or reference material for comparison; X = Not used in species description; U = Unknown (*e.g.*, sampled from a symbiotic habitat but not necessarily likely to be the numerically dominant symbiont); Y = Yes; N = No; NA = Not Applicable; ND* = Not diagnostic of species, but lack of elongated amphiesmal vesicles is diagnostic of *Cladocopium*; D** = Some ITS2 sequences may be diagnostic, but others in the in the same genome may not be; Y*** = Opportunistic and occurring at background levels unless host health is compromised. Note: for an extended version of the table that includes synonyms, key references, and authentic cultured strains for each species, see Table S1.

Genus (11 total)	Species (39 total)	ITS2 Type (majority sequence)	Symbiotic?	Cultured?	Host-specificity	Cell Size	Plate Tabulation	Nuclear			Mito		Chloro	
								ITS2	LSU	msat flanks	mtCOB	mtCOXI	cp23S	psbA ^{scr}
<i>Breviolum</i>	<i>B. aenigmaticum</i>	-	U	Y	U	ND	X	D	D	D	ND	X	D	X
<i>Breviolum</i>	<i>B. antillogorgium</i>	B1	Y	Y	D	ND	X	ND	ND	D	ND	X	D	X
<i>Breviolum</i>	<i>B. dendrogryum</i>	B1k, B1	Y	N	D	ND	X	D**	ND	D	ND	X	ND	D
<i>Breviolum</i>	<i>B. endomadracis</i>	B7	Y	N	D	ND	X	D	ND	D	D	X	ND	D
<i>Breviolum</i>	<i>B. favinorum</i>	B14, B14a, B1	Y	N	ND	ND	X	D**	ND	D	ND	X	ND	D
<i>Breviolum</i>	<i>B. meandrinum</i>	B20, B1	Y	N	ND	ND	X	D**	ND	D	ND	X	ND	D
<i>Breviolum</i>	<i>B. minutum</i>	B1	Y	Y	ND	ND	X	ND	ND	D	ND	X	ND	D
<i>Breviolum</i>	<i>B. pseudominutum</i>	B1	U	Y	U	ND	X	ND	ND	D	D	X	ND	X
<i>Breviolum</i>	<i>B. pygmaophilum</i>	B2	Y	Y	ND	ND	X	D	D	D	ND	X	D	X
<i>Cladocopium</i>	<i>C. goreauii</i>	C1	Y	Y	ND	Y	ND*	ND	ND	X	D	D	ND	D
<i>Cladocopium</i>	<i>C. infistulum</i>	C2	Y	Y	D	Y	ND*	D	D	X	D	D	D	X
<i>Cladocopium</i>	<i>C. latusorum</i>	C1b-c, C42a, C42a-b, C1c-ff	Y	N	D	Y	X	D	D	D	ND	X	X	D
<i>Cladocopium</i>	<i>C. pacificum</i>	C1d, C1d-t	Y	N	D	Y	X	D	D	D	ND	X	X	D
<i>Cladocopium</i>	<i>C. thermophilum</i>	C3	Y	N	ND	U	X	D	ND	X	D	D	ND	D
<i>Durusdinium</i>	<i>D. glynnii</i>	D1, D1-4-6	Y	N	ND	ND	X	D**	ND	D	ND	X	ND	D
<i>Durusdinium</i>	<i>D. boreum</i>	D15	Y	N	ND	ND	X	D	D	D	D	X	ND	X
<i>Durusdinium</i>	<i>D. eurythalpos</i>	D8, D12-13, D13	Y	N	ND	ND	X	D	D	D	ND	X	ND	X
<i>Durusdinium</i>	<i>D. trenchii</i>	D1a, D1-4, D1-4-6	Y	Y	ND	ND	X	D**	ND	D	ND	X	D	D
<i>Effrenium</i>	<i>E. voratum</i>	E1	N	Y	NA	M	M	M	M	X	M	X	M	M
<i>Freudenthalidium</i>	<i>Fr. endolithicum</i>	F3.8	U	Y	X	D	ND	R	R	X	R	X	R	X
<i>Freudenthalidium</i>	<i>Fr. heronense</i>	F3.7	U	Y	X	D	ND	R	R	X	R	X	R	X
<i>Fugacium</i>	<i>Fu. kawagutii</i>	F1	U	Y	U	M	X	M	M	X	X	X	X	X
<i>Gerakladium</i>	<i>G. endoclonum</i>	-	Y	N	D	ND	X	X	D	X	ND	X	R	R
<i>Gerakladium</i>	<i>G. spongolum</i>	-	Y	N	D	ND	X	X	ND	X	ND	X	R	R
<i>Halluxium</i>	<i>H. paucillum</i>	H7	U	Y	X	M	M	R	R	X	ND	X	R	X
<i>Miliolidium</i>	<i>M. leei</i>	D1.1	U	Y	U	M	X	R	R	X	R	R	X	X
<i>Philozoon</i>	<i>P. actiniarum</i>	A19	Y	N	D	D	X	D	ND	X	D	ND	ND	R
<i>Philozoon</i>	<i>P. adriaticum</i>	-	Y	N	D	D	X	ND	D	X	ND	ND	ND	R
<i>Philozoon</i>	<i>P. anthopleurum</i>	-	Y	N	D	D	X	ND	ND	X	ND	ND	ND	R
<i>Philozoon</i>	<i>P. balanophyllum</i>	-	Y	N	D	D	X	ND	ND	X	ND	D	ND	R
<i>Philozoon</i>	<i>P. colozsum</i>	-	Y	N	D	D	X	ND	D	X	ND	D	D	R
<i>Philozoon</i>	<i>P. geddesianum</i>	-	Y	N	D	D	X	ND	ND	X	ND	D	D	R
<i>Philozoon</i>	<i>P. medusarum</i>	-	Y	N	D	D	X	ND	ND	X	ND	ND	ND	R
<i>Philozoon</i>	<i>P. paranemonium</i>	-	Y	N	D	D	X	ND	ND	X	ND	D	ND	R
<i>Symbiodinium</i>	<i>S. microadriaticum</i>	A1	Y	Y	ND	ND	ND	D	D	X	D	X	ND	D
<i>Symbiodinium</i>	<i>S. natans</i>	-	N	Y	U	ND	ND	D	D	X	D	X	D	X
<i>Symbiodinium</i>	<i>S. necroappetens</i>	A13	Y***	N	ND	ND	ND	D	D	X	D	X	ND	D
<i>Symbiodinium</i>	<i>S. pilosum</i>	A2	N	Y	U	ND	X	D	D	X	D	X	D	X
<i>Symbiodinium</i>	<i>S. tridaenidorum</i>	A6, A3a, A3*	Y	Y	ND	ND	ND	D**	D	X	D	X	D	X

The rate of evolution of gene markers dictates their respective power to resolve distinct genetic entities and whether these entities are likely to represent distinct species (Table 1). In addition, genetic differentiation may vary among genera for the same marker region (Pochon et al., 2014). Efforts are underway to develop a taxonomic key for Symbiodiniaceae species based on genetic and ecological data, this is a long-term process whereby several hundred currently identified ITS2 sequence variants require individual evaluation with subsequent species assignments. We envision a dynamic dichotomous key that would guide users to the appropriate markers and characteristics for a particular host organism of interest, or alternatively, suggest combinations of markers and characteristics most likely to provide species-level resolution within specific sets of closely related Symbiodiniaceae.

1.4 How many Symbiodiniaceae species exist?

The current best estimate for the total number of symbiotic Symbiodiniaceae species is in the range of hundreds based on phylogenetic (*e.g.*, ITS2) sequence variants (Thornhill et al., 2014). However, these species numbers are likely a significant underestimate because sampling efforts have mainly focused on scleractinian coral hosts living at shallow depths in tropical and subtropical waters. It will be important to continue describing Symbiodiniaceae species in non-scleractinian hosts, including other cnidarians; *e.g.*, octocorals (Goulet et al., 2017; Ramsby et al., 2014), zoantharians (Fujiwara et al., 2021; Mizuyama et al., 2020), actinarians (Grajales et al., 2016), corallimorpharians (Kuguru et al., 2008), hydrocorals (Rodríguez et al., 2019), jellyfish (Vega de Luna et al., 2019); as well as sponges (Hill et al., 2011; Ramsby et al., 2017), acoelomorph flatworms (Kunihiro and Reimer, 2018), molluscs (Baillie et al., 2000; Banaszak et al., 2013; Lim et al., 2019), ciliates (Mordret et al., 2016), and foraminifera (Pochon et al., 2007). Further collections from undersampled habitats and sources such as benthic sediment and rubble (Fujise et al., 2021; Nitschke et al., 2020; Takabayashi et al., 2012), seagrasses and macroalgae (Porto et al., 2008; Yamashita and Koike, 2013), mesophotic depths (Frade et al., 2008; Goulet et al., 2019), the water column (Manning and Gates, 2008; Pochon et al., 2010), and predator feces (Castro-Sanguino and Sánchez, 2012; Grupstra et al., 2021; Parker, 1984) will likely yield many undiscovered species and possibly even novel genera (Yorifuji et al., 2021). These efforts should not be limited to subtropical and tropical waters, as Symbiodiniaceae have been reported in more temperate locations (LaJeunesse et al., 2021; Lien et al., 2012). Systematic and wide-ranging effort to better describe the genetic diversity of Symbiodiniaceae (such as the Tara Oceans expedition; Sunagawa et al., 2020) will lead to a better understanding of the drivers of taxonomic and functional diversity of Symbiodiniaceae.

1.5 What steps can be taken to enhance our understanding of Symbiodiniaceae species?

Expanding publicly accessible Symbiodiniaceae culture collections can drive not only taxonomic but also ecological, physiological, and genomic research (LaJeunesse et al., 2018; Voolstra et al., 2021b; Xiang et al., 2013). Most of the diversity in culture constitutes just a handful of species, predominantly from the *Symbiodinium* and *Breviolum* genera. More targeted and consistent funding to support further development, maintenance, and sharing of culture collections are critical to the field. Progress toward protocols for Symbiodiniaceae cryopreservation would help conserve biodiversity through the generation of cryogenic archives (Di Genio et al., 2021) and support research in laboratories that cannot maintain continuous cultures. Depositing live specimens in national and organizational archives can alleviate the burden on individual research groups. Examples of national archives include the Provasoli-Guillard National Center for Marine Algae and Microbiota at Bigelow Laboratory in the USA, (<https://ncma.bigelow.org/>), the Symbiont Culture Facility at the Australian Institute of Marine Science in Australia (<https://www.aims.gov.au/>), the National Institute for Environmental Studies, (<https://mcc.nies.go.jp/>) and Biological Resource Center at National Institute of Technology and Evaluation (<https://www.nite.go.jp/nbrc/catalogue/>) in Japan, the Central Collection of Algal Cultures in Germany (<https://www.uni-due.de/biology/ccac/>), the Roscoff Culture Collection in France (<https://roscoff-culture-collection.org/>), and the Culture Collection of Algae and Protozoa in the United Kingdom (<https://www.ccap.ac.uk/>).

Live cultures can benefit taxonomic studies by providing relatively homogeneous strains to establish baselines of diversity and morphology. Molecular data from cultured isoclonal strains are less diverse than those from host tissues, which may contain multiple Symbiodiniaceae genera, species, or strains (Figure 1; Voolstra et al., 2021b). Cultures are also superior for holotype depositions, and they facilitate morphometric analysis, for example, on swimming behavior (motility). However, live culture is not a prerequisite for formal species description, especially because many Symbiodiniaceae are currently culture recalcitrant (Krueger and Gates, 2012). Furthermore, many strains cultured from host tissue do not represent the dominant Symbiodiniaceae in a host species (Santos et al., 2001). Efforts toward testing new media and bringing new species into culture, as well as documenting and sharing successful and failed attempts are encouraged. "Culturability" itself may be a useful phenotype to track, as it may reflect the degree of host-specificity, and influence media choice (Ishikura et al., 2004; Nitschke et al., 2020; Reimer et al., 2010; Yorifuji et al., 2021). Motility, cell division rates (growth), bacterial communities (microbiomes) and viral consortia (viromes) are also informative characteristics that can vary within and among species (Grupstra et al., 2022a; Lawson et al., 2018; Levin et al., 2017; Parkinson and Baums, 2014; Yamashita and Koike, 2016). Constructing a global phenotypic

database for cultures, much like the Coral Trait Database (Madin et al., 2016) is another priority for Symbiodiniaceae research.

Finally, it would be advantageous to identify and culture model Symbiodiniaceae lineages to test species boundaries. For example, measuring DNA sequence differences between sibling species separated by a geological barrier (*e.g.*, the Isthmus of Panama; LaJeunesse et al., 2018; Pochon et al., 2006) would provide molecular-divergence cutoffs that could then be applied to better resolve sympatric lineages. Additionally, cultures of closely related, putative sibling species could be used to explore cytological evidence for sexual recombination (Figueroa et al., 2021), evaluate potential hybridization (Brian et al., 2019), and characterize the role symbiotic interactions play in genome evolution (González-Pech et al., 2019).

2. Guidance for Population-Level Assessment of Symbiodiniaceae

2.1 How can we design population-level studies?

Studies evaluating the distribution of genetic variation within species, often across spatiotemporal gradients or among host taxa, seek to understand how populations are influenced by evolutionary processes such as gene flow, genetic drift, and selection (Aichelman and Barshis, 2020; Davies et al., 2020; Forsman et al., 2020; Prada et al., 2014; Reich et al., 2021; Thornhill et al., 2017; Turnham et al., 2021). Here, we define a population as a group of individuals belonging to the same species that live and interbreed with each other in a given space and time. The study of Symbiodiniaceae populations is fundamental to improving the resolution at which phenotypes of interest are differentiated. Thus, this section focuses on allele-based identification and quantification of genetic variation.

Because a single host can contain a mixture of multiple species and/or genera, a first step in experimental design should include assessing sample sets for the presence of multiple distinct Symbiodiniaceae that may confound the interpretation of population-level genetic variation (see Section 3). Such assessment can be done pre- and post-population-level analysis with established genetic markers (*e.g.*, ITS2, *cp23S*) and may be guided by published literature for some regions or host species. Pre-screening is especially advantageous where information on the community composition of Symbiodiniaceae is also sought and especially for hosts which tend to associate with multiple genera or species. Quantitative PCR (qPCR) may be one potential technique to pre-screen Symbiodiniaceae samples for the presence of particular lineages (Correa et al., 2009; Mieog et al., 2007; Saad et al., 2020). After pre-screening, population-level studies typically target genetic variation from the numerically dominant symbiont genus associating with a particular host or set of hosts (Baums et al., 2014), while excluding any confounding genetic variation from additional species that may be

present within host samples (Baums et al., 2010; Thornhill et al., 2006). Post-screening of samples is also possible using tests of assignment to genetic clusters (Davies et al., 2020) or identifying and excluding samples with outlier allelic profiles. Post-screening may be more time- and cost-effective as verification can be performed on a subset of the total sample set.

The ideal number of samples to collect and analyze will depend on the particular aims of the study (*e.g.*, delineating populations versus characterizing the degree of admixture among them), and the scale of comparison (*e.g.*, reef, habitat, colony, intra-colony, etc.), and the markers being employed. However, studies leveraging more traditional markers, such as microsatellites, tend to benefit from robust sample sizes with minimum ranges of 20-30 individual hosts *per* level of interest (*e.g.*, habitat and location) (Hale et al., 2012). Although this is a good target, studies limited by permit authorizations, budgets, and other constraints are still informative in some contexts.

2.2 How can we best use microsatellite loci?

Microsatellite loci (or simple sequence repeats; SSRs) are segments of DNA where 1-6 base pairs are repeated in a tandem array; these loci are distributed abundantly across genomes of nearly all eukaryotic organisms (Tautz, 1989). Variations in the length of repeats are generated by polymerase slippage during DNA replication, resulting in homologous regions (*i.e.*, loci) of differing lengths (*i.e.*, alleles) among individuals. Microsatellites are generally thought to represent neutral loci with high mutation rates. Their single-locus, multiallelic, and codominant properties can yield valuable information regarding ploidy and reveal genetic structure among populations within and between species. Furthermore, microsatellite analyses are generally a PCR-based technique, making them cost-effective relative to other methods. With the advent of high-throughput sequencing and transcriptomics, the generation of hundreds of potential microsatellite loci is now comparatively straightforward. For species or lineages where numerous loci are available, costs and effort can remain low by multiplexing primer sets (Davies et al., 2013). Taken together, these features make microsatellites attractive for studying Symbiodiniaceae populations. These markers have been used to address questions related to overall diversity, population structure within and between reefs, gene flow, dispersal, and relatedness between symbionts (see Table 1 in Thornhill et al., 2017).

Once the target Symbiodiniaceae species or lineage has been identified within a dataset of interest, these samples can be tested for variability using previously developed microsatellite loci *via* PCR amplification (Figure 3). Primers for such loci have been developed for Symbiodiniaceae species across at least five genera: *Symbiodinium* (Pinzón et al., 2011), *Breviolum* (Andras et al., 2011; Grupstra et al., 2017; Pettay and

LaJeunesse, 2007; Santos et al., 2001; Santos et al., 2003; Wirshing et al., 2013), *Cladocopium* (Bay et al., 2009; Davies et al., 2020; Howells et al., 2009; Magalon et al., 2006; Wham and LaJeunesse, 2016), *Durusdinium* (Pettay and LaJeunesse, 2009; Wham et al., 2011), and *Philozoon* (Molecular Ecology Resources Primer Development Consortium et al., 2010). Importantly, these loci tend to have narrow phylogenetic ranges, with primers developed for a given species typically working only on other closely-related species within the same genus. Therefore, it is necessary to screen existing primers for utility with a given target species, to ensure that allelic variability among the chosen suite of microsatellite loci is sufficient, and to develop novel primer sets if existing primers fail or prove insufficiently specific. Microsatellite development has historically been costly and time-consuming. However, with the advent of "genome skimming by shotgun sequencing", where genome sequencing of thousands of short reads identifies microsatellites for development (e.g., Abdelkrim et al., 2009), microsatellite development is now relatively inexpensive and rapid. Ideally, new Symbiodiniaceae primers should be tested against monoclonal cultures of species within the same genus (positive controls) as well as against symbiont-free sperm or apo-symbiotic larvae (negative controls) to rule out off-target PCR amplification of host DNA. Although more loci will generally increase discriminatory power in population-level studies, as few as 2-3 loci have provided sufficient discriminatory power for some questions (Santos et al., 2003; Thornhill et al., 2009).




Figure 3 Recommendations for designing microsatellite-based Symbiodiniaceae population genetics experiments. Sample collection, marker choice, and analytical pipeline should be considered from the outset.

2.2.1 Analyses of Microsatellite Data: Given that Symbiodiniaceae are haploid in their vegetative life stage (Santos and Coffroth, 2003), a single allele *per* microsatellite locus is expected when a host harbors a single clonal strain of Symbiodiniaceae (represented by a single multi-locus genotype; MLG). When a single allele is recovered from nearly all loci, establishing MLGs is straightforward. However, recovery of multiple alleles at a given locus from a single sample is not uncommon (Figure 1). Instances of multiple alleles *per* locus can be interpreted as detection of cells from multiple clonal strains (multiple MLGs) within host tissues (Andras et al., 2009; Grupstra et al., 2017; Santos et al., 2003; Santos and Coffroth, 2003; Thornhill et al., 2017, 2009). Examples of multiple MLGs tend to be more common within Indo-Pacific corals hosting *Cladocopium* species (Bay et al., 2009; Davies et al., 2020; Wham et al., 2014), whereas they are less common in Caribbean corals hosting *Cladocopium* and other genera (Andras et al., 2009; Grupstra et al., 2017; Pettay et al., 2015; Santos et al., 2003; Santos and Coffroth, 2003; Thornhill et al., 2017, 2014a, 2009). Consistent patterns of multiple alleles for certain loci among a subset of monoclonal cultures has led to the proposal of whole or segmental genome duplication within certain Symbiodiniaceae. This scenario would make overestimation of symbiont genotype diversity within samples likely (Wham et al., 2014), and make the assignment of MLGs difficult, raising challenges for data analyses and interpretations.

Several approaches have been developed to accommodate instances of multiple MLGs within a sample (Figure 3; Andras et al., 2011; Davies et al., 2020; Howells et al., 2013b; Kirk et al., 2009; Magalon et al., 2006; Wham and LaJeunesse, 2016), including the exclusion of some samples/genotypes in certain cases. When multiple alleles for a given locus occur infrequently among samples, two data sets can be created: 1) a set where all microsatellite alleles within each sample are used and scored for presence or absence (*i.e.*, binary) within each sample, and 2) a curtailed data set omitting samples with multiple alleles at one or more loci, allowing MLGs to be assigned. Notably, studies using this approach have come to similar conclusions across the two data sets (*e.g.*, Andras et al., 2011; Davies et al., 2020; Howells et al., 2013b; Kirk et al., 2009; Magalon et al., 2006; Wham and LaJeunesse, 2016). In general, given that reported scales of genetic divergence are similar across studies using binary and MLG-based approaches, and excluding many samples can lead to underestimating genetic diversity (Howells et al., 2016), we suggest that the binary approach should be used when possible (*e.g.*, a high proportion of samples exhibit multiple alleles *per* locus).

2.2.2 Caveats: While microsatellite analyses have proven informative and valuable in population genetic studies of Symbiodiniaceae, they present challenges in data acquisition and interpretation. For example, the long repetitive regions of microsatellites are often difficult to reliably amplify, making it arduous to verify repeat length *via* fragment analysis. Microsatellites can suffer from allele dropout, and low specificity of PCR

primers, which can potentially lead to diversity underestimates within a sample. Microsatellites themselves are subject to more general criticisms including unclear mutation models and the potential for homoplasy (Putman and Carbone, 2014). Additionally, many analytical pipelines used to assess population genetic patterns make basic assumptions that Symbiodiniaceae do not follow (*e.g.*, that organisms are diploid and exhibit predominantly sexual reproduction). In light of this, researchers should be cautious about interpreting results from pipelines developed for organisms that exhibit more traditional population biology.

2.3 What other markers can resolve Symbiodiniaceae populations?

The ITS2 region of rDNA is repeated in tandem arrays within all known Symbiodiniaceae genomes. For population-level assessments, this universality presents an advantage over microsatellites, but the multi-copy nature of this marker poses unique challenges. As long as appropriate analytical frameworks are applied (see Section 3), ITS2 data can be used to resolve within species. Such assessments require consideration of similarities in the assemblages of ITS2 sequences and their relative abundances within each genome. For example, genetic structure among *Cladocopium thermophilum* strains in the Persian/Arabian Gulf has been characterized (Hume et al., 2019; Smith et al., 2017) and patterns of IGV obtained from amplicon sequencing data show fine-scale spatial structure among *C. thermophilum* populations separated by tens to hundreds of kilometers (Howells et al., 2020). However, recombination (*i.e.*, whether two populations are interbreeding) is often considered sufficient for operational recognition that those entities are members of the same species (Andras et al., 2009; Grupstra et al., 2017; Santos et al., 2003; Santos and Coffroth, 2003; Thornhill et al., 2017, 2009). Therefore, it is not possible to determine whether ITS2-based genotypes correspond to distinct populations of the same species or different species. Other markers are also able to resolve at the population-level, but their application to Symbiodiniaceae population biology is limited. Examples include the chloroplast *psbA* minicircle noncoding region (*psbA^{ncr}*; Moore et al., 2003), the chloroplast 23S ribosomal region (*cp23S*; Santos et al., 2003), and the mitochondrial cytochrome *b* region (*cob*; Takabayashi et al., 2004).

2.4 What are the next steps for understanding Symbiodiniaceae population biology?

Advancing our understanding of Symbiodiniaceae population biology will be greatly informed by leveraging samples that have single Symbiodiniaceae MLGs (Prada et al., 2014). For example, available monoclonal cultures of Symbiodiniaceae from several species could be leveraged to develop and test new technologies and markers (including validation of copy number, see Section 3.3.5) and these technologies could then be extended to more complex associations *in hospite* (within a host organism). For example, Frommlet and Iglesias-Rodriguez (2008) developed a method for multi-locus microsatellite genotyping of single cells in a free-living dinoflagellate species, providing a novel tool for high-resolution population genetic studies. To avoid the challenges of widespread gene duplication in Symbiodiniaceae genomes (González-Pech et al.,

2021; Pochon et al., 2012; Prada et al., 2014), efforts should be directed toward identifying new low copy markers (or preferably single copy markers). Discovery of single copy loci may be informed by screening for universal single copy markers collated in the Benchmarking Universal Single-Copy Orthologs (BUSCO) database (Seppey et al., 2019; Simão et al., 2015), although many BUSCOs are undetected in Symbiodiniaceae genomes (González-Pech et al., 2021). Restriction-associated DNA sequencing may serve as a low-cost method for generating single copy markers for population-level assessments in Symbiodiniaceae (Kitchen et al., 2020; Suyama and Matsuki, 2015); however, these methods require further development.

Whole-genome sequencing (WGS) is also becoming more affordable, especially at low coverage (< 5X), opening the possibility of evaluating genome-wide variation in Symbiodiniaceae (González-Pech et al., 2021; Reich et al., 2021), although Symbiodiniaceae genomes are large (> Gbp) and few chromosome-scale assemblies exist (Marinov et al., 2021; Nand et al., 2021). We suggest that WGS should first be applied to isoclonal cultures, where possible, to ensure reads derive from one genetic entity (*e.g.*, Voolstra et al., 2021a). Subsequently, this approach can be applied to multispecies assemblages where different Symbiodiniaceae lineages within the same genus could be mapped to these reference genomes. These types of analyses would allow for simultaneous quantification of gene flow and divergence among Symbiodiniaceae populations of co-occurring species and improve estimates of effective population sizes and clonality within and among species, hosts, and reefs. Another major advantage of genome-wide data is the potential to evaluate adaptive (non-neutral) genetic variation and signatures of selection across the genome. For example, identifying associations between traditional markers, genomic regions, Symbiodiniaceae functional traits, and/or environmental variables—including those that are important for the survivorship of corals under warmer, more acidic, and more eutrophic oceans—remains a research priority (see Sections 4 and 5).

3. Guidance for Community-Level Assessment of Symbiodiniaceae

3.1 What is a Symbiodiniaceae community?

Generally defined, ecological communities are composed of more than one species that live together and interact. However, what is meant by terms such as “together” and “interact” can vary (Konopka, 2009), particularly when considering free-living *versus* symbiotic Symbiodiniaceae. Typically one to two (but up to ten) Symbiodiniaceae cells reside in a coral gastrodermal cell (Davy et al., 2012; Muscatine et al., 1998), potentially restricting direct interactions between the endosymbiont cells within a coral host. Here, we use the term “local Symbiodiniaceae community” to refer to two or more Symbiodiniaceae species *within a single host*, whereas “macroscale Symbiodiniaceae community” (“phenomenological community” in Konopka, 2009) describes the total diversity of Symbiodiniaceae across some larger scale (*e.g.*, conspecific hosts or

multiple host species). Environments that include multiple free-living Symbiodiniaceae species also constitute macroscale communities; *e.g.*, benthic sediments (Nitschke et al., 2016; Quigley et al., 2017a), the water column (Fujise et al., 2021; Porto et al., 2008), and macro-algal surfaces (Fujise et al., 2021; Porto et al., 2008).

Macroscale Symbiodiniaceae communities contain more species and encompass higher genetic diversity than local Symbiodiniaceae communities because symbiotic diversity accumulates with increased host colony and habitat sampling (Swain et al., 2020). Environmental samples include cells of symbiotic Symbiodiniaceae expelled from hosts as well as non-symbiotic, free-living species. In contrast, a given adult host colony typically harbors only one or two dominant Symbiodiniaceae species (Goulet, 2006), often from distinct genera, as well as other species at low relative abundances (Hume et al., 2020; Silverstein et al., 2012). *In hospite* Symbiodiniaceae communities can be transmitted vertically (promoting higher fidelity), reassembled horizontally (allowing for greater flexibility), or some combination of both (mixed-mode transmission) with each host generation (Quigley et al., 2017b). The diversity of the free-living component of macroscale Symbiodiniaceae communities and the symbiotic component of local Symbiodiniaceae communities are each likely to be underestimated (*e.g.*, Baker and Romanski, 2007), but for different reasons. Free-living communities are relatively diffuse and are therefore more difficult to exhaustively sample. In contrast, local Symbiodiniaceae community assessments are prone to sampling bias (but see, *e.g.*, Goulet and Coffroth, 2003). Characterizations of local communities are often based on a single sample from a well-lit, “top” surface of a colony. Sampling across a host’s surface has revealed heterogeneous distributions of dominant Symbiodiniaceae within colonies of Caribbean stony corals such as *Colpophyllia*, *Montastraea*, *Orbicella*, *Porites*, and *Siderastrea* (*e.g.*, Correa et al., 2009; Kemp et al., 2008; Rowan et al., 1997; Ulstrup and van Oppen, 2003), as well as some Pacific stony corals (*e.g.*, Correa et al., 2009; Fifer et al., 2022; Innis et al., 2018; Kemp et al., 2008; Rowan et al., 1997; Ulstrup and van Oppen, 2003) and zoantharians such as *Zoanthus* (Fujiwara et al., 2021) and *Palythoa* (Wee et al., 2021). Whether local Symbiodiniaceae communities exhibit structure over smaller spatial scales *in hospite* (*e.g.*, oral versus aboral host surfaces) is unknown, but could be resolved with single-cell techniques (see Section 5).

3.2 Why study Symbiodiniaceae community diversity?

Studying macroscale communities can provide insights into cnidarian-Symbiodiniaceae dynamics along environmental gradients (Cunning et al., 2015; Rossbach et al., 2021; Silverstein et al., 2011; Terraneo et al., 2019). Regional macroscale Symbiodiniaceae community structure (*i.e.*, beta diversity) may also reflect chronic disturbance from anthropogenic activity (Claar et al., 2020a) and help identify more resilient or resistant reefs (Ziegler et al., 2015). Additionally, macroscale communities in reef seawater, sediments, feces,

and on macro-algal surfaces may be important sources of symbiotic Symbiodiniaceae that can be acquired horizontally by prospective hosts (Adams et al., 2009; Ali et al., 2019; Castro-Sanguino and Sánchez, 2012; Coffroth et al., 2006; Cumbo et al., 2013; Fujise et al., 2021; Granados-Cifuentes et al., 2015; Grupstra et al., 2022b, 2021; Nitschke et al., 2016; Porto et al., 2008; Quigley et al., 2018, 2017a; Umeki et al., 2020; Venera-Ponton et al., 2010). Symbiodiniaceae in a free-living mode may influence important processes, such as sexual reproduction, hybridization, and gene flow within Symbiodiniaceae (Figueroa et al., 2021).

Positive and negative species interactions can occur within local Symbiodiniaceae communities resulting in resource and niche partitioning (Davy et al., 2012; Howe-Kerr et al., 2020; Matthews et al., 2020). Quantifying these interactions may help disentangle the factors and processes governing Symbiodiniaceae community assembly in early host life history stages (McIlroy et al., 2019; Quigley et al., 2016), as well as successional dynamics (or stability) in adult hosts. Studying local Symbiodiniaceae communities can also identify conditions that trigger symbiotic breakdown (*i.e.*, dysbiosis). Dysbiosis has frequently been documented in the bacterial communities of stressed hosts (*e.g.*, Zaneveld et al., 2017; Ziegler et al., 2017), and may also be evident in local Symbiodiniaceae communities. Generally speaking, dysbiosis can manifest itself in the host as: 1) an increase in symbiont richness (invasion or proliferation of low abundance symbionts), 2) a decrease in symbiont richness (loss of symbionts), or 3) more complex community changes in community structure or beta diversity (Egan and Gardiner, 2016). For example, *Symbiodinium necroappetens* (LaJeunesse et al., 2015; Stat et al., 2008) and some symbionts in the genera *Durusdinium* (Bay et al., 2016; Manzello et al., 2018), *Breviolum* (LaJeunesse et al., 2010b), and *Cladocopium* (Wee et al., 2021) can opportunistically increase or decrease their abundance in bleached or stressed hosts. Stony coral juveniles in the field (Quigley et al., 2016) and adults in tank-based experiments (Howe-Kerr et al., 2020) have exhibited decreased survival in conjunction with more diverse local Symbiodiniaceae communities. Additional experiments to assess how frequently different types of dysbiosis occur in local Symbiodiniaceae communities are needed, including in non-scleractinian hosts, some of which can harbor up to 60 symbionts *per* host cell (Fitt, 2000). Testing the extent to which different types of dysbiosis are associated with specific cnidarian hosts, as well as specific environmental contexts, should also be prioritized.

Current challenges in understanding local Symbiodiniaceae community diversity and dynamics include: 1) determining actual and relative abundances of Symbiodiniaceae species given IGV and copy number issues (see Section 3.3.5); and 2) understanding the roles (if any) that low abundance Symbiodiniaceae play in holobiont survival and fitness (see Arif et al., 2014; Bay et al., 2016; Lee et al., 2016). This knowledge is key to connecting Symbiodiniaceae genotypes to phenotypes (see Section 4). Low abundance Symbiodiniaceae

may serve as a reservoir of *in hospite* algal genotypes that may increase to dominance (at least ephemerally) during or following a change in environmental conditions (Bay et al., 2016; Berkelmans and Van Oppen, 2006; Boulotte et al., 2016; Buddemeier and Fautin, 1993; Claar et al., 2020a; Jones et al., 2008; Lewis et al., 2019; Thornhill et al., 2006; Ziegler et al., 2018). The mechanisms controlling this turnover *in hospite* remain poorly understood, but involve host rewards/sanctions (Kiers et al., 2011, 2003) and competitive interactions among symbionts (Palmer et al., 2003). Competition among Symbiodiniaceae affects the initial uptake of symbionts in early coral ontogeny (McIlroy et al., 2019) and influences longer-term persistence in experimentally-generated symbioses (Gabay et al., 2019), but the relative importance of competition in shaping *in hospite* communities once they are established remains poorly understood. Beyond their potential to shift *in hospite* following bleaching events (Jones et al., 2008; Thornhill et al., 2006), low abundance Symbiodiniaceae could also contribute to emergent holobiont properties (suggested by Howe-Kerr et al., 2020; Ziegler et al., 2018). Quantification of holobiont traits with and without the addition of low abundance homologous (*i.e.*, lineages that typically enter into a symbiotic relationship with a given host taxon) Symbiodiniaceae from a range of inoculation sources constitutes a critical next step to understanding the functional role these symbionts play in the host.

3.3 How can we optimize the study of Symbiodiniaceae community diversity?

Improving our understanding of the processes shaping Symbiodiniaceae communities is critical to predicting their distributions and potentially mitigating the trajectories of coral reefs experiencing global change. The methods below constitute suggested approaches for analyzing diversity of macroscale and local Symbiodiniaceae communities. In some circumstances, identifying numerically dominant Symbiodiniaceae lineages (as opposed to the total diversity of a Symbiodiniaceae community) may be sufficient for the question at hand, since hosts are generally selective in the symbionts they harbor, and some are highly specific to particular symbiont lineages (*e.g.*, Hume et al., 2020; Thornhill et al., 2014). Whether quantifying numerically dominant or total Symbiodiniaceae diversity, molecular marker selection, as well as the approach to data generation and analyses, have implications for the interpretation of Symbiodiniaceae community diversity. Molecular markers available for assessing Symbiodiniaceae community diversity are multicopy, and thus, present the challenge of distinguishing intragenomic from intergenomic variation. Inclusion of symbiont taxa above or below the species level in community diversity analyses can obscure patterns and lead to under- or over-estimation of diversity metrics.

3.3.1 Markers that behave as if single copy: The Symbiodiniaceae *SSU* and *LSU* rDNA markers as well as the *cob* mitochondrial marker are multicopy but are considered to behave like single copy loci because the vast majority of copies present are a single sequence. The few intragenomic sequence differences that do occur

tend to be relatively straightforward to resolve in the context of identifying the dominant Symbiodiniaceae lineage within each genus. Many analysis algorithms produce amplicon sequence variants (ASVs), which are statistically inferred based on sequence variation within and among samples; the degree to which ASVs represent distinct genotypes may vary by marker and Symbiodiniaceae genus. For example, *LSU* consistently resolves species within *Symbiodinium* (Lee et al., 2015), but not for all of *Breviolum* (Table 1; Parkinson et al., 2015). Thus, it is important to keep in mind that when assessing total community diversity (across multiple Symbiodiniaceae genera) with *LSU*, the number of species within certain genera may be under-represented. Despite this, markers that behave as if single copy are arguably the best option currently available for assessing total community diversity in Symbiodiniaceae.

3.3.2 Multicopy markers: Among the commonly used markers, the hypervariable chloroplast *psbA* non-coding region (*psbA^{ncr}*) consistently resolves below the species level in Symbiodiniaceae (LaJeunesse et al., 2015; Lewis et al., 2019; Turnham et al., 2021; Wham et al., 2017), while the ITS2 region can resolve below, at, or above the species level depending on the lineage. Higher resolution comes at a considerable cost in terms of complexity of analyses.

***psbA^{ncr}*:** The *psbA^{ncr}* region can assess relatedness only among closely related Symbiodiniaceae lineages within the genus (LaJeunesse and Thornhill, 2011; Thornhill et al., 2014). It is helpful to have *a priori* knowledge of the genera being amplified when using this marker (see Section 3.4) as available primers have known biases for specific genera (*e.g.*, the Symbiodiniaceae *psbA^{ncr}* primers 7.4-Forw and 7.8-Rev (Moore et al., 2003) preferentially amplify *Cladocopium* in samples of mixed communities, whereas the more recent *psbAFor_1* and *psbARev_1* do not (LaJeunesse and Thornhill, 2011). Although *psbA^{ncr}* is multi-copy and can exhibit IGV in some species, drawing inferences from these sequence datasets is still relatively straightforward because – similar to markers that do not present IGV – large genetic distances exist even between sequences from closely related species (LaJeunesse et al., 2021). However, because the *psbA^{ncr}* region cannot be amplified across Symbiodiniaceae using a single set of primers, this marker is suboptimal for some types of community-level analyses, such as assessing total community diversity or beta diversity metrics. Nevertheless, it would be appropriate to pair *psbA^{ncr}* with other markers; *i.e.*, to resolve additional diversity within established ITS2 lineages (Noda et al., 2017; Reimer et al., 2017); and also to use this marker to verify ITS2 sequence variants generated *via* amplicon sequencing (Hume et al., 2019; Smith et al., 2020).

***ITS2*:** The ITS2 of Symbiodiniaceae rDNA resolves many species and some subspecies (Hume et al., 2019). ITS2 has a broader application for defining lineages since one set of primers amplifies all known

Symbiodiniaceae ITS2 sequences (note, however, that sequence variants only align well within-genus). These two favorable characteristics, in concert with its history of use within the field, make ITS2 a popular choice among researchers, even in situations when greater resolution might be achieved with alternative marker(s). Intragenomic sequence diversity is relatively high within Symbiodiniaceae ITS2 (Arif et al., 2014) and along with copy number, varies considerably across genera (Saad et al., 2020) and likely species (though no data are currently available at this resolution). This IGV severely restricts the inferences that can be made regarding the relative abundance of community members, in cases of multiple Symbiodiniaceae lineages *per* host (see Section 3.3.5). The central issue in using ITS2 to characterize symbiont diversity *in hospite* is differentiating intragenomic sequence variants (those that reflect differences within one genetic entity) from intergenomic sequence variants (those that reflect differences between two or more genetic entities). This is of particular importance because, unlike with *psbA^{ncr}*, Symbiodiniaceae ITS2 intragenomic distances can be larger than intergenomic distances. Practically, it can be challenging to determine if sequence variation comes from one species or multiple species. Varied awareness and treatment of this issue among Symbiodiniaceae researchers has generated significant debate, which has often played out in peer review, rather than being articulated, addressed, and resolved as a research community (see Section 6).

One technique to differentiate between intra- and inter-genomic sequence variants involves analyzing co-occurrence patterns. Sets of different sequences that co-occur across multiple biological replicates are more likely to be from the same genotype than to derive from multiple co-occurring lineages, with each lineage contributing a subset of the sequences. This is particularly true in cases where the relative abundances of each of the sequences of the set are similar across biological replicates. There are gel-based (Denaturing Gradient Gel Electrophoresis, DGGE; LaJeunesse, 2001) and bioinformatic methods (*e.g.*, Frøslev et al., 2017; Green et al., 2014; Hume et al., 2019) to detect these co-occurring sequences in both dominant and low abundance taxa. Gel-based and *in silico* approaches each have their advantages and disadvantages, which have been discussed elsewhere (Saad et al., 2020). Because these techniques rely on identifying banding profiles that correspond to references (gel-based) or other biological replicates (gel- and bioinformatic-based), their power to resolve generally increases with access to references or further biological replicates. For this purpose, reference sets of DGGE profiles as published in the literature (*e.g.*, LaJeunesse et al., 2010a; LaJeunesse and Thornhill, 2011; Silverstein et al., 2011), or online reference databases of *in silico* profiles (*e.g.*, at symportal.org) are available to researchers. However, strong inferences can still often be made from relatively small datasets for Symbiodiniaceae taxa that are sampled multiple times in the dataset. Both techniques rely on the same biological assumption: that coral hosts commonly associate with one numerically dominant Symbiodiniaceae taxon *per* genus. In cases where this assumption does not hold—when congeneric

Symbiodiniaceae co-occur in multiple biological replicates—diversity may be underestimated with multiple taxa being considered one. Identifying intergenomic and intragenomic variation is necessary for making conclusions about diversity when using multi-copy markers like ITS2. Differentiating between this variation can be challenging, particularly when dealing with less common genotypes, smaller numbers of biological replicates, lower sequencing depths, and complex communities; in these situations, sequencing of the samples in question with an additional marker may be necessary. Critically, such an additional marker must be able to resolve between the putative taxa. For example, if attempting to ascertain whether two closely related *Cladocodium* taxa (e.g., within the C3-radiation) are present in a sample, *psbA^{ncr}* would be more appropriate than *cp23S* as the former is highly likely to resolve between such taxa (Thornhill et al., 2014), whereas the latter may or may not (Pochon et al., 2019).

3.3.3 Assessing total Symbiodiniaceae diversity: When characterizing both dominant and low abundance Symbiodiniaceae *in hospite*, three general considerations need to be made. First, Symbiodiniaceae communities can exhibit spatial structure within an individual host (e.g., Correa et al., 2009; Fifer et al., 2022; Kemp et al., 2008; Rowan et al., 1997). Second, assessment of total Symbiodiniaceae diversity is recommended with high-throughput sequencing or qPCR (genera/species present must be known *a priori* and primers specific to these must be available or designed) as these approaches provide the resolution to detect both dominant and low abundance Symbiodiniaceae. Gel- or Sanger sequencing-based methods can provide qualitative information on diversity, but lack the resolution to detect Symbiodiniaceae present at very low abundances (i.e., <2-11% for restriction fragment length polymorphism-based (RFLP-based) methods, (Correa, 2009); <5-30% for denaturing gradient gel electrophoresis-based (DGGE-based) methods, (LaJeunesse et al., 2009; Lien et al., 2007; Loram et al., 2007). Third, all caveats for specific markers from above still apply (e.g., only diversity that can be resolved can be detected, and PCR biases may occur). Markers that behave as if single copy (e.g., *SSU*, *LSU*, *cob*) are putatively well suited to characterizing total Symbiodiniaceae diversity due to their taxonomic breadth; analyses of total diversity using these markers will often be more straightforward than with *psbA^{ncr}* (or ITS2). Despite this, *psbA^{ncr}* is also a reasonable choice when investigating total Symbiodiniaceae diversity due to its apparent low(er) copy number and intragenomic richness, as long as the community diversity in question does not exceed the taxonomic range of this marker. In these limited circumstances, *psbA^{ncr}* may resolve lineages well because genetic distances among taxa are relatively high with this marker.

When assessing the total diversity of macroscale Symbiodiniaceae communities, it is important to consider how molecular techniques and approaches apply to ‘free-living Symbiodiniaceae’. In the broadest sense, this

term refers to all cells external to metazoan (*e.g.*, coral, mollusc) or protistan (*e.g.*, ciliate, foraminifera) hosts. These cells may be found in the water column or associated with benthic substrates/biofilms. ‘Transiently free-living’ refers to Symbiodiniaceae cells that are recently released from nearby hosts but that are not adapted to proliferate outside of hosts (Yamashita and Koike, 2013). In contrast, ‘exclusively free-living’ refers to Symbiodiniaceae species with lifestyles entirely external to hosts (Jeong et al., 2014). Although exclusively free-living Symbiodiniaceae may be detected occasionally “within” host samples, such detections can be interpreted as contamination resulting from host ingestion (rather than symbiosis establishment) or adherence to mucus (Baker and Romanski, 2007; Lee et al., 2016; Silverstein et al., 2012). Because ‘free-living’ (*sensu lato*) Symbiodiniaceae communities on reefs are complex mixtures of these two categories, resolving this diversity presents specific challenges (Box 2).

Box 2: Characterizing free-living Symbiodiniaceae community diversity

Markers that behave as if they are single copy, as well as multicopy markers, may be applied to answering questions related to free-living Symbiodiniaceae communities. Although most commonly used, primer sets for ITS2 are problematic because they result in non-target amplification of other species (*e.g.*, fungi, other dinoflagellates) present in the reef environments (Hume et al., 2018; Nitschke et al., 2020). Despite this, relative to ITS2, other markers pose greater challenges to assessing free-living Symbiodiniaceae community diversity. Specifically, the *cp23S* marker frequently amplifies non-Symbiodiniaceae plastid-containing taxa when used in free-living systems (Nitschke unpublished data). Additionally, *cp23S*'s relatively coarse taxonomic resolution in some lineages (*e.g.*, *Breviolum*; Parkinson et al., 2015) may not be suited to some research questions. In contrast, *psbA^{ncr}* operates on a narrow taxonomic breadth (see Section 3.3.2). Thus, although there are no issues with non-target amplifications by *psbA^{ncr}*, multiple primer pairs would be required to amplify across all species of Symbiodiniaceae likely to be of interest; some of these primer pairs have yet to be developed.

ITS2 has its own challenges for assessing free-living Symbiodiniaceae communities because the process of looking for sets of sequences that co-occur among samples as a proxy for collapsing intragenomic variants (*e.g.*, Hume et al., 2019) is not a valid approach for free-living Symbiodiniaceae. This is because in the free-living environment, multiple Symbiodiniaceae species per genus are likely to be present in a single sample. A number of strategies exist to alleviate this problem. First, free-living Symbiodiniaceae communities, while interesting for their novel diversity, are likely to be studied alongside symbiotic Symbiodiniaceae on the same reef. For example, (Fujise et al., 2021) studied coral symbionts from the C15 and C3 radiations of *Cladocopium* and generated ITS2 defining intragenomic variant (DIV) profiles, or informative assemblies of within-sample intragenomic sequences (see (Hume et al., 2019) for details), for each. These sets of ITS2 DIV sequences were then searched for in water, macroalgae, and sediment samples from the same reef. Complete sets of sequences from the C15 and C3 profiles were successfully retrieved from water and macroalgae, however in sediments only partial DIV profiles were retrieved alongside a greater representation of sequences from additional genera (*e.g.*, *Symbiodinium*, *Freudenthalidium*, *Gerakladium*, and *Halluxium*). It is not possible to differentiate whether these partial profiles in sediments represent novel *Cladocopium* diversity not present in corals or other hosts, or if sequencing depth was exhausted due to the greater representation of diversity across the family. A second approach, analogous to the first, leverages the high culturability of Symbiodiniaceae from free-living environments (Hirose et al., 2008; Nitschke et al., 2020; Yamashita and Koike, 2013). Of 263 Symbiodiniaceae-like single cells isolated from sands of the same reef examined by (Fujise et al., 2021), 114 successfully established as novel cultures belonging to the family Symbiodiniaceae (Nitschke et al., 2020). ITS2 sequences of these isoclonal cultures were later used by (Fujise et al., 2021) as reference sequences and exact matches were found within the free-living communities. Again, both of these strategies rely upon building definitive sets of ITS2 sequences from Symbiodiniaceae cells of (ideally) a clonal population of a single strain within a single species, and then querying for these ITS2 sequence sets within communities of greater complexity.

Prior to the advent of high-throughput sequencing techniques, multiple markers were PCR amplified, cloned, and Sanger sequenced when examining free-living Symbiodiniaceae communities. For example, ITS2 and the short hypervariable region of *cp23S* (*cp23S-HVR*) have been used to study Symbiodiniaceae communities in the water column, sediments, and in stony corals in Hawaii and the Caribbean (Manning and Gates, 2008; Pochon et al., 2010). The *cp23S-HVR* primers were selected for their high specificity for Symbiodiniaceae; although the amplicons produced by these primers are of a size amenable to high-throughput sequencing workflows (~140 bp), this sequencing approach is not cost effective for these primers because the gene region appears to have less resolving power than ITS2 (Pochon et al., 2010; Santos et al., 2003). New, low copy number markers that resolve diversity at or

below the level of ITS2 are needed to study the diversity of free-living Symbiodiniaceae communities. *psbA^{ncr}* has yet to be applied to free-living communities in a high-throughput approach, but this gene region is an obvious candidate.

3.3.4 Assessing beta diversity: Beta diversity can be useful for measuring changes to Symbiodiniaceae community structure over space and time (Eckert et al., 2020; Epstein et al., 2019). Although beta diversity encompasses a range of metrics including dissimilarity, turnover, nestedness, and dispersion, it is dispersion that is most commonly used to assess Symbiodiniaceae communities (Arif et al., 2014; Claar et al., 2020b; Cunning et al., 2017; Green et al., 2014; Howe-Kerr et al., 2020; Hume et al., 2019; Quigley et al., 2014). An important consideration when analyzing Symbiodiniaceae beta diversity data is establishing whether the analysis focuses on *sequence* beta diversity (e.g., amplicon sequence variant data, which typically encompass copy number and intragenomic variability below the species level), or whether the analysis focuses on *ecological* beta diversity (e.g., species data). Either approach may be viable, but it is important to explicitly state which is being used, and to frame interpretations based on the potential pitfalls relevant to that approach.

3.3.5 Accounting for copy number variation: Copy number variation (CNV) is any genetic trait involving the number of copies of a gene in the genome of an individual. Efforts to quantify the absolute and relative abundances of different species in a local Symbiodiniaceae community are complicated by the presence of high CNV across taxa for key markers such as ITS2 (Correa et al., 2009; Mieog et al., 2007; Saad et al., 2020; Stat et al., 2006; Thornhill et al., 2007). Thus, relative rDNA-PCR amplicon abundance does not necessarily equate to actual abundance of Symbiodiniaceae cells in a sample (especially when inter-genus comparisons are being made; Arif et al., 2014; Correa et al., 2009; Quigley et al., 2014). For instance, some symbiont taxa that have been reported to possess a considerably higher rDNA copy number than others (e.g., *Cladocopium* spp.; Saad et al., 2020); these high copy number taxa can appear to be abundant in mixed communities even though they might represent a low fraction of cells *in hospite*, leading to inaccurate estimation of actual symbiont abundances (Figure 4). The incorrect classification of low abundance *versus* dominant taxa can impact interpretations related to biogeography and ecology. Such errors could be avoided if a correction factor is applied (e.g., dividing the abundance value by the number of copies present in the genome of the relevant species, Figure 4; Correa et al., 2009; Mieog et al., 2007; Rubin et al., 2021; Saad et al., 2020), but such corrections rely on accurate copy number reference values, which are not currently available for the majority of Symbiodiniaceae taxa. Most studies that have quantified CNV have been limited to comparisons between genera, and there is considerable variation in the values reported across studies (e.g., Gong and Marchetti, 2019; Loram et al., 2007; Mieog et al., 2007; Quigley et al., 2014; Saad et al., 2020; Thornhill et al., 2007). Inconsistencies in reported CNV values may be attributed to variation among strains or species (within-genus differences can be as large as between-genus differences) or to methodological differences between studies.

Therefore, the extent of CNV within Symbiodiniaceae genera or across populations largely remains to be established.

Lineage-specific qPCR assays have helped to quantitatively characterize mixed communities at the genus (Correa et al., 2009; Cunning et al., 2018; Cunning and Baker, 2013) and species (Fujiwara et al., 2021) levels. These targeted qPCR assays are more quantitative than sequencing approaches, but still require correction for CNV. When applied to systems with known symbiont diversity, qPCR can accurately and cost-effectively quantify local symbiont community structure and dynamics; however, these primer sets must be developed on a *per-taxon* basis. Another approach is the use of flow cytometry to quantify and/or physically separate cells of interest. While the natural variability in cell characteristics (e.g., size, shape, fluorescence) cannot distinguish taxa (Apprill et al., 2007), the use of fluorescent probes to tag taxa of interest has successfully quantified the absolute and relative abundance of co-occurring taxa (McIlroy et al., 2020, 2014). Importantly, these methods are also conducive to subsequent genetic and physiological analyses of sorted cells. The development of further resources to account for CNV is an important priority within the field (see Section 5).




Figure 4. An example of ITS2 rDNA copy number variation (CNV) between the genomes of two Symbiodiniaceae species from different genera (*Breviolum minutum* and *Cladocopium goreauai*). Bar graphs demonstrate how original, uncorrected values (lighter bars) can lead to inaccurate perceptions regarding the proportional representation and numerical dominance of a species. In this case, raw *C. goreauai* ITS2 counts need to be divided by ~10 to correct for CNV (darker bars). Modified from Saad *et al.* (2020).

3.4 When should researchers use multiple Symbiodiniaceae genetic markers for community-level analyses?

As each marker has its own evolutionary history and methodological bias (e.g., primer bias, CNV, *etc.*), congruence among multiple independent markers should enable more robust characterization of Symbiodiniaceae community ecology (Fujiwara et al., 2021; Kavousi et al., 2020; Noda et al., 2017; Pochon

et al., 2019; Reimer et al., 2017; Smith et al., 2020, 2017). Where possible, when multiple markers are used, the markers should complement each other's taxonomic scope and power to resolve. For example, it may be productive to pair the *cp23S* (larger taxonomic scope, lower resolving power) with the *psbA^{ncr}* (smaller taxonomic scope, higher resolving power). Instances where multiple markers produce conflicting results may help identify a Symbiodiniaceae lineage that cannot be accurately characterized with a single broad taxonomic marker. Additionally, combining multiple markers can provide greater resolution and improved interpretability compared to single-marker approaches. For example, the use of *psbA^{ncr}* can help overcome issues associated with interpretation of IGV and high copy number in ITS2 and provide support for ITS2-type profiles (Smith et al., 2020) by confirming which ITS2 IGVs are most likely part of a single lineage (LaJeunesse and Thornhill, 2011; Smith et al., 2017). Budget and logistics permitting, it is recommended to use multiple markers in some situations. For some ITS2 lineages (e.g., *Breviolum* B1 or *Cladocopium* C15; Hoadley et al., 2021; Parkinson et al., 2015), available markers that behave as if single copy do not distinguish ecologically relevant variants, but universal primers have yet to be designed for gene regions that capture this variation (e.g., *psbA^{ncr}*). Therefore, a marker that behaves as if single copy can first be applied to determine the dominant Symbiodiniaceae genera present to assist in the selection of the correct primer set for a higher resolution marker.

3.5 How can we interpret Symbiodiniaceae diversity while acknowledging the pitfalls of common markers?

Given the complexities associated with common methodological approaches and how they influence ecological interpretations of Symbiodiniaceae genetic information in community-level studies, it is critical to provide sufficient methodological details when reporting and interpreting results. We encourage the field to follow reproducible research standards, which include making analysis pipelines and raw data available after publication (see Lowndes et al., 2017) for a comprehensive guide to open science tools). At minimum, commented code (including filtering thresholds, analysis decision points, and processing steps) should be deposited in each article's supplemental materials or in a publicly accessible repository (e.g., GitHub) with a DOI (e.g., procured through GitHub and Zenodo). Raw sequencing data must be deposited in a dedicated archive such as NCBI SRA for NGS-generated data, or NCBI Genbank for single sequence data. Alongside the code and sequences, additional metadata (e.g., environmental and physiological parameters, as well as trackable information regarding the hosts' ID, if applicable; Voolstra et al., 2021b), should be deposited either with the publishing journal or with a data repository (e.g., Dryad, Zenodo, the National Science Foundation's BCO-DMO). Finally, all of these deposition options can be integrated. For example, a Zenodo deposition can be linked to a GitHub repository so that new code releases are automatically updated in the Zenodo repository.


By making published data widely available, we can accelerate our understanding of cnidarian-dinoflagellate symbioses and their responses to a changing environment.

There continues to be dialogue amongst members in the Symbiodiniaceae field regarding the interpretation of gene amplicon data produced by metabarcoding (*e.g.*, Illumina MiSeq). Specifically, there is an ongoing debate about if and how to incorporate Symbiodiniaceae taxa present at low abundances, and how certain parameters are built into existing analytical pipelines. For example, SymPortal will not attempt to predict profiles for a Symbiodiniaceae genus in a given sample if there are less than 200 reads for that genus/sample combination (Hume et al., 2019); this can potentially contribute to systematic underestimation of total Symbiodiniaceae community diversity. As such, we encourage authors to consider carefully what their data can and cannot discern (*e.g.*, Table 1), report assumptions associated with their data interpretation, acknowledge that other interpretations exist, and discuss whether or not these other interpretations change the biological or ecological conclusions of their study. Diversity metrics merit careful attention and gene sequence diversity should not be conflated with species diversity. Authors (and reviewers and editors) can weigh what results, tables, or figures might be included in the Supplementary Material to acknowledge and address these additional interpretations in order to facilitate the inclusion of diverse perspectives (see Section 6).

4. Beyond Genotype: Phenotyping Symbiodiniaceae

4.1 Why do we need to characterize Symbiodiniaceae phenotypic diversity?

Not all genetically distinct Symbiodiniaceae taxa exhibit physiological differences (*e.g.*, through functional convergence; Goyen et al., 2017; Suggett et al., 2015), whereas unique isolates of the same taxon may be functionally divergent (*e.g.*, Beltrán et al., 2021; Díaz-Almeyda et al., 2017; Hawkins et al., 2016; Howells et al., 2011; Mansour et al., 2018; Parkinson et al., 2016; Parkinson and Baums, 2014; Russnak et al., 2021). This is not surprising given the effects of strong local selection within reef habitats (Howells et al., 2011; Kriefall et al., 2022; Marhoefer et al., 2021; Suggett et al., 2017; van Oppen et al., 2018) and the role of acclimatization (Torda et al., 2017). Stringent functional interrogation is therefore critical to determining how healthy cnidarian-symbiont associations will survive the climate crisis. This goal rests on advancing physiological descriptions, increasing the number of cultured isolates from diverse hosts, and extending the methodological toolbox to characterize Symbiodiniaceae differences. Thus, a more comprehensive functional characterization can accompany taxonomic assignment, helping to build greater community consensus on methods and standards for describing phenotypes of interest (Figure 5).



~ Shared across axenic and xenic culture states

^ Complexity enhanced by the *in hospite* state compared to isolated culture

* Shared across all three states (axenic culture, xenic culture, and *in hospite*)

Figure 5 Considerations for efforts to measure Symbiodiniaceae phenotypes across three states (axenic monoclonal culture, xenic monoclonal culture, and *in hospite*).

4.2 What do we need to consider when assessing Symbiodiniaceae phenotypes *in hospite*?

An overwhelming interest among Symbiodiniaceae researchers to date has been identifying thermal threshold phenotypes based on bio-optics (*e.g.*, Goyen et al., 2017; Hennige et al., 2009; Voolstra et al., 2021d), targeted biochemistry (*e.g.*, Tchernov et al., 2004) or “-omics” metrics (*e.g.*, Olander et al., 2021; Roach et al., 2021). These varied foci illustrate that phenotypes are operationally defined. Consequently, the detected functional diversity (the extent and range of phenotypes resolved) may appear different depending on the metrics used. For example, descriptions of phenotype diversity for heat stress sensitivity based on photobiological properties may not align with those based on metabolic indicators (Goyen et al., 2017) or light adaptation (Suggett et al., 2015; Suggett and van Oppen, 2022). Thus, reconciling genetic diversity with functional diversity must be carefully contextualized based on the measurement criteria and scientific questions at hand.

While it is valuable to confirm symbiont traits when in symbiosis, the presence of local Symbiodiniaceae communities and “secondary” symbionts in cnidarian holobionts complicate this effort. In local Symbiodiniaceae communities, it can be difficult to determine the relative abundance of each lineage present. Although accounting for copy number can help determine symbiont cell number and density in such cases (see Section 3.3.5), other algal-centric physiological metrics, which reflect the combined average of all symbionts present within the host (Cunning et al., 2018), will be difficult to interpret. Single-cell sorting techniques may help assess unique phenotypic distinctions across different symbiont species from the same host (Snyder et al., 2020), but these techniques constitute additional effort and cost. Moreover, cnidarians host a variety of other microeukaryotic, prokaryotic, and viral symbionts (Ainsworth et al., 2017; Hernandez-Agreda et al., 2017; Thurber et al., 2017), some of which are associated with colony health and resilience to environmental stress (Bourne et al., 2016; Voolstra et al., 2021c; Ziegler et al., 2017). Some viruses even infect Symbiodiniaceae cells themselves (Grupstra et al., 2022a; Levin et al., 2017), with diverse potential impacts on Symbiodiniaceae phenotypes (Correa et al., 2021; van Oppen et al., 2009). The degree to which these “secondary” symbionts impact the observed phenotype of Symbiodiniaceae cells *in hospite* is an active area of research (Maire et al., 2021; Matthews et al., 2020). Finally, coral tissue thickness, pigmentation, skeletal reflectance, or other coral-associated microorganisms can affect irradiance levels reaching Symbiodiniaceae *in hospite* (Dimond et al., 2013, 2012; Enríquez et al., 2005; Marcelino et al., 2013; Smith et al., 2013; Titlyanov et al., 2009; Wangpraseurt et al., 2014, 2012). Variation in these physiological metrics can therefore affect symbiont phenotype, and lead to variable responses to climate stress (Hoadley et al., 2019).

Traits where variability across species exceeds that within populations are ideally suited for phenotypic analysis, but are presently unknown to the field or are challenging to measure in consistent and ecologically meaningful ways. Consequently, high-throughput approaches for assessing Symbiodiniaceae phenotypes need to consider tradeoffs that are constrained by end goals. For example, recent high-throughput approaches for assessing thermal tolerance at the whole coral level (*e.g.*, coral bleaching automated stress systems (CBASS; Woolstra et al., 2020) and single cell levels (Behrendt et al., 2020) have incorporated short thermal challenges followed by stress characterization through the measurement of 1-2 physiological variables such as maximum PSII photochemical efficiency (F_v/F_m) and cell density. While single-phenotype assays can be informative within the context of ecosystem service values (*e.g.*, identifying thermally tolerant corals for nursery propagation; Cunning et al., 2021), identification of functionally distinct Symbiodiniaceae phenotypes will benefit from measuring a broader spectrum of physiological metrics (Hoadley et al., 2021). Phenotypic characterization using multiple photosynthetic metrics can provide some species-specific resolution (Suggett et al., 2015), and the non-invasive nature of chlorophyll *a* fluorometry lends itself to high-throughput approaches. However, poor contextualization of photosynthetic parameters with respect to cnidarian resilience currently limits the use of these techniques alone for large-scale phenomic studies, and may ultimately require integration of fitness metrics influenced by resource availability such as elemental composition *via* nutrient acquisition. While specific consensus on measurement protocols is beyond the scope of this perspective, taking a multidisciplinary approach and transparently documenting important methodological choices will help move the field forward.

4.3 What do we need to consider when assessing Symbiodiniaceae phenotypes in culture?

Axenic or bacteria-depleted cultures are promising tools for connecting Symbiodiniaceae genotypes to phenotypes because their genetic identity is readily determined (see Section 1) and morphological, physiological, and behavioral diversity are readily discernible among such algal isolates (Costa et al., 2019; Xiang, 2018; Xiang et al., 2013). In terms of photo-physiology, fluorometry has become a convenient and accessible tool to gauge “culture health” (Hennige et al., 2009; Robison and Warner, 2006; Suggett et al., 2009). Fluorometry is also used in studies examining phenotypic variation focused on photosynthetic traits and how they are affected by resource availability (light, nutrients) and temperature (Díaz-Almeyda et al., 2017; Suggett et al., 2015), and has been inferred to reflect holobiont health (Woolstra et al., 2020). While photosynthetic traits are informative of cellular functioning, they are insufficient in isolation of other measurements to explain phenotypic variation in growth (*e.g.*, Brading et al., 2011; Hennige et al., 2009; Suggett et al., 2015). Recent data point to variable photo-physiological tolerance and thermal plasticity of genetically divergent Symbiodiniaceae grown in monoculture, which has contributed to a deeper understanding of the algal symbiont response to increasing sea surface temperatures (Grégoire et al., 2017;

Klueter et al., 2017; Russnak et al., 2021; Suggett et al., 2015). However, a large number of genetically distinct algal symbionts identified *in hospite* have resisted sustained growth in culture (*e.g.* Krueger and Gates, 2012; Santos et al., 2001). Furthermore, physiological and functional ‘omics data indicate that when in culture or freshly isolated, Symbiodiniaceae exhibit responses to thermal stress that differ from those of the same population *in hospite* (Bellantuono et al., 2019; Gabay et al., 2019; Goulet et al., 2005). These data suggest that some physiological traits measured from culture-based studies may not be easily extrapolated to the symbiotic state. Such issues are particularly pronounced when measuring nutrient-associated phenotypes, as most culture media are nutrient-replete while Symbiodiniaceae *in hospite* appear to be nutrient-limited (Maruyama and Weis, 2021).

Emergent properties are novel characteristics that smaller units of organization gain when they become part of a larger complex system. Research focusing on core emergent properties expressed in culture, that can also be easily assessed in nature (*in hospite*), is logical given that phenotypes will consistently be the result of specific environmental conditions operating on the underlying molecular machinery. However, in decades of studies on Symbiodiniaceae cultures, the environmental conditions imposed have not consistently been reported at the time of, or prior to, sampling. Examples of such metadata include the growth phase (steady state versus non-steady state; Tivey et al., 2020) or cell cycle phase (Fujise et al., 2018; Tivey et al., 2020), as well as the actual environments in the cultures (light quality/quantity, nutrients) as opposed to those measured in the incubators or assumed from the recipe of the medium used (*e.g.*, Camp et al., 2020; Reich et al., 2020), and the extent of bacterial loading. Consequently, developing guidelines for rigorous reporting of environmental (experimental) conditions when phenotypes are quantified is a key priority. Ensuring inter-comparability among studies in the future will similarly depend on operating under a more consistent set of measurement protocols for phenotypic traits.

5. Integrating Multiomic Technologies to Study Symbiodiniaceae

5.1 How can genomics and high-throughput sequencing be leveraged?

With continued cost reductions and increases in computational power and accessibility, advanced “-omic” technologies, including genomics, transcriptomics, proteomics, and metabolomics, are rapidly enhancing our ability to understand biological mechanisms (Krassowski et al., 2020). Coupled with powerful, multivariate statistical techniques and machine learning, ‘omics technologies have greatly refined our understanding of Symbiodiniaceae biology.

Whole-genome sequencing (WGS) remains the gold standard for capturing genetic diversity (Aranda et al., 2016; González-Pech et al., 2021; Lin et al., 2015; Liu et al., 2018; Reich et al., 2021; Shoguchi et al., 2018, 2013). While individual and concatenated genes can help resolve phylogenetic relationships and define taxonomic lineages (LaJeunesse et al., 2018; Parkinson et al., 2015), whole-genome data can provide more comprehensive phylogenomic signals, which can be used to investigate divergent selection. For example, while a Symbiodiniaceae phylogeny reconstructed using k -mers (short, sub-sequences of defined length k) derived from whole-genome sequences is largely consistent with the phylogeny reconstructed with *LSU* rDNA data (González-Pech et al., 2021), different genomic regions exhibit distinct phylogenetic signals (Lo et al., 2022). Further comparison of WGS data indicate that the similarity shared between different species within the genus *Symbiodinium* is comparable to that between different Symbiodiniaceae genera, revealing more extensive divergence than anticipated and suggesting a need for future revision (Dougan et al., 2022).

WGS efforts employ short- or long-read technologies, or a combination of both strategies. Short-read sequencing technologies (*e.g.*, Illumina) have typically offered a cost-effective approach for deep sequencing with low error rates (< 1%) and have provided valuable insights into Symbiodiniaceae diversity, including gene family expansions across different Symbiodiniaceae lineages (Aranda et al., 2016; Lin et al., 2015; Liu et al., 2018). Though short-read sequencing can identify lineage-specific divergence, short reads are difficult to assemble, especially with highly repetitive genomic regions. Long-read sequencing technologies such as Pacific Biosciences (PacBio) or Oxford Nanopore Technologies offer a viable alternative to improve the contiguity of highly fragmented short-read genomes. However, long-read sequencing technologies are more expensive and error-prone relative to short-read platforms, though these technologies are rapidly advancing (Karst et al., 2021). Long-read data allow us to observe chromosome structure, a greater number of genomic elements, such as DNA transposons, long terminal repeats, or chromosomal enrichment for genes with similar biological functions (González-Pech et al., 2021; Li et al., 2020; Nand et al., 2021). Chromosome-level assemblies represent a major milestone in dinoflagellate genomics as they confirm that many genes are encoded in unidirectional clusters which correspond to large topological domains (Marinov et al., 2021; Nand et al., 2021). Not only does this discovery provide insights into the structure of genes in Symbiodiniaceae, but the structure can be correlated with the encoded genes and their expression patterns to observe their interactions, elucidating novel insights into the evolution of diverse Symbiodiniaceae lineages at the chromosome level (Lin et al., 2021).

Efforts are underway to expand the number of high-quality short- and long-read assemblies for cnidarian-associated and free-living Symbiodiniaceae and incorporate these data into taxonomic descriptions (Dougan et al., 2022; McKenna et al., 2021; Voolstra et al., 2021b). Additionally, the two read types can be coupled

(*e.g.*, Illumina with PacBio HiFi) to incorporate both the contiguous sequences (> 20 kb) of long reads with the low error rate of short reads, allowing robust comparisons of sequence divergence within and across genomes (Ebert et al., 2021). Once more data become available, it may be feasible to incorporate whole-genome information into future taxonomic and systematic revisions to the family (Dougan et al., 2022). However, it will be crucial to achieve consensus on how to use these data to study Symbiodiniaceae diversity and taxonomy. A lack of consistency in methodology and quality standards persists, making cross-study analyses difficult (*e.g.*, Chen et al., 2020). First steps in such standardization have been taken (Voolstra et al., 2021b), but will need to be expanded as more genome data become available and the community using these data grows. Additionally, the current costs of completely sequencing the genomes of hundreds of potential Symbiodiniaceae species remains prohibitive. For the near future, feasible alternatives for WGS include using reduced representation phylogenomic approaches (such as those targeting ultraconserved elements; Cowman et al., 2020; Quattrini et al., 2020), full-length rDNA gene amplicons (Tedersoo et al., 2018), or entire organellar genome sequences (Liu et al., 2020). These alternatives may represent a compromise between WGS and phylogenetic marker studies by providing an intermediate amount of sequence information for taxonomic accuracy.

5.2 How can transcriptomics, proteomics, and single-cell techniques advance our knowledge?

Researchers are keen to make functional inferences about Symbiodiniaceae, which requires focusing on coding regions. To do so requires sequencing the collection of RNAs within the cells (*i.e.*, transcriptomics). Transcriptome sequencing characterizes molecular phenotypes, such as transient responses to the environment, and can reveal differential gene expression among taxa that could reflect selective pressures driving Symbiodiniaceae diversification at the functional level (Avila-Magaña et al., 2021; Bayer et al., 2012; Parkinson et al., 2016). However, the extent of gene expression changes among Symbiodiniaceae is often surprisingly subtle (*e.g.*, Barshis et al., 2014; Davies et al., 2018; Parkinson et al., 2016), although this is a matter of current debate (*e.g.*, Bellantuono et al., 2019; Voolstra et al., 2021d). Furthermore, transcription can be influenced through alternatively spliced transcripts (Lin, 2011; Méndez et al., 2015), RNA editing (Liew et al., 2017; Mungpakdee et al., 2014; Shoguchi et al., 2020), microRNA interactions (Baumgarten et al., 2018), and methylation of mRNAs (de Mendoza et al., 2018; Lohuis and Miller, 1998; Yang et al., 2020). These post-transcriptional modifications create variation in the transcriptome, which can complicate transcriptomic interpretation, but tracking the conservation and divergence of these variations across the Symbiodiniaceae phylogeny may elucidate novel insights into the evolution of diverse lineages.

The primary concern with bulk transcriptomic analysis is that methods often pool transcriptomes from all Symbiodiniaceae cells within a host sample, so only “average” expression profiles can be generated (Traylor-

Knowles, 2021). This approach may obscure nuances in the interactions between specific symbiont and host cells, especially for less abundant symbionts. Single-cell transcriptomics (*i.e.*, isolating individual cells and sequencing their transcriptomes) could solve this issue as gene expression could be explored within and among each symbiont cell *in hospite*. The generation of a cell atlas for the coral *Stylophora pistillata* enabled the characterization of fine-scale metabolic interactions between symbionts and host gastrodermal cells (Levy et al., 2021). Single-cell sequencing can also enable high-resolution interrogations of how Symbiodiniaceae and host cells interact during symbiosis establishment, maintenance, and breakdown, particularly when Symbiodiniaceae cells can be isolated from different parts of the host coral that exhibit contrasting physiologies. By comparing expression from symbiont cells derived from different positions in the coral colony, the location and ecological role of Symbiodiniaceae can be characterized, which is a major priority for improving our understanding of symbiont communities within cnidarians.

Proteomic analyses provide an alternative mechanism to explore Symbiodiniaceae physiology and the functional impacts of different symbionts on cnidarian-algal associations, such as metabolic mismatches that occur when hosts associate with atypical (heterologous) symbionts (Sproles et al., 2019). Proteomic analyses use liquid chromatography-mass spectrometry to identify and quantify proteins, which are a precursor to phenotype and more directly linked to phenotype than transcript abundance (Feder and Walser, 2005). “Bottom-up” or “shotgun” approaches are commonly employed to identify and quantify as many proteins as possible in a sample in an untargeted manner, and with modern instrumentation, 3,000-4,000 proteins are commonly quantified in established model systems as well as Symbiodiniaceae (Camp et al., 2022; Richards et al., 2015). Two features of proteomic analyses are particularly powerful. One is the characterization of post-translational modifications, such as protein phosphorylation, oxidation, acetylation, or ubiquitination (Witze et al., 2007). Post-translational modifications help regulate protein activity and are crucial in many biological processes, yet they cannot be detected by pre-translational analyses. The second is the ability to localize proteins to particular cellular compartments by sample fractionation, which is valuable in increasing proteome coverage depth and provides non-sequence-based data for inferring protein function (Tortorelli et al., 2022). This may be of particular use with Symbiodiniaceae, where sequence homology-based gene annotation is less effective due to their phylogenetic distance from well-studied organisms, and to the highly derived genomes of dinoflagellates. However, these methods still require high-quality protein model search databases derived from genomic or transcriptomic sequences as they cannot identify proteins from complex samples *de novo*. Care must be taken to use a sequence search database that is tailored to the organism in question (*e.g.*, congeneric sequences and spectra) to generate highly resolved functional data.

5.3 What other omic technologies are promising?

Epigenomics, genome editing, metabolomics, and volatilomics are emerging areas within Symbiodiniaceae research. Epigenomic mechanisms, such as DNA methylation or chromatin modification, can modulate gene expression *via* gene suppression, gene enhancement, alternative mRNA splicing, or the regulation of spurious transcription without requiring any changes to genomic sequences (Bossdorf et al., 2008; Feil and Fraga, 2012; Foret et al., 2012). DNA methylation is one epigenetic modification that occurs when methyl groups are added to DNA nucleotides, altering how transcriptional proteins bind to promoter regions thereby altering gene expression (Suzuki and Bird, 2008). Symbiodiniaceae have unusually high levels of genome methylation (Lohuis and Miller, 1998). Originally, the high level of methylation raised uncertainty about whether methylation actually played a role in gene regulation, but methylation has been linked to differential gene expression with varying irradiance (Yang et al., 2020). Thus far, epigenomic analyses have largely been focused on the host animal, and questions are often centered around how methylation contributes to environmental tolerance (Dixon et al., 2018, 2014; Dixon and Matz, 2021; Durante et al., 2019; Liew et al., 2018; Putnam et al., 2016; Putnam and Gates, 2015; Rodríguez-Casariago et al., 2021; Rodríguez-Casariago et al., 2020; Rodríguez-Casariago et al., 2018). Therefore, determining how methylation contributes to Symbiodiniaceae functional diversity requires further exploration.

Overall, Symbiodiniaceae genomes are very difficult to annotate. At present, dinoflagellate genome and transcriptome projects rarely manage to annotate > 50% of putative coding sequences *via* homology searches against genes that have been functionally characterized in other organisms (González-Pech et al., 2021; Stephens et al., 2018). In the future, genome editing could be better developed to knock out Symbiodiniaceae genes with unknown functions, making it easier to determine their biological roles. UV mutagenesis is a classic method for introducing mutations; it was used recently to create photosynthesis mutants *via* screening of colored mutants (Jinkerson et al., 2022), but its random nature is less than ideal for reverse genetics. RNA silencing is a more targeted approach that could potentially be exploited in Symbiodiniaceae studies (Zhang and Lin, 2019), but the rapidly advancing CRISPR/Cas9 technology is most desirable for its ability to knock out specific genes. Although genome editing efforts for protists have made encouraging progress (Faktorová et al., 2020), success with Symbiodiniaceae remains elusive (Chen et al., 2019).

Biochemical analyses of Symbiodiniaceae are also in the early stages. Characterization of metabolic products (metabolomics) and volatile organic compounds (volatilomics) can provide insights into molecular cross-talk between partners. Among Symbiodiniaceae, both metabolomic and volatilomic profiles are species-specific, but they also fluctuate with environmental conditions (Klueter et al., 2015; Lawson et al., 2019; Roach et al.,

2021). Distinct biochemical profiles reflect the interactions and coevolution (or lack thereof) between the host and symbiont (Matthews et al., 2017), so biochemical assays can lead to a greater understanding of the drivers of Symbiodiniaceae evolution. As with all the omics methods mentioned so far, if metabolomics and volatilomics are to be used to understand Symbiodiniaceae divergence, more data spanning the phylogeny will be required.

5.4 How can we integrate omic technologies?

Integrative approaches that use more than one type of technology may be required to answer intricate research questions about Symbiodiniaceae biology. For example, transcriptomics informs us of gene expression patterns, but cannot reveal protein end-products and how they are used for symbiosis, particularly since transcript abundance does not correlate well with protein levels (Cziesielski et al., 2018; Liang et al., 2021). However, when transcriptomics is integrated with proteomics or metabolomics, phenotypes can be directly observed. Then, tools that make use of multivariate statistics to combine these different types of biological data across studies, such as mixOmics (Rohart et al., 2017) or weighted gene coexpression network analysis (WGCNA; Langfelder and Horvath, 2008), are improving our ability to elucidate molecular mechanisms associated with phenotypes of interest. Thus, integration across several omic technologies (“multiomics”) holds great promise for advancing our understanding of cnidarian-algal symbiosis (e.g., Camp et al., 2022), yet the financial costs associated with using multiple technologies in parallel remains a limiting factor. Additionally, the expertise of one laboratory may be restricted to one major type of analysis or instrument. Therefore, collaborations among different research groups are essential (see Section 6). Logistical hurdles to collaborations include how to house and share samples and data, along with the major financial burden. Integrative approaches will remain constrained until these issues are resolved or facilitated through funding agencies.

6. Ensuring an Inclusive Symbiodiniaceae Research Community

6.1 How can we improve inclusivity in Symbiodiniaceae research?

Despite an increased recognition of the benefits and need for more diverse representation in science, systemic biases continue to persist in science at large, limiting our creativity and innovation potential (Ahmadia et al., 2021). In coral reef science, where actual reefs are mostly found in non-industrial nations, capacity-building through “leveling the playing field” is required to facilitate a more inclusive research community and advance novel and important discoveries (O’Brien et al., 2020). Marginalized groups within science have been and continue to be excluded from access to many opportunities, including funding, publishing, resources, collaborations, and networking. This exclusion is driven by limited resource availability and systemic racism

and sexism (Davies et al., 2021; Dzirasa, 2020; Ginther et al., 2011; Hoppe et al., 2019; Taffe and Gilpin, 2021). While some progress has been made in the scientific community more broadly, there are still many deeply entrenched biases and critical gender, race, and ethnicity gaps exist with respect to resource access; these need to be addressed by researchers, including those whose work focuses on Symbiodiniaceae, to ensure a more inclusive scientific community.

Research institutions, hiring committees, and organizers of panels, seminars, and conferences must actively work to change the demographics of scientists by increasing diversity at all levels—from trainees to senior research scientists in positions of power. Gender, race, and ethnicity biases are rampant in the scientific-hiring process (*e.g.*, Barber et al., 2020; Bennett et al., 2020; Huang et al., 2020); for example, in the United States these biases are particularly strong against Black and Latinx scholars (Eaton et al., 2020) and in New Zealand biases are stronger against people of Māori and Pasifika descent (Naepi et al., 2020). In addition to recruitment difficulties, if scholars from these backgrounds are hired, they often face continued challenges that hinder their retention. Recognizing this, people in positions of power in the Symbiodiniaceae scientific community should 1) invest in retaining a diverse workforce by promoting the academic work of minority scientists; 2) provide spaces where researchers can safely report aggressions and other challenges (Valenzuela-Toro and Viglino, 2021); and 3) create programs that provide strong multidimensional mentorship, which serve to support and retain these scholars throughout each career stage (Davies et al., 2021; Montgomery, 2017a; Montgomery, 2017b; Montgomery et al., 2014). Moving forward, Symbiodiniaceae researchers need to understand and implement the strategies and proposals that already exist and continue to be put forward regarding increasing recruitment and retention of marginalized scholars (*e.g.*, Barber et al., 2020; Chaudhary and Berhe, 2020; Greider et al., 2019). Increasing the diversity of perspectives at the decision-making table leads to more innovative discoveries (Hofstra et al., 2020; Nielsen et al., 2018), which are desperately needed to meet the formidable challenges of the coral reef crisis.

6.2 How can we ensure an equitable publication process for everyone?

Equity and diversity issues exist in the scholarly publication process at multiple levels and across different areas of research. Men are first authors more often than women (Casadevall et al., 2019), notably even when both authors are identified as having contributed equally to the work (Broderick and Casadevall, 2019). Such systematic and implicit gender biases are also evident in the peer-review processes (Calaza et al., 2021). Manuscript authors, irrespective of gender, are also less likely to suggest women reviewers (Fox et al., 2017). Unprofessional reviews disproportionately impact members of underrepresented groups, who report greater self-doubt after receiving such reviews, ultimately reducing scientific productivity overall (Silbiger and Stubler, 2019). Beyond peer evaluation, more men serve in editorial roles than women (Fox et al., 2019);

Grinnell et al., 2020; Hafeez et al., 2019; Palser et al., 2022; Pinho-Gomes et al., 2021), and editors tend to invite men more often than women to write invited reviews or perspectives. For example, the journal *Molecular Ecology*, which often publishes research from the Symbiodiniaceae community, found significant gender bias in authorship of invited ‘perspective’ articles, with women only authoring between 17.2-28.6% of these pieces (Baucom et al., 2019).

Language biases are also pervasive. English is currently the default language of science, which disadvantages scientists who do not consider English as their primary language (Gordin, 2015). Non-native English speakers spend on average 97 more writing hours than native English speakers on preparation for each manuscript (Ramírez-Castañeda, 2020). In addition, ideas may be lost in translation or are often challenging to explain in a secondary language (Flowerdew, 2001). There are also costs associated with publishing in a non-native language: for example, paying for translation and editorial services. Conversely, fluent English speakers publish more research articles at higher rates than non-English speakers (Taubert et al., 2021). To address English-centric journals, regional journals publish in their native languages (Bordons, 2004), but these publications are read by a smaller readership and are cited less (Di Bitetti and Ferreras, 2017), and thereby viewed as less impactful, and are less likely to be shared widely in the Symbiodiniaceae community.

General actions that can be taken within our community to ensure a more equitable publication process include: 1) increasing diversity on editorial boards, 2) increasing the diversity of invited reviewers, as well as the authors we review for; 3) promoting cost reduction strategies (as implemented in journals like *Frontiers*, *PeerJ*, and *PLoS One*) whereby publication fees are prorated by country or institution type, as well as other strategies that reduce editorial costs for non-native speakers (Taubert et al., 2021); 4) promoting double-blind review processes (Budden et al., 2008); 5) intentionally citing articles led by diverse colleagues (*e.g.*, from diverse gender identities and geographical areas), thereby increasing the diversity of perspectives in the field that contribute to discussion; and 6) gathering data about where and how these inequities exist and working together as a community to take actionable steps for equity in science.

6.3 How can we avoid parachute science?

Parachute science, sometimes referred to as “helicopter” or “colonial” science, is a common practice whereby members of the scientific community from higher-income countries fail to involve local/indigenous/native people in an equitable fashion when performing research in lower-income countries (Haelewaters et al., 2021; Stefanoudis et al., 2021). These practices tend to be more common in ecology and conservation research (de Vos, 2020), including coral reef studies. As a community, we need to understand the sometimes exploitative history of our discipline and avoid perpetuating it. We should stay informed regarding the history of the lands

and peoples who live in the areas in which we conduct our studies. Our specific recommendations include: 1) developing laboratory manuals that includes sections outlining values and best practices (including ethics approval and necessary permits); 2) adequately training students to conduct transparent research and develop equitable relationships with members of the host region; 3) supporting the establishment of long-term collaborations and exchange programs to involve local students in research; and 4) including the development of these relationships as important components of the tenure and promotion process in departments and institutions.

Importantly, researchers from institutions in high-income regions (*e.g.*, North America, Australia, Western Europe) should: 1) be sensitive to the many challenges their colleagues in lower-income regions experience, such as a lack of funding, infrastructure, and institutional support; 2) be respectful of these collaborators by treating them as peers and not as assistants, involving them in all steps of the science, and acknowledging their intellectual contributions during discussions, and; 3) be fair to these researchers and their contributions through continued involvement in planning, manuscripts, projects, and grants generated through these collaborations. Together, these strategies can facilitate a more inclusive and collaborative Symbiodiniaceae community (Armenteras, 2021; Belhabib, 2021). Importantly, integration of members of the local community provides a long-term context to the collected scientific data, such as anecdotal observations (*e.g.*, episodes of bleaching) that may be critically informative to the research.

6.4 How might we increase accessibility and collaboration?

There is an urgent need to increase the accessibility of Symbiodiniaceae science and foster collaboration, as innovation is necessary to address the coral reef crisis. One way to facilitate accessibility and collaboration would be the development of a database of all Symbiodiniaceae researchers and their research products and expertise. This database would connect and promote diverse researchers and also facilitate discussions of science from different perspectives. It could be queried based on topic, expertise, and methodology to aid scientists in diversifying their citations. The COVID-19 pandemic also showcased the effectiveness of virtual conferences (case in point: this perspective is the product of a virtual workshop). Maintaining hybrid conferences with reduced costs for virtual attendance would ensure that the sharing of scientific information is more inclusive, alongside continued efforts to drop conference charges for lower-income countries (*e.g.*, the 15th International Coral Reef Symposium in 2022). Therefore, we encourage conference organizers to facilitate virtual attendance and funding sources for those who have difficulties traveling to foster equitable networking opportunities across people from diverse backgrounds and academic stages. Additionally, incentivizing consortia of Symbiodiniaceae researchers across diverse career stages and locations, and explicitly engaging researchers from marginalized backgrounds, would lead to stronger capacity building and

greater transfer of knowledge. Lastly, we encourage sponsors to continue expanding their funding schemes to support international collaborations. Several examples of these efforts exist, including a new grant solicitation from the United States National Science Foundation, which calls for collaborations with Brazilian scientists through the São Paulo Research Foundation. Similar schemes (*e.g.*, Deutsche Forschungsgemeinschaft) also exist to foster collaboration between researchers in developing countries. The United States Fulbright Program and the European Union Marie Skłodowska-Curie Program are other examples that support collaborations across countries. The Japanese Society for the Promotion of Science provides funding for international exchange and research for graduate students, postdoctoral scholars, and early career scientists. These types of funding mechanisms are important because they promote wealth sharing across countries, encourage collaboration while thwarting parachute science, help with international challenges including permitting, and ultimately lead to more open sharing of data and ideas within our Symbiodiniaceae research community and beyond.

Conclusions

Addressing and ultimately solving the challenges associated with the coral reef crisis is increasingly urgent as climate change accelerates. Microalgae in the family Symbiodiniaceae play a critical role in determining coral bleaching outcomes; advancing our knowledge of the genetic diversity of these organisms, how their diversity functionally impacts coral bleaching, and how we can apply such knowledge to mitigate climate change consequences is vital. We identified consensus approaches for studies of Symbiodiniaceae genetic diversity at the species and population levels, while recognizing several outstanding issues regarding the characterization of community diversity. We highlight key paths forward for research including exploration of the phenotypic landscape and leveraging new technologies that are broadly applied in model systems. We also emphasize the need for increased collaboration and inclusivity among Symbiodiniaceae researchers. Overall, we acknowledge the dire need for advancing our understanding of Symbiodiniaceae ecology, physiology, and evolution, which will have the potential to expedite restoration practices and facilitate management decisions as we continue to push for political action on climate change.

Acknowledgements

Funding for the virtual workshop during which the content of this manuscript was developed was provided by the National Science Foundation Division of Biological Oceanography (OCE 2127506; 2127514; and 2127508 to SWD, AMSC, and JEP, respectively). We would like to thank Dan Thornhill, Allison Lewis, and Joanna Shisler at NSF for attending and facilitating the workshop, and Andrea Grottoli, Jody Harwood, Krista Liguori, Amy Pruden, and Rebecca Vega Thurber for advice on running workshops. Finally, we wish to thank all Symbiodiniaceae researchers past and present whose research has helped shape our understanding of Symbiodiniaceae diversity and biology.

Author Contributions

Funding (SWD, AMSC, JEP), conceptualization (SWD, AMSC, JEP), project management (SWD, AMSC, JEP), workshop organization (SWD, AMSC, JEP, MHG, NGK, LHK), post-workshop facilitation (SWD, AMSC, JEP, MHG, NGK, LHK), writing and editing (all authors).

References

- Abdelkrim J, Robertson B, Stanton J-A, Gemmell N. 2009. Fast, cost-effective development of species-specific microsatellite markers by genomic sequencing. *Biotechniques* **46**:185–192.
- Abrego D, Ulstrup KE, Willis BL, van Oppen MJH. 2008. Species-specific interactions between algal endosymbionts and coral hosts define their bleaching response to heat and light stress. *Proc Biol Sci* **275**:2273–2282.
- Adams LM, Cumbo VR, Takabayashi M. 2009. Exposure to sediment enhances primary acquisition of *Symbiodinium* by asymbiotic coral larvae. *Mar Ecol Prog Ser* **377**:149–156.
- Ahmadia GN, Cheng SH, Andradi-Brown DA, Baez SK, Barnes MD, Bennett NJ, Campbell SJ, Darling ES, Estradivari, Gill D, Gress E, Gurney GG, Horigue V, Jakub R, Kennedy EV, Mahajan SL, Mangubhai S, Matsuda SB, Muthiga NA, Navarro MO, Santodomingo N, Vallès H, Veverka L, Villagomez A, Wenger AS, Wosu A. 2021. Limited progress in improving gender and geographic representation in coral reef science. *Front Mar Sci* **8**:731037.
- Aichelman HE, Barshis DJ. 2020. Adaptive divergence, neutral panmixia, and algal symbiont population structure in the temperate coral along the Mid-Atlantic United States. *PeerJ* **8**:e10201.
- Ainsworth TD, Fordyce AJ, Camp EF. 2017. The other microeukaryotes of the coral reef microbiome. *Trends Microbiol* **25**:980–991.
- Ali A, Kriefall NG, Emery LE, Kenkel CD, Matz MV, Davies SW. 2019. Recruit symbiosis establishment and Symbiodiniaceae composition influenced by adult corals and reef sediment. *Coral Reefs* **38**:405–415.
- Andras JP, Kirk NL, Coffroth MA, Harvell CD. 2009. Isolation and characterization of microsatellite loci in *Symbiodinium* B1/B184, the dinoflagellate symbiont of the Caribbean sea fan coral, *Gorgonia ventalina*. *Mol Ecol Resour* **9**:989–993.
- Andras JP, Kirk NL, Drew Harvell C. 2011. Range-wide population genetic structure of *Symbiodinium* associated with the Caribbean Sea fan coral, *Gorgonia ventalina*. *Mol Ecol* **20**:2525–2542.
- Apprill AM, Bidigare RR, Gates RD. 2007. Visibly healthy corals exhibit variable pigment concentrations and symbiont phenotypes. *Coral Reefs* **26**:387–397.
- Apprill AM, Gates RD. 2007. Recognizing diversity in coral symbiotic dinoflagellate communities. *Mol Ecol* **16**:1127–1134.
- Aranda M, Li Y, Liew YJ, Baumgarten S, Simakov O, Wilson MC, Piel J, Ashoor H, Bougouffa S, Bajic VB, Ryu T, Ravasi T, Bayer T, Micklem G, Kim H, Bhak J, LaJeunesse TC, Voolstra CR. 2016. Genomes of coral dinoflagellate symbionts highlight evolutionary adaptations conducive to a symbiotic lifestyle. *Sci Rep* **6**:39734.
- Arif C, Daniels C, Bayer T, Banguera-Hinestroza E, Barbrook A, Howe CJ, LaJeunesse TC, Voolstra CR. 2014. Assessing *Symbiodinium* diversity in scleractinian corals via next-generation sequencing-based genotyping of the ITS2 rDNA region. *Mol Ecol* **23**:4418–4433.
- Armenteras D. 2021. Guidelines for healthy global scientific collaborations. *Nature Ecology & Evolution* **5**:1193–1194.
- Avila-Magaña V, Kamel B, DeSalvo M, Gómez-Campo K, Enríquez S, Kitano H, Rohlfs RV, Iglesias-Prieto R, Medina M. 2021. Elucidating gene expression adaptation of phylogenetically divergent coral holobionts under heat stress. *Nat Commun* **12**:5731.
- Baillie BK, Belda-Baillie CA, Maruyama T. 2000. Conspecificity and Indo-pacific distribution of *Symbiodinium* genotypes (Dinophyceae) from giant clams. *J Phycol* **36**:1153–1161.
- Baker AC. 2003. Flexibility and specificity in coral-algal symbiosis: Diversity, ecology, and biogeography of *Symbiodinium*. *Annu Rev Ecol Evol Syst.* **34**:661–689
- Baker AC, Romanski AM. 2007. Multiple symbiotic partnerships are common in scleractinian corals, but not in octocorals: Comment on Goulet (2006). *Mar Ecol Prog Ser* **335**:237–242.
- Banaszak AT, García Ramos M, Goulet TL. 2013. The symbiosis between the gastropod *Strombus gigas* and

- the dinoflagellate *Symbiodinium*: An ontogenic journey from mutualism to parasitism. *J Exp Mar Bio Ecol* **449**:358–365.
- Barber PH, Hayes TB, Johnson TL, Márquez-Magaña L, 10,234 signatories. 2020. Systemic racism in higher education. *Science* **369**:1440–1441.
- Barshis DJ, Ladner JT, Oliver TA, Palumbi SR. 2014. Lineage-specific transcriptional profiles of *Symbiodinium* spp. unaltered by heat stress in a coral host. *Mol Biol Evol* **31**:1343–1352.
- Baucom RS, Gerald AM, Rieseberg LH. 2019. Some perspective on Molecular Ecology perspectives: Are women being left out? *Mol Ecol* **28**:2451–2455.
- Baumgarten S, Cziesielski MJ, Thomas L, Michell CT, Esherick LY, Pringle JR, Aranda M, Voolstra CR. 2018. Evidence for miRNA-mediated modulation of the host transcriptome in cnidarian-dinoflagellate symbiosis. *Mol Ecol* **27**:403–418.
- Baums IB, Devlin-Durante MK, LaJeunesse TC. 2014. New insights into the dynamics between reef corals and their associated dinoflagellate endosymbionts from population genetic studies. *Mol Ecol* **23**:4203–4215.
- Baums IB, Johnson ME, Devlin-Durante MK, Miller MW. 2010. Host population genetic structure and zooxanthellae diversity of two reef-building coral species along the Florida Reef Tract and wider Caribbean. *Coral Reefs*. **29**:835–842.
- Bayer T, Aranda M, Sunagawa S, Yum LK, Desalvo MK, Lindquist E, Coffroth MA, Voolstra CR, Medina M. 2012. *Symbiodinium* transcriptomes: Genome insights into the dinoflagellate symbionts of reef-building corals. *PLoS One* **7**:e35269.
- Bay LK, Doyle J, Logan M, Berkelmans R. 2016. Recovery from bleaching is mediated by threshold densities of background thermo-tolerant symbiont types in a reef-building coral. *R Soc Open Sci* **3**:160322.
- Bay LK, Howells EJ, van Oppen MJH. 2009. Isolation, characterisation and cross amplification of thirteen microsatellite loci for coral endo-symbiotic dinoflagellates (*Symbiodinium* Clade C). *Conserv Genet Resour* **1**:199–203.
- Behrendt L, Salek MM, Trampe EL, Fernandez VI, Lee KS, Kühl M, Stocker R. 2020. PhenoChip: A single-cell phenomic platform for high-throughput photophysiological analyses of microalgae. *Sci Adv* **6**:eabb2754.
- Belhabib D. 2021. Ocean science and advocacy work better when decolonized. *Nat Ecol Evol* **5**:709–710.
- Bellantuono AJ, Dougan KE, Granados-Cifuentes C, Rodriguez-Lanetty M. 2019. Free-living and symbiotic lifestyles of a thermotolerant coral endosymbiont display profoundly distinct transcriptomes under both stable and heat stress conditions. *Mol Ecol* **28**:5265–5281.
- Beltrán VH, Puill-Stephan E, Howells E, Flores-Moya A, Doblin M, Núñez-Lara E, Escamilla V, López T, van Oppen MJH. 2021. Physiological diversity among sympatric, conspecific endosymbionts of coral (*Cladocopium* C1^{acro}) from the Great Barrier Reef. *Coral Reefs* **40**:985–997.
- Bennett CL, Salinas RY, Locascio JJ, Boyer EW. 2020. Two decades of little change: An analysis of U.S. medical school basic science faculty by sex, race/ethnicity, and academic rank. *PLoS One* **15**:e0235190.
- Berkelmans R, Van Oppen MJH. 2006. The role of zooxanthellae in the thermal tolerance of corals: A “nugget of hope” for coral reefs in an era of climate change. *Proceedings of the Royal Society B: Biological Sciences* **273**:2305–2312.
- Bordons M. 2004. Towards international recognition of Spanish scientific publications. *Rev Esp Cardiol* **9**:799–802.
- Bossdorf O, Richards CL, Pigliucci M. 2008. Epigenetics for ecologists. *Ecol Lett* **11**:106–115.
- Boulotte NM, Dalton SJ, Carroll AG, Harrison PL, Putnam HM, Peplow LM, van Oppen MJ. 2016. Exploring the *Symbiodinium* rare biosphere provides evidence for symbiont switching in reef-building corals. *ISME J* **10**:2693–2701.
- Bourne DG, Morrow KM, Webster NS. 2016. Insights into the coral microbiome: Underpinning the health and resilience of reef ecosystems. *Annu Rev Microbiol* **70**:317–340.
- Brading P, Warner ME, Davey P, Smith DJ, Achterberg EP, Suggett DJ. 2011. Differential effects of ocean acidification on growth and photosynthesis among phylotypes of *Symbiodinium* (Dinophyceae). *Limnol*

- Oceanogr* **56**:927–938.
- Brian JI, Davy SK, Wilkinson SP. 2019. Multi-gene incongruence consistent with hybridisation in *Cladocopium* (Symbiodiniaceae), an ecologically important genus of coral reef symbionts. *PeerJ* **7**:e7178.
- Broderick NA, Casadevall A. 2019. Meta-Research: Gender inequalities among authors who contributed equally. *eLife* **8**:e36399.
- Buddemeier RW, Fautin DG. 1993. Coral bleaching as an adaptive mechanism: A testable hypothesis. *Bioscience* **43**:320–326.
- Budden AE, Tregenza T, Aarssen LW, Koricheva J, Leimu R, Lortie CJ. 2008. Double-blind review favours increased representation of female authors. *Trends Ecol Evol* **23**:4–6.
- Calaza KC, Erthal FCS, Pereira MG, Macario KCD, Daflon VT, David IPA, Castro HC, Vargas MD, Martins LB, Stariolo JB, Volchan E, de Oliveira L. 2021. Facing racism and sexism in science by fighting against social implicit bias: A Latina and Black woman's perspective. *Front Psychol* **12**:671481.
- Camp EF, Kahlke T, Nitschke MR, Varkey D, Fisher NL, Fujise L, Goyen S, Hughes DJ, Lawson CA, Ros M, Woodcock S, Xiao K, Leggat W, Suggett DJ. 2020. Revealing changes in the microbiome of Symbiodiniaceae under thermal stress. *Environ Microbiol* **22**:1294–1309.
- Camp EF, Kahlke T, Signal B, Oakley CA, Lutz A, Davy SK, Suggett DJ, Leggat WP. 2022. Proteome metabolome and transcriptome data for three Symbiodiniaceae under ambient and heat stress conditions. *Sci Data* **9**:153.
- Casadevall A, Semenza GL, Jackson S, Tomaselli G, Ahima RS. 2019. Reducing bias: accounting for the order of co-first authors. *J Clin Invest* **129**:2167–2168.
- Castro-Sanguino C, Sánchez JA. 2012. Dispersal of *Symbiodinium* by the stoplight parrotfish *Sparisoma viride*. *Biol Lett* **8**:282–286.
- Chaudhary VB, Berhe AA. 2020. Ten simple rules for building an antiracist lab. *PLoS Comput Biol* **16**:e1008210.
- Chen JE, Barbrook AC, Cui G, Howe CJ, Aranda M. 2019. The genetic intractability of *Symbiodinium microadriaticum* to standard algal transformation methods. *PLoS One* **14**:e0211936.
- Chen Y, González-Pech RA, Stephens TG, Bhattacharya D, Chan CX. 2020. Evidence that inconsistent gene prediction can mislead analysis of dinoflagellate genomes. *J Phycol* **56**:6–10.
- Claar DC, Starko S, Tietjen KL, Epstein HE, Cunning R, Cobb KM, Baker AC, Gates RD, Baum JK. 2020a. Dynamic symbioses reveal pathways to coral survival through prolonged heatwaves. *Nat Commun* **11**:6097.
- Claar DC, Tietjen KL, Cox KD, Gates RD, Baum JK. 2020b. Chronic disturbance modulates symbiont (Symbiodiniaceae) beta diversity on a coral reef. *Sci Rep* **10**:4492.
- Coffroth MA, Lewis CF, Santos SR, Weaver JL. 2006. Environmental populations of symbiotic dinoflagellates in the genus *Symbiodinium* can initiate symbioses with reef cnidarians. *Curr Biol* **16**:R985–7.
- Coffroth MA, Poland DM, Petrou EL, Brazeau DA, Holmberg JC. 2010. Environmental symbiont acquisition may not be the solution to warming seas for reef-building corals. *PLoS One* **5**:e13258.
- Correa AMS. 2009. Molecular ecology of endosymbiotic dinoflagellates (genus *Symbiodinium*) in scleractinian corals. Columbia University. 252 pp.
- Correa AMS, Baker AC. 2009. Understanding diversity in coral-algal symbiosis: a cluster-based approach to interpreting fine-scale genetic variation in the genus *Symbiodinium*. *Coral Reefs* **28**:81–93.
- Correa AMS, Brandt ME, Smith TB, Thornhill DJ, Baker AC. 2009. *Symbiodinium* associations with diseased and healthy scleractinian corals. *Coral Reefs* **28**:437–448.
- Correa AMS, Howard-Varona C, Coy SR, Buchan A, Sullivan MB, Weitz JS. 2021. Revisiting the rules of life for viruses of microorganisms. *Nat Rev Microbiol* **19**:501–513.
- Correa AMS, McDonald MD, Baker AC. 2009. Development of Clade-specific *Symbiodinium* primers for quantitative PCR (qPCR) and their application to detecting Clade D symbionts in Caribbean corals. *Mar Biol* **156**:2403–2411.

- Costa RM, Fidalgo C, Cárdenas A, C. Frommlet J, R Voolstra C. 2019. Protocol for the generation of axenic/bacteria-depleted Symbiodiniaceae cultures v1. *protocols.io* doi:10.17504/protocols.io.87khzkw
- Cowman PF, Quattrini AM, Bridge TCL, Watkins-Colwell GJ, Fadli N, Grinblat M, Roberts TE, McFadden CS, Miller DJ, Baird AH. 2020. An enhanced target-enrichment bait set for Hexacorallia provides phylogenomic resolution of the staghorn corals (Acroporidae) and close relatives. *Mol Phylogenet Evol* **153**:106944.
- Cumbo VR, Baird AH, van Oppen MJH. 2013. The promiscuous larvae: flexibility in the establishment of symbiosis in corals. *Coral Reefs* **32**:111–120.
- Cunning R, Baker AC. 2013. Excess algal symbionts increase the susceptibility of reef corals to bleaching. *Nature Climate Change* **3**:259–262.
- Cunning R, Gates RD, Edmunds PJ. 2017. Using high-throughput sequencing of ITS2 to describe *Symbiodinium* metacommunities in St. John, US Virgin Islands. *PeerJ* **5**:e3472.
- Cunning R, Glynn PW, Baker AC. 2013. Flexible associations between *Pocillopora* corals and *Symbiodinium* limit utility of symbiosis ecology in defining species. *Coral Reefs*.
- Cunning R, Parker KE, Johnson-Sapp K, Karp RF, Wen AD, Williamson OM, Bartels E, D'Alessandro M, Gilliam DS, Hanson G, Levy J, Lirman D, Maxwell K, Million WC, Moulding AL, Moura A, Muller EM, Nedimyer K, Reckenbeil B, van Hoodonk R, Dahlgren C, Kenkel C, Parkinson JE, Baker AC. 2021. Census of heat tolerance among Florida's threatened staghorn corals finds resilient individuals throughout existing nursery populations. *Proc Biol Sci* **288**:20211613.
- Cunning R, Silverstein RN, Baker AC. 2018. Symbiont shuffling linked to differential photochemical dynamics of *Symbiodinium* in three Caribbean reef corals. *Coral Reefs* **37**:145–152.
- Cunning R, Yost DM, Guarinello ML, Putnam HM, Gates RD. 2015. Variability of *Symbiodinium* communities in waters, sediments, and corals of thermally distinct reef pools in American Samoa. *PLoS One* **10**:e0145099.
- Cziesielski MJ, Liew YJ, Cui G, Schmidt-Roach S, Campana S, Marondedze C, Aranda M. 2018. Multi-omics analysis of thermal stress response in a zooxanthellate cnidarian reveals the importance of associating with thermotolerant symbionts. *Proc Biol Sci* **285**:20172654.
- Davies SW, Moreland KN, Wham DC, Kanke MR, Matz MV. 2020. *Cladocopium* community divergence in two *Acropora* coral hosts across multiple spatial scales. *Mol Ecol* **29**:4559–4572.
- Davies SW, Putnam HM, Ainsworth T, Baum JK, Bove CB, Crosby SC, Côté IM, Duploux A, Fulweiler RW, Griffin AJ, Hanley TC, Hill T, Humanes A, Mangubhai S, Metaxas A, Parker LM, Rivera HE, Silbiger NJ, Smith NS, Spalding AK, Traylor-Knowles N, Weigel BL, Wright RM, Bates AE. 2021. Promoting inclusive metrics of success and impact to dismantle a discriminatory reward system in science. *PLoS Biol* **19**:e3001282.
- Davies SW, Rahman M, Meyer E, Green EA, Buschiazzi E, Medina M, Matz MV. 2013. Novel polymorphic microsatellite markers for population genetics of the endangered Caribbean star coral, *Montastraea faveolata*. *Mar Biodivers* **43**:167–172.
- Davies SW, Ries JB, Marchetti A, Castillo KD. 2018. *Symbiodinium* functional diversity in the coral *Siderastrea siderea* is influenced by thermal stress and reef environment, but not ocean acidification. *Front Mar Sci* **5**:150.
- Davy Simon K., Allemand Denis, Weis Virginia M. 2012. Cell biology of cnidarian-dinoflagellate symbiosis. *Microbiol Mol Biol Rev* **76**:229–261.
- de Mendoza A, Bonnet A, Vargas-Landin DB, Ji N, Li H, Yang F, Li L, Hori K, Pflueger J, Buckberry S, Ohta H, Rosic N, Lesage P, Lin S, Lister R. 2018. Recurrent acquisition of cytosine methyltransferases into eukaryotic retrotransposons. *Nat Commun* **9**:1341.
- De Queiroz K. 2007. Species concepts and species delimitation. *Syst Biol* **56**:879–886.
- de Vos A. 2020. The problem of “colonial science.” *Scientific American*.
- Díaz-Almeyda EM, Prada C, Ohdera AH, Moran H, Civitello DJ, Iglesias-Prieto R, Carlo TA, LaJeunesse TC, Medina M. 2017. Intraspecific and interspecific variation in thermotolerance and photoacclimation in *Symbiodinium* dinoflagellates. *Proc Biol Sci* **284**:20171767.

- Di Bitetti MS, Ferreras JA. 2017. Publish (in English) or perish: The effect on citation rate of using languages other than English in scientific publications. *Ambio* **46**:121–127.
- Di Genio S, Wang L-H, Meng P-J, Tsai S, Lin C. 2021. “Symbio-Cryobank”: Toward the development of a cryogenic archive for the coral reef dinoflagellate symbiont Symbiodiniaceae. *Biopreserv Biobank* **19**:91–93.
- Dimond JL, Holzman BJ, Bingham BL. 2012. Thicker host tissues moderate light stress in a cnidarian endosymbiont. *J Exp Biol* **215**:2247–2254.
- Dimond JL, Kerwin AH, Rotjan R, Sharp K, Stewart FJ, Thornhill DJ. 2013. A simple temperature-based model predicts the upper latitudinal limit of the temperate coral *Astrangia poculata*. *Coral Reefs* **32**:401–409.
- Dixon GB, Bay LK, Matz MV. 2014. Bimodal signatures of germline methylation are linked with gene expression plasticity in the coral *Acropora millepora*. *BMC Genomics* **15**:1109.
- Dixon G, Liao Y, Bay LK, Matz MV. 2018. Role of gene body methylation in acclimatization and adaptation in a basal metazoan. *Proc Natl Acad Sci USA* **115**:13342–13346.
- Dixon G, Matz M. 2021. Benchmarking DNA methylation assays in a reef-building coral. *Mol Ecol Resour* **21**:464–477.
- Dougan KE, González-Pech RA, Stephens TG, Shah S, Chen Y, Ragan MA, Bhattacharya D, Chan CX. 2022. Genome-powered classification of microbial eukaryotes: focus on coral algal symbionts. *Trends Microbiol*.
- Durante MK, Baums IB, Williams DE, Kemp DW, Vohsen S. 2019. What drives phenotypic divergence among coral clonemates of *Acropora palmata*? *Mol Ecol*. **28**:3208–3224.
- Dzirasa K. 2020. Revising the *a priori* hypothesis: Systemic racism has penetrated scientific funding. *Cell* **183**:576–579.
- Eaton AA, Saunders JF, Jacobson RK, West K. 2020. How gender and race stereotypes impact the advancement of scholars in STEM: Professors’ biased evaluations of physics and biology post-doctoral candidates. *Sex Roles* **82**:127–141.
- Ebert P, Audano PA, Zhu Q, Rodriguez-Martin B, Porubsky D, Bonder MJ, Sulovari A, Ebler J, Zhou W, Serra Mari R, Yilmaz F, Zhao X, Hsieh P, Lee J, Kumar S, Lin J, Rausch T, Chen Y, Ren J, Santamarina M, Höps W, Ashraf H, Chuang NT, Yang X, Munson KM, Lewis AP, Fairley S, Tallon LJ, Clarke WE, Basile AO, Byrska-Bishop M, Corvelo A, Evani US, Lu T-Y, Chaisson MJP, Chen J, Li C, Brand H, Wenger AM, Ghareghani M, Harvey WT, Raeder B, Hasenfeld P, Regier AA, Abel HJ, Hall IM, Flicek P, Stegle O, Gerstein MB, Tubio JMC, Mu Z, Li YI, Shi X, Hastie AR, Ye K, Chong Z, Sanders AD, Zody MC, Talkowski ME, Mills RE, Devine SE, Lee C, Korbel JO, Marschall T, Eichler EE. 2021. Haplotype-resolved diverse human genomes and integrated analysis of structural variation. *Science* **372**:eabf7117.
- Eckert RJ, Reaume AM, Sturm AB, Studivan MS, Voss JD. 2020. Depth influences Symbiodiniaceae associations among *Montastraea cavernosa* corals on the Belize Barrier Reef. *Front Microbiol* **11**:518.
- Egan S, Gardiner M. 2016. Microbial dysbiosis: Rethinking disease in marine ecosystems. *Front Microbiol* **7**:991.
- Enríquez S, Méndez ER, Iglesias-Prieto R. 2005. Multiple scattering on coral skeletons enhances light absorption by symbiotic algae. *Limnol Oceanogr* **50**:1025–1032.
- Epstein HE, Torda G, van Oppen MJH. 2019. Relative stability of the *Pocillopora acuta* microbiome throughout a thermal stress event. *Coral Reefs* **38**:373–386.
- Faktorová D, Nisbet RER, Fernández Robledo JA, Casacuberta E, Sudek L, Allen AE, Ares M Jr, Aresté C, Balestreri C, Barbrook AC, Beardslee P, Bender S, Booth DS, Bouget F-Y, Bowler C, Breglia SA, Brownlee C, Burger G, Cerutti H, Cesaroni R, Chiurillo MA, Clemente T, Coles DB, Collier JL, Cooney EC, Coyne K, Docampo R, Dupont CL, Edgcomb V, Einarsson E, Elustondo PA, Federici F, Freire-Beneitez V, Freyria NJ, Fukuda K, García PA, Girguis PR, Gomaa F, Gornik SG, Guo J, Hampl V, Hanawa Y, Haro-Contreras ER, Hehenberger E, Highfield A, Hirakawa Y, Hopes A, Howe CJ, Hu I, Ibañez J, Irwin NAT, Ishii Y, Janowicz NE, Jones AC, Kachale A, Fujimura-Kamada K, Kaur B, Kaye

- JZ, Kazana E, Keeling PJ, King N, Klobutcher LA, Lander N, Lassadi I, Li Z, Lin S, Lozano J-C, Luan F, Maruyama S, Matute T, Miceli C, Minagawa J, Moosburner M, Najle SR, Nanjappa D, Nimmo IC, Noble L, Novák Vanclová AMG, Nowacki M, Nuñez I, Pain A, Piersanti A, Pucciarelli S, Pyrih J, Rest JS, Rius M, Robertson D, Ruaud A, Ruiz-Trillo I, Sigg MA, Silver PA, Slamovits CH, Jason Smith G, Sprecher BN, Stern R, Swart EC, Tsaousis AD, Tsypin L, Turkewitz A, Turnšek J, Valach M, Vergé V, von Dassow P, von der Haar T, Waller RF, Wang L, Wen X, Wheeler G, Woods A, Zhang H, Mock T, Worden AZ, Lukeš J. 2020. Genetic tool development in marine protists: emerging model organisms for experimental cell biology. *Nat Methods* **17**:481–494.
- Feder ME, Walser J-C. 2005. The biological limitations of transcriptomics in elucidating stress and stress responses. *J Evol Biol* **18**:901–910.
- Feil R, Fraga MF. 2012. Epigenetics and the environment: emerging patterns and implications. *Nat Rev Genet* **13**:97–109.
- Fifer JE, Bui V, Berg JT, Kriefall N, Klepac C, Bentlage B, Davies SW. 2022. Microbiome structuring within a coral colony and along a sedimentation gradient. *Front Mar Sci* **8**:2071.
- Figuroa RI, Howe-Kerr LI, Correa AMS. 2021. Direct evidence of sex and a hypothesis about meiosis in Symbiodiniaceae. *Sci Rep* **11**:18838.
- Finney JC, Pettay DT, Sampayo EM, Warner ME, Oxenford HA, LaJeunesse TC. 2010. The relative significance of host–habitat, depth, and geography on the ecology, endemism, and speciation of coral endosymbionts in the genus *Symbiodinium*. *Microb Ecol* **60**:250–263.
- Fitt WK. 2000. Cellular growth of host and symbiont in a cnidarian-zooxanthellar symbiosis. *Biol Bull* **198**:110–120.
- Fitt WK, McFarland FK, Warner ME, Chilcoat GC. 2000. Seasonal patterns of tissue biomass and densities of symbiotic dinoflagellates in reef corals and relation to coral bleaching. *Limnol Oceanogr* **45**:677–685.
- Flowerdew J. 2001. Attitudes of journal editors to nonnative speaker contributions. *TESOL Quarterly* **35**:121–150.
- Foret S, Kucharski R, Pellegrini M, Feng S, Jacobsen SE, Robinson GE, Maleszka R. 2012. DNA methylation dynamics, metabolic fluxes, gene splicing, and alternative phenotypes in honey bees. *Proc Natl Acad Sci USA* **109**:4968–4973.
- Forsman ZH, Ritson-Williams R, Tisthammer KH, Knapp ISS, Toonen RJ. 2020. Host-symbiont coevolution, cryptic structure, and bleaching susceptibility, in a coral species complex (Scleractinia; Poritidae). *Sci Rep* **10**:1–12.
- Fox CW, Burns CS, Muncy AD, Meyer JA. 2017. Author-suggested reviewers: gender differences and influences on the peer review process at an ecology journal. *Funct Ecol* **31**:270–280.
- Fox CW, Duffy MA, Fairbairn DJ, Meyer JA. 2019. Gender diversity of editorial boards and gender differences in the peer review process at six journals of ecology and evolution. *Ecol Evol* **9**:13636–13649.
- Frade PR, De Jongh F, Vermeulen F, van Bleijswijk J, Bak RPM. 2008. Variation in symbiont distribution between closely related coral species over large depth ranges. *Mol Ecol* **17**:691–703.
- Freudenthal HD. 1962. *Symbiodinium* gen. nov. and *Symbiodinium microadriaticum* sp. nov., a Zooxanthella: taxonomy, life cycle, and morphology. *J Protozool* **9**:45–52.
- Frommlet JC, Iglesias-Rodríguez MD. 2008. Microsatellite genotyping of single cells of the dinoflagellate species *Lingulodinium polyedrum* (Dinophyceae): A novel approach for marine microbial population genetic studies. *J Phycol* **44**:1116–1125.
- Frøslev TG, Kjølner R, Bruun HH, Ejrnæs R, Brunbjerg AK, Pietroni C, Hansen AJ. 2017. Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nat Commun* **8**:1188.
- Fujise L, Nitschke MR, Frommlet JC, Serôdio J, Woodcock S, Ralph PJ, Suggett DJ. 2018. Cell cycle dynamics of cultured coral endosymbiotic microalgae (*Symbiodinium*) across different types (species) under alternate light and temperature conditions. *J Eukaryot Microbiol* **65**:505–517.
- Fujise L, Suggett DJ, Stat M, Kahlke T, Bunce M, Gardner SG, Goyen S, Woodcock S, Ralph PJ, Seymour JR, Siboni N, Nitschke MR. 2021. Unlocking the phylogenetic diversity, primary habitats, and abundances of free-living Symbiodiniaceae on a coral reef. *Mol Ecol* **30**:343–360.

- Fujiwara Y, Kawamura I, Reimer JD, Parkinson JE. 2021. Zoantharian endosymbiont community dynamics during a stress event. *Front Microbiol* **12**:1282.
- Gabay Y, Parkinson JE, Wilkinson SP, Weis VM, Davy SK. 2019. Inter-partner specificity limits the acquisition of thermotolerant symbionts in a model cnidarian-dinoflagellate symbiosis. *The ISME Journal* **13**:2489–2499.
- Ginther DK, Schaffer WT, Schnell J, Masimore B, Liu F, Haak LL, Kington R. 2011. Race, ethnicity, and NIH research awards. *Science* **333**:1015–1019.
- Glynn PW. 1996. Coral reef bleaching: Facts, hypotheses and implications. *Glob Chang Biol* **2**:495–509.
- Gong W, Marchetti A. 2019. Estimation of 18S gene copy number in marine eukaryotic plankton using a next-generation sequencing approach. *Front Mar Sci* **6**:219.
- González-Pech RA, Bhattacharya D, Ragan MA, Chan CX. 2019. Genome evolution of coral reef symbionts as intracellular residents. *Trends Ecol Evol* **34**:799–806.
- González-Pech RA, Stephens TG, Chen Y, Mohamed AR, Cheng Y, Shah S, Dougan KE, Fortuin MDA, Lagorce R, Burt DW, Bhattacharya D, Ragan MA, Chan CX. 2021. Comparison of 15 dinoflagellate genomes reveals extensive sequence and structural divergence in family Symbiodiniaceae and genus *Symbiodinium*. *BMC Biol* **19**:73.
- Gordin MD. 2015. Scientific Babel: How Science Was Done Before and After Global English. University of Chicago Press.
- Goulet T, Coffroth M. 2003. Genetic composition of zooxanthellae between and within colonies of the octocoral *Plexaura kuna*, based on small subunit rDNA and multilocus DNA fingerprinting. *Mar Biol* **142**:233–239.
- Goulet TL. 2006. Most corals may not change their symbionts. *Mar Ecol Prog Ser* **321**:1–7.
- Goulet TL, Cook CB, Goulet D. 2005. Effect of short-term exposure to elevated temperatures and light levels on photosynthesis of different host-symbiont combinations in the *Aiptasia pallida*/*Symbiodinium* symbiosis. *Limnol Oceanogr* **50**:1490–1498.
- Goulet TL, Lucas MQ, Schizas NV. 2019. Symbiodiniaceae genetic diversity and symbioses with hosts from shallow to mesophotic coral ecosystems In: Loya Y, Puglise KA, Bridge TCL, editors. Mesophotic Coral Ecosystems. Cham: Springer International Publishing. pp. 537–551.
- Goulet TL, Shirur KP, Ramsby BD, Iglesias-Prieto R. 2017. The effects of elevated seawater temperatures on Caribbean gorgonian corals and their algal symbionts, *Symbiodinium* spp. *PLoS One* **12**:e0171032.
- Goyen S, Pernice M, Szabó M, Warner ME, Ralph PJ, Suggett DJ. 2017. A molecular physiology basis for functional diversity of hydrogen peroxide production amongst *Symbiodinium* spp. (Dinophyceae). *Mar Biol* **164**:1–12.
- Grajales A, Rodríguez E, Thornhill DJ. 2016. Patterns of *Symbiodinium* spp. associations within the family Aiptasiidae, a monophyletic lineage of symbiotic of sea anemones (Cnidaria, Actiniaria). *Coral Reefs* **35**:345–355.
- Granados-Cifuentes C, Neigel J, Leberg P, Rodriguez-Lanetty M. 2015. Genetic diversity of free-living *Symbiodinium* in the Caribbean: The importance of habitats and seasons. *Coral Reefs* **34**:927–939.
- Green EA, Davies SW, Matz MV, Medina M. 2014. Quantifying cryptic *Symbiodinium* diversity within *Orbicella faveolata* and *Orbicella franksi* at the Flower Garden Banks, Gulf of Mexico. *PeerJ* **2**:e386.
- Grégoire V, Schmacka F, Coffroth MA, Karsten U. 2017. Photophysiological and thermal tolerance of various genotypes of the coral endosymbiont *Symbiodinium* sp. (Dinophyceae). *J Appl Phyc* **29**:1893–1905.
- Greider CW, Sheltzer JM, Cantalupo NC, Copeland WB, Dasgupta N, Hopkins N, Jansen JM, Joshua-Tor L, McDowell GS, Metcalf JL, McLaughlin B, Olivarius A, O’Shea EK, Raymond JL, Ruebain D, Steitz JA, Stillman B, Tilghman SM, Valian V, Villa-Komaroff L, Wong JY. 2019. Increasing gender diversity in the STEM research workforce. *Science* **366**:692–695.
- Grinnell M, Higgins S, Yost K, Ochuba O, Lobl M, Grimes P, Wysong A. 2020. The proportion of male and female editors in women’s health journals: A critical analysis and review of the sex gap. *Int J Womens Dermatol* **6**:7–12.
- Grupstra CGB, Coma R, Ribes M, Leydet KP, Parkinson JE, McDonald K, Catllà M, Voolstra CR, Hellberg

- ME, Coffroth MA. 2017. Evidence for coral range expansion accompanied by reduced diversity of *Symbiodinium* genotypes. *Coral Reefs* **36**:981–985.
- Grupstra CGB, Howe-Kerr LI, Veglia AJ, Bryant RL, Coy SR, Blackwelder PL, Correa AMS. 2022a. Thermal stress triggers productive viral infection of a key coral reef symbiont. *ISME J* **16**:1430–1441.
- Grupstra CGB, Lemoine NP, Cook C, Correa AMS. 2022b. Thank you for biting: Dispersal of beneficial microbiota through “antagonistic” interactions. *Trends Microbiol.* doi:10.1016/j.tim.2022.03.006
- Grupstra CGB, Rabbitt KM, Howe-Kerr LI, Correa AMS. 2021. Fish predation on corals promotes the dispersal of coral symbionts. *Anim Microbiome* **3**:25.
- Haelewaters D, Hofmann TA, Romero-Olivares AL. 2021. Ten simple rules for Global North researchers to stop perpetuating helicopter research in the Global South. *PLoS Comput Biol* **17**:e1009277.
- Hafeez DM, Waqas A, Majeed S, Naveed S, Afzal KI, Aftab Z, Zeshan M, Khosa F. 2019. Gender distribution in psychiatry journals’ editorial boards worldwide. *Compr Psychiatry* **94**:152119.
- Hale ML, Burg TM, Steeves TE. 2012. Sampling for microsatellite-based population genetic studies: 25 to 30 individuals per population is enough to accurately estimate allele frequencies. *PLoS One* **7**:e45170.
- Hawkins TD, Hagemeyer JCG, Warner ME. 2016. Temperature moderates the infectiousness of two conspecific *Symbiodinium* strains isolated from the same host population. *Environ Microbiol* **18**:5204–5217.
- Hennige SJ, Suggett DJ, Warner ME, McDougall KE, Smith DJ. 2009. Photobiology of *Symbiodinium* revisited: bio-physical and bio-optical signatures. *Coral Reefs* **28**:179–195.
- Hernandez-Agreda A, Gates RD, Ainsworth TD. 2017. Defining the core microbiome in corals’ microbial soup. *Trends Microbiol* **25**:125–140.
- Hill M, Allenby A, Ramsby B, Schönberg C, Hill A. 2011. *Symbiodinium* diversity among host clonoid sponges from Caribbean and Pacific reefs: Evidence of heteroplasmy and putative host-specific symbiont lineages. *Mol Phylogenet Evol* **59**:81–88.
- Hirose M, Reimer JD, Hidaka M, Suda S. 2008. Phylogenetic analyses of potentially free-living *Symbiodinium* spp. isolated from coral reef sand in Okinawa, Japan. *Mar Biol* **155**:105–112.
- Hoadley KD, Lewis AM, Wham DC, Pettay DT, Grasso C, Smith R, Kemp DW, LaJeunesse TC, Warner ME. 2019. Host-symbiont combinations dictate the photo-physiological response of reef-building corals to thermal stress. *Sci Rep* **9**:9985.
- Hoadley KD, Pettay DT, Lewis A, Wham D, Grasso C, Smith R, Kemp DW, LaJeunesse T, Warner ME. 2021. Different functional traits among closely related algal symbionts dictate stress endurance for vital Indo-Pacific reef-building corals. *Glob Chang Biol* **27**:5295–5309.
- Hofstra B, Kulkarni VV, Munoz-Najar Galvez S, He B, Jurafsky D, McFarland DA. 2020. The diversity-innovation paradox in science. *Proc Natl Acad Sci USA* **117**:9284–9291.
- Hoppe TA, Litovitz A, Willis KA, Meseroll RA, Perkins MJ, Hutchins BI, Davis AF, Lauer MS, Valentine HA, Anderson JM, Santangelo GM. 2019. Topic choice contributes to the lower rate of NIH awards to African-American/black scientists. *Sci Adv* **5**:eaaw7238.
- Howe-Kerr LI, Bachelot B, Wright RM, Kenkel CD, Bay LK, Correa AMS. 2020. Symbiont community diversity is more variable in corals that respond poorly to stress. *Glob Chang Biol* **26**:2220–2234.
- Howells EJ, Abrego D, Meyer E, Kirk NL, Burt JA. 2016. Host adaptation and unexpected symbiont partners enable reef-building corals to tolerate extreme temperatures. *Glob Chang Biol* **22**:2702–2714.
- Howells EJ, Bauman AG, Vaughan GO, Hume BCC, Voolstra CR, Burt JA. 2020. Corals in the hottest reefs in the world exhibit symbiont fidelity not flexibility. *Mol Ecol* **29**:889–911.
- Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH. 2011. Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat Clim Chang* **2**:116–120.
- Howells EJ, Berkelmans R, van Oppen MJH, Willis BL, Bay LK. 2013a. Historical thermal regimes define limits to coral acclimatization. *Ecology* **94**:1078–1088.
- Howells EJ, van Oppen MJH, Willis BL. 2009. High genetic differentiation and cross-shelf patterns of genetic diversity among Great Barrier Reef populations of *Symbiodinium*. *Coral Reefs* **28**:215–225.
- Howells EJ, Willis BL, Bay LK, van Oppen MJH. 2016. Microsatellite allele sizes alone are insufficient to

- delineate species boundaries in *Symbiodinium*. *Mol Ecol* **25**:2719–2723.
- Howells EJ, Willis BL, Bay LK, van Oppen MJH. 2013b. Spatial and temporal genetic structure of *Symbiodinium* populations within a common reef-building coral on the Great Barrier Reef. *Mol Ecol* **22**:3693–3708.
- Huang J, Gates AJ, Sinatra R, Barabási A-L. 2020. Historical comparison of gender inequality in scientific careers across countries and disciplines. *Proc Natl Acad Sci USA* **117**:4609–4616.
- Hughes TP, Barnes ML, Bellwood DR, Cinner JE, Cumming GS, Jackson JBC, Kleypas J, van de Leemput IA, Lough JM, Morrison TH, Palumbi SR, van Nes EH, Scheffer M. 2017. Coral reefs in the Anthropocene. *Nature* **546**:82–90.
- Hume BCC, D'Angelo C, Smith EG, Stevens JR, Burt J, Wiedenmann J. 2015. *Symbiodinium thermophilum* sp. nov., a thermotolerant symbiotic alga prevalent in corals of the world's hottest sea, the Persian/Arabian Gulf. *Sci Rep* **5**:8562.
- Hume BCC, Mejia-Restrepo A, Voolstra CR, Berumen ML. 2020. Fine-scale delineation of Symbiodiniaceae genotypes on a previously bleached central Red Sea reef system demonstrates a prevalence of coral host-specific associations. *Coral Reefs* **39**:583–601.
- Hume BCC, Smith EG, Ziegler M, Warrington HJM, Burt JA, LaJeunesse TC, Wiedenmann J, Voolstra CR. 2019. SymPortal: A novel analytical framework and platform for coral algal symbiont next-generation sequencing ITS2 profiling. *Mol Ecol Resour* **19**:1063–1080.
- Hume BCC, Ziegler M, Poulain J, Pochon X, Romac S, Boissin E, de Vargas C, Planes S, Wincker P, Voolstra CR. 2018. An improved primer set and amplification protocol with increased specificity and sensitivity targeting the *Symbiodinium* ITS2 region. *PeerJ* **6**:e4816.
- Innis T, Cunning R, Ritson-Williams R, Wall CB, Gates RD. 2018. Coral color and depth drive symbiosis ecology of *Montipora capitata* in Kāneʻohe Bay, Oʻahu, Hawaiʻi. *Coral Reefs* **37**:423–430.
- Ishikura M, Hagiwara K, Takishita K, Haga M, Iwai K, Maruyama T. 2004. Isolation of new *Symbiodinium* strains from tridacnid giant clam (*Tridacna crocea*) and sea slug (*Pteraeolidia ianthina*) using culture medium containing giant clam tissue homogenate. *Mar Biotechnol* **6**:378–385.
- IUCN. 2021. The IUCN Red List of Threatened Species. <https://www.iucnredlist.org/resources/tax-sources>
- Jeong HJ, Lee SY, Kang NS, Yoo YD, Lim AS, Lee MJ, Kim HS, Yih W, Yamashita H, LaJeunesse TC. 2014. Genetics and morphology characterize the dinoflagellate *Symbiodinium voratum*, n. sp., (Dinophyceae) as the sole representative of *Symbiodinium* Clade E. *J Eukaryot Microbiol* **61**:75–94.
- Jinkerson RE, Russo JA, Newkirk CR, Kirk AL, Chi RJ, Martindale MQ, Grossman AR, Hatta M, Xiang T. 2022. Cnidarian-Symbiodiniaceae symbiosis establishment is independent of photosynthesis. *Curr Biol*. doi:10.1016/j.cub.2022.04.021
- Jones AM, Berkelmans R, van Oppen MJH, Mieog JC, Sinclair W. 2008. A community change in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field evidence of acclimatization. *Proc Biol Sci* **275**:1359–1365.
- Kareiva P, Levin SA. 2015. The Importance of Species. Princeton University Press.
- Karst SM, Ziels RM, Kirkegaard RH, Sørensen EA, McDonald D, Zhu Q, Knight R, Albertsen M. 2021. High-accuracy long-read amplicon sequences using unique molecular identifiers with Nanopore or PacBio sequencing. *Nat Methods* **18**:165–169.
- Kavousi J, Denis V, Sharp V, Reimer JD, Nakamura T, Parkinson JE. 2020. Unique combinations of coral host and algal symbiont genotypes reflect intraspecific variation in heat stress responses among colonies of the reef-building coral, *Montipora digitata*. *Mar Biol* **167**:23.
- Kemp DW, Fitt WK, Schmidt GW. 2008. A microsampling method for genotyping coral symbionts. *Coral Reefs* **27**:289–293.
- Kiers ET, Denison RF, Kawakita A, Herre EA. 2011. The biological reality of host sanctions and partner fidelity. *Proc Natl Acad Sci USA* **108**:E7.
- Kiers ET, Rousseau RA, West SA, Denison RF. 2003. Host sanctions and the legume-rhizobium mutualism. *Nature* **425**:78–81.
- Kirk NL, Andras JP, Drew Harvell C, Santos SR, Coffroth MA. 2009. Population structure of *Symbiodinium*

- sp. associated with the common sea fan, *Gorgonia ventalina*, in the Florida Keys across distance, depth, and time. *Mar Biol* **156**:1609–1623.
- Kitchen SA, Von Kuster G, Kuntz KLV, Reich HG, Miller W, Griffin S, Fogarty ND, Baums IB. 2020. STAGdb: a 30K SNP genotyping array and Science Gateway for *Acropora* corals and their dinoflagellate symbionts. *Sci Rep* **10**:12488.
- Klueter A, Crandall JB, Archer FI, Teece MA, Coffroth MA. 2015. Taxonomic and environmental variation of metabolite profiles in marine dinoflagellates of the genus *Symbiodinium*. *Metabolites* **5**:74–99.
- Klueter A, Trapani J, Archer FI, McIlroy SE, Coffroth MA. 2017. Comparative growth rates of cultured marine dinoflagellates in the genus *Symbiodinium* and the effects of temperature and light. *PLoS One* **12**:e0187707.
- Konopka A. 2009. What is microbial community ecology? *ISME J* **3**:1223–1230.
- Krassowski M, Das V, Sahu SK, Misra BB. 2020. State of the field in multi-omics research: From computational needs to data mining and sharing. *Front Genet* **11**:610798.
- Kriefall NG, Kanke MR, Aglyamova GV, Davies SW. 2022. Reef environments shape microbial partners in a highly connected coral population. *Proc Biol Sci* **289**:20212459.
- Krueger T, Gates RD. 2012. Cultivating endosymbionts — Host environmental mimics support the survival of *Symbiodinium* C15 *ex hospite*. *J Exp Mar Bio Ecol* **413**:169–176.
- Kuguru B, Chadwick NE, Achituv Y, Zandbank K, Tchernov D. 2008. Mechanisms of habitat segregation between corallimorpharians: photosynthetic parameters and *Symbiodinium* types. *Mar Ecol Prog Ser* **369**:115–129.
- Kunihiro S, Reimer JD. 2018. Phylogenetic analyses of *Symbiodinium* isolated from *Waminoa* and their anthozoan hosts in the Ryukyu Archipelago, southern Japan. *Symbiosis* **76**:253–264.
- LaJeunesse TC. 2017. Validation and description of *Symbiodinium microadriaticum*, the type species of *Symbiodinium* (Dinophyta). *J Phycol* **53**:1109–1114.
- LaJeunesse TC. 2001. Investigating the biodiversity, ecology, and phylogeny of endosymbiotic dinoflagellates in the genus *Symbiodinium* using the ITS region: In search of a “species” level marker. *J Phycol* **37**:866–880.
- LaJeunesse TC, Lee SY, Gil-Agudelo DL. 2015. *Symbiodinium necroappetens* sp. nov. (Dinophyceae): an opportunist “zooxanthella” found in bleached and diseased tissues of Caribbean reef corals. *European Journal of Phycology*.
- LaJeunesse TC, Loh W, Trench RK. 2009. Do introduced endosymbiotic dinoflagellates “take” to new hosts? *Biol Invasions* **11**:995–1003.
- LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR. 2018. Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr Biol* **28**:2570–2580.
- LaJeunesse TC, Parkinson JE, Reimer JD. 2012. A genetics-based description of *Symbiodinium minutum* sp. nov. and *S. psygmophilum* sp. nov. (Dinophyceae), two dinoflagellates symbiotic with cnidaria. *J Phycol* **48**:1380–1391.
- LaJeunesse TC, Pettay DT, Sampayo EM, Phongsuwan N, Brown B, Obura DO, Hoegh-Guldberg O, Fitt WK. 2010a. Long-standing environmental conditions, geographic isolation and host-symbiont specificity influence the relative ecological dominance and genetic diversification of coral endosymbionts in the genus *Symbiodinium*. *J Biogeogr* **37**:785–800.
- LaJeunesse TC, Smith RT, Walther M, Pinzón JH, Pettay DT, McGinley MP, Aschaffenburg MD, Medina-Rosas P, Cupul-Magaña AL, Pérez AL, Reyes-Bonilla H, Warner ME. 2010b. Host-symbiont recombination versus natural selection in the response of coral-dinoflagellate symbioses to environmental disturbance. *Proc Bio Sci* **277**:2925–2934.
- LaJeunesse TC, Thornhill DJ. 2011. Improved resolution of reef-coral endosymbiont (*Symbiodinium*) species diversity, ecology, and evolution through *psbA* non-coding region genotyping. *PLoS One* **6**:e29013.
- LaJeunesse TC, Trench RK. 2000. Biogeography of two species of *Symbiodinium* (Freudenthal) inhabiting the intertidal sea anemone *Anthopleura elegantissima* (Brandt). *Biol Bull* **199**:126–134.

- LaJeunesse TC, Wham DC, Pettay DT, Parkinson JE. 2014. Ecologically differentiated stress-tolerant endosymbionts in the dinoflagellate genus *Symbiodinium* (Dinophyceae) Clade D are different species. *Phycologia* **53**:305–319.
- LaJeunesse TC, Wiedenmann J, Casado-Amezúa P, D’Ambra I, Turnham KE, Nitschke MR, Oakley CA, Goffredo S, Spano CA, Cubillos VM, Davy SK, Suggett DJ. 2021. Revival of *Philozoon* Geddes for host-specialized dinoflagellates, “zooxanthellae”, in animals from coastal temperate zones of northern and southern hemispheres. *Eur J Phycol* **57**:166-180.
- Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**:559.
- Lawson CA, Possell M, Seymour JR, Raina J-B, Suggett DJ. 2019. Coral endosymbionts (Symbiodiniaceae) emit species-specific volatiles that shift when exposed to thermal stress. *Sci Rep* **9**:17395.
- Lawson CA, Raina J-B, Kahlke T, Seymour JR, Suggett DJ. 2018. Defining the core microbiome of the symbiotic dinoflagellate, *Symbiodinium*. *Environ Microbiol Rep* **10**:7–11.
- Lee MJ, Jeong HJ, Jang SH, Lee SY, Kang NS, Lee KH, Kim HS, Wham DC, LaJeunesse TC. 2016. Most low-abundance “background” *Symbiodinium* spp. are transitory and have minimal functional significance for symbiotic corals. *Microb Ecol* **71**:771–783.
- Lee SY, Jeong HJ, Kang NS, Jang TY. 2015. *Symbiodinium tridacnidorum* sp. nov., a dinoflagellate common to Indo-Pacific giant clams, and a revised morphological description of *Symbiodinium microadriaticum*. *Eur J Phycol* **50**:155-172.
- Lee SY, Jeong HJ, LaJeunesse TC. 2020. *Cladocopium infistulum* sp. nov. (Dinophyceae), a thermally tolerant dinoflagellate symbiotic with giant clams from the western Pacific Ocean. *Phycologia* **59**:515–526.
- Leliaert F, Verbruggen H, Vanormelingen P, Steen F, López-Bautista JM, Zuccarello GC, De Clerck O. 2014. DNA-based species delimitation in algae. *Eur J Phycol* **49**:179–196.
- Levin RA, Voolstra CR, Weynberg KD, van Oppen MJH. 2017. Evidence for a role of viruses in the thermal sensitivity of coral photosymbionts. *The ISME Journal* **11**:808–812.
- Levy S, Elek A, Grau-Bové X, Menéndez-Bravo S, Iglesias M, Tanay A, Mass T, Sebé-Pedrós A. 2021. A stony coral cell atlas illuminates the molecular and cellular basis of coral symbiosis, calcification, and immunity. *Cell* **184**:2973–2987.e18.
- Lewis AM, Chan AN, LaJeunesse TC. 2019. New species of closely related endosymbiotic dinoflagellates in the greater Caribbean have niches corresponding to host coral phylogeny. *J Eukaryot Microbiol* **66**:469–482.
- Lewis C, Neely K, Rodriguez-Lanetty M. 2019. Recurring episodes of thermal stress shift the balance from a dominant host-specialist to a background host-generalist zooxanthella in the threatened pillar coral, *Dendrogyra cylindrus*. *Front Mar Sci* **6**:5.
- Li T, Yu L, Song B, Song Y, Li L, Lin X, Lin S. 2020. Genome improvement and core gene set refinement of *Fugacium kawagutii*. *Microorganisms* **8**:102.
- Liang J, Luo W, Yu K, Xu Y, Chen J, Deng C, Ge R, Su H, Huang W, Wang G. 2021. Multi-omics revealing the response patterns of symbiotic microorganisms and host metabolism in scleractinian coral *Pavona minuta* to temperature stresses. *Metabolites* **12**:18.
- Lien Y-T, Fukami H, Yamashita Y. 2012. *Symbiodinium* Clade C dominates zooxanthellate corals (Scleractinia) in the temperate region of Japan. *Zool Sci* **29**:173–180.
- Lien Y-T, Nakano Y, Plathong S, Fukami H, Wang J-T, Chen CA. 2007. Occurrence of the putatively heat-tolerant *Symbiodinium* phylotype D in high-latitude outlying coral communities. *Coral Reefs* **26**:35–44.
- Liew YJ, Li Y, Baumgarten S, Voolstra CR, Aranda M. 2017. Condition-specific RNA editing in the coral symbiont *Symbiodinium microadriaticum*. *PLoS Genet* **13**:e1006619.
- Liew YJ, Zoccola D, Li Y, Tambutté E, Venn AA, Michell CT, Cui G, Deutekom ES, Kaandorp JA, Voolstra CR, Forêt S, Allemand D, Tambutté S, Aranda M. 2018. Epigenome-associated phenotypic acclimatization to ocean acidification in a reef-building coral. *Sci Adv* **4**:eaar8028.
- Lim SSQ, Huang D, Soong K, Neo ML. 2019. Diversity of endosymbiotic Symbiodiniaceae in giant clams at

- Dongsha Atoll, northern South China Sea. *Symbiosis* **78**:251–262.
- Lin S. 2011. Genomic understanding of dinoflagellates. *Res Microbiol* **162**:551–569.
- Lin S, Cheng S, Song B, Zhong X, Lin X, Li W, Li L, Zhang Y, Zhang H, Ji Z, Cai M, Zhuang Y, Shi X, Lin L, Wang L, Wang Z, Liu X, Yu S, Zeng P, Hao H, Zou Q, Chen C, Li Y, Wang Y, Xu C, Meng S, Xu X, Wang J, Yang H, Campbell DA, Sturm NR, Dagenais-Bellefeuille S, Morse D. 2015. The *Symbiodinium kawagutii* genome illuminates dinoflagellate gene expression and coral symbiosis. *Science* **350**:691–694.
- Lin S, Song B, Morse D. 2021. Spatial organization of dinoflagellate genomes: Novel insights and remaining critical questions. *J Phycol* **57**:1674–1678.
- Liu B, Hu Y, Hu Z, Liu G, Zhu H. 2020. Taxonomic scheme of the order Chaetophorales (Chlorophyceae, Chlorophyta) based on chloroplast genomes. *BMC Genomics* **21**:442.
- Liu H, Stephens TG, González-Pech RA, Beltran VH, Lapeyre B, Bongaerts P, Cooke I, Aranda M, Bourne DG, Forêt S, Miller DJ, van Oppen MJH, Voolstra CR, Ragan MA, Chan CX. 2018. *Symbiodinium* genomes reveal adaptive evolution of functions related to coral-dinoflagellate symbiosis. *Commun Biol* **1**:95.
- Lohuis MR, Miller DJ. 1998. Hypermethylation at CpG-motifs in the dinoflagellates *Amphidinium carterae* (Dinophyceae) and *Symbiodinium microadriaticum* (Dinophyceae): Evidence from restriction analyses, 5-azacytidine and ethionine treatment. *J Phycol* **34**:152–159.
- Loram JE, Boonham N, O'Toole P, Trapido-Rosenthal HG, Douglas AE. 2007. Molecular quantification of symbiotic dinoflagellate algae of the genus *Symbiodinium*. *Biol Bull* **212**:259–268.
- Lo R, Dougan KE, Chen Y, Shah S, Bhattacharya D, Chan CX. 2022. Alignment-free analysis of Symbiodiniaceae whole-genome sequences reveals differential phylogenetic signals in distinct regions. *Front Plant Sci* **1233**:815714.
- Lowndes JSS, Best BD, Scarborough C, Afflerbach JC, Frazier MR, O'Hara CC, Jiang N, Halpern BS. 2017. Our path to better science in less time using open data science tools. *Nat Ecol Evol* **1**:160.
- Madin JS, Anderson KD, Andreasen MH, Bridge TCL, Cairns SD, Connolly SR, Darling ES, Diaz M, Falster DS, Franklin EC, Gates RD, Harmer A, Hoogenboom MO, Huang D, Keith SA, Kosnik MA, Kuo C-Y, Lough JM, Lovelock CE, Luiz O, Martinelli J, Mizerek T, Pandolfi JM, Pochon X, Pratchett MS, Putnam HM, Roberts TE, Stat M, Wallace CC, Widman E, Baird AH. 2016. The Coral Trait Database, a curated database of trait information for coral species from the global oceans. *Sci Data* **3**:160017.
- Magalon H, Baudry E, Husté A, Adjeroud M, Veuille M. 2006. High genetic diversity of the symbiotic dinoflagellates in the coral *Pocillopora meandrina* from the South Pacific. *Mar Biol* **148**:913–922.
- Maire J, Girvan SK, Barkla SE, Perez-Gonzalez A, Suggett DJ, Blackall LL, van Oppen MJH. 2021. Intracellular bacteria are common and taxonomically diverse in cultured and *in hospite* algal endosymbionts of coral reefs. *ISME J* **15**:2028–2042.
- Manning MM, Gates RD. 2008. Diversity in populations of free-living *Symbiodinium* from a Caribbean and Pacific reef. *Limnol Oceanogr* **53**:1853–1861.
- Mansour JS, Pollock FJ, Díaz-Almeyda E, Iglesias-Prieto R, Medina M. 2018. Intra- and interspecific variation and phenotypic plasticity in thylakoid membrane properties across two *Symbiodinium* Clades. *Coral Reefs* **37**:841–850.
- Manzello DP, Matz MV, Enochs IC, Valentino L, Carlton RD, Kolodziej G, Serrano X, Towle EK, Jankulak M. 2018. Role of host genetics and heat-tolerant algal symbionts in sustaining populations of the endangered coral *Orbicella faveolata* in the Florida Keys with ocean warming. *Glob Chang Biol* **25**:1016–1031.
- Marcelino LA, Westneat MW, Stoyneva V, Henss J, Rogers JD, Radosevich A, Turzhitsky V, Siple M, Fang A, Swain TD, Fung J, Backman V. 2013. Modulation of light-enhancement to symbiotic algae by light-scattering in corals and evolutionary trends in bleaching. *PLoS One* **8**:e61492.
- Marhoefer SR, Zenger KR, Strugnell JM, Logan M, van Oppen MJH, Kenkel CD, Bay LK. 2021. Signatures of adaptation and acclimatization to reef flat and slope habitats in the coral *Pocillopora damicornis*. *Front Mar Sci* **8**:704709.

- Marinov GK, Trevino AE, Xiang T, Kundaje A, Grossman AR, Greenleaf WJ. 2021. Transcription-dependent domain-scale three-dimensional genome organization in the dinoflagellate *Breviolum minutum*. *Nat Genet* 1–5.
- Maruyama S, Weis VM. 2021. Limitations of using cultured algae to study cnidarian-algal symbioses and suggestions for future studies. *J Phycol* 57:30–38.
- Matthews JL, Crowder CM, Oakley CA, Lutz A, Roessner U, Meyer E, Grossman AR, Weis VM, Davy SK. 2017. Optimal nutrient exchange and immune responses operate in partner specificity in the cnidarian-dinoflagellate symbiosis. *Proc Natl Acad Sci USA* 114:13194–13199.
- Matthews JL, Raina J-B, Kahlke T, Seymour JR, van Oppen MJH, Suggett DJ. 2020. Symbiodiniaceae-bacteria interactions: Rethinking metabolite exchange in reef-building corals as multi-partner metabolic networks. *Environ Microbiol* 22:1675–1687.
- Mayr E. 1942. Systematics and the Origin of Species. New York: Columbia University Press.
- McIlroy SE, Cuning R, Baker AC, Coffroth MA. 2019. Competition and succession among coral endosymbionts. *Ecol Evol* 9:12767–12778.
- McIlroy SE, Smith GJ, Geller JB. 2014. FISH-Flow: A quantitative molecular approach for describing mixed clade communities of *Symbiodinium*. *Coral Reefs* 33:157–167.
- McIlroy SE, Wong JCY, Baker DM. 2020. Competitive traits of coral symbionts may alter the structure and function of the microbiome. *ISME J* 14:2424–2432.
- McKenna V, Archibald JM, Beinart R, Dawson MN, Hentschel U, Keeling PJ, Lopez JV, Martín-Durán JM, Petersen JM, Sigwart JD, Simakov O, Sutherland KR, Sweet M, Talbot N, Thompson AW, Bender S, Harrison PW, Rajan J, Cochran G, Berriman M, Lawniczak M, Blaxter M. 2021. The Aquatic Symbiosis Genomics Project: Probing the evolution of symbiosis across the tree of life. *Wellcome Open Res* 6:254.
- Méndez C, Ahlenstiel CL, Kelleher AD. 2015. Post-transcriptional gene silencing, transcriptional gene silencing and human immunodeficiency virus. *World J Virol* 4:219–244.
- Mieog JC, van Oppen MJH, Cantin NE, Stam WT, Olsen JL. 2007. Real-time PCR reveals a high incidence of *Symbiodinium* clade D at low levels in four scleractinian corals across the Great Barrier Reef: implications for symbiont shuffling. *Coral Reefs* 26:449–457.
- Mizuyama M, Iguchi A, Iijima M, Gibu K, Reimer JD. 2020. Comparison of Symbiodiniaceae diversities in different members of a *Palythoa* species complex (Cnidaria: Anthozoa: Zoantharia)-implications for ecological adaptations to different microhabitats. *PeerJ* 8:e8449.
- Molecular Ecology Resources Primer Development Consortium, Abdoullaye D, Acevedo I, Adebayo AA, Behrmann-Godel J, Benjamin RC, Bock DG, Born C, Brouat C, Caccone A, Cao L-Z, Casado-Amezúa P, Catanéo J, Correa-Ramirez MM, Cristescu ME, Dobigny G, Egbosimba EE, Etchberger LK, Fan B, Fields PD, Forcioli D, Furla P, Garcia de Leon FJ, García-Jiménez R, Gauthier P, Gergs R, González C, Granjon L, Gutiérrez-Rodríguez C, Havill NP, Helsen P, Hether TD, Hoffman EA, Hu X, Ingvarsson PK, Ishizaki S, Ji H, Ji XS, Jimenez ML, Kapil R, Karban R, Keller SR, Kubota S, Li S, Li W, Lim DD, Lin H, Liu X, Luo Y, Machordom A, Martin AP, Matthysen E, Mazzella MN, McGeoch MA, Meng Z, Nishizawa M, O'Brien P, Ohara M, Ornelas JF, Ortu MF, Pedersen AB, Preston L, Ren Q, Rothhaupt K-O, Sackett LC, Sang Q, Sawyer GM, Shiojiri K, Taylor DR, Van Dongen S, Van Vuuren BJ, Vandewoestijne S, Wang H, Wang JT, Wang LE, Xu X-L, Yang G, Yang Y, Zeng YQ, Zhang Q-W, Zhang Y, Zhao Y, Zhou Y. 2010. Permanent genetic resources added to Molecular Ecology Resources Database 1 August 2009–30 September 2009. *Mol Ecol Resour* 10:232–236.
- Montgomery BL. 2017a. From deficits to possibilities: Mentoring lessons from plants on cultivating individual growth through environmental assessment and optimization. *Manifold*. doi:10.25335/m5/ppj.1.1-3
- Montgomery BL. 2017b. Mapping a mentoring roadmap and developing a supportive network for strategic career advancement. *SAGE Open* 7:2158244017710288.
- Montgomery BL, Dodson JE, Johnson SM. 2014. Guiding the way: Mentoring graduate students and junior faculty for sustainable academic careers. *SAGE Open* 4:2158244014558043.
- Moore RB, Ferguson KM, Loh WKW, Hoegh-Guldberg O, Carter DA. 2003. Highly organized structure in the non-coding region of the *psbA* minicircle from Clade C *Symbiodinium*. *Int J Syst Evol Microbiol*

53:1725–1734.

- Mordret S, Romac S, Henry N, Colin S, Carmichael M, Berney C, Audic S, Richter DJ, Pochon X, de Vargas C, Decelle J. 2016. The symbiotic life of *Symbiodinium* in the open ocean within a new species of calcifying ciliate (*Tiarina* sp.). *ISME J* **10**:1424–1436.
- Mungpakdee S, Shinzato C, Takeuchi T, Kawashima T, Koyanagi R, Hisata K, Tanaka M, Goto H, Fujie M, Lin S, Satoh N, Shoguchi E. 2014. Massive gene transfer and extensive RNA editing of a symbiotic dinoflagellate plastid genome. *Genome Biol Evol* **6**:1408–1422.
- Muscantine L, Ferrier-Pagès C, Blackburn A, Gates RD, Baghdasarian G, Allemand D. 1998. Cell-specific density of symbiotic dinoflagellates in tropical anthozoans. *Coral Reefs* **17**:329–337.
- Naepi S, McAllister TG, Thomsen P, Leenen-Young M, Walker LA, McAllister AL, Theodore R, Kidman J, Suaaliia T. 2020. The pakaru “pipeline”: Māori and Pasifika pathways within the academy. *N Z Annu Rev Educ* **24**:142.
- Nand A, Zhan Y, Salazar OR, Aranda M, Voolstra CR, Dekker J. 2021. Genetic and spatial organization of the unusual chromosomes of the dinoflagellate *Symbiodinium microadriaticum*. *Nat Genet* **53**:618–629.
- Nielsen MW, Bloch CW, Schiebinger L. 2018. Making gender diversity work for scientific discovery and innovation. *Nat Hum Behav* **2**:726–734.
- Nitschke MR, Craveiro SC, Brandão C, Fidalgo C, Serôdio J, Calado AJ, Frommlet JC. 2020. Description of *Freudenthalidium* gen. nov. and *Halluxium* gen. nov. to formally recognize Clades Fr3 and H as genera in the family Symbiodiniaceae (Dinophyceae). *J Phycol* **56**:923–940.
- Nitschke MR, Davy SK, Ward S. 2016. Horizontal transmission of *Symbiodinium* cells between adult and juvenile corals is aided by benthic sediment. *Coral Reefs* **35**:335–344.
- Noda H, Parkinson JE, Yang S-Y, Reimer JD. 2017. A preliminary survey of zoantharian endosymbionts shows high genetic variation over small geographic scales on Okinawa-jima Island, Japan. *PeerJ* **5**:e3740.
- O’Brien LT, Bart HL, Garcia DM. 2020. Why are there so few ethnic minorities in ecology and evolutionary biology? Challenges to inclusion and the role of sense of belonging. *Soc Psychol Edu* **23**:449–477.
- Olander A, Lawson CA, Possell M, Raina J-B, Ueland M, Suggett DJ. 2021. Comparative volatilomics of coral endosymbionts from one- and comprehensive two-dimensional gas chromatography approaches. *Mar Biol* **168**:76.
- Palmer TM, Stanton ML, Young TP. 2003. Competition and coexistence: Exploring mechanisms that restrict and maintain diversity within mutualist guilds. *Am Nat* **162**:S63–79.
- Palser ER, Lazerwitz M, Fotopoulou A. 2022. Gender and geographical disparity in editorial boards of journals in psychology and neuroscience. *Nat Neurosci* **25**:272–279.
- Parker GM. 1984. Dispersal of zooxanthellae on coral reefs by predators on cnidarians. *Biol Bull* **167**:159–167.
- Parkinson JE, Baumgarten S, Michell CT, Baums IB, LaJeunesse TC, Voolstra CR. 2016. Gene expression variation resolves species and individual strains among coral-associated dinoflagellates within the genus *Symbiodinium*. *Genome Biol Evol* **8**:665–680.
- Parkinson JE, Baums IB. 2014. The extended phenotypes of marine symbioses: ecological and evolutionary consequences of intraspecific genetic diversity in coral–algal associations. *Front Microbiol* **5**:445.
- Parkinson JE, Coffroth MA, LaJeunesse TC. 2015. New species of Clade B *Symbiodinium* (Dinophyceae) from the greater Caribbean belong to different functional guilds: *S. aenigmaticum* sp. nov., *S. antillogorgium* sp. nov., *S. endomadracis* sp. nov., and *S. pseudominutum* sp. nov. *J Phycol* **51**:850–858.
- Pettay DT, LaJeunesse TC. 2009. Microsatellite loci for assessing genetic diversity, dispersal and clonality of coral symbionts in “stress-tolerant” Clade D *Symbiodinium*. *Molecular Ecology Resources* **9**:1022–1025.
- Pettay DT, LaJeunesse TC. 2007. Microsatellites from Clade B *Symbiodinium* spp. specialized for Caribbean corals in the genus *Madracis*. *Mol Ecol Notes* **7**:1271–1274.
- Pettay DT, Wham DC, Smith RT, Iglesias-Prieto R, LaJeunesse TC. 2015. Microbial invasion of the Caribbean by an Indo-Pacific coral zooxanthella. *Proc Natl Acad Sci USA* **112**:7513–7518.
- Pinho-Gomes A-C, Vassallo A, Thompson K, Womersley K, Norton R, Woodward M. 2021. Representation

- of women among editors in chief of leading medical journals. *JAMA Netw Open* **4**:e2123026.
- Pinzón JH, Devlin-Durante MK, Weber MX, Baums IB, LaJeunesse TC. 2011. Microsatellite loci for *Symbiodinium* A3 (*S. fitti*) a common algal symbiont among Caribbean *Acropora* (stony corals) and Indo-Pacific giant clams (*Tridacna*). *Conserv Genet Resour* **3**:45–47.
- Pochon X, Garcia-Cuetos L, Baker AC, Castella E, Pawlowski J. 2007. One-year survey of a single Micronesian reef reveals extraordinarily rich diversity of *Symbiodinium* types in soritid foraminifera. *Coral Reefs* **26**:867–882.
- Pochon X, LaJeunesse TC. 2021. *Miliolidium* n. gen, a new Symbiodiniacean genus whose members associate with soritid foraminifera or are free-living. *J Eukaryot Microbiol* **68**:e12856.
- Pochon X, Montoya-Burgos JI, Stadelmann B, Pawlowski J. 2006. Molecular phylogeny, evolutionary rates, and divergence timing of the symbiotic dinoflagellate genus *Symbiodinium*. *Mol Phylogenet Evol* **38**:20–30.
- Pochon X, Putnam HM, Burki F, Gates RD. 2012. Identifying and characterizing alternative molecular markers for the symbiotic and free-living dinoflagellate genus *Symbiodinium*. *PLoS One* **7**:e29816.
- Pochon X, Putnam HM, Gates RD. 2014. Multi-gene analysis of *Symbiodinium* dinoflagellates: A perspective on rarity, symbiosis, and evolution. *PeerJ* **2**:e394.
- Pochon X, Stat M, Takabayashi M, Chasqui L, Chauka LJ, Logan DDK, Gates RD. 2010. Comparison of endosymbiotic and free-living *Symbiodinium* (Dinophyceae) diversity in a Hawaiian reef environment. *J Phycol* **46**:53–65.
- Pochon X, Wecker P, Stat M, Berteaux-Lecellier V, Lecellier G. 2019. Towards an in-depth characterization of Symbiodiniaceae in tropical giant clams via metabarcoding of pooled multi-gene amplicons. *PeerJ* **7**:e6898.
- Porto I, Granados C, Restrepo JC, Sánchez JA. 2008. Macroalgal-associated dinoflagellates belonging to the genus *Symbiodinium* in Caribbean reefs. *PLoS One* **3**:e2160.
- Prada C, McIlroy SE, Beltrán DM, Valint DJ, Ford SA, Hellberg ME, Coffroth MA. 2014. Cryptic diversity hides host and habitat specialization in a gorgonian-algal symbiosis. *Mol Ecol* **23**:3330–3340.
- Putman AI, Carbone I. 2014. Challenges in analysis and interpretation of microsatellite data for population genetic studies. *Ecol Evol* **4**:4399–4428.
- Putnam HM, Davidson JM, Gates RD. 2016. Ocean acidification influences host DNA methylation and phenotypic plasticity in environmentally susceptible corals. *Evol Appl* **9**:1165–1178.
- Putnam HM, Gates RD. 2015. Preconditioning in the reef-building coral *Pocillopora damicornis* and the potential for trans-generational acclimatization in coral larvae under future climate change conditions. *J Exp Biol* **218**:2365–2372.
- Quattrini AM, Rodríguez E, Faircloth BC, Cowman PF, Brugler MR, Farfan GA, Hellberg ME, Kitahara MV, Morrison CL, Paz-García DA, Reimer JD, McFadden CS. 2020. Palaeoclimate ocean conditions shaped the evolution of corals and their skeletons through deep time. *Nat Ecol Evol* **4**:1531–1538.
- Quigley KM, Baker AC, Coffroth MA, Willis BL. 2018. Bleaching resistance and the role of algal endosymbionts. In: van Oppen, MJH, Lough JM, editors. *Coral Bleaching: Patterns, Processes, Causes, and Consequences*. Heidelberg: Springer Berlin. pp. 111–151.
- Quigley KM, Bay LK, Willis BL. 2017a. Temperature and water quality-related patterns in sediment-associated *Symbiodinium* communities impact symbiont uptake and fitness of juveniles in the genus *Acropora*. *Front Mar Sci* **4**:401.
- Quigley KM, Davies SW, Kenkel CD, Willis BL, Matz MV, Bay LK. 2014. Deep-sequencing method for quantifying background abundances of *Symbiodinium* types: exploring the rare *Symbiodinium* biosphere in reef-building corals. *PLoS One* **9**:e94297.
- Quigley KM, Willis BL, Bay LK. 2017b. Heritability of the *Symbiodinium* community in vertically- and horizontally-transmitting broadcast spawning corals. *Sci Rep* **7**:8219.
- Quigley KM, Willis BL, Bay LK. 2016. Maternal effects and *Symbiodinium* community composition drive differential patterns in juvenile survival in the coral *Acropora tenuis*. *R Soc Open Sci* **3**:160471.
- Ramírez-Castañeda V. 2020. Disadvantages in preparing and publishing scientific papers caused by the

- dominance of the English language in science: The case of Colombian researchers in biological sciences. *PLoS One* **15**:e0238372.
- Ramsby BD, Hill MS, Thornhill DJ, Steenhuizen SF, Achlatis M, Lewis AM, LaJeunesse TC. 2017. Sibling species of mutualistic *Symbiodinium* Clade G from bioeroding sponges in the western Pacific and western Atlantic oceans. *J Phycol* **53**:951–960.
- Ramsby BD, Shirur KP, Iglesias-Prieto R, Goulet TL. 2014. *Symbiodinium* photosynthesis in Caribbean octocorals. *PLoS One* **9**:e106419.
- Reich HG, Kitchen SA, Stankiewicz KH, Devlin-Durante M, Fogarty ND, Baums IB. 2021. Genomic variation of an endosymbiotic dinoflagellate (*Symbiodinium* “fitti”) among closely related coral hosts. *Mol Ecol* **30**:3500–3514.
- Reich HG, Rodriguez IB, LaJeunesse TC, Ho T-Y. 2020. Endosymbiotic dinoflagellates pump iron: Differences in iron and other trace metal needs among the Symbiodiniaceae. *Coral Reefs* **39**:915–927.
- Reimer JD, Herrera M, Gatins R, Roberts MB, Parkinson JE, Berumen ML. 2017. Latitudinal variation in the symbiotic dinoflagellate *Symbiodinium* of the common reef zoantharian *Palythoa tuberculosa* on the Saudi Arabian coast of the Red Sea. *J Biogeogr* **44**:661–673.
- Reimer JD, Shah MMR, Sinniger F, Yanagi K, Suda S. 2010. Preliminary analyses of cultured *Symbiodinium* isolated from sand in the oceanic Ogasawara Islands, Japan. *Mar Biodivers* **40**:237–247.
- Richards AL, Merrill AE, Coon JJ. 2015. Proteome sequencing goes deep. *Curr Opin Chem Biol* **24**:11–17.
- Roach TNF, Dilworth J, H CM, Jones AD, Quinn RA, Drury C. 2021. Metabolomic signatures of coral bleaching history. *Nat Ecol Evol* **5**:495–503.
- Robison JD, Warner ME. 2006. Differential impacts of photoacclimation and thermal stress on the photobiology of four different phylotypes of *Symbiodinium* (Pyrrhophyta). *J Phycol* **42**:568–579.
- Rodríguez-Casariago JA, Cunning R, Baker AC, Eirin-Lopez JM. 2021. Symbiont shuffling induces differential DNA methylation responses to thermal stress in the coral *Montastraea cavernosa*. *Mol Ecol* **31**:588–602.
- Rodríguez-Casariago JA, Ladd MC, Shantz AA, Lopes C, Cheema MS, Kim B, Roberts SB, Fourqurean JW, Ausio J, Burkepile DE, Eirin-Lopez JM. 2018. Coral epigenetic responses to nutrient stress: Histone H2A.X phosphorylation dynamics and DNA methylation in the staghorn coral *Acropora cervicornis*. *Ecol Evol* **48**:224.
- Rodríguez-Casariago JA, Mercado-Molina AE, Garcia-Souto D, Ortiz-Rivera IM, Lopes C, Baums IB, Sabat AM, Eirin-Lopez JM. 2020. Genome-wide DNA methylation analysis reveals a conserved epigenetic response to seasonal environmental variation in the staghorn coral *Acropora cervicornis*. *Front Mar Sci* **7**:822.
- Rodríguez-Lanetty M. 2003. Evolving lineages of *Symbiodinium*-like dinoflagellates based on ITS1 rDNA. *Mol Phylogenet Evol* **28**:152–168.
- Rodríguez L, López C, Casado-Amezua P, Ruiz-Ramos DV, Martínez B, Banaszak A, Tuya F, García-Fernández A, Hernández M. 2019. Genetic relationships of the hydrocoral *Millepora alcicornis* and its symbionts within and between locations across the Atlantic. *Coral Reefs* **38**:255–268.
- Rohart F, Gautier B, Singh A, Lê Cao K-A. 2017. *mixOmics*: An R package for 'omics feature selection and multiple data integration. *PLoS Comput Biol* **13**:e1005752.
- Rosbach S, Hume BCC, Cárdenas A, Perna G, Voolstra CR, Duarte CM. 2021. Flexibility in Red Sea *Tridacna maxima*-Symbiodiniaceae associations supports environmental niche adaptation. *Ecol Evol* **11**:3393–3406.
- Rowan R, Knowlton N. 1995. Intraspecific diversity and ecological zonation in coral-algal symbiosis. *Proc Natl Acad Sci USA* **92**:2850–2853.
- Rowan R, Knowlton N, Baker A, Jara J. 1997. Landscape ecology of algal symbionts creates variation in episodes of coral bleaching. *Nature* **388**:265–269.
- Rowan R, Powers DA. 1992. Ribosomal RNA sequences and the diversity of symbiotic dinoflagellates (zooxanthellae). *Proc Natl Acad Sci USA* **89**:3639–3643.
- Rowan R, Powers DA. 1991. A molecular genetic classification of zooxanthellae and the evolution of animal-

- algal symbioses. *Science* **251**:1348–1351.
- Rubin ET, Enochs IC, Foord C, Mayfield AB, Kolodziej G, Basden I, Manzello DP. 2021. Molecular mechanisms of coral persistence within highly urbanized locations in the Port of Miami, Florida. *Front Mar Sci* **8**:695236.
- Russnak V, Rodriguez-Lanetty M, Karsten U. 2021. Photophysiological tolerance and thermal plasticity of genetically different Symbiodiniaceae endosymbiont species of Cnidaria. *Front Mar Sci* **8**:657348.
- Saad OS, Lin X, Ng TY, Li L, Ang P, Lin S. 2020. Genome size, rDNA copy, and qPCR assays for Symbiodiniaceae. *Front Microbiol* **11**:847.
- Sampayo EM, Dove S, LaJeunesse TC. 2009. Cohesive molecular genetic data delineate species diversity in the dinoflagellate genus *Symbiodinium*. *Mol Ecol* **18**:500–519.
- Sampayo EM, Ridgway T, Bongaerts P, Hoegh-Guldberg O. 2008. Bleaching susceptibility and mortality of corals are determined by fine-scale differences in symbiont type. *Proc Natl Acad Sci USA* **105**:10444–10449.
- Santos SR, Coffroth MA. 2003. Molecular genetic evidence that dinoflagellates belonging to the genus *Symbiodinium freudenthal* are haploid. *Biol Bull* **204**:10–20.
- Santos SR, Gutierrez-Rodriguez C, Coffroth MA. 2003. Phylogenetic identification of symbiotic dinoflagellates via length heteroplasmy in domain V of chloroplast large subunit (*cp23S*)-ribosomal DNA sequences. *Mar Biotechnol* **5**:130–140.
- Santos SR, Gutierrez-Rodriguez C, Lasker HR, Coffroth MA. 2003. *Symbiodinium* sp. associations in the gorgonian *Pseudopterogorgia elisabethae* in the Bahamas: High levels of genetic variability and population structure in symbiotic dinoflagellates. *Mar Biol* **143**:111–120.
- Santos SR, Shearer TL, Hannes AR, Coffroth MA. 2004. Fine-scale diversity and specificity in the most prevalent lineage of symbiotic dinoflagellates (*Symbiodinium*, Dinophyceae) of the Caribbean. *Mol Ecol* **13**:459–469.
- Santos SR, Taylor DJ, Coffroth MA. 2001. Genetic comparisons of freshly isolated versus cultured symbiotic dinoflagellates: implications for extrapolating to the intact symbionts. *J Phycol* **37**:900–912.
- Schoenberg DA, Trench RK. 1980. Genetic variation in *Symbiodinium* (= *Gymnodinium*) *microadriaticum* Freudenthal, and specificity in its symbiosis with marine invertebrates. I. Isoenzyme and soluble protein patterns of axenic cultures of *Symbiodinium microadriaticum*. *Proc R Soc Lond* **207**:405–427.
- Schoenberg DA, Trench RK. 1976. Specificity of symbioses between marine cnidarians and zooxanthellae In: Mackie GO, editor. *Coelenterate Ecology and Behavior*. Boston, MA: Springer US. pp. 423–432.
- Sepey M, Manni M, Zdobnov EM. 2019. BUSCO: Assessing genome assembly and annotation completeness In: Kollmar M, editor. *Gene Prediction: Methods and Protocols*. New York, NY: Springer New York. pp. 227–245.
- Shah S, Chen Y, Bhattacharya D, Chan CX. 2020. Sex in Symbiodiniaceae dinoflagellates: genomic evidence for independent loss of the canonical synaptonemal complex. *Sci Rep* **10**:9792.
- Shoguchi E, Beedessee G, Tada I, Hisata K, Kawashima T, Takeuchi T, Arakaki N, Fujie M, Koyanagi R, Roy MC, Kawachi M, Hidaka M, Satoh N, Shinzato C. 2018. Two divergent *Symbiodinium* genomes reveal conservation of a gene cluster for sunscreen biosynthesis and recently lost genes. *BMC Genomics* **19**:458.
- Shoguchi E, Shinzato C, Kawashima T, Gyoja F, Mungpakdee S, Koyanagi R, Takeuchi T, Hisata K, Tanaka M, Fujiwara M, Hamada M, Seidi A, Fujie M, Usami T, Goto H, Yamasaki S, Arakaki N, Suzuki Y, Sugano S, Toyoda A, Kuroki Y, Fujiyama A, Medina M, Coffroth MA, Bhattacharya D, Satoh N. 2013. Draft assembly of the *Symbiodinium minutum* nuclear genome reveals dinoflagellate gene structure. *Curr Biol* **23**:1399–1408.
- Shoguchi E, Yoshioka Y, Shinzato C, Arimoto A, Bhattacharya D, Satoh N. 2020. Correlation between organelle genetic variation and RNA editing in dinoflagellates associated with the coral *Acropora digitifera*. *Genome Biol Evol* **12**:203–209.
- Silbiger NJ, Stubler AD. 2019. Unprofessional peer reviews disproportionately harm underrepresented groups in STEM. *PeerJ* **7**:e8247.

- Silverstein RN, Correa AMS, Baker AC. 2012. Specificity is rarely absolute in coral–algal symbiosis: Implications for coral response to climate change. *Proc Biol Sci* **279**:2609–2618.
- Silverstein RN, Correa AMS, LaJeunesse TC, Baker AC. 2011. Novel algal symbiont (*Symbiodinium* spp.) diversity in reef corals of Western Australia. *Mar Ecol Prog Ser* **422**:63–75.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**:3210–3212.
- Smith EG, D'Angelo C, Salih A, Wiedenmann J. 2013. Screening by coral green fluorescent protein (GFP)-like chromoproteins supports a role in photoprotection of zooxanthellae. *Coral Reefs* **32**:463–474.
- Smith EG, Gurskaya A, Hume BCC, Voolstra CR, Todd PA, Bauman AG, Burt JA. 2020. Low Symbiodiniaceae diversity in a turbid marginal reef environment. *Coral Reefs* **39**:545–553.
- Smith EG, Ketchum RN, Burt JA. 2017. Host specificity of *Symbiodinium* variants revealed by an ITS2 metahaplotype approach. *ISME J* **11**:1500–1503.
- Snyder GA, Browne WE, Traylor-Knowles N, Rosental B. 2020. Fluorescence-activated cell sorting for the isolation of scleractinian cell populations. *J Vis Exp* **159**:e60446.
- Sproles AE, Oakley CA, Matthews JL, Peng L, Owen JG, Grossman AR, Weis VM, Davy SK. 2019. Proteomics quantifies protein expression changes in a model cnidarian colonised by a thermally tolerant but suboptimal symbiont. *ISME J* **13**:2334–2345.
- Stat M, Baker AC, Bourne DG, Correa AMS, Forsman Z, Huggett MJ, Pochon X, Skillings D, Toonen RJ, van Oppen MJH, Gates RD. 2012. Molecular delineation of species in the coral holobiont. *Adv Mar Biol* **63**:1–65.
- Stat M, Carter D, Hoegh-Guldberg O. 2006. The evolutionary history of *Symbiodinium* and scleractinian hosts—Symbiosis, diversity, and the effect of climate change. *Perspect Plant Ecol Evol Syst* **8**:23–43.
- Stat M, Morris E, Gates RD. 2008. Functional diversity in coral–dinoflagellate symbiosis. *Proc Natl Acad Sci USA* **105**:9256–9261.
- Stefanoudis PV, Licuanan WY, Morrison TH, Talma S, Veitayaki J, Woodall LC. 2021. Turning the tide of parachute science. *Curr Biol* **31**:R184–R185.
- Stephens TG, Ragan MA, Bhattacharya D, Chan CX. 2018. Core genes in diverse dinoflagellate lineages include a wealth of conserved dark genes with unknown functions. *Sci Rep* **8**:17175.
- Suggett DJ, Goyen S, Evenhuis C, Szabó M, Pettay DT, Warner ME, Ralph PJ. 2015. Functional diversity of photobiological traits within the genus *Symbiodinium* appears to be governed by the interaction of cell size with cladal designation. *New Phytol* **208**:370–381.
- Suggett DJ, Moore CM, Hickman AE, Geider RJ. 2009. Interpretation of fast repetition rate (FRR) fluorescence: signatures of phytoplankton community structure versus physiological state. *Mar Ecol Prog Ser* **376**:1–19.
- Suggett DJ, van Oppen MJH. 2022. Horizon scan of rapidly advancing coral restoration approaches for 21st century reef management. *Emerg Top Life Sci* **6**:125–136.
- Suggett DJ, Warner ME, Leggat W. 2017. Symbiotic dinoflagellate functional diversity mediates coral survival under ecological crisis. *Trends Ecol Evol* **32**:735–745.
- Sunagawa S, Acinas SG, Bork P, Bowler C, Tara Oceans Coordinators, Eveillard D, Gorsky G, Guidi L, Iudicone D, Karsenti E, Lombard F, Ogata H, Pesant S, Sullivan MB, Wincker P, de Vargas C. 2020. Tara Oceans: Towards global ocean ecosystems biology. *Nat Rev Microbiol* **18**:428–445.
- Suyama Y, Matsuki Y. 2015. MIG-seq: An effective PCR-based method for genome-wide single-nucleotide polymorphism genotyping using the next-generation sequencing platform. *Sci Rep* **5**:16963.
- Suzuki MM, Bird A. 2008. DNA methylation landscapes: provocative insights from epigenomics. *Nat Rev Genet* **9**:465–476.
- Swain TD, Lax S, Backman V, Marcelino LA. 2020. Uncovering the role of Symbiodiniaceae assemblage composition and abundance in coral bleaching response by minimizing sampling and evolutionary biases. *BMC Microbiol* **20**:124.
- Taffe MA, Gilpin NW. 2021. Racial inequity in grant funding from the US National Institutes of Health. *eLife* **10**:e65697.

- Takabayashi M, Adams LM, Pochon X, Gates RD. 2012. Genetic diversity of free-living *Symbiodinium* in surface water and sediment of Hawai'i and Florida. *Coral Reefs* **31**:157–167.
- Takabayashi M, Santos SR, Cook CB. 2004. Mitochondrial DNA phylogeny of the symbiotic dinoflagellates (*Symbiodinium*, Dinophyta). *J Phycol* **40**:160–164.
- Takishita K, Ishikura M, Koike K, Maruyama T. 2003. Comparison of phylogenies based on nuclear-encoded SSU rDNA and plastid-encoded *psbA* in the symbiotic dinoflagellate genus *Symbiodinium*. *Phycologia* **42**:285–291.
- Taubert N, Bruns A, Lenke C, Stone G. 2021. Waiving article processing charges for least developed countries: a keystone of a large-scale open access transformation. *Insights Imaging* **34**:1–10.
- Tautz D. 1989. Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Res* **17**:6463–6471.
- Tchernov D, Gorbunov MY, de Vargas C, Narayan Yadav S, Milligan AJ, Häggblom M, Falkowski PG. 2004. Membrane lipids of symbiotic algae are diagnostic of sensitivity to thermal bleaching in corals. *Proc Natl Acad Sci USA* **101**:13531–13535.
- Tedersoo L, Tooming-Klunderud A, Anslan S. 2018. PacBio metabarcoding of Fungi and other eukaryotes: errors, biases and perspectives. *New Phytol* **217**:1370–1385.
- Terraneo TI, Fusi M, Hume BCC, Arrigoni R, Voolstra CR, Benzoni F, Forsman ZH, Berumen ML. 2019. Environmental latitudinal gradients and host-specificity shape Symbiodiniaceae distribution in Red Sea Porites corals. *J Biogeogr* **46**:2323–2335.
- Thornhill DJ, Howells EJ, Wham DC, Steury TD, Santos SR. 2017. Population genetics of reef coral endosymbionts (*Symbiodinium*, Dinophyceae). *Mol Ecol* **26**:2640–2659.
- Thornhill DJ, LaJeunesse TC, Kemp DW, Fitt WK, Schmidt GW. 2006. Multi-year, seasonal genotypic surveys of coral-algal symbioses reveal prevalent stability or post-bleaching reversion. *Mar Biol* **148**:711–722.
- Thornhill DJ, LaJeunesse TC, Santos SR. 2007. Measuring rDNA diversity in eukaryotic microbial systems: how intragenomic variation, pseudogenes, and PCR artifacts confound biodiversity estimates. *Mol Ecol* **16**:5326–5340.
- Thornhill DJ, Lewis AM, Wham DC, LaJeunesse TC. 2014a. Host-specialist lineages dominate the adaptive radiation of reef coral endosymbionts. *Evolution* **68**:352–367.
- Thornhill DJ, Xiang Y, Fitt WK, Santos SR. 2009. Reef endemism, host specificity and temporal stability in populations of symbiotic dinoflagellates from two ecologically dominant Caribbean corals. *PLoS One* **4**:e6262.
- Thurber RV, Payet JP, Thurber AR, Correa AMS. 2017. Virus-host interactions and their roles in coral reef health and disease. *Nat Rev Microbiol* **15**:205–216.
- Titlyanov EA, Kiyashko SI, Titlyanova TV, Yakovleva IM. 2009. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in tissues of reef building corals and the endolithic alga *Ostreobium quekettii* under their symbiotic and separate existence. *Galaxea* **11**:169–175.
- Tivey TR, Parkinson JE, Weis VM. 2020. Host and symbiont cell cycle coordination is mediated by symbiotic state, nutrition, and partner identity in a model cnidarian-dinoflagellate symbiosis. *mBio* **11**:e02626.
- Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L, Berumen ML, Bourne DG, Cantin N, Foret S, Matz M, Miller DJ, Moya A, Putnam HM, Ravasi T, van Oppen MJH, Thurber RV, Vidal-Dupiol J, Voolstra CR, Watson S-A, Whitelaw E, Willis BL, Munday PL. 2017. Rapid adaptive responses to climate change in corals. *Nat Clim Chang* **7**:627–636.
- Tortorelli G, Oakley CA, Davy SK, van Oppen MJH, McFadden GI. 2022. Cell wall proteomic analysis of the cnidarian photosymbionts *Breviolum minutum* and *Cladocopium goreau*. *J Eukaryot Microbiol* **69**:e12870.
- Traylor-Knowles N. 2021. Unlocking the single-cell mysteries of a reef-building coral. *Cell* **184**:2802–2804.
- Trench RK, Blank RJ. 1987. *Symbiodinium microadriaticum* Freudenthal, *S. goreau* sp. nov., *S. kawagutii* sp. nov. and *S. pilosum* sp. nov.: Gymnodinioid dinoflagellate symbionts of marine invertebrates. *J Phycol* **23**:469–481.

- Turland N, Kusber W-H, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Marhold K, McNeill J. 2018. International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code), Regnum vegetabile. Schmittgen, Germany: Koeltz Botanical Books.
- Turnham KE, Wham DC, Sampayo E, LaJeunesse TC. 2021. Mutualistic microalgae co-diversify with reef corals that acquire symbionts during egg development. *ISME J* **15**:3271–3285.
- Ulstrup KE, van Oppen MJH. 2003. Geographic and habitat partitioning of genetically distinct zooxanthellae (*Symbiodinium*) in *Acropora* corals on the Great Barrier Reef. *Mol Ecol* **12**:3477–3484.
- Umeki M, Yamashita H, Suzuki G, Sato T, Ohara S, Koike K. 2020. Fecal pellets of giant clams as a route for transporting Symbiodiniaceae to corals. *PLoS One* **15**:e0243087.
- Valenzuela-Toro AM, Viglino M. 2021. How Latin American researchers suffer in science. *Nature* **598**:374–375.
- van Oppen MJH, Bongaerts P, Frade P, Peplow LM, Boyd SE, Nim HT, Bay LK. 2018. Adaptation to reef habitats through selection on the coral animal and its associated microbiome. *Mol Ecol* **27**:2956–2971.
- van Oppen MJH, Leong J-A, Gates RD. 2009. Coral-virus interactions: A double-edged sword? *Symbiosis* **47**:1–8.
- van Oppen MJH, Mieog JC, Sánchez CA, Fabricius KE. 2005. Diversity of algal endosymbionts (zooxanthellae) in octocorals: the roles of geography and host relationships. *Mol Ecol* **14**:2403–2417.
- Vega de Luna F, Dang K-V, Cardol M, Roberty S, Cardol P. 2019. Photosynthetic capacity of the endosymbiotic dinoflagellate *Cladocopium* sp. is preserved during digestion of its jellyfish host *Mastigias papua* by the anemone *Entacmaea medusivora*. *FEMS Microbiol Ecol* **95**:fiz141.
- Venera-Ponton DE, Diaz-Pulido G, Rodriguez-Lanetty M, Hoegh-Guldberg O. 2010. Presence of *Symbiodinium* spp. in macroalgal microhabitats from the southern Great Barrier Reef. *Coral Reefs* **29**:1049–1060.
- Voolstra CR, Aranda M, Zhan Y, Dekker J. 2021a. *Symbiodinium microadriaticum* (coral microalgal endosymbiont). *Trends Genet* **37**:1044–1045.
- Voolstra CR, Buitrago-López C, Perna G, Cárdenas A, Hume BCC, Rädicker N, Barshis DJ. 2020. Standardized short-term acute heat stress assays resolve historical differences in coral thermotolerance across microhabitat reef sites. *Glob Chang Biol* **26**:4328–4343.
- Voolstra CR, Quigley KM, Davies SW, Parkinson JE, Peixoto RS, Aranda M, Baker AC, Barno AR, Barshis DJ, Benzoni F, Bonito V, Bourne DG, Buitrago-López C, Bridge TCL, Chan CX, Combosch DJ, Craggs J, Frommlet JC, Herrera S, Quattrini AM, Röthig T, Reimer JD, Rubio-Portillo E, Suggett DJ, Villeda H, Ziegler M, Sweet M. 2021b. Consensus guidelines for advancing coral holobiont genome and specimen voucher deposition. *Front Mar Sci* **8**:1029.
- Voolstra CR, Suggett DJ, Peixoto RS, Parkinson JE, Quigley KM, Silveira CB, Sweet M, Muller EM, Barshis DJ, Bourne DG, Aranda M. 2021c. Extending the natural adaptive capacity of coral holobionts. *Nat Rev Earth Environ* **2**:747–762.
- Voolstra CR, Valenzuela JJ, Turkarslan S, Cárdenas A, Hume BCC, Perna G, Buitrago-López C, Rowe K, Orellana MV, Baliga NS, Paranjape S, Banc-Prandi G, Bellworthy J, Fine M, Frias-Torres S, Barshis DJ. 2021d. Contrasting heat stress response patterns of coral holobionts across the Red Sea suggest distinct mechanisms of thermal tolerance. *Mol Ecol* **30**:4466–4480.
- Wangpraseurt D, Larkum AWD, Franklin J, Szabó M, Ralph PJ, Kühl M. 2014. Lateral light transfer ensures efficient resource distribution in symbiont-bearing corals. *J Exp Biol* **217**:489–498.
- Wangpraseurt D, Larkum AWD, Ralph PJ, Kühl M. 2012. Light gradients and optical microniches in coral tissues. *Front Microbiol* **3**:316.
- Wee HB, Kobayashi Y, Reimer JD. 2021. Effects of temperature, salinity, and depth on Symbiodiniaceae lineages hosted by *Palythoa tuberculosa* near a river mouth. *Mar Ecol Prog Ser* **667**:43–60.
- Wham DC, Carmichael M, LaJeunesse TC. 2014. Microsatellite loci for *Symbiodinium goreau* and other Clade C *Symbiodinium*. *Conserv Genet Resour* **6**:127–129.
- Wham DC, LaJeunesse TC. 2016. *Symbiodinium* population genetics: testing for species boundaries and analysing samples with mixed genotypes. *Mol Ecol* **25**:2699–2712.

- Wham DC, Ning G, LaJeunesse TC. 2017. *Symbiodinium glynnii* sp. nov., a species of stress-tolerant symbiotic dinoflagellates from pocilloporid and montiporid corals in the Pacific Ocean. *Phycologia* **56**:396–409.
- Wham DC, Pettay DT, LaJeunesse TC. 2011. Microsatellite loci for the host-generalist “zooxanthella” *Symbiodinium trenchi* and other Clade D *Symbiodinium*. *Conserv Genet Resour* **3**:541–544.
- Wirshing HH, Baker AC. 2016. On the difficulty of recognizing distinct *Symbiodinium* species in mixed communities of algal symbionts. *Mol Ecol* **25**:2724–2726.
- Wirshing HH, Feldheim KA, Baker AC. 2013. Vectored dispersal of *Symbiodinium* by larvae of a Caribbean gorgonian octocoral. *Mol Ecol* **22**:4413–4432.
- Witze ES, Old WM, Resing KA, Ahn NG. 2007. Mapping protein post-translational modifications with mass spectrometry. *Nat Methods* **4**:798–806.
- Xiang T. 2018. Isolation of axenic *Symbiodinium* cultures v1 (protocols.io.qxzdxp6). *protocols.io*. doi:10.17504/protocols.io.qxzdxp6
- Xiang T, Hambleton EA, DeNofrio JC, Pringle JR, Grossman AR. 2013. Isolation of clonal axenic strains of the symbiotic dinoflagellate *Symbiodinium* and their growth and host specificity. *J Phycol* **49**:447–458.
- Yamashita H, Koike K. 2016. Motility and cell division patterns among several strains of *Symbiodinium*. *Galaxea* **18**:13–19.
- Yamashita H, Koike K. 2013. Genetic identity of free-living *Symbiodinium* obtained over a broad latitudinal range in the Japanese coast. *Phycological Res* **61**:68–80.
- Yang F, Li L, Lin S. 2020. Methylation pattern and expression dynamics of methylase and photosystem genes under varied light intensities in *Fugacium kawagutii* (Symbiodiniaceae). *J Phycol* **56**:1738–1747.
- Yorifuji M, Yamashita H, Suzuki G, Kawasaki T, Tsukamoto T, Okada W, Tamura K, Nakamura R, Inoue M, Yamazaki M, Harii S. 2021. Unique environmental Symbiodiniaceae diversity at an isolated island in the northwestern Pacific. *Mol Phylogenet Evol* **161**:107158.
- Zaneveld JR, McMinds R, Vega Thurber R. 2017. Stress and stability: Applying the Anna Karenina principle to animal microbiomes. *Nat Microbiol* **2**:17121.
- Zhang C, Lin S. 2019. Initial evidence of functional siRNA machinery in dinoflagellates. *Harmful Algae* **81**:53–58.
- Ziegler M, Eguíluz VM, Duarte CM, Voolstra CR. 2018. Rare symbionts may contribute to the resilience of coral-algal assemblages. *ISME J* **12**:161–172.
- Ziegler M, Roder C, Büchel C, Voolstra CR. 2015. Niche acclimatization in Red Sea corals is dependent on flexibility of host-symbiont association. *Mar Ecol Prog Ser* **533**:149–161.
- Ziegler M, Seneca FO, Yum LK, Palumbi SR, Voolstra CR. 2017. Bacterial community dynamics are linked to patterns of coral heat tolerance. *Nat Commun* **8**:14213.