

Review

Promising Strategies for Preserving Adult Endothelium Health and Reversing Its Dysfunction: From Liquid Biopsy to New Omics Technologies and Noninvasive Circulating Biomarker

Carmela Rita Balistreri

Cellular and Molecular Laboratory, Department of Biomedicine, Neuroscience and Advanced Diagnostics (Bi.N.D.), University of Palermo, 90134, Palermo, Italy

Correspondence: Prof CR Balistreri, Cellular and Molecular Laboratory, Department of Biomedicine, Neuroscience and Advanced Diagnostics (Bi.N.D.), University of Palermo, 90134, Palermo, Italy,
Email: carmelarita.balistreri@unipa.it

Abstract: The endothelium has multiple functions, ranging from maintaining vascular homeostasis and providing nutrition and oxygen to tissues to evoking inflammation under adverse conditions and determining endothelial barrier disruption, resulting in dysfunction. Endothelial dysfunction represents the typical condition associated with the pathogenesis of all diseases of the cardiovascular system, as well as of diseases of all the other human body's systems, including sepsis, acute respiratory distress syndrome, and COVID-19 respiratory distress. Such evidence is leading to the identification of potential biomarkers and therapeutic targets for preserving, reverting, or restoring endothelium integrity and functionality by promptly treating its dysfunction. Here, some strategies for achieving these goals are explored, despite the diverse challenges that exist, necessitating significant bench work associated with an increased number of clinical studies.

Keywords: Endothelium; Endothelium dysfunction; Newer omics technologies; network medicine; biomarkers and therapeutic targets.

1. Introduction

The term “endothelium” indicates the cellular layer that acts as barrier between blood flow and the inner lining of the vessel wall. Considered, for numerous years, as an inert vascular component, the endothelium is recognized today as the hub of the homeostasis of the entire cardiovascular system, including the heart valves, where endothelial cells (EC), defined as valvular ECs (VECs), can control the proliferation, differentiation, apoptosis, and remodeling of valvular interstitial cells (VICs) [1]. Thus, the endothelium represents a vital organ of the cardiovascular system and the principal element of the stroma of all body tissues, executing diverse regulatory activities thanks to the production and release of several endothelial-derived vasoactive factors, acting via endocrine, paracrine, or autocrine signaling. Accordingly, endothelial control includes not only the regulation of vascular tone, but also angiogenesis, inflammation, platelet aggregation, leukocyte adhesion, vascular cell proliferation, and oxidative stress [2-6] (see **Figure 1**). Such essential functions of the endothelium are already marked during fetal life. In the fetus, the endothelium has the fundamental role of ensuring the correct ontogenesis of immune cells (i.e., tissue-resident macrophages, microglia cells, and epidermal $\gamma\delta$ T cells) and cells of the hematopoietic system, as well as the development of other systems, such as the nervous and cardiovascular systems [6-8].




Figure 1. Architecture, EC pathways, and their heterogeneous presence and complexity in various body tissues. (a). Endothelium architecture and expression of some crucial pathways involved in its homeostasis, as well as in its activation, determining changes in the expression of specific genes and in the consequent phenotypic EC changes related to evocation of dysfunction, caused by inflammation and endothelial barrier disruption (increased vascular permeability, edema formation, release of proinflammatory cytokines, and leukocyte extravasation). (b) Endothelium as an essential element of not only of the entire cardiovascular system, but also as the “heart” of all the tissues of the body, consequently contributing to their physiological homeostasis and function.

Consistent with these observations, the current evidence reports a close relationship between biochemical or structural injury, altered genesis, homeostasis, and function of the endothelium, and the onset and progression of pathological conditions correlated to inflammation, thrombosis, and atherosclerosis, as well as alterations in the vascular resistance and blood pressure [2,4,9-11]. Consequently, endothelial dysfunction is assumed as the axis of activation of all pathophysiological mechanisms and related disease pathways linked to excessive morbidity and mortality, such as hypertension, myocardial infarction, aneurysm, stroke, diabetes, metabolic syndrome, rheumatic diseases, renal failure, neurodegenerative diseases, cancer, and inflammatory diseases [2,4,5,8, 9, 11-20] (see **Figure 2**). For instance, tumor neovascularization is characterized by a highly disordered vascular endothelium network and the generation of new blood vessels. In addition, the abovementioned pathologies are characterized to coexist in patients with a median age of about 65 years old and over, by increasing the risk of adverse events [21,22]. Furthermore, a significant association between typical EC plasma proteins and individual risk of cardiovascular diseases (CVD), detected using the Framingham risk score, has also been evidenced in recent studies confirming the close relationship between CVD onset and endothelium dysfunction, as well as corresponding biomarker profiles [23]. Interestingly, alterations in the vascular endothelium have also been associated with organ dysfunction in patients affected by severe SARS-CoV-2 infection, belonging to the 70–89 age group [24-27] (see **Figure 2**).




Figure 2. Endothelial dysfunction and its key involvement in many human diseases.

Considering these observations, it is of crucial importance to develop strategies, with the aim of not only warding off aggressive agents or risk factors for preventing endothelium dysfunction, but also identifying therapies which can reverse or mitigate the damaging effects generated by such conditions. Certainly, regenerative strategies via induced pluripotent stem cells (IPSCs) and direct reprogramming targeting ECs via transcription factors (TFs) involved in endothelial cell fate are optimal examples. TFs, such as OCT4, SOX2, KLF4, c-MYC, ETV2, and ERG, have recently become the subject of many studies, and the direct reprogramming strategy appears to be promising for the treatment of diseases such as CVD. However, limitations in their efficiency linked principally to epigenetic resistance and the methods used have been evidenced [28] (see **Figure 3**). Thus, the clinical applicability and translational nature of such strategies do not appear be evident. Here, attention is focused on new promising methodologies and biological matrices for use in achieving the suggested objectives. Accordingly, the use of blood liquid biopsy and omics technologies is described in the next section, in addition to their application in studies on endothelial dysfunction aimed at identifying biomarkers and emerging therapeutic targets.

Given the complexity of the mechanisms and related pathways related to the responses of the endothelium to physical or biochemical stimuli in carrying out its roles, diverse biomarkers and therapeutic targets are described and discussed with respect to their likely prevention and treatment of endothelium dysfunction and its pathological complications, while advantages and limitations are also reported.




Figure 3. Some strategies for obtaining mature EC.

2. Omics Technologies and Liquid Biopsy as Emerging Tools for Identifying Endothelial Biomarkers and Therapeutic Targets

In proposing new strategies for preserving the functionality and efficiency of adult endothelium tissue, it is imperative to identify molecules as optimal biomarkers and therapeutic targets. Biomarkers represent crucial indicators of both physiological and pathological processes, such as endothelium dysfunction. Specific changes in molecular and cellular mechanisms of physiological processes result in biochemical alterations at both the tissue and the systemic level, which can give us comprehensive information regarding the healthy status of tissues (in our case, the endothelium), as well as the nature of any pathological tissue dysfunction and disease. In addition, any disease biomarker should be specific and reliable, able to distinguish between the physiological condition of a tissue, organ, or system and a dysfunctional condition or disease, and diverse across various diseases or their subgroups and phenotypes. Accordingly, biomarkers can predict the risk of disease, facilitate early diagnosis, and enable setting guidelines for the development of new therapies for treating diseases. New-generation omics technologies and their platforms, including next-generation sequencing (NGS), epigenomics, proteomics, metabolomics, and microbiomics [29,30] (see **Figure 4**), can detect molecules involved in the close relationship between genotype and the environment, nutritional habits, and endothelial health or dysfunction, at an unprecedented grade of importance and resolution [31-33]. They also allow obtaining an enormous quantity of data, albeit accompanied by consequent computational difficulties in their interpretation [31-33]. Today, such a challenge has been resolved thanks to potent bioinformatics algorithms, which can easily interpret and combine big data, by offering the possibility of both analyzing them at diverse multilevel networks and identifying genes, proteins, metabolites, and microbiomes, as well as their interactions in the complex human interactome, which can be used as emerging biomarkers and drug targets [31-33] (see **Figure 4**). Such a strategy exemplifies network medicine with the feature of combining molecular and bioinformatic data, which can improve current medical practices in the management of endothelial dysfunction-related pathologies by overcoming the limits of the obsolete points of view regarding endothelial dysfunction, considered the direct consequence of a single molecular alteration. An example of such an approach is machine learning (see **Figure 4**). The Abasht group [34] used this approach for evaluating vascular

endothelial dysfunction in wooden breast syndrome, while the Treder group [35] used this approach for detecting graft detachment in Descemet membrane endothelial keratoplasty. Furthermore, it is currently being applied to develop algorithms for vascular endothelial senescence, as recently evidenced by Sun and Feinberg [36].




Figure 4. Network medicine and its features in endothelium studies

Furthermore, omics technologies enable evaluating changes at a single-cell level, i.e., the evaluation of gene expression profiles of the endothelium at the transcriptome and epigenome level, when exposed to one or more stressors, such as disturbed blood flow, high-glucose status, or inflammation. Precisely, it is possible to evaluate the transcriptomic and chromatin accessibility variations at the resolution of a single-EC. This allows comparing the effects of stressors in healthy ECs vs. stressed ECs in the same vessel or in diverse vessels, thereby following the changes in physiological status of the endothelium as a result of dysfunction or other conditions, e.g., hyperplasia (see below). A similar study design was recently adopted by Kumar and coworkers [37] for evaluating the gene expression profile in ECs isolated, via an innovative protocol, from the lumen of carotid arteries from mice subjected to a partial carotid ligation (PCL), which induces disturbed blood flow (d-flow) in the left carotid artery, while maintaining stable laminar flow in the right carotid artery, used as a control. Such a mouse PCL model allows comparing the effects induced in ECs by blood flow under disturbed conditions or with stable laminar flow (s-flow). ECs were obtained from the carotid arteries by dissection 2 days or 2 weeks post PCL surgery, before subjecting their lumen to collagenase digestion to acquire endothelial-enriched single cells or single nuclei. These single-cell and single-nuclei preparations were utilized for RNA preparation, library generation, single-cell RNA sequencing (scRNA-seq), and single-cell transposase-accessible chromatin sequencing (scATACseq), which were consented to determine the transcriptomic and chromatin accessibility changes. The data obtained consequently allowed evaluating the genes expressed under the effects induced by d-flow or s-flow at a single-cell level, resulting in the activation of proatherogenic or physiological responses. These proatherogenic genes and their molecules were then suggested as targets for developing therapies. Precisely, it was confirmed that Krüppel-like factors 4 and 2 (KLF4/KLF2) act as s-flow-sensitive transcription factor (TF)-binding sites. The identified TFs sensitive to d-flow were RELA, AP1, STAT1, and TEAD1, which were suggested to induced d-flow EC reprogramming from an athero-protective to proatherogenic phenotypes, including endothelial-mesenchymal transition [37,38]. Furthermore, thanks to the integration of diverse omics assays, including acetylation of histone 3 lysine 27 (H3K27ac) ChIP-seq (chromatin immunoprecipitation followed by high-throughput sequencing), ATAC-seq (an assay for transposase-accessible chromatin sequencing), and RNA-seq

(RNA-sequencing), it was possible to detect the genome-wide epigenetic regulations in ECs in response to athero-protective pulsatile shear stress (PS). [39]. The combined data obtained showed that inositol 1,4,5-trisphosphate receptor 3 (ITPR3) was upregulated under PS conditions via KLF4 or statins, in ECs isolated from mouse aorta, lung ECs isolated from EC-KLF4-transgenic versus EC-KLF4-knockout mice, and atorvastatin-treated ECs. The use of KLF4 ATAC-qPCR (quantitative polymerase chain reaction) and ChIP-qPCR also led to the identification of a specific locus in the promoter region of the ITPR3 gene indispensable for KLF4 binding, H3K27ac enrichment, chromatin accessibility, RNA polymerase II recruitment, and ITPR3 transcriptional activation. [39]. By evocating the deletion of KLF4 binding locus in EC thanks to the use of a clustered regularly interspaced short palindromic repeats-associated protein 9 (CRISPR/Cas9) system, a blunted calcium influx, reduced expression of endothelial nitric oxide synthase, and diminished nitric oxide bioavailability were evidenced. In the complex, such interesting results obtained thanks to novel multi-omics investigations suggested that KLF4 is crucial for PS-modulated H3K27ac, which allows the transcriptional activation of ITPR3. Such a mechanism contributes to Ca²⁺-dependent eNOS (endothelial nitric oxide synthase) activation and EC homeostasis, and these molecules are potential candidates for biomarkers and therapeutic targets [39].

Another group integrated investigations of in vivo epigenomic mapping with conditional knockout, gene transfer, and pharmacology with the aim of studying, in rodent models, the endothelial hyperplasia/proliferation (IH), which represents the primary etiology of vascular stenosis. The data from injured (IH-prone) rat arteries evidenced a rise in genome-wide occupancy by histone 3 lysine 27 trimethylation (H3K27me3), a gene-repression indicator. This result was unexpected when considering the traditional point of view on prevailing post-injury gene activation rather than repression. Additional evaluation has also demonstrated a shift in H3K27me3 enrichment to anti-proliferative genes from pro-proliferative genes, characterized by gene activation via H3K27ac (acetylation) accumulation. H3K27ac and its reader BRD4 (bromodomain protein) showed co-enrichment of *Ezh2*, while conditional BRD4 knockout in injured mouse arteries resulted in a decrease in H3K27me3 and its writer EZH2, which positively regulates another pro-IH chromatin modulator UHRF1. Thus, these results reveal an injury-induced locus-specific H3K27me3 redistribution in the epigenomic landscape involving BRD4→EZH2→UHRF1 hierarchical regulation, while suggesting that these players may be pharmaceutical targets [40].

Another interesting study was conducted in 2021 by a group in Napoli examining the glucose-dependent and dose-responsive alterations in endothelial DNA methylation to detect a putative epigenetic mechanism underlying diabetic vasculopathy. Specifically, they discovered the disproportionate glucose-dependent methylation and gene expression of VEGF and NO signaling cascades, constituting a physiologic imbalance known to evoke endothelial dysfunction in diabetes. Consequently, they assumed that epigenetic mechanisms encode glycemic memory within EC, and the pathways involved can be used as therapeutic targets [41]. Specifically, their novel evidence demonstrated that hyperglycemia triggers dose-responsive variations in DNA methylation dynamics, affecting key physiological processes involved in the maintenance of endothelial function, including a glucose-dependent physiologic uncoupling of VEGF and NO signaling, which causes endothelial dysfunction.

Additionally, recent advances in molecular genetics and omics technologies also showed a significant molecular heterogeneity within brain endothelial and perivascular cell types [42]. The blend of these conventional and modern approaches has enabled detecting phenotypical variations between healthy and abnormal conditions at the single-cell level. Accordingly, a deep understanding of brain vascular cell states during physiological, pathological, and aging processes has rapidly emerged, and therapeutic approaches have been developed, including vascularization models of human brain organoids on a chip [widely quoted in 42], self-assembling multicellular blood-brain barrier (BBB) spheroids [widely quoted in 42], and improved endothelial BBB differentiation

protocols using human pluripotent or induced pluripotent stem cells [widely quoted in 8 and 42]. Human organoid and cell reprogramming technologies have rapidly improved, allowing the simulation of human brain development and disease modeling with the construction of a functional vasculature. These in vitro models will become potent instruments for exploring human-specific vascular traits absent in animal models, for investigating drug delivery across the BBB, and for screening drugs targeting BBB dysfunction in neurological diseases. Consequently, this research recently led our group, along with others, to hypothesize a close relationship between endothelial BBB conditions and neurodegenerative diseases [widely quoted in 8 and 42].

Therefore, the literature is seeing an increase in the number of multi-omics investigations on endothelium, and the studies reported above constitute only some examples. However, many of these studies were complex with regard to their design, often invasive or necessitating the sacrifice of animals, along with the intricate approach to collecting the biological study samples and the related costs. Alternatively, studies on other types of samples, whose collection attracts a minor cost and a simple technique, are growing in number. Indeed, studies aimed at identifying noninvasive indicators of endothelial tissue status, simply using liquid biopsy, are increasing. Liquid biopsy is attractive as it facilitates the collection of an optimal biological sample, thereby enabling the extrapolation of big data via multi-omics technologies [43-45]. Liquid biopsy consists of an easy sampling of diverse bodily fluids, principally including blood, saliva, and urine, for the evaluation of various potential circulating noninvasive biomarkers [43-45]. As a biopsy, it also enables obtaining circulating intact cells, cell-free nucleic acids, circulating epigenetic-sensitive molecules (methylated DNA, modified histones, and noncoding RNAs), circulating metabolites, and other cell products, such as micro-vesicles and exosomes [43-45] (see **Figure 5**). However, unlike tissue biopsy, liquid biopsy offers the advantage of collecting heterogeneous cellular phenotypes by giving information on many circulating cell types and their products at specific times, thereby allowing a real-time monitoring of the evolution of a pathological condition or of the switch of the endothelium from physiological to dysfunctional [46]. In addition, emerging NGS platforms executed on body fluid samples, principally blood, can detect novel circulating noninvasive biomarkers, which may further the development of precision medicine and personalized therapy in the field of endothelium dysfunction-related pathologies, by preserving the health of the endothelium or reverting its alterations [43-46]. Furthermore, liquid biopsy shows several benefits compared to tissue biopsy; it is (a) noninvasive with negligible pain and risk, (b) more quickly executed, (c) easier to collect, (d) able to provide a real-time systemic profile, (e) able to give spatiotemporal information, and (f) able to monitor physiological or pathological conditions over time.




Figure 5. Liquid biopsy, describing some components and laboratory analysis techniques. Liquid biopsy involves the collection and analysis of diverse components from peripheral blood samples, including cell-free nucleic acids (cfDNA/ctDNA and cfRNA) and exosomes. cfDNA: circulating free DNA, ctDNA: circulating tumor DNA, cfRNA: cell-free RNA, CTCs: circulating tumor cells, TEPs: tumor-educated platelets, NGS: next-generation sequencing, qPCR: quantitative polymerase chain reaction.

As mentioned above, fluid-based assays may provide noninvasive indicators that indicate the endothelium status or allow sequential monitoring of the evolution of dysfunction. Interesting evidence can be gained by assessing the number, function, and senescence grade of endothelial progenitor cells (EPCs), as well as by evaluating the effects induced by stem-cell therapies (i.e., human umbilical cord-derived mesenchymal stem cells (MSCs)) or by detecting the quantity and quality of circulating metabolites and other cell products, such as micro-vesicles and exosomes or molecules able to modulate the gene expression at transcriptional, post-transcriptional, and translational levels, including circular RNAs (circRNAs) and ADAR enzymes. The next section reports the description of some of these metabolites and cell products, stressing their emerging potential as biomarkers and therapeutic targets of endothelium dysfunction, as well as their related limitations.

3. Endothelial Progenitor Cells (EPCs) as Potential Biomarkers of Endothelium Dysfunction and Therapeutic Agents

Circulating EPCs represent the progenitors of ECs, and they originate from bone marrow (BM)-derived hematopoietic stem cells (HSCs) [47-49]. EPCs constitute a real resource of ECs, thus maintaining vascular and tissue homeostasis and appropriate oxygen regulation and transport. Accordingly, EPCs represent a reservoir of circulating cells, able to target injury sites, restore endothelium integrity, and enable physiological activities [47-49]. The impact of EPCs in the vascularization process has been proven in both animal models and humans. Such evidence has also led to the assumption that reductions in EPC circulating number and/or alterations in their functions related to different causes could impact endothelium function and architecture, as well as the onset and complications of endothelium dysfunction and, consequently, the survival of affected persons. Increased or decreased circulating EPC levels, as well as alterations in their function (for their detailed description it invites to consult our book <https://link.springer.com/book/10.1007/978-3-319-55107-4>; references 48), have indeed been associated with vascular endothelium aging and diverse endothelium dysfunction pathologies, including coronary artery disease, stroke, diabetes, systemic sclerosis, autoimmune disorders, and aneurysms [50-58]. Our group, for instance, recently evidenced that subjects affected by bicuspid aorta valve syndrome show a significant decrease in

both tissue and circulating levels of the Notch pathway, as well as a decrease in blood EPC number, compared to subjects with a tricuspid physiological valve, whether in the presence or absence of aorta aneurysm (AAA) [51,59]. In addition, we also evidenced, using both blood and human tissue aorta samples, that the unique independent risk predictors for vascular aging are age and a reduced EPC number, as well as reduced EPC migratory activity and senescence, mediated by high SA- β -Gal activity and high levels of TP53, p21, p16, and inflammatory genes. [59] Furthermore, we reported in another study [60] that the decreased circulating number of EPCs, associated with increased RDW values, augmented blood levels of high-sensitivity C-reactive protein, and reduced mean values of both leukocyte telomere length and telomerase activity, can predict both vascular aging and ascending aorta aneurysm (AAA) onset and prognosis. Consequently, we affirm that these players might be used as an ideal biomarker profile for vascular aging, as well as for the diagnosis and outcome of sporadic AAA.

Such evidence might be of clinical relevance, and possible new recommendations and preventive measures might be applied. Accordingly, Hill and colleagues [61] already showed that the number of circulating EPCs represents a better predictor of vascular reactivity than conventional cardiovascular risk factors. Furthermore, a significant correlation between *in vitro* EPC senescence and endothelium dysfunction pathologies risk was also detected in blood donors [59]. Consequently, EPCs might be considered as an optimal predictive biomarker, as well as a diagnostic and prognostic biomarker of endothelium dysfunction and related pathologies. Certainly, ulterior studies are needed. However, such evidence has also led to the use of EPCs as therapeutic agents in the autologous or heterologous treatment of several endothelium dysfunction diseases, as well as in clinical trials, despite contradictory results being obtained. In particular, a favorable improvement in left-ventricular (LV) function in a rat model of myocardial infarction (MI) after intravenous injection of *ex vivo* expanded human CD34⁺ cells was reported [47-49]. Another study examined the effect of catheter-based intramyocardial transplantation in a swine model of MI, providing encouraging outcomes in favoring the application of EPCs as a potential cell therapy in clinical trials [47-49]. Naruse and colleagues conducted a study to assess the therapeutic treatment of diabetic neuropathy using *in vivo* expanded human EPC and streptozotocin-induced diabetic nude rats [62]. They observed numerous micro-vessels at the site of EPC injection [63]. Another group evidenced an improvement of neurological functions in chronic cerebral ischemic rats injected with CD34⁺ HSC cells, including EPCs [63]. The ability of EPCs to expand in culture under *in vitro* conditions represents another barrier for their therapeutic use. Genetically modified and *ex vivo* expanded EPCs may become promising agents able to appropriately rescue the impaired neovascularization process under disease conditions. In a rhesus model, *ex vivo* CD34⁺ cell transfection with recombinant non-replicative herpes virus vector and subsequent cell transplantation resulted in the expression of vector genes in angiogenic areas of skin autografts of rhesus macaques. Since CD34⁺ cells possess a natural angiogenic tropism to the injured endothelium, they may serve as ideal candidates for the delivery of genes into areas of angiogenesis [64]. These results encouraged the execution of clinical trials for evaluating the potential of EPCs to enhance endothelial integrity and vascularization at ischemia sites in patients with CVDs. Three different strategies were mainly applied: (a) the *administration of granulocyte-colony stimulating factor (G-CSF)* for verifying the recruitment of the patient's own bone marrow resident progenitors, with two preliminary studies demonstrating increased LV function [65]; b) *the intracoronary infusion of BM progenitor cells in patients with MI*, which demonstrated positive effects on LV function in three smaller studies [47-49]. Subsequently, two prospective large trials assessed significant LV function after 4–6 months of administration of BM progenitor cells. Ten recent and large trials confirmed the success and safety of this procedure with a follow-up over 1.5 years [widely quoted in 47-49]. In addition, intramyocardial and intracoronary administration has recently been suggested as a suitable strategy for the treatment of patients with refractory angina [widely quoted in 47-49].]; (c) a more invasive strategy involving *the direct injection of cells into target tissues* [59]. This treatment

(precisely, *transepical or transendocardial injection of unfractionated BM cells*) has been performed in patients with diffuse coronary artery disease and intractable angina with no option of recanalization. Ventricular function and physical capacity were observed to increase, but the small sample size of these studies necessitates confirmation in larger studies [widely quoted in 47-49].

Studies using *autologous cell therapy* are also interesting. Accordingly, Yamamoto's group executed an intramuscular injection of autologous BM-derived mononuclear cells containing 1% CD34⁺ cells in patients with chronic limb ischemia [66]. They quantitatively assessed the expression of EPCs and endothelial markers (i.e., CD133 and VE-cadherin) before the experiment and after the injection. Before investigation, the transcription of these molecules was undetectable. Autologous injection determined an elevation of EPC marker transcription. Thus, they established that autologous BM cells may be used in the therapy of patients with arterial diseases. A replication of these results was obtained by Lenk and colleagues [67]. Erbs and colleagues also used this autologous treatment in patients who underwent recanalization of chronic coronary total occlusion [68]. Autologous treatment with EPCs, expanded by 4 days in endothelium growth medium, improves coronary endothelium function and wall motion abnormalities, and it has a beneficial effect on metabolism in the target area in patients with symptomatic coronary atherosclerosis.

Despite these promising data, the clinical application of EPCs in exogenous or autologous cell therapy remains unclear for different reasons. Accordingly, the validity of their results is influenced by different factors: (a) the insignificant number of patients in these studies; (b) their missing randomization; (c) participation of a limited number of centers; (d) the imprecise phenotypic features of EPC utilized in the treatments, (e) the different methods of administration applied, and (f) the security and viability of the treatments [47-49]. Teratoma formation, immunoreactivity, or arrhythmias constitute the major unfavorable effects, which have been detected after such treatments. In addition, other limitations have not been considered, including the limited number of EPCs in circulation, which usually requires their cultured expansion in a sufficient number of subpopulations from peripheral blood. The *in vitro* quantification of progenitor cells to obtain a quantity sufficient to utilize in therapeutic treatment can be modulated by phenotypic changes during their differentiation *in vitro* with the risk of obtaining senescent cells, requiring artificial cell pre-activation or stimulation for this approach. In addition to such evidenced limitations, another aspect influencing the results of such study is the lack of standardized criteria and a consensus for defining, characterizing, and identifying EPCs with well-established surface biomarkers, protocols, and methodologies. Consequently, other investigations are imperative [47-49].

3.1. Other Candidates for Therapeutic Agents of Endothelium Dysfunction

Promising evidence has recently been reported on the great potential of human umbilical cord-derived MSCs (hucMSCs) in repairing diabetic vascular endothelial damage, by applying assays of resazurin staining, MTT cell viability, wound healing, transwell migration, and Matrigel tube formation on human umbilical vein endothelial cells (HUVECs), and by evaluating how hucMSCs work through the assessment of RNA sequencing (RNA-seq) and molecular experiments [69]. Precisely, the use of a conditioned medium of hucMSCs (MSC-CM) revealed that hucMSCs improved the cell viability, wound healing, migration, and angiogenesis of HUVECs damaged by high glucose via paracrine signaling, and the altered gene expressions of IL-6, TNF- α , ICAM-1, VCAM-1, BAX, P16, P53, and ET-1 were significantly restored by MSC-CM. RNA-seq incorporated with real-time PCR and Western blot clarified that high glucose activates MAPK/ERK signaling in HUVECs, while MSC-CM reverses the abnormal phosphorylation of ERK and overexpression of MKNK2, ERBB3, MYC, and DUSP5 in the MAPK/ERK signaling pathway [69]. Certainly, other studies are needed.

4. CircRNAs and Their Potential as Biomarkers and Therapeutic Targets

CircRNAs are an endogenous group of noncoding RNA molecules [70-72], originating from pre-mRNAs and produced by back splicing [72]. They were first observed in plant viroids [73] and eukaryotic cells [74], and eventually discovered in all almost all organisms across the eukaryotic tree of life [75]. Their expression appears to be tissue [76], and cell-type specific and correlated to the specific stage of development or physiological and pathological conditions [75]. Accordingly, growing recent evidence sustains diverse regulator roles and versatile cell actions of such molecules [71, 75]. For example, circRNAs can act as miRNA decoys [77], RNA binding protein (RBP) sponges [78] and protein scaffolds [79], and they can also be translated into proteins [80,81]. In addition, it was also evidenced that circRNAs play roles in a large range of biological processes, including cell proliferation, apoptosis, and senescence, showing an abnormal expression in diverse human diseases, such as cancers [82, 83], neurodegenerative diseases, cardiovascular diseases [84] and immune diseases [85]. Such evidence has led to the consideration of circRNAs as promising biomarkers and therapeutic targets for human diseases [86], due to their stability, elevated levels, and tissue-cell-specificity [86]. Moreover, they may be easily detected in body fluids, such as urine, saliva, cerebrospinal fluid, or blood. Precisely, they are collected using liquid biopsy, which as mentioned above represent a revolutionary device in the management of diseases, facilitating diagnosis, prognosis, and treatments [44,45]. Specifically, peripheral blood represents the major fluid used in liquid biopsy investigations, where circRNAs can be cell-free or in blood cells [45]. Cell-free circRNAs can be observed in the plasma, serum, or blood endogenous vesicles, resulting from their secretion released into blood from diverse tissues. In addition, abovementioned, they are characterized by a large stability and abundance, facilitating their status as biomarkers of physiological processes or diseases.

Furthermore, recent studies have suggested the role of circRNAs in preventing or protecting the endothelium from further injury via senescence, remodeling, and progression in atherosclerotic tissue. Below, we report important data regarding this aspect.

4.1. *CircRNAs and the Endothelium: Recent Literature Evidence*

A growing number of studies have demonstrating the close association between the expression of specific profiles of circRNAs and various physiological and pathological endothelium conditions. Here, we report some recent studies. For example, a study by Wu and coworkers [87] demonstrated that circGNAQ shows a significantly low expression in senescent endothelial cells. Accordingly, circGNAQ silencing was revealed to evocate cell senescence, confirmed by the increase in senescence-associated β -galactosidase activity, decrease in cell proliferation, and inhibition of angiogenesis. Mechanistic investigations demonstrated that circGNAQ works in the endothelium as an endogenous miR-146a-5p sponge, enhancing the expression of its target gene *PLK2* by decoying miR-146a-5p and, consequently, delaying the endothelial cell senescence. Through in vivo studies, Wu 's group also revealed a significant association between circGNAQ overexpression and the inhibition of both endothelium senescence and potential progression in atherosclerosis [87]. Consequently, such data suggest that circGNAQ has important functions in endothelial cells, particularly in controlling senescence and the pathogenesis of atherosclerosis [87]. This also led to evidence that the management of circGNAQ might represent a potential therapeutic approach for limiting senescence of the endothelium and progression of atherosclerosis. In agreement with these data, it was also observed that circANRIL, derived from antisense noncoding RNA in the INK4 locus (ANRIL), can interact with PES1 to alter exonuclease-mediated pre-rRNA processing and ribosome biogenesis. Consequently, circANRIL can provoke apoptosis and impede cell proliferation in vascular smooth muscle cells and macrophages, promoting antiatherogenic functions [88]. Given the stability of circRNAs, circANRIL might also be used as a potential therapeutic target for the treatment of atherosclerosis. Li and coworkers reported that circRNAs ABCA1 and KHDRBS1 were significantly upregulated in atherosclerotic aortic vessels and H₂O₂-treated mouse aortic endothelial cells (MAECs). Consequently, they impacted atherosclerosis and vascular en-

endothelial injury by targeting the miR-30d-3p/TP53RK and miR-140-3p/MKK6 axes, as well as their downstream signaling pathways [89]. Likewise, Wang and coworkers [90] using human umbilical vascular endothelial cells (HUVECs) treated with oxidized low-density lipoprotein (ox-LDL), observed the role of circ_0124644 in promoting ox-LDL-induced endothelial injury of HUVECs through the miR-149-5p/PAPP-A axis, emphasizing its diagnostic and therapeutic application in atherosclerosis. In contrast, Yu's group [91] demonstrated in human oxidized low-density lipoprotein (ox-LDL)-induced brain microvascular endothelial cells that circ_0003423 improved ox-LDL-induced injury by regulating the miR-589-5p/TET2 axis. In HUVEC cells treated with high-glucose-levels, Zhou and coworkers [92] observed an overexpression of circ_0008360. Accordingly, circ_0008360 knockdown reduced the high glucose-induced vascular endothelial dysfunction by regulating miR-186-5p and cyclin D2, suggesting that circ_0008360 might act as a target for the treatment of vascular endothelial dysfunction [92].

Furthermore, Huang and coworkers [93] discovered that circ-RELL1 has a proinflammatory role in ox-LDL-induced HUVECs using high-throughput circRNA microarray assays. Accordingly, knockdown of circ-RELL1 decreased ICAM1 and VCAM1 expression in ox-LDL induced endothelium inflammation, by directly targeting miR-6873-3p in the cytoplasm. This reduction in the miR-6873-3p determines decreases in MyD88 (myeloid differentiation primary response 88) protein expression and MyD88 mediated NF- κ B activation. Furthermore, circ-RELL1 could abrogate the inhibition of inflammation response by acting on miR-6873-3p. Thus, circ-RELL1 modulates endothelium inflammation via miR-6873-3p/MyD88/NF- κ B axis in ox-LDL induced endothelial cells, evidencing its role as a potential therapeutic candidate for endothelium inflammation in atherosclerotic cardiovascular disease [93].

4.2 Limitations and Future Perspectives

The elevated stability, great quantity, and spatiotemporal specific expression of blood circRNAs suggest such molecules as perfect biomarkers for endothelium investigations using liquid biopsy. However, a biomarker with broad clinical application must have proven analytical validity, clinical validity, and clinical utility [94]. Consequently, several questions need to be studied before peripheral blood circRNA biomarkers can be translated into clinical practice. First, a blood circRNA-based gene analysis should prove its analytical validity within clinically relevant conditions. Even if significant improvements have been made in the past several years [95], the methods for identifying profiles of circRNAs are far from ideal. Forthcoming studies need to test the analytical performance of various circRNA profiling methods in clinical blood samples, such as RNA-seq, circRNA microarray, reverse transcription quantitative PCR (RT-qPCR) and RTddPCR [95]. In estimating analytical sensitivity and specificity, reference standards that can be specifically applied in circRNA discovery and profiling are needed (Hardwick et al., 2017). Furthermore, the procedure to uncover and confirm blood circRNA biomarkers needs to be standardized, including blood collection and preservation, circRNA extraction, library construction, and computational analysis [95, 96]. With the use of a standardized procedure, future studies need to estimate the technical robustness and reproducibility of the proposed biomarkers within and between laboratories. In addition, the blood circRNA biomarkers identified in current studies are only preliminary biomarker signatures for endothelium dysfunction [96]. Lastly, to examine their clinical validity, more clinical samples are necessary to prove their sensitivity and specificity in a larger sample size.

5. ADAR Protein Family as Biomarkers and Therapeutic Targets

Adenosine to inosine (A-to-I) RNA editing represents a post-transcriptional process that allows for selectively converting adenosines to inosines in double-stranded RNA (dsRNA) substrates [97]. Such a process is executed by a highly conserved group of enzymes, i.e., members of the adenosine deaminase acting on RNA (ADAR) family. Precisely, ADARs can alter the splicing and translation machineries, double-stranded RNA

structures, and binding affinity between RNA and RNA-binding proteins. All ADAR components are characterized by a structure with a common domain having a variable number of amino-terminal dsRNA-binding domains (dsRBDs) and a carboxy-terminal catalytic deaminase domain. In humans, there are three ADAR enzymes defined as ADAR1, ADAR2, and ADAR3 [97]. ADAR1 and ADAR2 are expressed in a ubiquitous manner in the tissues and act on numerous RNA substrates. ADAR3 is expressed only in the brain and appears to be inactive, as assessed on various RNA substrates [98]. In addition, two isoforms of ADAR1 have been identified, defined as ADAR1p150 and ADAR1p110, with cytoplasmatic and nuclear localization, respectively. The cytoplasmatic location of ADAR1p150 is unusual, because ADAR proteins generally map in the nucleus, where their presence is dependent on functional dsRBDs. dsRNAs originating from ribosomal RNAs (rRNAs) and small nucleolar RNAs (snoRNAs) can guide their location [97,98].), which seems to be a regulatory mechanism for the editing of premRNAs through enzyme sequestration. However, additional ADAR roles are emerging, linked to a large spectrum of genes and proteins, such as diverse cell pathways regulated by ADAR family proteins and related to phenotypic changes in tissue cells. Accordingly, it is possible to underline that ADAR activity is an essential mechanism in mammals, and altered editing has been associated with several human diseases, ranging from neurological and neurodegenerative diseases (schizophrenia, Alzheimer's disease, depression, epilepsy, amyotrophic lateral sclerosis (ALS), and systemic lupus erythematosus (SLE)) to cancers and cardiovascular diseases [99,100]. Such evidence has led to numerous efforts to modify ADAR activity in vivo and correct eventual RNA editing dysfunction.

5.1. ADAR Enzymes in the Endothelium Literature

Among the multiple roles of ADAR enzymes, it was evidenced by Zhang group [101] that ADAR1 shows the capacity to regulate endothelial cytokine activation. Precisely, they used HUVECs as an in vitro model to examine the role of ADAR1 in interleukin (IL)-1 β -stimulated endothelial activation. They observed that stimulation with IL-1 β affected the expression of ADAR1 and the adherence molecules VCAM-1 and ICAM-1 in HUVECs. Specifically, ADAR1 overexpression reduced the expression of VCAM-1 and ICAM-1, while ADAR1 knockdown augmented the expression of ICAM-1 and VCAM-1, as well as the recruitment of THP-1 monocytes. Furthermore, using a luciferase reporter assay, it was proven that ADAR1 can act as a direct target of miR-143 in HUVECs, and that miR-143 overexpression can exacerbate this condition, while miR-143 knockdown can stop IL-1 β -induced activation of HUVECs. Moreover, ADAR1 overexpression was observed to mediate the augmenting effect of miR-143 overexpression on IL-1 β -induced HUVEC activation. Specifically, it was demonstrated that ADAR1 overexpression could reduce this effect, while ADAR1 knockdown could increase the PKR, I κ B α , and NF- κ B phosphorylation generated by IL-1 β . In addition, the authors demonstrated that blocking NF- κ B signaling with specific NF- κ B inhibitor PDTC (pyrrolidine dithiocarbamate) could block the IL-1 β -induced HUVEC activation augmented by ADAR1 knockdown. Thus, these data suggest that ADAR1 can be targeted by miR-143 in regulating IL-1 β -induced HUVEC activation, and that the NF- κ B pathway can act as the downstream mediator of ADAR1. In conclusion, miR-143 and ADAR1 may represent therapeutic targets for endothelium dysfunction [101].

Another European group [102] discovered that EC showed an increased microRNA editing under ischemia in *vivo* in a murine hindlimb ischemia model and *ex vivo* in human veins, associated with a differential expression of ADAR1 and ADAR2. Likewise, the Stellos group demonstrated an increased ADAR activity in EC, that can control inflammation in atherosclerotic cardiovascular disease [103]. In addition, ADAR1's function was also found to be associated with editing on specific RNA substrates such as cathepsin S (CTSS) and miRNA487b in an atherosclerotic state [104, 105].

5.2 Limitations and Future Perspectives

The recoding and editing events mediated by ADAR enzymes can likely impact the functions and activities of relevant proteins, RNAs, and miRNAs. RNA editing can play a part in endothelium dysfunction as reported above. Such A-to-I changes are part of an epigenetic process that can potentially be regulated and corrected. A feasible approach might be the modulation of ADAR expression (by overexpression or silencing) at a single-cell level. Alternatively, specific editing sites with a key role in endothelial dysfunction and complications could be targets of therapeutic strategies aimed at preventing such conditions, by rescuing the correct editing version. In this regard, it is worth noting that the inhibition of RNA editing at a specific site is possible (Schirle et al., 2010) thanks to small molecules (such as antisense oligonucleotides or helix-threading peptides) that can abrogate the duplex structure required for editing [106]. A recent revolutionary study by Montiel-Gonzalez and coworkers [107] reported innovative therapeutic strategies possible thanks to ADAR enzymes. The increasing importance of ADARs in the context of endothelium dysfunction and complications provides newfound hope of inducing important changes in the landscape of future biomedicine.

6. Vascular Endothelial Glycocalyx and Its Components as Emerging Endothelial Biomarkers and Therapeutic Targets of Endothelium Dysfunction

ECs owe their physiological, anti-inflammatory, and anticoagulant properties to the complex carbohydrate-rich layer covering their luminal surface, named the glycocalyx. [108, 109]. Recent evidence has suggested that glycocalyx shedding affects the endothelium, causing dysfunction and inflammation, which emphasizes the value of glycocalyx maintenance for preventing disease induction and progression [108, 109]. Accordingly, the endothelial glycocalyx (EG) realizes diverse functions, including mechanotransduction, maintenance of vascular integrity and vascular tone, support of the production of nitric oxide (NO), and anti-inflammatory and anticoagulant properties by interacting with plasma proteins and plasma cells [108-110]. In addition, it is composed of a scaffolding of proteoglycans (PGs), glycoproteins (GPs), and glycosaminoglycans (GAGs) associated with the underlying ECs. PGs are composed of a core protein and covalently attached GAG chains, represented by syndecans and glypicans. Syndecans, observed on the surface of most body cells, are transmembrane proteins with extracellular, transmembrane, and cytosolic domains [108-110]. The extracellular domain binds GAGs and detects extracellular signals, such as shear stress, which are transduced to the intracellular environment via the transmembrane portion and the cytoplasmic tail. There are four well-known syndecans in humans, syndecans 1, 2, 3, and 4; however, the EG primarily contains syndecan 1 (SDC1). Glypicans are not transmembrane proteins; they are instead attached to the EC luminal membrane via a glycosylphosphatidylinositol anchor. There are six well-known glypicans, with glypican-1 expressed on the endothelium [108-110].

Diverse factors, including disturbed or reduced flow profiles, vascular aging, diabetes, and obesity determine biochemical and structural alterations in EG associated with the onset of endothelium dysfunction [108-111]. In addition, recent evidence has shown that EG is considered as one of pathological conditions commonly accompanying an ischemia/reperfusion injury (IRI), along with tissue damage due to ROS produced upon reperfusion, mitochondrial dysfunction with opening of the mitochondrial transition pore and loss of ATP production, activation of the complement system, and endothelial dysfunction [112, 113].). Animal models of cardiac IRI have demonstrated an early reduction in the thickness of EG, precisely 5 min after reperfusion, whereby EG shedding causes a reduced vasodilation of the endothelium mediated by NO. In addition, it was demonstrated that the EG can regenerate over time; specifically, some in vivo studies have revealed that about 7 days are sufficient to reestablish the EG layer. However, the mechanism underlying EG restoration and whether or not its physiological composition is maintained has not yet been explored [113]. However, various preventive and restorative approaches, with the aim of recreating physiological EG functions, have been proposed as potential therapies for endothelial dysfunction and its complications. Among

these, the application of diverse molecules, i.e., angiopoietin-1 [114], hydrocortisone [115], ATIII [116], and SOD [117] increased the vascular integrity and prevented endothelial dysfunction during an IRI by inhibiting the degradation of glycocalyx components syndecan-1, heparan sulfate and hyaluronic acid; therefore, they have been proposed as potential therapeutic candidates. However, how these molecules maintain EG integrity is still ambiguous.

Furthermore, many studies have demonstrated that some drugs [118], restoration with plasma [119], heparin [120, 121], sulodexide [122], and hydroxyethyl starch can downregulate heparinase, hyaluronidase and neuraminidase [123], during inflammation or sepsis, consequently preventing endothelial dysfunction by preserving EG integrity. However, in this case, the corresponding mechanism of action also remains unclear. Thus, further studies are needed to confirm such a hypothesis.

7. Circulating Metabolites

Numerous metabolic dysregulations accompany the dysfunctional endothelium, as well as its related pathologies. Metabolites constitute both small molecules of the human metabolome, including amino acids, organic acids, nucleic acids, fatty acids, amines, sugars, vitamins, cofactors, pigments, and antibiotics (i.e., endogenous metabolites) and exogenous chemical metabolites (i.e., drugs, environmental contaminants, food additives, toxins, and other xenobiotics) produced by metabolic reactions. Thanks to diverse advanced metabolomics platforms, including nuclear magnetic resonance (NMR) spectroscopy, mass spectrometry (MS), liquid chromatography (LC), gas chromatography, and capillary electrophoresis, it is possible to both quantify and separate these compounds.

Some research groups have suggested the presence in circulation of diverse profiles of metabolites in the presence or absence of endothelial dysfunction or related pathologies. The Kots group [124] evaluated the associations of circulating levels of stable metabolites of nitric oxide, nitrate, nitrite (NOx), and endothelin-1, as well as the endothelin-1/NOx ratio with blood pressure, in 177 asymptomatic subjects without signs of coronary atherosclerosis, and in 457 patients with presence of coronary lesions and suspected to have coronary heart disease with or without coronary lesions confirmed by coronary angiography. They observed that men had NOx levels inversely correlated with blood pressure and did women with a low (0-4%) European Systematic Coronary Risk Estimation (SCORE). Furthermore, a significant negative association was evidenced between high systolic blood pressure over 140 mm Hg and NOx levels in asymptomatic men ($p=0.05$) but not in women. In addition, in male patients, NOx ($p=0.05$), endothelin ($p=0.01$), and the endothelin/NOx ratio ($p=0.04$) were significantly associated with coronary lesions.

Beckman and coworkers [125] recently demonstrated in patients affected by type 2 diabetes and healthy controls the presence of different metabolites in plasma samples, including homocysteine, dimethylguanidino valeric acid and β -alanine (all $p<0.05$). Furthermore, they evidenced the presence of changes in the levels of ischemia-induced metabolites between the two groups including 5-hydroxyindoleacetic acid (healthy: -27%; DM +14%), orotic acid (healthy: +5%; DM -7%), trimethylamine-N-oxide (healthy: -51%; DM +0.2%), and glyoxylic acid (healthy: +19%; DM -6%) (all $p<0.05$). Plasma amounts of serine, betaine, β -aminoisobutyric acid and anthranilic acid were significantly associated with vessel diameter at baseline, but only in T2DM cases (all $p<0.05$). The results have suggested that dysregulated muscle metabolism in T2DM may have direct effects on vascular function.

The Wang group [126] evidenced that healthy endothelium releases metabolites able to modulate inflammatory responses and protect against systemic inflammation. Precisely, they detected, in conditioned endothelium extracts using liquid chromatography-mass spectrometry, the presence of 5-methoxytryptophan (5-MTP), but not other related tryptophan metabolites. Furthermore, it was observed that endothelial cell-derived 5-MTP can suppress lipopolysaccharide-induced inflammatory responses

and signaling in macrophages and endotoxemic lung tissues. Thus, it was suggested that 5-MTP may be considered as part of a novel class of endothelium-derived protective metabolites that defend against endothelial barrier dysfunction and excessive systemic inflammatory responses.

Such results are promising; however, a larger number of studies are needed to confirm the role of such or other metabolites as biomarkers related to a healthy or dysfunctional endothelium. In addition, it is known that metabolomics deals with the end-products of gene expression in cells; consequently, this aspect needs to be strongly clarified, as well as the relationship among genetic variations, environmental factors, and metabolite endothelial profiles. Thus, further longitudinal studies may reveal novel metabolic pathways and biomarkers for improving personalized therapy of endothelial-related diseases.

8. Suggestions and Recommendations for Further Investigations Needed to Develop Biomarker Endothelial Profiles

The choice of and progress in identifying endothelial biomarkers should be based on a clear knowledge of the mechanisms and pathways related to the maintenance of endothelial health or its dysfunction. With that in mind, the several pathways and molecules described above have not completely achieved such a purpose. They can clarify only part of the strong relationships between endothelium dysfunction and the onset of related diseases in various body tissues. For example, they do not allow for revealing the severity of injury, which would help practitioners to immediately intervene with personalized treatments or predict complications. In addition, given the long-term functional impairments of the endothelium throughout adult life caused by multiple factors and related inflammation, identifying diagnostic biomarkers to correlate with the severity of injury, prognostic biomarkers for early prediction of related diseases, and diagnostic biomarkers for monitoring the progression of the dysfunctional endothelium, along with regression following appropriate treatments, could serve to strongly reduce the high incidence of frequent endothelial-related diseases in the elderly. Furthermore, understanding the dynamic and temporal biomarker levels might enable the development of a potent algorithm and reveal novel putative therapeutic targets.

However, these observations suggest that further scientific efforts are needed for the identification of biomarkers and targets for the development of therapies. To achieve this goal, further advances should be realized using a new technological appraisal based on innovative approaches and systems, as mentioned above. The integration of multi-omics analyses (genomics, epigenomics, transcriptomics, and proteomics, along with metabolomics, microbiomics, and nutrigenomics) and the introduction of new-generation multi-omics technologies are encouraged [30]. In the next section, we describe some of these promising technologies.

9. Innovative Technologies for Targeted Endothelium Treatments: Nanomedicine and Its Benefits and Limitations

The term nanomedicine indicates a specific targeted treatment, which can enhance the delivery of targeted drugs (i.e., natural antioxidant compounds or pharmacological drugs) and their bioavailability, as well as reduce the associated toxicity or side-effects and costs, using nanoparticles (NPs) [127]. NPs comprise liposomes, niosomes, polymers, lipid and organic polymer hybrids and precursors, carbon nanotubes, quantum dots, metals, and metal oxides (see Figure 6).




Figure 6. Innovative technologies for targeted endothelium treatments: nanomedicine + CRISPR/Cas9 editing.

Thus, they constitute specialized carriers with the potential to facilitate the delivery of drugs and efficient molecular targets into desired tissues, such as the endothelium. Furthermore, NPs show another benefit in improving the effectiveness of therapeutic compounds, i.e., by decreasing their toxicity or side-effects. To this aim, NPs feature two aspects: (1) a versatile preparation method and a structure that allows encapsulating therapeutic agents, and (2) a functional surface able to interact with antibodies, aptamers, small molecules, etc. to further facilitate the delivery of therapeutics or diagnostics to pathological sites of the endothelium [127, 128]. Accordingly, the linking of NP surfaces with molecular fractions mimicking natural ligands can provide selective targeting and precise delivery of therapeutics to a diseased tissue. Nevertheless, only a small percentage of these NPs can successfully accumulate in the selected tissues. For example, Wilhelm and coworkers [129] that fewer than 1% of nanoparticles manage to accomplish their role, specifically related to tumoral tissues. The reasons are diverse and mostly related to multiple physiological barriers and the high degree of stochasticity involved in NP activity. Accordingly, a high percentage of NPs are phagocytized by the mononuclear phagocytic system (MPS), while some remain physically “entrapped” in the sinusoids of the liver and others are taken up by hepatocytes and Kupffer cells [127]. In addition, NPs are incapable of adequately negotiating other biological barriers. This has hampered their clinical translation. Consequently, innovative biomimetic strategies [127] have been developed. They are mainly based on two approaches: (1) top-down approaches, including the bioengineering of pathogens (bacteria and viruses) or cells (leukocytes, erythrocytes, platelets, and stem cells, [130] or (2) bottom-up approaches, such as conjugating NP surfaces with analogs of bioactive molecules that bind CAMs and selectins [131, 132] or coating synthetic particles with cell membranes as reported by Parodi and coworkers [133]. Additional approaches include poly (lactic-co-glycolic acid) (PLGA) NPs cloaked with platelet cell membrane that preferentially bind the denuded artery in a rat model of coronary restenosis [131] Other biomimetic NPs exhibit endogenous surface molecules, such as LDL and HDL, which have been used for incorporating nucleic material and delivering both therapeutic and diagnostic molecules [134].

Another important issue is the assessment of NPs biodistribution following *in vivo* administration in animals and humans [135, 136]. This constitutes a great challenge despite the large range of techniques available for detecting nanoparticle biodistribution, including histology, electron microscopy, liquid scintillation counting (LSC), indirect measurement of drug concentrations, *in vivo* optical imaging, computed tomography (CT), magnetic resonance imaging (MRI), and nuclear medicine imaging.

However, a biomimetic innovative approach has been utilized to improve the bio-availability, stability, and consequent advantageous actions of curcumin on endothelial related diseases [136]. Similarly, pH and ROS dual-responsive NPs fabricated by incor-

porating pH- and ROS-responsive cyclodextrin materials with resveratrol were utilized as an efficient and secure nanoplatform for therapeutic delivery to areas of vascular inflammation, in the presence of acidosis and oxidative stress [137,138]. Cho and co-workers also used biomimetic NPs with synthetic HDL (liposomal formulation with dimyristoyl phosphatidylcholine (DPMC)), demonstrating that they could significantly enhance the blood HDL levels, as well as reduce the plaque development in an animal study [139]. In another study, using a murine model of inflammation, the researchers have targeted E-selectins by conjugating anti-E-selectin monoclonal antibodies to ultra-small-SPION *in vivo* for imaging of the endothelium to detect vascular inflammation [31]

9.1. Consideration of NPs as New Endothelium Treatment Strategy

Currently, newer treatment strategies, with reduced risks of evoking adverse effects, are being investigated, among which nanomedicine shows a more targeted specificity and efficacy. Thanks to the discovery of various endothelium-targeting molecules, efforts to design NP treatments targeting specific endothelium sites and ligands using appropriate nano-carriers are increasing. However, there are several challenges, including the shear stress under continuous flow, the variable size of arteries and their marginalization from the bloodstream, and the elimination of NPs by macrophages, which affect NP internalization in the endothelium. Consequently, numerous delivery strategies have been developed to improve the adequate transport of medication into specific endothelial regions, as mentioned above, including the production of biomimetic NPs. However, the number of translational studies is inadequate for elucidating the safety and efficacy of such NPs, and their delivery strategies remain a challenge. To progress to clinical application, NPs should interact with and internalize in the endothelium effectively, and they should be proven to be nontoxic, more successful than existing options, easy to use, and cost- and design-effective. Nevertheless, significant benchwork associated with an increased number of clinical studies may provide the necessary hope for treating endothelium dysfunction with nanotechnology [140].

Below, we evidence how the integration of NPs with other newer omics technologies can be used to successfully recovering endothelium function in adults.

9.2. Integrating NPs with CRISPR/Cas9 Editing and Base Primers

In the case of the endothelium, NP application shows several limitations, as largely stressed above. While developing such treatments, it is imperative to consider the high degree of endothelium heterogeneity and complexity, owing to its diverse morphology, location, and function shown in various body tissues. Endothelial alterations are directly or indirectly involved in a wide variety of human disorders, encompassing several mechanisms and pathways (including inflammation), and they are caused by and promote the progression of endothelial dysfunction. This can severely affect other cellular types, in physiological or pathological conditions, impacting the function of organs and systems, as well as physiological processes, such as regeneration, repair, cardiovascular genesis, neurogenesis, and cardiovascular and neural survival. This is significantly related to the altered gene expression of cells linked or not to genetic variants and/or genomic modifications. Consequently, it is hypothesized that a strong genetic editing technique might be a solution, as well as the integration of gene-editing techniques with nanotechnology. Accordingly, the clustered regularly interspaced short palindromic repeats-associated protein 9 (CRISPR/Cas9) system and techniques to chemically modify RNAs are also being incorporated in NPs to study human pathologies and/or examine potential disease correction through restoration [141,142]. The CRISPR/Cas9 system is significantly utilized for both editing the genome of zygotes, thereby generating genetically modified animal species, and treating human diseases. In fact, the development of safe and efficient treatments is becoming a reality by delivering the CRISPR/Cas9 system into body tissues and targeting specific cells [143, 144]. A delivery system with optimal *in vivo* genome-editing efficacy is based on the use of the recombinant adeno-associated

virus (AAV), but it can cause adverse effects. Thus, lipid NPs represent an alternative. For example, lipid NPs were used to target macrophages in mice using the CRISPR/Cas9 system, although a reduced genome-editing percentage (20%) was obtained. This suggests that it is a challenge to induce robust genome editing in the targeted endothelium. However, hope can be derived from a recent study conducted by the Zhang group on ECs obtained from the lung, heart, aorta, and peripheral vessels of adult mice. Precisely, Zhang and coworkers [145] developed poly(ethylene glycol) methyl ether-*block*-poly(lactide-*co*-glycolide) (PEG-*b*-PLGA; PP)-based NPs with excellent bio-distribution for vascular delivery and demonstrated that the polyethyleneimine (PEI)-formulated PP NP-mediated delivery of the all-in-one CRISPR plasmid DNA expressing Cas9 under the control of the human CDH5 promoter and a guide RNA (gRNA) driven by the U6 promoter determined highly efficient genome editing in ECs of various vascular locations with a single administration. A reduction in protein expression of about 80% was evidenced in ECs. The potential demonstrated by such an approach could allow efficiently modulating or eliminating the expression of genes encoding proteins strongly associated with phenotypic changes related to the development of endothelium dysfunction and its related pathologies [145]. Today, several new CRISPR-based modalities have been developed that are capable of facilitating gene editing without the requirement of a DNA double-strand break (DSB), including base editors, prime editors, and RNA-targeting CRISPR-associated protein (Cas)13 effectors. These modalities can potentially be integrated with NPs. However, this will require completely optimizing individual base editor proteins for specific therapeutic targets to guarantee maximally efficient formation of the on-target product, as well as minimize potentially counter-productive nontarget editing outcomes. In addition to developing methods to further refine the capabilities of these increasingly sophisticated gene-editing machines, special attention must be given to their delivery. Thus, further studies are needed.

10. Conclusions

Individualized profiling of molecular patterns identified in bodily fluids represents a revolutionary approach in the framework of detecting biomarkers and developing therapeutic targeted treatments for preserving, reverting, and restoring endothelium barrier integrity and function. Currently, EPCs, circulating metabolites, EG components, circRNAs, and ADAR enzymes are attracting substantial attention for diagnostic and treatment purposes (see **Figure 7**).




Figure 7. Innovative non-invasive biomarkers and therapeutic targets

Altered profiles of such molecules and cells have been detected under both physiological and pathological conditions of the endothelium. However, advanced implementation of such research is imperative for achieving these purposes independently of the application area.

This article highlighted the involvement of such molecules and cells in the healthy and diseased endothelium by addressing whether specific patterns of such molecules and cells in fluid biopsy and the use of newer omics technologies or their combinations can be further applied to accompany diagnosis, targeted prevention, creation of individualized therapy algorithms, therapeutic monitoring, and prognosis. The presented considerations correspond with the principles of network and personalized medicine, and they can be ameliorated to increase individual long-term survival outcomes, thereby guaranteeing a high quality of life and improved cost efficacy of medical services provided to the population.

Conflicts of Interest: The authors declare no conflicts of interest.

Funding: This research was funded by an Italian University Minister FFABR grant 2017 to C.R.B. and University of Palermo FFR 2019/2020 grants to C.R.B.

References

1. Inai K. Atrioventricular Valve Abnormalities: From Molecular Mechanisms Underlying Morphogenesis to Clinical Perspective. 2016 Jun 25. In: Nakanishi T, Markwald RR, Baldwin HS, Keller BB, Srivastava D, Yamagishi H, editors. Etiology and Morphogenesis of Congenital Heart Disease: From Gene Function and Cellular Interaction to Morphology [Internet]. Tokyo: Springer; 2016. Chapter 17.
2. Vita JA. Endothelial function. *Circulation*. 2011 Dec 20;124(25):e906-12. doi: 10.1161/CIRCULATIONAHA.111.078824.
3. Sabe SA, Feng J, Sellke FW, Abid MR. Mechanisms and Clinical Implications of Endothelium-dependent Vasomotor Dysfunction in Coronary Microvasculature. *Am J Physiol Heart Circ Physiol*. 2022 Mar 25. doi: 10.1152/ajpheart.00603.2021.
4. Boulanger CM. Endothelium. *Arterioscler Thromb Vasc Biol*. 2016 Apr;36(4):e26-31. doi: 10.1161/ATVBAHA.116.306940.
5. Balistreri CR, Madonna R, Melino G, Caruso C. The emerging role of Notch pathway in ageing: Focus on the related mechanisms in age-related diseases. *Ageing Res Rev*. 2016 Aug;29:50-65. doi: 10.1016/j.arr.2016.06.004.
6. Balistreri CR, Garagnani P, Madonna R, Vaiserman A, Melino G. Developmental programming of adult haematopoiesis system. *Ageing Res Rev*. 2019 Sep;54:100918. doi: 10.1016/j.arr.2019.100918.
7. Allegra A, Giarratana RM, Scola L, Balistreri CR. The close link between the fetal programming imprinting and neurodegeneration in adulthood: The key role of "hemogenic endothelium" programming. *Mech Ageing Dev*. 2021 Apr;195:111461. doi: 10.1016/j.mad.2021.111461.
8. Schirò G, Balistreri CR. The close link between brain vascular pathological conditions and neurodegenerative diseases: Focus on some examples and potential treatments. *Vascul Pharmacol*. 2022 Feb;142:106951. doi: 10.1016/j.vph.2021.106951.
9. Pober JS, Min W, Bradley JR. Mechanisms of endothelial dysfunction, injury, and death. *Annu Rev Pathol*. 2009;4:71-95. doi: 10.1146/annurev.pathol.4.110807.092155.
10. Regina C, Panatta E, Candi E, Melino G, Amelio I, Balistreri CR, Annicchiarico-Petruzzelli M, Di Daniele N, Ruvolo G. Vascular ageing and endothelial cell senescence: Molecular mechanisms of physiology and diseases. *Mech Ageing Dev*. 2016 Oct;159:14-21. doi: 10.1016/j.mad.2016.05.003. Epub 2016 May 4. PMID: 27155208.
11. Madonna R, Novo G, Balistreri CR. Cellular and molecular basis of the imbalance between vascular damage and repair in ageing and age-related diseases: As biomarkers and targets for new treatments. *Mech Ageing Dev*. 2016 Oct;159:22-30. doi: 10.1016/j.mad.2016.03.005. Epub 2016 Mar 15. PMID: 26993150.
12. Wang YF, Hsu YJ, Wu HF, Lee GL, Yang YS, Wu JY, Yet SF, Wu KK, Kuo CC. Endothelium-Derived 5-Methoxytryptophan Is a Circulating Anti-Inflammatory Molecule That Blocks Systemic Inflammation. *Circ Res*. 2016 Jul 8;119(2):222-36. doi: 10.1161/CIRCRESAHA.116.308559.

13. Miura S, Morimoto Y, Furihata T, Takeuchi S. Functional analysis of human brain endothelium using a microfluidic device integrating a cell culture insert. *APL Bioeng*. 2022 Mar 9;6(1):016103. doi: 10.1063/5.0085564.
14. Jalnapurkar S, Landes S, Wei J, Mehta PK, Shufelt C, Minissian M, Pepine CJ, Handberg E, Cook-Wiens G, Sopko G, Bairey Merz CN. Coronary endothelial dysfunction appears to be a manifestation of a systemic process: A report from the Women's Ischemia Syndrome Evaluation - Coronary Vascular Dysfunction (WISE-CVD) study. *PLoS One*. 2021 Sep 27;16(9):e0257184. doi: 10.1371/journal.pone.0257184.
15. Meilhac A, Cautela J, Thuny F. Cancer Therapies and Vascular Toxicities. *Curr Treat Options Oncol*. 2022 Mar;23(3):333-347. doi: 10.1007/s11864-022-00964-2.
16. Camilli M, La Vecchia G, Lillo R, Iannaccone G, Lamendola P, Montone RA, Hohaus S, Aspromonte N, Massetti M, Lanza GA, Crea F, Graziani F, Lombardo A. Cardiovascular involvement in patients affected by multiple myeloma: a comprehensive review of recent advances. *Expert Rev Hematol*. 2021 Nov 29;1-14. doi: 10.1080/17474086.2021.2003704.
17. Love KM, Barrett EJ, Malin SK, Reusch JEB, Regensteiner JG, Liu Z. Diabetes pathogenesis and management: the endothelium comes of age. *J Mol Cell Biol*. 2021 Oct 21;13(7):500-512. doi: 10.1093/jmcb/mjab024.
18. Trejo-Lopez JA, Yachnis AT, Prokop S. Neuropathology of Alzheimer's Disease. *Neurotherapeutics*. 2021 Nov 2. doi: 10.1007/s13311-021-01146-y.
19. Olivieri F, Prattichizzo F, Grillari J, Balistreri CR. Cellular Senescence and Inflammaging in Age-Related Diseases. *Mediators Inflamm*. 2018 Apr 17;2018:9076485. doi: 10.1155/2018/9076485.
20. Peng H, Wang X, Du J, Cui Q, Huang Y, Jin H. Metabolic Reprogramming of Vascular Endothelial Cells: Basic Research and Clinical Applications. *Front Cell Dev Biol*. 2021 Feb 18;9:626047. doi: 10.3389/fcell.2021.626047. PMID: 33681205; PMCID: PMC7930387.
21. Jaul E, Barron J. Characterizing the Heterogeneity of Aging: A Vision for a Staging System for Aging. *Front Public Health*. 2021 Oct 12;9:513557. doi: 10.3389/fpubh.2021.513557.
22. Pisano C, Poliso D, Balistreri CR, Altieri C, Nardi P, Bertoldo F, Trombetti D, Asta L, Ferrante MS, Buioni D, Foti C, Ruvolo G. Role of Cachexia and Fragility in the Patient Candidate for Cardiac Surgery. *Nutrients*. 2021 Feb 5;13(2):517. doi: 10.3390/nu13020517.
23. Iglesias MJ, Kruse LD, Sanchez-Rivera L, Enge L, Dusart P, Hong MG, Uhlén M, Renné T, Schwenk JM, Bergstrom G, Odeberg J, Butler LM. Identification of Endothelial Proteins in Plasma Associated With Cardiovascular Risk Factors. *Arterioscler Thromb Vasc Biol*. 2021 Dec;41(12):2990-3004. doi: 10.1161/ATVBAHA.121.316779.
24. Amraei R, Rahimi N. COVID-19, Renin-Angiotensin System and Endothelial Dysfunction. *Cells*. 2020 Jul 9;9(7):1652. doi: 10.3390/cells9071652.
25. Pons S, Fodil S, Azoulay E, Zafrani L. The vascular endothelium: the cornerstone of organ dysfunction in severe SARS-CoV-2 infection. *Crit Care*. 2020 Jun 16;24(1):353. doi: 10.1186/s13054-020-03062-7.
26. Zhang J, Tecson KM, McCullough PA. Endothelial dysfunction contributes to COVID-19-associated vascular inflammation and coagulopathy. *Rev Cardiovasc Med*. 2020 Sep 30;21(3):315-319. doi: 10.31083/j.rcm.2020.03.126.
27. Lio D, Scola L, Giarratana RM, Candore G, Colonna-Romano G, Caruso C, Balistreri CR. SARS CoV2 infection _The longevity study perspectives. *Ageing Res Rev*. 2021 May;67:101299. doi: 10.1016/j.arr.2021.101299.
28. Wingo M, Rafii S. Endothelial reprogramming for vascular regeneration: Past milestones and future directions. *Semin Cell Dev Biol*. 2022 Feb;122:50-55. doi: 10.1016/j.semcdb.2021.09.003.
29. Leon-Mimila P, Wang J, Huertas-Vazquez A. Relevance of multi-omics studies in cardiovascular diseases. *Front Cardiovasc Med*. 2019;6.
30. Scola L, Giarratana RM, Torre S, Argano V, Lio D, Balistreri CR. On the Road to Accurate Biomarkers for Cardiometabolic Diseases by Integrating Precision and Gender Medicine Approaches. *Int J Mol Sci*. 2019 Nov 29;20(23):6015. doi: 10.3390/ijms20236015.
31. Greene JA, Loscalzo J. Putting the Patient Back Together - Social Medicine, Network Medicine, and the Limits of Reductionism. *N Engl J Med*. 2017;377:2493-9.
32. Cheng F, Lu W, Liu C, et al. A genome-wide positioning systems network algorithm for in silico drug repurposing. *Nat Commun*. 2019;10:3476.

33. Lee LY-H, Loscalzo J. Network medicine in pathobiology. *Am J Pathol* 2019;189:1311–26.
34. Abasht B, Papah MB, Qiu J. Evidence of vascular endothelial dysfunction in Wooden Breast disorder in chickens: Insights through gene expression analysis, ultra-structural evaluation, and supervised machine learning methods. *PLoS One*. 2021 Jan 4;16(1):e0243983. doi: 10.1371/journal.pone.0243983.
35. Treder M, Lauermann JL, Alnawaiseh M, Eter N. Using Deep Learning in Automated Detection of Graft Detachment in Descemet Membrane Endothelial Keratoplasty: A Pilot Study. *Cornea*. 2019 Feb;38(2):157-161. doi: 10.1097/ICO.0000000000001776.
36. Sun X, Feinberg MW. Vascular Endothelial Senescence: Pathobiological Insights, Emerging Long Noncoding RNA Targets, Challenges and Therapeutic Opportunities. *Front Physiol*. 2021 Jun 16;12:693067. doi: 10.3389/fphys.2021.693067.
37. Kumar S, Andueza A, Kim J, Kang DW, Jo H. Isolation of Endothelial Cells from the Lumen of Mouse Carotid Arteries for Single-cell Multi-omics Experiments. *J Vis Exp*. 2021 Oct 4;(176). doi: 10.3791/63128. PMID: 34661575.
38. Andueza A, Kumar S, Kim J, Kang DW, Mumme HL, Perez JL, Villa-Roel N, Jo H. Endothelial Reprogramming by Disturbed Flow Revealed by Single-Cell RNA and Chromatin Accessibility Study. *Cell Rep*. 2020 Dec 15;33(11):108491. doi: 10.1016/j.celrep.2020.108491. PMID: 33326796; PMCID: PMC7801938.
39. He M, Huang TS, Li S, Hong HC, Chen Z, Martin M, Zhou X, Huang HY, Su SH, Zhang J, Wang WT, Kang J, Huang HD, Zhang J, Chien S, Shyy JY. Atheroprotective Flow Upregulates ITPR3 (Inositol 1,4,5-Trisphosphate Receptor 3) in Vascular Endothelium via KLF4 (Krüppel-Like Factor 4)-Mediated Histone Modifications. *Arterioscler Thromb Vasc Biol*. 2019 May;39(5):902-914. doi: 10.1161/ATVBAHA.118.312301. PMID: 30917677; PMCID: PMC6536300.
40. Zhang M, Urabe G, Ozer HG, Xie X, Webb A, Shirasu T, Li J, Han R, Kent KC, Wang B, Guo LW. Angioplasty induces epigenomic remodeling in injured arteries. *Life Sci Alliance*. 2022 Feb 15;5(5):e202101114. doi: 10.26508/lsa.202101114. PMID: 35169042; PMCID: PMC8860099.
41. Pepin ME, Schiano C, Miceli M, Benincasa G, Mansueto G, Grimaldi V, Soricelli A, Wende AR, Napoli C. The human aortic endothelium undergoes dose-dependent DNA methylation in response to transient hyperglycemia. *Exp Cell Res*. 2021 Mar 15;400(2):112485. doi: 10.1016/j.yexcr.2021.112485. Epub 2021 Jan 27. PMID: 33515594; PMCID: PMC8038422.
42. Matsuoka RL, Buck LD, Vajrала KP, Quick RE, Card OA. Historical and current perspectives on blood endothelial cell heterogeneity in the brain. *Cell Mol Life Sci*. 2022 Jun 20;79(7):372. doi: 10.1007/s00018-022-04403-1. PMID: 35726097.
43. Benincasa G, Mansueto G, Napoli C. Fluid-based assays, and precision medicine of cardiovascular diseases: the 'hope' for Pandora's box? *J Clin Pathol*. 2019 Dec;72(12):785-799. doi: 10.1136/jclinpath-2019-206178.
44. Mattox AK, Bettgowda C, Zhou S, Papadopoulos N, Kinzler KW, Vogelstein B (2019) Applications of liquid biopsies for cancer. *Sci Transl Med* 11:eaay1984
45. Rubis G, Krishnan S, Bebawy M (2019) Liquid biopsies in cancer diagnosis, monitoring, and prognosis. *Trends Pharmacol Sci* 40:172–186
46. Perakis S, Speicher MR. Emerging concepts in liquid biopsies. *BMC Med* 2017;15:75.
47. Balistreri CR, Buffa S, Pisano C, Lio D, Ruvoło G, Mazzesi G. Are Endothelial Progenitor Cells the Real Solution for Cardiovascular Diseases? Focus on Controversies and Perspectives. *Biomed Res Int*. 2015;2015:835934. doi: 10.1155/2015/835934.
48. Balistreri, C. Endothelial progenitor cells: a new real hope or only an unrealizable dream? Springer International Publishing, Dordrecht pgg. 1–80 (2017)
49. Olivieri F, Pompilio G, Balistreri CR. Endothelial progenitor cells in ageing. *Mech Ageing Dev*. 2016 Oct;159:1-3. doi: 10.1016/j.mad.2016.09.002.
50. Altabas V, Altabas K, Kirigin L. Endothelial progenitor cells (EPCs) in ageing and age-related diseases: How currently available treatment modalities affect EPC biology, atherosclerosis, and cardiovascular outcomes. *Mech Ageing Dev*. 2016 Oct;159:49-62. doi: 10.1016/j.mad.2016.02.009.
51. Balistreri CR, Crapanzano F, Schirone L, Allegra A, Pisano C, Ruvoło G, Forte M, Greco E, Cavarretta E, Marullo AGM, Sciarretta S, Frati G. Deregulation of Notch1 pathway and circulating endothelial progenitor cell (EPC) number in patients with bicuspid aortic valve with and without ascending aorta aneurysm. *Sci Rep*. 2018 Sep 14;8(1):13834. doi: 10.1038/s41598-018-32170-2.
52. Balistreri CR, Pisano C, Bertoldo F, Massoud R, Dolci S, Ruvoło G. Red Blood Cell Distribution Width, Vascular Aging Biomarkers, and Endothelial Progenitor Cells for Predicting Vascular Aging and Diagnosing/Prognosing Age-Related Degenerative Arterial Diseases. *Rejuvenation Res*. 2019

- Oct;22(5):399-408. doi: 10.1089/rej.2018.2144. Epub 2019 Feb 11. PMID: 30572793.
53. Ferrante A, Guggino G, Di Liberto D, Ciccio F, Cipriani P, Balistreri CR, Sireci G, Giacomelli R, Triolo G. Endothelial progenitor cells: Are they displaying a function in autoimmune disorders? *Mech Ageing Dev.* 2016 Oct;159:44-48. doi: 10.1016/j.mad.2016.05.001.
 54. Loiola RA, García-Gabilondo M, Grayston A, Bugno P, Kowalska A, Duban-Deweer S, Rizzi E, Hachani J, Sano Y, Shimizu F, Kanda T, Mysiorek C, Mazurek MP, Rosell A, Gosselet F. Secretome of endothelial progenitor cells from stroke patients promotes endothelial barrier tightness and protects against hypoxia-induced vascular leakage. *Stem Cell Res Ther.* 2021 Oct 26;12(1):552. doi: 10.1186/s13287-021-02608-y.
 55. Morrone D, Picoi MEL, Felice F, De Martino A, Scatena C, Spontoni P, Naccarato AG, Di Stefano R, Bortolotti U, Dal Monte M, Pini S, Abelli M, Balbarini A. Endothelial Progenitor Cells: An Appraisal of Relevant Data from Bench to Bedside. *Int J Mol Sci.* 2021 Nov 28;22(23):12874. doi: 10.3390/ijms222312874.
 56. Ota Y, Kuwana M. Endothelial cells and endothelial progenitor cells in the pathogenesis of systemic sclerosis. *Eur J Rheumatol.* 2020 Oct;7(Suppl 3):S139-S146. doi: 10.5152/eurjrheum.2019.19158.
 57. Pulito-Cueto V, Remuzgo-Martínez S, Genre F, Atienza-Mateo B, Mora-Cuesta VM, Iturbe-Fernández D, Lera-Gómez L, Pérez-Fernández R, Prieto-Peña D, Portilla V, Blanco R, Corrales A, Gualillo O, Cifrián JM, López-Mejías R, González-Gay MA. Endothelial Progenitor Cells: Relevant Players in the Vasculopathy and Lung Fibrosis Associated with the Presence of Interstitial Lung Disease in Systemic Sclerosis Patients. *Biomedicines.* 2021 Jul 20;9(7):847. doi: 10.3390/biomedicines9070847.
 58. Xiao ST, Kuang CY. Endothelial progenitor cells and coronary artery disease: Current concepts and future research directions. *World J Clin Cases.* 2021 Oct 26;9(30):8953-8966. doi: 10.12998/wjcc.v9.i30.8953.
 59. Buffa S, Borzi D, Chiarelli R, Crapanzano F, Lena AM, Nania M, Candi E, Triolo F, Ruvolo G, Melino G, Balistreri CR. Biomarkers for vascular ageing in aorta tissues and blood samples. *Exp Gerontol.* 2019 Dec;128:110741. doi: 10.1016/j.exger.2019.110741. Epub 2019 Oct 21. PMID: 31648011.
 60. Balistreri CR, Pisano C, Bertoldo F, Massoud R, Dolci S, Ruvolo G. Red Blood Cell Distribution Width, Vascular Aging Biomarkers, and Endothelial Progenitor Cells for Predicting Vascular Aging and Diagnosing/Prognosing Age-Related Degenerative Arterial Diseases. *Rejuvenation Res.* 2019 Oct;22(5):399-408. doi: 10.1089/rej.2018.2144. Epub 2019 Feb 11. PMID: 30572793
 61. Hill JM, Zalos G, Halcox JP, Schenke WH, Waclawiw MA, Quyyumi AA, Finkel T. Circulating endothelial progenitor cells, vascular function, and cardiovascular risk. *N Engl J Med.* 2003 Feb 13;348(7):593-600. doi: 10.1056/NEJMoa022287. _
 62. Naruse K, Hamada Y, Nakashima E, Kato K, Mizubayashi R, Kamiya H, Yuzawa Y, Matsuo S, Murohara T, Matsubara T, Oiso Y, Nakamura J. Therapeutic neovascularization using cord blood-derived endothelial progenitor cells for diabetic neuropathy. *Diabetes.* 2005 Jun;54(6):1823-8. doi: 10.2337/diabetes.54.6.1823. Erratum in: *Diabetes.* 2006 May;55(5):1534. PMID: 15919805.
 63. Shyu WC, Lin SZ, Chiang MF, Su CY, Li H. Intracerebral peripheral blood stem cell (CD34+) implantation induces neuroplasticity by enhancing beta1 integrin-mediated angiogenesis in chronic stroke rats. *J Neurosci.* 2006 Mar 29;26(13):3444-53. doi: 10.1523/JNEUROSCI.5165-05.2006. PMID: 16571751; PMCID: PMC6673856.
 64. Gómez-Navarro J, Contreras JL, Arafat W, Jiang XL, Krisky D, Oligino T, Marconi P, Hubbard B, Glorioso JC, Curiel DT, Thomas JM. Genetically modified CD34+ cells as cellular vehicles for gene delivery into areas of angiogenesis in a rhesus model. *Gene Ther.* 2000 Jan;7(1):43-52. doi: 10.1038/sj.gt.3301054. PMID: 10680015.
 65. Ohtsuka M, Takano H, Zou Y, Toko H, Akazawa H, Qin Y, Suzuki M, Hasegawa H, Nakaya H, Komuro I. Cytokine therapy prevents left ventricular remodeling and dysfunction after myocardial infarction through neovascularization. *FASEB J.* 2004 May;18(7):851-3. doi: 10.1096/fj.03-0637fje. Epub 2004 Mar 4. PMID: 15001565.
 66. Yamamoto K, Kondo T, Suzuki S, Izawa H, Kobayashi M, Emi N, Komori K, Naoe T, Takamatsu J, Murohara T. Molecular evaluation of endothelial progenitor cells in patients with ischemic limbs: therapeutic effect by stem cell transplantation. *Arterioscler Thromb Vasc Biol.* 2004 Dec;24(12):e192-6. doi: 10.1161/01.ATV.0000147730.78953.60. Epub 2004 Oct 14. PMID: 15486312.
 67. Lenk K, Adams V, Lurz P, Erbs S, Linke A, Gielen S, Schmidt A, Scheinert D, Biamino G, Emmrich F, Schuler G, Hambrecht R. Therapeutical potential of blood-derived progenitor cells in patients with peripheral arterial occlusive disease and critical limb ischaemia. *Eur Heart J.* 2005 Sep;26(18):1903-9. doi: 10.1093/eurheartj/ehi285. Epub 2005 Apr 26. PMID: 15855189.
 68. Erbs S, Linke A, Adams V, Lenk K, Thiele H, Diederich KW, Emmrich F, Kluge R, Kendziorra K, Sabri O, Schuler G, Hambrecht R.

- Transplantation of blood-derived progenitor cells after recanalization of chronic coronary artery occlusion: first randomized and placebo-controlled study. *Circ Res.* 2005 Oct 14;97(8):756-62. doi: 10.1161/01.RES.0000185811.71306.8b. Epub 2005 Sep 8. PMID: 16151021.
69. Liu Y, Chen J, Liang H, Cai Y, Li X, Yan L, Zhou L, Shan L, Wang H. Human umbilical cord-derived mesenchymal stem cells not only ameliorate blood glucose but also protect vascular endothelium from diabetic damage through a paracrine mechanism mediated by MAPK/ERK signaling. *Stem Cell Res Ther.* 2022 Jun 17;13(1):258. doi: 10.1186/s13287-022-02927-8. PMID: 35715841; PMCID: PMC9205155.
 70. Chen L-L. The biogenesis and emerging roles of circular RNAs. *Nat Rev Mol Cell Biol* 2016, 17:205–211
 71. Chen L-L. The expanding regulatory mechanisms and cellular functions of circular RNAs. *Nat Rev Mol Cell Biol*, 2020, 21:1–16
 72. Li X, Yang L, Chen L-L The biogenesis, functions, and challenges of circular RNAs. *Mol Cell*, 2018 71:428–442
 73. Sanger HL, Klotz G, Riesner D, Gross HJ, Kleinschmidt AK. Viroids are single-stranded covalently closed circular RNA molecules existing as highly base-paired rod-like structures. *Proc Natl Acad Sci USA*, 1976, 73:3852–3856
 74. Hsu M-T, Coca-Prados M (1979) Electron microscopic evidence for the circular form of RNA in the cytoplasm of eukaryotic cells. *Nature* 280:339–340
 75. Wang PL, Bao Y, Yee M-C, Barrett SP, Hogan GJ, Olsen MN, Dinnyen JR, Brown PO, Salzman J. Circular RNA is expressed across the eukaryotic tree of life. *PLoS ONE*, 2014, 9(6):e90859
 76. Zhou T, Xie X, Li M, Shi J, Zhou JJ, Knox KS, Wang T, Chen Q, Gu W. Rat BodyMap transcriptomes reveal unique circular RNA features across tissue types and developmental stages. *RNA*, 2018, 24:1443–1456
 77. Hansen TB, Jensen TI, Clausen BH, Bramsen JB, Finsen B, Damgaard CK, Kjems J. Natural RNA circles function as efficient microRNA sponges. *Nature*, 2013, 495:384–388
 78. Huang A, Zheng H, Wu Z, Chen M, Huang Y. Circular RNA– protein interactions: functions, mechanisms, and identification. *Theranostics*, 2020, 10:3503–3517
 79. Li S, Zhang J, Tan X, Deng J, Li Y, Piao Y, Li C, Yang W, MoW, Sun J et al. Microarray expression profile of circular RNAs and mRNAs in children with systemic lupus erythematosus. *Clin Rheumatol*, 2019, 38:1339–1350
 80. Legnini I, Timoteo GD, Rossi F, Morlando M, Briganti F, Sthandier O, Fatica A, Santini T, Andronache A, Wade M et al. Circ-ZNF609 is a circular RNA that can be translated and functions in myogenesis. *Mol Cell*, 2017, 66:22–37
 81. Pamudurti NR, Bartok O, Jens M, Ashwal-Fluss R, Stottmeister C, Ruhe L, Hanan M, Wyler E, Perez-Hernandez D, Ramberger E et al. Translation of circRNAs. *Mol Cell*, 2017, 66:9–21
 82. Shang Q, Yang Z, Jia R, Ge S. The novel roles of circRNAs in human cancer. *Mol Cancer*, 2019, 18:6
 83. Vo JN, Cieslik M, Zhang Y, Shukla S, Xiao L, Zhang Y, Wu Y-M, Dhanasekaran SM, Engelke CG, Cao X et al. The landscape of circular RNA in cancer. *Cell*, 2019, 176:869-881.e813
 84. Aufiero S, Reckman YJ, Pinto YM, Creemers EE. Circular RNAs open a new chapter in cardiovascular biology. *Nat Rev Cardiol*, 2019, 16:503–514
 85. Zhou Z, Sun B, Huang S, Zhao L. Roles of circular RNAs in immune regulation and autoimmune diseases. *Cell Death Dis*, 2019, 10:503
 86. Zhang Z, Yang T, Xiao J. Circular RNAs: promising biomarkers for human diseases. *EBioMedicine*, 2018, 34:267–274
 87. Wu WP, Zhou MY, Liu DL, Min X, Shao T, Xu ZY, Jing X, Cai MY, Xu S, Liang X, Mo M, Liu X, Xiong XD. circGNAQ, a circular RNA enriched in vascular endothelium, inhibits endothelial cell senescence and atherosclerosis progression. *Mol Ther Nucleic Acids*. 2021 Aug 8;26:374-387. doi: 10.1016/j.omtn.2021.07.020.
 88. Holdt LM, Stahnger A, Sass K, Pichler G, Kulak NA, Wilfert W, Kohlmaier A, Herbst A, Northoff BH, Nicolaou A, Gäbel G, Beutner F, Scholz M, Thiery J, Musunuru K, Krohn K, Mann M, Teupser D. Circular non-coding RNA ANRIL modulates ribosomal RNA maturation and atherosclerosis in humans. *Nat Commun.* 2016 Aug 19;7:12429. doi: 10.1038/ncomms12429.

89. Li H, Liu X, Sun N, Wang T, Zhu J, Yang S, Song X, Wang R, Wang X, Zhao Y, Zhang Y. Differentially Expressed Circular Non-coding RNAs in Atherosclerotic Aortic Vessels and Their Potential Functions in Endothelial Injury. *Front Cardiovasc Med*. 2021 Jul 7;8:657544. doi: 10.3389/fcvm.2021.657544.
90. Wang G, Li Y, Liu Z, Ma X, Li M, Lu Q, Li Y, Lu Z, Niu L, Fan Z, Lei Z. Circular RNA circ_0124644 exacerbates the ox-LDL-induced endothelial injury in human vascular endothelial cells through regulating PAPP-A by acting as a sponge of miR-149-5p. *Mol Cell Biochem*. 2020 Aug;471(1-2):51-61. doi: 10.1007/s11010-020-03764-0.
91. Yu H, Pan Y, Dai M, Xu H, Li J. Circ_0003423 Alleviates ox-LDL-Induced Human Brain Microvascular Endothelial Cell Injury via the miR-589-5p/TET2 Network. *Neurochem Res*. 2021 Nov;46(11):2885-2896. doi: 10.1007/s11064-021-03387-x.
92. Zhu QQ, Pu XB, Chen TC, Qiu CY, Wu ZH, Tian L, He YY, Wang XH, Shang T, Wang X, Xiang YL, Li DL, Zhang HK. Hsa_circ_0008360 sponges miR-186-5p to target CCND2 to modulate high glucose-induced vascular endothelial dysfunction. *Cell Cycle*. 2021 Jul;20(14):1389-1401. doi: 10.1080/15384101.2021.1918877.
93. Huang HS, Huang XY, Yu HZ, Xue Y, Zhu PL. Circular RNA circ-RELL1 regulates inflammatory response by miR-6873-3p/MyD88/NF- κ B axis in endothelial cells. *Biochem Biophys Res Commun*. 2020 Apr 30;525(2):512-519. doi: 10.1016/j.bbrc.2020.02.109. Epub 2020 Feb 26. Erratum in: *Biochem Biophys Res Commun*. 2020 Nov 19;532(4):682.
94. Byron SA, Van Keuren-Jensen KR, Engelthaler DM, Carpten JD, Craig DW. Translating RNA sequencing into clinical diagnostics: opportunities and challenges. *Nat Rev Genet*. 2016 May;17(5):257-71. doi: 10.1038/nrg.2016.10.
95. Szabo L, Salzman J. Detecting circular RNAs: bioinformatic and experimental challenges. *Nat Rev Genet*. 2016 Oct 14;17(11):679-692. doi: 10.1038/nrg.2016.114.
96. Nielsen AF, Bindereif A, Bozzoni I, Hanan M, Hansen TB, Irimia M, Kadener S, Kristensen LS, Legnini I, Morlando M, Jarlstad Olesen MT, Pasterkamp RJ, Preibisch S, Rajewsky N, Suenkel C, Kjems J. Best practice standards for circular RNA research. *Nat Methods*. 2022 May 26. doi: 10.1038/s41592-022-01487-2.
97. Mannion N, Arieti F, Gallo A, Keegan LP, O'Connell MA. New Insights into the Biological Role of Mammalian ADARs; the RNA Editing Proteins. *Biomolecules*. 2015 Sep 30;5(4):2338-62. doi: 10.3390/biom5042338.
98. Hogg M, Paro S, Keegan LP, O'Connell MA. RNA editing by mammalian ADARs. *Adv Genet*. 2011;73:87-120. doi: 10.1016/B978-0-12-380860-8.00003-3..
99. Song C, Sakurai M, Shiromoto Y, Nishikura K. Functions of the RNA Editing Enzyme ADAR1 and Their Relevance to Human Diseases. *Genes (Basel)*. 2016 Dec 17;7(12):129. doi: 10.3390/genes7120129. PMID: 27999332
100. Cayir A. RNA A-to-I editing, environmental exposure, and human diseases. *Crit Rev Toxicol*. 2021 May;51(5):456-466. doi: 10.1080/10408444.2021.1953438
101. Chen Y, Peng H, Zhou S, Zhuang Y. ADAR1 is targeted by miR-143 to regulate IL-1 β -induced endothelial activation through the NF κ B pathway. *Int J Biochem Cell Biol*. 2017 Aug;89:25-33. doi: 10.1016/j.biocel.2017.05.021
102. van der Kwast RVCT, Parma L, van der Bent ML, van Ingen E, Baganha F, Peters HAB, Goossens EAC, Simons KH, Palmen M, de Vries MR, Quax PHA, Nossent AY. Adenosine-to-Inosine Editing of Vasoactive MicroRNAs Alters Their Targetome and Function in Ischemia. *Mol Ther Nucleic Acids*. 2020 Sep 4;21:932-953. doi: 10.1016/j.omtn.2020.07.020.
103. Vlachogiannis NI, Sachse M, Georgiopoulos G, Zormpas E, Bampatsias D, Delialis D, Bonini F, Galyfos G, Sigala F, Stamatiopoulos K, Gatsiou A, Stellos K. Adenosine-to-inosine Alu RNA editing controls the stability of the pro-inflammatory long noncoding RNA NEAT1 in atherosclerotic cardiovascular disease. *J Mol Cell Cardiol*. 2021 Nov;160:111-120. doi: 10.1016/j.yjmcc.2021.07.005.
104. Fei J, Cui XB, Wang JN, Dong K, Chen SY (2016) ADAR1-mediated RNA editing, a novel mechanism controlling phenotypic modulation of vascular smooth muscle cells. *Circ Res* 119: 463–469. 10.1161/CIRCRESAHA.116.309003
105. van der Kwast RVCT, van Ingen E, Parma L, Peters HAB, Quax PHA, Nossent AY. Adenosine-to-Inosine editing of MicroRNA-487b alters target gene selection after ischemia and promotes neovascularization. *Circ Res*, 2018, 122: 444–456.
106. Schirle NT, Goodman RA, Krishnamurthy M, Beal PA. Selective inhibition of ADAR2-catalyzed editing of the serotonin 2c receptor pre-mRNA by a helix-threading peptide. *Org Biomol Chem*. 2010 Nov 7;8(21):4898-904. doi: 10.1039/c0ob00309c. Epub 2010 Aug 31. PMID: 20820662

107. Montiel-González MF, Vallecillo-Viejo IC, Rosenthal JJ. An efficient system for selectively altering genetic information within mRNAs. *Nucleic Acids Res.* 2016 Dec 1;44(21):e157. doi: 10.1093/nar/gkw738. Epub 2016 Aug 23. PMID: 27557710; PMCID: PMC5137428.
108. Suzuki A, Tomita H, Okada H. Form follows function: The endothelial glycocalyx. *Transl Res.* 2022 Apr 11:S1931-5244(22)00069-X. doi: 10.1016/j.trsl.2022.03.014. Epub ahead of print. PMID: 35421613.
109. Hu Z, Cano I, D'Amore PA. Update on the Role of the Endothelial Glycocalyx in Angiogenesis and Vascular Inflammation. *Front Cell Dev Biol.* 2021 Aug 31;9:734276. doi: 10.3389/fcell.2021.734276. PMID: 34532323; PMCID: PMC8438194.
110. Gaudette S, Hughes D, Boller M. The endothelial glycocalyx: Structure and function in health and critical illness. *J Vet Emerg Crit Care (San Antonio).* 2020 Mar;30(2):117-134. doi: 10.1111/vec.12925. Epub 2020 Feb 18. PMID: 32067360.
111. Dogné S, Flamion B. Endothelial Glycocalyx Impairment in Disease: Focus on Hyaluronan Shedding. *Am J Pathol.* 2020 Apr;190(4):768-780. doi: 10.1016/j.ajpath.2019.11.016. Epub 2020 Feb 6. PMID: 32035885.
112. Duni A, Liakopoulos V, Koutlas V, Pappas C, Mitsis M, Dounousi E. The Endothelial Glycocalyx as a Target of Ischemia and Reperfusion Injury in Kidney Transplantation-Where Have We Gone So Far? *Int J Mol Sci.* 2021 Feb 22;22(4):2157. doi: 10.3390/ijms22042157. PMID: 33671524; PMCID: PMC7926299.
113. Milusev A, Rieben R, Sorvillo N. The Endothelial Glycocalyx: A Possible Therapeutic Target in Cardiovascular Disorders. *Front Cardiovasc Med.* 2022 May 13;9:897087. doi: 10.3389/fcvm.2022.897087. PMID: 35647072; PMCID: PMC9136230.
114. Salmon AHJ, Neal CR, Sage LM, Glass CA, Harper SJ, Bates DO. Angiopoietin-1 alters microvascular permeability coefficients in vivo via modification of endothelial glycocalyx. *Cardiovasc Res.* (2009) 83:24–33. doi: 10.1093/cvr/cvp093
115. Chappell D, Jacob M, Hofmann-Kiefer K, Bruegger D, Rehm M, Conzen P, et al. Hydrocortisone preserves the vascular barrier by protecting the endothelial glycocalyx. *Anesthesiology.* (2007) 107:776–84. doi: 10.1097/01.anes.0000286984.39328.96
116. Chappell D, Jacob M, Hofmann-Kiefer K, Rehm M, Welsch U, Conzen P, et al. Antithrombin reduces shedding of the endothelial glycocalyx following ischaemia/reperfusion. *Cardiovasc Res.* (2009) 83:388–96. doi: 10.1093/cvr/cvp097
117. Rubio-Gayosso I, Platts SH, Duling BR. Reactive oxygen species mediate modification of glycocalyx during ischemia-reperfusion injury. *Am J Physiol.* (2006) 290:H2247–56. doi: 10.1152/ajpheart.00796.2005
118. Torres Filho IP, Torres LN, Salgado C, Dubick MA. Novel adjunct drugs reverse endothelial glycocalyx damage after hemorrhagic shock in rats. *Shock.* (2017) 48:583–9. doi: 10.1097/SHK.0000000000000895
119. Kozar RA, Peng Z, Zhang R, Holcomb JB, Pati S, Park P, et al. Plasma restoration of endothelial glycocalyx in a rodent model of hemorrhagic shock. *Anesth Analg.* (2011) 112:1289–95. doi: 10.1213/ANE.0b013e318210385c
120. Lipowsky HH, Lescanic A. Inhibition of inflammation induced shedding of the endothelial glycocalyx with low molecular weight heparin. *Microvasc Res.* (2017) 112:72–8. doi: 10.1016/j.mvr.2017.03.007
121. Huang X, Han S, Liu X, Wang T, Xu H, Xia B, et al. Both UFH and NAH alleviate shedding of endothelial glycocalyx and coagulopathy in LPS-induced sepsis. *Exp Ther Med.* (2020) 19:913–22. doi: 10.3892/etm.2019.8285
122. Masola V, Onisto M, Zaza G, Lupo A, Gambaro G. A new mechanism of action of sulodexide in diabetic nephropathy: inhibits heparanase-1 and prevents FGF-2-induced renal epithelial-mesenchymal transition. *J Transl Med.* (2012) 10:213. doi: 10.1186/1479-5876-10-213
123. Zhao H, Zhu Y, Zhang J, Wu Y, Xiang X, Zhang Z, et al. The beneficial effect of HES on vascular permeability and its relationship with endothelial glycocalyx and intercellular junction after hemorrhagic shock. *Front Pharmacol.* (2020) 11:597. doi: 10.3389/fphar.2020.00597
124. Gumanova NG, Gorshkov AU, Klimushina MV, Kots AY. Associations of endothelial biomarkers, nitric oxide metabolites and endothelin, with blood pressure and coronary lesions depend on cardiovascular risk and sex to mark endothelial dysfunction on the SCORE scale. *Horm Mol Biol Clin Investig.* 2020 Nov 6;41(4). doi: 10.1515/hmbci-2020-0024.
125. Beckman JA, Hu JR, Huang S, Farber-Eger E, Wells QS, Wang TJ, Gerszten RE, Ferguson JF. Metabolomics reveals the impact of Type 2 diabetes on local muscle and vascular responses to ischemic stress. *Clin Sci (Lond).* 2020 Sep 18;134(17):2369-2379. doi: 10.1042/CS20191227.

126. Wang YF, Hsu YJ, Wu HF, Lee GL, Yang YS, Wu JY, Yet SF, Wu KK, Kuo CC. Endothelium-Derived 5-Methoxytryptophan Is a Circulating Anti-Inflammatory Molecule That Blocks Systemic Inflammation. *Circ Res*. 2016 Jul 8;119(2):222-36. doi: 10.1161/CIRCRESAHA.116.308559.
127. Adamczyk Z, Morga M, Nattich-Rak M, Sadowska M. Nanoparticle and bioparticle deposition kinetics. *Adv Colloid Interface Sci*. 2022 Apr;302:102630. doi: 10.1016/j.cis.2022.102630.
128. Chacko AM, Hood ED, Zern BJ, Muzykantov VR.. Targeted nano-carriers for imaging and therapy of vascular inflammation. *Curr Opin Colloid Interface Sci*. 2011. June; 16(3): 215– 27.
129. Wilhelm S, et al. Analysis of nanoparticle delivery to tumours. *Nat. Rev. Mater*. 2016;1:16014. doi: 10.1038/natrevmats.2016.14.
130. Yoo JW, Irvine DJ, Discher DE, Mitragotri S.. Bio-inspired, bio-engineered and biomimetic drug delivery carriers. *Nat Rev Drug Discov*. 2011. July 1; 10(7): 521– 35
131. Hu CM, Fang RH, Wang KC,. et al. Nanoparticle biointerfacing by platelet membrane cloaking. *Nature*. 2015. October 1; 526(7571): 118– 21.
132. Robbins GP, Saunders RL, Haun JB, Rawson J, Therien MJ, Hammer DA.. Tunable leuko-polymersomes that adhere specifically to inflammatory markers. *Langmuir*. 2010. September 7; 26(17): 14089– 96
133. Parodi A, Quattrocchi N, van de Ven AL, . et al. Synthetic nanoparticles functionalized with biomimetic leukocyte membranes possess cell-like functions. *Nat Nanotechnol*. 2013. January; 8(1): 61– 8.
134. McMahon KM, Mutharasan RK, Tripathy S, . et al. Biomimetic high density lipoprotein nanoparticles for nucleic acid delivery. *Nano Lett*. 2011. March 9; 11(3): 1208– 14.
135. Taneja, G.; Sud, A.; Pendse, N.; Panigrahi, B.; Kumar, A.; Sharma, A.K. Nano-medicine and vascular endothelial dysfunction: Options and delivery strategies. *Cardiovasc. Toxicol*. 2019, 19, 1–12
136. Quispe, C.; Cruz-Martins, N.; Manca M.L.; Manconi, M.; Sytar, O.; Hudz, N.; Shanaida, M.; Kumar, M.; Taheri, Y.; Martorell, M.; Sharifi-Rad, J.; Pintus, G.; Cho, W.C. Nano-Derived Therapeutic Formulations with Curcumin in Inflammation-Related Diseases. *Oxid Med Cell Longev*. 2021 Sep 15, 2021:3149223
137. Zhang, R.; Liu, R.; Liu, C.; Pan, L.; Qi, Y.; Cheng, J.; Guo, J.; Jia, Y.; Ding, J.; Zhang, J.; Hu H. A pH/ROS dual-responsive and targeting nanotherapy for vascular inflammatory diseases. *Biomaterials*. 2020 Feb, 230:119605.
138. Jeandet, P.; Sobarzo-Sánchez, E.; Uddin, M.S.; Bru, R.; Clément, C.; Jacquard, C.; Nabavi, S.F.; Khayat Kashani, M.; Batiha, G.E.; Khan, H.; Morkunas, I.; Trotta, F.; Matencio, A.; Nabavi, S.M. Resveratrol and cyclodextrins, an easy alliance: Applications in nanomedicine, green chemistry and biotechnology. *Biotechnol Adv*. 2021 Oct 7, 53:107844
139. Cho, B. H., Park, J. R., Nakamura, M. T., Odintsov, B. M., Wallig, M. A., & Chung, B. H. (2010). Synthetic dimyristoylphosphatidylcholine liposomes assimilating into high-density lipoprotein promote regression of atherosclerotic lesions in cholesterol-fed rabbits. *Experimental Biology and Medicine (Maywood)*, 235, 1194–1203
140. Pillai SC, Borah A, Jacob EM, Kumar DS. Nanotechnological approach to delivering nutraceuticals as promising drug candidates for the treatment of atherosclerosis. *Drug Deliv*. 2021 Dec;28(1):550-568. doi: 10.1080/10717544.2021.1892241. PMID: 33703990; PMCID: PMC7954496.
141. Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini, L.A., et al. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science* 339, 819–823
142. Doudna, J.A., and Charpentier, E. (2014). Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science* 346, 1258096.
143. Cox, D.B., Platt, R.J., and Zhang, F. (2015). Therapeutic genome editing: prospects and challenges. *Nat. Med.* 21, 121–131.
144. Varshney, G.K., Pei, W., LaFave, M.C., Idol, J., Xu, L., Gallardo, V., Carrington, B., Bishop, K., Jones, M., Li, M., et al. (2015). High-throughput gene targeting and phenotyping in zebrafish using CRISPR/Cas9. *Genome Res*. 25, 1030–1042.
145. Zhang X, Jin H, Huang X, Chaurasiya B, Dong D, Shanley TP, Zhao YY. Robust genome editing in adult vascular endothelium by nanoparticle delivery of CRISPR-Cas9 plasmid DNA. *Cell Rep*. 2022 Jan 4;38(1):110196. doi: 10.1016/j.celrep.2021.110196.

146. Botto C, Dalkara D, El-Amraoui A. Progress in Gene Editing Tools and Their Potential for Correcting Mutations Underlying Hearing and Vision Loss. *Front Genome Ed.* 2021 Oct 28;3:737632. doi: 10.3389/fgeed.2021.737632. PMID: 34778871; PMCID: PMC8581640.