

Article

Evaluating the sensitivity of heat wave definitions among North Carolina physiographic regions

Jagadeesh Puvvula ¹, Azar M. Abadi ¹, Kathryn C. Conlon ², Jared J. Rennie ³, Hunter Jones⁴, and Jesse E. Bell ^{1,5,6*}

¹ Department of Environmental, Agricultural and Occupational Health, College of Public Health, University of Nebraska Medical Center, Omaha, NE, 68198

² Department of Public Health Sciences, University of California Davis, One Shields Ave, Davis, CA, 95616

³ NOAA/National Centers for Environmental Information, Asheville, NC, 28801

⁴ Medical Sciences Interdepartmental Area, Office of Graduate Studies, University of Nebraska Medical Center, NE, USA

⁵ School of Natural Resources, University of Nebraska-Lincoln, Lincoln, NE, 68583

⁶ Daugherty Water for Food Global Institute, University of Nebraska, Lincoln, NE, 68583

* Correspondence: jesse.bell@unmc.edu; 402-552-7237

Abstract: Exposure to extreme heat is a known risk factor that is associated with increased heat-related illness (HRI) outcomes. The relevance of heat wave definitions could change across the health conditions and geographies due to the heterogeneous climate profile. This study compared the sensitivity of 28 heat wave definitions associated with HRI emergency department visits over five summer seasons (2011-2016), stratified by two physiographic regions (Coastal and Piedmont) in North Carolina. The HRI rate ratios associated with heat waves were estimated using the generalized linear regression framework assuming a negative binomial distribution. We compared the Akaike Information Criterion (AIC) values across the heat wave definitions to identify an optimal heat wave definition. In the Coastal region, heat wave definition based on daily maximum temperature with a threshold >90th percentile for two or more consecutive days had the optimal model fit. In the Piedmont region, heat wave definition based on the daily minimum temperature with a threshold value >90th percentile for two or more consecutive days was optimal. Additionally, we observed that the optimal heat wave definitions from this study captured moderate and frequent heat episodes than the national weather service (NWS) heat products that worked best for extreme heat episodes. This study compared the HRI morbidity risk associated with epidemiologic-based heat wave definitions and with NWS heat products. Our findings could be used for public health education and suggest recalibrating NWS heat products.

Keywords: Heat Wave; Heat-related illness; Early heat-health warning systems.

1. Introduction

A heat wave is often described as an acute episode of one or more consecutive days with temperatures or heat indices exceeding a threshold value [1]. However, there is no standard definition to identify heat waves [2]. Heat waves are typically classified using a synoptic (e.g., air mass, temperature-humidity index), physiologic (e.g., Environmental Stress Index, Wet Bulb Global Temperature), or epidemiologic approach [3]. Hajat et al. (2010) reported that epidemiologic-based algorithms (temperature-mortality relationship) identified the days with higher heat-related mortality.

Heat waves are associated with an increased risk of HRI outcomes [4]. In the United States (US), roughly 700 heat-related deaths per year are attributable to ambient temperature exposure [5]. The frequency and intensity of heat waves have been on the rise since the industrial revolution and are likely to increase in the future due to climate change [6, 7]. In Philadelphia, heat-related risk communication, along with the NWS warnings, played a crucial role in minimizing up to 3 heat-related deaths per day that are associated with extreme heat exposure [8, 9]. In the US, public health departments rely on heat

products (e.g., excessive heat watch, heat advisory, and excessive heat warning) from the local National Weather Service Weather Forecast Office (NWS-WFO) to communicate heat-health risks. According to Weinberger et al. (2018), the NWS heat products moderately reduced the impact of extreme heat on human health, but the human health risk associated with ambient temperature is not a resolved issue [10].

Despite the early heat warnings, the average annual percentage of HRI emergency department visits in North Carolina has increased by 19% over the past decade [11, 12]. The rise in HRI risk could be due to population vulnerabilities (age, economic status, and occupation) and higher thresholds for temperature and humidity set by NWS for issuing heat alerts [13]. Multiple researchers attempted to characterize human health risks associated with heat wave definitions [3, 14-17]. Epidemiologic studies evaluating the association between outdoor temperature exposure (e.g., heat waves or ambient temperature) and human health are generally focused on extreme events, where the human health outcome is typically measured as cause-specific or all-cause mortality [17, 18]. A meta-analysis on the heat wave definition evaluation studies summarized that the studies included in this review generalized the warnings to a larger geographic area, such as a state or a group of states [17]. Generalizing heat warning systems over larger geography may not be ideal due to heterogeneity in exposures, population vulnerability, and exposure-outcome associations [19]. Physiographic or sub-regional scale heat warning systems that account for meteorological heterogeneity and specific to health conditions were found to play a role in minimizing the human health risks associated with heat waves [8, 16, 20, 21].

This study assessed the association between frequent and moderate heat waves and HRI emergencies in North Carolina physiographic regions. The objective of this study is to compare the statistical model performance of heat wave definitions and assess their association with HRI emergency visits in North Carolina. Additionally, we aim to compare the heat wave definition with the best model performance from our study with the NWS extreme heat alerts.

2. Materials and Methods

This study is focused on five summer seasons (May 01 – September 30) from 2011 to 2016 among the three physiographic regions (Coastal, Piedmont & Mountain) in North Carolina. The year 2013 was excluded in this study due to data availability constraints.

2.1. Data

2.1.1. Heat metrics

Daily mean, minimum, and maximum temperatures were obtained from the Global Historical Climate Network – Daily (GHCN-D) database [7, 22]. Dew point data was obtained from the Parameter-elevation Regression on Independent Slopes Model (PRISM) database [23]. The station-based temperature measurements and gridded dew point data were aggregated by physiographic regions. The daily maximum apparent temperature and relative humidity were estimated using daily maximum temperature and dew point using `heat.index.function` and `dewpoint.to.humidity` functions available from the `weathermetrics` package in R [24].

2.1.2. Heat Wave definitions

Twenty-eight heat wave definitions (Table 1) associated with human health outcomes were adopted from the existing literature and were included in this study [14, 15, 25, 26]. These heat wave definitions were classified based on four factors: 1) a heat metric (daily mean, minimum, maximum, and apparent temperatures), 2) duration (number of days), 3) threshold type (relative/absolute), and 4) threshold intensity. Among the two threshold types, relative threshold-based definitions account for cumulative heat exposure (2+ or 3+ consecutive days), and definitions based on absolute threshold are based on single heat day exposure. The heat wave definitions based on daily temperature as metric and a relative threshold were classified using four percentile values (99, 98, 95, 90) as

threshold intensity. The percentile threshold values were calculated using historical observations over the summer season from 1895-2016. The definitions based on the apparent temperature as metric and a relative threshold were classified using three percentile threshold values (95, 90, 85) as threshold intensity.

Among the 28 heat wave definitions (HW_01 to HW_28) included in this study, 24 are based on relative threshold values, and four heat wave definitions are based on the absolute threshold value. Three of the 27 heat wave definitions using relative threshold values were based on maximum apparent temperature as the heat metric. The remaining 24 heat wave definitions were based on the daily mean (HW_01-HW_08), maximum (HW_09-HW_16), and minimum (HW_18-HW_25) temperature values as the heat metric. Additionally, one heat wave definition using absolute threshold value is based on daily maximum temperature as the heat metric (Table 1). Using the 28 heat wave definitions, we categorized the summer days during the study period as heat waves and non-heat wave days using a binary variable to indicate heat waves.

2.1.3. National Weather Service – Heat products

The heat products released by the NWS during the study period were retrieved from the Iowa Environmental Mesonet [27]. During the study period, the NWS heat products (heat advisories and excessive heat warnings) were released by the three NWS-WFOs (ILM-Wilmington; MHX-Newport/Morehead city; and RAH-Raleigh) located in North Carolina were included in this study. The WFOs ILM and MHX cover most of the Coastal region, and RAH covers the Piedmont region [28]. Among the three WFOs, heat products released by the ILM and MHX follow the NWS procedural directive. The RAH WFO is the only center in North Carolina that collaborated with health partners to revise the heat products [29]. The heat products from RAH are based on local conditions such as maximum temperature, sunlight, nighttime temperature, heterogeneity between rural and urban temperatures, and knowledge from historical weather conditions [29]. The heat products used in the three WFOs in North Carolina are based on the following criteria. A heat advisory is released during the days when the daytime heat index value is between 100-105°F [30]. An excessive heat warning is released if the daytime heat index forecast value is between 105-110°F [30].

We extracted the start and end dates and County information from the Iowa Environmental Mesonet heat product archives. The County information was aggregated to the North Carolina physiographic region scale to match the spatial resolution of the health data included in this study. The NWS heat products were represented using a binary variable (NWS_HW) to identify the days with NWS alerts on a daily scale by physiographic region.

2.1.4. Heat-related illness

Daily HRI-related emergency department visit data were obtained as an aggregate count per day per physiographic region from the North Carolina Disease Event Tracking and Epidemiologic Collection Tool (NC DETECT) surveillance program maintained by the North Carolina Division of Public Health (NC DPH) [31]. Heat-related illnesses were defined using ICD-9 CM codes with E992/E900.0/E900.0/E900; ICD-10 CM codes within T67/X30/X32; and various keywords from the chief complaint/triage notes [31, 32]. The days with HRI emergency department visits fewer than five were censored, amounting to 28.81% (219) of observations from the Coastal and 28.94% (220) in the Piedmont region. The days with censored HRI emergency visits in the Coastal and Piedmont regions were imputed by the median value of 3 visits per day. The Mountain region was excluded from the analysis as 50.13% of the data was censored due to low HRI emergencies.

2.2. Statistical analysis

The sensitivity of 28 heat wave definitions was compared using the Akaike Information Criterion (AIC) value corresponding to the model fit [33, 34] evaluating the HRI morbidity rate associated with heatwaves included in this study. AIC is a metric that is a balance between model accuracy and penalty due to complexity, commonly used to

measure the optimal model fit (equation 1) [35]. Smaller the AIC value (close to $-\infty$) represent an optimal fit [36].

$$AIC = \text{goodness of fit} + \text{penalty} \dots (\text{equation 1})$$

The HRI rate ratios corresponding to the 28 heat wave definitions included in this study were estimated using the Generalized Linear Model (GLM) and assuming negative binomial distribution to account for outcome overdispersion. To compare the HRI risk across physiographic regions, the regression model was adjusted for population density by using the 2010 decennial population by region as an offset term [37]. To estimate the direct effect of heat wave definitions, the statistical models using heat wave definitions based on temperature as a heat metric were adjusted for relative humidity, and NWS heat wave alert days were added as covariates to adjust for potential confounding effects and effect modification. Similarly, the statistical models with heat wave definitions using apparent temperature as a heat metric were adjusted for NWS heat alerts. Additionally, to account for temporal autocorrelation, we adjusted the statistical models mentioned above for the day of the week (weekday/weekend (binary)), month (factor), and year (factor) (Figure 1 and equation 2). In equation 2., HW is a binary variable that represents heat wave definitions, RH represents relative humidity, NWS-HW represents NWS heat wave alerts, and TS represents the time series variables (day of week, month, and year).

$$\log\left(\frac{E(\text{HRI count})}{\text{population}}\right) = \beta_0 + \beta_{HW} + \beta_{RH} + \beta_{NWS_HW} + \beta_{TS} + \varepsilon \dots (\text{equation 2})$$

We processed 28 statistical models stratified by physiographic region to obtain the rate ratio (RR) and 95% confidence intervals. The sensitivity of heat wave definitions was evaluated by comparing the AIC values across heat wave definitions by physiographic region that were generated from the GLM output. The statistical model with the lowest AIC value among the 28 heat wave definitions was considered the optimal heat wave definition. We then compared the overlap between the days considered as heat waves from this study and the NWS heat products using the Chi-Square test [38]. The analysis was conducted using R version 4.0.3 and MASS package version 7.3 [39].

3. Results

3.1. Heat wave definition – Sensitivity

Among the 28-heat metrics included in this study, the heat wave definition using maximum temperature had the best fit with HRI morbidity in the Coastal region and mean temperature for the Piedmont region. The heat wave definitions based on a moderate (90th) percentile threshold for Coastal and Piedmont regions had an optimal model fit than the heat wave definitions based on extreme threshold (99th, 98th, and 95th percentile) values.

In the Coastal region, the heat wave definition based on daily maximum temperature as a heat metric with a threshold value > 90th percentile for two or more consecutive days (HW_15) had the optimal model fit (lowest AIC value) to estimate the HRI morbidity compared to the heat wave definitions included in this study. We did not observe a similar result for the Piedmont region. In the Piedmont region, the heat wave definition based on daily mean temperature as a heat metric with a threshold value > 90th percentile for two or more consecutive days (HW_07) had the optimal model fit to estimate the HRI morbidity. In the Coastal region, the heat wave definition HW_15 is associated with 2.75 (95% CI 2.40-3.08) times higher HRI morbidity rate during heat wave days than the non-heat wave days. In the Piedmont region, the heat wave definition HW_07 is associated with a 2.72 (95% CI 2.46-3.01) times higher HRI morbidity rate compared to the non-heat wave days (Figure 2).

Using the HW_15 definition, 27% (190/704) of the days in the Coastal and using the HW_07 heat wave definition, 34% (241/719) of the days in the Piedmont region were

flagged as heat wave days during the study period (Table 2). There are an average of 6 HRI ED visits per day in the Coastal region during the heat wave days based on HW_15 and an average of 8 HRI ED visits per day during the heat wave days based on HW_07 in the Piedmont region (Table 2). The frequency of heat wave days in the Piedmont region using the HW_07 definition was 24% higher than in the Coastal region using the HW_15 definition. About 72% of the heat wave days from the Coastal region matched with the Piedmont region. During the study period, we observed a lower number of heat wave days during the summer of 2014. The frequency of heat wave days was higher in July than in other summer months, based on the epidemiologic relationship-based heat wave definition (Coastal: HW_15; Piedmont: HW_07).

3.2. Comparing epidemiologic-based heat wave definition and NWS heat products

During the study period, NWS flagged 26 days in the Coastal and 18 days in the Piedmont regions as heat waves. The NWS heat wave days overlapped with the optimal heat wave definition identified in this study (HW_15 for Coastal and HW_07 for Piedmont). In the Coastal region, there were a significantly higher (6-times) number of heat wave days based on HW_15 than the NWS heat alerts (McNemar $\chi^2=158.15$, $df=1$; $P<0.05$). Similarly, the Piedmont region had a significantly higher (13 times) number of heat waves based on HW_07 than the NWS heat alerts (McNemar $\chi^2=219.04$, $df=1$, $P<0.05$) (Figure. 3).

3.2. Figures, Tables, and Schemes




Figure 1. Conceptualization of evaluating the direct effect of temperature or apparent temperature on HRI ED visits. We assumed that the association between temperature and HRI is mediated through NWS heat products. Additionally, relative humidity is influenced by temperature. To evaluate the association between heat wave definitions based on temperature and HRI, we adjusted for relative humidity and NWS heat products. We adjusted for NWS heat products while evaluating the association between heat wave definitions based on apparent temperature and HRI ED visits.




Figure 2. HRI rate ratio corresponding to heat wave definitions. The rate ratios and the corresponding 95% confidence intervals were generated using the generalized linear model (GLM), assuming a negative binomial distribution. The X-axis represents distinct heat wave definitions, stratified by North Carolina physiographic regions, and grouped by metric and threshold type. The Y-axis represents the HRI morbidity rate ratio, which could be interpreted as an increase/decrease in HRI morbidity rate during a heat wave day compared to a non-heat wave day.




Figure 3. Comparison of the effective heat wave definition with the NWS-Heat alerts. The X-axis represents daily temperature in degrees Celsius (Coastal: Maximum temperature & Piedmont: Minimum temperature). The Y-axis represents the daily rate of heat-related illness morbidity per 100,000 population. Each dot represents an observation corresponding to daily temperature and the rate of HRI morbidity during the study period. Panel A represents the Coastal region, and panel B represents the Piedmont region. The dots in the scatter plot are color-coded with three possible combinations: 1. Red: Categorized as heat wave day from our results (HW_15 or HW_07) and the NWS; 2. Blue: Categorized as heat wave day only based on our result; 3. Beige: Not flagged as a heat wave day from our results nor the NWS.

Table 1. Description of the heat wave definitions

Defini- tion	Heat met- ric	Duration (No. of days)	Thresh- old type	Threshold intensity	Coastal			Piedmont		
					Thresh- old (°C)	HW days	AIC	Thresh- old (°C)	HW days	AIC
HW_01	Mean temperature	2+consecutive	Relative	>99th percentile	29.16	33	4475.7	28.40	29	4817.9
HW_02		3+consecutive	Relative	>99th percentile		25	4480.7		21	4843.0
HW_03		2+consecutive	Relative	>98th percentile	28.57	58	4362.9	27.66	61	4778.4
HW_04		3+consecutive	Relative	>98th percentile		46	4396.7		49	4796.9
HW_05		2+consecutive	Relative	>95th percentile	27.65	105	4287.3	26.70	136	4629.7
HW_06		3+consecutive	Relative	>95th percentile		97	4293.5		118	4656.9
HW_07		2+consecutive	Relative	>90th percentile	26.65	217	4216.2	25.63	241	4547.9
HW_08		3+consecutive	Relative	>90th percentile		195	4211.8		227	4551.8
HW_09	Maximum temperature	2+consecutive	Relative	>99th percentile	35.04	22	4475.4	35.52	20	4842.6
HW_10		3+consecutive	Relative	>99th percentile		16	4479.7		14	4852.4
HW_11		2+consecutive	Relative	>98th percentile	34.31	38	4413.7	34.69	38	4799.8
HW_12		3+consecutive	Relative	>98th percentile		30	4435.3		32	4813.2
HW_13		2+consecutive	Relative	>95th percentile	33.13	98	4283.5	33.30	109	4665.9
HW_14		3+consecutive	Relative	>95th percentile		80	4318.0		85	4716.0
HW_15		2+consecutive	Relative	>90th percentile	31.97	190	4192.6	31.94	194	4583.4
HW_16		3+consecutive	Relative	>90th percentile		168	4234.4		180	4615.9
HW_17		1-day	Absolute	> 35°C	35.00	26	4423.1	35.00	34	4801
HW_18	Minimum temperature	2+consecutive	Relative	>99th percentile	23.86	29	4488.1	21.86	45	4810.3
HW_19		3+consecutive	Relative	>99th percentile		25	4479.1		31	4831.9
HW_20		2+consecutive	Relative	>98th percentile	23.36	56	4401.1	21.40	83	4783.6
HW_21		3+consecutive	Relative	>98th percentile		46	4433.5		65	4799.5
HW_22		2+consecutive	Relative	>95th percentile	22.57	108	4352.0	20.63	156	4722.0
HW_23		3+consecutive	Relative	>95th percentile		94	4353.0		136	4722.6
HW_24		2+consecutive	Relative	>90th percentile	21.64	223	4305.5	19.72	265	4670.3
HW_25		3+consecutive	Relative	>90th percentile		199	4303.0		235	4693.7
HW_26	Maximum apparent temperature	1-day	Absolute	>95th percentile	37.21	36	4254.0	35.26	27	4659.0
HW_27		1-day	Absolute	>90th percentile	36.20	71	4319.9		58	4749.8
HW_28		1-day	Absolute	>85th	35.47	106	4415.7		98	4799.1

HW (Heat wave) days per Coastal and Piedmont region represent the cumulative number of days during the study period that are categorized as heat wave days corresponding to the heat wave definitions.

Table 2. Frequency of heat wave days and HRI ED visits in North Carolina physiographic regions

		Month									
		May		Jun		Jul		Aug		Sep	
		a	b	a	b	a	b	a	b	a	b
2011	HW	3	1	21	14	22	25	10	15	0	0
	ED	65	37	309	334	225	417	195	302	0	0
2012	HW	0	0	5	6	23	27	0	8	2	6
	ED	0	0	45	95	306	520	0	112	35	52
2014	HW	0	0	6	7	7	11	0	5	5	6
	ED	0	0	101	141	103	149	0	55	68	92
2015	HW	0	0	15	16	14	19	5	9	2	3
	ED	0	0	817	897	306	569	117	177	25	54
2016	HW	0	0	5	8	24	26	19	24	2	5
	ED	0	0	80	204	721	950	366	499	30	106

a-Coastal; b-Piedmont; HW- number of heat wave days using the definition from this study [Coastal: HW_15; Piedmont: HW_07] and excluding the days that overlapped with the heat wave days flagged by the NWS; ED-number of HRI emergency department visits corresponding to the heat wave days.

4. Discussion

This study compared the optimal model fit using AIC values across the 28 heat wave definitions. Additionally, compared the heat wave days flagged using the optimal definition identified from this study to the NWS heat products. We observed that the heat wave definition based on the maximum temperature (HW_15) had an optimal performance for the Coastal region and the mean temperature-based heat wave definition (HW_07) for the Piedmont region. The heat wave definitions mentioned above were associated with 2.75 (95% CI 2.40-3.08) times higher rate of HRI morbidity in the Coastal region and 2.72 (95% CI 2.46-3.01) times higher HRI morbidity rate in Piedmont than the non-heat wave days. During the study period, our results suggest an excess of 33 heat wave days per summer season in the Coastal and 45 in the Piedmont region based on the heat wave definitions HW_15 for Coastal and HW_07 for Piedmont. During the summer, most days in July were flagged as vulnerable to heat-related emergencies while using the heat wave definitions from this study.

Heterogeneity while evaluating the heat wave definitions across the US climate regions or sub-regions was well established in the literature [15, 16, 21]. To address the meteorological heterogeneity within North Carolina, we evaluated the heat waves stratified

by physiographic delineations with clustering the administrative boundaries into physiographic regions. In North Carolina, there are two Weather Forecast Offices (WFOs) covering the Coastal region and a WFO covering the Piedmont region. The WFO covering the Piedmont region collaborated with the regional health partners to revise the heat thresholds. The WFO that overlaps the Piedmont region considered revising its heat threshold based on the local conditions for optimization [29]. Independent WFOs setting heat product thresholds at a sub-regional scale would be beneficial for the climate-related heterogeneity across the administrative boundaries. However, a study discussing the heat products across the US reported that the three WFOs across the Coastal and Piedmont regions follow the same threshold value and criteria to release heat products [40]. In contrast to the homogenous heat product thresholds across the physiographic regions, our results suggest heterogeneity of the heat wave definitions between the Coastal and Piedmont regions.

The results from our study overlapped with the observations from a previous study that compared the sensitivity of heat wave definitions across San Diego climate zones, using heat-related hospitalizations as an outcome [21]. McElroy et al. (2020) reported using daily maximum temperature above 90th percentile (29.11°C) in a day as a criterion for heat wave definitions in the Coastal region to be most efficient, using heat-related hospitalizations as an outcome. In this study, we observed that the heat wave definition based on daily maximum temperature above 90th percentile (31.97°C) for two or more days had an optimal fit with HRI morbidity in the Coastal region. Additionally, we observed that the heat wave definition based on daily mean temperature was optimal for the Piedmont region. In contrast, McElroy et al. (2020) reported that heat wave definitions based on daily maximum and the minimum temperature had the most impact in the Inland and Desert regions of San Diego. The major difference between our findings and McElroy et al. (2020) is focused on the duration criterion for defining heat waves. We observed that the heat wave definitions using two or more days as a duration criterion had an optimal fit, compared to McElroy et al. (2020) reported absolute thresholds were most efficient.

Early heat health warning systems play a crucial role in systematically minimizing the risks associated with outdoor temperature exposure [8]. Multiple studies attempted to characterize a gold standard heat wave definition, where most of these studies compared the sensitivity of the heat wave definitions in the context of mortality [14, 15, 17]. Vaidyanathan et al. (2016) evaluated the sensitivity of several heat wave definitions and their association with heat-related deaths by comparing the effect estimates (extreme heat effect). Similarly, Anderson and Bell (2009) evaluated the sensitivity of heat wave definitions based on the percent increase in relative risk associated with heat-related mortality. McElroy et al. (2020) evaluated heat waves by climate zones, using the attributable risk associated with heat-related hospitalizations. These studies assessed the optimal heat wave definition by comparing the effect estimates/relative or attributable risks associated with health outcomes. We compared and identified an optimal heat wave definition per North Carolina physiographic regions using the model fit metric – lowest AIC value (model fit) instead of the effect estimates (strength of association).

Using AIC as a metric, we compared 28 heat wave definitions and identified a heat wave definition with an optimal model fit. The heat wave definition (HW_15) using daily maximum temperature with a percentile value > 90th percentile for two or more consecutive days had an optimal fit for the Coastal region. Similarly, using daily mean temperature with a threshold value > 90th percentile for two or more consecutive days was optimal for the Piedmont region compared to the heat wave definitions included in this study. During the study period, 27% of the summer days in the Coastal and 34% in the Piedmont region were flagged as vulnerable to HRI emergencies by our definition. In contrast, the NWS released heat products during ~2.5% of the summer days. During the study period, there were an average of 783 (6 per day) HRI emergency visits per summer season during the days flagged as vulnerable based on the HW_15 definition in the Coastal region and 1,152 (8 per day) HRI emergencies per summer season in the Piedmont region during the days flagged as vulnerable to heat-related emergencies using the HW_07 definition.

Abasilim and Friedman (2021) reported about 16 heat-related hospitalizations per day during the summer days without NWS excess heat warnings in Illinois [41].

Our results could be influenced by the interaction between vulnerability factors and risk perception. Additionally, our results are subjective to a variety of unmeasured biases driven by human vulnerabilities such as co-existing medical conditions, occupational vulnerabilities, demographics (age, gender, race, education, urbanicity), and socioeconomic factors (wealth, employment, housing) that were identified to exacerbate the risk of heat-related illnesses [42-48]. Additionally, our results could be influenced by effect modifiers such as human behavioral factors that include knowledge on heat risk sensitivity, external locus of control, and emotional and cognitive factors that heavily alter the risk perception of heat warnings [49-51]. Further studies evaluating heat wave definitions using mixed methods by considering quantitative information from human vulnerability characteristics and qualitative information from heat risk perception could strengthen the heat health risk ascertainment.

5. Conclusions

Our results showed heterogeneity of the optimal heat wave definitions among the Coastal and Piedmont regions in North Carolina. Additionally, the threshold values associated with the optimal heat wave definitions were smaller compared to the NWS thresholds for the North Carolina physiographic regions. Our results suggest recalibrating the heat wave definitions used by the NWS WFOs in North Carolina.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1.

Author Contributions: Conceptualization, J.P., and J.E.B.; methodology, J.P., and J.E.B.; software, J.P.; validation, J.E.B., and K.C.C.; formal analysis, J.P.; resources, J.R., A.M.A., and J.E.B.; data curation, J.R., A.M.A., and J.P.; writing—original draft preparation, J.P.; writing—review and editing, J.P., J.E.B., K.C.C., A.M.A., and H.J.; visualization, J.P.; supervision, J.E.B.; project administration, J.E.B.; funding acquisition, J.E.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Society of Actuaries.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and Ethical review and approval were waived for this study, due to the REASON the use of data aggregation at the physiographic level.

Informed Consent Statement: Not applicable.

Data Availability Statement: Temperature data is available the Global Historical Climatology Network daily (GHCNd) database, [Dataset]. Available at: <https://www.ncei.noaa.gov/products/land-based-station/global-historical-climatology-network-daily>; Precipitation and dew point data is available at the PRISM Climate Group database, [Dataset]. Available at: <https://www.prism.oregonstate.edu/re-cent/>; Heat-related emergency department visit data could be requested from the North Carolina Department of Health and Human Services, Available at: <https://ncdetect.org/data/>.

Acknowledgments: We would like to acknowledge the Society of Actuaries, Climate Central, Claire M. Hubbard Foundation and the Robert B. Daugherty Water for Food Global Institute at the University of Nebraska for supporting this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Mazdiyasni, O.; AghaKouchak, A., Substantial increase in concurrent droughts and heatwaves in the United States. *Proceedings of the National Academy of Sciences* **2015**, 112, (37), 11484-11489.
2. USGCRP, The impacts of climate change on human health in the United States : A scientific assessment. In U.S Global Change Research Program: Washington, D.C., 2016.
3. Hajat, S.; Sheridan, S. C.; Allen, M. J.; Pascal, M.; Laaidi, K.; Yagouti, A.; Bickis, U.; Tobias, A.; Bourque, D.; Armstrong, B. G.; Kosatsky, T., Heat-health warning systems: a comparison of the predictive capacity of different approaches to identifying dangerously hot days. *Am J Public Health* **2010**, 100, (6), 1137-44.
4. Knowlton, K.; Rotkin-Ellman, M.; King, G.; Margolis, H. G.; Smith, D.; Solomon, G.; Trent, R.; English, P., The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* **2009**, 117, (1), 61-7.
5. Vaidyanathan, A.; Malilay, J.; Schramm, P.; Saha, S., Heat-Related Deaths - United States, 2004-2018. *MMWR Morb Mortal Wkly Rep* **2020**, 69, (24), 729-734.
6. IPCC *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*; Cambridge University Press, 2021.
7. Rennie, J.; Bell, J. E.; Kunkel, K. E.; Herring, S.; Cullen, H.; Abadi, A. M., Development of a Submonthly Temperature Product to Monitor Near-Real-Time Climate Conditions and Assess Long-Term Heat Events in the United States. *Journal of Applied Meteorology and Climatology* **2019**, 58, (12), 2653-2674.
8. Ebi, K. L.; Teisberg, T. J.; Kalkstein, L. S.; Robinson, L.; Weiher, R. F., Heat Watch/Warning Systems Save Lives: Estimated Costs and Benefits for Philadelphia 1995–98. *Bulletin of the American Meteorological Society* **2004**, 85, (8), 1067-1074.
9. Kalkstein, S. L., Jamason, F.P., Greene, S.J., Philadelphia Hot Weather Health Watch Warning System. <https://www1.udel.edu/SynClim/phl.html>
10. Weinberger, K. R.; Zanutti, A.; Schwartz, J.; Wellenius, G. A., Effectiveness of National Weather Service heat alerts in preventing mortality in 20 US cities. *Environ Int* **2018**, 116, 30-38.
11. CDC, National Environmental Public Health Tracking Network. In 2020.
12. NCDHHS Heat data and prevention. <https://publichealth.nc.gov/chronicdiseaseandinjury/heat.htm> (Sep 11, 2020),
13. Wellenius, G. A.; Eliot, M. N.; Bush, K. F.; Holt, D.; Lincoln, R. A.; Smith, A. E.; Gold, J., Heat-related morbidity and mortality in New England: Evidence for local policy. *Environ Res* **2017**, 156, 845-853.
14. Anderson, G. B.; Bell, M. L., Heat waves in the United States: mortality risk during heat waves and effect modification by heat wave characteristics in 43 U.S. communities. *Environ Health Perspect* **2011**, 119, (2), 210-8.
15. Vaidyanathan, A.; Kegler, S. R.; Saha, S. S.; Mulholland, J. A., A Statistical Framework to Evaluate Extreme Weather Definitions from A Health Perspective: A Demonstration Based on Extreme Heat Events. *Bull Am Meteorol Soc* **2016**, 97, (10), 1817-1830.
16. Vaidyanathan, A.; Saha, S.; Vicedo-Cabrera, A. M.; Gasparrini, A.; Abdurehman, N.; Jordan, R.; Hawkins, M.; Hess, J.; Elixhauser, A., Assessment of extreme heat and hospitalizations to inform early warning systems. *Proc Natl Acad Sci U S A* **2019**, 116, (12), 5420-5427.
17. Xu, Z.; Fitzgerald, G.; Guo, Y.; Jalaludin, B.; Tong, S., Impact of heatwave on mortality under different heatwave definitions: A systematic review and meta-analysis. *Environ Int* **2016**, 89-90, 193-203.
18. Song, X.; Wang, S.; Hu, Y.; Yue, M.; Zhang, T.; Liu, Y.; Tian, J.; Shang, K., Impact of ambient temperature on morbidity and mortality: An overview of reviews. *Sci Total Environ* **2017**, 586, 241-254.
19. Reid, C. E.; O'Neill, M. S.; Gronlund, C. J.; Brines, S. J.; Brown, D. G.; Diez-Roux, A. V.; Schwartz, J., Mapping community determinants of heat vulnerability. *Environ Health Perspect* **2009**, 117, (11), 1730-6.
20. Fechter-Leggett, E. D.; Vaidyanathan, A.; Choudhary, E., Heat Stress Illness Emergency Department Visits in National Environmental Public Health Tracking States, 2005-2010. *J Community Health* **2016**, 41, (1), 57-69.

21. McElroy, S.; Schwarz, L.; Green, H.; Corcos, I.; Guirguis, K.; Gershunov, A.; Benmarhnia, T., Defining heat waves and extreme heat events using sub-regional meteorological data to maximize benefits of early warning systems to population health. *Sci Total Environ* **2020**, 721, 137678.
22. NCEI, Global Historical Climatology Network daily (GHCNd). In 2019.
23. PRISM, Parameter-elevation Regression on Independent Slopes Model. In PRISM Climate Group: 2019.
24. Anderson, B.; Peng, R.; Ferreri, J. *Functions to Convert Between Weather Metrics*, 1.2.2; CRAN: 2016.
25. Smith, T. T.; Zaitchik, B. F.; Gohlke, J. M., Heat waves in the United States: definitions, patterns and trends. *Clim Change* **2013**, 118, (3-4), 811-825.
26. Kent, S. T.; McClure, L. A.; Zaitchik, B. F.; Smith, T. T.; Gohlke, J. M., Heat waves and health outcomes in Alabama (USA): the importance of heat wave definition. *Environ Health Perspect* **2014**, 122, (2), 151-8.
27. University, I. S., Archived NWS Watch, Warnings, Advisories. In 2021.
28. Service, N. W. NWS Weather Forecast Offices. <https://www.weather.gov/srh/nws/offices>
29. Ferrell, J.; Brown, V.; Hawkins, M. D., Assessment of NOAA National Weather Service Methods to Warn for Extreme Heat Events. *Weather Clim. Soc.* **2017**, 9, (1), 5-13.
30. NC-DPS Extreme Heat. <https://www.readync.gov/stay-informed/north-carolina-hazards/extreme-heat>
31. NC-DETECT, North Carolina Disease Event Tracking and Epidemiologic Collection Tool. In 2021.
32. Harduar Morano, L.; Waller, A. E., Evaluation of the Components of the North Carolina Syndromic Surveillance System Heat Syndrome Case Definition. *Public Health Rep* **2017**, 132, (1_suppl), 40S-47S.
33. Hibble, M. J., Poisson Regression. In *Modeling count data*, Cambridge University Press: New York, NY, USA, 2014; pp 35-73.
34. Peng, D. R.; Dominici, F., Statistical models. In *Statistical Methods for Environmental Epidemiology with R: A Case Study in Air Pollution and Health*, Gentleman, R., Hornik, K., Parmigiani, G., Ed. Springer: New York, NY, USA, 2008; pp 69-95.
35. Kamo, K. I.; Yanagihara, H.; Satoh, K., Bias-corrected AIC for selecting variables in poisson regression models. *Commun Stat Theory Methods* **2013**, 42, (11), 1911-1921.
36. Dunn, K. P., Smyth K.G., *Generalized Linear Models With Examples in R*. Springer: New York, NY, USA, 2018.
37. Bureau, U. C., Decennial Census. In 2010.
38. Walker, G., Common Statistical Methods for Clinical Research with SAS Examples, Third Edition, 3rd Edition. In 3rd edition ed.; Shostak, J., Ed. SAS Institute: 2010.
39. Ripley, B.; Venables, B.; Bates, D. M.; Hornik, K.; Gebhardt, A.; Firth, D.; Ripley, M. B., Package 'mass'. *Cran r* **2013**, 538, 113-120.
40. Hawkins, M. D.; Brown, V.; Ferrell, J., Assessment of NOAA National Weather Service methods to warn for extreme heat events. *Weather Clim. Soc.* **2017**, 9, (1), 5-13.
41. Abasilim, C.; Friedman, L. S., Comparison of health outcomes from heat-related injuries by National Weather Service reported heat wave days and non-heat wave days - Illinois, 2013-2019. *Int J Biometeorol* **2022**, 66, (3), 641-645.
42. Danzl, D. F., Heat-Related Illnesses. In *Harrison's Principles of Internal Medicine, 20e*, Jameson, J. L.; Fauci, A. S.; Kasper, D. L.; Hauser, S. L.; Longo, D. L.; Loscalzo, J., Eds. McGraw-Hill Education: New York, NY, 2018.
43. Sorensen, C. J.; Cook-Shimane, M.; Newman, L. S., Climate Change and Worker Health: Implications for Clinical Practice. In *CURRENT Diagnosis & Treatment: Occupational & Environmental Medicine, 6e*, LaDou, J.; Harrison, R. J., Eds. McGraw Hill: New York, NY, 2021.
44. Jung, J.; Uejio, C. K.; Kintziger, K. W.; Duclos, C.; Reid, K.; Jordan, M.; Spector, J. T., Heat illness data strengthens vulnerability maps. *BMC Public Health* **2021**, 21, (1), 1999.
45. Jalalzadeh Fard, B.; Mahmood, R.; Hayes, M.; Rowe, C.; Abadi, A. M.; Shulski, M.; Medcalf, S.; Lookadoo, R.; Bell, J. E., Mapping Heat Vulnerability Index Based on Different Urbanization Levels in Nebraska, USA. *Geohealth* **2021**, 5, (10), e2021GH000478.

46. Niu, Y.; Li, Z.; Gao, Y.; Liu, X.; Xu, L.; Vardoulakis, S.; Yue, Y.; Wang, J.; Liu, Q., A Systematic Review of the Development and Validation of the Heat Vulnerability Index: Major Factors, Methods, and Spatial Units. *Curr Clim Change Rep* **2021**, *7*, (3), 87-97.
47. Gronlund, C. J., Racial and socioeconomic disparities in heat-related health effects and their mechanisms: a review. *Curr Epidemiol Rep* **2014**, *1*, (3), 165-173.
48. Guirguis, K.; Basu, R.; Al-Delaimy, W. K.; Benmarhnia, T.; Clemesha, R. E. S.; Corcos, I.; Guzman-Morales, J.; Hailey, B.; Small, I.; Tardy, A.; Vashishtha, D.; Zivin, J. G.; Gershunov, A., Heat, Disparities, and Health Outcomes in San Diego County's Diverse Climate Zones. *Geohealth* **2018**, *2*, (7), 212-223.
49. Toloo, G. S.; Fitzgerald, G.; Aitken, P.; Verrall, K.; Tong, S., Are heat warning systems effective? *Environ Health* **2013**, *12*, (1), 27.
50. Kalkstein, A. J.; Sheridan, S. C., The social impacts of the heat-health watch/warning system in Phoenix, Arizona: assessing the perceived risk and response of the public. *Int J Biometeorol* **2007**, *52*, (1), 43-55.
51. Beckmann, S. K.; Hiete, M., Predictors Associated with Health-Related Heat Risk Perception of Urban Citizens in Germany. *Int J Environ Res Public Health* **2020**, *17*, (3).