

Review

# Dilemmas in the Choice of Adequate Therapeutic Treatment in Patients with Acute Pulmonary Embolism - from Modern Recommendations to Clinical Application

Ratko Lasica<sup>1,\*</sup>, Milika Asanin<sup>1</sup>, Lazar Djukanovic<sup>1</sup>, Nebojsa Radovanovic<sup>1</sup>, Lidija Savic<sup>1</sup>, Marija Polovina<sup>1</sup>, Sanja Stankovic<sup>2</sup>, Arsen Ristic<sup>3</sup>, Marija Zdravkovic<sup>4</sup>, Andjelka Lasica<sup>5</sup>, Jelena Kravic<sup>1</sup> and Jovan Peruničić<sup>1</sup>

<sup>1</sup>Department of Cardiology, Emergency Center, Clinical Center of Serbia, 11000 Belgrade, Serbia; drlasica@gmail.com (R.L.); masanin2013@gmail.com (M.A.); lazardjukanovic08@gmail.com (L.D.); radovanovicn@sbb.rs (N.R.); lidijasavic2007@gmail.com (L.S.); maki.marijapolovina@gmail.com (M.P.); jelena.kravic@gmail.com (J.K.); perun.j@ptt.rs (J.P.)

<sup>2</sup>Clinical Center of Serbia, Center for Medical Biochemistry, 11000 Belgrade, Serbia; sanjast2013@gmail.com (S.S.)

<sup>3</sup>Department of Cardiology, Clinical Center of Serbia, 11000 Belgrade, Serbia; arsen.ristic@med.bg.ac.rs (A.R.)

<sup>4</sup>Clinical Center Bezanjska Kosa, 11000 Belgrade, Serbia; sekcija.kardioloska@gmail.com (M.Z.)

<sup>5</sup>Health center New Belgrade, 11000 Belgrade, Serbia; aleksandralasica763@gmail.com (A.L.)

\* Correspondence: drlasica@gmail.com

**Abstract:** Pulmonary thromboembolism is a very common cardiovascular disease, with a still high mortality rate. Despite the clear guidelines, this disease still represents a great challenge both in diagnosis and treatment. Heterogeneous clinical picture, often without pathognomonic signs and symptoms, represents a huge differential diagnostic problem even for experienced doctors. The decision on the therapeutic regimen also represents a major dilemma in the group of patients who are hemodynamically stable at initial presentation and have signs of right ventricular (RV) dysfunction proven by echocardiography and positive biomarker values (pulmonary embolism of intermediate-high risk). Studies have shown conflicting results about the benefit of using fibrinolytic therapy in this group of patients until hemodynamic decompensation, due to the risk of major bleeding. The latest recommendations give preference to new oral anticoagulants (NOACs) compared to vitamin K antagonists (VKA), except for certain categories of patients (patients with antiphospholipid syndrome, mechanical valves, pregnancy). When using oral anticoagulant therapy, special attention should be paid to drug-drug interactions, which can lead to many complications, even to the death of the patient. Special population groups such as pregnant women, obese patients, patients with antiphospholipid syndrome and cancer represent a great therapeutic challenge in the application of anticoagulant therapy. In these patients, not only the effectiveness of the drugs must be taken into account, but great attention must be paid to their safety and possible side effects, which is why a multidisciplinary approach is emphasized in order to provide the best therapeutic option.

Keywords: acute pulmonary embolism; dilemmas; therapeutic treatment; recommendations; clinical application

## 1. Frequency of pulmonary embolism

Venous thromboembolism (VTE) is the third most common cardiovascular disease (after myocardial infarction and cerebrovascular insult), with an estimated annual incidence in epidemiological studies of 1-2 cases per 1000 population [1, 2]. The frequency of VTE largely depends on age, gender, and associated diseases. With increasing age, the incidence of pulmonary embolism (PE) increases, so that in patients aged 40-49 it amounts to 1.4/1000 patients, and in patients over 80 years old it increases to 11.3/1000 patients [3,4]. The incidence of PE is much higher in women younger than 45 years or older than 80 years compared to men [5]. In patients with cancer, the annual frequency of the first VTE differs depending on the type of cancer (3% in patients with bladder and breast cancer; 4-7% in patients with colon and prostate cancer; 10-12% in lung, stomach, ovarian cancer and brain; 15% in pancreatic cancer) [6]. Recurrent VTE occurs in 30% of patients in the first 10 years from the appearance of the first episode of the disease. It is most often associated with persistent VTE risk factors. More than half a million people die annually in Europe as a result of PE [7].

## 2. Diagnostic doubts and missed diagnoses of VTE

In recent times, the tendency to reduce the unnecessary costs of testing patients with suspected PE on the one hand and overlooking non-specific signs and symptoms of the disease on the other hand, are the causes of the largest number of missed diagnoses of PE. By standardizing algorithms for the diagnosis and treatment of patients with suspected PE, the diagnostic accuracy of this disease has been improved [8]. These diagnostic algorithms include pre-test probability assessment for the existence of PE, evaluation of D-dimer values as well as non-invasive radiographic techniques. However, in clinical practice, the non-specific clinical picture of PE represents a great challenge for the diagnosis of this disease. The initial presentation of patients with PE varies from asymptomatic cases (discovered incidentally during a diagnostic examination to detect other diseases) to high-risk PE resulting in RV dysfunction and subsequent development of shock. The symptoms of the disease in PE can vary and gain in dynamics. Thus, patients initially misdiagnosed with PE may deteriorate clinically to such an extent that re-evaluation will lead to a definitive diagnosis of PE. On the other hand, patients who are hemodynamically unstable with cardiogenic shock may raise the suspicion of an initially cardiac cause of such a condition.

The diagnosis of this sometimes insidious disease is also complicated by the fact that the specificity and sensitivity of symptoms associated with PE is very low, so the presence or absence of any symptom is not enough to confirm or exclude the existence of PE [9]. The most common symptoms and signs of PE are dyspnea (30-88%) [7, 10, 11], pleuritic type of chest pain (from 39-70%) [7, 9,12], leg swelling suspicious for deep vein thrombosis (DVT) (24%) [4], hemoptysis (2-11.8%) [13, 14], cough (9-23%) [11, 14] syncope (6-39%) [7, 9, 14], tachycardia (40%) [7, 9, 10], hypoxemia (70%) [6, 15], new-onset atrial fibrillation (24%)

[16]. The onset of dyspnea is most often sudden, but it can also occur gradually. Patients with emboli present in the segmental and lobar branches of the pulmonary artery rarely experience dyspnea, and it may even be absent in patients with circulatory collapse [12]. The absence of pleuritic pain in PE most often occurs in patients with embolization in the more proximal segment of the pulmonary vasculature. On the contrary, these patients may have anginal pain due to myocardial ischemia as a result of increased RV wall tension and reduced flow through the right coronary artery, which makes differential diagnosis from acute coronary syndrome (ACS) difficult [9]. Even DVT is not a common finding in patients with PE, which further complicates the diagnosis of this disease. About 40% of patients with symptomatic PE have proximal DVT and 25% have distal DVT. Only half of patients with PE and proven DVT have characteristic symptoms for DVT [6]. Syncope and/or shock indicate cardiac insufficiency in patients with high-risk PE. Although syncope is considered a rare manifestation of PE, a cross-sectional study conducted by Prandoni et al. showed a prevalence of PE in patients hospitalized for syncope of about 17% [17]. The incidence of syncope is about 39% in the International Cooperative Pulmonary Embolism Registry (ICOPER), where the symptoms of patients who developed massive pulmonary embolism were analyzed [14].

Most episodes of VTE are provoked by the presence of a number of risk factors, and in some episodes of VTE it is not possible to establish a clear risk factor (unprovoked PE). The presence of strong transient risk factors for the development of VTE (major surgery, prolonged immobilization, and major trauma) that are mentioned at the initial presentation of the patient may reduce the risk of misdiagnosis. Strong transient risk factors are responsible for approximately 20% of all VTE episodes [16]. Also, just knowing that the patient has an active malignant disease reduces the possibility of misdiagnosis or missed diagnosis. It has been shown that 20% of patients with VTE have an active malignant disease [18].

It is recommended that in patients with a low or moderate pre-test probability for PE, D dimer values are also determined, while in patients with a high probability, sophisticated diagnostic methods are immediately implemented [8, 19]. However, both decreased and elevated D dimer values can lead the doctor to misdiagnose. Elevated D-dimer values are also found in patients with inflammatory processes, who suffer from chronic renal failure, cancer during pregnancy, injuries, surgeries [20, 21]. Patients with elevated D-dimer levels due to its low positive predictive value should undergo further diagnostic testing to confirm and/or rule out the diagnosis of PE. Given that normal D-dimer values increase with age, it is necessary to take the patient's age into account when interpreting the results, which is achieved with age-adjusted D-dimer values [22]. Studies have shown that an age-adjusted D-dimer cutoff may be effective and safe in ruling out PE compared with conventional D-dimer (500  $\mu\text{g/L}$ ) [23, 24]. Elevated values of D dimer are physiologically present in pregnancy (they increase in relation to the trimester of pregnancy) which can lead to a wrong diagnosis [25]. Clinical suspicion of the existence of PE and elevated D dimer values are often the reason for

performing CT pulmonary angiography and/or ventilation-perfusion scanning, which exposes both the mother and the fetus to the harmful effects of radiation [26]. In the case of negative D-dimer values combined with a low clinical probability, PE can be ruled out without further diagnostic procedures [19]. A study that analyzed data from the national collaborative study (PIOPED II) showed that despite a low clinical probability for the presence of PE (according to the Wells score), patients had a CT-proven embolus in the main or lobar branches of the PA [12].

Despite advances in medicine, Alonso-Martínez et al. showed that misdiagnosis occurred in 50% (95% CI 44–55) of patients. Higher age, more days of delay and the absence of syncope or sudden onset dyspnea were factors associated with misdiagnosis [27]. To reduce the number of missed diagnoses of PE, the YEARS rule can be used, which consists of three items: clinical suspicion of PE, presence of clinical signs of DVT, presence of hemoptysis [8]. Pulmonary embolism can be ruled out if D-dimer is less than 500 ng/ml, with the presence of one or more of the above signs, or if it is less than 1000 ng/ml without the presence of these signs [9]. The Pulmonary Embolism Rule-out Criteria (PERC) score is also used, where it is possible to rule out PE in patients with a low clinical probability who, in addition, have fulfilled all eight criteria of the PERC rule. The impossibility of its widespread use is reflected in the fact that it can only be applied in clinical settings with a low (<5%) prevalence of PE [8, 6]. Tome van der Hulle and colleagues showed in a prospective cohort study that D-dimer testing in combination with clinical pretest probability assessment using the YEARS criteria or the revised Geneva score can exclude pulmonary embolism [28].

In a systematic review by Kwok Chun S and associates (analysis of 18 studies), it was shown that the most common diagnoses with which PE is mixed are pneumonia, bronchitis, exacerbation of chronic obstructive pulmonary disease, heart failure and acute coronary syndrome [29]. Out of all the misdiagnosed conditions, the proportion with a wrong diagnosis of pneumonia, bronchitis or exacerbation of COPD represented 37.4%, while those wrongly diagnosed with heart failure and coronary artery disease or acute coronary syndrome were 18.2% and 12.4%, respectively [29].

The outcome of patients with PE largely depends not only on the correct diagnosis, but also on the delay in its diagnosis. Approximately 10% of patients with PE have a delay longer than 21 days in establishing the correct diagnosis [29, 30]. The importance of prompt diagnosis and initiation of anticoagulant therapy is reflected in the high risk of early mortality in untreated PE [31].

### **3. Making a diagnosis of PE**

The presence of individual signs and symptoms for PE has a low sensitivity and specificity, while combining them increases the probability that a patient suspected of PE really has this disease. The latest European recommendations for the diagnosis of PE suggest the use of the revised Geneva score and the Wells

score [8]. Regardless of which score is used, the expected percentage of patients with PE is about 10% in the low-probability category, 30% in the intermediate category, and 65% in the high-probability category.

Although changes in the **electrocardiogram (ECG)** in patients with PE are considered insufficiently specific and sensitive for establishing a diagnosis, they can point us to this disease. Most often, sinus tachycardia can be registered in the ECG (present in about 40% of patients); sign S1Q3T3 (Mc Ginn White sign - in about 10% of patients); complete or incomplete right bundle branch block; deviation of the electric shaft to the right; P pulmonale; ST segment depression; T wave inversion; atrial fibrillation [7, 32]. Qr configuration in V1 (Weber and Phillips sign) is specific for PE but has a low prevalence (11-19%). This sign is a predictor of DK dysfunction. Right bundle branch block in patients with PE is associated with a poor prognosis [32].

**Teleradiography of the heart and lungs**, although a non-specific diagnostic method for establishing the diagnosis of PE, can be useful in the differential diagnosis from other diseases that can present with a similar clinical picture.

**Echocardiography** is a non-invasive method with a huge role in the diagnosis and clinical assessment of patients with PE. Enlargement of DK, hypokinesia of the free wall of DK and interventricular septal flattening were found in 27.4%, 26.6%, and 18.4% of patients, respectively [33]. An enlarged DK with akinesia of the basal segment of the free wall (McConnell's sign) can be seen in approximately 20% of patients with PE [33]. The presence of this sign has 77% sensitivity and 94% specificity for the diagnosis of acute PE [34]. Casazza et al. demonstrated that McConnell's sign can also be seen in cases of RV infarction and thus cannot be considered pathognomonic for acute PE [35, 36]. The only sure sign for the existence of PE with this method is the visualization of thrombus masses in the right heart cavities, which occurs in 4-18% of patients with acute PE [37]. **Transesophageal echocardiography**, when it comes to central PE, has a sensitivity of 90-95% and a specificity of 100%. Thromboembolus is very difficult to visualize in the middle part of the left pulmonary artery because the interposition of the left main bronchus interferes with the ultrasound beam.

Thanks to the results of The Prospective Investigation On Pulmonary Embolism Diagnosis (PIOPED) II study, **CT pulmonary angiography (CTPA)** has become the method of choice for the diagnosis of PE (sensitivity of 83% and specificity of 96%) [12].

#### **4. Risk stratification in patients with PE**

Risk stratification in patients with acute PE is necessary to determine the initial therapeutic approach. Determining the correct initial therapeutic regimen is very important because the risk of early mortality in normotensive patients is still high and amounts to 2-8%, while in patients presenting with cardiogenic shock it is up to 30%, and in the case of the need for cardiopulmonary resuscitation, the risk of early mortality


increases up to 65% [38]. Initial stratification is based on clinical symptoms and signs: hemodynamic status at initial presentation, PESI score (Pulmonary Embolism Severity Index) and simplified PESI score (sPESI), presence of RV dysfunction based on transthoracic echocardiography results and biomarker values (Troponin and BNP- a) [8, 39]. Cardiac troponin, a marker of myocardial damage, was shown to be a significant predictor of early mortality in patients with PE, and the presence of positive troponin even in the case of low risk assessed by the PESI score indicates higher early mortality [40]. RV pressure overload due to PE is associated with increased myocardial distension causing the release of BNP and N-terminal pro BNP (NT pro BNP). Natriuretic peptide levels reflect the severity of RV dysfunction in acute PE [41].

All patients with PE can be classified into three risk categories: low, intermediate and high. Patients who are hypotensive (systolic blood pressure less than 90 mmHg) or in shock (systolic blood pressure less than 80 mmHg) on admission belong to the category of patients with high-risk PE (about 5% of PE cases) [8, 42]. Such patients should be treated with thrombolytic therapy and/or embolectomy. (Figure 1.)

In the group of patients in whom hemodynamic instability is not present, further risk stratification is based on the assessment of prognostic criteria, namely clinical, visualization and laboratory indicators (mostly related to proving the presence of RV dysfunction). Patients with a low risk of early mortality are normotensive on admission, have a sPESI score of less than 1, have no RV dysfunction on echocardiographic examination and have negative biomarker values and are treated with anticoagulant therapy [43].

Patients with a intermediate risk of early mortality are hemodynamically stable on admission and have a sPESI score of high than one. If they have positive or biomarker values or echocardiography findings (presence of signs of RV dysfunction), they belong to the intermediate-low risk group and are treated with anticoagulant therapy [8]. If these patients have positive biomarker values and the presence of RV dysfunction, they belong to the intermediate- -high risk group and are initially treated with anticoagulant therapy, and in case of hemodynamic decompensation with thrombolytic therapy [44].

It is also important to note that the risk of pulmonary embolism is a dynamic category, so the initially set therapeutic decision can be changed depending on the patient's clinical condition. Choosing the right initial treatment for PE patients not only affects their survival, but also reduces the frequency of post-thromboembolic pulmonary hypertension, which worsens the long-term prognosis of these patients and reduces the quality of life.



**Figure 1. Early mortality risk and initial treatment in patients with acute pulmonary embolism.**

Legend: **ACT**- anticoagulant therapy; **RT**-reperfusion therapy; **PESI**- The Pulmonary Embolism Severity Index Score; **sPESI**- simplified Pulmonary Embolism Severity Index Score.

## 5. Initial therapeutic regimen in the acute phase of PE

### 5.1. Anticoagulant therapy

Initiation of anticoagulant therapy without delay is necessary in patients with a high and intermediate clinical probability for PE, while the diagnostic procedure is still in progress. Hemodynamically stable patients should be treated with anticoagulant therapy: unfractionated heparin (UFH), low molecular weight heparin (LMWH) or new oral anticoagulants ((NOACs: apixaban, dabigatran, edoxaban or rivaroxaban)) [45]. NOACs are administered depending on whether oral therapy can be administered, depending on kidney function and other circumstances. Parenteral anticoagulant therapy is based on the use of indirect and direct anticoagulant drugs. The activity of indirect anticoagulant drugs is based on binding to plasma cofactors, while direct anticoagulant drugs do not need a plasma cofactor to exert their effect. Subcutaneous administration of low molecular weight heparin or fondaparinux, or intravenous administration of unfractionated heparin is common.

**5.1.1 Unfractionated Heparin** - an indirect anticoagulant that achieves its effect by binding to antithrombin III. The resulting complex inactivates thrombin and factors Xa, IXa, XIa and XIIa [46]. It also inhibits the aggregation of platelets, activates osteoclasts, and inhibits the formation of osteoblasts [47]. It achieves its anticoagulant effect by reducing the propagation of thrombus and preventing new embolic events. The main limitation of heparin stems from its tendency to bind to positively charged plasma proteins, proteins released from platelets and endothelial cells, which can result in a variable anticoagulant response and the phenomenon of heparin resistance [48, 49]. Due to its largely dose-dependent clearance, the plasma half-life

of UFH ranges from 30-60 min. [46]. One of the side effects of heparin is the occurrence of heparin-induced thrombocytopenia (HIT). The frequency of HIT ranges from 1 to 5% of patients receiving heparin [50]. Type I is the most common form of thrombocytopenia and is self-limiting [51]. Mild thrombocytopenia occurs in the first 2 days after heparin initiation and normalizes even with continued heparin therapy [50]. A more serious form (HIT, Type II) is an immune-mediated disorder characterized by the formation of antibodies against the heparin-PF4 complex. Typically, the platelet count declines by >50% between days 5 and 14 of heparin treatment [52]. When compared with low molecular weight heparin (LMWH), UFH is much more likely to cause HIT [53, 54]. Prandoni P et al. showed in a prospective study that the incidence of HIT in patients who received LMWH was about 0.8%. It was also shown that patients were more likely to get HIT if they were treated with UFH before LMWH (1.7 vs. 0.3%, OR, 4.9; 95% CI, 1.5 – 16) [55]. By activating osteoclasts, heparin can affect the occurrence of osteopenia. The initial dose of UFH depends on body weight and is 80 IU/kg in an intravenous bolus, and then continues with 18 IU/kg intravenous in a continuous infusion [8]. The therapeutic effect of heparin is controlled by determining the activated partial thromboplastin time (aPTT), which should be maintained in the range of 1.5 to 2.5 times higher than the starting aPTT.

The use of UFH is currently mostly limited to patients with obvious hemodynamic instability or immediate hemodynamic decompensation, in whom primary reperfusion treatment will be necessary. Intravenous unfractionated heparin is also recommended for patients with severe renal impairment (creatinine clearance <30 ml/min) and/or severe obesity [8, 56].

In case of HIT development, it is necessary to use direct thrombin inhibitors (lepirudin, argatroban, bivalirudin) as well as factor Xa inhibitors (eg fondaparinux).

**5.1.2 Low molecular weight heparin** - its anticoagulant action is mostly achieved by inhibiting activated factor X. LMWH has advantages over heparin in that it has better bioavailability and longer half-life, simplified dosing, predictable anticoagulant response, lower risk of HIT, and lower risk of osteoporosis [56]. LMWH is administered in fixed doses. Administration of LMWH does not require aPTT monitoring and can be administered subcutaneously [57, 58]. The usual dose of LMWH for therapeutic anticoagulation is 1 mg/kg given every 12 hours, although another option is 1.5 mg/kg given once daily for VTE. Dose efficacy of LMWH should also be monitored in patients with renal dysfunction, as LMWH is excreted by the kidneys, so dose adjustments are sometimes necessary in relation to creatinine clearance. Unlike UFH, the clearance of LMWH is dose-independent, and the half-time in plasma is approximately 4 hours. AntiXa level monitoring is not routinely recommended, but its therapeutic range is 0.6-1.0 mmol/l. The following LMWHs are approved for the treatment of PE: enoxaparin, tinzaparin, dalteparin, nadroparin [8].

The main complications during UFH and LMWH treatment are bleeding. In patients treated for VTE, a lower risk of bleeding was registered when using LMWH compared to UFH (OR=0.68, p=0.05). In the

Systematic Review of the Cochrane Collaboration on VTE, LMWH were shown to be significantly safer than UFH (incidence of major bleeding, 1% vs 2.1%, OR 0.57; 95% CI 0.39–0.83) [57]. The advantage of intravenous administration of UFH is that the overdose can be easily neutralized by the administration of protamine sulfate. Protamine sulfate in a dose of 1 mg neutralizes 100 IU of UFH. Although protamine sulfate can be used as an antidote to LMWH, it does not completely neutralize the anticoagulant activity of LMWH because it binds only to the longer chains of LMWH [59].

**5.1.3 Fondaparinux** – is also a factor Xa inhibitor. Its half-life is 17 h. Because fondaparinux is cleared unchanged via the kidney, it is contraindicated in patients with a creatinine clearance below 30 mL/min and should be used with caution in those with a creatinine clearance below 50 mL/min. [60]. Unlike LMWH, there is no cross-reactivity between Fondaparinux and HIT antibodies [61]. Lori-Ann Linkins et al. within the framework of a systematic review conducted on 9 studies, she concluded that the use of Fondaparinux is effective and safe for the treatment of HIT, although according to the recommendations, this drug is not included for the treatment of this condition [62]. The major side effect of fondaparinux is bleeding. There is no antidote for this drug. Anticoagulant effect is achieved when administered subcutaneously in a dose of 2-8 mg, or intravenously in a dose of 2 mg to 20 mg. Due to its almost complete bioavailability after subcutaneous administration, long half-life, lack of variability in anticoagulant response, fondaparinux can be administered in a single daily dose without laboratory monitoring [63]. In the treatment of DVT and PE, it is given in a dose of 7.5 mg for patients with a body weight of 50-100 kg; 5 mg for patients under 50 kg and 10 mg for patients over 100 kg body weight [8]. Fondaparinux does not bind to protamine sulfate. If bleeding occurs, recombinant factor VII may be effective [64].

**5.1.4 Direct thrombin inhibitors** are used in patients who develop HIT. Direct thrombin inhibitors are a class of anticoagulants that inhibit thrombin generation by binding to the active site on thrombin. They are classified as univalent or bivalent inhibitors. Univalent inhibitors such as argatroban interact with the active, catalytic site of thrombin itself compared to bivalent inhibitors (hirudin and its derivatives), which recognize the binding site on fibrinogen. Three prospective studies, HAT-1, HAT-2 and HAT-3 demonstrated the effectiveness of **lepirudin** as an anticoagulant in reducing the risk of thromboembolic complications in patients with HIT [65, 66, 67]. The pharmacodynamic effect of lepirudin on thrombin activity was assessed using the aPTT. Prolongation of aPTT was linear over the range of plasma concentrations tested (up to 0.5 mg/kg i.v.). Lepirudin can be successfully used subcutaneously in patients with HIT. The mean half-life was around 1.7 h. Dose was shown to correlate linearly with aPTT response with lepirudin doses of 0.75 – 2 mg/kg s.c. in two daily doses. The elimination of Lepirudin is predominantly through the kidneys, so in patients with renal failure, the half-time of elimination can be significantly prolonged, so dose adjustment is recommended

[68]. The recommended dose of lepirudin in patients with normal renal functions is 0.4 mg/kg i.v. push (over 15 – 20 s), followed by continuous infusion at 0.15 mg/(kg/h). **Argatroban** - has a half-life of 40-50 minutes and requires dose reduction both in patients with liver damage and in critically ill patients. For the prophylaxis or treatment of thrombosis in HIT, the recommended starting dose of argatroban is 2 mg/kg/min for patients without hepatic impairment [69]. The starting dose should be reduced in patients with moderate hepatic impairment to 0.5 mg/kg/min. The initial dose can be adjusted up to 10 mg/kg/min to achieve a stable aPTT 1.5–3 times longer than basal. **Bivalirudin** is also a direct thrombin inhibitor that is injected at a dose of 0.05–0.1 mg/kg/h. Argatroban may be advantageous compared to bivalirudin in achieving initial therapeutic anticoagulation goals among patients with suspected or confirmed HIT [70]. American Society of Hematology 2018 guidelines for management of venous thromboembolism: heparin-induced thrombocytopenia suggest the possible use of NOAC in HIT [76]. With respect to the choice of NOAC, most of the published experience in HIT is with rivaroxaban [71]. (Table 1)

Drug	Dose	Special consideration	Specific patient characteristics	Reversal agents
<b>UFH</b>	80 unit/kg IV bolus, followed by an 18-unit/kg/h infusion.	<b>Avoid</b> HIT; Osteopenia Pronounced drug-drug interactions	Overt haemodynamic instability; (CrCl) $\leq$ 30 mL/min; Pregnancy; Severe obesity	Protamine sulfate
<b>LMWH</b>	1 mg/kg twice daily 1.5mg/kg once daily	<b>Avoid</b> with severe renal impairment	Pregnancy; Obesity;	Protamine sulfate
<b>VKA</b>	<b>Warfarin</b> 5 mg/day once daily 4 mg/day once day-patients > 70 years	<b>Cross the placenta- contraindicated in pregnancy</b>	Antiphospholipid syndrome; Mechanical heart valves; Extremely reduced renal function; Severe mitral stenosis;	4F-PCC 4 or FFP
<b>Apixaban</b>	10 mg twice daily for 7 days followed by 5 mg twice daily	<b>Avoid in</b> CrCl < 15 mL/min Severe hepatic impairment	Previous GI bleeding or high risk of bleeding; Patients with CA; Elderly patients;	Andexanet
<b>Rivaroxaban</b>	15 mg- twice daily (3 weeks) then 20 mg once daily (at least 6 months)	<b>Avoid in</b> CrCl < 30 mL/min; (FDA) CrCl < 15 mL/min (EMA).	Low risk of bleeding and without gastrointestinal tumours; Patient preference – a single dose regimen;	Andexanet
<b>Dabigatran</b>	150 mg - twice daily 110 mg- twice daily for patients $\geq$ 80 years	<b>Avoid in</b> CrCl < 30 mL/min.; Concomitant treatment with P-gp inhibitors in patients with CrCl < 50 mL/min; Reduce dose to 110 mg for patients $\geq$ 80 years or $\geq$ 75 years with at least one bleeding risk factor.	Can be removed by hemodialysis in patients with severe renal impairment.	Idarucizumab
<b>Edoxaban</b>	60 mg - once daily 30 mg- once daily if body weight $\leq$ 60 kg	<b>Avoid</b> CrCl < 15 mL/min. Severe hepatic dysfunction	Low risk of bleeding and without gastrointestinal tumours	Andexanet
<b>Fondaparinux</b>	5 mg subQ daily < 50 kg 50–100 mg: 7.5 mg subQ daily 10 mg subQ daily > 100 kg	<b>Avoid</b> CrCl < 30 mL/min	HIT ( off label); allergy of LMWH	Factor VIIa
<b>Lepirudin</b>	0.4 mg/kg i.v over 15-20 min, followed by 0.15mg/kg/h infusion	<b>Avoid</b> Severe renal impairment	HIT	
<b>Argatroban</b>	2 $\mu$ g/kg/min Liver dysfunction (bilirubin > 1.5 mg/dL) $\rightarrow$ 0.5-1.2 $\mu$ g/kg/min	<b>Avoid</b> in severe hepatic dysfunction	HIT	
<b>Bivalirudin</b>	0.15 mg/kg/h	<b>Avoid</b> Severe renal impairment	HIT ( off label)	

**Table 1. Treatment with anticoagulant therapy in patients with pulmonary embolism.** Legend: **UFH**- unfractionated heparin ; **LMWH**- low molecular weight heparin ; **VKA**- vitamin K antagonists ; **HIT**- heparin-induced thrombocytopenia; **CrCl**- creatinine clearance ; **4F-PCC**- 4-factor prothrombin complex concentrate; **FFP**- fresh-frozen plasma ; **GI**- gastrointestinal; **CA**-cancer; **P-gp inhibitors**- P glycoprotein inhibitors; **subQ**- subcutaneously;

## 5.2. Thrombolytic therapy

The application of thrombolytic therapy, systemic or directed by a catheter into the pulmonary artery, is used in patients with high-risk PE, which is accompanied by hemodynamic instability [42,72]. Thrombolytic therapy in these patients leads to recanalization of the blood vessel and normalization of the hemodynamic status, which is accompanied by a reduction in RV dilatation on echocardiography. Thrombolytic drugs dissolve blood clots by activating plasminogen, which forms a product called plasmin. Plasmin is a proteolytic enzyme capable of breaking the cross-links between fibrin molecules, which ensure the structural integrity of blood clots. The greatest benefit is seen if therapy is started within the first 48 hours of symptom onset, although thrombolysis may also be beneficial in patients who have had symptoms for 6-14 days [42, 72]. Medicines that are used from this group are: streptokinase, urokinase and recombinant tissue plasminogen activator (rtPA) [8]. Contraindications to the use of thrombolytic therapy can be divided into **absolute** (history of hemorrhagic stroke, ischemic stroke in the last 6 months, neoplasm of the central nervous system, major trauma or surgery, head injury in the previous 3 weeks, hemorrhagic diathesis, active bleeding) and **relative** (transient ischemic attack in the previous 6 months, use of oral anticoagulant drugs, pregnancy, refractory hypertension, advanced liver disease, infective endocarditis, active peptic ulcer) [73]. The potential benefits are often complicated by the relatively high incidence of hemorrhagic complications. Within the meta-analysis of Saurav Chatterjee et al. the frequency of major bleeding during the use of thrombolytic therapy was shown to be 9.24% [74].

### 5.3. Embolectomy

**5.3.1. Surgical embolectomy** with cardiopulmonary bypass can be performed in patients with acute PE accompanied by hemodynamic instability and a contraindication to thrombolytic therapy [44]. Lee T et al. did not show a significant difference in 30-day mortality when using medical thrombolytic therapy and surgical embolectomy (15 vs 13% respectively)[75]. Stein et al. reported that the mortality of pulmonary embolectomy for acute PE was 20% in a meta-analysis of 1,300 cases [76]. Case series have also been published that have shown variable results with perioperative mortality ranging from 2.3% to 27.2% [77]. Extracorporeal membrane oxygenation (ECMO) in combination with surgical embolectomy can be used in case of hemodynamic instability of the patient with contraindications for the use of thrombolytic therapy [78].

**5.3.2. Percutaneous catheter embolectomy** is the removal of a thrombus from the pulmonary circulation and involves: 1) fragmentation of the thrombus using a balloon catheter or loop; 2) rheolytic thrombectomy using hydrodynamic catheters; 3) suction thrombectomy using aspiration catheters and 4) rotary thrombectomy [79]. For patients who do not have absolute contraindications to thrombolytic treatment, percutaneous catheter embolectomy or pharmacomechanical thrombolysis is preferred [8].

### 5.4. Initial therapeutic regimen in patients with intermediate-high risk pulmonary embolism

Patients with moderate-high risk PE are a very heterogeneous group of patients, in whom there is still a dilemma regarding the overall therapeutic benefit of using fibrinolytic therapy compared to heparin treatment. According to study data: Moderate pulmonary embolism treated with thrombolysis (MOPET); Management Strategies and Prognosis of Pulmonary Embolism-3 (MAPPET-3); Treatment of submassive pulmonary embolism with tenecteplase or placebo: Cardiopulmonary Outcomes at Three months (TOPCOAT); Fibrinolysis for Patients with Intermediate-Risk Pulmonary Embolism (PEIHTO) patients with moderate-high risk PE had a significant clinical benefit from the application of thrombolytic therapy [80, 81, 82]. In the PEIHTO study in patients with intermediate-high risk PE, fibrinolytic therapy with anticoagulant therapy compared to placebo prevented death and hemodynamic decompensation of the patient (2.6% vs 5.6% respectively) but increased major bleeding (6.3 vs 1.2% ) and stroke [82]. A meta-analysis by Saurav Chatterjee et al. showed that in intermediate-risk pulmonary embolism trials, thrombolysis was associated with lower mortality (OR, 0.48; 95% CI, 0.25-0.92) and more major bleeding events (OR, 3.19; 95% CI, 2.07-4.92) compared with anticoagulant therapy [74]. A meta-analysis by Nakamuras et al. showed that thrombolytic therapy significantly reduced the incidence of the composite endpoint of all-cause death or clinical deterioration (3.9% vs. 9.4%; RR, 0.44;  $P < 0.001$ ) [83]. For now, the decision to apply thrombolytic therapy is left to the prescribing physician, who should summarize the risk of clinical deterioration (age, comorbidities, degree of DK dysfunction, biomarker values, respiratory status) and the risk of hemorrhagic complications. In addition, different combinations of clinical and laboratory parameters were used to construct prognostic scores, which allow semiquantitative assessment of the early risk of death associated with PE. One of those risk scores is the Bova score, which includes: RV dysfunction, arterial blood pressure values, heart rate, and troponin values. The model identified three stages (I, II and III) with 30-day PE-related complication rates of 4.2%, 10.8% and 29.2%, respectively [84]. In 2021, the CHEST recommendations and the report of the expert panel on the indications for the use of thrombolytic therapy in patients with moderate-risk PE were published [85]. These recommendations also suggest that hemodynamically stable patients be treated aggressively with anticoagulant and not thrombolytic therapy with careful monitoring, and that only if hemodynamic decompensation occurs, thrombolysis should be administered. Apart from hypotension, other types of deterioration such as progressive increase in heart rate, drop in systolic blood pressure ( $>90$  mm Hg), decrease in saturation or partial pressure of oxygen in arterial blood, worsening of RV dysfunction and increase in biomarker values are indications for the use of thrombolytic therapy. For a long time, the question was raised as to what dose of thrombolytics would be effective in these patients without increasing the hemorrhagic risk. For a long time, the question was raised as to what dose of thrombolytics would be effective in these patients without increasing the hemorrhagic risk. Results of a randomized pilot study Wang et al. in 2010, they showed that the administration of half the therapeutic dose of rtPA (50 mg) in patients who are

hemodynamically triggered is not inferior to the full dose in terms of RV function recovery [86]. However, the results of the mentioned before study are not supported by current guidelines for the treatment of intermediate-high risk PE [8].

## **6. Prolonged treatment with anticoagulant therapy of patients with PE**

The development of awareness that PE is often recurrent has led to a more precise consideration of the duration of anticoagulant therapy after the initial event. According to the current recommendations of the European Association of Cardiologists for the diagnosis and treatment of PE, extended treatment with anticoagulant therapy should last at least three months, and in a large number of patients, longer than three months [8]. Continuation of anticoagulant therapy for longer than 3-6 months leads to a reduction of 80% or more in the occurrence of recurrent VTE, but an increase in the risk of major bleeding by 1-3 times with NOACs and 4-5 times with vitamin K antagonists [87]. Patients with VTE associated with transient risk factors have a low recurrence risk of 3%. In this group of patients, oral anticoagulant therapy for three or more months is justified (assuming that the risk factor has been resolved). Patients with idiopathic (unprovoked) thromboembolism have a significantly higher recurrence rate, corresponding to an annual event rate of 7.9%. In these patients, prolonged use of anticoagulant therapy should last longer than 3 months [88]. Oral anticoagulant treatment of indefinite duration is recommended for patients who have recurrent VTE not associated with a reversible risk factor. In these patients, the interval of re-evaluation of the continuation of oral anticoagulant therapy must be determined, depending on the risk/benefit ratio. Patients who have any of the thrombophilias (Factor V Leiden mutation, MTHFR mutation and prothrombin 20210 mutation) are candidates for prolonged use of anticoagulant therapy. Patients with antiphospholipid syndrome, active cancer, and protein S, C, and antithrombin III deficiency are at significant risk for thromboembolic recurrence and lifelong anticoagulation should be considered [89]. Aspirin is not recommended for prolonged treatment of PE, given its lower efficacy and similar safety profile compared to NOACs [90]. In all patients who had a first episode of VTE, it is important to assess the risk of repeated episodes of thromboembolism. This can be done using several different scores such as Vienna prediction model (VPM), DASH score and HERDOO2 scoring system, which use different patient parameters such as gender, d dimer values, location of the thrombus during the first event [91, 92]. The significance of these scores is that they can help clinicians decide on the duration of NOACs after the first VTE.

### **6.1 Vitamin K antagonists**

For more than 60 years, vitamin K antagonists were the only oral anticoagulants available for clinical use. Vitamin K antagonists are rapidly absorbed from the gut and 90% are bound to circulating albumin. The half-life of VKA is 36 to 48 hours, and hepatic metabolism by microsomal enzymes, including cytochrome P450 variants, partially explains the marked interindividual variation in the doses required for drug effect. A

number of factors influence the biological activity of VKA, including patient's age, patient's activity level, diet, drug absorption, albumin levels, vitamin K intake, general health status, and use of various medications.

Drug-drug interactions are very important when it comes to the safety of drug administration, especially in patients with multiple comorbidities, who are treated with a large number of drugs. Cholestyramine and sucralfate are thought to reduce the gastrointestinal absorption of warfarin [93]. On the other hand, drugs such as amiodarone, fluvastatin, lovastatin, fluconazole and sertaline can potentiate the effect of warfarin by inhibiting CYP2C9 [93]. By binding to plasma proteins, drugs such as ibuprofen, quinidine, fenofibrate, losartan, valsartan, amlodipine and felodipine can take an important place for warfarin and thus potentiate its anticoagulant effect [94]. Metronidazole, fluconazole, trimethoprim-sulfamethoxazole, miconazole can affect the effect of warfarin by inhibiting CYP 1A or CYP 3A4, and macrolide antibiotics only by inhibiting CYP3A4 [94]. The use of antibiotics can reduce the absorption of vitamin K by affecting the intestinal flora [95]. Concomitant use of antiplatelet drugs such as aspirin and P2Y<sub>12</sub> receptor inhibitors with warfarin greatly increases the risk of bleeding [96]. The interaction of warfarin with selective serotonin reuptake inhibitors (SSRI) leads to a decrease in serotonin in platelets and a decrease in platelet aggregation [97]. Nonsteroidal anti-inflammatory drugs (NSAIDs) affect the activity of warfarin by inhibiting cytochrome p450, and by inhibiting the digestion of prostaglandins, they can lead to the formation of erosions on the stomach, increasing the risk of gastrointestinal bleeding [98]. The most recent meta-analysis showed a higher degree of clinically relevant bleeding with the combined use of warfarin and antiplatelet drugs (OR=1.74; 95% CI 1.56, 1.94), antibiotics (OR=1.63; 95% CI 1.45, 1.83), NSAIDs (OR=1.83; 95% CI 1.29, 2.59), SSRIs (OR=1.62; 95% CI 1.42, 1.85), compared with warfarin alone [99]. Also, within this meta-analysis, no increase in clinically relevant bleeding was demonstrated with the combined use of amiodarone, a beta blocker. In this study, no drug combined with warfarin had a significant effect on thromboembolic events or mortality. Many statins (fluvastatin, lovastatin, simvastatin, atorvastatin) bind to plasma proteins and affect the prolongation of the INR value in patients with concomitant use of these drugs [100, 101]. This is confirmed by a study by Anna E and collaborators who measured INR values in patients who used warfarin with some of the statins (simvastatin, rosuvastatin, atorvastatin). Simvastatin initiation led to an increase in mean INR from 2.40 to 2.71, with INRs peaking after 4 weeks, corresponding to a mean change of 0.32 (95%CI 0.25-0.38). Initiation of atorvastatin and rosuvastatin lead to INR increases of 0.27 (95%CI 0.12-0.42) and 0.30 (95%CI -0.09-0.69) [102].

Many studies have shown that various types of supplements and herbs can affect CYP2C9 activity, so caution should always be exercised in their combined use with warfarin [103]. When it comes to supplements, it has been shown that combined intake of grapefruit, mango, ginkgo biloba, cranberry with warfarin can prolong the INR and increase the likelihood of bleeding, while the use of St. John's wort and green tea can

reduce its effect [104-109]. Concomitant consumption of foods rich in vitamin K (lettuce, broccoli, spinach, green peas) was associated with a risk of reduced therapeutic effect of VKA [104].

The varying effectiveness of the drug is expressed in varying INR. Because of occasionally high INR values, patients are exposed to the risk of hemorrhagic complications, and because of low INR values, to the risk of thromboembolic events. The risk of bleeding is directly related to the INR; it begins to rise quite significantly when the INR increases to more than 3.5, and an INR greater than 5 requires the administration of fresh frozen plasma or Vitamin K [110]. The average oral dose of Warfarin is 5 mg/day, and for patients over 70 years old it is 4 mg /day. The optimal target INR for treatment and prophylaxis of VTE is 2.0–3.0. Given the long half-life of both VKA (36 hours) and factor II (60 hours), to achieve a full anticoagulant effect the INR must be in the therapeutic range for at least 4 to 6 days to achieve a 20-30% reduction in vitamin K-dependent coagulation factors compared to normal [8]. Frequent INR controls, constant INR variations and the risk of thrombosis/bleeding made NOACs preferable except in specific indications (antiphospholipid syndrome, presence of mechanical valves, extremely reduced renal function, severe mitral stenosis).

## 6.2 New oral anticoagulant drugs

In recent years, novel treatment strategies with direct oral anticoagulants (NOACs) have been increasing in popularity and availability. NOACs are preferred over vitamin K antagonist (VKA) therapy. This is due to a similar reduction in the risk of recurrent VTE, a reduced risk of bleeding and easier monitoring of their effect compared to vitamin K antagonists [111]. Each of the NOACs showed similar results compared to VKA when it comes to recurrent PE [112, 113]. In a meta-analysis by Mohammad Alhousani and colleagues, NOACs and other anticoagulants (VKA and LMWH) showed no statistical difference in preventing recurrent VTEs among patients with chronic renal failure, but NOACs had significantly lower risk of major and non-major clinically relevant bleeding regardless of the level of renal impairment compared to VKAs [114].

### 6.1. Apixaban

The oral bioavailability is approximately 50%, with most of the drug absorption occurring in the small intestine [115]. Drug elimination occurring via the metabolism through the CYP3A4 systems in the intestine and liver and the P-glycoprotein system can be enhanced through drug-drug interactions Apixaban [116]. Apixaban is given at a dose of 10 mg twice a day for 7 days and then 5 mg twice a day [8]. Andexanet alfa can be used to reverse apixaban (off-label) in life-threatening or uncontrollable bleeding. The dosing is based on the specific factor Xa agent-inhibitor to be reversed, dose, and the time since the last dose was administered [116]. Andexanet alfa 400 mg intravenous bolus is administered at a rate of 30 mg/min, followed by 4 mg/min

via continuous infusion for up to 120 min, to reverse apixaban (5 mg or less) or rivaroxaban (10 mg or less), administered within 8 h or if the time is unknown [116].

## 6.2. Rivaroxaban

Rivaroxaban is an oral direct factor Xa inhibitor approved for the prevention and treatment of DVT and PE [8]. Rivaroxaban is given in a dose of 2x15 mg for 3 weeks and then 20 mg once a day. Renal elimination accounts for approximately 36% of unchanged drug, its use in patients with a CrCL <30 mL/min is not advised, and <15 mL/min employment is contraindicated [117]. Rivaroxaban is not dialyzable. A significant increase in rivaroxaban exposure was demonstrated with the strong CYP3A4 inhibitors, ketoconazole (158% increase [95% CI 136%, 182%] for a 400 mg once daily dose) and ritonavir (153% increase [95% CI 134%, 174%]). Therefore, the use of rivaroxaban should be avoided with strong combined CYP3A4 inhibitors (mainly antifungals (except fluconazole) and HIV protease inhibitors) [118].

## 6.3. Dabigatran

Dabigatran is a direct thrombin (IIa) inhibitor approved for the prevention and treatment of DVT and PE [8]. Caution must be used in the elderly, as the risk of stroke and bleeding increases with age, as seen in an analysis of the RE-LY (Randomized Evaluation of Long-Term Anticoagulant Therapy) trial [119]. Dabigatran is contraindicated for use in patients with mechanical heart valves. The RE-ALIGN (Randomized, Phase II Study to Evaluate the Safety and Pharmacokinetics of oral dabigatran Etxilate in Patients after Heart Valve replacement) trial was terminated early due to thromboembolic events (valve thrombosis, stroke, and myocardial infarction), and major bleeding was observed in the dabigatran group compared to the warfarin group in heart valve patients [116, 120]. Dabigatran is a substrate for P-glycoprotein 1 (P-gp) and its concomitant use with strong P-gp inhibitors such as verapamil and ketoconazole or P-gp inducers such as rifampicin, carbamazepine and phenytoin may affect its effect [94]. Although co-administration of Dabigatran with most Pgp inhibitors does not require dose adjustment, it should be administered at least 2 hours before these drugs when co-administered. In the documents of the US Food and Drug Administration, verapamil increases the exposure ie the effectiveness of dabigatran. The European Medicines Agency recommendations and clinical guidance suggest a dabigatran dose reduction to 110 mg when given with verapamil regardless of kidney function. No dose adjustment is needed with the concomitant use of P-gp inhibitor amiodarone [121]. The use of Dabigatran with ketoconazole, itraconazole should be avoided because its bioavailability increases with their simultaneous use [122]. Care should also be taken when using St. John's wort and dabigatran because their simultaneous use may affect the reduced activity of the drug. No interaction was observed when dabigatran was coadministered with atorvastatin, an HMG-CoA reductase inhibitor. When the granules are taken without a protective capsule, its bioavailability increases by as much as 75% compared to an intact capsule [123]. Therefore, it is not advisable to crush or chew the medicine. Due to the fact that the capsules

are designed to be released in the stomach, they must not be used in patients receiving medication through a nasogastric or jejunostomy tube. It is mainly eliminated by the kidneys and is the only NOAC that can be removed by hemodialysis [124].

Dabigatran is given in a dose of 150 mg twice a day and for patients older than 80 years in a dose of 110 mg twice a day. The agent named idarucizumab is a monoclonal antibody with 350 times more affinity for dabigatran than thrombin.

#### **6.4. Edoxaban**

Approval was based primarily on the Hokusai VTE study, which evaluated 3319 patients with PE. The trial showed that edoxaban was not inferior to warfarin but had a lower bleeding risk [125]. The oral bioavailability is approximately 60%, and renal elimination accounts for approximately 50% of unchanged drug. Edoxaban is given 60 mg once a day and 30 mg once a day if the body weight is  $\leq 60$  kg. The simultaneous use of edoxaban and ritonavir, erythromycin, azithromycin, clarithromycin, ketoconazole should be avoided due to a significant increase in its plasma concentration [94, 126]. If macrolide antibiotics and edoxaban are necessary, its dose can be reduced by 50% [127]. Amiodarone may increase edoxaban exposure by 40%. In the case of cyclosporine and tacrolimus administration, dose adjustment is necessary [128]. A study by Mainbourg S and associates that examined the use of NOACs (apixaban, edoxaban, rivaroxaban and dabigatran) in different dosage regimens and analyzed whether the dosage method correlates with the occurrence of thrombogenic events. There was no difference for major thrombotic event (RR BID/QD = 1.06, 95%IC 0.86–1.30) nor for major bleeding (RR BID/QD = 1.02, 95%IC 0.84–1.23) between NOACs that were dosed twice daily or once a day, without heterogeneity ( $I^2 = 0\%$ ) [129]. If oral anticoagulation continues after 3 months of PE in a patient without cancer, after 6 months of therapy, consider reducing the NOAC dose: apixaban (2.5 mg daily) or rivaroxaban (10 mg daily). When using all NOACAs, the use of proton pump inhibitors (PPIs) is also suggested. No interactions between PPIs, apixaban, rivaroxaban and edoxaban have been shown. Administration of dabigatran and PPIs reduced the absorption of dabigatran by 20%, because an acidic environment is necessary for the absorption of dabigatran.

### **7. Treatment of PE in specific patient populations**

#### **7.1. Treatment of PE in pregnancy**

Acute PE remains one of the leading causes of maternal death. The risk of VTE is higher in pregnant women compared to women of a similar age who are not pregnant. Given that they do not cross the placental barrier, the use of UFH and LMWH has proven to be the safest during pregnancy [40]. Due to the ease of administration and known pharmacodynamics, LMWHs are used most often. In pregnant women in whom PE occurred a month before delivery, it is possible to schedule induced delivery with suspension of LMWH 24 hours before delivery. According to ESC recommendations in high-risk situations, for it is recommended that

LMWH be converted to UFH  $\geq 36$  h prior to delivery [8]. The UFH infusion should be stopped 4 – 6 h prior to anticipated delivery [40]. The use of VKA is not recommended for pregnant women considering that they pass through the placenta and their use is associated with embryopathies, fetal and neonatal bleeding [130]. Cases of placental abruption are also reported during their use. The use of NOAC is contraindicated in pregnant women [8]. Data on the use of NOACs in pregnancy indicate an increased rate of miscarriage (31%) compared to the rate of miscarriage in the general population. Bone and facial abnormalities in the fetus were observed in 4% of pregnancies during rivaroxaban use, although as many as 85% of women discontinued NOACs within the first two months of pregnancy [131]. The decision to use anticoagulants during pregnancy, especially before delivery, requires a multidisciplinary team of anesthesiologists, gynecologists and cardiologists in order to reduce unwanted hemorrhagic side effects. After delivery, anticoagulant treatment in lactating women includes the use of UFH, LMWH, vitamin K antagonists and fondaparinux. NOACs are concentrated in breast milk and are contraindicated but may be considered in women who are not breastfeeding or after breastfeeding in those who have an indication for longer-term treatment.

### **7.2. Treatment of PE in antiphospholipid syndrome**

Antiphospholipid syndrome is associated with a high risk of recurrent venous thrombosis. The persistent presence of elevated values of antiphospholipid antibodies during the first manifestation of VTE is an indication for an unlimited duration of anticoagulant therapy. Recommendations of the European Association of Cardiologists do not support the use of NOACs instead of VKA [8]. According to a study by Vittorio Pengo et al. the use of rivaroxaban was associated with an increased rate of thromboembolic and major bleeding events compared with warfarin [132]. Such data were confirmed by the I Trial of Rivaroxaban in AntiPhospholipid Syndrome (TRAPS) study. Rivaroxaban was compared with warfarin in patients with high-risk antiphospholipid syndrome. Data from this study support earlier studies on the superiority of warfarin over rivaroxaban in high-risk patients with antiphospholipid syndrome [133]. Given that the majority of studies examined the effect of NOACs in patients with high-risk antiphospholipid syndrome, further studies could examine the effectiveness in lower-risk antiphospholipid syndrome and thus identify special groups of patients in whom NOAC therapy would be efficient.

### **7.3. Treatment of PE in patients with cancer**

Cancer patients have a four to seven times higher risk of VTE, three times the risk of recurrent VTE, two times the risk of bleeding when using anticoagulants, and ten times the risk of death compared to non-cancer patients. The presence of comorbidities in cancer patients, such as anemia, kidney failure, and reduced mobility can further complicate the treatment of these patients. The first line of therapy for the treatment of PE in patients with cancer is LMWH, and among the NOACs, apixaban and rivaroxaban are also approved for the treatment of these patients. NOACs have shown non-inferiority in reducing VTE recurrence compared to LMWH in the most recent meta-analysis and randomized controlled trials [134]. However, for edoxaban

and dabigatran, a high bleeding risk was found compared to LMWH, especially for patients with gastrointestinal tract tumors [135, 136]. However, the latest evidence on apixaban has suggesting that apixaban is the best alternative, among NOACs, in patients with gastrointestinal cancers due to a lower risk of major bleeding [137]. The ADAM VTE trial [99], which randomized 300 patients to apixaban or LMWH for six months, reported higher recurrent thrombosis in the LMWH group without any difference in the safety outcomes of major bleeding or clinically relevant bleeding [138]. NOACs have frequent interactions with other drugs that raise concerns in the setting of polypharmacotherapy or when administered with drugs that are metabolized by cytochrome P450 such, as drugs used in chemotherapy. For these reasons, LMWH remains the treatment of choice in these patients [8].

#### **7.4 Treatment of PE in elderly patients**

The use of NOACs in elderly patients is associated with a lower risk of bleeding with the same efficacy as VKA. The possibility of using lower doses of NOACs in this population should be considered in view of the presence of many comorbidities [139].

#### **7.5. Treatment of PE in obese patients**

Although the guidelines (ISTH SSC) from 2016 did not advise the use of NOACs in extremely obese people (body mass index [BMI]  $>40$  kg/m<sup>2</sup> or weight  $>120$  kg), new recommendations from 2021. suggest that rivaroxaban and apixaban can be adequate for treatment of VTE in patients with obesity regardless of body weight and BMI [140]. In a meta-analysis by Mohamed Nabil Elshafei et al., which was conducted on five observational studies with 6585 patients with severe obesity (BMI  $\geq 40$  kg/m<sup>2</sup> or weight  $\geq 120$  kg), NOACs (rivaroxaban, apixaban, and dabigatran) were compared with warfarin for the treatment of VTE. There are found similar efficacy of NOACs (recurrent VTE OR 1.07; 0.93–1.23) and a nonsignificant trend toward reduced risk of major bleeding (OR 0.80; 0.54–1.17) [141]. According to a post hoc analysis of the EINSTEIN study including 861 patients with a BMI  $\geq 35$  kg/m<sup>2</sup> there was no significant difference in recurrent VTE in those using rivaroxaban compared with warfarin at even 21 days (2.1% vs. 0.9%, respectively; HR 2.22; 0.68–7.26) nor after 12 months (3% vs. 2.1%, respectively; HR 1.45; 0.62–3.39). There were no differences in the occurrence of major bleeding in the studied groups [142].

#### **7.6. Treatment of chronic thromboembolic pulmonary hypertension (CTEPH)**

Chronic thromboembolic pulmonary hypertension (CTEPH) occurs in about 3% of patients who survive PE. The exact pathophysiology of why CTEPH occurs in fewer patients after PE remains unknown. The following reasons are most often cited: delay in diagnosis, presence of thrombosis in most branches of the pulmonary artery, recurrent symptomatic pulmonary embolism, failure to apply thrombolytic therapy. The diagnosis is established by the existence of a mean pressure in the pulmonary artery above 25 mmHg with the

presence of proven thrombus masses in the branches of the pulmonary artery. Bilateral pulmonary endarterectomy is a curative treatment for CTEPH, but life-long anticoagulant therapy is indicated in most patients.

### **8. Conclusions**

Both the survival of patients with PE and their future quality of life depends on a quick and correct diagnosis of PE and an adequate therapeutic regimen.

Timely administration of thrombolytic therapy and/or embolectomy is life-saving in patients with high-risk pulmonary embolism. Most therapeutic doubts still exist in patients who are hemodynamically stable on admission and who have signs of right heart dysfunction on echocardiography and positive values of biomarkers of both myocardial necrosis and right heart dysfunction. Within that group of patients, how to single out those who are at risk of deterioration and who would benefit unequivocally from the initial application of thrombolytic therapy? For now, the current guidelines of the European Association of Cardiologists for PE do not single out such a group of patients but suggest the initial application of anticoagulant therapy. We are not sure that when hemodynamic destabilization occurs, the application of thrombolytic therapy will always be effective. Also, the adequate dose of thrombolytic therapy in patients with intermediate-high risk embolism is not clearly defined. Defining the dose is very important both for survival and for reducing the risk of resistant pulmonary hypertension, which is a permanent problem and requires prolonged use of anticoagulant therapy. For the same reasons, it is necessary to introduce new studies and recommendations that will clearly define the initial therapeutic regimen in patients with intermediate-high risk pulmonary embolism.

In the treatment of PE, NOACs are recommended before VKA except in patients with antiphospholipid syndrome, mechanical valves, severe mitral stenosis and in pregnancy. Special attention should be paid to interactions of NOACs with other drugs.

Our work could help doctors who encounter EP on a daily basis, not only in understanding this disease but also in the therapeutic approach to it. The duration of anticoagulant therapy is very important considering the high rate of recurrent VTE events.

Our work clearly indicates that PE is a disease that we must think about and where therapeutic challenges still exist.

**Author Contributions:** Conceptualization, methodology, investigation, data curation, writing, and original draft preparation, R.L. and L.D.; original draft preparation, N.R., L.S., M.P. and S.S.; visualization, M.Z.; investigation, J.K. and A.L.; supervision, A.R., J.P. and M.A. All authors have read and agreed to the published version of the manuscript

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare that there is no conflict of interest.

#### References:

1. Heit, J.A.; Spencer, F.A.; White, R.H. The epidemiology of venous thromboembolism. *J Thromb Thrombolysis*. **2016**; 41(1), 3-14.
2. Raskob, G.E.; Angchaisuksiri, P.; Blanco, A.N.; Buller, H.; Gallus, A.; Hunt, B.J.; Hylek, E.; Kakkar, A.; Konstantinides, S.V.; McCumber, M.; et al. ISTH Steering Committee for World Thrombosis Day. Thrombosis: a major contributor to global disease burden. *Arterioscler Thromb Vasc Biol*. **2014**, 34(11), 2363-71.
3. Wendelboe, A.M.; Raskob, G.E.; Global burden of thrombosis: epidemiologic aspects. *Circ Res*. **2016**, 118:1340–1347.
4. Stein, P.D.; Hull, R.D.; Kayali, F.; Ghali, W.A.; Alshab, A.K.; Olson, R.E. Venous thromboembolism according to age: the impact of an aging population. *Arch Intern Med*. **2004**, 164(20), 2260-5.
5. Barco, S.; Mahmoudpour, S.H.; Valerio, L.; Klok, F.A.; Münzel, T.; Middeldorp, S.; Ageno, W.; Cohen, A.T.; Hunt, B.J.; Konstantinides, S.V. Trends in mortality related to pulmonary embolism in the European Region, 2000-15: analysis of vital registration data from the WHO Mortality Database. *Lancet Respir Med*. **2020**, 8(3), 277-287.
6. Khan, F.; Tritschler, T.; Kahn, S.R.; Rodger, M.A. Venous thromboembolism. *Lancet*. **2021**, 398(10294), 64-77.
7. Morrone, D.; Morrone, V. Acute Pulmonary Embolism: Focus on the Clinical Picture. *Korean Circ J*. **2018**, 48(5), 365-381.
8. Konstantinides, S.V.; Meyer, G.; Becattini, C.; Bueno, H.; Geersing, G.J.; Harjola, V.P.; Huisman, M.V.; Humbert, M.; Jennings, C.S.; Jiménez, D. ESC Scientific Document Group. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J*. **2020**, 41(4), 543-603.
9. Righini, M.; Robert-Ebadi, H.; Le Gal, G. Diagnosis of acute pulmonary embolism. *J Thromb Haemost*. **2017**, 15(7), 1251-1261.
10. Nilsson, L.T.; Andersson, T.; Larsen, F.; Lang, I.M.; Liv, P.; Söderberg, S. Dyspnea after pulmonary embolism: a nation-wide population-based case-control study. *Pulm Circ*. **2021**, 11(4), 20458940211046831.

11. Pollack, C.V.; Schreiber, D.; Goldhaber, S.Z.; Slattery, D.; Fanikos, J.; O'Neil, B.J.; Thompson, J.R.; Hiestand, B.; Briese, B.A.; Pendleton, R.C.; et al. Clinical characteristics, management, and outcomes of patients diagnosed with acute pulmonary embolism in the emergency department: initial report of EMPEROR (Multicenter Emergency Medicine Pulmonary Embolism in the Real World Registry). *J Am Coll Cardiol.* **2011**, *57*(6), 700-6.
12. Stein, P.D.; Beemath, A.; Matta, F.; Weg, J.G.; Yusen, R.D.; Hales, C.A.; Hull, R.D.; Leeper, K.V.; Jr Sostman, H.D.; Tapson, V.F.; et al. Clinical characteristics of patients with acute pulmonary embolism: data from PIOPED II. *Am J Med.* **2007**, *120*(10), 871-9.
13. Ji, Q.Y.; Wang, M.F.; Su, C.M.; Yang, Q.F.; Feng, L.F.; Zhao, L.Y.; Fang, S.Y.; Zhao, F.H.; Li, W.M. Clinical symptoms and related risk factors in pulmonary embolism patients and cluster analysis based on these symptoms. *Sci Rep.* **2017**, *7*(1), 14887.
14. Kucher, N.; Rossi, E.; De Rosa, M.; Goldhaber, S.Z. Massive pulmonary embolism. *Circulation.* **2006**, *113*(4), 577-82.
15. Tarbox, A.K.; Swaroop, M. Pulmonary embolism. *Int J Crit Illn Inj Sci.* **2013**, *3*(1), 69-72.
16. Kukla, P.; McIntyre, W.F.; Koracevic, G.; Kutlesic-Kurtovic, D.; Fijorek, K.; Atanaskovic, V.; Krupa, E.; Mirek-Bryniarska, E.; Jastrzębski, M.; Bryniarski, L.; et al. Relation of atrial fibrillation and right-sided cardiac thrombus to outcomes in patients with acute pulmonary embolism. *Am J Cardiol.* **2015**, *115*(6), 825-30.
17. Prandoni, P.; Lensing, A.W.; Prins, M.H.; Ciammaichella, M.; Perlati, M.; Mumoli, N.; Bucherini, E.; Visona, A.; Bova, C.; Imberti, D.; et al. PESIT Investigators. Prevalence of pulmonary embolism among patients hospitalized for syncope. *N Engl J Med.* **2016**, *375*, 1524–1531.
18. Tsuyuki, H.; Yamamoto, N.; Unno, N.; Inuzuka, K.; Sano, M.; Katahashi, K.; Yata, T.; Kayama, T.; Yamanaka, Y.; Endo, Y.; et al. Characteristics and Prognostic Factors of Venous Thromboembolism in Cancer Patients. *Ann Vasc Dis.* **2022** *25*, *15*(2), 107-112.
19. Kearon, C.; de Wit, K.; Parpia, S.; Schulman, S.; Afilalo, M.; Hirsch, A.; Spencer, F.A.; Sharma, S.; D'Aragon, F.; Deshaies, J.F.; et al. PEGeD Study Investigators. Diagnosis of Pulmonary Embolism with d-Dimer Adjusted to Clinical Probability. *N Engl J Med.* **2019**, *381*, 2125-34.
20. Ziegler, T.; Murzik, M.; Schau, A.; Winkler, C.; Fünfstück, R. Interpretation erhöhter D-Dimerkonzentration [Interpretation of increased D-dimer values]. *Hamostaseologie.* **2004** *24*(2),144-6.
21. Shafiee, M.A.; Hosseini, S.F.; Mortazavi, M.; Emami, A.; Mojtahed Zadeh, M.; Moradi, S.; Shaker, P. Anticoagulation therapy in COVID-19 patients with chronic kidney disease. *J Res Med Sci.* **2021**, *30*, 26:63.
22. Righini, M.; Goehring, C.; Bounameaux, H.; Perrier, A. Effects of age on the performance of common diagnostic tests for pulmonary embolism. *Am J Med.* **2000**, *109*:357-61.
23. Righini, M.; Van Es, J.; Den Exter, P.L.; Roy, P.M.; Verschuren, F.; Ghuysen, A.; Rutschmann, O.T.; Sanchez, O.; Jaffrelot, M.; Trinh-Duc, A.; et al. Age-adjusted D-dimer cutoff levels to rule out pulmonary embolism: the ADJUST-PE study. *JAMA.* **2014** *311*(11), 1117-24.
24. van Es, N.; Kraaijpoel, N.; Klok, F.A.; Huisman, M.V.; Den Exter, P.L.; Mos, I.C.; Galapienzo, J.; Büller, H.R.; Bossuyt, P.M. The original and simplified Wells rules and age-adjusted D-dimer testing to rule out pulmonary embolism: an individual patient data meta-analysis. *J Thromb Haemost.* **2017** *15*(4), 678-684.
25. Righini, M.; Robert-Ebadi, H.; Elias, A.; Sanchez, O.; Le Moigne, E.; Schmidt, J.; Le Gall, C.; Cornuz, J.; Aujesky, D.; et al. CT-PE-Pregnancy Group. Diagnosis of Pulmonary Embolism During Pregnancy: A Multicenter Prospective Management Outcome Study. *Ann Intern Med.* **2018**, *169*(11), 766-773.

- 
26. Tromeur, C.; van der Pol, L.M.; Le Roux, P.Y.; Ende-Verhaar, Y.; Salaun, P.Y.; Leroyer, C.; Couturaud, F.; Kroft, L.J.M.; et al. Computed tomography pulmonary angiography versus ventilation-perfusion lung scanning for diagnosing pulmonary embolism during pregnancy: a systematic review and meta-analysis. *Haematologica*. **2019** 104(1), 176-188.
27. Alonso-Martínez, J.L.; Sánchez, F.J.; Echezarreta, M.A. Delay and misdiagnosis in sub-massive and non-massive acute pulmonary embolism. *Eur J Intern Med*. **2010** 21(4), 278-82.
28. van der Hulle, T.; Cheung, W.Y.; Kooij, S.; Beenen, L.F.M.; van Bommel, T.; van Es, J.; Faber, L.M.; Hazelaar, G.M.; Heringhaus, C.; Hofstee, H.; et al. YEARS study group. Simplified diagnostic management of suspected pulmonary embolism (the YEARS study): a prospective, multicentre, cohort study. *Lancet*. **2017** 15, 390(10091), 289-297.
29. Kwok, C.S.; Wong, C.W.; Lovatt, S.; Myint, P.K.; Loke, Y.K. Misdiagnosis of pulmonary embolism and missed pulmonary embolism: A systematic review of the literature, *Health Sciences Review*, **2022**,3,100022.
30. Walen, S.; Damoiseaux, R.A.; Uil, S.M.; van den Berg, J.W.; Diagnostic delay of pulmonary embolism in primary and secondary care: a retrospective cohort study. *Br J Gen Pract*. **2016** 66(647), e444-50.
31. Smith, S.B.; Geske, J.B.; Maguire, J.M.; Zane, N.A.; Carter, R.E.; Morgenthaler, T.I. Early anticoagulation is associated with reduced mortality for acute pulmonary embolism. *Chest*. **2010**, 137(6), 1382-90.
32. Novicic, N.; Dzudovic, B.; Subotic, B.; Shalinger-Martinovic, S.; Obradovic, S. Electrocardiography changes and their significance during treatment of patients with intermediate-high and high-risk pulmonary embolism. *Eur Heart J Acute Cardiovasc Care*. **2020**; 9(4), 271-278.
33. Kurnicka, K.; Lichodziejewska, B.; Goliszek, S.; Dzikowska-Diduch, O.; Zdończyk, O.; Kozłowska, M.; Kostrubiec, M.; Ciurzyński, M.; Palczewski, P.; Grudzka, K.; et al. Echocardiographic Pattern of Acute Pulmonary Embolism: Analysis of 511 Consecutive Patients. *J Am Soc Echocardiogr*. **2016** 29(9), 907-13.
34. McConnell, M.V.; Solomon, S.D.; Rayan, M.E.; Lee, R.T.; Come, P.C.; Goldhaber, S.Z.; Lee, R.T. Regional right ventricular dysfunction detected by echocardiography in acute pulmonary embolism. *Am J Cardiol*, **1996**, 78(4), 469-73.
35. Casazza, F.; Bongarzone, A.; Capozzi, A.; Agostoni, O. Regional right ventricular dysfunction in acute pulmonary embolism and right ventricular infarction. *Eur J Echocardiogr*. **2005**, , 6(1):11-4.
36. Shafiq, Q.; Assaly, R.; Kanjwal, Y.; McConnell Sign in a Patient with Massive Acute Pulmonary Embolism. *Case Rep Cardiol*. **2011**, 2011, 201097.
37. Naeem, K. Floating thrombus in the right heart associated with pulmonary embolism: The role of echocardiography. *Pak J Med Sci*. **2015**; 31(1): 233-5.
38. Bělohávek, J.; Dytrych, V.; Linhart, A. Pulmonary embolism, part I: Epidemiology, risk factors and risk stratification, pathophysiology, clinical presentation, diagnosis and nonthrombotic pulmonary embolism. *Exp Clin Cardiol*. **2013**; 18(2):129-38.
39. Elias, A.; Mallett, S.; Daoud-Elias, M.; Poggi, J.N.; Clarke, M. Prognostic models in acute pulmonary embolism: a systematic review and meta-analysis. *BMJ*. **2016**, 6, e010324.
40. Duffett, L.; Castellucci, L.A.; Forgie, M.A. Pulmonary embolism: update on management and controversies. *BMJ*. **2020**, 370, m2177.
41. Mirambeaux, R.; León, F.; Bikdeli, B.; Morillo, R.; Barrios, D.; Mercedes, E.; Moores, L.; Tapson, V.; Yusen, R.D.; et al. Intermediate-High Risk Pulmonary Embolism. *TH Open*. **2019** , 3(4), e356-e363.
42. Yamamoto, T. Management of patients with high-risk pulmonary embolism: a narrative review. *J Intensive Care*. **2018**, 6, 16.
43. Bledsoe, J.R.; Woller, S.C.; Stevens, S.M.; Aston, V.; Patten, R.; Allen, T.; Horne, B.D.; Dong, L.; et al. Management of Low-Risk Pulmonary Embolism Patients Without Hospitalization: The Low-Risk Pulmonary Embolism Prospective Management Study. *Chest*. **2018** ,154(2), 249-256.

- 
44. Piazza, G. Advanced Management of Intermediate- and High-Risk Pulmonary Embolism: **JACC Focus Seminar. *J Am Coll Cardiol.* 2020, 3, 76(18), 2117-2127.**
45. Konstantinides, S.; Mavromanoli, A.; Hobohm, L. Diagnostik und Therapie der Lungenembolie [Diagnosis and treatment of pulmonary embolism]. *Herz.* **2021**, 46(6), 589-604.
46. Hirsh, J.; Anand, S.S.; Halperin, J.L.; Fuster, V. Mechanism of action and pharmacology of unfractionated heparin. *Arterioscler Thromb Vasc Biol.* **2001**, 21(7), 1094-6.
47. Li, B.; Lu, D.; Chen, Y.; Zhao, M.; Zuo, L. Unfractionated Heparin Promotes Osteoclast Formation in Vitro by Inhibiting Osteoprotegerin Activity. *Int J Mol Sci.* **2016**, 22, 17(4), 613.
48. Capila, I.; Linhardt, R.J. Heparin-protein interactions. *Angew Chem Int Ed Engl.* 2002, 41(3), 391-412.
49. Chan, A.K.; Paredes, N.; Thong, B.; Chindemi, P.; Paes, B.; Berry, L.R.; Monagle, P. Binding of heparin to plasma proteins and endothelial surfaces is inhibited by covalent linkage to antithrombin. *Thromb Haemost.* **2004**, 91(5), 1009-18.
50. Lovecchio, F. Heparin-induced thrombocytopenia. *Clinical Toxicology*, **2014**, 52(6), 579–583.
51. Patriarcheas, V.; Pikoulas, A.; Kostis, M.; Charpidou, A.; Dimakakos, E. Heparin-induced Thrombocytopenia: Pathophysiology, Diagnosis and Management. *Cureus.* **2020**, 12(3), e7385.
52. Warkentin, T.E.; Greinacher, A.; Koster, A.; Lincoff, A.M. Treatment and prevention of heparin-induced thrombocytopenia: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest.* **2008**, 133, 340S-380S.
53. Arepally, G.M. Heparin-induced thrombocytopenia. *Blood.* **2017**, 129(21), 2864-2872.
54. Junqueira, D.R.; Perini, E.; Penholati, R.R.; Carvalho, M.G. Unfractionated heparin versus low molecular weight heparin for avoiding heparin-induced thrombocytopenia in postoperative patients. *Cochrane Database Syst Rev.* **2012**, 9, CD007557.
55. Prandoni, P.; Siragusa, S.; Girolami, B.; Fabris, F. BELZONI Investigators Group. The incidence of heparin-induced thrombocytopenia in medical patients treated with low-molecular-weight heparin: a prospective cohort study. *Blood.* **2005**, 106(9), 3049-54.
56. Leentjens, J.; Peters, M.; Esselink, A.C.; Smulders, Y.; Kramers, C. Initial anticoagulation in patients with pulmonary embolism: thrombolysis, unfractionated heparin, LMWH, fondaparinux, or DOACs? *Br J Clin Pharmacol.* **2017**, 83(11), 2356-2366.
57. Costantino, G.; Ceriani, E.; Rusconi, A.M.; Podda, G.M.; Montano, N.; Duca, P.; Cattaneo, M.; Casazza, G. Bleeding risk during treatment of acute thrombotic events with subcutaneous LMWH compared to intravenous unfractionated heparin; a systematic review. *PLoS One.* **2012**, 7(9), e44553.
58. van Rein, N.; Biedermann, J.S.; van der Meer, F.J.M.; Cannegieter, S.C.; Wiersma, N.; Vermaas, H.W.; Reitsma, P.H.; Kruip, M.J.H.A.; Lijfering, W.M. Major bleeding risks of different low-molecular-weight heparin agents: a cohort study in 12 934 patients treated for acute venous thrombosis. *J Thromb Haemost.* **2017**, 15(7), 1386-1391.
59. Schroeder, M.; Hogwood, J.; Gray, E.; Mulloy, B.; Hackett, A.M.; Johansen, K.B. Protamine neutralisation of low molecular weight heparins and their oligosaccharide components. *Anal Bioanal Chem.* **2011**, 399(2), 763-71.
60. Saheb Sharif-Askari, F.; Syed Sulaiman, S.A.; Saheb Sharif-Askari, N. Anticoagulation Therapy in Patients with Chronic Kidney Disease. *Adv Exp Med Biol.* **2017**, 906, 101-114.
61. Warkentin, T.E. Fondaparinux: does it cause HIT? Can it treat HIT? *Expert Rev Hematol.* **2010** 3(5), 567-81.
62. Linkins, L.A.; Hu, G.; Warkentin, T.E. Systematic review of fondaparinux for heparin-induced thrombocytopenia: When there are no randomized controlled trials. *Res Pract Thromb Haemost.* **2018**, 9, 2(4), 678-683.
63. Cumbo-Nacheli, G.; Samavati, L.; Guzman, J.A. Bioavailability of fondaparinux to critically ill patients. *J Crit Care.* **2011**, 26(4), 342-6.

64. Bijsterveld, N.R.; Moons, A.H.; Boekholdt, S.M.; van Aken, B.E.; Fennema, H.; Peters, R.J.; Meijers, J.C.; Büller, H.R.; Levi, M. Ability of recombinant factor VIIa to reverse the anticoagulant effect of the pentasaccharide fondaparinux in healthy volunteers. *Circulation*. **2002**, *12*, 106(20), 2550-4.
65. Greinacher, A.; Völpel, H.; Janssens, U.; Hach-Wunderle, V.; Kemkes-Matthes, B.; Eichler, P.; Mueller-Velten, H.G.; for the HIT Investigators Group. Recombinant hirudin (lepirudin) provides safe and effective anticoagulation in patients with heparin-induced thrombocytopenia: a prospective study. *Circulation* **1999**; *99*, 73–80.
66. Greinacher, A.; Janssens, U.; Berg, G.; Böck, M.; Kwasny, H.; Kemkes-Matthes, B.; Eichler, P.; Völpel, H.; Pötzsch, B.; Luz, M. for the Heparin-Associated Thrombocytopenia Study (HAT) Investigators. Lepirudin (recombinant hirudin) for parenteral anticoagulation in patients with heparin-induced thrombocytopenia. *Circulation*, **1999**, *100*: 587–93.
67. Lubenow, N.; Eichler, P.; Lietz, T.; Greinacher, A.; Hit Investigators Group. Lepirudin in patients with heparin-induced thrombocytopenia - results of the third prospective study (HAT-3) and a combined analysis of HAT-1, HAT-2, and HAT-3. *J Thromb Haemost*. **2005**, *3*(11), 2428-36.
68. Dager, W.E.; White, R.H. Use of lepirudin in patients with heparin-induced thrombocytopenia and renal failure requiring hemodialysis. *Ann Pharmacother*. **2001**, *35*(7-8):885-90.
69. Bamrah, R.K.; Pham, D.C.; Rana, F. Argatroban in heparin-induced thrombocytopenia: rationale for use and place in therapy. *Ther Adv Chronic Dis*. **2013**, *4*(6), 302-4.
70. Duewell, B.E.; Briski, M.J.; Feih, J.T.; Rinka, J.R.G.; Tawil, J.N. Argatroban Versus Bivalirudin in the Treatment of Suspected or Confirmed Heparin-Induced Thrombocytopenia. *J Pharm Pract*. **2021** *34*(4), 529-534.
71. Farasatinasab, M.; Zarei, B.; Moghtadaei, M.; Nasiripour, S.; Ansarinejad, N.; Zarei, M. Rivaroxaban as an Alternative Agent for Heparin-Induced Thrombocytopenia. *J Clin Pharmacol*. **2020**, *60*(10), 1362-1366.
72. Leidi, A.; Bex, S.; Righini, M.; Berner, A.; Groscurin, O.; Marti, C. Risk Stratification in Patients with Acute Pulmonary Embolism: Current Evidence and Perspectives. *J Clin Med*. **2022**, *11*(9), 2533.
73. Ucar, E.Y. Update on Thrombolytic Therapy in Acute Pulmonary Thromboembolism. *Eurasian J Med*. **2019**, *51*(2), 186-190.
74. Chatterjee, S.; Chakraborty, A.; Weinberg, I.; Kadakia, M.; Wilensky, R.L.; Sardar, P.; Kumbhani, D.J.; Mukherjee, D.; Jaff, M.R.; Giri, J. Thrombolysis for pulmonary embolism and risk of all-cause mortality, major bleeding, and intracranial hemorrhage: a meta-analysis. *JAMA*. 2014 Jun 18; *311*(23):2414-21.
75. Lee, T.; Itagaki, S.; Chiang, Y.P.; Egorova, N.N.; Adams, D.H.; Chikwe, J. Survival and recurrence after acute pulmonary embolism treated with pulmonary embolectomy or thrombolysis in New York State, 1999 to 2013. *J Thorac Cardiovasc Surg*. **2018**, *155*(3):1084-1090.e12.
76. Stein, P.D.; Alnas, M.; Beemath, A.; Patel, N.R. Outcome of pulmonary embolectomy. *Am J Cardiol*. **2007**, *1*, 99(3), 421-3.
77. QiMin, W.; LiangWan, C.; DaoZhong, C.; HanFan, Q.; ZhongYao, H.; XiaoFu, D.; XueShan, H.; Feng, L.; HuaBin, C. Clinical outcomes of acute pulmonary embolectomy as the first-line treatment for massive and submassive pulmonary embolism: a single-centre study in China. *J Cardiothorac Surg*. **2020**, *15*(1), 321.
78. Martinez Licha, C.R.; McCurdy, C.M.; Maldonado, S.M.; Lee, L.S. Current Management of Acute Pulmonary Embolism. *Ann Thorac Cardiovasc Surg*. **2020**, *26*(2), 65-71.
79. Li, K.; Cui, M.; Zhang, K.; Liang, K.; Liu, H.; Zhai, S. Treatment of acute pulmonary embolism using rheolytic thrombectomy. *EuroIntervention*. **2021**, *11*, 17(2), e158-e166.
80. Sharifi, M.; Bay, C.; Skrocki, L.; Rahimi, F.; Mehdipour, M. "MOPETT" Investigators. Moderate pulmonary embolism treated with thrombolysis (from the "MOPETT" Trial). *Am J Cardiol*. **2013**, *111*(2), 273-7.

- 
81. Konstantinides, S.; Geibel, A.; Heusel, G.; Heinrich, F.; Kasper, W. Management Strategies and Prognosis of Pulmonary Embolism-3 Trial Investigators. Heparin plus alteplase compared with heparin alone in patients with submassive pulmonary embolism. *N Engl J Med.* **2002**, 347(15), 1143-50.
82. Meyer, G.; Vicaut, E.; Danays, T.; Agnelli, G.; Becattini, C.; Beyer-Westendorf, J.; Bluhmki, E.; Bouvaist, H.; et al. PEITHO Investigators. Fibrinolysis for patients with intermediate-risk pulmonary embolism. *N Engl J Med.* **2014**, 10; 370(15), 1402-11.
83. Nakamura, S.; Takano, H.; Kubota, Y.; Asai, K.; Shimizu, W. Impact of the efficacy of thrombolytic therapy on the mortality of patients with acute submassive pulmonary embolism: a meta-analysis. *J Thromb Haemost.* **2014**, 12(7),1086-95.
84. Bova, C.; Sanchez, O.; Prandoni, P.; Lankeit, M.; Konstantinides, S.; Vanni, S.; Jiménez, D. Identification of intermediate-risk patients with acute symptomatic pulmonary embolism. *Eur Respir J.* **2014**, 44(3), 694-703.
85. Stevens, S.M.; Woller, S.C.; Kreuziger, L.B.; Bounameaux, H.; Doerschug, K.; Geersing, G.J.; Huisman, M.V.; Kearon, C.; King, C.S.; et al. Antithrombotic Therapy for VTE Disease: Second Update of the CHEST Guideline and Expert Panel Report. *Chest.* **2021**, 160(6), e545-e608.
86. Wang, C.; Zhai, Z.; Yang, Y.; Wu, Q.; Cheng, Z.; Liang, L.; Dai, H.; Huang, K.; Lu, W.; Zhang, Z.; et al. China Venous Thromboembolism (VTE) Study Group. Efficacy and safety of low dose recombinant tissue-type plasminogen activator for the treatment of acute pulmonary thromboembolism: a randomized, multicenter, controlled trial. *Chest.* **2010**, 137(2), 254-62.
87. Alexander, P.; Visagan, S.; Issa, R.; Gorantla, V.R. Thomas SE. Current Trends in the Duration of Anticoagulant Therapy for Venous Thromboembolism: A Systematic Review. *Cureus* **2021**, 13(10), e18992.
88. Chopard, R.; Albertsen, I.E.; Ecarnot, F.; Guth, S.; Besutti, M.; Falvo, N.; Piazza, G.; Meneveau, N. Extended Anticoagulation After Pulmonary Embolism: A Multicenter Observational Cohort Analysis. *J Am Heart Assoc.* **2022**, 11(13), e024425.
89. Ghembaza, A.; Saadoun, D. Management of Antiphospholipid Syndrome. *Biomedicines.* **2020**, 8 (11), 508.
90. Weitz, J.I.; Lensing, A.W.A.; Prins, M.H.; Bauersachs, R.; Beyer-Westendorf, J.; Bounameaux, H.; Brighton, T.A.; Cohen, A.T.; Davidson, B.L.; Decousus, H.; et al. EINSTEIN CHOICE Investigators. Rivaroxaban or Aspirin for Extended Treatment of Venous Thromboembolism. *N Engl J Med.* **2017**, 376(13), 1211-1222.
91. Tritschler, T.; Méan, M.; Limacher, A.; Rodondi, N.; Aujesky, D. Predicting recurrence after unprovoked venous thromboembolism: prospective validation of the updated Vienna Prediction Model. *Blood.* **2015**, 126(16), 1949-51.
92. Rodger, M.A.; Le Gal, G.; Langlois, N.J.; Gin, B.; Mallick, R.; Giulivi, A.; Freedman, M.; Kovacs, M.J.; REVERSE II investigators. "HERDOO2" clinical decision rule to guide duration of anticoagulation in women with unprovoked venous thromboembolism. Can I use any d-Dimer? *Thromb Res.* **2018**, 169, 82-86.
93. Wang, Y.; Bajorek, B. New oral anticoagulants in practice: pharmacological and practical considerations. *Am J Cardiovasc Drugs.* **2014**, 14(3):175-89.
94. Di Minno, A.; Frigerio, B.; Spadarella, G.; Ravani, A.; Sansaro, D.; Amato, M.; Kitzmiller, J.P.; Pepi, M.; Tremoli, E.; Baldassarre, D. Old and new oral anticoagulants: Food, herbal medicines and drug interactions. *Blood Rev.* **2017**, 31(4):193-203.
95. Yan, H., Chen, Y.; Zhu, H.; Huang, W.H.; Cai, X.H.; Li, D.; Lv, Y.J.; Si-Zhao, Zhou, H.H; Luo, F.Y.; Zhang, W.; Li, X. The Relationship Among Intestinal Bacteria, Vitamin K and Response of Vitamin K Antagonist: A Review of Evidence and Potential Mechanism. *Front Med (Lausanne).* **2022**, 9, 829304.
96. Xu, K.; Chan, N.C. Bleeding in patients with atrial fibrillation treated with combined antiplatelet and anticoagulant therapy: time to turn the corner. *Ann Transl Med.* **2019**, 7, S198.
97. Schalekamp, T.; Klungel, O.H.; Souverein, P.C.; de Boer, A. Increased bleeding risk with concurrent use of selective serotonin reuptake inhibitors and coumarins. *Arch Intern Med.* **2008** 168(2),180-5.

98. Sostres, C.; Gargallo, C.J.; Lanas, A. Nonsteroidal anti-inflammatory drugs and upper and lower gastrointestinal mucosal damage. *Arthritis Res Ther.* **2013**, *15*, S3.
99. Wang, M.; Zeraatkar, D.; Obeda, M.; Lee, M.; Garcia, C.; Nguyen, L.; Agarwal, A.; Al-Shalabi, F.; Benipal, H.; Ahmad, A.; Drug-drug interactions with warfarin: A systematic review and meta-analysis. *Br J Clin Pharmacol.* **2021**, *87*(11), 4051-4100.
100. Westergren, T.; Johansson, P.; Molden, E. Probable warfarin-simvastatin interaction. *Ann Pharmacother.* **2007**, *41*(7), 1292-5.
101. Andrus, M.R. Oral anticoagulant drug interactions with statins: case report of fluvastatin and review of the literature. *Pharmacotherapy.* **2004**, *24*(2), 285-90.
102. Engell, A.E.; Svendsen, A.L.O.; Lind, B.S.; Andersen, C.L.; Andersen, J.S.; Willadsen T.G.; Persson, F.; Pottgård, A. Drug-drug interaction between warfarin and statins: A Danish cohort study. *Br J Clin Pharmacol.* **2021**, *87*(2), 694-699.
103. Fan, Y.; Adam, T.J.; McEwan, R.; Pakhomov, S.V.; Melton, G.B.; Zhang, R. Detecting Signals of Interactions Between Warfarin and Dietary Supplements in Electronic Health Records. *Stud Health Technol Inform.* **2017**, *245*, 370-374.
104. Brandin, H.; Myrberg, O.; Rundlöf, T.; Arvidsson, A.K.; Brenning, G. Adverse effects by artificial grapefruit seed extract products in patients on warfarin therapy. *Eur J Clin Pharmacol.* **2007**, *63*(6), 565-70.
105. Monterrey-Rodríguez, J. Interaction between warfarin and mango fruit. *Ann Pharmacother.* **2002**, *36*(5): 940-1.
106. Ulbricht, C.; Chao, W.; Costa, D.; Rusie-Seamon, E.; Weissner, W.; Woods, J. Clinical evidence of herb-drug interactions: a systematic review by the natural standard research collaboration. *Curr Drug Metab.* **2008**, *9*(10), 1063-120.
107. Hamann, G.L.; Campbell, J.D.; George, C.M. Warfarin-cranberry juice interaction. *Ann Pharmacother.* **2011**, *45*(3), e17.
108. Jiang, X.; Williams, K.M.; Liauw, W.S.; Ammit, A.J.; Roufogalis, B.D.; Duke, C.C.; Day, R.O.; McLachlan, A.J. Effect of St John's wort and ginseng on the pharmacokinetics and pharmacodynamics of warfarin in healthy subjects. *Br J Clin Pharmacol.* **2004**, *57*(5), 592-9.
109. Taylor, J.R.; Wilt, V.M. Probable antagonism of warfarin by green tea. *Ann Pharmacother.* **1999**, *33*(4), 426-8.
110. Shikdar, S.; Vashisht, R.; Bhattacharya, P.T. International Normalized Ratio (INR) [Updated 2022 May 8]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; **2022** Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK507707/>
111. Bauer, K.A. Pros and cons of new oral anticoagulants. *Hematology Am Soc Hematol Educ Program.* **2013**, *2013*, 464-70.
112. Cohen, A.T.; Sah, J.; Dhamane, A.D.; Lee, T.; Rosenblatt, L.; Hlavacek, P.; Emir, B.; Keshishian, A.; Yuce, H.; Luo, X. Effectiveness and Safety of Apixaban Versus Warfarin Among Older Patients with Venous Thromboembolism with Different Demographics and Socioeconomic Status. *Adv Ther.* **2021**, *38*(11), 5519-5533.
113. Coleman, C.I.; Bunz, T.J.; Turpie, A.G.G. Effectiveness and safety of rivaroxaban versus warfarin for treatment and prevention of recurrence of venous thromboembolism. *Thromb Haemost.* **2017**, *117*(10), 1841-1847.
114. Alhousani, M., Malik, S.U.; Abu-Hashyeh, A.; Poznanski, N.J.; Al-Hasan, S.; Roth, D.F.; Alsharedi, M.; Mustafa, B. Using oral anticoagulants among chronic kidney disease patients to prevent recurrent venous thromboembolism: A systematic review and meta-analysis. *Thromb Res.* **2021**, *198*, 103-114.
115. Wang, L.; Zhang, D.; Raghavan, N.; Yao, M.; Ma, L.; Frost, C.A.; et al. In vitro assessment of metabolic drug-drug interaction potential of apixaban through cytochrome P450 phenotyping, inhibition, and induction studies. *Drug Metabolism and Disposition.* **2010**, *38*:448-458.
116. Panahi, L.; Udeani, G.; Horseman, M.; Weston, J.; Samuel, N.; Joseph, M.; Mora, A.; Bazan, D. Review of Medical Therapies for the Management of Pulmonary Embolism. *Medicina (Kaunas).* **2021**, *57*(2), 110.

117. Mueck, W.; Stampfuss, J.; Kubitz, D.; Becka, M.; Clinical pharmacokinetic and pharmacodynamic profile of rivaroxaban. *Clin Pharmacokinet.* **2014**, 53(1), 1-16.
118. Mueck, W.; Kubitz, D.; Becka, M. Co-administration of rivaroxaban with drugs that share its elimination pathways: pharmacokinetic effects in healthy subjects. *Br J Clin Pharmacol.* **2013**, 76(3), 455-66.
119. Reilly, P.A.; Lehr, T.; Haertter, S.; Connolly, S.J.; Yusuf, S.; Eikelboom, J.W.; Ezekowitz, M.D.; Nehmiz, G.; Wang, S.; Wallentin, L. RE-LY Investigators. The effect of dabigatran plasma concentrations and patient characteristics on the frequency of ischemic stroke and major bleeding in atrial fibrillation patients: the RE-LY Trial (Randomized Evaluation of Long-Term Anticoagulation Therapy). *J Am Coll Cardiol.* **2014**, 63(4), 321-8.
120. Eikelboom, J.W.; Connolly, S.J.; Brueckmann, M.; Granger, C.B.; Kappetein, A.P.; Mack, M.J.; Blatchford, J.; Devenny K.; Friedman, J.; Guiver, K.; et al. RE-ALIGN Investigators. Dabigatran versus warfarin in patients with mechanical heart valves. *N Engl J Med.* **2013**, 369(13), 1206-14..
121. Eriksson, B.I.; Quinlan, D.J.; Weitz, J.I.; Comparative pharmacodynamics and pharmacokinetics of oral direct thrombin and factor xa inhibitors in development. *Clin Pharmacokinetic* **2009**, 48 (1), pp. 1
122. Walenga, J. M.; Adiguzel, C. Drug and dietary interactions of the new and emerging oral anticoagulants. *International Journal of Clinical Practice* **2010**, 64(7), 956–967.
123. Stangier, J.; Rathgen, K.; Stähle, H.; Gansser, D.; Roth, W. The pharmacokinetics, pharmacodynamics and tolerability of dabigatran etexilate, a new oral direct thrombin inhibitor, in healthy male subjects. *Br J Clin Pharmacol.* **2007**, 64(3), 292-303.
124. Chang, D.N.; Dager, W.E.; Chin, A.I. Removal of dabigatran by hemodialysis. *Am J Kidney Dis.* **2013**, 61(3), 487-9.
125. Hokusai-VTE Investigators, Büller, H.R.; Décousus, H.; Grosso, M.A.; Mercuri, M.; Middeldorp, S.; Prins, M.H.; Raskob, G.E.; Schellong, S.M.; Schwocho, L.; Segers A, et al. Edoxaban versus warfarin for the treatment of symptomatic venous thromboembolism. *N Engl J Med.* **2013**, 369(15), 1406-15.
126. Grześk, G.; Rogowicz, D.; Wołowicz, L.; Ratajczak, A.; Gilewski, W.; Chudzińska, M.; Sinkiewicz, A.; Banach, J. The Clinical Significance of Drug–Food Interactions of Direct Oral Anticoagulants. *Int. J. Mol. Sci.* **2021**, 22, 8531.
127. Bounameaux, H.; Camm, A.J. Edoxaban: An Update on the New Oral Direct Factor Xa Inhibitor. *Drugs* **2014**, 74, 1209–1231.
128. Mendell, J.; Zahir, H.; Matsushima, N.; Noveck, R.; Lee, F.; Chen, S.; Zhang, G.; Shi, M. Drug-drug interaction studies of cardiovascular drugs involving P-glycoprotein, an efflux transporter, on the pharmacokinetics of edoxaban, an oral factor Xa inhibitor. *Am J Cardiovasc Drugs.* **2013**;13(5):331-42.
129. Mainbourg, S.; Cucherat, M.; Provencher, S.; Bertolotti, L.; Nony, P.; Gueyffier, F.; Mismetti, P.; Grange, C.; Durieu, I.; Kilo, R.; et al. META-EMBOL group. Twice- or Once-Daily Dosing of Direct Oral Anticoagulants, a systematic review and meta-analysis. *Thromb Res.* **2021**, 197, 24-32.
130. Rutz, T.; Eggel-Hort B.; Alberio, L.; Bouchardy, J.; Anticoagulation of women with congenital heart disease during pregnancy. *International Journal of Cardiology Congenital Heart Disease* **2021**, 5, 100210.
131. Lameijer, H.; Aalberts, J.J.J.; van Veldhuisen, D.J.; Meijer, K.; Pieper, P.G. Efficacy and safety of direct oral anticoagulants during pregnancy; a systematic literature review. *Thromb Res.* **2018**, 169, 123-127.
132. Pengo, V.; Denas, G.; Zoppellaro, G.; Jose, S.P.; Hoxha, A.; Ruffatti, A.; Andreoli, L.; Tincani, A.; Cenci, C.; Prisco, D.; et al. Rivaroxaban vs warfarin in high-risk patients with antiphospholipid syndrome. *Blood.* **2018**, 132(13), 1365-1371.
133. Pengo, V.; Hoxha, A.; Andreoli, L.; Tincani, A.; Silvestri, E.; Prisco, D.; Fierro, T.; Gresele, P.; Cafolla, A.; De Micheli, V.; et al. Trial of Rivaroxaban in AntiPhospholipid Syndrome (TRAPS): Two-year outcomes after the study closure. *J Thromb Haemost.* **2021** 19(2), 531-535.

- 
134. Camilli, M.; Lombardi, M.; Vescovo, G.M.; Del Buono, M.G.; Galli, M.; Aspromonte, N.; Zoccai, G.B.; Niccoli, G.; Montone, R.A.; Crea, F.; et al. Efficacy and safety of novel oral anticoagulants versus low molecular weight heparin in cancer patients with venous thromboembolism: A systematic review and meta-analysis. *Crit Rev Oncol Hematol.* **2020**, *154*, 103074.
135. Raskob, G.E.; van Es, N.; Verhamme, P.; Carrier, M.; Di Nisio, M.; Garcia, D.; Grosso, M.A.; Kakkar, A.K.; Kovacs, M.J.; Mercuri, M.F.; et al. Hokusai VTE Cancer Investigators. Edoxaban for the Treatment of Cancer-Associated Venous Thromboembolism. *N Engl J Med.* **2018**, *378*(7), 615-624.
136. Clemens, A.; Strack, A.; Noack, H.; Konstantinides, S.; Brueckmann, M.; Lip, G.Y. Anticoagulant-related gastrointestinal bleeding--could this facilitate early detection of benign or malignant gastrointestinal lesions? *Ann Med.* **2014**, *46*(8), 672-8.
137. Mokadem, M.E.; Hassan, A.; Algaby, A.Z. Efficacy and safety of apixaban in patients with active malignancy and acute deep venous thrombosis. *Vascular.* **2021**, *29*(5), 745-750.
138. McBane, R.D.; Wysokinski, W.E.; Le-Rademacher, J.G.; Zemla, T.; Ashrani, A.; Tafur, A.; Perepu, U.; Anderson, D.; Gundabolu, K.; Kuzma, C.; et al. Apixaban and dalteparin in active malignancy-associated venous thromboembolism: The ADAM VTE trial. *J Thromb Haemost.* **2020**, *18*(2), 411-421.
139. Ayalon-Dangur, I.; Vega, Y.; Israel, M.R.; Grossman, A.; Spectre, G.; Shochat, T.; Leibovici, L.; Gafter-Gvili, A. Outcome of Elderly Patients with Venous Thromboembolism Treated with Direct Oral Anticoagulants-A Retrospective Cohort Study. *J Clin Med.* **2021**, *10*(23), 5673.
140. Martin, K.A.; Beyer-Westendorf, J.; Davidson, B.L.; Huisman, M.V.; Sandset, P.M., Moll S. Use of direct oral anticoagulants in patients with obesity for treatment and prevention of venous thromboembolism: Updated communication from the ISTH SSC Subcommittee on Control of Anticoagulation. *J Thromb Haemost.* **2021**, *19*(8), 1874-1882.
141. Elshafei, M.N.; Mohamed, M.F.H.; El-Bardissy, A.; Ahmed, M.B.; Abdallah, I.; Elewa, H.; Danjuma, M. Comparative effectiveness and safety of direct oral anticoagulants compared to warfarin in morbidly obese patients with acute venous thromboembolism: systematic review and a meta-analysis. *J Thromb Thrombolysis.* **2021**, *51*(2), 388-396.
142. Di Nisio, M.; Vedovati, M.C.; Riera-Mestre, A.; Prins, M.H.; Mueller, K.; Cohen, A.T.; Wells, P.S.; Beyer-Westendorf, J.; Prandoni, P.; Bounameaux, H.; et al. Treatment of venous thromboembolism with rivaroxaban in relation to body weight. A sub-analysis of the EINSTEIN DVT/PE studies. *Thromb Haemost.* **2016**, *116*(4), 739-746.