

Article

The Impact of Foehn Wind on Mental Distress among Patients in a Swiss Psychiatric Hospital

Christian A. Mikutta ^{1,2} [†], Charlotte Pervilhac ^{1,3} [†], Hansjörg Znoj³, Andrea Federspiel ^{1,2} and Thomas J. Müller ^{1,2*}

¹ Private Clinic Meiringen, Meiringen, Switzerland

² Translational Research Center, University Hospital of Psychiatry and Psychotherapy, University of Bern, Bern, Switzerland

³ Institute of Psychology, Department of Health Psychology and Behavioural Medicine, University of Bern, Bern, Switzerland

[†] Both authors contributed equally

^{*} Corresponding author: thomas.mueller@privatklinik-meiringen.ch

Abstract: Psychiatric patients are particularly vulnerable to strong weather stimuli, such as foehn, a hot wind that occurs in the alps. However, there is a dearth of research regarding its impact on mental health. This study investigated the impact of foehn wind among patients of a psychiatric hospital located in a foehn area in the Swiss Alps. Analysis was based on anonymized datasets obtained from routine records on admission and discharge, including the Brief Symptom Checklist (BSCL) questionnaire, as well as sociodemographic parameters (age, sex, and diagnosis). Between 2013 and 2020 a total of 10,456 admission days and 10,575 discharge days were recorded. All meteorological data were extracted from the database of the Federal Office of Meteorology and Climatology of Switzerland. We estimated the effect of foehn on the BSCL items using a distributed lag model. Significant differences were found between foehn and non-foehn admissions in obsession-compulsion, Interpersonal Sensitivity, depression, Anxiety, Phobic Anxiety, Paranoid Ideation, and General Severity Index (GSI) ($p < .05$). Our findings suggest that foehn wind events may negatively affect specific mental health parameters in patients. More research is needed to fully understand the impact of foehn's events on mental health.

Keywords: foehn wind; psychopathology; BSCL; mental health; weather; meteorological factors; climate change

1. Introduction

Global climate change has far-reaching implications considering the effects that are already apparent and those that are to be expected in the future (i.e., IPCC [1]). In addition to the diverse ecological, political, socioeconomic, and demographic aspects associated with climate change, the individual effects of climate change on the population's health must also be considered.

Public health research increasingly indicates the crucial and multifaceted impact of climate change on mental health and emotional well-being [2,3]. Recent literature reveals a correlation between experiencing climate change implications, such as extreme weather events or an increase in ambient temperature, and deterioration in mental health [3–9].

Data particularly show negative consequences related to ambient temperature and heat, with an increase in temperature being associated with an increase in mental distress, hospitalizations for mental disorders (including bipolar disorder, schizophrenia, and dementia) [8,10–12], and increased suicide rates [6,8,13–16].

In Switzerland, Bundo and colleagues [17] showed that the hospitalization risk for mental disorders increased linearly by 4.0% for every 10 °C increase in mean daily temperature in a 45-year study period. Higher risks have been identified in patients with developmental disorders and schizophrenia. Furthermore, heatwaves and extremely high

temperatures have been associated with increased mortality in people suffering from mental illness [3,18–22]. Hence, research has shown that strong meteorological stimuli appear to exacerbate symptoms in people with mental disorder.

Despite the growing interest in the relationship between climate change events and mental well-being in public health research, little attention has been paid to weather conditions other than temperature—such as foehn [23]. Foehn wind is a well-known phenomenon in the Swiss Alpine region, generating a warm, dry, and often severe wind on the leeward downwind side of the mountain, descending from the crest into the valley, and reaching hurricane force wind speeds of 117 km/h (maximum exceeding 200 km/h) [24]. This meteorological phenomenon can be found all over the world in high mountain regions, such as “Halny” in Poland or “Chinook” in North America [25].

Although research remains scarce, foehn wind has been associated with a large variety of worsening health conditions, frequently labeled as ‘foehn illness’ [25,26]. These symptoms include migraine and headaches, joint pain, wound pain, sleep disorders, fatigue, concentration problems, anxiety, nausea, paresthesia, an increase in aggressiveness, agitation, depressive states, and suicide attempts [27–34]. Recently, Greve, et al. [25] detected an increase in trauma-related hospital admissions on days with foehn. Some studies have found no effects on health conditions [29,35,36] or even relief of neuropathic pain during Chinook winds [37].

Bundo et al. [38] found that episodes of foehn were associated with a 5% increase in mental health hospitalizations over a 35-year study period in Switzerland. A recent study by Lickiewicz et al. [33] showed that lower barometric pressure and foehn wind increased aggressive behavior in psychiatric hospital patients in Poland. Despite these two studies, to the best of our knowledge, no other study has explored the impact of foehn on the mental distress of psychiatric inpatients—a particularly vulnerable group.

With mental illnesses affecting a billion people worldwide, causing 7% of the global burden of disease as measured in DALYs and 19% of all years lived with disability [39], mental disorders constitute a significant health and socioeconomic burden. The effects of climate change and people affected by mental illnesses are both projected to increase [3,40–44]. Therefore, a profound understanding of the potential risk factors related to mental disorders and the interplay between climate change and mental health must be strengthened to reduce this burden. Specifically, different psychopathological symptoms (and a combination thereof) seem to be appropriate targets for estimating the impact of foehn winds on mental health. However, to date, there are very limited data correlating specific psychopathology symptoms with foehn winds.

To relate self-reported clinically relevant psychological symptoms to the occurrence of foehn wind, we used the Brief Symptom Checklist (BSCL) in a sample of admissions and discharges from a psychiatric hospital located in a well-known foehn area in Switzerland. Specifically, based on the available literature, we expected foehn to increase ratings of aggressiveness and somatization.

2. Materials and Methods

The analysis was based on anonymized individual datasets obtained from routine ANQ records (Swiss National Association for Quality Development in Hospitals and Clinics) on the admission and discharge days of patients from the Private Clinic Meiringen, a psychiatric hospital located in a foehn area in the Swiss Alps. The collected data included the Brief Symptom Checklist (BSCL) questionnaire [45] and sociodemographic parameters (age, sex, and diagnosis). All admission and discharge data between 1st of January 2013 and 31st of December 2020 were included. During this interval a total of 10,456 admission and 10,575 discharge days were recorded. An exemption from the ethics committee of the University of Berne was obtained due to the anonymised data format.

2.1. Brief Symptom Checklist (BSCL)

The Brief Symptom Checklist (BSCL; known as the Brief Symptom Inventory [BSI] in English) is the short German version of the well-validated SCL-90 -R assessment designed

by Derogatis [45–47]. The assessment is an important standardized multidimensional self-report scale designed to assess psychopathological symptoms and psychological distress experienced over the previous seven days. The BSCL is a 53-item instrument that measures nine dimensions: somatization (seven items), obsession-compulsion (six items), interpersonal sensitivity (four items), depression (six items), anxiety (six items), hostility (five items), phobic anxiety (five items), paranoid ideation (five items), and psychoticism (five items). These dimensions are assessed using three global indices: the General Severity Index (GSI), Positive Symptom Distress Index, and Positive Symptom Total. In our study, we used only the GSI as a global index. The original questionnaire reported an internal consistency coefficients range from 0.71 to 0.85 [48]. The test-retest reliability of the GSI after one week was shown to have a reliability of 0.87 [47]. Satisfactory convergent reliability has been reported for the depression subscale with Beck's Depression Inventory [49], as well as a variety of other clinical questionnaires ($r = .36$, $r = .83$) [50].

2.2. Meteorological Data

All meteorological datasets were extracted from the database of the Federal Office of Meteorology and Climatology of Switzerland [51]. The location of the meteorological station is Meiringen (coordinates: $46^{\circ}44' N$ $8^{\circ}10' E$ at 588 m above mean sea level), in climate zone *Cfb* according to the Köppen and Geiger Classification (KGC) [52,53]. The observations from this station comprised the following variables: daily index of the presence of foehn, recorded every 10 min. The Foehn Index classifies the wind as follows: 0 [no foehn], 1 [moderate foehn], and 2 [strong foehn]. Hence, the Foehn index was a categorical variable, with one day represented by 144 Foehn index values.

To account for the fact that the Foehn index values have a different time scale than the BSCL data, we decided to co-register both datasets based on the BSCL data. Therefore, we assigned a single Foehn index value for one day. Specifically, although we had a total of 144 time points for the Foehn index, we compressed this information to one single Foehn index value for the whole day: 0 [no foehn], 1 [moderate foehn], and 2 [strong foehn]. Therefore, we included of 2,922 days of meteorological data. Only a subsample of 259 days had a Foehn index value of 1 or 2, hereafter labeled as 'Foehn events'.

2.3. Definition of the Foehn Group

The BSCL dataset included 10,456 admissions and 10,575 discharges. We identified all BSCL data of patients that fulfilled the following criteria: 1) present admission and discharge records have BSCL data, 2) admission and discharge BSCL records must be unique to one patient (i.e., no double or triple counting of the same patients), 3) Foehn index > 0 (i.e., foehn must be present). Using these criteria, 182 datasets were included in the foehn group.

2.4. Statistical Methods

1. The foehn group analysis was done (a) comparing the John group to the admission group and (b) comparing the foehn group to the discharge group.
2. To evaluate the potential short-term associations between Foehn index events and BSCL values, we first aggregated meteorological (Foehn index) data to BSCL data defined for all patients at admission and discharge. Therefore, the data retained the typical time-series structure. This model assumed stationarity in the association and aimed to control for the long-term effects of the Foehn index data on the BSCL data.
3. We chose a distributed lag model (DLM) [75] to assess the potential impact of days characterized as 'foehn' on patients' psychological distress symptoms, according to the BSCL, on hospitalization and discharge dates. The impact of this meteorological variable could be instantaneous and temporally lagged (e.g., Foehn index days over multiple previous days were used as predictors in a regression model for the specific outcome of interest).

-
4. In the DLM, we included lags of the explanatory time-series as independent variables (i.e., Foehn index days). With this flexible framework, we aimed to compare different parametrizations of the exposure-response association and account for lagged dependencies. We used functions modeling the lag-response dimension with a maximum lag of seven days. Therefore, our model consisted of an unconstrained distributed lag linear model (i.e., assuming a linear association) with a maximum lag of seven days.

Moreover, we used generalized linear models (GLM) with conditional quasi-Poisson regression accounting for overdispersion, with the daily number of BSCL entities for each of that day's patients as the dependent variable. Each day with events (i.e., Foehn index values >0) was matched with the BSCL values for each patient. To evaluate the dependency of both variables, we used a 'finite impulse response' model within the GLM framework. For each Foehn index event, we constructed an event convolution matrix (ECM) that would account for potential lag dependencies. Therefore, this ECM consisted of a 7×7 matrix, with columns representing the time lag. For each row we placed a '1' that was shifted for each column to the right (this is done with the commands: `eye(7)` in MATLAB or `diag(7)` in R). This method does not make any assumptions regarding the exact shape of the response function.

Finally, we computed the paired Student's t-tests against zero to identify significant time periods within each BSCL variable. P-values were corrected using false discovery rate correction (FDR). All statistical analyses were conducted in R (version 4.1.1 [54]) using the package 'dLagM' [of Haydar Demirhan] and using MATLAB 'glmFit' (MathWorks, Natick, Massachusetts, USA). It should be noted that we decided to display not only the potential lag dependencies after the events with Foehn index values of >0 , but also to show the effect of this dependency for four days before this event in Figure 2.

3. Results

3.1. Description of the foehn group and non-foehn group (total admission and total discharge sample)

In the foehn group ($n = 182$), 47% of the patients were female and 52 % were male. There was no significant difference between the sexes ($p = .35$). In the non-foehn admission group ($n = 10,274$) 52.1% of the patients were female, and 47.9% were male. In the non-foehn discharge group ($n = 10,575$) 52% of the patients were female and 48% were male. There was a significant difference in sex ($p < .001$) between the non-foehn's admission and discharge groups. The Chi² test revealed no significant sex difference between the foehn and non-foehn groups ($\text{Chi}^2 = 1.303$, $p = .520$; $\text{Chi}^2 = 1.307$, $p = .521$).

The mean age of the foehn group was 47.9 (SD: 13.2, range: 18-78), while the mean age of the non-foehn group was 47.86 (SD: 15.5, range 16-95) for the admission group and 47.80 (SD: 15.6, range 16-95) for the discharge group. No significant differences were found in the age of the groups ($p > .05$ for all).

Tables 1-3 depict the most common ICD-10 diagnosis codes found in the foehn group and the non-foehn admission and discharge groups, respectively. The Chi² test did not reveal significant differences in the percentages of ICD-10 codes between the foehn group and the admission and discharge samples ($\text{Chi}^2 = 209.4$, $p = .582$; $\text{Chi}^2 = 212.4$, $p = .574$). Supplementary Tables 1 and 2 provide a comprehensive overview of all present ICD-10 diagnoses codes present in the sample.

Table 1. Frequencies of ICD Codes in the foehn group.

ICD Code	Frequency	Percent
F32.1	56	30.8%
F33.1	40	22%
F10.2	19	10.4%
F33.2	13	7.1%

Table 2. Frequencies of ICD Codes in the non-foehn admission group.

ICD Code	Frequency	Percent
F33.1	2001	19.1%
F32.1	1792	17.1%
F33.2	1210	11.6%
F10.2	1079	10.4%

Table 3. Frequencies of ICD Codes in the non-foehn discharge group.

ICD Code	Frequency	Percent
F33.1	1996	19.2%
F32.1	1792	17.1%
F33.2	1210	11.6%
F10.2	1087	10.4%

3.2. Distributed Lag Model of foehn

First, we calculated the GLM with time lags to estimate the impact of foehn on the mental health variables of the BSCL. Figure 1 depicts the unconstrained distributed lag linear model (i.e., with a maximum lag of seven days).




Figure 1. Depicts the distributed lag functional form for the foehn data. Zero equals the foehn day.

3.3. *T-Test of the BSCL variables vs. Zero*

Finally, we identified significant time lags for each BSCL variable. We found significant results on the following scales when comparing the foehn group with the admission sample: Obsession-Compulsion ($t_{\max}=2.67$, $p=.010$; time lag = day 2), interpersonal sensitivity ($t_{\max}=2.59$, $p=.012$, time lag = day 2), depression ($t_{\max}=2.30$, $p=.027$, time lag = day 3), anxiety ($t_{\max}=2.32$, $p=.026$, time lag = day 3), phobic anxiety ($t_{\max}=3.00$, $p=.004$, time lag = day 3), paranoid ideation ($t_{\max}=2.20$, $p=.034$, time lag = day 2), and GSI ($t_{\max}=2.24$, $p=.031$, time lag = day 3). Figure 2 depicts the development of the t-values (vs. zero) as a function of time lags for each BSCL variable with significant results. Supplementary figure 1 shows the non significant BSCL items of the admission group.




Figure 2. Depicts the t-values of the different significant BSCL variables (at admission) as a function of the time lags. Red dots indicate the significant time lags (t-test vs. Zero $p<.05$).

When comparing the foehn group with the discharge sample, we found significant results exclusively during negative time lags in obsession-compulsion ($t_{max}=2.03$, $p=.048$; time lag = day - 2), depression ($t_{max}=2.15$ $p=.037$, time lag = day - 2), and psychotism ($t_{max}=2.29$ $p=.027$, time lag = day - 2) (see Supplementary Figure 2).

4. Discussion

This study investigated the impact of foehn wind events on the mental distress of patients in a psychiatric hospital in Switzerland. Our results suggest that foehn winds might increase distress, negatively impacting the psychological state of patients with mental disorders before obtaining inpatient psychiatric treatment. These patients might be considered unstable, leading to hospitalization in psychiatric clinics. Thus, they appear particularly vulnerable to external stressors, such as meteorological factors, including foehn winds. Moreover, our findings indicate that at the time of admission, foehn episodes might have a negative influence on several specific psychopathological dimensions, namely obsession-compulsion, interpersonal sensitivity, depression, anxiety, phobic anxiety, and paranoid ideation.

The negative health effects of foehn events at admission generally appeared to have a lag effect of two days, replicating similar lag effects of previous studies, which examined the relationship between weather events and mental health outcomes [17].

Our results are generally in line with the – so far limited – literature, indicating that the foehn wind might represent a risk factor for particularly vulnerable groups. Bundo et al. [38] showed that foehn episodes led to a higher risk of mental health hospitalizations in Bern. Lickiewicz et al. [33] demonstrated an increase in aggressive among psychiatric inpatients in Poland relating to foehn winds and barometric pressure changes.

Interestingly, no significant results after the foehn wind on mental distress outcomes were observed on the day of discharge. We theorise, that patients might have acquired some stability in the course of their inpatient treatment and thus are less affected by external environmental stressors, such as meteorological stressors. The shape of the potential dependency of all BSCL psychopathology variables and foehn events on the day of discharge suggest a stochastic. In fact, it resembles a zig-zag curve with arbitrary significant negative time lags for obsession-compulsion, depression, and psychotism. Therefore, the mental distress outcome at admission, measured by the BSCL variables (compared to discharge), may be regarded as a sensitive marker that links subjective symptoms to objective meteorological measures.

The biological mechanisms underlying the influence of meteorological factors on mental health remain largely unknown. Regarding heatwave-related exacerbations, several postulations have been put forth regarding the failure of thermoregulation, leading to an imbalance of serotonin levels, neuroinflammation in the brain, neurotoxicity, and alterations in metabolic functions [55–57]. Additionally, the dysfunctional effects of certain psychiatric medicines inhibiting adaptive thermoregulation of the body, a deficit of weather dependent behavioral adaptations by people suffering from mental disorders, and disrupted sleep have been proposed [10,21,22,57,58]. However, to date, no specific pathophysiological hypothesis concerning foehn winds and mental health outcomes has been proposed.

A strength of this study is that, to the authors knowledge, no other study has explored the relationship between foehn episodes on specific psychopathological dimensions of people affected by mental disorders; thus, for the first time enabling insight into what particular dimensions of psychological distress (e.g. Depression, Anxiety or Somatization) might negatively be affected by foehn winds. Contrary to recent literature, our results suggest that somatization and aggressiveness are not negatively affected by foehn winds. However, in line with Bundo et al. [38] our data suggest that symptoms of depression and anxiety are negatively altered by foehn winds. This is of particular importance since anxiety/fear-related and depressive disorders are the most prevalent mental disorders worldwide [59–61].

Finally, with obsession-compulsion, interpersonal sensitivity, phobic anxiety, and paranoid ideation, further psychopathological entities were connected to foehn's winds. Of note is paranoid ideation, since patients with psychotic symptoms appear to be particularly vulnerable to the influence of environmental factors [17], and so often need hospitalization in situations with symptom exacerbation.

The literature on climate change and mental health has been limited by research gaps and various limitations [3,23,62]. Therefor, the importance of studying climate-vulnerable groups is emphasized [3,23,62]. Another strength of the current study, therefore, lies in the investigation of a climate-vulnerable and marginalized group of psychiatric inpatients [23]. People suffering from mental disorders have been postulated to be particularly at risk for the effects of climate change on physical as well as mental health [3,18,63]. Extreme weather events elevate the risk of symptom exacerbation (cf. Löhmus [57]), suicide [8,13–15], and a three times higher risk of mortality during heatwaves compared to people without mental disorders [18]. Other marginalized groups constitute women (especially pregnant or postnatal women), children, groups with less social support, insufficient medical care, people suffering from financial difficulties, and minoritized ethnic groups [3,23,64–69].

Regarding limitations, the study did not encompass confounding effects, for example, simultaneous exposure to intense foehn wind noise, with chronic noise having been

identified as leading to negative mental health outcomes [70–73], and temporal noise exposure leading to feelings of annoyance [74]. Additionally, the sudden rise in ambient temperature is a phenomenon of foehn and was not considered in our study. However, it should be noted that the rise in mental health hospitalizations related to foehn winds over a 35-year study period has been shown to be independent of temperature [17].

As this study was conducted in a high-income country, looking at inpatient data, the interpretability and generalizability of our findings are somewhat limited [23].

Another limitation lies in the individual-level variability in foehn exposure, which confines the validity and precision of the exposure assessment of our population. Additionally, following a retrospective analysis, the study design did not allow for causal assumptions. In the future prospective, personal exposure studies based on meteorological and geographical high-resolution data should be used in the risk assessment of weather stimuli.

As foehn winds are a frequent phenomenon of the Swiss Alps, affecting a large number of people, our results indicate the significance of future research examining the specifics of foehn winds. As such, it is critical to comprehend if (and why) certain foehn-affected communities and individuals demonstrate resilience in the light of foehn events.

Preliminary clinical prevention efforts and public health implications should consider building climate resilience by, among other things, increasing knowledge related to the climate-change-mental health risks for clinicians, individuals with mental health disorders, as well as their relatives or care takers; using psychoeducational approaches to increase patients' knowledge and insight of their mental disorder, including climate-related risk factors; constructing specific (behavioral) adaptation interventions and evaluating these empirically; and importantly, identifying protective factors and coping mechanisms related to foehn winds, including learning from affected foehn communities [3,23,62].

5. Conclusions

Significant and positive associations were found between foehn wind events and specific psychological distress dimensions among patients at a Swiss psychiatric hospital at the time of admission. However, at the end of inpatient treatment, foehn wind events did not appear to represent a significant risk factor for patients. These results suggest that external meteorological stressors, such as foehn winds, could negatively affect the most vulnerable populations.

Climate-related stressors will increase in the future, and the number of people affected by mental health issues is projected to increase. Thus, a profound understanding of the possible risk as well as protective factors related to mental health disorders and the interplay between climate change and mental health must be strengthened to reduce the burden of mental health disorders in the future.

Supplementary Materials: Supplementary Figures 1 and 2, Supplementary Table 1

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used “Conceptualization, CP, CM, AF and TJM.; methodology, CM, AF.; software, AF.; validation, CM.; formal analysis, AF.; data curation, CP, CM, AF. TJM.; writing—original draft preparation, CM, CP, AF, HZ, TJM.; writing—review and editing, CM CP.; visualization, AF.; supervision, HZ & TJM.; All authors have read and agreed to the published version of the manuscript.”

Funding: This research received no external funding

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and an exemption by the Ethics Committee of the University of Berne was obtained (protocol code: Req-2021-01387).

Informed Consent Statement: Patient consent was waived due to the retrospective nature of this study.

Data Availability Statement: Data is available from the authors with reasonable request

Conflicts of Interest: The authors declare no conflict of interest.

Reference

1. Change, I.P.o.C. Global Warming of 1.5°C: an IPCC Special Report on the Impacts of Global Warming of 1.5°C above Preindustrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty; Intergovernmental Panel on Climate Change, 2018.
2. Bourque, F.; Wilcox, A. C. Climate Change: The Next Challenge for Public Mental Health? *Int. Rev. Psychiatry*. **2014**, *26*, 415–422. DOI:10.3109/09540261.2014.925851.
3. Lawrence, E.; Thompson, R.; Fontana, G.; Jennings, N. **2021**. *The Impact of Climate Change on Mental Health and Emotional Well-being: Current Evidence and Implications for Policy and Practice*. Available at: <https://www.imperial.ac.uk/grantham/publications/the-impact-of-climate-change-on-mentalhealth-and-emotional-wellbeing-current-evidence-and-implications-for-policy-and-practice.php>.
4. Cruz, J.; White, P.C.L.; Bell, A.; Coventry, P.A. Effect of extreme weather events on mental health: A narrative synthesis and meta-analysis for the UK. *Int. J. Environ. Res. Public Health*. **2020**, *17*, 8581. DOI:10.3390/ijerph17228581.
5. Ding, N.; Berry, H.L.; Bennett, C.M. The importance of humidity in the relationship between heat and population mental health: Evidence from Australia. *PLOS ONE*. **2016**, *11*, e0164190. DOI:10.1371/journal.pone.0164190.
6. Frangione, B.; Rodríguez Villamizar, L.A.R.; Lang, J.J.; Colman, I.; Lavigne, E.; Peters, C.; Anisman, H.; Villeneuve, P.J.; Villeneuve, P.J. Short-term changes in meteorological conditions and suicide: a systematic review and meta-analysis. *Environ. Res.* **2022**, *207*, 112230. DOI:10.1016/j.envres.2021.112230.
7. Noelke, C.; McGovern, M.; Corsi, D.J.; Jimenez, M.P.; Stern, A.; Wing, I.S.; Berkman, L. Increasing ambient temperature reduces emotional well-being. *Environ. Res.* **2016**, *151*, 124–129. DOI:10.1016/j.envres.2016.06.045.
8. Thompson, R.; Hornigold, R.; Page, L.; Waite, T. Associations between high ambient temperatures and heat waves with mental health outcomes: A systematic review. *Public Health*. **2018**, *161*, 171–191. DOI:10.1016/j.puhe.2018.06.008.
9. Trombley, J.; Chalupka, S.; Anderko, L. Climate Change and Mental Health. *Am. J. Nurs.* **2017**, *117*, 44–52. DOI:10.1097/01.NAJ.0000515232.51795.fa.
10. Hansen, A.; Bi, P.; Nitschke, M.; Ryan, P.; Pisaniello, D.; Tucker, G. Effect of heat waves on mental health in a temperate Australian city. *Environ. Health Perspect.* **2008**, *116*, 1369–1375. DOI:10.1289/ehp.11339.
11. Vaneckova, P.; Bambrick, H. Cause-specific Hospital Admissions on Hot Days in Sydney, Australia. *PLOS ONE*. **2013**, *8*, e55459. DOI:10.1371/journal.pone.0055459.
12. Wang, X.; Lavigne, E.; Ouellette-Kuntz, H.; Chen, B.E. Acute impacts of extreme temperature exposure on emergency room admissions related to mental and behavioral disorders in Toronto, Canada. *J. Affect. Disord.* **2014**, *155*, 154–161. DOI:10.1016/j.jad.2013.10.042.
13. Burke, M.; González, F.; Baylis, P.; Heft-Neal, S.; Baysan, C.; Basu, S.; Hsiang, S. Higher temperatures increase suicide rates in the United States and Mexico. *Nat. Clim. Change*. **2018**, *8*, 723–729. DOI:10.1038/s41558-018-0222-x.
14. Pervilhac, C.; Schoilew, K.; Znoj, H.; Müller, T.J. Wetter und Suizid[Weather and Suicide: Association between Meteorological Variables and Suicidal Behavior-a Systematic Qualitative Review Article]. *Nervenarzt*. **2020**, *91*, 227–232. DOI:10.1007/s00115-019-00795-x.
15. Cheng, S.; Plouffe, R.; Nanos, S.M.; Qamar, M.; Fisman, D.N.; Soucy, J.-P.R. The Effect of Average Temperature on Suicide Rates in Five Urban California Counties, 1999–2019: An Ecological Time Series Analysis. *BMC Public Health*. **2021**, *21*, 974. DOI:10.1186/s12889-021-11001-6.
16. Bär, S.; Bundo, M.; de Schrijver, E.; Müller, T.J.; Vicedo-Cabrera, A.M. Suicides and Ambient Temperature in Switzerland: A Nationwide Time-Series Analysis. *Swiss Med. Wkly*. **2022**, *152*, w30115. DOI:10.4414/smw.2022.w30115.
17. Bundo, M.; de Schrijver, E.; Federspiel, A.; Toreti, A.; Xoplaki, E.; Luterbacher, J.; Franco, O.H.; Müller, T.; Vicedo-Cabrera, A.M.; Müller, T.; Vicedo-Cabrera, A.M. Ambient Temperature and Mental Health Hospitalizations in Bern, Switzerland: A 45-Year Time-Series Study. *PLOS ONE*. **2021**, *16*, e0258302. DOI:10.1371/journal.pone.0258302.
18. Bouchama, A.; Dehbi, M.; Mohamed, G.; Matthies, F.; Shoukri, M.; Menne, B. Prognostic Factors in Heat Wave Related Deaths: a Meta-Analysis. *Arch. Intern. Med.* **2007**, *167*, 2170–2176. DOI:10.1001/archinte.167.20.ira70009.
19. Hoffmann, M.S.; Oliveira, L.M.; Lobato, M.I.R.; Belmonte-de-Abreu, P. Heat stroke during long-term clozapine treatment: should we be concerned about hot weather? *Trends Psychiatry Psychother.* **2016**, *38*, 56–59. DOI:10.1590/2237-6089-2015-0066.
20. Mann, S.C.; Boger, W.P. Psychotropic drugs, summer heat and humidity, and hyperpyrexia: A danger restored. *Am. J. Psychiatry*. **1978**, *135*, 1097–1100. DOI:10.1176/ajp.135.9.1097.
21. Martin-Latry, K.; Goumy, M.P.; Latry, P.; Gabinski, C.; Bégaud, B.; Faure, I.; Verdoux, H. Psychotropic drug use and risk of heat-related hospitalization. *Eur. Psychiatry*. **2007**, *22*, 335–338. DOI:10.1016/j.eurpsy.2007.03.007.
22. Stöllberger, C.; Lutz, W.; Finsterer, J. Heat-related side-effects of neurological and non-neurological medication may increase heatwave fatalities. *Eur. J. Neurol.* **2009**, *16*, 879–882. DOI:10.1111/j.1468-1331.2009.02581.x.
23. Charlson F., Ali S., Benmarhnia T., Pearl M., Massazza A.; Augustinavicius J., Scott J.G.. Climate change and mental health: a scoping review. *Int. J. Environ. Res. Public Health*. **2021**, *18*, 4486. DOI:10.3390/ijerph18094486.
24. WMO, Geneva & OMM Geneva. *International Meteorological Vocabulary*, 1966.

25. Greve, F.; Kanz, K.-G.; Zyskowski, M.; von Matthey, F.; Biberthaler, P.; Muthers, S.,... Huber-Wagner, S. Influence of Foehn Winds on the Incidence of Severe Injuries in Southern Bavaria: An Analysis of the Trauma Register DGU®. *BMC Musculoskelet. Disord.* **2020**, *21*, 1–9.
26. Richner, H.; Hächler, P. Understanding and forecasting alpine foehn. Ecol. In. *Springer Atmospheric Sciences* Springer. **2013**, (219–260). DOI:10.1007/978-94-007-4098-3_4.
27. Amiel M. Geography of Health. *MD*. 1963, *7*, 109–114.
28. Berg, H. Die Wirkung des Föhns auf den menschlichen Organismus. *Geofisica Pura e Applicata*. **1950**, *17*, 104–111. DOI:10.1007/BF02018347.
29. Fletcher, R.J. “Föhn illness” and Human Biometeorology in the Chinook Area of Canada. *Int. J. Biometeorol.* **1988**, *32*, 168–175. DOI:10.1007/BF01045275.
30. Kals, W. Y. *Your Health, Our Moods, and the Weather* (Vol. vol. 1) Doubleday Publishing: New York, NY, U.S.A., **1982**.
31. Koszewska, I.; Walawender, E.; Baran, A.; Zieliński, J.; Ustrnul, Z. Foehn Wind as a Seasonal Suicide Risk Factor in a Mountain Region. *Psychiatr. Psychol. Klin.* **2019**, *19*, 48–53. DOI:10.15557/PiPK.2019.0007.
32. Koszewska, I.; Walawender, E.; Baran, A.; Zieliński, J.; Ustrnul, Z. The influence of foehn winds (Halny) on the occurrence of suicide in the Tatra Mountains, 1999–2014. *European Psychiatry*. **2016**, *33*(S 1), S597–S597. DOI:10.1016/j.eurpsy.2016.01.2230.
33. Lickiewicz, J.; Piotrowicz, K.; Hughes, P. P.; Makara-Studzińska, M. Weather and Aggressive Behavior among Patients in Psychiatric Hospitals-An Exploratory Study. *Int. J. Environ. Res. Public Health*. **2020**, *17*, 9121. DOI:10.3390/ijerph17239121.
34. Rosen, S. *Weathering. How the Atmosphere Conditions Your Body, Your Mind, Your Moods, and Your Health*; M. Evans & Company: New York, **1979**.
35. Field, T.S.; Hill, M.D., Weather, Chinook, and stroke occurrence. *Stroke*. **2002**, *33*, 1751–1757. DOI:10.1161/01.STR.0000020384.92499.59.
36. Maciejczak, A.; Guzik, A.; Wolan-Nieroda, A.; Wójcik, M.; Pop, T. Impact of Foehn Wind and Related Environmental Variables on the Incidence of Cardiac Events. *Int. J. Environ. Res. Public Health*. **2020**, *17*, 2638. DOI:10.3390/ijerph17082638.
37. Ngan, S.; Toth, C. Influence of Chinook winds and other weather patterns on neuropathic pain. *Pain Med.* **2011**, *12*, 1523–1531. DOI:10.1111/j.1526-4637.2011.01227.x.
38. Bundo, M.; de Schrijver, E.; Federspiel, A.; Luterbacher, J.; Franco, O.; Müller, T.; Vicedo-Cabrera, A. Impact of Foehn Winds on Mental Health Hospitalizations in Bern, Switzerland: A 35-Year Time-Series Study. *ISEE Conference Abstracts*. Paper presented at the International Society for Environmental Epidemiology (ISEE) Annual Conference, Virtual Conference. 2020, **2020**. DOI:10.1289/isee.2020.virtual.P-0356.
39. Rehm, J.; Shield, K. D. Global burden of disease and the impact of mental and addictive disorders. *Curr. Psychiatry Rep.* **2019**, *21*, 10. DOI:10.1007/s11920-019-0997-0.
40. Bundesamt für Umwelt BAFU; Bericht des Bundesrates. **2018**. *Umwelt Schweiz 2018*. Retrieved from <https://www.bafu.admin.ch/bafu/de/home/zustand/publikationen-zum-umweltzustand/umwelt-schweiz-2018.html>.
41. Coverdale, J.; Balon, R.; Beresin, E.V.; Brenner, A.M.; Guerrero, A.P.S.; Louie, A.K.; Roberts, L.W. Climate Change: A Call to Action for the Psychiatric Profession. In. *Acad. Psychiatry*; Springer. 2018, *42*. DOI:10.1007/s40596-018-0885-7.
42. Guerreiro, S.B., Dawson, R.J., Kilsby, C., Lewis, E., Ford, A. Future heat wWaves, droughts, and floods in 571 European cities. *Environ. Res. Lett.* **2018**, *13*, 034009. DOI:10.1088/1748-9326/aaaad3.
43. Nie, H.; Qin, T.; Yang, H.; Chen, J.; He, S.; Lv, Z.; Shen, Z. Trends Analysis of Temperature and Precipitation Extreme during Winter Wheat Growth Period in the Major Winter Wheat Planting Area of China. *Atmosphere*. **2019**, *10*, 240. DOI:10.3390/atmos10050240.
44. Watts, N.; Arnell, N.; Ayeb-Karlsson, S.; Belesova, K.; Boykoff, M.; Byass, P.; Cai, W.; Campbell-Lendrum, D.; Capstick, S.; et al. The 2019 report of the Lancet Countdown on Health and Climate Change: Ensuring that the health of a child born today is not defined by a changing climate. *Lancet*. **2019**, *394*, 1836–1878. DOI:10.1016/S0140-6736(19)32596-6.
45. Franke, G.H. *SCL-90®-S. Symptom-Checklist-90®-Standard-Manual*; Hogrefe: Göttingen, 2014.
46. Derogatis L. Brief Symptom Inventory (Baltimore, Clinical Psychometric Research). *Psychopathology*. **1975**, *27*, 14–18.
47. Franke, G.H. Symptom-Checklist-90®-Standard. Paper presented at the Diagnostische Verfahren in der Psychotherapie, 2016.
48. Franke, G.H., *Brief Symptom Inventory, German version*; Beltz: Göttingen, Germany, 2000.
49. Hautzinger, M.; Bailer, M.; Worall, H.; Keller, F. *BDI Beck-Depressions-Inventory*; Pearson Assessment: Frankfurt and Information GmbH, 1995.
50. Beck, A.; Steer, R. 2015. BHS. Beck Hopelessness Scale. Beck-Hoffnungslosigkeits-Skala. *Deutsche Fassung*; Pearson: Frankfurt am main.
51. MeteoSwiss. Available online: www.meteoswiss.ch, 2022.
52. Köppen, W. Die Wärmezonen der Erde, nach der Dauer der heissen, gemässigten und kalten Zeit und nach der Wirkung der Wärme auf die organische Welt betrachtet. *Meteorol. Z.* **1884**, *1*, 5–226.
53. Köppen, W. *Das geographische System der Klimate: Mit 14 Textfiguren*; Borntraeger: Berlin, 1936.
54. R Core Team. *R: A Language and Environment for Statistical Computing*, 2013.
55. Chong, T.W.; Castle, D.J. Layer upon layer: thermoregulation in schizophrenia. *Schizophr. Res.* **2004**, *69*, 149–157. DOI:10.1016/s0920-9964(03)00222-6.
56. Heo S., Lee W., Bell M.L.. Suicide and associations with air pollution and ambient temperature: a systematic review and meta-analysis. *Int. J. Environ. Res. Public Health*. **2021**, *18*, 7699. DOI:10.3390/ijerph18147699.

-
57. Lõhmus M. Possible Biological Mechanisms Linking Mental Health and Heat-A Contemplative Review. *Int. J. Environ. Res. Public Health.* **2018**, *15*, 1515. DOI:10.3390/ijerph15071515.
58. Martinez M., Devenport L., Saussy J., Martinez J. Drug-associated heat stroke. *South. Med. J.* **2002**, *95*, 799–802.
59. Gutiérrez-Rojas, L.; Porras-Segovia, A.; Dunne, H.; Andrade-González, N.; Cervilla, J. A. Prevalence and correlates of major depressive disorder: A systematicSystematic review. *Braz. J. Psychiatry.* **2020**, *42*, 657–672. DOI:10.1590/1516-4446-2020-0650.
60. Scott, K.M.; de Jonge P., Stein D.J., Kessler R.C. *Mental Disorders around the World: Facts and Figures from the WHO World Mental Health Surveys*; Cambridge University Press, 2018.
61. Solmi, M.; Radua, J.; Olivola, M.; Croce, E.; Soardo, L.; Salazar de Pablo, G.; Il Shin, J.; Kirkbride, J.B.; Jones, P.; Kim, J.H.; et al. Age at onset of mental disorders worldwide: A large-scale meta-analysis of 192 epidemiological studies. *Mol. Psychiatry.* **2022**, *27*, 281–295. DOI:10.1038/s41380-021-01161-7.
62. Charlson F., Ali S., Augustinavicius J., Benmarhnia T., Birch S.; Clayton S., et al. Global priorities for climate change and mental health research. *Environ. Int.* **2022**, *158*, 106984. DOI:10.1016/j.envint.2021.106984.
63. Kenny, G.P.; Yardley, J.; Brown, C.; Sigal, R.J.; Jay, O. Heat Stress in Older Individuals and Patients with Common Chronic Diseases. *CMAJ.* **2010**, *182*, 1053–1060. DOI:10.1503/cmaj.081050.
64. Mills M.A., Edmondson D., Park C.L. Trauma and stress response among hurricane Katrina evacuees. *Am. J. Public Health.* **2007**, *97*(supplement 1)(Supplement1), S116–S123. DOI:10.2105/AJPH.2006.086678.
65. Obrien, L.V.; Berry, H.L.; Coleman, C.; Hanigan, I.C. Drought as a mental health exposure. *Environ. Res.* **2014**, *131*, 181–187. DOI:10.1016/j.envres.2014.03.014.
66. Perceval, M.; Kõlves, K.; Ross, V.; Reddy, P.; De Leo, D. Environmental factors and suicide among Australian farmers: A qualitative study. *Arch. Environ. Occup. Health.* **2019**, *74*, 279–286. DOI:10.1080/19338244.2018.1453774.
67. Ruskin J., Rasul R., Schneider S., Bevilacqua K., Taioli E., Schwartz R.M.. Lack of access to medical care during hurricane sandy and mental health symptoms. *Prev. Med. Rep.* **2018**, *10*, 363–369. DOI:10.1016/j.pmedr.2018.04.014.
68. Sirey, J. A., Berman, J., Halkett, A., Giunta, N., Kerrigan, J., Raeifar, E., Artis, A., Banerjee, S., Rau, P.J., Banerjee, S., Rau, P.J., Storm impact and depression among older adults living in hurricane sandy-affected areas. *Disaster Med. Public Health Prep.* **2017**, *11*, 97–109. DOI:10.1017/dmp.2016.189.
69. Picou, J.S.; Hudson, Hurricane Katrina, Mental Health: A Research Note on Mississippi Gulf Coast Residents. *Sociol. Inq.* **2010**, *80*, 513–524. DOI:10.1111/j.1475-682x.2010.00345.x.
70. Clark, C.; Paunovic, K. WHO Health Organization Environmental Noise Guidelines for the European Region: A systematicSystematic review on environmental noise and quality of life, well-being, and mental health. *Int. J. Environ. Res. Public Health.* **2018**, *15*, 2400. DOI:10.3390/ijerph15112400.
71. Hegewald, J., Schubert, M., Freiberg, A., Romero Starke, K., Augustin, F., Riedel-Heller, S.G., Zeeb, H., Seidler, A., Seidler, A. Traffic noise and mental health: a systematic review and meta-analysis. *Int. J. Environ. Res. Public Health.* **2020**, *17*, 6175. DOI:10.3390/ijerph17176175.
72. Monazzam Esmaelpour, M.R.; Zakerian, S.A.; Abbasi, M.; Balochkhaneh, F.; Kordmiri, S.H. Investigating the effect of noise exposure on mental disorders and the work ability index among industrial workers. *Noise Vib. Worldwide.* **2022**, *53*, 3–11. DOI:10.1177/09574565211052690.
73. Zare Sakhvidi, F.; Zare Sakhvidi, M. J.; Mehrparvar, A. H.; Dzhambov, A. M. Environmental Noise Exposure and Neurodevelopmental and Mental Health Problems in Children: A Systematic Review. *Curr. Environ. Health Rep.* **2018**, *5*, 365–374. DOI:10.1007/s40572-018-0208-x.
74. Brink, M.; Schäffer, B.; Vienneau, D.; Foraster, M.; Pieren, R.; Eze, I.C.; Cajochen, C.; Probst-Hensch, N.; Röösli, M.; Wunderli, J.M.; Probst-Hensch, N.; Röösli, M.; Wunderli, J.M. A Survey on Exposure-Response Relationships for Road, Rail, and Aircraft Noise Annoyance: Differences between Continuous and Intermittent Noise. *Environ. Int.* **2019**, *125*, 277–290. DOI:10.1016/j.envint.2019.01.043.
75. Granger, C.W.J.; Newbold, P. Spurious Regressions in Econometrics. *J. Econ.* **1974**, *2*, 111–120. DOI:10.1016/0304-4076(74)90034-7.