
Review

Buffalo on the Edge: Factors Affecting Historical Distribution and Restoration of *Bison bison* in the Western Cordillera, North America

Jonathan J. Farr^{1,*} and Clifford A. White²

¹ Department of Biological Sciences, University of Alberta

² CW and Associates, 1018 14 St. Canmore, AB. T1W 1V5 2; cliffawhite@gmail.com

* Correspondence: jfarr@ualberta.ca

Abstract: The historic western edge of bison (*Bison bison*) range and the ecological processes that caused its formation are frequently debated with important implications for bison restoration across North America. We test the hypothesis that a combination of bottom-up habitat suitability and top-down harvest pressure from humans were important processes in forming the western edge of bison distribution. Using 9,384 historical journal observations from 1691 – 1928, we employ MaxEnt ecological niche modelling to identify suitable bison habitat across the Western Cordillera from bottom-up climatic, land cover, and topographic factors. We then use mixed-effect logistic regression to test if bison occurrence in journal records can be in part explained by the abundance of Indigenous humans, wolves, or grizzly bears, in addition to MaxEnt-derived habitat suitability. We find support for our hypothesis because of the limited suitable habitat in the Rocky Mountains that likely prevented westward bison dispersal from core habitat, and there was a negative relationship between bison occurrence and human harvest pressure. On this basis, we propose that intensive human harvest from large populations in the Western Cordillera, subsidized by other wildlife, salmon, and vegetation resources, is an underappreciated socioecological process that needs to be restored alongside bison populations. Co-managing bison with Indigenous people will also mitigate the adverse effects of overabundant bison and maximize the ecological and cultural benefits of bison restoration.

Keywords: bison; restoration; socio-ecological processes; Indigenous harvest; maximum entropy modelling

1. Introduction

In the period ~CE 1750 to ~ CE 1880 Eurasian colonization of western North America caused one of the greatest near extinctions ever documented. Within a century, *Bison bison*, commonly called the buffalo, once numbering in the millions across the grasslands and woodlands of Great Plains were reduced to less than 1000 animals [1]. Humans' role in the bison overkill are well-documented [2], and researchers mapped the spatial pattern of extirpation [3,4] within years of its occurrence (Figure 1). The political ramifications of this slaughter of the continent's largest land-mammal helped stimulate conservation by the national governments of both United States and Canada, including designating wildlife a public resource [5,6], and withholding massive areas as public lands mandated for the sustainable use of wildlife and plant resources [7–9].




Figure 1. Range contractions of the American bison during the period pre-1800 to 1875 as mapped by Joel Allen of the Kentucky Geological Survey in 1876 [3] showing locations of Western Cordillera restoration projects described in this paper. The figure shows the extreme edge of historic bison range before 1800 (blue), range contraction 1800 to 1825 (pink), 1825 to 1850 (green), 1850 to 1875 (yellow), and the range in 1875 (orange). The dotted line delineates where the mountains of the Western Cordillera meet the Great Plains.

Today, the parks, protected areas, and public lands of North America's Western Cordillera, stretching from the Rocky Mountains westward to the Pacific from Yellowstone to the Yukon, provide one of the greatest opportunities for restoration and conservation of bison and many other species. Yellowstone in the United States and Banff in Canada are birthplaces of the world's first national park systems, and both played roles in the initial efforts to save the American bison from extinction. Moreover, these parks are the cores of an area that now constitutes one of the planet's largest areas of public lands—a network that provides ecological connectivity from along the Cordillera from Yellowstone to the Yukon [10]. These parks and other public lands are also the homelands of Indigenous peoples and provide ecosystem services, natural resources and recreational opportunities for millions of local and regional residents. Furthermore, they are heavily visited by domestic and international travellers, providing an opportunity to observe, hunt, and study in the wilds of world-renowned scenery.

However, many of the parks and protected areas of the Western Cordillera are located on the historic edge of core bison distribution, and bison restoration in this area requires consideration of the factors creating this dynamic. These factors are still debated and several researchers have provided hypotheses for the scarcity of bison west of the Rocky Mountain [11–14], including lower forage quality, heavy winter snows, discontinuous habitat, and high harvest-caused mortality from large populations Indigenous

peoples. These explanations are not mutually exclusive and likely interacted to limit low-density bison populations in this area. Empirical support for these hypotheses is limited, although one recent study used archeological data and bioclimatic models to conclude that during the late Holocene a source-sink dynamic likely existed between abundant bison populations in the great plains and limited populations in the Western Cordillera [13]. Although they did not explicitly examine the role of Indigenous hunting, these authors proposed that both marginal habitat and hunting pressure may have created this dynamic.

There is extensive evidence of intensive human bison harvest in the Rocky Mountains [14–17], and if this process affected bison abundance in the Western Cordillera then bison restoration projects need to consider the effects of two interacting phenomena. First is the potential historic socio-ecological system resulting from human involvement in creating low-density areas where bison abundance was limited. In these areas with historically no, or very few bison, vegetation and other ecosystem attributes may be potentially ill-adapted to high numbers of a large and gregarious herbivore [18]. The second phenomenon is socio-political. The early establishment and management of national parks and other public lands followed a socially and politically-driven tradition of removal of Indigenous and other human influences [19,20], and in parks, ongoing programs of “natural regulation” to allow wildlife populations to rise and fall unimpeded by human influences [21,22]. These decisions effectively eliminated any socio-ecological roles of Indigenous humans. In situations where harvest might have maintained low bison densities, creating few ecological impacts, the recent and relatively novel human perspective of modern management may allow high numbers of bison, elk and other herbivores that greatly impact ecosystems [23–26].

In this paper we further inform bison restoration by examining the role that human harvest of bison may have historically played in creating socio-ecological systems that contained limited bison populations in the Western Cordillera. Using a large database of historic journal observations from northwest North America, we empirically test the hypothesis that bison populations in the Western Cordillera were limited by both low bottom-up habitat suitability and high top-down human harvest. First, we present historic information on bison and its potential primary predators [27–29]: humans, wolves (*Canis lupus*) and grizzly bears (*Ursus arctos horribilis*), using abundance and distribution data from a biome-scale record of the first-person journals of early non-Indigenous travellers. Second, we conduct analyses of this data to test how bottom-up habitat suitability based on Maxent modelling and mixed-effect modelling of top-down processes, namely wolf and bear predation and human harvest, could have limited bison occurrence in the Western Cordillera. Third, we review the demographics and range areas of bison restored on the edge, or outside of historic range, to further test whether the growth of these populations is limited by their location. We then discuss restoration of bison in accordance with bottom-up and top-down ecological processes, and long-term human influences.

2. Materials and Methods

2.1. First Person Journal Observations

Our analysis required specific observations on bison locations and abundance, human cultural practices, and the abundance of other predator species present in historic landscapes. Historic bison, human, wolf, and grizzly bear occurrence and abundance was indexed using the first-person daily wildlife observations obtained from the journals of European mariners, fur traders, trappers, and government mappers for the period of historical range contraction (Figure 1): CE 1691-1860 in southern Canada and northern United States, and CE 1770-1928 in northern Canada and eastern Alaska (Figure 2). Our study area encompasses the northwestern portion of historic bison distribution in North America (Figure 1). While this excludes a large portion of historic bison range in the southwest and east portions of North America, we chose to focus on restoration initiatives within the Western Cordillera and thus delineated our study area accordingly. We

collected journal observations from the Pacific Ocean and the Alaska-Yukon border in the west to lake Winnipeg and the Hudson Bay in the east, and from the Arctic Ocean in the north to approximately the state boundaries of California, Nevada, Utah and Colorado in the south.

Methods followed Kay's procedures for tallying observations from the Lewis and Clark journals [30]. For bison, wolves and bears, three measures quantify the observations of journalists. First is animals seen where a value of 1 was assigned if journalists reported old sign, 2 if the sign was fresh, and 3 if they actually saw animals. The second index was animals killed, where either the exact number killed is recorded, or where "some" or "a few" is recorded as 3, "several" as 7, and "many" as 10. Third, is herd or group size. Where journalists report sighting large numbers of a species, a value of 10 is assigned, 5 for moderate amounts. Animals seen, killed, and herd/group size are then added together to obtain a measure of abundance. Observations made by journalists at long-term camps or trading posts may be tallied by specified periods of 4 to 30 days with total kill numbers for the period. For quantifying human abundance, if old sign was observed this is assigned a 1, fresh sign a 2, and if the journalists actually saw people a 3, or a 10 if the human group size was greater than ten. Further, the quality of the journal observation was rated as "ND" or no data for day/period, or low, moderate or high depending on the level of detail. The location was plotted as the nightly campsite, and again from low to high quality depending on the journalist's description of the location. For all analyses, we excluded observations where wildlife or location data quality were rated as no data. The complete database for these observations (in spread sheet and Google Earth format) is available in the Supplementary Data, and is available at:

<https://lensoftimenorthwest.com/themes/lens-northwest-files/google-earth-map-journal-wildlife-observations/>

To visualize large-scale trends in bison, human, wolf, and grizzly bear abundance based on historic journal observations, we averaged abundance indices for North American ecoregions [31,32] mapped for our study area. We used an ecoregion scale because they are useful for delineating terrestrial biodiversity patterns for global land-use planning and conservation across taxa [33] and have recently been used in studies of large mammal restoration [34]. Where ecoregion boundaries extend beyond the study area, the mean resource index includes observations from across the ecoregion.

2.2. Journal Data Analysis

2.2.1. Bottom-Up Effects on Bison

We first accounted for the effect of bottom-up factors on bison distribution by constructing an ecological niche model for bison using maximum entropy modelling (MaxEnt) [35]. We predicted that bison distribution would be largely explained by bottom-up modelling, but expected that the addition of top-down factors (see below) would also predict bison occurrence and thus provide evidence of their importance in defining historic bison range. MaxEnt has been widely used to estimate habitat suitability and predict the geographic distribution of a variety of species [36], including both European and North American bison [13,37]. The output of MaxEnt modelling provides an estimated probability of species occurrence based on environmental variables, and we join others in referring to this as synonymous to habitat suitability [36,38,39].

To represent bottom-up factors that could affect habitat suitability, we collected a series of land cover, topographic and climatic predictors in ArcGIS Pro version 2.4.0 [40]. The temporal span during which land cover and climatic predictors were measured does not reach back to time periods when bison were observed and reported by journalists, and land cover and climate have both changed significantly over recent centuries [41]. Our analysis assumes that these variables have remained static over time, and, in part to account for this assumption, we applied current data measured at a large scale (raster pixel size = 200 square kilometers) to describe the general land cover and climatic characteristics of locations where bison were reported. Furthermore, we included a covariate

representing human modification of terrestrial systems to account for changes in land cover since historical times (e.g., the development of cities) [42,43].

For land cover variables, we reclassified the MODIS 2005 land cover map into grass and treed areas at a 250m cell size [44], and calculated their proportional area within each pixel. To capture topographic variation, we calculated terrain ruggedness using the USDS North America Elevation 1-km resolution grid [45]. We calculated a curvature grid in ArcGIS, then used a moving window analysis to determine the standard deviation of curvature within a 3 km radius of each 1-km cell to produce a measure of topographic variability, with high values indicating more rugged terrain [46]. We then averaged ruggedness values within each pixel. Lastly, we collected five climatic variables from a global climate model at a 1km pixel size, which were quantified from climate normal between 1961 and 1990 [47]. We considered annual precipitation as snow and mean temperature of the coldest month to be indicative of snow depth and winter severity, which have been previously shown to affect bison distribution [48,49]. The other three climatic variables were mean annual solar radiation, mean summer precipitation, and mean temperature of the warmest month, which we considered indicative of primary productivity and thus forage availability for bison. Again, we caution that our analysis assumes that modern climates are representative of historic trends, and given the occurrence of the Little Ice Age during the journal observation period this may affect our conclusions as climates have warmed substantially.

We randomly split bison journal observations ($N = 1,654$) into training (75%) and testing (25%), and used the *kuenm* package in R [50] to calibrate 17 candidate models using the *kuenm_cal* function, select the best models with *kuenm_eval*, and then create final models using the full set of occurrences and selected parameters using *kuenm_mod*. We created candidate models including all possible combinations of environmental variables, 14 values of regularization multiplier (0.1–1.0 at intervals of 0.1, 2–5 at intervals of 1, 8 and 10), and linear, quadratic and product feature classes. We excluded hinge or threshold feature classes in an attempt to minimize overfitting by limiting model complexity [51]. We evaluated models based on statistical significance (partial ROC) [52], omission rates ($E = 5\%$) [53], and model complexity ($AICc \leq 2$ from top model) [54]. To evaluate the relative contributions of our environmental predictors, we determined the percent contribution of each variable toward model fit [13], and generated marginal response curves to evaluate the effect of these variables on bison habitat suitability. Finally, we generated 10 replicates of the single top selected model using k-fold cross validation [50], averaged their logistic outputs, binned the values into 10 equal interval categories, and projected this across our study area to quantify bison habitat suitability on a scale from 1 (low suitability) to 10 (high suitability).

2.2.2. Top-Down Effects on Bison

To test the hypothesis that top-down processes acted in addition to bottom-up habitat suitability to limit bison distribution in the Western Cordillera, we modelled the relationship between the presence of bison and the abundance of three species that consume bison (humans, wolves and grizzly bears) using mixed-effect logistic regression [55]. If bottom-up factors were adequate at explaining historic bison distribution, we would expect that top-down factors would fail to significantly explain any variation in bison distribution. We assume that the abundance of humans, wolves and grizzly bears is representative of the harvest or predation intensity that they pose towards bison at each location.

For each journal observation for which data was available, we rated bison presence as 1 if any evidence of bison was observed, or 0 if not. We then used the *glmer* function from package *lme4* [56] to model presence/absence as a function of habitat suitability as determined from the MaxEnt logistic output value at the observation location, and the observed abundance index value for humans, wolves, and grizzly bears. All predictors were mean centered and scaled to a standard deviation of 1, and we verified that no

predictors were highly correlated ($r^2 > 0.6$). We also included the journal observer as a random effect to account for spatial and temporal autocorrelation within journal expeditions. We then used the *dredge* function from package MuMIn to identify the top model(s) based on the most parsimonious combination of predictor variables ($\Delta AIC_c < 2$ from top model [57]). We evaluated model fit of our top selected model(s) based on conditional and marginal R^2 values using the *r2* function from the package performance [58]. We conducted all statistical analyses in *Rstudio* version 2022.2.3.492 [59,60] and considered results significant if p values < 0.05 .

2.3. Cordilleran Bison Restoration Project Demographics and Outcomes

Finally, we test whether the historic edge of current bison range has limited the success of current restoration projects by summarizing the range area and demographics (current population estimates, carrying capacity, growth rates) of bison in populations outside of historic core distribution. We predicted that the current outcomes of these projects may provide further insight to the processes delineating historic bison distribution.

3. Results

3.1. Historical Range and Abundance of Bison

Of 9,384 journal observations where wildlife and locational data were robust, bison were reported 1,841 times. Bison were mostly reported as sightings (81.2 %), but other evidence of their presence was also recorded (18.4 %) based on traditional ecological knowledge acquired from guides or other Indigenous people, feces, tracks, or wallows. Humans were reported as seen or encountered on 3,272 journal observations, and in another 438 records evidence of humans was observed without encountering people. Evidence or visual sightings of wolves or grizzly bears were less frequently reported in 183 and 243 journal observations, respectively.




Figure 2. Study area boundary and location of historic journal observations from CE 1691-1860 in southern Canada and northern United States, and CE 1770-1928 in northern Canada and eastern Alaska (A) and the mean abundance indices per ecoregion of (B) bison, humans (C), grizzly bears (D) and wolves (E). Abundance was indexed following on the methods of Kay (2007), based on sign, harvest and group size.

Across ecoregions (Figure 2), bison abundance largely followed the proposed historic distribution (Figure 1) and centered around the Great Plains ecoregions. Bison were rarely observed in the Rocky Mountain ecoregions and only extended west of the Rocky Mountains in the south end of the study area. In contrast, humans were most abundant west of the Rocky Mountains. The abundance of wolves and grizzly bears was reportedly highest in the same area as bison, the prairie ecoregions, although both were also reported farther north in the region.

3.2. Factors Affecting Bison Distribution

As predicted, bottom-up factors were strong predictors of bison occurrence based on MaxEnt modelling. All 17 candidate models were statistically better than the null model and 16 of these met omission rate criteria, but examination of model complexity revealed a single top model that included all the predictor variables (average area under ROC curve = 0.864, omission rate at 5 % = 0.048, AIC = 28312.04). The predictor with the highest contribution to model fit was the proportion of grass (43.5 %), followed by the mean coldest month temperature (19.8 %), mean annual radiance (16.3 %), human modification (6 %), the proportional tree cover (5.6 %), ruggedness (3.7 %), mean warmest month temperature (2.7 %), precipitation as snow (1.3 %) and mean summer precipitation (1.2%). Marginal response curves (Figure A1) indicated that habitat suitability was positively related to mean annual solar radiance and human modification level, negatively related to mean summer precipitation, precipitation as snow, mean warmest month temperature and the proportional tree cover, and that intermediate values of mean coldest month temperature, ruggedness, and proportion of grass yielded the highest habitat suitability values.

Habitat suitability varied across the study area with the highest values occurring on the Great Plains east of the Rocky Mountains (Figure 3). The Rocky Mountains and higher elevation areas had much lower habitat suitability, as did areas in the far north and near the coasts. However, several areas in the Western Cordillera exhibited some habitat suitability, including the larger valleys of the Cordillera in Montana and Wyoming, the grassland interior plateau of British Columbia, and the plains of Idaho and Washington state.




Figure 3. Bison habitat suitability (1 = low, 10 = high) as determined through MaxEnt modelling of bison occurrences based on historic journal observations. Models were constructed in R using the *kuenm* package and used climatic, land cover, and topographic variables as bottom-up predictors of bison occurrence.

Top-down factors were important predictors of bison occurrence, as mixed-effect logistic regression analysis of bison presence revealed a clear top model (>5 AIC_c from the next model) that included all predictor variables (Table 1). The conditional R² value was 0.520 while the marginal R² was 0.322, suggesting reasonably good model fit. Bottom-up habitat suitability, quantified based on MaxEnt output values, was the strongest predictor of bison presence, as expected. Of the top-down effects on bison distribution, only human abundance was negatively related to bison occurrence, while a positive relationship existed between the abundance of wolves and grizzly bears and bison presence (Table 1).

Table 1. Top mixed-effect logistic regression model predicting bison presence or absence in historic journal records as a function of MaxEnt habitat suitability and the abundance of humans, wolves, and grizzly bears.

<i>Predictors</i>	<i>Estimate</i>	<i>Std. Error</i>	<i>p</i>
(Intercept)	-2.76	0.192	< 0.001
MaxEnt habitat suitability	1.33	0.0575	< 0.001
Human abundance index	-0.224	0.0404	< 0.001
Grizzly bear abundance index	0.0827	0.0291	0.0045
Wolf abundance index	0.201	0.0361	<0.001

3.3. Cordilleran Bison Restoration Project Outcomes

Table 2 summarizes demographics for seven ongoing bison restoration projects in the Cordillera. Although all native non-human predators occur in all areas, annual population rates of increase are generally high (>10%) when populations are <50% of carrying capacity. The exceptions are those projects such as Nordquist and Nahanni that are bisected with highways causing high bison mortality [61]. Generally, where carrying capacity is estimated from habitat quality, the estimated density is higher than for those where it is determined from social criteria. There is no apparent relationship between demographics of projects either on the edge, or those outside historic bison range. For example, although the Aishihik project in the Yukon is several hundred kilometers NW of the historical bison range boundary, both population growth rates and potential population totals comparable to those in or near historic range.

Table 2. Demographics for select current bison restoration projects in the Western Cordillera, North America.

Location and ecotype [references]	Estimated Range (km ²)	Population Estimate (year)	Estimated carrying capacity (K, bison/ km ²)	Observed annual rate of increase (%) at $\leq 50\% K$	Remarks
Yellowstone, WY, MT Plains Bison [22,62,63]	9,400	4,500 (2018)	.405	12	All native predators except humans. K determined by habitat quality and demographics, population c. 2010 >75% of K (counts of ~4000), large impacts on grasslands, woody plants and hydrologic indicators, emigration rate out of park is increasing
Banff, AB Plains Bison [64–66]	1,200	80 (2020)	.480	30	All native predators except humans. K determined by habitat quality, current increase rate very high due to high female-to-male ratio of the founder herd, current population <100 but increasing rapidly
Halfway-Sikkanni, BC Plains Bison [67–69]	3,200	1,013 (2014)	.500	10-38	All native predators present including humans. 48 plains bison escaped in 1971, population ~1300 in 2006, K socially determined to minimize range expansion
Etthithun, BC & AB Wood Bison [61,68,70–72]	5,000	193 (2010)	No information	15-20	All native predators present. First Nations allocated permits to take bison to discourage range expansion to conflicts with industry and agriculture
Nordquist, BC Wood Bison [61,68,71,73–75]	1,400	124-142 (2010)	No information	<5%	All native predators present including humans. A total of 49 bison released in 1995 at Aline Lake, by 2010, range mainly along 170 km Alaska Highway, population limited by traffic accidents
Nahanni, NT & BC Wood Bison [76,77]	11,700	431 (2011)	No information	~8%	All native predators present including humans. Slow growth since establishment, limiting factors appear to be drowning, traffic accidents, hunting, and perhaps tooth wear due to diet high in silica
Aishihik, YT Wood Bison [78–82]	11,000	1,106-1,385 (2011)	.125	10-20	All native predators present. Socially determined carrying capacity to minimize conflicts with Indigenous land uses and competition with other species

4. Discussion

Our analysis of historical journal observations informs bison restoration in the Western Cordillera by supporting the hypothesis that top-down human harvest pressure, in addition to bottom-up habitat suitability, acted to influence bison population dynamics in this region. First, we found that bottom-up factors largely explained bison distribution and highlighted a large band of low-suitability habitat in the Rocky Mountains that isolated otherwise suitable areas in the Cordillera from the core bison range on the Great Plains. Second, we identified a negative relationship between bison occurrence and the abundance of humans even when bottom-up habitat suitability was included in models, suggesting that harvest pressure may have also been an important determinant of the western limit of bison distribution. The rapid expansion of bison populations restored to these areas of historically low habitat suitability provides further support of our hypothesis. Based on our findings, we propose that maximizing the ecological and cultural benefits of bison restoration may require the re-establishment of the role of Indigenous people

as a key socio-ecological process to regulate bison populations in the Western Cordillera [83,84].

4.1. Processes Limiting Historic Bison Distribution

With a MaxEnt ecological niche modelling approach, we identified a large core area of highly suitable bison habitat from central Alberta, Canada, southward along the Rocky Mountains into the Great Plains, falling within the historic bison distribution mapped by Allen [3] and Roe [4] (Fig 1, 3). Of the two recognized ecotypes of North American bison [85,86], the MaxEnt model largely encompasses the range of the migratory “plains” variant that occurred in high numbers on the grasslands of the southern Great Plains. Traditional ecological knowledge and numerous accounts of early journalists provide evidence of plains bison herds numbering in the tens of thousands [30,87,88]. The range of “wood bison”, a more sedentary ecotype that existed at low densities in the northern mixed and boreal forest, was identified as less-suitable habitat by MaxEnt. Both bottom-up factors as identified by the model, and top-down pressure may have acted to regulate bison distribution and abundance.

Within the core range, a combination of factors likely made bison abundant including high forage availability from expansive grasslands [89,90], favorable climatic conditions [13], a lack of geographic barriers to formation, movements, and dispersal of large herds [90,91], and areas that provided refuge from human harvest [89,91]. Indigenous groups in this area were heavily dependent on bison [2,92,93], but over-harvest was likely limited because bison found refuge both spatially, in intertribal buffer zones between conflicting Indigenous groups [30,94], and temporally during winter when large areas were mostly inaccessible to human hunters because of unavailable firewood and shelter on the exposed prairie [91,95,96]. Further north and west in forested areas, bison likely persisted despite lower habitat suitability due to spacing in low densities, low human populations, and high connectivity leading to periodic dispersals from large bison herds to the south and east [30,97]. Source-sink wildlife population dynamics likely operated within this core area, both between and within plains and wood subspecies, to allow for persistent bison populations in many areas [30,88,98,99]. In contrast to the high numbers of bison found throughout the core distribution, few bison were reported west of the Rocky Mountains, and we suggest that this was caused by both more limited connected habitat and greater top-down harvest pressure from humans.

Outside of the core range, bottom-up habitat connectivity was likely in-part responsible for this dynamic in the northern part of bison range, where the model identified a large band of low-suitability habitat in the northern Rocky Mountains that likely limited colonization. The topography of the foothills and mountains, and deep winter snowpack, also provided opportunities for Indigenous people to harvest bison using natural traps and jumps. There is considerable archeological evidence of bison harvest in this region [14–16,100,101], and this likely further limited bison dispersal westward. Connectivity was less of a barrier in the south, as habitat corridors existed in Wyoming and southwest Montana [102]. Indeed, low numbers of bison were historically observed west of the Rockies on the grasslands of southern Idaho, northern Utah, eastern Oregon, and south-eastern Washington (Figure 2). Archeological evidence also shows periodic bison presence in this region throughout the Holocene period [103,104]. Source-sink dynamics may have existed here with low-density herds occasionally being replenished by larger herds dispersing from core bison range on the Great Plains. During this period, bison body mass in these small herds declined in a pattern similar to those on the Great Plains, suggesting their origin was possibly from periodic dispersals through these corridors from the heartland of bison habitat to the east [105]. Even without re-colonization from larger populations, some low-density populations may have been self-sustaining in the Western Cordillera despite low connectivity to the Great Plains populations. Future research should aim to characterize bison movement corridors [102], migration patterns [95], and source-

sink population dynamics to better understand the historic distribution and abundance of bison populations.

Top-down harvest pressure from human populations was likely also higher both in the north and south portion of the Western Cordillera than in core bison range, because they contained large human populations that were supported by substantial plant and salmon resources [106–108]. Our analysis and other research supports the hypothesis that high human abundance is negatively related to bison occurrence due to the socio-political pressure to over-exploit bison populations where possible [30,88,98,99]. For example, the relatively recent extirpation of bison (<3000 BP) in Alaska and Yukon may be associated with increasing human numbers [109,110], and development of hide trade routes from the Liard and Yukon rivers to the Pacific coast [111,112]. The recent population growth of bison restored both at the edge and outside of their core historic range (Table 2) emphasizes that human harvest may have been as important, or potentially more limiting, than bottom-up factors in these areas. We also emphasize that the effect of human harvest on bison interacted with various bottom-up factors that had more prevalent impacts in the Western Cordillera such as deep snow, rugged topography with potential natural traps, and abundant wood resources from forests.

We acknowledge several limitations of our analysis that should be considered when interpreting our findings. First, as mentioned in the methods, covariates used in MaxEnt models assume that climatic and land cover variables measured in the 21st century are representative of historical patterns. While we believe that this assumption is appropriate for modelling broad-scale trends in bison distribution, changes in local climate and land cover [41] make more detailed interpretations of our habitat suitability model potentially perilous. Also, climate warming since the period during which observations were made may cause directional bias in our MaxEnt model to suggest bison habitat occurred more in cooler climates than was the case, shifting our model northward. Second, journal observations were largely located along historic travel routes that were influenced by climate, terrain, and food availability. This introduces another potential source of bias in this dataset towards river valleys and other low-resistant travel routes, which we attempted to account for by using a large (200 square kilometer) raster pixel size. Additionally, with the more intensive sampling in the south part of our study area, our estimates may be biased towards this area. Third, historical journal observations are almost exclusively written from a male, colonial perspective. While the records of some journals describe traditional ecological knowledge shared by Indigenous guides or groups encountered when travelling, they often do not describe plant use, or female perspectives on seasonal resource availability. Future research requires further Indigenous knowledge of historic bison and other resources to better understand the nuance of historic human-bison interactions [6,101,106,113–115].

4.2. Implications for Bison Restoration in the Western Cordillera

Despite these bottom-up and top-down limitations imposed on historical bison distribution, recent restoration projects have achieved rapid bison population growth in areas that were historically on the fringe or outside of what we identified as suitable habitat (Figure 3, Table 2). The success of these projects demonstrates that bison can thrive in a broad range of climates, vegetation conditions, and predators, even if snow depth [48], forage quality and availability [116] and non-human predation [117] influence their localized spatial distribution. Recent studies have also demonstrated remarkable plasticity in bison diets across North America [118,119], and MaxEnt ecological niche modelling based on fossil records of bison and historic climates demonstrated that much of North America was suitable bison habitat over the past 20,000 years [13].

Bison can clearly thrive outside of their historical range with minimal human harvest pressure, but their potential adverse effects on areas that have not recently sustained high numbers of bison may present a challenge for bison restoration. These impacts can occur from intense herbivory on ill-adapted vegetation species [23–25,63,83], direct and

apparent competition with other species [120], disease transmission [1,25], and detrimental cultural effects through human injury and damage to cultural resources [79,121]. However, restoring the top-down process posed by Indigenous people's harvest of bison may benefit bison populations from reduced disease prevalence in low-density herds under top-down pressure [122], higher social tolerance from more involved local communities [121,123], reduced risk of excessive herbivory that degrades their ecosystems [23,24,84], and, as an additional but substantial benefit, Indigenous people may restore deep cultural connections [124]. Our analysis supported the hypothesis that this top-down process may have limited bison populations in the Western Cordillera, adding to a large body of recent evidence demonstrating that Indigenous people were a key ecological force shaping North American landscapes by affecting the abundance of large mammals [30,88], cultivating and gathering plant and animal species [114,125], and manipulating fire regimes across the continent [113,126–128]. Restoring these socio-ecological dynamics may help restore bison across the Western Cordillera.

Of the bison restoration projects discussed in this paper, Indigenous harvest as a socio-ecological process has been partially restored in Aishihik, Nahanni, Nordquist, Etthithun, and Halfway-Sikanni. In the national parks of Banff and Yellowstone, Indigenous harvest has not yet been restored and this top-down process remains de-coupled from these bison populations. Free-roaming bison have only recently been restored to Banff in 2018 [64,65], and implementing Indigenous harvest and other practices early in the project may allow it to quickly gain the cultural benefits of bison harvest while proactively mitigating adverse ecosystem impacts [83,124]. In contrast, Yellowstone has had free-roaming bison for over a century [63], and populations have reached a point where ecosystem recovery is challenged by overabundant bison as Indigenous people remains excluded from harvesting or burning the park [19,23,25]. Without policy adjustments and management changes, both Banff and Yellowstone may risk of the ecological and socio-ecological consequences of overabundant bison populations in historically low abundance areas [23,79,83,121].

Adjusting 19th and 20th century socio-political decisions that excluded Indigenous people [19,20] and restoring their socio-ecological role in managing both bison and their ecosystems may have numerous benefits [6]. In addition, human harvesting of bison is only one process in complex Cordilleran ecosystems and cannot be restored in isolation—it must be done in the context of a holistic ecosystem restoration method that recognizes long-term human roles as not only harvesters, but also as gatherers, cultivators, and prescribed burners [30,88,113,114,125–128]. There are several examples of successful co-management models built on strong relationships between science and traditional ecological knowledge [6,129,130], and we encourage bison restoration projects to follow suite. Co-managing bison with Indigenous peoples and restoring the top-down harvest pressure that they provide may be the key to restoring bison and their full range of ecological and cultural benefits to the Western Cordillera and across North America.


Funding: Parks Canada provided initial funding for historical journal database compilation.

Data Availability Statement: Geospatial data and R code supporting the reported results of this study are openly available on GitHub at https://github.com/jfarr99/Farr_and_White_Historic_Bison_Distribution. The full dataset of historical journal observations is available at <https://lensofthenorthwest.com/themes/lens-northwest-files/google-earth-map-journal-wildlife-observations/> or after 2025 in the Whyte Museum of the Canadian Rockies, Banff, Alberta Clifford A White fonds

Acknowledgments: We appreciate comments from M. Hebblewhite, K. Heuer, and C Lewis on earlier drafts of this manuscript. We respectfully acknowledge that this work was conducted on Treaty 7 territory, a traditional gathering place for diverse Indigenous peoples including the Blackfoot Confederacy (comprising the Siksika, Piikani, and Kainai First Nations), as well as the Tsuut'ina First Nation, and the Stoney Nakoda (including the Chiniki, Bearspaw, and Wesley First Nations). We hope that our future work on bison will be conducted in collaboration with these peoples.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A



Appendix A Figure A1. Marginal response curves revealing the relationship between predictor variables (x axes) and bison habitat suitability (y axes) as determined by the top selected MaxEnt model.

References

- Gates, C.C., Freese, G.H., Gogan, P., Kotzman, M. (eds. and comps.). *American Bison: Status Survey and Conservation Guidelines*; International Union Conservation Nature: Gland, Switzerland 2011.
- Isenberg, A.C. *The destruction of the bison an environmental history, 1750-1920*; Cambridge University Press: Cambridge, United States, 2000.
- Allen, J.A. *The American Bisons, Living and Extinct*; Harvard College Memoirs of the Museum of Comparative Zoology; University Press, Cambridge: Cambridge, MA, 1876; Vol. 10.
- Roe, F.G. *The North American Buffalo*; University of Toronto Press: Toronto, 1972.
- Wildlife Conservation Policy: A Reader*; Geist, V., McTaggart-Cowan, I. (eds.); Detselig Enterprises: Calgary, 1998.
- Hessami, M.A.; Bowles, E.; Popp, J.N.; Ford, A.T. Indigenizing the North American Model of Wildlife Conservation. *FACETS* **2021**, *6*, 1285–1306, doi:10.1139/facets-2020-0088.
- Gates, P.W. *History of Public Land Law Development*; US Government Printing Office: Washington, United States; 1968; Vol. 62.
- Donahue, D.L. *The Western Range Revisited: Removing Livestock from Public Lands to Conserve Native Biodiversity*; University of Oklahoma Press, 1999; Vol. 5.
- Bryner, G. Balancing preservation and logging: Public lands policy in British Columbia and the Western United States. *Policy Studies Journal* **1999**, *27*, 307–327.
- Hebblewhite, M.; Hilty, J.A.; Williams, S.; Locke, H.; Chester, C.; Johns, D.; Kehm, G.; Francis, W.L. Can a large-landscape conservation vision contribute to achieving biodiversity targets? *Conservat Sci and Prac* **2022**, *4*, 1-11, doi:10.1111/csp2.588.
- Daubenmire, R. The western limits of the range of the American Bison. *Ecology* **1985**, *66*, 622–624.
- Van Vuren, D. Bison west of the Rocky Mountains: An alternative explanation. *Northwest science (USA)* **1987**, *61(2)*, 65-69.
- Wendt, J.A.F.; McWethy, D.B.; Widga, C.; Shuman, B.N. Large-scale climatic drivers of Bison distribution and abundance in North America since the Last Glacial Maximum. *Quaternary Science Reviews* **2022**, *284*, 107472, doi:10.1016/j.quasci-rev.2022.107472.
- Bailey, J.A. Historic distribution and abundance of Bison in the Rocky Mountains of the United States. *Intermountain Journal of Sciences* **2016**, *22*, 36–53.
- Reeves, B.O.K. Bison killing in the Southwestern Alberta Rockies. *Plains Anthropologist* **1978**, *23*, 63–78, doi:10.1080/2052546.1978.11908902.
- Zedeño, M.N. Bison hunters and the Rocky Mountains: An evolving partnership. *Quaternary International* **2017**, *461*, 80–101, doi:10.1016/j.quaint.2017.06.043.
- Zedeño, M.N.; Ballenger, J.A.M.; Murray, J.R. Landscape engineering and organizational complexity among late prehistoric Bison hunters of the Northwestern Plains. *Current Anthropology* **2014**, *55*, 23–58, doi:10.1086/674535.

18. Mack, R.N.; Thompson, J.N. Evolution in Steppe with few large, hooved mammals. *The American Naturalist* **1982**, *119*, 757–773.
19. Nabokov, P.; Loendorf, L. *Restoring a Presence: American Indians and Yellowstone National Park*; University of Oklahoma Press: Norman, United States, 2004.
20. Binnema, T.; Niemi, M. 'Let the line be drawn now:' Wilderness, conservation, and the exclusion of Aboriginal people from Banff National Park in Canada. *Environmental History* **2006**, *11*, 724–750.
21. Boyce, M.S. Natural regulation or control of nature? In *The Greater Yellowstone Ecosystem: Redefining America's Wilderness Heritage*; Keiter, B., Boyce, M.S., Eds.; Yale University Press: New Haven, United States, 1991; pp. 183–207.
22. Wagner, F.H. *Yellowstone's Destabilized Ecosystem: Elk Effects, Science, and Policy Conflict*; Oxford University Press: Oxford, United States, 2006.
23. Beschta, R.L.; Ripple, W.J.; Kauffman, J.B.; Painter, L.E. Bison limit ecosystem recovery in Northern Yellowstone. *Food Webs* **2020**, *23*, e00142, doi:10.1016/j.fooweb.2020.e00142.
24. Kay, C.E. The condition and trend of Aspen, Willows, and associated species on the Northern Yellowstone Range. *Rangelands* **2018**, *40*, 202–211, doi:10.1016/j.rala.2018.10.005..
25. Yonk, R.M.; Mosley, J.C.; Husby, P.O. Human influences on the Northern Yellowstone Range. *Rangelands* **2018**, *40*, 177–188, doi:10.1016/j.rala.2018.10.004.
26. White, C.A.; Olmsted, C.E.; Kay, C.E. Aspen, Elk, and Fire in the Rocky Mountain National Parks of North America. *Wildlife Society Bulletin* **1998**, *26*, 449–462.
27. Wyman, T. Grizzly Bear Predation on a Bull Bison in Yellowstone National Park. *Ursus* **2002**, *13*, 375–377.
28. Carbyn, L.N.; Trotter, T. Descriptions of Wolf Attacks on Bison Calves in Wood Buffalo National Park. *ARCTIC* **1988**, *41*, 297–302, doi:10.14430/arctic1736.
29. Smith, D.W.; Mech, L.D.; Meagher, M.; Clark, W.E.; Jaffe, R.; Phillips, M.K.; Mack, J.A. Wolf-Bison Interactions in Yellowstone National Park. *Journal of Mammalogy* **2000**, *81*(4), 1128–1135, doi:10.1644/1545-1542(2000)081<1128:WBIIYN>2.0.CO;2.
30. Kay, C.E. Were Native people keystone predators? A continuous-time analysis of wildlife observations made by Lewis and Clark. *Canadian Field Naturalist* **2007**, *121*, 1–16.
31. Wiken, E.; Nava, F.J.; Griffith, G. *North American Terrestrial Ecoregions—Level III*; Commission for Environmental Cooperation: Montreal, Canada, 2011.
32. Dinerstein, E.; Olson, D.; Joshi, A.; Vynne, C.; Burgess, N.D.; Wikramanayake, E.; Hahn, N.; Palminteri, S.; Hedao, P.; Noss, R.; et al. An ecoregion-based approach to protecting half the terrestrial realm. *BioScience* **2017**, *67*, 534–545, doi:10.1093/bio-sc/bix014.
33. Smith, J.R.; Hendershot, J.N.; Nova, N.; Daily, G.C. The biogeography of ecoregions: Descriptive power across regions and taxa. *J Biogeogr* **2020**, *47*, 1413–1426, doi:10.1111/jbi.13871.
34. Vynne, C.; Gosling, J.; Maney, C.; Dinerstein, E.; Lee, A.T.L.; Burgess, N.D.; Fernández, N.; Fernando, S.; Jhala, H.; Jhala, Y.; et al. An ecoregion-based approach to restoring the world's intact large mammal assemblages. *Ecography* **2022**, *2022*, doi:10.1111/ecog.06098.
35. Phillips, S.J.; Dudík, M.; Schapire, R.E. A maximum entropy approach to species distribution modeling. 665–662.
36. Venne, S.; Currie, D.J. Can habitat suitability estimated from MaxEnt predict colonizations and extinctions? *Divers Distrib* **2021**, *27*, 873–886, doi:10.1111/ddi.13238.
37. Bleyhl, B.; Sipko, T.; Trepel, S.; Bragina, E.; Leitão, P.J.; Radeloff, V.C.; Kuemmerle, T. Mapping seasonal European Bison habitat in the Caucasus Mountains to identify potential reintroduction sites. *Biological Conservation* **2015**, *191*, 83–92, doi:10.1016/j.bio-con.2015.06.011.
38. Kuemmerle, T.; Perzanowski, K.; Chaskovskyy, O.; Ostapowicz, K.; Halada, L.; Bashta, A.-T.; Kruhlov, I.; Hostert, P.; Waller, D.M.; Radeloff, V.C. European Bison habitat in the Carpathian Mountains. *Biological Conservation* **2010**, *143*, 908–916, doi:10.1016/j.biocon.2009.12.038.
39. Su, H.; Bista, M.; Li, M. Mapping habitat suitability for Asiatic Black Bear and Red Panda in Makalu Barun National Park of Nepal from Maxent and GARP models. *Sci Rep* **2021**, *11*, 14135, doi:10.1038/s41598-021-93540-x.
40. Esri Inc. ArcGIS Pro (Version 2.4.0) 2019.
41. Rockström, J.; Steffen, W.; Noone, K.; Persson, Å.; Chapin, F.S.; Lambin, E.F.; Lenton, T.M.; Scheffer, M.; Folke, C.; Schellnhuber, H.J.; et al. A safe operating space for humanity. *Nature* **2009**, *461*, 472–475, doi:10.1038/461472a.
42. Kennedy, C.M.; Oakleaf, J.R.; Theobald, D.M.; Baruch-Mordo, S.; Kiesecker, J. Global Human Modification of Terrestrial Systems. *NASA Socioeconomic Data and Applications Center (SEDAC)* **2020**, doi:https://doi.org/10.7927/edbc-3z60.
43. Kennedy, C.M.; Oakleaf, J.R.; Theobald, D.M.; Baruch-Mordo, S.; Kiesecker, J. Managing the Middle: A Shift in Conservation Priorities Based on the Global Human Modification Gradient. *Glob Change Biol* **2019**, *25*, 811–826, doi:10.1111/gcb.14549.
44. Canada Centre for Remote Sensing (CCRS), Earth Sciences Sector, Natural Resources Canada. 2005 Land Cover of North America at 250 Meters. *Canada Centre for Remote Sensing (CCRS)* **2013**, *3*.
45. U.S. Geological Survey North America Elevation 1-Kilometer Resolution GRID 2007.
46. Poole, K.G.; Stuart-Smith, K.; Teske, I.E. Wintering strategies by Mountain Goats in interior mountains. *Can. J. Zool.* **2009**, *87*, 273–283, doi:10.1139/Z09-009.

47. Mahony, C.R.; Wang, T.; Hamann, A.; Cannon, A.J. A global climate model ensemble for downscaled monthly climate normals over North America. *Intl Journal of Climatology* **2022**, joc.7566, doi:10.1002/joc.7566.
48. Sheppard, A.H.C.; Hecker, L.J.; Edwards, M.A.; Nielsen, S.E. Determining the influence of snow and temperature on the movement rates of Wood Bison (*Bison Bison Athabascae*). *Can. J. Zool.* **2021**, *99*, 489–496, doi:10.1139/cjz-2020-0280.
49. Meagher, M. Range expansion by Bison of Yellowstone National Park. *Journal of Mammalogy* **1989**, *70*, 670–675, doi:10.2307/1381449.
50. Cobos, M.E.; Peterson, A.T.; Barve, N.; Osorio-Olvera, L. Kuenm: An R package for detailed development of Ecological Niche Models using Maxent. *PeerJ* **2019**, *7*, 2–15, doi:10.7717/peerj.6281.
51. Merow, C.; Smith, M.J.; Edwards, T.C.; Guisan, A.; McMahon, S.M.; Normand, S.; Thuiller, W.; Wüest, R.O.; Zimmermann, N.E.; Elith, J. What do we gain from simplicity versus complexity in Species Distribution Models? *Ecography* **2014**, *37*, 1267–1281, doi:10.1111/ecog.00845.
52. Peterson, A.T.; Papeş, M.; Soberón, J. Rethinking Receiver Operating Characteristic Analysis Applications in Ecological Niche Modeling. *Ecological Modelling* **2008**, *213*, 63–72, doi:10.1016/j.ecolmodel.2007.11.008.
53. Anderson, R.P.; Lew, D.; Peterson, A.T. Evaluating Predictive Models of Species' Distributions: Criteria for Selecting Optimal Models. *Ecological Modelling* **2003**, *162*, 211–232, doi:10.1016/S0304-3800(02)00349-6.
54. Warren, D.L.; Seifert, S.N. Ecological Niche Modeling in Maxent: The Importance of model complexity and the performance of model selection criteria. *Ecological Applications* **2011**, *21*, 335–342, doi:10.1890/10-1171.1.
55. Hosmer, D.W.; Lemeshow, S. *Applied Logistic Regression: Hosmer/Applied Logistic Regression*; John Wiley & Sons, Inc.: Hoboken, United States, 2000; ISBN 978-0-471-72214-4.
56. Bates, D.; Machler, M.; Bolker, B.; Walker, S. Fitting Linear Mixed-Effects Models using Lme4. *Journal of Statistical Software* **2015**, *67*, 1–48, doi:10.18637/jss.v067.i01.
57. Barton, K. MuMIn: Multi-Model Inference 2022.
58. Lüdtke, D.; Ben-Shachar, M.; Patil, I.; Waggoner, P.; Makowski, D. Performance: An R package for assessment, comparison and testing of statistical Mmodels. *JOSS* **2021**, *6*, 3139, doi:10.21105/joss.03139.
59. RStudio Team RStudio: Integrated Development Environment for R 2022.
60. R Core Team R: A Language and Environment for Statistical Computing 2022.
61. Harper, W.L.; Elliot, J.P.; Hatter, I.; Schwantje, H. *Management Plan for Wood Bison in British Columbia*; Wildlife Bulletin; British Columbia, Ministry of Environment, Lands, and Parks, Wildlife Branch: Victoria, Canada, 2000; p. 43; ISBN 0-7726-4200-1.
62. Plumb, G.E.; White, P.J.; Coughenour, M.B.; Wallen, R.L. Carrying capacity, migration, and dispersal in Yellowstone Bison. *Biological Conservation* **2009**, *142*, 2377–2387, doi:10.1016/j.biocon.2009.05.019.
63. Mosley, J.C.; Munding, J.G. History and Status of Wild Ungulate Populations on the Northern Yellowstone Range. *Rangelands* **2018**, *40*, 189–201.
64. Laskin, D.N.; Watt, D.; Whittington, J.; Heuer, K. Designing a fence that enables free passage of wildlife while containing reintroduced Bison: A multispecies evaluation. *Wildlife Biology* **2020**, *2020*, 1–14, doi:10.2981/wlb.00751.
65. Heuer, K. Update to Stakeholders on Banff Bison Restoration Project 2021.
66. Steenweg, R.; Hebblewhite, M.; Gummer, D.; Low, B.; Hunt, B. Assessing potential habitat and carrying capacity for reintroduction of Plains Bison (*Bison Bison Bison*) in Banff National Park. *PLoS ONE* **2016**, *11*, doi:10.1371/journal.pone.0150065.
67. Rowe, M. *Halfway-Sikanni Plains Bison Inventory*; BC Ministry of Environment: Fort St. John, Canada, 2006.
68. COSEWIC COSEWIC Assessment and Status Report on the Plains Bison (*Bison Bison Bison*) and the Wood Bison (*Bison Bison Athabascae*) in Canada.; Committee on the Status of Endangered Wildlife in Canada: Ottawa, Canada, 2013.
69. Hoar, B.R. Does the Pink Mountain Region Contain Sufficient Habitat Suitable for Plains Bison in Summer and Winter?; BC Ministry of Forests, Lands, Natural Resource Operations and Rural Development, 2021.
70. Rowe, M.; Backmeyer, R. *Ethithun Wood Bison Inventory*; BC Ministry of Environment: Fort St. John, BC, 2006.
71. Leverkus, S.E.R. Conservation of Biodiversity in Northern Canada through Ecological Processes and Cultural Landscapes; Dissertation; Oklahoma State University: Stillwater, United States, 2015.
72. Lewis, C.J.; Das Gupta, S. *Wood Bison - Ethithun Herd Minimum Population Survey Results: February 22-25th, 2021*; British Columbia Ministry of Forests, Lands, Natural Resource Operations and Rural Development: Peace Region, Canada, 2021; pp. 1–30.
73. Rowe, M. *Nordquist Wood Bison Inventory*; Peace Region Technical Report; BC Ministry of Environment: Fort St. John, Canada, 2007.
74. Jung, T.S. Extralimital movements of reintroduced Bison (*Bison Bison*): Implications for potential range expansion and human-wildlife conflict. *Eur J Wildl Res* **2017**, *63*, 35, doi:10.1007/s10344-017-1094-5.
75. Theissen, C. *Peace Wood Bison Project: Annual Report 2009/2010*; BC Ministry of Environment: Fort St. John, Canada, 2010.
76. Larter, N.C.; Allaire, D.G. *History and Current Status of the Nahanni Wood Bison Population*; File Report; Government of the Northwest Territories: Yellowknife, Canada, 2007.
77. Larter, N.C.; Allaire, D.G. *Population Survey of the Nahanni Wood Bison Population*; Manuscript Report; Environment and Natural Resources, Government of the Northwest Territories: Yellowknife, Canada, 2013.
78. Markel, C.; Clark, D. Developing policy alternatives for the management of Wood Bison (*Bison Bison Athabascae*) in Kluane National Park. *The Northern Review* **2012**, *36*, 53–75.

79. Clark, D.A.; Workman, L.; Jung, T.S. Impacts of reintroduced bison on First Nations People in Yukon, Canada: Finding common ground through participatory research and social learning. *Conservation and Society* **2016**, *14*, 1–12, doi:10.4103/0972-4923.182798.
80. Jung, T.S.; Czetwertynski, S.M.; Schmiegelow, F.K.A. Boreal Forest titans do not clash: Low overlap in winter habitat selection by Moose (*Alces Americanus*) and reintroduced Bison (*Bison Bison*). *Eur J Wildl Res* **2018**, *64*, 25, doi:10.1007/s10344-018-1184-z.
81. Jung, T.S.; Hegel, T.M.; Stotyn, S.A.; Czetwertynski, S.M. Co-occurrence of reintroduced and resident ungulates on a shared winter range in Northwestern Canada. *Écoscience* **2015**, *22*, 7–16, doi:10.1080/11956860.2015.1047133.
82. Fischer, L.A.; Gates, C.C. Competition potential between sympatric Woodland Caribou and Wood Bison in Southwestern Yukon, Canada. *Can. J. Zool.* **2005**, *83*, 1162–1173, doi:10.1139/z05-117.
83. White, C.A.; Langemann, E.G.; Gates, C.C.; Kay, C.E.; Shury, T.; Hurd, T.E. Plains Bison restoration in the Canadian Rocky Mountains: Ecological and management considerations. In Proceedings of the George Wright Biannual Conference on Research and Resource Management in National Parks and on Public Lands 11; George Wright Society: Washington United States, 2001; Vol. 11, pp. 152–160.
84. Mosley, J.C.; Fidel, J.; Hunter, H.E.; Husby, P.O.; Kay, C.E.; Mundinger, J.G.; Yonk, R.M. An ecological assessment of the Northern Yellowstone Range: Synthesis and call to action. *Rangelands* **2018**, *40*, 224–227, doi:10.1016/j.rala.2018.10.007.
85. Cronin, M.A.; MacNeil, M.D.; Vu, N.; Leesburg, V.; Blackburn, H.D.; Derr, J.N. Genetic Variation and Differentiation of Bison (*Bison Bison*) Subspecies and Cattle (*Bos Taurus*) Breeds and Subspecies. *Journal of Heredity* **2013**, *104*, 500–509, doi:10.1093/jhered/est030.
86. Wilson, G.A.; Strobeck, C. Genetic Variation within and Relatedness among Wood and Plains Bison Populations. **1999**, *42*, 14.
87. Geist, V. Buffalo Nation: History and Legend of the North American Bison; Fifth House: Saskatoon, Canada, 1996.
88. Laliberte, A.S.; Ripple, W.J. Wildlife encounters by Lewis and Clark: A spatial analysis of interactions between Native Americans and wildlife. *BioScience* **2003**, *53*, 994–1003.
89. Binnema, T. Common and Contested Ground: A Human and Environmental History of the Northwestern Plains; University of Oklahoma Press: Norman, United States, 2001.
90. Beaudoin, A.B. A Bison's view of landscape and the paleoenvironment. In *Bison and people on the North American Great Plains*; Cunfer, G., Waiser, W.A., Eds.; Texas A&M University Press: College Station, United Statesm 2016; pp. 51–89.
91. Binnema, T. A Fur trade historian's view of seasonal Bison movements on the Northern Great Plains. In *Bison and people on the American Great Plains: a deep environmental history*; Cunfer, G., Waiser, W.A., Eds.; Texas A&M University Press: College Station, United States, 2016; pp. 159–177.
92. *Bison and People on the North American Great Plains: A Deep Environmental History*; Cunfer, G., Waiser, W.A., Eds.; Connecting the greater west; First edition.; Texas A&M University Press: College Station, 2016.
93. Carlson, K.; Bement, L. Organization of bison hunting at the Pleistocene/Holocene transition on the Plains of North America. *Quaternary International* **2013**, *297*, 93–99, doi:10.1016/j.quaint.2012.12.026.
94. Bayham, J.; Cole, K.E.; Bayham, F.E. Social boundaries, resource depression, and conflict: A bioeconomic model of the Intertribal Buffer Zone. *Quaternary International* **2017**, *518*, 69–82, doi:10.1016/j.quaint.2017.11.007.
95. Moodie, D.W.; Ray, A.J. Buffalo migrations on the Canadian Plains. *Plains Anthropologist* **1976**, *21*, 45–46.
96. Morgan, R.G. Bison movement patterns on the Canadian Plains: An ecological analysis. *Plains Anthropologist* **1980**, *25*, 143–160.
97. Geist, V. Phantom Subspecies: The Wood Bison (*Bison Bison* "Athabasca" Rhoads 1897) Is Not a Valid Taxon, but an Ecotype. *Arctic* **1991**, *44*, 283–300.
98. Kay, C.E. Aboriginal overkill: The role of Native Americans in structuring Western Ecosystems. *Human Nature* **1994**, *5*, 359–398.
99. Martin, P.S.; Szuter, C.R. Megafauna of the Columbia Basin, 1800-1840, Lewis and Clark in a game Sink. In *Northwest Lands, Northwest Peoples: Readings in Environmental History*; Goble, D.D., Hirt, P.W., Eds.; University of Washington Press: Seattle, United States, 1999.
100. Arthur, G. An introduction to the ecology of early historic communal Bison hunting among the Northern Plains Indians. *Canada, National Museums of Canada, National Museum of Man, Mercury Series, Archaeological Survey of Canada Paper* **1975**, *37*, 144.
101. Brink, Jack. Imagining Head-Smashed-In: Aboriginal Buffalo Hunting on the Northern Plains; Athabasca University Press: Edmonton, Canada, 2008.
102. White, C.A. Historical Ecology of Bison Movement Corridors, Western Cordillera, North America: A Cross Watershed Comparison; CW and Associates: Canmore, AB, 2018.
103. Grayson, D.K. Holocene Bison in the Great Basin, Western USA. *The Holocene* **2006**, *16*, 913–925, doi:10.1191/0959683606hol982fa.
104. Lupo, K.; Schmitt, D.N. On Late Holocene Variability in Bison Populations in the Northeastern Great Basin. *Journal of California and Great Basin anthropology* **1997**, *19*, 55–69.
105. Lyman, R.L. Late-Quaternary Diminution and Abundance of Prehistoric Bison (*Bison* Sp) in Eastern Washington State, USA. *Quat. res.* **2004**, *62*, 76–85, doi:10.1016/j.yqres.2004.04.001.
106. Hunn, E.S.; Selam, J. *Nch'i-Wana, "The Big River": Mid-Columbia Indians and Their Land*; University of Washington Press: Seattle, United States, 1990.
107. Prentiss, W.C.; Kuijt, I. Complex Hunter Gatherers Evolution Organization of Prehistoric Communities Plateau of Northwestern North America; Utah State University Press: Salt Lake City, United States, 2004.

108. Prentiss, A.M.; Kuijt, I. *People of the Middle Fraser Canyon an Archaeological History*; University of British Columbia Press, Vancouver, Canada, 2014.
109. Tremayne, A.H.; Winterhalder, B. Large mammal biomass predicts the changing distribution of hunter-gatherer settlements in Mid-Late Holocene Alaska. *Journal of Anthropological Archaeology* **2017**, *45*, 81–97, doi:10.1016/j.jaa.2016.11.006.
110. Doering, B.N. Subarctic landscape adaptations and paleodemography: A 14,000-year history of climate change and human settlement in Central Alaska and Yukon. *Quaternary Science Reviews* **2021**, *268*, 107139, doi:10.1016/j.quascirev.2021.107139.
111. Klein, L. Demystifying the opposition: The Hudson's Bay Company and the Tlingit. *Arctic Anthropology* **1987**, *24*, 101–114.
112. Cloud, J.; Williams, L. The Tlingit Map of 1869. *Expedition Magazine* **2012**, *54*, 10–18.
113. Christianson, A.C.; Sutherland, C.R.; Moola, F.; Gonzalez Bautista, N.; Young, D.; MacDonald, H. Centering Indigenous voices: The role of fire in the Boreal Forest of North America. *Curr Forestry Rep* **2022**, doi:10.1007/s40725-022-00168-9.
114. Joseph, L.; Turner, N.J. "The old foods are the new foods!": Erosion and revitalization of Indigenous Food Systems in Northwestern North America. *Front. Sustain. Food Syst.* **2020**, *4*, 596237, doi:10.3389/fsufs.2020.596237.
115. Peacock, S.M.; Turner, N.J. 'Just like a garden,' traditional resource management and biodiversity conservation on the Interior Plateau of British Columbia. In *Biodiversity and Native America*; Minnis, P.E., Elisens, W.J., Eds.; University of Oklahoma Press: Norman, United States, 2000; pp. 133–179.
116. Fortin, D.; Fryxell, J.M.; O'Brodovich, L.; Frandsen, D. Foraging ecology of Bison at the landscape and plant community levels: The applicability of energy maximization principles. *Oecologia* **2003**, *134*, 219–227, doi:10.1007/s00442-002-1112-4.
117. Harvey, L.; Fortin, D. Spatial heterogeneity in the strength of plant-herbivore interactions under predation risk: The tale of Bison foraging in Wolf country. *PLoS ONE* **2013**, *8*, 73324, doi:10.1371/journal.pone.0073324.
118. Craine, J.M. Seasonal patterns of Bison diet across climate gradients in North America. *Sci Rep* **2021**, *11*, 6829, doi:10.1038/s41598-021-86260-9.
119. Hecker, L.J.; Coogan, S.C.P.; Nielsen, S.E.; Edwards, M.A. Latitudinal and seasonal plasticity in American Bison *Bison Bison* diets. *Mam Rev* **2021**, *51*, 193–206, doi:10.1111/mam.12229.
120. Holt, R.D.; Bonsall, M.B. Apparent competition. *Annu. Rev. Ecol. Evol. Syst.* **2017**, *48*, 447–71.
121. Jung, T.S. Investigating local concerns regarding large mammal restoration: Group size in a growing population of reintroduced Bison (*Bison Bison*). *Global Ecology and Conservation* **2020**, *24*, 01303, doi:10.1016/j.gecco.2020.e01303.
122. Packer, C.; Holt, R.D.; Hudson, P.J.; Lafferty, K.D.; Dobson, A.P. Keeping the herds healthy and alert: Implications of predator control for infectious disease. *Ecol Letters* **2003**, *6*, 797–802, doi:10.1046/j.1461-0248.2003.00500.x.
123. Schusler, T.M.; Chase, L.C.; Decker, D.J. Community-based comanagement: Sharing responsibility when tolerance for wildlife is exceeded. *Human Dimensions of Wildlife* **2000**, *5*, 34–49, doi:10.1080/10871200009359186.
124. Redford, K.H.; Aune, K.; Plumb, G. Hope is a Bison: Editorial. *Conservation Biology* **2016**, *30*, 689–691, doi:10.1111/cobi.12717.
125. Thornton, T.; Deur, D.; Kitka, H. Cultivation of Salmon and other marine resources on the Northwest Coast of North America. *Hum Ecol* **2015**, *43*, 189–199, doi:10.1007/s10745-015-9747-z.
126. Kimmerer, R.W.; Lake, F.K. Maintaining the mosaic - The role of Indigenous burning in land management. *Journal of Forestry* **2001**, *99*, 36–41, doi:https://doi.org/10.1093/jof/99.11.36.
127. Hoffman, K.M.; Davis, E.L.; Wickham, S.B.; Schang, K.; Johnson, A.; Larking, T.; Lauriault, P.N.; Quynh Le, N.; Swerdfager, E.; Trant, A.J. Conservation of Earth's biodiversity is embedded in Indigenous fire stewardship. *Proc. Natl. Acad. Sci. U.S.A.* **2021**, *118*, doi:10.1073/pnas.2105073118.
128. White, C.A.; Perrakis, D.D.B.; Kafka, V.G.; Ennis, T. Burning at the edge: Integrating biophysical and eco-cultural fire processes in Canada's Parks and Protected Areas. *Fire Ecology* **2011**, *7*, 74–106, doi:10.4996/fireecology.0701074.
129. Reid, A.J.; Eckert, L.E.; Lane, J.; Young, N.; Hinch, S.G.; Darimont, C.T.; Cooke, S.J.; Ban, N.C.; Marshall, A. "Two-Eyed Seeing": An Indigenous framework to transform fisheries research and management. *Fish Fish* **2021**, *22*, 243–261, doi:10.1111/faf.12516.
130. Artelle, K.A.; Zurba, M.; Bhattacharyya, J.; Chan, D.E.; Brown, K.; Housty, J.; Moola, F. Supporting resurgent Indigenous-led governance: A nascent mechanism for just and effective conservation. *Biological Conservation* **2019**, *240*, 108284, doi:10.1016/j.bi-con.2019.108284.