


Article

Metamorphic Ages of the Jurassic Accretionary Complexes in the Kanto Mountains, Central Japan, Determined by K–Ar Dating of Illite: Implications for the Tectonic Relationship Between the Chichibu and Sanbagawa Belts

Zhiqiang Lu¹, Ichiko Shimizu^{1,2*} , and Tetsumaru Itaya^{3,4,5}

¹ Geological Institute, Faculty of Science, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

² Division of Earth and Planetary Sciences, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan

³ Hiruzen Institute for Geology & Chronology, 2-5 Nakashima, Naka-ku, Okayama 703-8252, Japan

⁴ Japan Geochronology Network, 2-12 Nakashima, Naka-ku, Okayama 703-8252, Japan

⁵ Institute of GeoHistory, Japan Geochronology Network, 1599 Susai, Akaiwa, 701-2503, Japan

* Correspondence: shimizu.ichiko.8c@kyoto-u.ac.jp

Abstract: To determine the metamorphic ages of the accretionary complexes in the Northern Chichibu Belt in SW Japan, K–Ar dating was conducted using weakly metamorphosed sedimentary rocks collected from the Kanto Mountains, Central Japan. Whole-rock ages were obtained for chert and red shale samples, and the mineral ages of fine-grained illite with the grain size less than 4 μm were obtained for chert, red shale, mudstone, acidic tuff, and basic tuff. The K–Ar ages of chert and red shale presented large variations, with systematically older ages compared to those of mudstone and tuff in the same strata. The influence of submarine hydrothermal activities on chert and red shale before subduction is a possible cause of this deviation. The illite samples, which were fractionated into four grain-size classes using a suspension method, yielded older ages and higher illite crystallinity (i.e., smaller values of Kübler's crystallinity index) for larger grain-size classes. The peak metamorphic ages were determined from the K–Ar ages of the 3–4 μm class illite in mudstone and tuff. The Late Jurassic to the Earliest Cretaceous accretionary complex of the lowest structural unit (Kashiwagi Unit) was dated within a small range between 117–110 Ma, which is distinctly older than the K–Ar ages of white mica reported from the Sanbagawa Belt. The peak metamorphic age of acidic tuff (113 Ma) at the type locality of the Mikabu Greenstones indicates that the subducted Mikabu seamount is a constituent of the Kashiwagi Unit. The peak metamorphic ages of the Manba and Kamiyoshida Units were obtained as 132–107 Ma and 163–144 Ma, respectively. Major structural discontinuity is suggested within the Middle Jurassic accretionary complexes.

Keywords: K–Ar dating; illite crystallinity; Northern Chichibu Belt; Mikabu Greenstones; Sanbagawa (Sambagawa) metamorphism; accretionary complex

1. Introduction

The Chichibu Belt, which is located to the south of the Sanbagawa (or Sambagawa) Belt in the Outer Zone of SW Japan (Figure 1), is constituted of coherent or chaotic units of the Jurassic to Early Cretaceous accretionary complex. The rocks in the Northern Chichibu Belt are weakly metamorphosed up to pumpellyite–actinolite facies in the northern areas, and it was traditionally treated as a southern extension of the 'Sanbagawa metamorphic belt' [6], whereas the 'Sanbagawa Belt' has been used for tectonic division in a narrower sense, wherein crystalline schists are exposed. In the 1980s, a different view on the tectonic relationships between the Sanbagawa and Chichibu Belts had been presented by Faure and Charvet [10,11] in Shikoku, SW Japan. They considered that the

Northern Chichibu complex is a superficial nappe emplaced on the crystalline schists of the Sanbagawa Belt after syn-metamorphic deformation. Guidi et al. [14–16] have applied a similar nappe model to the type localities of the Sanbagawa and Chichibu Belts in the Kanto Mountains, Central Japan.

In central Shikoku, where high-grade metamorphic rocks of the Sanbagawa Belt are largely exposed, K–Ar and Ar–Ar dating methods have been extensively used to determine the muscovite and subduction–uplifting histories of the Sanbagawa schists, as has been previously discussed [7,31,61]. These dating methods have also been applied to fine-grained white mica in weakly metamorphosed sedimentary rocks in the Northern Chichibu Belt, and a systematic change in the metamorphic ages from the northern to the southern parts of the belt was recognized [7,35]. Isozaki and Itaya [25] have considered that the northern unit (ca. 115 Ma, according to K–Ar dating of white mica) was metamorphosed contemporaneously with the main suite of the Sanbagawa Belt, whereas the southern unit (ca. 140 Ma, *ibid*) is a nappe rooted in the Mino–Tanba Belt in the Inner Zone of SW Japan. According to this model, the actual boundary between the Sanbagawa metamorphic belt and the ‘Chichibu belt’, which is free from the Cretaceous Sanbagawa metamorphism, can be drawn within the ‘traditional’ Chichibu Belt [24,27]. The basic assumption of this tectonic model is that the deposition and accretion ages of the host rocks of the Sanbagawa schists and those of the northern unit of the Chichibu Belt were contemporaneous (from the Late Jurassic to Early Cretaceous). However, U–Pb dating of detrital zircons in the past decade has revealed that the protoliths of the Sanbagawa Belt were deposited in the Late Cretaceous, which is comparable with the sedimentation ages in the Northern Shimanto Belt [3,37,38]. Accordingly, the tectonic model of the Sanbagawa and Chichibu Belts must be revised.

The Mikabu Greenstones (or Mikabu greenrocks) [39] is a volcanic complex with distinct lithology, including pillow lava and breccia, hyaroclastite, dolerite, gabbro, and ultramafic cumulates, which are exposed at the boundary region between the Sanbagawa and Chichibu Belts. The Mikabu Greenstone and surrounding meta-sedimentary rocks have been treated as an independent tectonic division, with names such as the ‘Mikabu belt’, ‘Mikabu greenstone belt’ [7], and ‘Mikabu unit’ [9]; a sub-unit of the Sanbagawa Belt [47]; or a member or sub-unit of the Northern Chichibu Belt [15,57]. The origin of the Mikabu Greenstones is considered as a seamount chain [48] or an oceanic plateau [62], which records volcanic activity from Early Jurassic (ca. 200 Ma, based on K–Ar dating of hornblende) [50] to Late Jurassic (155 Ma for basalt [53] and 157 Ma for anorthosite, based on U–Pb dating of igneous zircon [62]). Here, the term ‘Mikabu seamount’ is conventionally used for the original volcanic bodies of the Mikabu Greenstones. Determining the subduction history of the Mikabu seamount is the key to understanding the tectonic evolution of the Mesozoic accretionary complex in the Chichibu–Sanbagawa Belts. Previously, white mica K–Ar ages of 124 Ma [68], 110 Ma [62], and 82 Ma [47], as well as whole-rock Ar–Ar ages of 98–96 Ma [7], have been reported from mudstone and tuff (including phyllite, tuffaceous phyllite, and pelitic schist) in the ‘Mikabu greenstone belt’ or ‘Mikabu unit’. However, their geotectonic meanings were not clear, as these rock types are not characteristic of the Mikabu Greenstones.

In this paper, we apply the K–Ar dating method to weakly metamorphosed accretionary complexes in the northern Kanto Mountains, central Japan, where the original structural sequence of the Sanbagawa–Chichibu Belts is well-preserved. The K–Ar ages of white mica in the Sanbagawa schists have been previously reported in this area [20,47], but little is known about metamorphic ages in the Northern Chichibu Belt and the Mikabu Greenstones. One of the problems when conducting K–Ar dating using low-grade metasedimentary rocks is the sample heterogeneity due to incomplete recrystallization. To examine the effects of detrital mica [60] and establish a reliable dating method for weakly metamorphosed materials, we measured whole-rock and white mica mineral ages for samples of various lithologic types, including mudstone, acidic tuff, and chert. In low-grade metasedimentary rocks, illite is generally the most dominant phase of

86 white mica, and its crystallinity has been used to estimate the maximum temperature
87 they attained [5,8,17,18,40]. In this study, we quantified the illite crystallinity of the
88 dated samples, in order to constrain their thermal histories and factors affecting the
89 illite K–Ar ages. Finally, we estimate the peak metamorphic age of each geological unit,
90 based on most reliable data set of the K–Ar ages, and discuss the tectonic evolution of
91 the Sanbagawa–Chichibu Belts.

92 **2. Outline of Geology**

93 The Northern Chichibu Belt in the study area (Figure 2) comprises three lithologi-
94 cally distinct units: the Kashiwagi, Manba, and Kamiyoshida Units, in ascending order
95 of structural level [49,56,57].

96 The Kashiwagi Unit is characterized by coherent layers of red chert, red shale, and
97 basic tuff in the lower sub-unit, and by alteration of acidic tuff, mudstone, and sandstone
98 in the upper sub-unit [56]. Upper Triassic limestone occasionally occurs as lenses and
99 beds [52]. Based on the radiolarian assemblages in mudstone, the sedimentation age of
100 the upper sub-unit of the Kashiwagi Unit has been constrained to during the Late Jurassic
101 and the Early Cretaceous [43,44,55]. The Manba and Kamiyoshida Units are considered
102 as Middle Jurassic accretionary complexes characterized by block-in-matrix structures.
103 The Manba Unit comprises chaotic blocks of pillow lava, pillow breccia, and basic tuff
104 associated with Carboniferous–Permian limestone [49,51], as well as chert blocks of
105 the Middle Triassic age [33]. Early to Middle Jurassic radiolaria have been reported
106 from mudstone (shale) in the western area of the Northern Kanto Mountains [33,44,54].
107 The Kamiyoshida Unit consists of mudstone of the early Middle Jurassic age [22,23,44],
108 sandstone, siliceous shale, and chert of the Triassic age, as well as Permian limestone
109 [33,51]. The type locality of the Mikabu Greenstones [39,66] is located at Mts. Mikabu
110 (Mikabo), where the meta-basaltic body associated with pillow lava, hyaloclastite, and
111 doleritic intrusives are exposed. The red shale on the top of hyaloclastite comprises late
112 Middle Jurassic to Late Jurassic radiolaria [45].

113 The metamorphic grade of the basic rocks in the Northern Chichibu Belt and the
114 Mikabu Greenstones increases toward the north (i.e., toward the lower structural level)
115 up to the pumpellyite–actinolite facies zones [19] (Figure 2). The metamorphic grade of
116 the Kamiyoshida Unit remains undetermined, due to a lack of basic rocks. Magnitudes
117 of strain measured by the shapes of radiolarian fossils and amygdules in pillow lava
118 increase toward higher metamorphic grade zones [56,57].

119 The Manba and Kamiyosida Units are sometimes treated as a single stratigraphic
120 unit, re-defined together as the ‘Kamiyoshida Unit’ [21,63], which is supposed to overlie
121 the Kashiwagi Unit with a fault contact [28,34]. Guidi et al. [14,15] have considered
122 that the Manba and Kamiyosida Units, together with the Mikabu Greenstones around
123 Mts. Mikabu, constitute a superficial nappe that covers the Sanbagawa Belt and the
124 Kashiwagi Unit. However, Hirajima and Banno [19] have contradicted this idea, based
125 on the findings of high-pressure minerals such as sodic-amphiboles and lawsonite in the
126 region where the superficial nappe was postulated. Shimizu [56] has considered that
127 the Kamiyoshida Unit constitutes a post-metamorphic nappe, which overlies both the
128 Manba and Kashiwagi Units. Herein we follow the classical devision of the Manba and
129 Kamiyoshida Units and investigate the relationships between each structural units.

130 **3. Samples**

131 Rock samples of mudstone, acidic tuff, basic tuff, chert, and red shale were collected
132 from the outcrops shown in Figure 2. Mudstone was considered to include black shale
133 (in the Manba and Kamiyoshida Units) and phyllite (in the Kashiwagi Unit). In two
134 outcrops along the Kannagawa River, samples were collected across the boundary region
135 of the Manba–Kashiwagi Units. As the exact location of the boundary between the
136 two units is controversial [34], the samples taken from the boundary zone between the
137 Manba and Kashiwagi Units are marked with asterisks (*) in Table 1 and Table 2. Sample

138 27-01 and 27-02 are mudstone (black shale) of the Manba Unit and bedded chert of the
139 Kamiyoshida Unit taken from the hanging wall and the footwall sides of a thrust fault
140 (Fig. 5 of Shimizu [56]), respectively.

141 At the type locality of the Mikabu Greenstones around Mt. Nishi-Mikaboyama,
142 basic lava and pyroclastic rocks are conformably overlain by red shale with a thickness
143 of more than 20 meters, containing a large number of deformed radiolarian fossils [56].
144 Sample 28-01 was collected from the roadside outcrop of this red shale. The red shale
145 layer extends to the west around Mt. Odokeyama [15], where Middle Jurassic to Late
146 Jurassic radiolaria have been reported [45]. In this outcrop, red shale was intercalated
147 with a whitish acidic tuff layer with a thickness of 30 cm. The sample Mi-10 of acidic tuff
148 was taken from this outcrop (Figure 3).

149 All the samples contained quartz, white mica, and chlorite in thin sections. White
150 mica and chlorite grains with length of about a few microns constitute slaty cleavage or
151 schistosity planes. In mudstones, detrital mica with larger grain sizes (20–50 μm) was
152 also observed. No detrital mica was identified in chert and red shale under an optical
153 microscope. Acidic tuff was found to contain larger feldspar grains in fine-grained
154 quartz matrix.

155 **4. Analytical Methods**

156 Rock samples were broken into fragments and powders using a jaw crusher and
157 a mortar mill, and then washed with ion-exchanged water several times. Rock frag-
158 ments of ca. 2 mm size were separated by hand-picking and used for the whole-rock
159 analysis. White mica was extracted from crushed samples in ion-exchanged water using
160 supersonic waves and fractionated into four grain-size classes using a centrifuge (0.5–1
161 μm , 1–2 μm , 2–3 μm , and 3–4 μm , according to Stokes law), then were freeze-dried. In
162 this paper, white mica separated in this way are conventionally referred to as ‘illite’,
163 although other mica species of detrital origin might have been included in the powders.

164 K–Ar dating was conducted with a mass spectrometer and associated analytical
165 system at the Research Institute of Natural Sciences, Okayama University of Science
166 [30]. An argon 38 spike was used in the isotopic dilution method. The K-content
167 was determined by flame photometry (HITACHI130-80), where Cs was added as a
168 buffer to reduce interference from other elements. The standard sample was a weakly-
169 metamorphosed rock, named 82A, whose potassium contents were previously measured
170 in the same laboratory. As the K-contents of some samples are very low, the potassium
171 weight percent was analyzed using ultra low blank chemical lines [29]. The radiometric
172 ages were calculated using the decay constants given by Staiger and Jäger [58].

173 The illite crystallinity was investigated using the Mac Science MXP3 X-ray diffrac-
174 tometer (XRD) at the University of Tokyo. All X-ray scans were conducted with $\text{CuK}\alpha$
175 radiation at 40 kV and 20 mA under the following machine settings: Divergent slit angle
176 of 1° , scan range of $7.5^\circ\text{--}10^\circ 2\theta$, step size ($\Delta 2\theta$) of 0.02° , and 2θ scan rate of 1.2° per
177 minute. The illite crystallinity index, CI, is defined by a full width at half maximum
178 (FWHM) of an X-ray diffraction peak for the basal (001) plane of illite with 10\AA spacing
179 [40]. Large CI indicates low crystallinity of the illite. The FWHM values were determined
180 after smoothing and background correction of the diffraction data.

181 **5. Results**

182 *5.1. K–Ar ages*

183 *5.1.1. Chert and red shale*

184 The results of K–Ar dating for the chert and red shale are summarized in Table
185 1 and Figure 5. The measurement errors in each sample were about ± 3 Ma with 2σ
186 confidence level. Hereafter, we describe the K–Ar age data in terms of the average values.
187 The whole-rock ages of chert and red shale in the Kashiwagi Unit were 133 Ma and
188 115 Ma, respectively. The red shale on the top of the Mikabu Greenstones yielded a
189 whole-rock age of 124 Ma. The whole-rock ages of chert varied from 168 Ma to 121 Ma

190 in the Manba Unit, and from 156 Ma to 125 Ma in the Kamiyoshida Unit. The Illite K–Ar
191 ages of chert and red shale yielded younger ages for smaller grain-size classes, most of
192 which were younger than the whole-rock age of the same sample. The mineral ages of
193 illite in chert and red shale varied in the ranges of 165–107 Ma in the Kshiwagi Unit,
194 154–127 Ma in the Manba Unit, and 150–120 Ma in the Kamiyoshida Unit. The ages
195 obtained for smaller grain-size classes yielded systematically younger ages.

196 5.1.2. Mudstone and tuff

197 The results of K–Ar dating of mudstone, acidic tuff, and basic tuff are summarized
198 in Table 2 and Figure 6. The measurement errors for each fractionated sample were
199 within ± 3 Ma with 2σ confidence level. All data, obtained for fractionated illite samples,
200 indicated younger ages for decreasing grain size (as observed for chert and red shale).
201 The illite ages of the acidic tuff associated with the Mikabu Greenstones were obtained
202 as 113–89 Ma. The illite K–Ar ages of the Kashiwagi and Manba Units were determined
203 in small ranges for each samples (within 20 My). Overall, the mudstone and acidic tuff
204 in the Kashiwagi Unit were dated as 117–100 Ma, and mudstone, acidic tuff, and basic
205 tuff in the Manba Unit were dated as 132–97 Ma. The two mudstone samples taken from
206 the Kamiyoshida Unit exhibited larger age variations (up to 33 My), ranging from 163
207 Ma to 121 Ma.

208 5.2. Illite Crystallinity

209 The CI values were determined for all fractionated samples of illite (Tables 1 and
210 2). Each sample exhibited smaller CI for larger grain size classes (Figure 7), although
211 there were some exceptions. The minimum CI in each sample was mostly around 0.4,
212 but acidic tuff samples taken from the Mikabu Greenstones and the Kashiwagi Unit
213 displayed smaller values (i.e., high crystallinity), around 0.3. No remarkable difference
214 was found between the CI values obtained for mudstone and chert.

215 5.3. Mechanical crushing test

216 To examine possible effects of Ar-loss during sample preparation, we conducted
217 a mechanical crushing test on a powdered and fractionated sample of acidic tuff (L-7).
218 The illite sample of the 3–4 μm class, which originally yielded a K–Ar age of 116 Ma,
219 was ground with an agate mortar, then fractionated into 1–2 μm , 2–3 μm , and 3–4 μm
220 size classes. The results of the K–Ar dating and the CI measurement are listed as L-7R
221 in Table 2. Figure 4 compares the K–Ar ages and CI values of illite before and after
222 refinement. Notably, no significant loss of Ar was detected in this test. Hence, the
223 possibility of Ar loss and change in illite crystallinity during mechanical crushing and
224 gliding in the laboratory are not considered in the following discussion.

225 6. Discussion

226 6.1. Interpretation of K–Ar ages

227 6.1.1. Chert and red shale

228 The K–Ar ages of chert and red shale exhibited large variation (Table 1). The
229 chert and red shale originated from pelagic sediments, which are expected to be almost
230 free from terrestrial materials. Hence, contamination of detrital mica would not have
231 seriously affected the K–Ar dating of these rocks. Nevertheless, the whole-rock ages of
232 chert in the Kamiyoshida and Manba Units showed large variation, and the oldest age
233 obtained from the Manba Unit (168 Ma) was very close to the timing of accretion at the
234 Middle Jurassic (>165 Ma; Figure 5). The illite samples separated from chert and red
235 shale also showed large variations, and the whole-rock measurements yielded relatively
236 old ages, compared with the illite mineral ages of the same sample (Figure 7a). The
237 K–Ar ages determined for chert and red shale in the Kashiwagi Unit also presented large
238 variations. The oldest one obtained for the largest grain-size class of illite yielded an age
239 before the Late Jurassic accretion age. The mineral ages of illite determined for chert

blocks (28-03A, 28-03B) in the Manba Unit were older than those of the surrounding mudstone matrix (28-04) at the same outcrop (Figure 8a). Similarly, the mineral ages for the largest grain size group of illite separated from red shale (27-07) displayed older ages than those separated from acidic tuff (L-7) in the same outcrops (Figure 8b).

The micro-fossil data in chert blocks of the Kashiwagi, Manba, and Kamiyoshida Units indicate deposition ages from Carboniferous to Triassic, whereas those in mudstone indicated younger ages of accretion, during the Middle to Late Jurassic, as summarized in the previous section. One possible explanation is that the K–Ar ages were affected by submarine processes at the time of deposition; for example, hydrothermal activity and submarine metasomatism-associated volcanism of ridges or seamounts may lead to the precipitation of illite or its precursor K-bearing minerals.

6.1.2. Mudstone and tuff

Compared with the ranges of K–Ar ages for chert and red shale, the illite ages of mudstone, acidic tuff, and basic tuff in each samples, and for each structural unit (Figure 6), were concentrated in narrow ranges (except for mudstone in the Kamiyoshida Unit). In the following discussion, we focus on the illite ages obtained from these rocks. Among the illite ages determined for different grain size classes of the same rock sample, larger illite grains yielded older ages and smaller CI values, although there were some exceptions (Figure 7b). Contamination of detrital mica, formation of new clay minerals, and brittle deformation in the retrograde stages are possible causes of the systematic changes in the K–Ar ages and illite crystallinity. In the present case, the mechanical crushing test exhibited no detectable changes in the K–Ar ages and CI values (Figure 4). Furthermore, the illite ages and CI values of the mudstone of the Manba Unit beneath the fault boundary (27-01) were similar to those obtained from other mudstone samples of the Manba Unit. Thus, we consider that the K–Ar age data and CI values were not strongly affected by brittle deformation in the retrograde stages. Grain-size dependence of the illite K–Ar ages and CI was observed not only in mudstone, but also in pelagic sediments (chert and red shale) and tuffs. Acidic tuff of the Kashiwagi Unit and Mikabu Greenstones had Illite grains with small CI (i.e., high crystallinity), which are thought to have crystallized from volcanic glass and feldspars during metamorphism. The illite K–Ar ages determined for mudstone and acidic tuff in the same outcrops (26-01A and 26-01B; 27-05 and 27-06) yielded similar ages. This suggests that the influence of detrital mica on the K–Ar dating was negligible.

The white mica in the low-grade metamorphic rocks was actually illite–smectite interstratified clay minerals, with CI values mainly controlled by the layer thickness of illite crystallite domains and the percentage of the smectite layers [32]. Kübler [40] has proposed to use the CI of the $<2\ \mu\text{m}$ sized fraction of clay samples as an indicator of metamorphic grade. Although illite crystallinity could be affected by many kinetic factors other than temperature, such as duration of metamorphism and chemical environment of the host rock, Kübler's index has been widely applied to estimate the peak temperature of the incipient to low-grade metamorphic rocks. Underwood et al. [67] have measured the Kübler's CI in the Cretaceous accretionary complex of the Shimanto Belt, SW Japan, and correlated CI and the peak metamorphic temperature T_{vtr} (°C), quantified by the vitrinite reflectance as

$$CI = 1.197 - 0.0029 T_{vtr}. \quad (1)$$

Using this equation, we define the apparent metamorphic temperature T_{CI} (°C) as

$$T_{CI} = (4.13 - 3.45 CI) \times 10^2. \quad (2)$$

The T_{CI} values calculated for each fractionated samples of illite are also listed in Tables 1 and 2. Taking the 1–2 μm size class illite samples, the CI values of mudstone in the Kamiyosida Unit (0.44–0.45) corresponded to T_{CI} of 256–260°C, while those of the Manba

Unit (0.40–0.45) corresponded to 259–277°C. Mudstone at the uppermost part of the Kashiwagi Unit yielded a relatively large CI (0.45), corresponding to 257°C; whereas acidic tuff in the Kashiwagi Unit (CI = 0.30–0.34) and the Mikabu Greenstones (CI = 0.31) yielded T_{CI} of 295–310°C. These results are roughly consistent with the mineral assemblages of the Kashiwagi Unit and the Mikabu Greenstones, which belong to the pumpellyite–actinolite facies zone (ca. 250°C–350°C) [19]. The peak metamorphic temperatures of mudstones in the eastern area of the Kanto Mountains estimated by the Raman spectroscopy of carbonaceous materials (315 °CC for the Mikabu Greenstones, 252–268 °CC for the Kashiwagi Unit, and ca. 260 °CC for the Kamiyoshida Unit) [64] are in reasonable agreement with the T_{CI} values of each units. In the south-east extension of the Kashiwagi Unit (Hashidate Unit), CI values of 0.23–0.33, corresponding to T_{CI} of 299–334 °C, have been reported from mudstone [17]. In the preparation of the XRD samples, we did not strictly follow the recommended procedure for the Kübler’s index [41]. The relatively smaller values of CI in our results might be partly due to the difference in sample preparation method.

As discussed above, the grain-size dependence of the K–Ar ages and CI observed in the samples of various lithology were not attributable to contamination of detrital muscovite. The younger ages and larger CI values observed for fine-grained illite would be more reasonably explained by their formation in the retrograde stages. The ranges of the illite ages in each sample possibly reflect the cooling history of each structural unit. Figure 9 tentatively plots the change in apparent metamorphic temperature of each fractionated sample of illite against its K–Ar age. It should be noted that kinetic factors other than temperature, such as duration of metamorphism and chemical environments, might have been affected the CI values. The illite K–Ar ages of acidic tuff (Mi-10) at the type locality of the Mikabu Greenstones showed deviation toward younger ages in the small grain-size classes (<2 µm). As the sample was located at a small outcrop along a mountain pass, small illite grains might have been influenced by Ar loss due to weathering on the ground surface.

6.1.3. Peak metamorphic ages

In previous works, the illite K–Ar ages obtained from the Northern Chichibu belt have been interpreted as the crystallization ages at the peak metamorphism [25], as the peak metamorphic temperatures were lower than the closure temperature of muscovite (ca. 350°C) [25]. The illite ages of the Mikabu Greenstones, which belongs to the pumpellyite–actinolite facies zone (ca. 250°C–350°C), were also considered as the peak metamorphic ages. In the present study, however, different K–Ar ages were obtained for illite in different size classes. The variation in K–Ar age likely reflects the cooling history of each structural unit, as discussed above. Hence, the illite ages obtained for the 3–4 µm class were taken to determine the timing of peak metamorphism (Figure 10).

Using the illite mineral ages obtained from mudstone, acidic tuff, and basic tuff, the Kashiwagi Unit was dated to 117–110 Ma, and the Manba Unit was dated to 132–107 Ma. The age of mudstone previously reported from the low-grade sub-zone of the pumpellyite–actinolite facies zone (117 Ma), using a conventional K–Ar dating method [20] is consistent with the determined peak metamorphic ages of the Kashiwagi Unit. The K–Ar age of tuffaceous phyllite (110 Ma) in the ‘Mikabu Unit’ in the eastern area the Kanto Mountains [62] can be correlated with the ages of acidic tuff in the Kashiwagi Unit determine in the present study. The peak metamorphic age of the red shale on the top of the ancient Mikabu seamount is 113 Ma according to the K–Ar dating of the intercalation of acidic tuff.

Two mudstone samples taken from the Kamiyoshida Unit were dated to 163 Ma and 144 Ma, based the illite ages of the 3–4 µm class. These ages were remarkably older than the illite ages of the Kashiwagi and Manba Units. This supports the tectonic interpretation of Shimizu [56], that the Kamiyoshida Unit is a different tectonic unit

covering the Kashiwagi and Manba Units with a major discontinuity. It should be noted, however, that the illite ages presented very large variation, depending on the grain size, and that the oldest age (163 Ma) was very close to the accretion age of the early Middle Jurassic [33] (Figure 11). Influence of contamination of older white mica might be relatively important in these very low-grade sedimentary rocks, but the CI values of these samples did not show markedly different values for different grain sizes (Figure 7b). As the rock samples of mudstone taken from the Kamiyoshida Unit comprised small fragments of chert breccia, contamination of older materials of chert fragments possibly affected the K–Ar ages. Thus, more detailed study is required to specify the peak metamorphic age of the Kamiyoshida Unit.

6.2. Tectonic evolution of the Sanbagawa–Chichibu Belts

6.2.1. Subduction of the ancient Mikabu seamount

In the Sanbagawa Belt to the north-east of the study area, the K–Ar ages of white mica monotonously increase toward the upper structural unit; the K–Ar ages of white mica in pelitic, psammitic, and basic schists in the chlorite, garnet, and biotite zones of the Sanbagawa schists in the Kanto Mountains range from 84–72 Ma, 82–58 Ma, and 67–53 Ma, respectively [20] [47] (Figure 11). The illite K–Ar age of the Mikabu Greenstones (115 Ma) determined in the present work was definitely older than the muscovite ages cited above. As the peak metamorphic temperatures in the high-grade zones exceed the closure temperature of muscovite, the younger ages in the biotite and garnet zones might be partly attributed to the difference in cooling ages. However, the metamorphic temperature achieved in the chlorite zone would not significantly differ from that in the Mikabu Greenstones (up to 325°C, according to T_{CI}). Hence, a chronological gap between the Mikabu Greenstones and the Sanbagawa Belt can be inferred. The K–Ar age of the ‘Mikabu unit’ (82 Ma), previously reported from the Kanto Mountains by Miyashita and Itaya [47] (Figure 2), likely represents the metamorphic age of the Sanbagawa Belt. The peak metamorphic age of 115 Ma determined at the type locality of the Mikabu Greenstones indicated that the timing of underplating of the Mikabu seamount was contemporaneous with the peak metamorphism of the Kashiwagi Unit (117–110 Ma) in the Early Cretaceous. This supports the idea that the Mikabu Greenstones and the oceanic and terrigenous sediments of the Kashiwagi Unit constitute the same oceanic plate stratigraphy [46,56].

The peak metamorphic ages estimated from the K–Ar ages of illite in the Manba Unit have bimodal concentrations around 115 Ma and 130 Ma. Notably, the younger ages ca. 115 Ma overlap with the peak metamorphic age of the Kashiwagi Unit, although there is a large gap in the accretion ages determined from radiolarian fossils and detrital zircon (134–127 Ma) [63] between the Kashiwagi Unit (the Early Cretaceous) and the Manba Unit (the Middle Jurassic). These include data from samples taken from the boundary region of the Kashiwagi and Manba Units. A possible scenario is that the basal part of the Middle Jurassic strata that formerly constituted the accretionary wedge was scratched out, subducted with the Kashiwagi Unit, and metamorphosed together at the deeper part of the subduction zone. Tectonic erosion of the basal part of the accretionary wedge due to subduction of the Mikabu seamount may possibly account for the distribution of the Manba Unit.

6.2.2. Comparison with chronological structures in central Shikoku

The chronological structures of the Sanbagawa Belt in Shikoku have been extensively studied using K–Ar and Ar–Ar dating methods (Figure 11). One of the conspicuous features addressed in the NS cross-section of the Sanbagawa Belt in Shikoku is the positive correlation between metamorphic grades and muscovite mineral ages [25]. Takasu and Dallmeyer [61] have explained chronological gaps by nappe structures, and considered that the main suite (Besshi Unit) of the Sanbagawa Belt thrusts over a younger unit (Oboke Unit). Aoki et al. [1–3] have divided the Besshi Unit into two

sub-units, based on protolith ages, but their chronological model was contradicted by recent U–Pb zircon dating evidence [37,38].

Aoki et al. [2,28] have correlated the Sanbagawa schists in the Kanto Mountains with the Oboke Unit in a broader sense (i.e., the ‘Shimanto metamorphic belt’, in their definition). However, unlike the Oboke Unit, coarse-grained turbiditic deposits are not observed in the Kanto Mountains, and the K–Ar and Ar–Ar ages of the Oboke Unit (70–63 Ma) [1,31,61] are significantly younger than the K–Ar ages reported from the same metamorphic grade zone in the Kanto Mountains (i.e., 84–76 Ma in the chlorite zone). Herein, we define the Sanbagawa schists in the Kanto Mountains as the ‘Ayukawa Unit’ of the Sanbagawa Belt, and distinguish it from the Besshi and Oboke Units in Shikoku.

The main constituent of the Ayukawa Unit are pelitic and basic schists, which are intercalated with siliceous and psammitic schists. These lithological features are similar to those of the Bessi Unit, except for eclogite bodies in the highest grade zone. Furthermore, the range of the muscovite K–Ar ages (83–53 Ma) in the Ayukawa Unit overlaps with those reported in the Bessi Unit (88–72 Ma); however, the inverted thermal structure is absent and the correlation between the metamorphic grades and K–Ar and Ar–Ar ages has opposite sense in the Kanto Mountains. The Ar–Ar ages in ‘the southern marginal belt’ to the south of the Kiyomizu Tectonic Line in Shikoku [4,7] range from 87 to 97 Ma [7], and the K–Ar ages have also been found to be concentrated around 90 Ma [25]. Previously reported phyllite ages of the ‘Mikabu greenstone belt’ (98–96 Ma) [7] may represent the metamorphic age of the southern marginal belt.

In the Northern Chichibu Belt in Shikoku, the western extensions of the Kashiwagi and Kamiyoshida Units have been identified based on lithology and radiolarian fossil ages [9,44]. The illite K–Ar ages reported from mudstone in the Kamiyoshida Unit in central Shikoku range from 159 Ma to 128 Ma, concentrated around 140 Ma; while those reported from mudstone and basic tuff in the Kashiwagi Units in central Shikoku range from 124 Ma to 104 Ma and are concentrated around 115 Ma [35]. These chronological data are consistent with the present results in the Kanto Mountains at the type localities of the Kamiyoshida and Kashiwagi Units. Dallmeyer et al. [7] have applied the Ar–Ar dating method to mudstone (phyllite) in the Northern Chichibu Belt. They have reported whole-rock ages of ca. 110 Ma in the northern part, 120–140 Ma in the middle part, and ca. 215 Ma in the southern part; however, no clear relationships could be recognized between the chronological data and the stratigraphic units [4]. Isozaki et al. [25,27] have considered that the northern part of the Northern Chichibu complex, comprising the Late Jurassic to Early Cretaceous radiolaria, is a relatively in-situ unit which underwent the Sanbagawa metamorphism at ca. 115 Ma; meanwhile, the southern part is a nappe of the Middle Jurassic accretionary complex, which originated in the Mino–Tanba Belt in the Inner Zone of SW Japan. The geological unit equivalent with the Manba Unit is missing (or overlooked) in Shikoku. The present results obtained from mudstone and tuff in the Kashiwagi and Kamiyoshida Units in the Kanto Mountains were consistent with the K–Ar ages obtained in Shikoku; however, there were no large gaps in the metamorphic ages between the Late Jurassic complex (Kashiwagi Unit) and the Middle Jurassic complex (Manba and Kamiyoshida Unit). The presented results indicated that the chronological discontinuity defined by K–Ar and Ar–Ar dating did not strictly correlate with a difference in accretion age. It is possible that the Manba Unit underplated at a deep part of the accretionary wedge, while the Kamiyoshida Unit accreted in a shallow part and later juxtaposed with the Manba Unit with an out-of-sequence thrust.

Regarding the protolith ages of the Sanbagawa schists, Aoki et al. [1] have reported Late Cretaceous detrital zircon ages (82 ± 11 Ma and 92 ± 4 Ma) from psammitic schists in the Oboke Unit, indicating that its deposition age is no older than 93 Ma. Detrital zircon of similar age (90–80 Ma) has also recently been reported in the Besshi Unit [37,38]. In the Ayukawa Unit, Tsutsumi et al. [65] have reported detrital zircon ages of 91.4 ± 1.4 Ma and 95.3 ± 1.5 Ma from psammitic schists in the chlorite zone, and 78.8 ± 1.3 Ma from

those in the biotite zone. Hence, the subduction of the host rock of the chlorite zone occurred no earlier than 97 Ma, while that of the biotite zone was no earlier than 80 Ma. Compared with the K–Ar ages of white mica reported in the Ayukawa Unit [20,47], which represents the timing of the peak metamorphism (for the chlorite zone) or cooling ages (for the biotite zones), the duration from deposition to the peak metamorphism in each zone was estimated to be 20–21 My, at most. The deposition of the protolith of the biotite zone occurred during or after the peak metamorphism of the chlorite zone (84–72 Ma). In the Northern Chichibu Unit in the Kanto Mountains, Tominaga et al. [63] have reported the youngest detrital zircon age of 128.2 ± 1.4 Ma and 126.7 ± 2.0 Ma from sandstone, and 134.2 ± 1.5 Ma from acidic tuff in the Kashiwagi Unit. Accordingly, the duration between deposition and the peak metamorphic age (117–109 Ma) was estimated to be 20 My, at most.

In the Kanto Mountains, K–Ar age data between 107 Ma and 84 Ma are missing, and the deposition of the protolith sandstone of the chlorite zone (<93 Ma) post-dated the peak metamorphism of the Kashiwagi Unit (ca. 117–109 Ma). Hence, there exists a chronological gap between the Northern Chichibu Belt (including the Mikabu Greenstones) and the Sanbagawa Belt. However, regarding the Sanbagawa marginal belt as the upper part of the Ayukawa Unit, the chronological gap between the Chichibu–Sanbagawa Belt, in the classical sense, can be reduced to less than 15 My. Thus, the major discontinuity between the Kashiwagi and Ayukawa Units, as proposed by Isozaki et al. [28], is questionable—in their terminology, the Kashiwagi Unit belongs to the ‘Sanbagawa metamorphic belt *sensu strictu*’ and the Ayukawa Unit belongs to the ‘Shimanto metamorphic belt’. Rather, the evolution of the Cretaceous metamorphic zone can be understood as the progressive growth of accretionary complexes with multiple cycles of subduction and metamorphism events; the ‘Sanbagawa metamorphic belt’ can be re-defined in this sense. The Kamiyoshida Unit likely experienced older metamorphism (before 130 Ma), and may be excluded from the Sanbagawa metamorphic belt.

7. Conclusions

K–Ar dating was conducted using low-grade meta-sedimentary rocks from the Northern Chichibu Belt and the Mikabu Greenstones in the Kanto Mountains. The measurement results revealed the following:

1. The whole-rock and illite mineral ages of chert and red shale showed large variations. Overall, the K–Ar ages of chert and red shale were significantly older than those of mudstone, acidic tuff, and basic tuff in the same stratigraphic units, although pelagic materials of chert and red shales were almost free of detrital mica. Thus, it is possible that the K–Ar ages of chert and red shale reflect hydrothermal activities on the sea floor before subduction.
2. The fractionated illite samples of chert, red shale, mudstone, acidic tuff, and basic tuff displayed older ages for larger grain size. The observed changes in CI can be explained by formation of illite–smectite mixed-layer clays during retrograde stages, rather than contamination with detrital mica. The illite K–Ar ages of the largest grain size class (3–4 μm) obtained from mudstone and tuff likely represent the peak metamorphic ages.
3. Based on the K–Ar dating of mudstone, acidic tuff, and basic tuff, the peak metamorphic age of the Kashiwagi Unit was estimated as 117–110 Ma (Early Cretaceous), and a similar age (113 Ma) was obtained for an acidic tuff sample taken from the type locality of the Mikabu Greenstones. Hence, the Mikabu Greenstones can be considered as constituents of the Kashiwagi Unit in the Northern Chichibu Belt.
4. Among the Middle Jurassic accretionary complexes, the lower structural unit (Manba Unit) yielded peak metamorphic age in the Early Cretaceous (132–107 Ma); whereas the upper structural unit (Kamiyoshida Unit) showed apparently older ages, in the Late Jurassic (163–144 Ma). As such, the lower part of the Manba Unit can be interpreted as a re-worked part of the Middle Jurassic accretionary

500
501
502

complex, which acted as a shadow zone of the subducting Mikabu seamount. The
discontinuity in the peak metamorphic age between the Manba and Kamiyoshida
Units supports the post-metamorphic nappe model proposed by Shimizu [56].

503 8. Figures, Tables, and Schemes





Figure 1. An index map showing the tectonic division of the Chichibu and Sanbagawa Belts and distribution of the Mikabu Greenstones. The study area is indicated by an orange rectangle. The Median Tectonic Line (M.T.L.) and the Itoigawa–Shizuoka Tectonic Line (I.-S.T.L.) are denoted by a red line and a red dotted line, respectively.






Figure 3. A route map to the north-east of Mt. Odokeyama in Figure 2, showing the detailed sampling locality of acidic tuff (Mi-10) deposited on the top of the Mikabu Greenstones. The sampling locality of the radiolarian fossils in literature [45] is also plotted.




Figure 4. The illite K–Ar ages and CI values of a sample (L-7) of acidic tuff before (blue) and after (red) the crushing test. See text for details.




Figure 5. The peak metamorphic ages determined from the bulk and fractionated illite samples of chert and red shale. Green, red, blue, and brown symbols indicate the Mikabu Greenstones and the Kashiwagi, Manba, and Kamiyoshida Units, respectively.





Figure 6. The peak metamorphic ages determined from the fractionated illite samples of mudstone, acidic tuff, and basic tuff. Green, red, blue, and brown symbols indicate the Mikabu Greenstones and the Kashiwagi, Manba, and Kamiyoshida Units, respectively.

(a) chert and red shale



(b) mudstone and tuff




Figure 7. The K-Ar ages and crystallinity index (CI) of illite in chert and red shale (a), and in mudstone, acidic tuff, and basic tuff (b), determined for each grain-size class. Green, red, blue, and brown symbols indicate the data of the Mikabu Greenstones and the Kashiwagi, Manba, and Kamiyoshida Units, respectively.





Figure 8. The illite K–Ar ages (circles) and CI (triangle) of selected samples. (a) Comparison between chert blocks and the surrounding mudstone matrix in the Manba Unit. (b) Comparison between red shale and acidic tuff in the same outcrop of the Kashiwagi Unit.






Figure 10. The geological map of the Northern Chichibu Belt in the study area plotted with the peak metamorphic ages of mudstone, acidic tuff and basic tuff, determine from the K–Ar dating of 3–4 μm fractionated samples of illite. Symbols of sampling localities are the same as those in Figure 2.




Figure 11. Sedimentation, accretion and metamorphic histories of the Sanbagawa and Chichibu Belts in the Kanto Mountains and Shikoku. Rectangular areas and symbols show the ranges and individual age data, respectively. The K–Ar and Ar–Ar ages in the Sanbagawa Belt after [20,47] (Kanto Mountains) and [25,61] (Shikoku). K–Ar ages in the Northern Chichibu Belt in Shikoku after [35]. Radiolarian ages of mudstone and siliceous shales in the Kanto Mountains are based on [43,55] (Kashiwagi Unit), [33,54] (Manba Unit), and [22,23] (Kamiyoshida Unit) with re-interpretation by [44], and those in Shikoku based on [44]. Radiolarian ages of red shale are based on [45] (Kanto Mountains) and [12] (Shikoku). The volcanic ages of the Mikabu Greenstones refer to [50] (K–Ar age) and [62] (U–Pb age). The youngest detrital zircon ages refer to [65] (Ayukawa Unit) and [62] (Kashiwagi Unit) in the Kanto Mountains, and [1] (Oboke Unit), [37,38] (Besshi Unit), and [9] (Kashiwagi Unit) in Shikoku.

Table 1. The K–Ar ages of whole-rock samples and fractionated illite samples from chert and red shales taken in the Northern Chichibu Belt in the Kanto Mountains, Central Japan. The crystallinity index (CI) of illite and the apparent temperature calculated from the CI are also listed.

Geologic Unit	Sample	Rock type	Grain size (μm)	K (wt.%)	Rad. ^{40}Ar (10^{-8} cc STP/g)	Age (Ma)	CI ($^{\circ}\Delta 2\theta$)	T_{CI} ($^{\circ}\text{C}$)
Mikabu Greenstones Kashiwagi Unit	28-01	red shale	whole rock	0.43 ± 0.02	216 ± 2	123.6 ± 2.6		
	27-07	red shale	whole rock	0.60 ± 0.02	277 ± 3	115.3 ± 2.3		
			0.5–1	6.03 ± 0.12	2641 ± 26	106.9 ± 2.3	0.474	249.2
			1–2	5.26 ± 0.11	2574 ± 25	121.8 ± 2.5	0.452	257.1
			2–3	3.46 ± 0.07	1978 ± 17	129.3 ± 2.6	0.456	255.7
			3–4	1.95 ± 0.03	1304 ± 14	164.7 ± 3.3	0.422	267.4
	27-08	chert	whole rock	0.14 ± 0.01	75 ± 1	133.2 ± 2.7		
			0.5–1	1.35 ± 0.03	617 ± 18	114.4 ± 2.3	0.441	260.6
			1–2	1.46 ± 0.03	705 ± 15	120.5 ± 2.4	0.407	272.5
			2–3	1.54 ± 0.03	747 ± 16	121.2 ± 2.4	0.389	278.6
Manba Unit			3–4	1.09 ± 0.02	632 ± 8	143.3 ± 2.9	0.372	284.3
	26-05	chert	whole rock	0.17 ± 0.01	94 ± 1	141.3 ± 2.8		
	27-10	chert	whole rock	0.22 ± 0.01	116 ± 2	130.6 ± 2.6		
	27-12B	chert	whole rock	0.12 ± 0.01	60 ± 1	120.8 ± 2.4		
	28-03A	chert	whole rock	0.24 ± 0.01	163 ± 2	168.2 ± 3.8		
			0.5–1	4.82 ± 0.09	2495 ± 25	128.7 ± 2.5	0.411	270.9
			1–2	4.27 ± 0.08	2206 ± 22	128.5 ± 2.5	0.422	267.2
			2–3	3.09 ± 0.06	1782 ± 17	142.8 ± 2.8	0.455	256.0
			3–4	1.28 ± 0.03	787 ± 8	152.5 ± 3.1	0.421	267.7
	28-03B	chert	whole rock	0.35 ± 0.01	212 ± 2	148.4 ± 2.9		
			0.5–1	5.91 ± 0.18	3019 ± 31	127.1 ± 2.5	0.467	251.8
			1–2	5.6 ± 0.12	2997 ± 29	132.9 ± 2.6	0.389	278.6
			2–3	4.3 ± 0.08	2477 ± 24	142.4 ± 2.8	0.377	282.6
			3–4	2.38 ± 0.06	1483 ± 15	154.1 ± 3.1	0.372	284.4
Kamiyoshida Unit	27-02	chert	whole rock	0.33 ± 0.02	205 ± 2	152.7 ± 3.1		
			0.5–1	5.81 ± 0.12	2787 ± 27	119.6 ± 2.4	0.419	268.2
			1–2	5.30 ± 0.11	2860 ± 27	128.8 ± 2.5	0.388	278.9
			2–3	3.81 ± 0.07	2119 ± 21	136.3 ± 2.7	0.395	276.6
			3–4	1.64 ± 0.03	1003 ± 10	149.6 ± 2.9	0.375	283.5
	27-03	chert	whole rock	0.25 ± 0.01	161 ± 0.1	155.9 ± 3.1		
Unknown	28-02	chert	whole rock	0.13 ± 0.01	65 ± 0.9	124.7 ± 2.4		

Table 2. The K–Ar ages of fractionated illite samples of mudstone and tuff taken from the Northern Chichibu Belt in the Kanto Mountains, Central Japan. The crystallinity index (CI) of illite and the apparent temperature T_{CI} calculated from the CI are also listed. * Samples taken from the boundary zone between the Kashiwagi and Manba Units. † Sample for the crushing test.

Geologic Unit	Sample	Rock type	Grain size (μm)	K (wt.%)	Rad. ^{40}Ar (10^{-8} cc STP/g)	Age (Ma)	CI ($^{\circ}\Delta 2\theta$)	T_{CI} ($^{\circ}\text{C}$)
Mikabu Greenstones	Mi-10	acidic tuff	0.5–1	4.62 \pm 0.10	1639 \pm 17	89.2 \pm 2.0	0.312	305.3
			1–2	4.92 \pm 0.10	1889 \pm 20	96.4 \pm 2.1	0.305	307.4
			2–3	4.83 \pm 0.10	2050 \pm 21	106.2 \pm 2.5	0.287	313.7
			3–4	3.48 \pm 0.07	1582 \pm 17	113.4 \pm 2.5	0.256	324.6
Kashiwagi Unit	26-03	acidic tuff	0.5–1	6.61 \pm 0.13	2763 \pm 29	104.0 \pm 2.1	0.403	273.8
			1–2	6.75 \pm 0.13	3032 \pm 30	111.7 \pm 2.2	0.297	310.3
			2–3	6.21 \pm 0.12	2863 \pm 28	114.1 \pm 2.3	0.294	311.3
			3–4	5.38 \pm 0.11	2356 \pm 23	109.8 \pm 2.2	0.255	324.9
	27-05	acidic tuff	0.5–1	6.16 \pm 0.12	2455 \pm 32	99.9 \pm 2.0	0.321	302.2
			1–2	6.11 \pm 0.12	2515 \pm 25	103.0 \pm 2.1	0.297	310.3
			2–3	5.53 \pm 0.11	2326 \pm 23	105.3 \pm 2.1	0.281	315.8
			3–4	3.97 \pm 0.08	1727 \pm 17	108.9 \pm 2.2	0.362	287.9
	27-06	mudstone	0.5–1	6.69 \pm 0.13	2912 \pm 28	108.8 \pm 2.2	0.460	254.1
			1–2	5.88 \pm 0.11	2604 \pm 26	110.7 \pm 2.2	0.453	256.5
			2–3	4.66 \pm 0.10	2116 \pm 21	113.5 \pm 2.3	0.444	259.7
			3–4	3.63 \pm 0.07	1700 \pm 17	116.8 \pm 2.3	0.436	262.4
	L-7	acidic tuff	0.5–1	6.75 \pm 0.13	2853 \pm 30	105.8 \pm 2.1	0.406	272.7
			1–2	6.86 \pm 0.14	3120 \pm 32	111.9 \pm 2.2	0.342	294.7
			2–3	6.38 \pm 0.13	2901 \pm 29	113.5 \pm 2.3	0.312	305.1
			3–4	5.23 \pm 0.11	2522 \pm 25	116.2 \pm 2.3	0.337	296.5
	L-7R [†]	acidic tuff	1–2	6.86 \pm 0.14	3173 \pm 33	115.4 \pm 2.5	0.333	
			2–3	6.33 \pm 0.13	3085 \pm 31	119.1 \pm 2.6	0.329	
			3–4	4.87 \pm 0.10	2302 \pm 24	118.0 \pm 2.6	0.314	
Manba Unit	27-01	mudstone	0.5–1	7.08 \pm 0.14	2883 \pm 30	100.5 \pm 2.0	0.470	250.7
			1–2	7.04 \pm 0.14	3116 \pm 32	110.6 \pm 2.2	0.415	269.7
			2–3	6.77 \pm 0.13	3006 \pm 31	111.0 \pm 2.2	0.390	278.2
			3–4	4.75 \pm 0.09	2192 \pm 22	115.2 \pm 2.3	0.399	275.3
	26-01A*	acidic tuff	0.5–1	6.88 \pm 0.13	2832 \pm 28	103.1 \pm 2.0	0.426	266.0
			1–2	6.92 \pm 0.14	2928 \pm 28	105.9 \pm 2.1	0.421	267.6
			2–3	6.12 \pm 0.12	2743 \pm 26	111.9 \pm 2.2	0.424	266.5
			3–4	4.33 \pm 0.09	1966 \pm 19	113.4 \pm 2.2	0.320	302.5
	26-01B*	mudstone	27-01	6.67 \pm 0.14	2407 \pm 27	97.0 \pm 1.9	0.456	255.5
			1–2	6.26 \pm 0.12	2537 \pm 35	101.3 \pm 2.0	0.447	258.5
			2–3	5.02 \pm 0.11	2172 \pm 21	107.6 \pm 2.1	0.422	267.1
			3–4	4.49 \pm 0.09	1915 \pm 18	106.6 \pm 2.1	0.431	264.2
	26-04	mudstone	0.5–1	6.88 \pm 0.14	3334 \pm 35	120.8 \pm 2.4	0.452	256.9
			1–2	6.40 \pm 0.12	3204 \pm 33	124.7 \pm 2.5	0.439	261.3
			2–3	5.21 \pm 0.10	2724 \pm 28	129.9 \pm 2.6	0.398	275.7
			3–4	3.28 \pm 0.06	1743 \pm 18	132.2 \pm 2.6	0.428	265.3
	27-04	basic tuff	0.5–1	7.59 \pm 0.15	3775 \pm 36	123.8 \pm 2.7	0.468	251.6
			1–2	7.20 \pm 0.14	3684 \pm 40	127.3 \pm 2.6	0.429	264.9
			2–3	6.92 \pm 0.14	3563 \pm 34	128.1 \pm 2.6	0.385	279.9
			3–4	6.67 \pm 0.13	3466 \pm 33	129.1 \pm 2.7	0.398	275.7
	27-11	mudstone	0.5–1	6.44 \pm 0.12	3225 \pm 25	121.4 \pm 2.4	0.443	259.9
			1–2	6.35 \pm 0.12	3149 \pm 23	123.5 \pm 2.6	0.394	277.0
			2–3	5.24 \pm 0.10	2796 \pm 19	127.7 \pm 2.5	0.385	279.9
			3–4	4.78 \pm 0.09	2484 \pm 25	129.1 \pm 2.6	0.363	287.6
	28-04	mudstone	0.5–1	5.60 \pm 0.11	2311 \pm 24	103.4 \pm 2.1	0.425	266.3
			1–2	4.98 \pm 0.10	2153 \pm 22	108.1 \pm 2.2	0.402	274.2
			2–3	3.80 \pm 0.07	1684 \pm 17	110.9 \pm 2.2	0.394	277.0
			3–4	2.47 \pm 0.05	1126 \pm 12	113.8 \pm 2.4	0.368	285.8
Kamiyoshida Unit	L-11	mudstone	0.5–1	6.71 \pm 0.13	3219 \pm 34	129.6 \pm 2.6	0.464	252.6
			1–2	5.34 \pm 0.10	2948 \pm 30	136.9 \pm 2.8	0.454	256.2
			2–3	3.95 \pm 0.08	2363 \pm 24	148.1 \pm 2.9	0.436	262.6
			3–4	2.75 \pm 0.05	1814 \pm 19	162.7 \pm 3.1	0.440	260.9
	L-13	mudstone	0.5–1	6.11 \pm 0.12	2969 \pm 31	121.1 \pm 2.4	0.455	255.9
			1–2	5.56 \pm 0.12	2911 \pm 30	130.1 \pm 2.6	0.444	259.8
			2–3	4.26 \pm 0.08	2303 \pm 24	134.2 \pm 2.7	0.420	268.0
			3–4	3.09 \pm 0.06	1797 \pm 19	144.1 \pm 2.8	0.440	261.0

Author Contributions: Field survey and geological interpretation, I.S.; sample collection, Z.L., I.S., and T.I.; methodology, T.I.; investigation and data processing, Z.L.; writing, I.S. and Z.L.; visualization, I.S.; supervision, I.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: We are grateful to Shizuo Yoshida for valuable discussions and comments in the field work. We also thank Hideo Takayama for technical assistance in the XRD measurements, and Atsusi Miyashita for helpful discussions.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Aoki, K., Iizuka, T., Hidaka, T., Maruyama, S. and Terabayashi, M., Tectonic boundary between the Sanbagawa belt and the Shimanto belt in central Shikoku, Japan. *J. Geol. Soc. Jap.*, **2007**, *113*, 171–183.
2. Aoki, K., Maruyama, S., Isozaki, Y., Otoh, S. and Yanai, S., Recognition of the Shimanto HP metamorphic belt within the traditional Sanbagawa HP metamorphic belt: New perspectives of the Cretaceous–Paleogene tectonics in Japan. *J. Asian Earth Sci.*, **2011**, *42*, 355–369.
3. Aoki, K., Seo, Y., Sakata, S., Obayashi, H., Tsuchiya, Y., Imayama, T., Yamamoto, S. and Hirata, T., U–Pb zircon dating of Sanbagawa metamorphic rocks in the Besshi–Asemi-gawa region, central Shikoku, Japan, and tectonostratigraphic consequences. *J. Geol. Soc. Jap.*, **2019**, *125*, 183–194. doi: 10.5575/geosoc.2018.0051
4. Aoya, M. Geology of the Hibihara district. Chapter 2, Outline of geology. Quadrangle Series, 1:50,000, Geological Survey of Japan, AIST, p. 5–9, 2009 (in Japanese).
5. Awan, M. A. and Kimura, K. Thermal structure and uplift of the Cretaceous Shimanto Belt, Kii Peninsula, Southwest Japan: An illite crystallinity and illite b_0 lattice spacing study. *The Island Arc*, **1996**, *5*, 69–88.
6. Banno, S. The high-pressure metamorphic belts of Japan: A review. *J. Soc. America, Mem.* **1986**, *164*, 365–385.
7. Dallmeyer, R.D., Takasu, A. and Yamaguchi, K., Mesozoic tectonothermal development of the Sanbagawa, Mikabu and Chichibu belts, South-west Japan: evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ whole-rock phyllite ages. *J. Metamorph. Geol.* **1995**, *13*, 271–286.
8. DiTullio, L., Laughland, M. M., and Byrne, T. Thermal maturity and constraints on deformation from illite crystallinity and vitrinite reflectance in the shallow levels of an accretionary prism: Eocene-Oligocene Shimanto Belt, southwest Japan. *Geol. Soc. Am. Spec. Pap.*, **1993**, *273*, 63–82.
9. Endo, S., Wallis, S.R., Structural architecture and low-grade metamorphism of the Mikabu-Northern Chichibu accretionary wedge, SW Japan. *J. Metamorph. Geol.* **2017**, *35*, 695–716.
10. Faure, M., 1985, The pre-Cretaceous structure of the Outer belt of southwest Japan. *Tectonophysics*, **1983**, *113*, 139–162.
11. Faure, M. and Charvet, J., 1983. Tangential tectonics in the Chichibu zone from the example of E. Shikoku. *Proc. of Japan Acad.*, **1983**, *59*, 117–120.
12. Faure, M., Iwasaki, M., Ichikawa, K. and Yao, A., The significance of Upper Jurassic radiolarians in high pressure metamorphic rocks of SW Japan. *Jour. SE Asian Earth Sci. Lett.*, **1991**, *62*, 131–136.
13. Geospatial Information Authority of Japan, Digital Topographic Map 25000. Available online: <https://maps.gsi.go.jp/vector/> (accessed on 29th July 2019).
14. Guidi, A. and Charvet, J., A new structural interpretation of the Kanto Mountains, northern-west of Tokyo, Japan. *Bull. Soc. Geol. France* **1987**, *8*, 843–853.
15. Guidi, A., Charvet, J. and Sato, T., 1984. Finding of granitic olistolith and pre-Cretaceous radiolarians in the northwestern Kanto Mountains, Gunma Prefecture, Central Japan. *J. Geol. Soc. Jap.*, **1984**, *90*, 853–856.
16. Guidi, A., Charvet, J., Sato, T., Takizaka, S. Pre-Cretaceous tangential structures in the Kanto Mountains, (SW Japan): Preliminary results. *Bull. Soc. Geol. France*, **1984**, *298*, 307–312. (In French with English abstract).
17. Hara, H., Hisada, K., Kimura, K. Paleo-geothermal structure based on illite crystallinity of the Chichibu and Shimanto Belts in the Kanto Mountains, central Japan. *J. Geol. Soc. Jap.*, **1998**, *104*, 705–717.
18. Hara, H., Kimura, K. Estimation of errors in measurement of illite crystallinity: the limits and problems of application to accretionary complexes. *J. Geol. Soc. Jap.*, **2000**, *106*, 264–279 (in Japanese with English abstract).
19. Hirajima T and Banno S. 1989. Records of high pressure metamorphism in the so-called "superficial nappe" in the Chichibu belt, Japan. *Bull. Soc. Geol. France* **1989**, *3*, 661–664.
20. Hirajima, T., Isono, T. and Itaya, T., K-Ar age and chemistry of white mica in the Sanbagawa metamorphic rocks in the Kanto Mountains, Central Japan. *J. Geol. Soc. Jap.* **1992**, *98*, 445–455.
21. Hisada, K., Tominaga, K., Sekine, K., Matsuoka, L., Kato, K., Geology of the northern Chichibu belt in the Kanto Mountains, central Japan. *J. Geol. Soc. Japan* **2016**, *122*, 325–342.
22. Iijima H., Sekine K., Saito Y., Jurassic Radiolarians from the clastic rock unit of the northern part of the Chichibu Belt, Kanto Mountains, Central Japan. *Bull. National Sci. Mus., Ser. C*, **1993**, *19*, 81–89.

23. Iijima H., Sekine K., Saito Y., Geologic age of the Kamiyoshida Unit of the Chichibu Belt, Kanto Mountains, Japan. *Bull. National Sci. Mus., Ser. C*, **1994**, 22, 119–131.
24. Isozaki, Y., Jurassic accretion tectonics of Japan. *The Island Arc* **1997**, 6, 25–51.
25. Isozaki, Y. and Itaya, T. Chronology of Sanbagawa metamorphism. *J. Metamorph. Geol.* **1990**, 8, 401–411.
26. Isozaki, Y., Itaya, T. and Kawato, K. Metamorphic age of Jurassic accretionary complex in the Northern Chichibu belt, Southwest Japan. *J. Geol. Soc. Jap.*, **1990**, 96, 557–560. (In Japanese with English abstract)
27. Isozaki, Y., Maruyama, S., Studies on Orogeny based on plate tectonics in Japan and new geotectonic subdivision of the Japanese islands *J. Geography (Chigaku Zasshi)* **1991**, 100, 697–761. (In Japanese with English abstract)
28. Isozaki, Y., Maruyama, S., Aoki, K., Nakama, T., Miyashita, A., Otoh, S. Geotectonic subdivision of the Japanese islands revisited: Categorization and definition of elements and boundaries of Pacific-type (Miyashiro-type) orogen *J. Geography (Chigaku Zasshi)* **2010**, 119, 999–1053. (In Japanese with English abstract).
29. Itaya, T., Doi, M. and Ohira, T., Very low potassium analysis by flame photometry using ultra low blank chemical lines: an application of K-Ar method to ophiolites. *Geochem. J.* **1996**, 30, 31–39.
30. Itaya, T., Nagao, K., Inoue, K., Honjou, Y., Okada, T. and Ogata, A., (1991) Argon isotope analysis by newly developed mass spectrometric system for K-Ar dating. *Mineral J.* **1991**, 15, 203–221.
31. Itaya, T. and Takasugi, H., Muscovite K-Ar ages of the Sanbagawa Schists, Japan and Argon depletion during cooling and deformation. *Contrib. Mineral. and Petrol.* **1988**, 100, 281–290.
32. Jaboyedoff, M., Bussy, F., Kübler, B., Thelin, P. Illite “crystallinity” revisited. *Clays Clay Min.*, **2001**, 49, 156–167.
33. Kamikawa, Y., Hisada, K., Sashida, K. and Igo, H., 1997. Geology of the Nanmoku area in the Chichibu Terrane, the northwest part of the Kanto Mountains, central Japan. *Sci. Rep. Inst. Geosci. Tsukuba, Ser. B*, 18, 19–38.
34. Kanto Mountains Research Group. The northern subbelt of the Chichibu Belt along the Kanna River in the Kanto Mountains, Central Japan. *Earth Sci. (Chikyu Kagaku)*, **1994**, 48, 83–101. (In Japanese)
35. Kawato, K. Isozaki, Y., Itaya, T. Geotectonic boundary between the Sanbagawa and Chichibu belts in central Shikoku, Southwest Japan *J. Geol. Soc. Jap.* **1991**, 97, 959–975. (In Japanese with English abstract)
41. Kisch H.J. (1991) Mite crystallinity: recommendations on sample preparation, X-ray diffraction settings, and interlaboratory samples *J. Metamorph. Geol.*, **1991**, 9, 665–670.
37. Knittel, U., Walia, M., Suzuki, S., and Lee, Y. H.. U-Pb single grain zircon ages for Sanbagawa Metamorphic Rocks in central Shikoku (Japan): the Sanbagawa Belt re-united. *Okayama Univ. Earth Sci. Rep.*, **2018**, 25, 39–48.
38. Knittel, U., Walia, M., Suzuki, S., and Lee, Y. H. Late Cretaceous age of eclogite facies metamorphism of the Sanbagawa belt in the Asemi River area, Shikoku (SW Japan): Evidence from detrital zircon. *J. Asian Earth Sci.*, **2019**, 77, 48–51.
39. Koto, B. On the so-called crystalline schists of Chichibu. *J. Coll. Sci. Imp. Univ. (Japan)*, **1988**, 2, 77–141.
40. Kübler B. La cristallinité de l’illite et les zones tout a fait superieures du métamorphisme. In: étages tectoniques, *Colloque de Neuchâtel 1966, a La Baconniere, Neuchâtel* 105–121, **1967**.
41. Kisch, H. J. . Illite crystallinity: recommendations on sample preparation, X-ray diffraction settings, and interlaboratory samples. *J. Metamorph. Geol.*, **1991** 9, 665–670.
42. Matsuoka, K. Discovery of Jurassic radiolarians from an unknown formation of the Northern subbelt of the Chichibu Belt in Manba town, Gumma Prefecture, Japan. *Earth Science (Chikyu Kagaku)*, **1995**, 49, 346–351. (In Japanese).
43. Matsuoka, K. Late Jurassic radiolarians from the northern subbelt of the Chichibu Belt in Ogawa town, Saitama Prefecture, Central Japan. *Earth Science (Chikyu Kagaku)*, **1996**, 50, 251–255. (In Japanese).
44. Matsuoka, A., Yamakita, S., Sakakibara, M., Hisada K., Unit division for the Chichibu Composite Belt from a view point of accretionary tectonics and geology of western Shikoku, Japan., *J. Geol. Soc. Jap.*, **1998**, 104, 634–653. (In Japanese with English abstract)
45. Matsuoka, K., Late Jurassic radiolarians from red shale on the Mikabu greenstones in the northern margin of Kanto Mountains, Japan. *Earth Science (Chikyu Kagaku)*, **1999**, 53, 71–74. (In Japanese)
46. Matsuoka, K.: The Kashiwagi Unit if the Northern Chichibu Belt in the northeastern Kanto Mountains — Lithofacies, geologic age and oceanic plate stratigraphy —, *Earth Science (Chikyu Kagaku)*, **2013**, 67, 101–112.
47. Miyashita, A. and Itaya, T., K-Ar age and chemistry of phengite from the Sanbagawa schists in the Kanto Mountains, central Japan, and their implication for exhumation tectonics. *Gondwana Res.*, **2002** 5, 837–848.
48. Ozawa, H., Motoyama, S., Inoue, S., Kato, Y., Murata, M., Petrology of basic volcanics of the Mikabu greenstone complex in the eastern Shikoku. *Mem. Geol. Soc. Japan*, No. 52, 217–228, **1999** (In Japanese with English abstract).
49. Okubo, M., Horiguchi, M. Geology of the Mamba district. Quadrangle Sries, 1:50,000, Geological Survey of Japan, 66p., 1969. (in Japanese).
50. Ozawa, H., Murata M. and Itaya, T. Early Jurassic volcanism of the Mikabu belt: Evidence from K-Ar age of picritic basalt, Kurouch ultramafic mass, Kanto Mountains, Japan, *J. Geol. Soc. Jap.*, **1997**, 103, 1089–1092.
51. Sashida, K., Igo, H., Igo, H., Kakizawa, S., Hisada, K., Shibata, T., Tsukada, K., Nishimura, H.: On the Jurassic radiolarian assemblages in the Kanto district. *News of Osaka Micropaleontologists Spec. Vol.* **1982**, 5, 51–64 (In Japanese with English abstract).
52. Sato, T., Kakizawa, S., Kodato T.: Revision of stratigraphy and structure of the Sakahara Formation at its type locality. *J. Geol. Soc. Jap.* **1977**, 83, 631–637 (In Japanese with English abstract).

53. Sawada, H., Isozaki, Y., Aoki, S., Sakata, S., Sawaki, Y., Hasegawa, R., Nakamura, Y., . The late Jurassic magmatic protoliths of the Mikabu greenstones in SW Japan: a frag- ment of an oceanic plateau in the Paleo-Pacific Ocean. *J. Asian Earth Sci.* **2019**, *169*, 228–236.
54. Sekine K., Iijima H., Saito Y., Jurassic radiolarians from the Manba Unit of the Northern Chichibu Belt, Kanto Mountains, Japan. *Bull. National Sci. Mus., Ser. C*, **1995**, *21*, 1–10.
55. Sekine K., Iijima H., Saito Y., Jurassic Radiolarians from the low-grade metamorphic zone of the northern part of the Chichibu Belt, Kanto Mountains, Central Japan. *Bull. National Sci. Mus., Ser. C*, **2001**, *27*, 101–108.
56. Shimizu, I. Ductile deformation in the low-grade part of the Sambagawa metamorphic belt in the Kanto Mountains, Central Japan. *J. Geol. Soc. Jap.* **1988**, *94*, 609–628.
57. Shimizu, I. and Yoshida S., Strain geometries in the Sanbagawa Metamorphic Belt inferred from deformation structures in metabasite. *The Island Arc*, **2004**, *13*, 95–109.
58. Steiger, R.H., Jäger, 1977. Subcommission on geochronology: convention on the use of decay constants in geo- and cosmo- chronology. *Earth Planet. Sci. Lett.*, **2004**, *36*, 359–362.
59. Suzuki, S., and H. Ishizuka. Low-grade metamorphism of the Mikabu and northern Chichibu belts in central Shikoku, SW Japan: Implications for the areal extent of the Sanbagawa low-grade metamorphism. *J. Metamorph. Geol.*, **1998**, *16*, 107–116.
60. Takami, M., Isozaki, Y., Nishimaru, Y. and Itaya, T., Effect of detrital white mica and contact metamorphism to K-Ar dating of weakly metamorphosed accretionary complex — an example of Jurassic accretionary complex in eastern Yamaguchi Prefecture, Southwest Japan —. *J. Geol. Soc. Jap.*, **1993**, *99*, 545–563. (In Japanese with English abstract)
61. Takasu, A. and Dallmeyer, R.D., $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age constraints for the tectonothermal evolution of the Sanbagawa metamorphic belt, central Shikoku, Japan: A Cretaceous accretionary prism. *Tectonophysics* **1990**, *185*, 111–139.
62. Tominaga, K., Hara, H., Paleogeography of Late Jurassic large-igneous-province activity in the Paleo-Pacific Ocean: Constraints from the Mikabu greenstones and Chichibu accretionary complex, Kanto Mountains, Central Japan. *Gondwana Res.* **2021**, *89*, 177–192.
63. Tominaga, K., Hara, H., Tokiwa, T., Zircon U–Pb ages of the Kashiwagi Unit of the accretionary complex in the Northern Chichibu Belt, Kanto Mountains, central Japan. *Bull. Geol. Surv. Japan* **2019**, *70*, 299–314. (In Japanese with English abstract).
64. Hara,H.;Mori,H.; Tominaga, K.; Nobe, Y. Progressive low-grade metamorphism reconstructed from the Raman spectroscopy of Carbonaceous material and an EBSD analysis of quartz in the Sanbagawa metamorphic event, Central Japan. *Minerals* **2021**, *11*, 854. [https:// doi.org/10.3390/min11080854](https://doi.org/10.3390/min11080854)
65. Tsutsumi, Yukiyasu, et al. "SHRIMP U–Pb dating of detrital zircons from the Sanbagawa Belt, Kanto Mountains, Japan: Need to revise the framework of the belt." *J. Mineral. Petrol. Sci.*, **2009**, *104*,12–24.
66. Uchida, N. Major-element petrochemistry of lavas and tuffs from the Sanbagawa and Chichibu terrane — The northern Kanto Mountains, Central Japan—, *Seikei Ronso* **1981**, *20*, 1–138.
67. Underwood, M. B., Laughland, M. M., and Kang, S. M. A comparison among organic and inorganic indicators of diagenesis and low-temperature metamorphism, Tertiary Shimanto Belt, Shikoku, Japan. *Spec. Pap. Geol. Soc. Am.* *273*, **1993**, 45–45.
68. Watanabe, T., K–Ar ages of white mica fraction from the Susunai metamorphic rocks in Sakhalin, Far East Russian, *Fac. Sci., Hokkaido Univ., Ser. 5* **1992**, *23*, 282–286.