

Article

Elemental Laser-Plasma Analysis of Pointed Gourd Leaves for Diabetes Management

Tejmani Kumar¹, Prashant K. Rai², Abhishek K. Rai³, Nilesh K. Rai⁴, Awadhesh K. Rai¹, Christian G. Parigger^{5,*}, Geeta Watal², Suman Yadav⁶

¹ Laser Spectroscopy Laboratory, Department of Physics, University of Allahabad, Prayagraj- 211 002, India; tejmani38@gmail.com, awadheshkrai@gmail.com

² Alternative Therapeutics Unit, Drug Development Division, Medicinal Research Laboratory, Department of Chemistry; University of Allahabad, Prayagraj – 211 002, India; pksherpuri@gmail.com, geetawatal@gmail.com

³ Department of Earth and Planetary Sciences, University of Allahabad, Prayagraj – 211 002, India; abhishekraigeology@gmail.com

⁴ Department of Physics, Chowdhary Mahadev Prasad Degree College, Prayagraj - 211 002, India; nileshkrai62@gmail.com

⁵ Physics and Astronomy Department, University of Tennessee, University of Tennessee Space Institute, Center for Laser Applications, 411 B.H. Goethert Parkway, Tullahoma, TN 37388-9700, USA; cparigge@tennessee.edu

⁶ Department of Chemistry, Swami Shraddhanand College, University of Delhi, Alipur, New Delhi - 110 036, India; sumanyadav@ss.du.ac.in

* Correspondence: cparigge@tennessee.edu; Tel.: +1 (931) 841 5690

Abstract: This interdisciplinary work communicates identification and quantification of elements responsible for the bioactive potency of leaves from pointed gourd, *trichosanthes dioica*, using laser induced breakdown spectroscopy (LIBS). Calibration-free LIBS determines the presence of various trace and major elements, their concentrations, and ratios in which they are present in the leaves. The presence of specific elemental ratios of magnesium/sodium and magnesium/potassium could be promising for managing diabetes mellitus. Variable doses of 500, 750, 1000, and 1250 mg per kg body weight of aqueous extract from *trichosanthes dioica* leaves are administered. A dose of 1000 mg per kg body weight is found to be the most effective. Based on encouraging results, the extract could be harvested to serve as anti-diabetic medication for diabetes and associated symptoms.

Keywords: optical emission spectroscopy; laser induced optical breakdown; *trichosanthes dioica*; pointed gourd; diabetes; biomedicine; biophysics

1. Introduction

Natural remedies offer alternative or complementary treatments for various diseases and are usually considered to be safe. For specific diseases that require lifelong pharmaceutical medication, therapeutic safety is important. For example, diabetes mellitus can be a lifelong disease and is globally of serious concern due to its complications. Several medicinal plants have been studied and scientifically proven to be beneficial for the treatment of diabetes mellitus [1–4]. Plant-based natural medicines are safe and cost-effective. In fact, in addition to bioactive phytochemicals, medicinal plants do have bioactive elements in appropriate ratios, and thus, are helpful in maintaining a balance of trace elements in disturbed metabolic processes diabetic mellitus [5]. However, medicinal plants need to be scientifically explored for assessment of efficacy and safety. In addition to phytochemical analysis, the detection of phyto-elemental composition and elemental ratios of medicinal plants is expected to play a pivotal role in evaluation of probable medicinal activity.

Recently, laser-induced breakdown spectroscopy (LIBS) has emerged as a promising analytical spectroscopy for elemental analysis of samples of interest [6]. Although other conventional analytical spectroscopic techniques such as inductively-coupled plasma mass

spectroscopy (ICP-MS), atomic absorption spectroscopy (AAS), inductively-coupled plasma Optical emission spectrometry (ICP-OES), and flame atomic absorption spectroscopy (FAAS) are available for elemental analysis, LIBS shows several advantages. Particularly, LIBS can be rapid, shows minimal environmental loads, requires minimal efforts in sample preparation, and provides in-situ real-time analysis of the sample in solid, liquid, or gas phases [7]. Furthermore, LIBS is capable of simultaneous multi elements detection and evaluation of their concentration and intensity ratios, which help for establishing a correlation of a particular bioactivity of any part of a plant with their specific elemental composition.

The current study addresses the generation of the bio-active profile of leaves of *trichosanthes dioica* from evaluation of elemental ratios with laser-plasma spectroscopy, and the interpretation of the results in view of the anti-diabetic potency for *in vivo* models. *Trichosanthes Dioica*, or *T. dioica*, is a vegetable of the *cucurbitaceae* family. It is widely grown throughout India and to a lesser extent in other parts of South Asia [8]. It is also known as 'Parwal' in Hindi, or colloquially, pointed gourd is called green potato. In addition to the fruit, it's leaves are also used as a vegetable. 100 g leaves contain 9.0 mg of magnesium, 2.6 mg of sodium, 83.0 mg of potassium, 1.1 mg of copper, and 17.0 mg of sulphur [9]. Pointed gourd leaves and delicate shoots of are applied in traditional medicinal therapy. The literature reveals that the effect of feeding shade-dried leaves to animals impacts their blood glucose level and lipid profile [10,11]. The plant seeds are commonly used to treat acid reflux symptoms and display antifungal and antibacterial activities [12]. The present study was conducted to evaluate the glycemic potency of aqueous extract of *T. dioica* leaves on blood glucose levels (BGL) of normal and streptozotocin-induced sub-, mild-, and severe- diabetic models. For a complete biomedical profile of the aqueous extract of *T. dioica* leaves, several other parameters were monitored, including TC, HDL, LDL, VLDL, TG, AST, ALT, ALKP, and CRTN. Based on the parameter ranges, and especially LFT (liver function test) and KFT (kidney function test), one can infer that *T. dioica* leaves should be developed as a promising anti-diabetic agent for management of diabetes type II and associated complications.

The present study includes the LIBS-based detection of the elemental composition of *trichosanthes dioica* leaves. A calibration-free LIBS, or CF-LIBS, algorithm was applied for the recorded LIBS spectra of *t. diocia* leaves to determine their elemental ratios of Na/K, Mg/Na, and Mg/K. The anti-diabetic behavior of *T. diocia* leaves has been correlated with these elemental ratios. The same intensity ratios of the elements Na, K and Mg were observed in LIBS spectra as for ayurvedic medicines prescribed for the treatment of diabetes mellitus. The anti-diabetic effect of *t. diocia* leaves extract was reconfrmed by conducting in-vivo studies on normal-, sub-, mild- and severe- diabetic models. Variable doses of the extract were given orally, and the impact on the BGL of selected animals is investigated in this work.

2. Materials and Methods

2.1. Experimental Arrangement for LIBS Instrumentation

Figure 1 illustrates The design of LIBS experiments. The Nd:YAG laser device (Continuum Surelite III-10, USA) is operated at a repetition rate of 10 Hz, delivering pulses of full-width-half-maximum 4 ns, and maximum laser energy up to 425 mJ at the second harmonic wavelength of 532 nm. The laser beam is focused onto the sample with a 30 cm focal length lens to achieve an irradiance of the order of 1 TW/cm². The light emitted from the plasma was collected by a small lens on the tip of the optical fiber that is connected to the entrance slit of a echelle-grating spectrometer equipped with an intensified charge-coupled device (ICCD). The lens at the tip of the fiber is positioned 1 cm distance from the laser-plasma. To avoid craters formed by an individual laser-sample interaction, the sample is mounted on a translational stage and moved so that a fresh sample surface is available for each laser-plasma generation.




Figure 1. Schematic diagram of the experimental setup for LIBS.

2.2. Sample Preparation

Fresh leaves of *t. dioica* in the amount of 5 kg were purchased at a local market in Allahabad, India, and were subsequently verified by Prof. Satya Narayan, Taxonomist, Department of Botany, University of Allahabad. A voucher specimen (Allahabad University 321) was also submitted. Leaves were washed, chopped, shade dried, and then crushed into powder. The Soxhlet technique was employed with distilled water for 36 hours for extraction of dried powder at 100 degree Celcius. The aqueous extract was filtered and concentrated in a rotatory evaporator at a temperature of 35 degree Celcius under reduced pressure to produce semi-solid material, which was subsequently lyophilized to obtain the powder with a 25.1 percent weight-to-weight yield.

2.3. Animals for the Study

Healthy male albino, Wistar rats at an age range of 6 to 8 weeks and at a weight in the range of 0.15 to 0.2 kg, served as the subjects in the experiments. Animals were obtained from the Luknow, India, central drug research institute. The rats were housed in standard ambient conditions, namely, 25 degree Celcius temperature, 50 % humidity, 12 hours of darkness and light per day, and with free access to water and standard laboratory diet composed of carbohydrates: 30%, proteins: 22%, lipids: 12%, and vitamins: 3%). The study was approved by the Allahabad University Institutional Ethical Committee, 83a/a/04/CPCSEA - Committee for the Purpose of Control and Supervision of Experiments on Animals.

2.4. Induction of Diabetes

In a group of overnight fasted rats, a single intra-peritoneal injection of freshly produced streptozotocin (bought from Sigma Aldrich Chemical Company, USA) at a dose of 55 mg per kg body-weight was given to induce diabetes [13]. The FBG level was checked after 3 days, and the PPG was continuously monitored up to stable hyperglycemia, which occurred a week later. Animals with significant hyperglycemia (FBG > 250 mg/dl) were selected for the study.

2.5. Blood Glucose Level Measurements

Standard kits from Bayer Diagnostics India, Limited, were used to estimate blood glucose level (BGL) based on the glucose oxidase method [14] and serum levels of TC (total cholesterol), TG (triglycerides), and HDL (high-density lipoprotein) spectrophotometrically in accordance with the manufacturer's instructions based on enzymatic studies [15,16]. However, Friedwald's formula, $VLDL = TG/5$, was used to determine the amount of VLDL (very low-density lipoprotein), and similarly, another Friedwald's formula, $LDL = TC - VLDL + HDL$, was used to determine the amount of LDL (low-density lipoprotein) [17].

Serum levels of LFT (liver function tests) [18] viz. AST(aspartate transferase), ALT(alanine transferase), ALKP (alkaline phosphatase)[19] and KFT (kidney function tests) viz. CRTN (creatinine) [20] and TPR (total protein) [21] were also estimated using standard kits from Bayer Diagnostics India Limited. Total Hb (total hemoglobin) was also measured before and after the therapy [22]. Reagent-based Uristix from Bayer Diagnostics was used to identify US (urine sugar). All parameters were assessed regularly throughout the long-term treatment.

2.6. Experimental Design

FBG and GTT experiments were performed with variable doses of aqueous extract of leaves in normal-, sub-, and mild- diabetic rats to assess hypoglycemic and anti-diabetic effects. The most effective dose identified was used for long-term treatment of severely diabetic rats to assess hypo-lipidemic-, hepato-protective-, and renal-protective- efficacies of the extract.

2.6.1. Assessment of Hypoglycemic Activity in Normal Healthy Rats

Five groups of six rats each were employed in the experiments. Animals in groups II, III, IV, and V received variable doses of aqueous extract viz. 500, 750, 1000, and 1250 mg/kg bw, respectively. Whereas, group I served as control group and received vehicle distilled water only. After administering the extract, blood samples were taken from the tail vein after 1.5, 3, 4.5, and 6 hours.

2.6.2. Assessment of Hypoglycemic Activity with the Glucose Tolerance Test in Normal Healthy Rats

A different group of healthy, normal animals received the aqueous extract orally in the same manner as above, and the effects on FBG were observed hourly for up to two hours. The BGL value at 2 h was treated as '0' h value for the Glucose Tolerance Test (GTT). The animals were then orally administrated with 4g/kg of glucose and their glucose tolerance was studied at 1 h interval for another 3 h. Thus, the total time for collecting the blood was up to 5 hours.

2.6.3. Assessment of Anti-Diabetic Activity with the Glucose Tolerance Test in Sub- and Mild- Diabetic Rats

The improvement of glucose tolerance in sub- and mildly diabetic rats was another way to measure the anti-diabetic impact of the aqueous extract of *t. dioica* leaves. The rats were divided in to six groups. Group I served as the control received vehicle (distilled water) only, whereas groups II, III, IV, and V received variable doses of the Leave extract—500, 750, 1000, and 1250 mg/kg bw, respectively. Blood glucose levels were checked firstly after 90 minutes of treatment considered as '0' h value and then 2 g/kg glucose was given orally to all the groups. Blood glucose levels were further checked upto three hours at regular intervals of 1 h each, considered as 1 h, 2 h, and 3 h values. The results were compared with the VI group of rats, which were treated, with 250 mg/kg of Tolbutamide (hypoglycemic agent).

2.6.4. Assessment of Anti-Diabetic Activity in Severely Diabetic Rats

For normal, subdiabetic and mild diabetic models, the most effective dose was determined to be 1000 mg/kg. A number of biochemical parameters related to antidiabetic attribute viz. FBG, PPG, Lipid profiles, Total Protein, Haemoglobin and Enzymatic assays were taken in to consideration for getting a complete biomedical profile of aqueous extract of *t. dioica* leaves. Two groups of 6 rats each were used in the experiment, group I served as control group, whereas group II was treated daily with a single oral administration of above mentioned dose. Body weight and Urine sugar were also estimated weekly along with other biochemical parameters up to 30 days.

2.6.5. Lethal Dose 50 Experiment

The lethal dose 50, LD₅₀, experiment was designed to assess the extract's toxicity. Two variable doses of 10 g and 15 g of the aqueous extract of *t. dioica* leaves were given once, orally to two different groups of rats of both sexes, weighing between 180 and 200 g. After which, rats were continuously monitored for gross behavioural, neurologic, autonomic, and toxic consequences. Food intake, face, and urine were also analyzed every two hours for the first six hours during a period of 24 hours.

2.7. Statistical Analysis

The one-way analysis of variance, ANOVA, was utilized in the statistically analysis the data. The Statistical Package for the Social Sciences, SPSS, version 7.5 assessed the *post hoc* Scheffe test. When the *P* values were less than 0.05, $P < 0.05$, results were deemed significant.

3. Results

3.1. LIBS Experiment

In the present experiment, quantitative LIBS analysis evaluated the elemental composition of the prepared samples of *t. dioica* leaves that were prepared as discussed in Section 2.2.

3.1.1. Determination of the Elemental Composition

For the evaluation of the elemental composition of *t. dioica* leaves, the recorded LIBS spectra illustrated in Figure 2 were analyzed. The wavelengths of spectral lines in the LIBS data were identified using the National Institute of Standards and Technology (NIST) atomic spectroscopic database [23]. The LIBS spectra reveal the presence of the spectral signature of a volley of elements in the *t. dioica* leaves. For analysis and discussion, the LIBS spectra of the sample are displayed in four different spectral regions. The spectra clearly reveal the presence of spectral elements including C, H, N, O, Na, K, Mg, Fe, Si, Sr, and Ca. Table 1 collates identified elements and their wavelengths.





Figure 2. LIBS spectra of *t. dioica* leaves for different spectral regions. (a) 230 to 300 nm, (b) 300 to 400 nm, (c) 400 to 500 nm, and (d) 550 to 900 nm.

Table 1. List of identified elements in *t. dioica* samples.

element	wavelength (nm)
carbon (C)	247.8
nitrogen (N)	744.2, 746.8, 818.4, 818.8, 821.6
oxygen (O)	777.3, 844.6
hydrogen (H)	656.3
calcium (Ca)	317.8, 318.2, 370.6, 373.7, 393.4, 396.3, 422.7, 430.2, 430.7, 442.5, 443.4, 443.5, 445.5, 445.6, 445.7, 487.8, 610.2, 612.2, 616.2, 643.9, 646.2, 649.3, 671.7, 714.8, 720.2
magnesium (Mg)	279.6, 280.2, 285.2, 383.0, 383.9,
sodium (Na)	588.9, 589.5
potassium (K)	404.5, 404.8, 766.4, 769.8
iron (Fe)	238.2, 239.5, 240.4, 252.3, 259.8, 263.1, 273.9, 274.9, 275.5, 278.8, 293.6

3.1.2. Semi-Quantitative Analysis

It is important to determine the concentration of trace elements present in any dietary supplement for the evaluation of the ratios of their consumable amounts. Knowing dietary supplement ratios is helpful for maintaining a healthy balance and proper regulation of physiological processes in the human body. Therefore, it is essential to evaluate the actual concentration and ratios of elements present in any medicine in order to correlate their specific bioactivity with specific elemental ratios. Kumari *et al.* evaluated the ratio of minerals helpful in diabetic management and concluded that in most ayurvedic anti-diabetic medicine the ratio of Mg/Na and Mg/K are approximately the same [24]. The present work also confirms, as shown in Figure 3, that the ratios of spectral intensity of Mg/Na and Mg/K are the same in *t. dioica* sample. The elements Ca, Fe, and Mg, are present in carbohydrate metabolism and also play a vital role in cellular uptake of the glucose molecule. A low Mg level could be risky for diabetes patients because of the Mg involvement in insulin activation. Potassium, K, activates beta cells of the pancreas and helps in the production of insulin in the human body. Thus, low potassium levels could be risky as well for developing diabetes due to persistent high blood glucose levels. Hence, the presence of elements like Mg, K, and Na in the sample could be correlated well with the observed significant efficacy of *t. dioica* leaves in insulin-dependent diabetes mellitus (IDDM) [24,25].

**Figure 3.** Intensity ratios Mg/Na, K/Na, and Mg/K of *t. dioica* leaves.

3.1.3. Quantitative Analysis

The importance of various bio-elements in the medicinal activity of plants cannot be ignored and thus it becomes imperative to detect their presence, concentrations, and ratios in which they are present in *T. dioica* leaves sample. Hence, the anti-diabetic activity of *T. dioica* leaves could be correlated not only with the presence of Na, K, and Mg but also with the assessment of specific ratios of Mg/Na and Mg/K, which are just the same

as in the earlier results of other anti-diabetic ayurvedic medicines [24]. For CF-LIBS one utilizes recorded intensities of different spectral lines of the elements, however, the laser-induced plasma needs to fulfill three conditions that must be verified: (i) plasma must be stoichiometric, (ii) plasma must be optically thin, and (iii) plasma must show local thermal equilibrium [26–28]. In the laser-plasma experiments, all three conditions are satisfied as further discussed below.

The calibration curve method is applied in a traditional approach for determination of concentrations of the constituents present in any materials. In this method, laser-induced breakdown spectra are recorded for each reference material with known element concentrations, viz. one measures the intensity of a spectral line present in each reference material. A graph is drawn for spectral intensities versus elemental concentrations in the reference materials. Calibration curves are established for each element of interest from reference material with known ingredient concentrations and the same matrix as the unknown target material. Usually it is challenging to find a reference material with the same matrix as the material in which concentrations of the constituents are to be determined. In turn, CF-LIBS method is well-established to determine each constituent's concentration simultaneously without requiring reference materials. The results obtained by CF-LIBS are reliable with a typical uncertainty of less than 10% [29].

Stoichiometric ablation

The ablation to be stoichiometric, the laser ablated part of the sample which is converted into the plasma must truly represent the composition of the sample. In this regard, it is said [26–28] that the laser irradiance greater than 1 GW/cm^2 leads to stoichiometric ablation irrespective of the state of matter and yields a plasma truly representative of the sample's elemental composition. Here, in our experiment, the pellet of the sample is homogeneous. Laser irradiance can be calculated by determining the laser spot size diameter, D , on the sample surface,

$$D = 4\lambda f / \pi d, \quad (1)$$

where λ is the wavelength of the laser beam, f is the focal length of the lens, and d is the beam diameter of the laser beam. In the present experiment, laser energy per pulse, $E = 15 \text{ mJ}$, $\lambda = 532 \text{ nm}$, $f = 30 \text{ cm}$, and $d = 0.9 \text{ cm}$. Therefore, the laser spot size diameter on the sample's surface amounts to $D = 22.6 \text{ }\mu\text{m}$. Consequently, the fluence and the irradiance are $3.8 \times 10^3 \text{ J/cm}^2$ and 0.94 TW/cm^2 , respectively. Clearly, in the present experiment, the laser irradiance is greater than 1 GW/cm^2 and fulfills the condition for stoichiometric ablation.

Optically Thin Plasma

For the validation of optically thin plasma, we have selected the pair of two interference-free lines of Ca II and Mg II having the same upper energy level and estimated the spectral intensity ratio followed by Lorentzian fitting. The theoretical intensity ratio has been estimated by using the other spectroscopic parameters in NIST data. Both the theoretical and Experimental ratios have been listed in table (2). We observed from the table that the experimental ratio (I_1/I_2) is approximately equal to the theoretical ratio $A_1g_1\lambda_2/A_2g_2\lambda_1$ for the pair spectral lines Ca-II (393.3/396.8) and MgII (279.5/280.2). It is clear from Table 2 that the intensity ratio (I_1/I_2) of two spectral lines is approximately equal to the ratio $A_1g_1\lambda_2/A_2g_2\lambda_1$. These results demonstrate that the laser-induced plasma in the present study is optically thin.

Table 2. Experimental and theoretical ratio of the spectral line Ca II and Mg II.

element	experimental ratio (I ₁ /I ₂)	theoretical ratio A ₁ g ₁ λ ₂ /A ₂ g ₂ λ ₁
Ca II (393.3/396.8)	1.92	1.86
Mg II (279.5/280.2)	1.98	2.05

Local Thermodynamic Equilibrium

251

The necessary McWhirter criterion,

$$N_e(\text{cm}^{-3}) > 1.6 \times 10^{12}(T(\text{K}))^{1/2}(\Delta E(\text{eV}))^3, \quad (2)$$

where N_e is the electron density, T the plasma temperature, and ΔE the energy difference for the corresponding transition, needs to be met for the laser-plasma to be considered in a state of local thermodynamic equilibrium (LTE). In this work, emission lines of Ca I and Mg I are selected for graphical display in form of a Boltzmann plot that utilizes emission intensity and the standard logarithm form,

$$I_\lambda^{ki} = C_s F A_{k_i} g_k \frac{e^{-E_k/(k_B T)}}{U_s(T)}, \quad (3)$$

$$\ln \left[\frac{I_\lambda^{k_i}}{A_{k_i} g_k} \right] = -\frac{E_k}{k_B T} + \ln \left[\frac{C_s}{U_s(T)} \right]. \quad (4)$$

Here, k_B is the Boltzmann constant, $U_s(T)$ the partition function, C_s the concentration, F an experimental factor. The plasma temperature of each constituent in laser-produced plasma was determined by using the slope and intercept of the above equation. The average temperature of the plasma is found to be 10,570 K, as displayed in Figure 4. The sufficient condition for plasma to be considered in LTE requires that the ionization and excitation temperatures determined by the Saha-Boltzmann equation and by the Boltzmann plot, respectively, must be in agreement to within 15 per cent. The Saha-Boltzmann equation [29],

$$\ln \left[\frac{I_{ij}^{II} A_{mn}^I g_m^I}{I_{ij}^I A_{ij}^{II} g_i^{II}} \right] = -\frac{E_{ion} - dE + E_i^{II} - E_m^I}{k_B T} + \ln \left[\frac{2(2\pi m_e k_B T)^{3/2}}{N_e h^3} \right], \quad (5)$$

for the ionic and atomic emission ratio serves as sufficient condition for confirmation of LTE. Here, dE is the lowering correction parameter, i.e., the correction term in the first ionization potential emerging due to high pressure in the plasma plume. m_e is the electron mass, h Planck's constant, E_{ion} the element's first ionization potential. E_i^{II} and E_m^I are the ionic and atomic species of an element's upper energy levels with transition probabilities of A_{ij}^{II} and A_{mn}^I and statistical weights g_i^{II} and g_m^I , respectively. Fig. 4 also illustrates the Saha-Boltzmann plot. The slope indicates an ionization temperature of 11,651 K. Excitation and ionization temperature are within 11 %, consequently, the laser-plasma is in LTE.

For determination of electron density, the Stark-broadened 422.6-nm Ca I line is well-isolated from other atomic lines, see Figure 4 (a). The electron density,

$$N_e = \frac{\Delta \lambda_{FWHM}}{2w_S}, \quad (6)$$

is proportional to the full-width-at half-maximum (FWHM), $\Delta \lambda_{FWHM}$, and inversely proportional to the Stark-broadening parameter, w_S . From the FWHM of the line profile, the average, line-of-sight electron number density amounts to $N_e = 2.2 \times 10^{18} \text{ cm}^{-3}$. The electron density is larger than the necessary McWhirter criterion of $1.6 \times 10^{15} \text{ cm}^{-3}$ for the 422.6-nm Ca I line and for an average, line-of-sight temperature of 10,570 K.




Figure 4. (a) Lorentzian plot for determination of electron density. (b) Boltzmann plot for determination of plasma excitation temperature. (c) Saha-Boltzmann plot for determination ionization temperature. Excitation and ionization temperature agree within 11 %, confirming local thermodynamic equilibrium, see text.

Application of CF-LIBS algorithm [30] leads to the following concentrations: 38.2% Ca, 2.6% Fe, 24.6% Na, 18.4% K, 6.3% Mg, and 0.24% Sr, 7% C, 2.8% N, 0.4% O, and 1.8% H. Figure 5 displays concentration ratios of Mg/Na, K/Na, and Mg/K that are calculated with CF-LIBS. These results indicate good agreement with the ratios of the integrated intensities of Mg/Na, K/Na, Mg/K of *t. dioica* leaves samples illustrated in Fig. 3. The experiments reveal that that spectral intensities of the elements can be used for evaluation of actual concentrations and concentration ratios of the elements.




Figure 5. Concentration ratios of Mg/Na, K/Na, and Mg/K, evaluated with CF-LIBS.

3.2. Biochemical Analysis

3.2.1. Effect on Normal Healthy Rats

Table 3 indicates the hypoglycemic impact of a single oral administration of variable doses of 500, 750, 1000, and 1250 mg/kg bw aqueous leave extract in normal healthy rats, including the control dose of 0, i.e., distilled water (dw) Treated rats showed a regular fall of 8.56, 20.36 and 23.87 % from the doses of 500, 750 and 1000 mg/kg bw, respectively, after 6 h. However, a fall of only 14.04% was observed with the dose of 1250 mg/kg bw after the same time interval.

Table 3. Effect of graded dose of *t. dioica* leave aqueous extract on blood glucose leveles (mg/dl) of normoglycemic rats (mean \pm standard deviation) for pre- and post- treatments.

experiment groups	treatment (mg/kg bw)	pre-treatment FBG	post (h) 1.5	post (h) 3.0	post (h) 4.5	post (h) 6.0
control	0 (dw)	71.5 \pm 3.2	69.3 \pm 3.2	73.0 \pm 3.6	71.5 \pm 3.3	71.2 \pm 4.0
extract	500	72.0 \pm 3.5	69.7 \pm 4.3	65.4 \pm 4.6	61.4 \pm 4.2	65.1 \pm 3.4 ^a
extract	750	69.9 \pm 3.9	64.5 \pm 4.2	63.1 \pm 4.4	57.9 \pm 4.2	56.7 \pm 3.9
extract	1000	70.3 \pm 3.7	67.1 \pm 4.1	64.8 \pm 4.3	57.8 \pm 4.7	54.6 \pm 3.5 ^a
extract	1250	71.1 \pm 4.6	70.4 \pm 4.2	67.1 \pm 4.8	65.7 \pm 3.6 ^a	61.2 \pm 4.9 ^a

^a $P < 0.01$ compared with control

3.2.2. Effect on Normal Healthy Rats during Glucose Tolerance Test

Table 4 summarizes the study results of aqueous extract of *t. dioica* leaves on blood glucose level (BGL) and glucose tolerance test (GTT) of normal healthy rats. Different doses of 500, 750, 1000 and 1250 mg/kg bw of extract were given orally to overnight fasted healthy rats. The observed decrease with the doses of 500, 750, 1000 and 1250 mg/kg bw in BGL after 2 h of administration was 12.28, 9.54, 15.21, and 12.15 %, respectively, and these BGL's are considered as the '0' h value. However, the decrease was observed up to 3 h after glucose administration, and at 1 h intervals. The results show that the percentage decrease in BGL's was regular up to the dose of 1000 mg/kg bw and reaches its maximum of 17.28 %. Moreover, a decrease of 14.77 % was observed with the dose of 1250 mg/kg bw at the same time.

Table 4. Effect of graded dose of *t. dioica* leaves aqueous extract on BGL during GTT of normoglycemic rats (mean \pm standard deviation) for pre- and post- treatments.

experiment groups	treatment (mg/kg bw)	pre-treatment FBG	post (h) 1	post (h) 2	post (h) 3	post (h) 4	post (h) 5
control	0 (dw)	73.3 \pm 4.1	75.9 \pm 3.2	76.5 \pm 4.5	107.3 \pm 4.1	99.6 \pm 3.7	91.4 \pm 4.5
treated 1	500	69.1 \pm 4.6	68.1 \pm 5.4	67.1 \pm 4.3	691.3 \pm 4.1	83.2 \pm 4.3 ^a	78.1 \pm 3.9
treated 2	750	66.9 \pm 5.5	67.8 \pm 4.2	69.2 \pm 3.7	591.4 \pm 4.4	81.4 \pm 4.4 ^a	75.1 \pm 3.9
treated 3	1000	72.5 \pm 4.1	65.4 \pm 4.1	64.8 \pm 5.1	592.1 \pm 4.4	81.3 \pm 4.4 ^a	75.6 \pm 4.1
treated 4	1250	69.9 \pm 4.6	69.1 \pm 3.9	67.2 \pm 3.8	690.1 \pm 4.3 ^a	684.1 \pm 3.8 ^a	77.9 \pm 3.5

^a $P < 0.01$ compared with control

3.2.3. Effect on diabetic rats during Glucose Tolerance Test

Figure 6 and Figure 7 demonstrate the antidiabetic effect of aqueous extract of *t. dioica* leaves on sub- and mild- diabetic animals, respectively. Different doses of aqueous extract as mentioned above along with the standard drug tolbutamide 250 mg/kg bw was given orally to the groups as defined in the experimental design. The decrease of 13.7, 17.01, 22.0 and 16.3 % in BGL's of sub-diabetic animals was observed after 3 h of glucose administration with the doses of 500, 750, 1000 and 1250 mg/kg bw, respectively. However, the dose of 250 mg/kg of tolbutamide reduced the BGL by 24.19 % at 3 h during GTT in sub-diabetic rats, which is comparable with the dose of 1000 mg/kg bw. Moreover, the maximum decrease of 22.0 % with 1000 mg/kg bw dose of aqueous extract and 24.19 % with 250 mg/kg bw doses of tolbutamide was observed after 3 h of glucose administration during GTT. The decrease of 15.7, 28.79, 31.39 and 30.86 % in BGL's of mild-diabetic animals was observed after 3 h of glucose administration with the doses of 500, 750, 1000 and 1250 mg/kg bw, respectively. However, the dose of 250 mg/kg of tolbutamide reduced BGL by 30.89 % after 3 h during GTT in mild-diabetic rats, which is almost same as with the dose of 1000 mg/kg bw. Moreover, the maximum decrease of 30.73 % with 1000 mg/kg bw dose of aqueous extract and 31.95 % with 250 mg/kg bw dose of tolbutamide was observed after 3 h of glucose administration during GTT.




Figure 6. Effect of grade doses of *t. dioica* leaves aqueous extract on BGL during GTT in sub-diabetic rats. Control: Distilled water, Treated 1: 500 mg/kg bw, Treated 2: 750 mg/kg bw, Treated 3: 1000 mg/kg bw, Treated 4: 1250 mg/kg bw, tolbutamide: 250 mg/kg bw.




Figure 7. Effect of grade doses of *t. dioica* leaves aqueous extract on BGL during GTT in mild-diabetic rats. Control: Distilled water, Treated 1: 500 mg/kg bw, Treated 2: 750 mg/kg bw, Treated 3: 1000 mg/kg bw, Treated 4: 1250 mg/kg bw, tolbutamide: 250 mg/kg bw.

3.2.4. Effect on Fasting Blood Glucose, Post Prandial Glucose, and Lipid Profile of Severely Diabetic Rats

The long-term treatment's antidiabetic and antilipemic effects on the blood sugar levels and lipid profiles of severely diabetic rats are shown in Table 5. Rats were treated with optimum effective dose of 1000 mg/kg bw of aqueous extract once a day at noon for one month. At the end of the treatment, the animals were compared with their own initial values and showed a significant reduction of 27.17 % in fasting blood glucose (FBG) and 31.17 % in post prandial glucose (PPG) levels. The enhanced levels of TG, TC, LDL, and VLDL cholesterol were brought down significantly to 59.21, 19.15, 9.13 and 56.92 %, respectively after the treatment period. The aqueous extract has gained much attention due to their promising improvement of 32.18 % in HDL level in severely diabetic treated group. Moreover, in untreated group there was a slight fall in HDL levels and slight increment in all the other above mentioned parameters.

3.2.5. Effect on Aspartate Aminotransferase, Alanine Aminotransferase, Alkaline Phosphatase, Creatinin, Total Protein, and Total Hemoglobin Levels of Severely Diabetic Rats

The hepatoprotective impact of long-term therapy is demonstrated in Table 6 on blood serum levels of aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALKP), and creatinine (CRTN). These indicators are usually measured in units per liter (u/l). The affected levels of total protein and hemoglobin are also recorded. In diabetic control rats there was a gradual increase in these all these parameters except total protein (TPR) and total haemoglobin (Hb) with vehicle (water only). Moreover, the most effective dose of 1000 mg/kg bw of the aqueous extract produced a fall of 26.19, 39.03, 38.43 and 41.17 %, respectively, in AST, ALT, ALKP and CRTN levels after 30 days treatment.

Table 5. Effect of graded dose of *t. dioica* leaves aqueous extract on blood glucose level and lipid profile of severely diabetic (SD) rats (mean \pm s.d.) for pre- and post- treatment levels.

experiment animals	treatment (aqueous extract)	pre-treatment levels	post (days) 7	post (days) 14	post (days) 21	post (days) 28
FBG (mg/dl)						
Normal (control)	0 (dw)	82.3 \pm 3.4	85.7 \pm 2.9	84.5 \pm 4.5	83.8 \pm 4.9	85.3 \pm 3.8
SD (control)	0 (dw)	287.4 \pm 5.6	290.8 \pm 3.5 ^a	302.5 \pm 4.7 ^a	299.8 \pm 4.8 ^a	298.8 \pm 4.8 ^a
SD (treated)	1000 mg/kg bw	297.3 \pm 4.6	273.5 \pm 4.5 ^a	255.6 \pm 4.4 ^a	223.9 \pm 4.8 ^a	211.2 \pm 4.9
PPG (mg/dl)						
Normal (control)	0 (dw)	161.6 \pm 5.1	161.9 \pm 4.8	162.1 \pm 4.8	161.4 \pm 5.2	162.6 \pm 4.2
SD (control)	0 (dw)	446.9 \pm 4.7	455.4 \pm 5.2 ^a	457.8 \pm 3.7 ^a	453.5 \pm 4.1 ^a	462.9 \pm 3.7 ^a
SD (treated)	1000 mg/kg bw	437.3 \pm 3.4	385.2 \pm 5.0 ^a	353.4 \pm 4.2 ^a	321.3 \pm 4.1 ^a	303.2 \pm 4.1 ^a
total cholesterol (mg/dl)						
Normal (control)	0 (dw)	94.5 \pm 4.1	96.0 \pm 3.9	97.9 \pm 3.7	97.2 \pm 3.1	97.0 \pm 3.3
SD (control)	0 (dw)	117.9 \pm 4.1	115.8 \pm 5.1 ^a	115.3 \pm 5.1 ^a	114.6 \pm 4.1 ^a	115.2 \pm 5.1 ^a
SD (treated)	1000 mg/kg bw	116.5 \pm 4.9	101.7 \pm 4.3 ^a	99.9 \pm 4.5 ^a	93.5 \pm 5.3 ^a	94.9 \pm 5.0 ^a
HDL cholesterol (mg/dl)						
Normal (control)	0 (dw)	27.6 \pm 4.1	28.9 \pm 4.6	29.7 \pm 4.5	28.8 \pm 3.8	29.9 \pm 4.1
SD (control)	0 (dw)	23.9 \pm 2.9	23.6 \pm 3.1 ^b	22.5 \pm 3.6 ^b	23.2 \pm 3.8 ^b	23.3 \pm 3.1 ^b
SD (treated)	1000 mg/kg bw	21.1 \pm 3.6	25.3 \pm 4.8 ^b	25.4 \pm 5.2 ^b	27.8 \pm 5.4 ^b	28.1 \pm 6.2 ^b
LDL cholesterol (mg/dl)						
Normal (control)	0 (dw)	53.1 \pm 3.6	53.3 \pm 3.7	53.8 \pm 4.0	55.6 \pm 4.1	54.7 \pm 3.6
SD (control)	0 (dw)	51.8 \pm 3.3	52.4 \pm 6.4 ^a	55.6 \pm 5.4 ^a	53.6 \pm 4.2 ^b	53.9 \pm 4.1 ^b
SD (treated)	1000 mg/kg bw	54.6 \pm 2.8	48.2 \pm 3.2 ^a	51.1 \pm 4.1 ^a	47.9 \pm 5.3 ^b	49.3 \pm 4.2 ^a
VLDL cholesterol (mg/dl)						
Normal (control)	0 (dw)	14.2 \pm 2.9	14.3 \pm 2.3	14.4 \pm 2.5	14.5 \pm 2.7	14.6 \pm 2.4
SD (control)	0 (dw)	41.7 \pm 5.2	39.4 \pm 4.5 ^a	36.2 \pm 4.2 ^a	38.3 \pm 4.8 ^a	38.9 \pm 4.8 ^a
SD (treated)	1000 mg/kg bw	40.6 \pm 3.3	32.3 \pm 4.3 ^a	20.4 \pm 5.3 ^a	17.4 \pm 4.3 ^a	17.5 \pm 4.5 ^a
Triglycerides (mg/dl)						
Normal (control)	0 (dw)	91.9 \pm 4.7	91.4 \pm 4.1	92.8 \pm 4.2	91.2 \pm 3.1	92.8 \pm 4.3
SD (control)	0 (dw)	198.3 \pm 4.1	196.7 \pm 4.8 ^a	184.2 \pm 4.4 ^a	190.5 \pm 3.6 ^a	199.7 \pm 3.1 ^b
SD (treated)	1000 mg/kg bw	204.9 \pm 4.3	160.5 \pm 4.8 ^a	100.9 \pm 4.1 ^b	88.5 \pm 4.6 ^a	87.4 \pm 4.8 ^a

^a $P < 0.001$ compared with control^b $P < 0.01$ compared with control

The unique observation indicated a promising improvement of 36.36 and 15.78 % in protein and haemoglobin levels, respectively.

Table 6. Effect of graded dose of *t. dioica* leaves aqueous extract on serum enzymes, total protein, and total hemoglobin of severely diabetic (SD) rats (mean \pm s.d.) for pre- and post- treatment levels.

experiment animals	treatment (aqueous extract)	pre-treatment levels	post (days) 7	post (days) 14	post (days) 21	post (days) 28
AST (u/l)						
Normal (control)	0 (dw)	21.1 \pm 3.4	21.6 \pm 2.9	21.4 \pm 3.4	21.6 \pm 4.1	21.8 \pm 3.5
SD (control)	0 (dw)	37.5 \pm 4.6	35.2 \pm 3.7 ^a	35.7 \pm 3.5 ^a	36.8 \pm 3.9 ^a	36.3 \pm 3.9 ^a
SD (treated)	1000 mg/kg bw	37.8 \pm 4.6	36.4 \pm 3.9 ^a	35.2 \pm 3.6 ^a	33.4 \pm 4.2 ^a	27.9 \pm 4.4 ^a
ALT (u/l)						
Normal (control)	0 (dw)	21.6 \pm 3.4	21.8 \pm 4.3	21.2 \pm 4.2	21.9 \pm 4.8	21.6 \pm 4.1
SD (control)	0 (dw)	37.3 \pm 3.1	36.5 \pm 3.2 ^a	38.1 \pm 3.6 ^a	37.3 \pm 4.2 ^a	37.8 \pm 3.1 ^a
SD (treated)	1000 mg/kg bw	39.2 \pm 4.6	38.3 \pm 4.3 ^a	34.1 \pm 4.2 ^a	29.7 \pm 4.6 ^a	23.9 \pm 3.7 ^a
Alkaline Phosphatase (u/l)						
Normal (control)	0 (dw)	91.9 \pm 5.1	90.3 \pm 4.7	91.2 \pm 4.4	91.9 \pm 3.9	91.9 \pm 3.4
SD (control)	0 (dw)	163.6 \pm 4.5	164.9 \pm 4.4 ^a	165.2 \pm 4.2 ^a	166.1 \pm 4.9 ^a	168.7 \pm 4.2 ^a
SD (treated)	1000 mg/kg bw	166.5 \pm 4.7	169.9 \pm 4.4 ^a	145.9 \pm 4.1 ^b	114.6 \pm 4.9 ^a	102.5 \pm 5.1 ^b
Serum Creatinine (u/l)						
Normal (control)	0 (dw)	0.9 \pm 0.1	0.8 \pm 0.3	0.8 \pm 0.5	0.9 \pm 0.6	0.8 \pm 0.3
SD (control)	0 (dw)	1.4 \pm 0.5	1.5 \pm 0.7 ^a	1.7 \pm 0.5 ^a	1.6 \pm 0.4 ^a	1.8 \pm 0.6 ^a
SD (treated)	1000 mg/kg bw	1.7 \pm 0.6	1.7 \pm 0.5 ^a	1.3 \pm 0.4 ^a	1.2 \pm 0.4 ^a	1.0 \pm 0.4 ^a
total haemoglobin (g/dl)						
Normal (control)	0 (dw)	13.9 \pm 2.4	13.6 \pm 2.4	14.1 \pm 2.4	14.0 \pm 2.3	14.2 \pm 2.3
SD (control)	0 (dw)	10.5 \pm 3.2	10.5 \pm 2.4 ^a	10.1 \pm 3.6 ^a	10.6 \pm 2.5 ^a	10.0 \pm 2.4 ^a
SD (treated)	1000 mg/kg bw	11.4 \pm 2.9	12.4 \pm 2.5 ^a	14.2 \pm 2.3 ^a	13.1 \pm 2.6 ^a	13.2 \pm 2.7 ^a
total protein (mg/dl)						
Normal (control)	0 (dw)	8.8 \pm 1.9	8.6 \pm 1.3	8.6 \pm 1.5	8.9 \pm 2.4	8.9 \pm 2.3
SD (control)	0 (dw)	5.4 \pm 2.5	5.2 \pm 2.6 ^a	5.4 \pm 1.8 ^a	5.6 \pm 2.5 ^a	5.5 \pm 1.4 ^a
SD (treated)	1000 mg/kg bw	5.5 \pm 2.3	5.7 \pm 2.3 ^a	6.5 \pm 2.1 ^a	7.2 \pm 1.8 ^a	7.5 \pm 2.3 ^a

^a $P < 0.001$ compared with control

^b $P < 0.01$ compared with control

3.2.6. Effect on Urine Sugar and Body Weight of Severely Diabetic Rats

The effects of the most effective dose of the aqueous extract, 1000 mg/kg bw, on the body weight and urine sugar levels of severely diabetic rats and the control group are shown in Table 7. Urine sugar results of 0 are usually indicated as nil, and strip-measurements showed +, ++, +++, or ++++ as noted in the table. The urine sugar levels decreased 75 % and the body weight increased by 16.7 % after treatment with the effective dose. In turn,

enhanced levels of urine sugar accompanied with weight loss were observed in the control group. 341
342

Table 7. Impact of the most effective dose of *t. dioica* aqueous extract on urine sugar and body weight in normal and severely diabetic (SD) rats (mean \pm s.d.) for pre- and post- treatment levels.

experiment animals	treatment extract)	pre-treatment levels	post (days) 7	post (days) 14	post (days) 21	post (days) 28
urine sugar						
Normal (control)	0 (dw)	nil	nil	nil	nil	nil
SD (control)	0 (dw)	+++	+++	++++	++++	++++
SD (treated)	1000 mg/kg bw	++++	++++	+++	++	+
body weight (g)						
Normal (control)	0 (dw)	210 \pm 10	210 \pm 10	210 \pm 10	200 \pm 10	210 \pm 10
SD (control)	0 (dw)	260 \pm 10	250 \pm 10	240 \pm 10	240 \pm 10	230 \pm 10
SD (treated)	1000 mg/kg bw	250 \pm 10	240 \pm 10	240 \pm 10	230 \pm 10	230 \pm 10

3.2.7. Lethal Dose 50 343

The experiment was conducted with normal healthy rats. The behavior of the treated rats appeared normal. No toxic effect was reported at doses up to 10 and 15 times of the effective dose of the aqueous extract as no mortality was observed in any of these groups. 344
345
346

4. Discussion 347

Application of LIBS delivers the elemental composition of *t. dioica* leaves. These leaves are instrumental for the lowering of an elevated BGL, and consequently, can be used for therapy and management of diabetes. Elements including Na, K, Ca, Mg and Fe, and elemental ratios plays a vital role in managing diabetes. Noteworthy is that the ratios of Mg/Na and Mg/K, determined by CF-LIBS in aqueous extract of *t. dioica* leaves, indicate similar trends as observed for other anti-diabetic medicines. The *t. dioica* plant is traditionally used as a vegetable in today's Indian food system and as well in the ayurvedic system of medicine [31]. 348
349
350
351
352
353
354
355

The communicated work demonstrates considerable hypoglycemic and antidiabetic efficacy of *t. dioica* leaves corroborated by short- and long- term studies. FBG studies on normal rats show the greatest hypoglycemic impact (23.8 percent) in less than 6 hours, see Table 3. Whereas, the GTT studies on normal rats demonstrate that the hypoglycemic impact was initiated within 1 hour and intensified later, see Tab. 4. This implies that within about one hour the active components of aqueous extract induces the hypoglycemic effect after reaching the target tissues through circulation. The data also confirms that a significant effect persists for the next three hours after the glucose administration. 356
357
358
359
360
361
362
363

The GTT results reveal an improvement in glucose tolerance of sub- and mild- diabetic animals reflecting a significant drop in high blood glucose levels. The results with a dose of 250 mg/kg bw of tolbutamide, a synthetic drug, are comparable with the dose of 1000 mg/kg bw of the aqueous extract. This would indicate a similar mechanisms of action. STZ induces type II diabetes in which pancreatic insulin-secreting cells get destroyed [32,33], consequently, the elevated blood glucose levels in the diabetic animals used in this study are consistent with type II diabetes. Hence, *t. dioica* aqueous extract, which supposedly acts as an insulin secretion enhancer by activating beta cells could be considered an important and useful treatment in the context of type II diabetes. 364
365
366
367
368
369
370
371
372

In cases of normal-, sub-, and mild- diabetic animals, the dose of 1000 mg/kg bw was found to be the most effective dose, therefore, this dose was used to treat the animals with severe diabetes for further studies. It has been noted frequently that hyperlipidemia is 373
374
375

always associated with hyperglycemia. As indicated in medical literature, hypertriglyceridemia is the most prevalent lipid abnormality [33]. Numerous studies have linked elevated blood triglyceride levels to a higher risk of stroke. If high levels of TC and LDL cholesterol persist, then LDL has a propensity to adhere to the blood vessel walls, stimulating atherosclerosis, which in turn causes heart attacks and strokes. Consequently, high levels of LDL cholesterol increases the danger of developing heart disease that may induce stroke [34]. There is currently a volley of data supporting the notion that HDL is a beneficial lipid, as high levels are linked to low rates of heart disease while low levels are linked to higher rates. After 30 days of treatment, the animals in the current study with severe diabetes had increased HDL cholesterol levels and considerably decreased levels of TC, LDL, VLDL, and TG.

Other biochemical parameters taken into consideration, for assessing the improvement in LFT and KFT of severely diabetic animals, include AST, ALT, ALKP, CRTN, and TPR. AST is located in the liver and it is released into the serum when the liver suffers from damage. Increased levels of AST in serum cause heart attacks and muscle disorders. Although ALT is not just found in the liver, liver damage causes it to be released into the bloodstream. As a result, it acts as a pretty accurate measure of liver health. Although ALKP is present in a variety of organs, its significant presence in the liver cannot be ignored. Hence, elevated levels of ALKP are indicative of liver disease, or bile tract obstruction. However, high levels of CRTN cause renal failure. The reduction of raised levels of AST, ALT, ALKP, and CRTN by the treatment of aqueous extract of *t. dioica* leaves suggests that it could be explored as an anti-hyperglycemic agent and as an hepato-protective and renal protective agent to manage diabetic complications as well. The additional advantageous results in this study include a reduction in urine sugar level, an increase in body weight and total protein level. It has considerable value for human subjects too due to its traditional use as an Indian vegetable and its high LD₅₀, showing a great margin of safety.

5. CONCLUSIONS

The present study reveals the fact that LIBS could be used as an advanced analytical tool to identify and quantify the elements and their ratios responsible for the medicinal profile of herbs that are part of the ayurvedic system of medicine. The elemental ratio of Mg/Na and Mg/K in the aqueous extract of *t. dioica* leaves are the same and also follow the same trend as reported in traditional medicine that is prescribed for treatment of diabetic patients. Based on the results of all parameters specially from LFT (Liver Function Test) and KFT (Kidney Function Test), it could be said that the aqueous extract of *t. dioica* leaves could be developed as an anti-diabetic drug for the management of diabetes and its complications. Further characterizations of active components in *t. dioica* leaves are of future interest.

Acknowledgments: One of the authors, CGP, acknowledges the support in part by the Center for Laser Applications at the University of Tennessee Space Institute.

Author Contributions: All authors contributed equally to this work. Material preparation, data collection and analysis of LIBS experiment were performed by Tejmani Kumar, Biochemical analysis were performed by Prasant Kumar Rai. The first draft of the manuscript was written by Tejmani Kumar. All authors edited and commented on previous versions of the manuscript. All authors read and approved the submitted manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Funding: This research received no external funding

Institutional Review Board Statement: The study was approved by the Institutional Ethical Committee (83a/a/04/CPCSEA).

Data Availability Statement: The datasets obtained and analyzed in the current study are available from the corresponding author on request.

Conflicts of Interest: The author declares no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Abbreviations

The following abbreviations are used in this manuscript:

LIBS	Laser Induced Breakdown Spectroscopy	428
ALKP	Alkaline Phosphate	429
ALT	Alanine Transferase	430
AST	Serum Separate Transferase	
BGL	Blood Glucose Level	
bw	body weight	
CRTN	Creatinine	
dw	distilled water	
FBG	Fasting Blood Glucose	
GTT	Glucose Tolerance Test	
h	hours	
Hb	Haemoglobin	
HDL	High Density Lipoprotein	431
kg	kilogram	
LDL	Low Density Lipoprotein	
mg	milligram	
PPG	Post Prandial Glucose	
SD	Severly Diabetic	
s.d.	standard deviation	
STZ	Streptozotocin	
<i>t. dioica</i>	<i>trichosanthes dioica</i>	
TC	Total Cholesterol	
TPR	Total Protein	
TG	Tri-Glyceride	
US	Urine Sugar	

References

- Kesari, A.N.; Gupta R.K.; Watal G. Hypoglycemic effects of *Murraya koenigii* on normal and alloxan diabetic rabbits. *Journal of Ethnopharmacology*, **2005**, *97*, 247–251. 432
- Gupta, R.K.; Kesari, A.N.; Watal. G.; Murthy, P.S.; Chandra, R.; Tandan, V. Hypoglycemic and anti-diabetic effect of aqueous extract of leaves of *Annona squamosa* (L). in experimental animals. *Current Science*, **2005**, *88*, 1244–1254. 433
- Grover, J.K.; Yadav S.; Vats V. Medicinal plants of India with antidiabetic potential. *Journal of Ethnopharmacology*, **2002**, *81*, 81–100. 434
- Irrova, M.D.; Paya, M.; Villar, A. A review of natural products and plants as potential antidiabetic drugs. *Journal of Ethnopharmacology*, **1989** *27*, 243–275. 435
- Rai, P.K.; Jaiswal, D.; Mehta, S.; Watal, G. Anti-hyperglycaemic potential of *Psidium guajava* raw fruit peel. *Indian Journal of Medical Research*, **2009**, *129*(5), 561. 436
- Pathak, A.K.; Kumar, R.; Singh, V.K.; Agrawal, R.; Rai, S.; Rai, A.K. Assessment of LIBS for Spectrochemical Analysis: A Review, *Applied Spectroscopy Reviews*, **2012**, *47*, 14–40. 437
- Trevizan, L.C., Santos, D.Jr.; Samad, R.E.; Vieira, N.D.Jr.; Nomura, C.S.; Nunes, L.C.; Rufini, I.A.; Krug, F.J. Evaluation of laser induced breakdown spectroscopy for the determination of macronutrients in plant materials. *Spectrochimica Acta Part B: Atomic Spectroscopy*, **2008**, *63*, 1151–1158. 438
- Chakravarthy, H.M. (1982) Fascicles of flora of India-11 Cucurbitaceae. *Botanical survey of India*, **1982**, p. 136. 439
- Singh, K. Pointed ground. *Indian Horticulture*, **1989** *33* & *34*, 33–37. 440
- Sharma, G.; Pant, M.C. Effect of feeding *T. dioica* (Parval) whole Leaves on blood glucose, serum triglycerides, phospholipid, cholesterol and high-density lipoprotein-cholesterol levels in the normal albino rabbits. *Current science*, **1988**, *57*, 1085–1087. 441
- Mukharjee, S.K. International Seminar on free radicals mediated disease. Scope and use of indigenous disease. Scope and use of indigenous herbal drugs in Madhumeha. 2-4 September 1996, Varansi, India. *Indian scenario*, Abstract-11. 442
- Harit, M.; Rathee, P.S. Antifungal activity of unsaponifiable fraction of the fixed oil of (*Trichosanthes*) seed. *Asian journal of Chemistry*, **1996**, *8*, 180–182. 443
- Singh, S.K.; Rai, P.K; Jaiswal, D; Watal, G. Evidence-based Critical Evaluation of Glycemic Potential of *Cynodon dactylon*. *Evid. Based Complement Alternat. Med.* **2008**, *5*, 415–420. 444

14. Brahm, D.; Trinder, P. Estimation of glucose by glucose oxidase method. *Analyst*, **1972**, *97*, 142–145. 457
15. Allian, C.C.; Poon, L.S.; Chan, C.S.; Richmond, W.; Fu, P.C. Enzymatic determination of total cholesterol. *Clinical Chemistry*, **1974**, *20*, 470–475. 458
16. Buccolo, G.; David, M. Quantitative determination of Serum Triglycerides by use of enzyme. *Clinical Chemistry*, **1973**, *19*, 476–482. 460
17. Friedwald, W.T.; Ley, R.I.; Fradrickson, D.S. Estimation of low-density lipoprotein cholesterol in plasma without the use of preparative ultracentrifuge. *Clinical Chemistry*, **1972**, *18*, 499–502. 461
18. Henry, R.J.; Canon, D.C.; Winkelman, J.W. *Clinical Chemistry Principles and Techniques 2nd ed.*; Harper and Row: New York, NY, USA, 1974. 463
19. Hørder, M.; Elser, R.C.; Gerhardt, W.; Mathieu, M.; Sampson, E.J. International Federation of Clinical Chemistry (IFCC): Scientific Division, Committee on Enzymes. IFCC methods for the measurement of catalytic concentration of enzymes. Part 7. IFCC method for creatine kinase (ATP: creatine (N-phosphotransferase, EC 2.7.3.2). IFCC Recommendation, *Journal of Analytical Methods in Chemistry*, **1990**, *12*, 628560, 19 pp., <https://doi.org/10.1155/S1463924690000049>. 464
20. Meites, S. and American Association of Clinical Chemists. *Standard methods of Clinical Chemistry*; Academic Press: New York, NY, USA, 1965. 466
21. Strickland, R.D.; Freeman, M.L.; Gurule, F.F. Copper binding by proteins in Alkaline solution. *Analytical Chemistry*, **1961**, *33*, 125–128. 470
22. Nonfon, M.; Lieb, F.; Moeschler, H.; Wendish, D. Four anonins from *Annona squamosa*. *Phytochemistry*, **1990**, *29*, 1951–1954. 472
23. NIST: National Institute of Standards and Technology USA, electronic database <http://physics.nist.gov/PhysRefData/ASD/linesform.html> (accessed on 28 April 2022). 473
24. Kumari, R.; Kumar, R.; Rai, A.; Rai, A.K. Evaluation of Na and K in anti-diabetic ayurvedic medicine using LIBS. *Lasers Med. Sci.*, **2021**, *37*, 513–522. 474
25. Singh, V.K.; Rai, A.K.; Rai, P.K.; Jindal, P.K. Cross-sectional study of kidney stones by laser-induced breakdown spectroscopy. *Lasers Med. Sci.*, **2009**, *24*, 749–759. 476
26. Cremers, D.A.; Radziemski, L.J. *Handbook of laser induced breakdown spectroscopy*; Wiley: New York, NY, USA, 2006. 477
27. Griem, H.R. Validity of local thermal equilibrium in plasma spectroscopy. *Physical Review*, **1963**, *131*, 1170–1176. 478
28. Griem, H.R. *Plasma spectroscopy*; McGraw-Hill: New York, NY, USA, 1964. 479
29. Kumar, T.; Rai, A.K.; Dwivedi, A.; Kumar, R.; Azam, M.; Singh, V.; Yadav, N.; Rai, A.K. Chemical Characterization for the Detection of Impurities in Tainted and Natural *Curcuma longa* from India Using LIBS Coupled with PCA. *Atoms*, **2022**, *10*, 91. <https://doi.org/10.3390/atoms10030091>. 480
30. Kumar, R.; Rai, A.K.; Alamelu, D.; Aggarwal, S.K. Monitoring of toxic elements present in sludge of industrial waste using CF-LIBS. *Environ. Monit. Assess.*, **2013**, *185*, 171–180, <https://doi.org/10.1007/s10661-012-2541-0>. 481
31. Brahmavarchas. *Ayurveda ka Pran: Vanaushidhi Vigyan. 3rd ed*; Brahmavarchas Shodh Sansthan: Haridwar, India, 2003, pp. 450–451. 482
32. Hoftiezer, V. Comparison of streptozotocin-induced diabetes in the rat inducing volumetric quantitation of the pancreatic islets. *Diabetologia*, **1973**, *9*, 178–184. 483
33. Sharma, S.R.; Dwivedi, S.K.; Swarup, D. Hypoglycemic and hypolipidemic effects of *Cinnamomum tamala* Nees leaves. *Indian Journal of Experimental Biology*, **1996**, *34*, 372–374. 484
34. Sachdeva, A.; Khemani, L.D. Effect of *Hibiscus rosa sinesis* Linn. ethanol flower extract on blood glucose level and lipid profile in streptozotocin induced diabetic rats. *Journal of Ethnopharmacology*, **2003**, *89*, 61–66. 485