

## Review

# Potential Inflammatory Biomarker in Patients with Attention Deficit Hyperactivity Disorder

<sup>1,\*</sup>Ji Hyun Park<sup>1</sup> College of Pharmacy, Duksung Women's University, Seoul, 01369, Republic of Korea

\* Correspondence: pharmerpark@duksung.ac.kr

**Abstract:** Attention deficit hyperactivity disorder (ADHD) is a prevalent neurodevelopmental disorder that can diminish the quality of life of both children and adults in academic, occupational, and social contexts. The kynurenine pathway (KP) contains a set of enzymatic reactions involved in tryptophan (TRP) degradation. It is known to be associated with the risk of developing ADHD. This review will address the KP and underlying mechanism of inflammation in ADHD. Potential inflammatory biomarkers reported in the most recent studies are summarized. Although a strong neuro-immunological basis has been established due to advances of recent neurobiological research, the pathophysiology of ADHD remains unclear.

**Keywords:** kynurenine pathway; inflammatory biomarkers; inflammation; ADHD; neurobiology; kynurenine


## 1. Introduction

Attention deficit hyperactivity disorder (ADHD), one of the most commonly found neurodevelopmental disorders in children and adolescents, is characterized by impairing symptoms of inattention, hyperactivity, and impulsivity [1]. ADHD affects 3-10% of school-aged children, with reported prevalence rate of 5.9 to 8.6% in Korean population [2]. Its prevalence varies among studies. In a study in Taiwan, a local prevalence was estimated to be 7.5% [3]. In a Norwegian study of children aged 8 to 10 years, its prevalence was 1.7% [4]. A meta-analysis reported that the worldwide prevalence of ADHD in children and adolescents was 5.3% (95% CI: 5.01-5.56) in 2007 [5]. It has been shown that 60-85% of ADHD cases diagnosed in children persist in adulthood [6]. Consistently, a meta-analysis has reported that the prevalence of ADHD in adults is 2.4% [7].

ADHD has been associated with distinctively worse school performance independent from socioeconomic factors among children [8]. It is also associated with significant impairment in occupational, academic, and social functioning in adults [9]. According to DSM-5 criteria, the diagnosis of ADHD in children is based on the presence of at least six out of nine symptoms in two areas of inattention and hyperactivity-impulsivity with observed behavioral issues [10, 11]. The current diagnostic criteria of ADHD rely on subjective reporting from patients or other informants (patients, teachers) and clinical observations, which might not be able to differentiate definite illness from normal variation [12]. Meanwhile, public awareness and widespread recognition of this disorder have led to an obvious increase in the diagnosis and treatment rate of adult ADHD over the last decade [13]. Overdiagnosis and misdiagnosis are also concerned as they might result in unnecessary labeling, extra costs for excessive tests, unneeded therapies, and increased healthcare cost [14].

The precise mechanisms underlying ADHD have not been clearly established yet. However, it has been shown that reduced volume or functionality of the brain's gray and white matter might lead to deficits of attention, cognition, processing response speed, motor planning, and other behavioral problems shown in ADHD [15, 16]. Recent research

studies have proposed that cerebellum, caudate, and prefrontal cortex (PFC) as primary areas related to deficits in ADHD are interconnected together as a neuronal network for regulating attention, thoughts, behavior, actions, and emotions [17, 18]. The network activity between these areas is mediated by neurotransmitters (NTs), norepinephrine (NE), and dopamine (DA) via multiple receptors in presynaptic or postsynaptic neurons [17, 19-22]. Figure 1 demonstrates the integration of hyper- and hypo-active catecholamine hypothesis of ADHD [23].



**Figure 1. Integration of Catecholamine Hypothesis of ADHD.** ① Dopamine (yellow circles) acts on postsynaptic D<sub>1</sub>-D<sub>5</sub> receptors. D<sub>2</sub> and D<sub>3</sub> receptors are locally positioned on the presynaptic neuron. Small amounts of dopamine stored in vesicles are released into the synapse from the presynaptic terminal if there is no action potential. ② This composes a tonic pool that acts on D<sub>2</sub> and D<sub>3</sub> presynaptic receptors, providing a feedback to inhibit the release of DA. ③ Action potential is arrived at the presynaptic terminal. ④ A large amount of vesicular DA is released into the synapse. This composes the phasic pool acting on postsynaptic receptors. The amount of DA released in the phasic pool depends on feedback inhibition, which is fired by the stimulation of D<sub>2</sub> and D<sub>3</sub> receptors of the tonic pool. ⑤ DAT-1 on the presynaptic terminal reuptakes DA and the action of DA on the postsynaptic receptor is terminated. Similar action of NE and NE receptors ( $\alpha$  and  $\beta$ ) is shown in the PFC. It is postulated that when an individual is fatigued or bored, activation of postsynaptic D<sub>1</sub> and  $\alpha$ 2A receptors is insufficient due to the lack of DA or NE release. This mechanism leads to an individual being easily disturbed and impulsive. If too much of those NTs are released under stressful conditions, it can lead to overstimulation of receptors, resulting in inattention and responses. Focused, organized, and attentive actions and thoughts are produced after a modest stimulation of postsynaptic receptors of DA and NE. In ADHD individuals, tonic pool is thought to be reduced, which can result in more phasic release of DA, causing behavioral issues including inattention, hyperactivity and impulsivity. By antagonizing DAT-1 and NET, stimulants can boost the tonic pool and lessen the phasic release fired by action potential, which might be the underlying issue in ADHD. AP, action potential; DA, dopamine; NE, norepinephrine; PFC, prefrontal cortex, DAT-1, dopamine transporter-1; NET, norepinephrine transporter; D<sub>n</sub> receptor, Dopamine-n receptor (n=number); PFC, prefrontal cortex.

Several studies have measured plasma levels of monoamines in an attempt to explicate underlying biological mechanisms involved in ADHD [24]. However, the results are inconclusive, so further investigation is required. Another possible way to explain the underlying pathophysiology of ADHD is alteration of the tryptophan metabolic pathway

(TMP) [25, 26]. Tryptophan (TRP) is the most prevalent amino acid with a significant role in biosynthesis of proteins in humans and animals. The metabolic fate of TRP is bifurcated into two pathways: the kynurenine pathway (KP) and the serotonin pathway (SP) [27, 28]. Several studies using animal/human models have shown the role of tryptophan (TRP) metabolism in ADHD by demonstrating levels of TRP metabolites such as kynurenine (KYN), kynurenic acid (KYNA), and 3-hydroxykynurenin (3-HK) [29]. TRP and its metabolites play an important role in mitigating diverse disorders ranging from cancer to psychiatric or neurological diseases [30-32]. Recent studies have demonstrated that KP can affect diverse biological systems. Therefore, the potential of using metabolites in KP as promising therapeutic biomarkers has attracted great interest of biomedical researchers [33]. Furthermore, mechanisms linking KP to ADHD provide an excellent opportunity to establish new treatment for neuropsychiatric disorders [33, 34].


Currently, there has been no clinically reliable biomarker for the diagnosis of ADHD, although several plausible inflammatory biomarker candidates have been suggested based on recent advances in biochemical and molecular biology [12]. Determining stable and reliable biomarkers to separate definite ADHD from normal behaviors that are not etiologically ADHD is highly desired to make less misdiagnosis.

From this point of view, the aim of the present review article was to summarize potential inflammatory biomarkers and the underlying mechanisms, including detailed description of the kynurenine pathway, inflammatory-cytokine-mediated regulation of kynurenine metabolism, dysregulation of kynurenine pathway in ADHD, and potential biomarkers in ADHD.

## 2. The Kynurenine Pathway

The KP is involved in catabolic TRP degradation. It acts on glutamate receptors in the central nerve system and potentially regulates essential mechanisms of ADHD by generating a number of neuro-active compounds collectively called kynurenines that could interact with neurotransmitter (NT) receptors in the central nerve system (CNS) [35]. This pathway takes place in the liver, kidney, and brain of mammals such as human [36]. Although the liver and kidney show the highest concentrations of enzymes, all primary enzymes are also found in the brain [37, 38]. Kynurenine metabolism happens all over brain cells despite various annexes of the pathway are segregated into specific cell types [39].

The KP pathway is initiated either by tryptophan-2,3-dioxygenase (TDO) in the liver to open the TRP indole ring, or indole-2,3-dioxygenase (IDO) in the brain to produce an instable metabolite of *N*-formylkynurenine [40, 41] (**Figure 2**). Conversion of *N*-formylkynurenine to L-kynurenine (KYN) is then followed. KYN is a substrate of various enzymes, including kynureninase (KYNU) for catalyzing the production of anthranilic acid (AA), kynurenine aminotransferases (KATs I-IV) for kynurenine acid (or kynurenic acid, KYNA), and kynurenic 3-monooxygenase (KMO) for 3-hydroxykynurenine (3-HK). From L-KYN, the pathway bifurcates into two distinct branches often called “neuroprotective” and “neurotoxic” arms. The neuroprotective arm is modulated by KAT, whereas the neurotoxic part is modulated by KMO.



**Figure 2.** Schematic connection between the kynurenine pathway and inflammation. The kynurenine pathway (KP) bifurcates into two distinct branches modulated by the availability of L-kynurenine in the brain: KATs, and KMO. In addition, a variety of inflammation-related mediators known to affect enzyme expression can regulate the metabolism of kynurenine by adjusting substrate availability and metabolite formation, preferring the KMP direction of the pathway with immune-related pathological conditions. TRP, tryptophan or L-tryptophan; TDO, TRP 2,3-dioxygenase; 5-HT, serotonin; KYN, kynurenine or L-kynurenine; 5-HTP, 5-hydroxytryptophan; HPA-axis, hypothalamic-pituitary-adrenal axis; KYNA, kynurenine acid; 3-HK, 3-hydroxykynurenine; AA, anthranilic acid; XA, xanthurenic acid; NAD<sup>+</sup>, nicotinamide adenine dinucleotide; PIC, picolinic acid; 3-HAA, 3-hydroxyanthranilic acid; ACMSD, aminocarboxymuconate semialdehyde decarboxylase; QUIN, quinolinic acid; NMDA-R, N-methyl-D-aspartate receptor; IDO, indoleamine-2,3-dioxygenase; KAT, kynurenine aminotransferase; KMO, kynurenine 3-monooxygenase; KYNU, kynureninase; HAAO, 3-hydroxyanthranilic acid oxidase; LPS, lipopolysaccharide; BCG, bacillus Calmette-Guerin; IFNs, interferons; TNF, tumor necrosis factor; IL, interleukin; TNF- $\alpha$ , tumor necrosis factor alpha, IFN- $\gamma$ , interferon gamma; COX2, cyclooxygenase 2.

Brain kynurenine metabolism occurs mostly in glial cells. KMO, KYNU, and 3-hydroxyanthranilic acid oxidase (3-HAAO) can regulate the formation of L-KYN in microglia, resulting in formation of AA, 3-HK, 3-HAA, and quinolinic acid (QUIN). QUIN is excitotoxic at NMDA glutamate receptors. It has a synergistic effect with 3-HK in generating oxidative stress [42]. In astrocytes, L-KYN can be metabolized by KATs alternately. KAT II is the predominant subtype in brains of humans and rats [43]. KATs can catalyze L-KYN to KYNA, a glutamate neurotransmission inhibitor and a possible antagonist at nicotinic  $\alpha 7$  receptors. In conjunction with these roles, KYNA can interact with arylhydrocarbon receptors and GPR35 [19, 44]. In such a way, kynurenine-derive neuro-active compounds have multiple receptor targets. Further research is needed to clearly demonstrate their endogenous function. A third possible degradation modulated by both KATs and KMO is xanthurenic acid (XA). Although not much is known about XA, it might play a

role in modulating glutamatergic neurotransmission by activating Group II metabotropic glutamate receptors (mGlu2 and mGlu3) or inhibiting vesicular glutamate transporters, indicating that it could also modulate glutamate neurotransmission by impacting presynaptic release [45, 46].

During the last decades, the underlying regulatory mechanism of kynurenine metabolism has been extensively investigated as it has a role in CNS disorders [47-49]. As “neuroprotective” and “neurotoxic” branches of the pathway, KATs and KMO, respectively, can balance the production of KYNA:QUIN which is essential in psychiatric and neuropsychiatric diseases. Some kynurenine metabolites can pass through the blood brain barrier, indicating that CNS levels of kynurenines are mostly regulated by peripheral enzyme activities [50]. However, a large neutral amino acid transporter can actively convey kynurenine into the brain [51]. Most kynurenine which is metabolized into KYNA and QUIN is from the periphery under normal physiological conditions [52]. After systemic inflammation, nearly all kynurenine in the CNS is derived from the periphery, at which IDO expression is greatly increased [53]. However, direct induction of CNS inflammation can result in over 98% of kynurenine from local synthesis in the brain [52].

To the best of our knowledge, only three research groups have investigated serum levels of kynurenines in patients with ADHD. Evangelisti et al. (2017) have reported the most recent measurements [10, 25, 54], showing that serum concentrations of kynurenic acid (KA), xanthurenic acid, and anthranilic acid are lower, while tryptophan and kynurenine concentrations are significantly higher in children with ADHD than in healthy controls. They found that the existence of ADHD was significantly related to low AA levels and high TRP levels in a human logistic regression model [10]. The following section summarizes the interaction between inflammatory mediators and their underlying mechanisms regulating the KP.

### 3. Inflammatory Cytokine-Mediated Regulation of Kynurenine Metabolism

#### 2.1. Indoleamine 2,3-dioxygenase (IDO) and inflammation mediators

The first step of TRP catabolism takes place with IDO and TDO generally known to be differently modulated. While IDO is induced by pro-inflammatory cytokines during immune response, TDO is induced by glucagon and corticosteroids [55]. There are some evidences showing that other enzymes in neurotoxic branches of the KP can be also induced by pro-inflammatory cytokines. However, IDO regulation by interferon (IFN)- $\gamma$  has been investigated most widely.

While IFN- $\gamma$  is regarded as the primary IDO inducer, there is evidence showing that the expression of IDO can be induced independently of IFN- $\gamma$  [56-58]. *In vitro* data using HTP-a cells, a human monocytic cell line, have shown that LPS-induced IPO activation is mediated by an IFN- $\gamma$ -independent mechanism, including synergistic effects of TNF- $\beta$ , IL-6, and IL- $\beta$  1 [57]. Connor et al. [56] have also suggested that IFN- $\gamma$  might not be necessary for LPS-induced IDO expression in mixed glia cultures harvested from neonatal rat cortex. In human progenitor cells of hippocampus, IL- $\beta$  treatment does not upregulate TDO, but greatly increases the expression of IDO transcript. This supports that IL- $\beta$  can elevate functional levels of IDO enzyme [58].

Experiments investigating the role of anti-inflammatory cytokines in IDO expression have shown limited and often conflicting results. This might be due to differences in models used and experimental conditions applied. For instance, IL-10 as one of the major anti-inflammatory cytokines decreased LPS-mediated IDO protein expression in a dose-dependent manner. However, IFN- $\gamma$ -mediated IDO protein expression was increased by IL-10 in mouse bone marrow-derived dendritic cells (BMDCs) [59]. This inconsistency may propose that anti-inflammatory cytokines such as IL-10 can differentially regulate distinct mechanisms of IDO induction. However, it has not been demonstrated whether this occurs in the brain. Notably, IFN- $\gamma$ -treated IDO expression in transformed mouse neuronal cell line was suppressed by IL-10 [60]. In addition to the case of IL-10, studies on



human monocytes and fibroblasts have suggested that IL-4 can inhibit IDO mRNA induction and IDO activity by IFN- $\gamma$ . Opposed to this, a study using mouse microglia cells has reported that IL-4 can enhance IFN- $\gamma$  induced IDO mRNA expression, which is diminished by IL-4 antiserum addition [61]. Along with IL-4, IL-13 which utilizes the same receptor subunit in signaling can potentiate IFN- $\gamma$  treated IDO mRNA expression in mouse microglia cultures [61]. Collectively, these findings indicate that responses to anti-inflammatory cytokines in microglia and peripheral myeloid cells are different.

Other than those anti-inflammatory cytokines, pro-inflammatory cytokines (such as TNF- $\alpha$  and IL-1 $\beta$ ) and toll-like receptor (TLR) agonist (such as LPS) can synergistically potentiate IFN- $\gamma$ -treated IDO expression [62]. Moreover, TNF- $\alpha$  can synergistically induce IDO expression with IFN- $\gamma$  by increasing NF- $\kappa$ B-dependent IRF-1 expression and STAT-nym1 activation [62]. Synergistic IDO induction by TNF- $\alpha$  and IFN- $\gamma$  occurs in primary mouse microglial cells. The mechanism is utilized in a research model of inflammation-related depression [53].

IFN- $\gamma$ -independent IDO induction is supported by studies using primary mouse microglial cells demonstrating that IFN- $\gamma$  mRNA is not detectable whereas IDO mRNA levels are increased after LPS stimulation [19, 56]. Other studies using THP-1 cells have indicated that LPS-boosted L-KYN production does not occur with IRF-1 or STAT-1 binding activation, but is lessened by p38 and NF- $\kappa$ B inhibitors [57]. To sum up, IDO induction stimulated by LPS in monocytes is IFN- $\gamma$ -independent and related to NF- $\kappa$ B as well as stress-activated mitogen-activated protein kinases (MAPK) including JNK and p38 [19, 56, 57]. Although subsequent mechanisms between JNK or p38 and IDO induction in response to LPS stimulation have not been clearly established, AP-1 factors are conventional substrates of MAPKs. They have critical roles as inflammation-related gene transcription regulators [19].

## 2.2. Kynurenine-3-monooxygenase (KMO) and inflammation mediators

Similar to IDO, pro-inflammatory stimuli may activate KMO enzymes downstream of the pathway. After systemic inoculation of LPS, KMO expression is induced in rat brain [56]. KMO is also induced in both IFN- $\gamma$  treated immortalized murine microglia (N11) and macrophage (MTs) cells. However, KYNU is induced only in MT2 whereas 3-HAAO is not affected [63]. In human progenitor cells of hippocampus, transcriptional levels of KMO and KYMU are upregulated following IL-1 $\beta$  [58].

## 2.3. Kynurenine aminotransferases (KATs) and inflammation mediators

Compared to expression levels of IDO and other kynurenine enzymes in the neurotoxic branch of the KP, KAT expression is neither elevated nor changed in response to pro-inflammatory stimuli. Systematic LPS inoculation of LPS causes no change of KAT II in rat brain cells [56]. In immortalized murine microglia (N11) and macrophage (MTs) cells, KAT shows constitutive expression. IFN- $\gamma$  treatment shows no effect on KAT activity [63]. In human progenitor cells of hippocampus, IL-1 $\beta$  treatment downregulates only KAT I and III, showing no effect on KAT II [58].

## 4. Genetic Links Between Inflammation and Kynurenine Metabolism in ADHD

Genetic studies have supported that gene polymorphisms are linked to the inflammatory pathway in ADHD. In a total of 398 subjects, Smith et al. [30] have evaluated a set of 164 single-nucleotide polymorphisms (SNPs) from 31 candidate genes and found that two SNPs in the ciliary neurotrophic factor receptor (*CNTFR*) are associated with the severity of ADHD inattentive symptom. Odell et al. [64] have conducted a population-based association study with 546 ADHD patients vs. 546 controls and proposed an association between *CNTFR* and ADHD in both children and adults. They also reported an associa-

tion of ADHD and major histocompatibility complex genes, demonstrating the role of inflammation and autoimmunity in this disorder. However, recent findings of genome-wide association meta-analysis have failed to replicate these results [65].

Another genome-wide association study for 478 ADHD patients and 880 controls has suggested no significant SNPs [66]. However, a pathway analysis has revealed an association of ADHD with SNPs involved in gene expression regulation, cell adhesion, and inflammation [30]. One study has inspected the genomic overlap between ADHD and other psychiatric disorders in 318 individuals, including 93 who are diagnosed with ADHD, and found a similar inflammation related genetic signature between ADHD and depression [67]. Segman et al. [68] have evaluated IL-1 receptor antagonist gene variable number tandem repeat polymorphism in a risk population of ADHD. As IL-1 was known to regulate murine catecholaminergic transmission, it was selected for the study. Segman et al. [68] have found an association between a 4-repeat allele and an increased risk for ADHD and an association between a 2-repeat allele and a decreased risk. However, they failed to reproduce the same results with a larger sample later [69].

While there is a good number of genetic studies supporting the association between inflammation and ADHD, there are high variations for methodologies applied in each study. Highly heterogeneous genetic features and clinical manifestation might take part in the observed variation. Currently, there is no consensus about which inflammatory-related genes precede to ADHD. Future research on populations with heterogeneous features might provide more conclusive findings.

## 5. Dysregulation of the Kynurenine Pathway in ADHD

A delay in the development of cortical maturation may cause evident deficits in neuropsychological performances in ADHD [16]. Although the etiology of this delay is unknown, impaired glial supply to support energy for neuronal activity has been suggested to have a contribution. A recent study on ADHD has proposed that patients may carry subsyndromal immunological imbalances such as increased serum IFN- $\gamma$  and IL-13 levels. It also demonstrates a decreased 3-HK despite normal levels of L-KYN [26]. Compared to medicated subjects, the alteration of pro-inflammatory cytokine production level and kynurenine metabolism showed a trend toward normalizing in medication naïve subjects. An impaired 3-HK production might be predisposed to reduced activation of microglia and hence impaired neuronal pruning that could bring in developmental delays. These reports might be congruous with early postulations about an imbalance of TRP metabolism in ADHD, suggesting that patients can produce excess serotonin, at least in peripheral compartments [70].

Although no report has directly explored cytokine and kynurenine profiles at the CNS level in ADHD, a few studies have tried to establish the association between these markers and behavioral endophenotypes by measuring their serum levels. Oades et al. have demonstrated that levels of S100b are negatively associated with oppositional and conduct problems in ADHD [71]. Their study also demonstrated an inverse relationship between S100b and IL-10/IL-16 in ADHD children. A subsequent study has reported that hyperactivity is strongly correlated with reduced S100b, while attention capacity may be related to IL-13 [26]. Increased kynurenine and IFN- $\gamma$  (though reduced TNF- $\alpha$ ) are related to faster reaction time, whereas TRP metabolism shows no relation with symptoms. Another study conducted by Oades et al. [54] has demonstrated that increases of 3-HK and IFN- $\gamma$  are linked to lower birth weight and shorter pregnancy in ADHD individuals, which are associated with the severity of symptoms. This result was only partially congruent with former reports [26] of dysregulated cytokine production and kynurenine metabolism, where a decrease in 3-HF was found. Although these findings of peripheral cytokine and kynurenine system alterations are impressive, further research is required to elucidate whether these peripheral measurements might be interpreted as changes in the

CNS compartment. In addition, a detailed analysis of cytokine levels and their relationship to the KP in the brain throughout the disease might be beneficial to research on developmental delay reported in ADHD patients.

6. Potential Inflammatory Biomarkers in ADHD

Recently, psychiatry research studies have measured and examined how individual cytokines known to be related to inflammatory processes are related to particular diagnostic categories and related phenotypes. Individual relationships of these markers with various mental disorders in perinatal and offspring outcomes, chronic states, and pre/post-treatment have been examined based on cytokines, C-reactive protein, hormones, neurotrophins, and so on. Some commonly investigated cytokine measures are summarized in **Table 1** [12, 40, 72-88]. When interpreting reports of individual studies, it is crucial to consider the extent to which other factors affecting peripheral cytokines are accounted for in specific analyses. Compounding factors including age, sex, weight, smoking, childhood trauma, the timing of blood sampling, medical comorbidities, concurrent medication use, and severity of illness are example variables that should be but are not always indicated in studies. They are possible sources of discrepancy in results [82, 89].

Table 1. Commonly Investigated Inflammation-Related Cytokines in Neuropsychiatric Disorders

Category	Protein Designation	Name	Major Function
General	CRP	C-reactive protein	Acute-phase protein produced in response to acute and chronic inflammation Produced as a result of increasing pro-inflammatory cytokines (IL-1 and IL-6) and lipopolysaccharides
	IFN- $\gamma$	Interferon gamma	Secreted by lymphocytes and is a potent activator of macrophages Critical to both innate and adaptive immunity
	IL-1 $\beta$	Interleukin 1 beta	Induces prostaglandin synthesis, neutrophil and T-cell activation, cytokine production, B-cell activation and antibody production Utilized as a biological response modifier in cancer therapy
Pro-inflammatory cytokine	IL-6	Interleukin 6	Pyrogenic, acute-phase response mediator, stimulating acute-phase protein synthesis and production of neutrophils Supports the growth of B-cell
	IL-8	Interleukin 8	Chemotactic factor in the recruitment of neutrophils and other immune cells to the site of inflammation
	IL-10	Interleukin 10	A potent anti-inflammatory cytokine that playing central role in limiting pro-inflammatory cytokines and maintaining tissue homeostasis
	IL-18	Interleukin 18	A pleiotropic cytokine produced mainly by antigen-presenting cells, playing key role in autoimmune, inflammation, and infection Enhances cytotoxic activity and proliferation of CD8+ T and NK cells



		Stimulates the production of other cytokines including IL-13
TNF- $\alpha$	Tumor necrosis factor alpha	Produced by macrophages during acute inflammation and plays a role for cell necrosis or apoptosis
TNF- $\beta$	Tumor necrosis factor beta	Involved in autoimmune disorders, mediating the inflammatory demyelination process

Adapted from Zlotnik (2012) [90] and Bishop et al. (2022) [91].

It has been reported that neuroinflammation might underlie neurodevelopment of the immune system, resulting in changes in normal microglia, astrocytes, chemokines, cytokines, oxidative stress, and related metabolism in the first months or early years of life [92]. In children with maternal inflammatory and immune system disruption, an increased profile of ADHD risk has been observed [93]. Gustafsson et al. [94] have suggested that maternal serum levels of IL-6, TNF- $\alpha$ , and monocyte chemoattractant protein-1 (MCP-1) are possible markers of ADHD risk, which is the first human study providing evidence of an association between inflammation and brain developmental/behavioral defects.

Consistently, mechanistic evidence has been demonstrated by animal studies, which show maternal immune activation in the offspring with ADHD [29]. During perinatal development, neurodevelopmental high inflammatory responses triggered by several mechanisms of environmental factors such as heavy metal exposures may increase ADHD risk conditionally [54].

Peripheral pro-inflammatory cytokines can cross the brain through humoral and neural pathways and maintain inflammatory responses via neuroimmune systems. Inflammation-related cytokine changes in the brain are known to cause neurotransmission changes in TRP metabolism and dopaminergic pathways in the brain, similar to those seen in patients with ADHD [71]. At this point, prenatal exposure to inflammation may restrain brain development resulting from structural changes in the volume of gray matter that can cause permanent neural circuits to fail to mature or bring neuroendocrine changes, thus elevating the risk of ADHD [95]. Also, interaction of the HPA axis with chronic increase of cytokine signals in the immune system during developmental processes is associated with the pathogenesis of ADHD [96]. However, these serological changes of inflammatory cytokines tend to wane or become reversed along with advances in age of ADHD patients. Therefore, the age of subjects in studies measuring potential inflammatory biomarkers of ADHD should be included when interpreting study results.

In recent studies, various inflammatory cytokines in the central and peripheral samples have been proposed as feasible potential biomarkers of ADHD risk. **Table 2** lists potential inflammatory biomarkers of ADHD risks suggested for youth and adult ADHD patients.

**Table 2.** Potential inflammatory biomarkers in youth and adult ADHD patients

Biomarkers	Youth	References	Adult <sup>1</sup>	References
CRP	↑	[75, 76]	↑	[87]
IL-1 $\beta$	↓	[71]	↔	[87]
IL-6	↑	[71, 97, 98]	↔	[86, 99, 100]
IL-10	↑	[71, 75, 98]	↔	[86]
IL-13	↑	[71, 75]	↔	[86]
IL-16	↑	[54, 71]		
TNF- $\alpha$	↓	[71, 86]	↓	[86]
Cortisol <sup>2</sup>	↓	[75, 76, 101, 102]	↓	[103]

<sup>1</sup> Compared to controls.  
<sup>2</sup> Cortisol: Morning salivary cortisol.

Chang et al. [75, 76] have evaluated CRP levels and found that its levels are elevated in youth with ADHD compared to those in healthy youth. Concurrently, TNF- $\alpha$  levels have been reported to be lower in ADHD children since IL-6 is known to have a suppressive effect on TNF- $\alpha$  production [104]. They mutually counter-regulate each other during early immune responses. Yang et al. [87] have evaluated serum CRP and other cytokines of individuals aged 5 to 55 years and provided sub-analysis results, showing that reported CRP levels are elevated in the adult group. They analyzed scores of Difficulties in Emotion Regulation Scale (DERS-16) and CRP levels and found that higher CRP levels were associated with lower DERS-16 scores.

Oades et al. [71] have investigated serum levels IL-1 $\beta$ , IL-6, IL-10, IL-13, IL-16, and TNF- $\alpha$  in 21 children (mean age: 8.9 $\pm$ 1.4 years) with ADHD who are treatment-naïve compared to the same number of controls (mean age: 12.6 $\pm$ 2.1) years. Lower IL-13 levels were seen in the ADHD medication-naïve group compared to those in the control. With the same sample, the authors further analyzed the correlation between levels of cytokines and symptom scores. An increase in IL-13 was associated with increased inattention symptoms. High IL-16 level was associated with an increase of hyperactivity-impulsive symptoms and positively related to motor activity. Another study [54] has reported that increased IL-16 level in the ADHD group is related to the poor infant health. Numerous reports have shown that increased level IL-6 levels in ADHD are associated with an increased risk of ADHD in children [71, 75, 97]. However, this evidence is diluted in advanced age groups of ADHD patients [99, 100].

Misiak et al. [86] have conducted a systematic review on possible peripheral blood inflammatory markers in ADHD patients and found no association with IL-6 or IL-10 in ADHD. However, individual case-control studies conducted by Donfrancisco et al. [98] have found that IL-6 and IL-10 levels are elevated in children with ADHD. The heterogeneity of subjective research in systematic review might be one reason for the diluted result. A systematic review conducted by Misiak et al. [86] has also found that lower TNF- $\alpha$  is meaningful in both adult and youth ADHD patients.

Chang et al. have investigated inflammatory biomarkers in ADHD and revealed that morning salivary cortisol levels are lower in ADHD youth than in health controls [75, 76]. This result is reproduced in the study of Llorens et al. [101] recently, which also reports lower morning cortisol levels in ADHD youths with consistently high levels of hyperactivity and inattention symptoms since childhood. A previous research by Isaksson et al. [102] has found a similar result between childhood trauma and ADHD symptoms regarding morning cortisol levels, with ADHD children showing a positive correlation between childhood adversity and cortisol increase after awakening. Scassellati et al. [103] have performed a systematic review on potential biomarkers in adults with ADHD and reported lower salivary cortisol levels from a meta-analysis.

Although the most recent research on ADHD biomarkers has suggested the possibility of finding more objective forms of diagnostics compared to the current diagnostic criteria in clinical use today, it remains unclear how these discrete markers are associated with diverse clinical manifestations and different populations that persistently confound research on ADHD. A few observational studies have evaluated different variables without adjusting or controlling confounders to investigate the role of inflammation in the pathophysiology of ADHD, yielding conflicting results. Further research overcoming these limitations of previous research studies performed so far is needed.

## 7. Summary

The goal of this review is to explore the possible underlying etiopathophysiology of KP in the regulation of ADHD. This reviewed summarized current knowledge on the range of possible peripheral inflammatory biomarkers. Uncertainty remains as to whether neuroinflammation is a cause or consequence of the risk of developing ADHD. However, some researchers have demonstrated that the KP and subsequent neuroinflammation

have a certain degree of association with the state of ADHD. Searching for reliable peripheral inflammatory biomarkers are of great interest in terms of eliminating uncertainty and overcoming diagnostic and treatment difficulties in the clinic. Although a solid neuroimmunological basis has been established through recent neuroimmunological advances, there is a need for further studies to determine how changes in inflammatory markers are related to disease status or whether these markers can be used to detect the development of ADHD and predict its progression and response to treatment.

**Author Contributions:** Conceptualization, writing, and constructing figures and tables, Ji Hyun Park.

**Data Availability Statement:** Not applicable.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** not applicable.

**Conflicts of Interest:** The author has no conflicts of interest relevant to this study to disclose.

## References

1. Riglin, L.; Collishaw, S.; Thapar, A. K.; Dalsgaard, S.; Langley, K.; Smith, G. D.; Stergiakouli, E.; Maughan, B.; O'Donovan, M. C.; Thapar, A., Association of genetic risk variants with attention-deficit/hyperactivity disorder trajectories in the general population. *JAMA psychiatry* **2016**, *73*, (12), 1285-1292.
2. Kim, K. M.; Ha, M.; Lim, M. H.; Kwon, H. J.; Yoo, S. J.; Kim, E.; Paik, K. C., The Symptom Trajectory of Attention-Deficit Hyperactivity Disorder in Korean School-Age Children. *Psychiatry Investig* **2018**, *15*, (5), 470-475.
3. Gau, S. S.; Chong, M.-Y.; Chen, T. H.; Cheng, A. T., A 3-year panel study of mental disorders among adolescents in Taiwan. *American Journal of Psychiatry* **2005**, *162*, (7), 1344-1350.
4. Heiervang, E.; Stormark, K. M.; Lundervold, A. J.; Heimann, M.; Goodman, R.; Posserud, M. B.; Ullebø, A. K.; Plessen, K. J.; Bjelland, I.; Lie, S. A.; Gillberg, C., Psychiatric disorders in Norwegian 8- to 10-year-olds: an epidemiological survey of prevalence, risk factors, and service use. *J Am Acad Child Adolesc Psychiatry* **2007**, *46*, (4), 438-447.
5. Polanczyk, G.; de Lima, M. S.; Horta, B. L.; Biederman, J.; Rohde, L. A., The worldwide prevalence of ADHD: a systematic review and metaregression analysis. *Am J Psychiatry* **2007**, *164*, (6), 942-8.
6. Agnew-Blais, J. C.; Polanczyk, G. V.; Danese, A.; Wertz, J.; Moffitt, T. E.; Arseneault, L., Evaluation of the persistence, remission, and emergence of attention-deficit/hyperactivity disorder in young adulthood. *JAMA psychiatry* **2016**, *73*, (7), 713-720.
7. Faraone, S. V.; Asherson, P.; Banaschewski, T.; Biederman, J.; Buitelaar, J. K.; Ramos-Quiroga, J. A.; Rohde, L. A.; Sonuga-Barke, E. J.; Tannock, R.; Franke, B., Attention-deficit/hyperactivity disorder. *Nat Rev Dis Primers* **2015**, *1*, 15020.
8. de Zeeuw, E. L.; van Beijsterveldt, C. E. M.; Ehli, E. A.; de Geus, E. J. C.; Boomsma, D. I., Attention Deficit Hyperactivity Disorder Symptoms and Low Educational Achievement: Evidence Supporting A Causal Hypothesis. *Behavior Genetics* **2017**, *47*, (3), 278-289.
9. Jangmo, A.; Stålhandske, A.; Chang, Z.; Chen, Q.; Almqvist, C.; Feldman, I.; Bulik, C. M.; Lichtenstein, P.; D'Onofrio, B.; Kuja-Halkola, R.; Larsson, H., Attention-Deficit/Hyperactivity Disorder, School Performance, and Effect of Medication. *J Am Acad Child Adolesc Psychiatry* **2019**, *58*, (4), 423-432.
10. Evangelisti, M.; De Rossi, P.; Rabasco, J.; Donfrancesco, R.; Lionetto, L.; Capi, M.; Sani, G.; Simmaco, M.; Nicoletti, F.; Villa, M. P., Changes in serum levels of kynurenine metabolites in paediatric patients affected by ADHD. *Eur Child Adolesc Psychiatry* **2017**, *26*, (12), 1433-1441.
11. Association, A. P., Diagnostic and statistical manual for mental disorders. *DSM-IIIIR 精神障害の分類と診断の手引* **1987**, 3-24.

12. Takahashi, N.; Ishizuka, K.; Inada, T., Peripheral biomarkers of attention-deficit hyperactivity disorder: Current status and future perspective. *J Psychiatr Res* **2021**, 137, 465-470.
13. Paris, J.; Bhat, V.; Thombs, B., Is adult attention-deficit hyperactivity disorder being overdiagnosed? *The Canadian Journal of Psychiatry* **2015**, 60, (7), 324-328.
14. Moynihan, R.; Doust, J.; Henry, D., Preventing overdiagnosis: how to stop harming the healthy. *Bmj* **2012**, 344, e3502.
15. Cortese, S., The neurobiology and genetics of Attention-Deficit/Hyperactivity Disorder (ADHD): what every clinician should know. *Eur J Paediatr Neurol* **2012**, 16, (5), 422-33.
16. Shaw, P.; Eckstrand, K.; Sharp, W.; Blumenthal, J.; Lerch, J. P.; Greenstein, D.; Clasen, L.; Evans, A.; Giedd, J.; Rapoport, J. L., Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *Proc Natl Acad Sci U S A* **2007**, 104, (49), 19649-54.
17. Arnsten, A. F.; Pliszka, S. R., Catecholamine influences on prefrontal cortical function: relevance to treatment of attention deficit/hyperactivity disorder and related disorders. *Pharmacol Biochem Behav* **2011**, 99, (2), 211-6.
18. Kesner, R. P.; Churchwell, J. C., An analysis of rat prefrontal cortex in mediating executive function. *Neurobiol Learn Mem* **2011**, 96, (3), 417-31.
19. Wang, M.; Ramos, B. P.; Paspalas, C. D.; Shu, Y.; Simen, A.; Duque, A.; Vijayraghavan, S.; Brennan, A.; Dudley, A.; Nou, E.; Mazer, J. A.; McCormick, D. A.; Arnsten, A. F., Alpha2A-adrenoceptors strengthen working memory networks by inhibiting cAMP-HCN channel signaling in prefrontal cortex. *Cell* **2007**, 129, (2), 397-410.
20. Bowton, E.; Saunders, C.; Erreger, K.; Sakrikar, D.; Matthies, H. J.; Sen, N.; Jessen, T.; Colbran, R. J.; Caron, M. G.; Javitch, J. A.; Blakely, R. D.; Galli, A., Dysregulation of dopamine transporters via dopamine D2 autoreceptors triggers anomalous dopamine efflux associated with attention-deficit hyperactivity disorder. *J Neurosci* **2010**, 30, (17), 6048-57.
21. Arnsten, A. F.; Paspalas, C. D.; Gamo, N. J.; Yang, Y.; Wang, M., Dynamic Network Connectivity: A new form of neuroplasticity. *Trends Cogn Sci* **2010**, 14, (8), 365-75.
22. Robbins, T. W., Dopamine and cognition. *Curr Opin Neurol* **2003**, 16 Suppl 2, S1-2.
23. Sharma, A.; Couture, J., A Review of the Pathophysiology, Etiology, and Treatment of Attention-Deficit Hyperactivity Disorder (ADHD). *Ann Pharmacother* **2014**, 48, (2), 209-225.
24. Bonvicini, C.; Faraone, S. V.; Scassellati, C., Attention-deficit hyperactivity disorder in adults: A systematic review and meta-analysis of genetic, pharmacogenetic and biochemical studies. *Molecular Psychiatry* **2016**, 21, (7), 872-884.
25. Aarsland, T. I.; Landaas, E. T.; Hegvik, T. A.; Ulvik, A.; Halmøy, A.; Ueland, P. M.; Haavik, J., Serum concentrations of kynurenines in adult patients with attention-deficit hyperactivity disorder (ADHD): a case-control study. *Behav Brain Funct* **2015**, 11, (1), 36.
26. Oades, R. D.; Myint, A.-M.; Dauvermann, M. R.; Schimmelmann, B. G.; Schwarz, M. J., Attention-deficit hyperactivity disorder (ADHD) and glial integrity: an exploration of associations of cytokines and kynurenine metabolites with symptoms and attention. *Behavioral and Brain Functions* **2010**, 6, (1), 32.
27. Comai, S.; Cavalletto, L.; Chemello, L.; Bernardinello, E.; Ragazzi, E.; Costa, C. V. L.; Bertazzo, A., Effects of PEG-interferon alpha plus ribavirin on tryptophan metabolism in patients with chronic hepatitis C. *Pharmacological Research* **2011**, 63, (1), 85-92.
28. Badawy, A. A., Kynurenine pathway of tryptophan metabolism: regulatory and functional aspects. *International Journal of Tryptophan Research* **2017**, 10, 1178646917691938.
29. Murakami, Y.; Imamura, Y.; Saito, K.; Sakai, D.; Motoyama, J., Altered kynurenine pathway metabolites in a mouse model of human attention-deficit hyperactivity/autism spectrum disorders: A potential new biological diagnostic marker. *Scientific Reports* **2019**, 9, (1), 13182.
30. Smith, T. F.; Anastopoulos, A. D.; Garrett, M. E.; Arias-Vasquez, A.; Franke, B.; Oades, R. D.; Sonuga-Barke, E.; Asherson, P.; Gill, M.; Buitelaar, J. K., Angiogenic, neurotrophic, and inflammatory system SNPs moderate the association between

- birth weight and ADHD symptom severity. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics* **2014**, 165, (8), 691-704.
31. Li, S. L.; Kam, K. W.; Chee, A. S. H.; Zhang, X. J.; Chen, L. J.; Yip, W. W. K.; Tham, C. C.; Young, A. L.; Wong, I. C. K.; Ip, P.; Yam, J. C., The association between attention-deficit/hyperactivity disorder and retinal nerve fiber/ganglion cell layer thickness measured by optical coherence tomography: a systematic review and meta-analysis. *Int Ophthalmol* **2021**, 41, (9), 3211-3221.
  32. Saccaro, L. F.; Schilliger, Z.; Perroud, N.; Piguet, C., Inflammation, Anxiety, and Stress in Attention-Deficit/Hyperactivity Disorder. *Biomedicines* **2021**, 9, (10), 1313.
  33. Benedetti, F.; Aggio, V.; Pratesi, M. L.; Greco, G.; Furlan, R., Neuroinflammation in bipolar depression. *Frontiers in Psychiatry* **2020**, 11, 71.
  34. Price, J. B.; Bronars, C.; Erhardt, S.; Cullen, K. R.; Schwieler, L.; Berk, M.; Walder, K.; McGee, S. L.; Frye, M. A.; Tye, S. J., Bioenergetics and synaptic plasticity as potential targets for individualizing treatment for depression. *Neuroscience & Biobehavioral Reviews* **2018**, 90, 212-220.
  35. Elia, J.; Glessner, J. T.; Wang, K.; Takahashi, N.; Shtir, C. J.; Hadley, D.; Sleiman, P. M.; Zhang, H.; Kim, C. E.; Robison, R., Genome-wide copy number variation study associates metabotropic glutamate receptor gene networks with attention deficit hyperactivity disorder. *Nature genetics* **2012**, 44, (1), 78-84.
  36. Moroni, F.; Russi, P.; Carlá, V.; Lombardi, G., Kynurenic acid is present in the rat brain and its content increases during development and aging processes. *Neuroscience letters* **1988**, 94, (1-2), 145-150.
  37. Campbell, B.; Charych, E.; Lee, A.; Möller, T., Kynurenines in CNS disease: regulation by inflammatory cytokines. *Frontiers in Neuroscience* **2014**, 8.
  38. Heyes, M. P.; Chen, C. Y.; Major, E. O.; Saito, K., Different kynurenine pathway enzymes limit quinolinic acid formation by various human cell types. *Biochemical Journal* **1997**, 326, (2), 351-356.
  39. Amori, L.; Guidetti, P.; Pellicciari, R.; Kajii, Y.; Schwarcz, R., On the relationship between the two branches of the kynurenine pathway in the rat brain in vivo. *Journal of neurochemistry* **2009**, 109, (2), 316-325.
  40. Ullah, I.; Awan, H. A.; Aamir, A.; Diwan, M. N.; de Filippis, R.; Awan, S.; Irfan, M.; Fornaro, M.; Ventriglio, A.; Vellante, F.; Pettorruso, M.; Martinotti, G.; Di Giannantonio, M.; De Berardis, D., Role and Perspectives of Inflammation and C-Reactive Protein (CRP) in Psychosis: An Economic and Widespread Tool for Assessing the Disease. *Int J Mol Sci* **2021**, 22, (23).
  41. Hirata, F.; Hayaishi, O., Possible participation of superoxide anion in the intestinal tryptophan 2, 3-dioxygenase reaction. *Journal of Biological Chemistry* **1971**, 246, (24), 7825-7826.
  42. Verlaet, A. A. J.; Breynaert, A.; Ceulemans, B.; De Bruyne, T.; Fransen, E.; Pieters, L.; Savelkoul, H. F. J.; Hermans, N., Oxidative stress and immune aberrancies in attention-deficit/hyperactivity disorder (ADHD): a case-control comparison. *European Child & Adolescent Psychiatry* **2019**, 28, (5), 719-729.
  43. Guidetti, P.; Amori, L.; Sapko, M. T.; Okuno, E.; Schwarcz, R., Mitochondrial aspartate aminotransferase: a third kynurenate-producing enzyme in the mammalian brain. *Journal of neurochemistry* **2007**, 102, (1), 103-111.
  44. DiNatale, B. C.; Murray, I. A.; Schroeder, J. C.; Flaveny, C. A.; Lahoti, T. S.; Laurenzana, E. M.; Omiecinski, C. J.; Perdew, G. H., Kynurenic acid is a potent endogenous aryl hydrocarbon receptor ligand that synergistically induces interleukin-6 in the presence of inflammatory signaling. *Toxicological sciences* **2010**, 115, (1), 89-97.
  45. Sathyaikumar, K. V.; Tararina, M.; Wu, H. Q.; Neale, S. A.; Weisz, F.; Salt, T. E.; Schwarcz, R., Xanthurenic Acid Formation from 3-Hydroxykynurenine in the Mammalian Brain: Neurochemical Characterization and Physiological Effects. *Neuroscience* **2017**, 367, 85-97.
  46. DE CARVALHO, L. P.; Bochet, P.; Rossier, J., The endogenous agonist quinolinic acid and the non endogenous homoquinolinic acid discriminate between NMDAR2 receptor subunits. *Neurochemistry international* **1996**, 28, (4), 445-452.



47. Goff, D. C.; Romero, K.; Paul, J.; Mercedes Perez-Rodriguez, M.; Crandall, D.; Potkin, S. G., Biomarkers for drug development in early psychosis: Current issues and promising directions. *Eur Neuropsychopharmacol* **2016**, 26, (6), 923-37.
48. Peter, F.; Andrea, S.; Nancy, A., Forty years of structural brain imaging in mental disorders: is it clinically useful or not? *Dialogues Clin Neurosci* **2018**, 20, (3), 179-186.
49. Skorobogatov, K.; De Picker, L.; Verkerk, R.; Coppens, V.; Leboyer, M.; Müller, N.; Morrens, M., Brain Versus Blood: A Systematic Review on the Concordance Between Peripheral and Central Kynurenine Pathway Measures in Psychiatric Disorders. *Front Immunol* **2021**, 12, 716980.
50. Gal, E.; Sherman, A., Synthesis and metabolism of L-kynurenine in rat brain. *Journal of neurochemistry* **1978**, 30, (3), 607-613.
51. Fukui, S.; Schwarcz, R.; Rapoport, S. I.; Takada, Y.; Smith, Q. R., Blood-brain barrier transport of kynurenines: implications for brain synthesis and metabolism. *Journal of neurochemistry* **1991**, 56, (6), 2007-2017.
52. Kita, T.; Morrison, P. F.; Heyes, M. P.; Markey, S., Effects of systemic and central nervous system localized inflammation on the contributions of metabolic precursors to the L-kynurenine and quinolinic acid pools in brain. *Journal of neurochemistry* **2002**, 82, (2), 258-268.
53. Moreau, M.; André, C.; O'Connor, J. C.; Dumich, S. A.; Woods, J. A.; Kelley, K. W.; Dantzer, R.; Lestage, J.; Castanon, N., Inoculation of Bacillus Calmette-Guerin to mice induces an acute episode of sickness behavior followed by chronic depressive-like behavior. *Brain, behavior, and immunity* **2008**, 22, (7), 1087-1095.
54. Oades, R. D., An exploration of the associations of pregnancy and perinatal features with cytokines and tryptophan/kynurenine metabolism in children with attention-deficit hyperactivity disorder (ADHD). *Atten Defic Hyperact Disord* **2011**, 3, (4), 301-18.
55. Lestage, J.; Verrier, D.; Palin, K.; Dantzer, R., The enzyme indoleamine 2, 3-dioxygenase is induced in the mouse brain in response to peripheral administration of lipopolysaccharide and superantigen. *Brain, behavior, and immunity* **2002**, 16, (5), 596-601.
56. Connor, T. J.; Starr, N.; O'Sullivan, J. B.; Harkin, A., Induction of indoleamine 2, 3-dioxygenase and kynurenine 3-monooxygenase in rat brain following a systemic inflammatory challenge: a role for IFN- $\gamma$ ? *Neuroscience letters* **2008**, 441, (1), 29-34.
57. Fujigaki, H.; Seishima, M.; Saito, K., Posttranslational modification of indoleamine 2, 3-dioxygenase. *Analytical and bioanalytical chemistry* **2012**, 403, (7), 1777-1782.
58. Zunszain, P. A.; Anacker, C.; Cattaneo, A.; Choudhury, S.; Musaelyan, K.; Myint, A. M.; Thuret, S.; Price, J.; Pariante, C. M., Interleukin-1 $\beta$ : a new regulator of the kynurenine pathway affecting human hippocampal neurogenesis. *Neuropsychopharmacology* **2012**, 37, (4), 939-949.
59. Jung, I. D.; Lee, C.-M.; Jeong, Y.-I.; Lee, J. S.; Park, W. S.; Han, J.; Park, Y.-M., Differential regulation of indoleamine 2, 3-dioxygenase by lipopolysaccharide and interferon gamma in murine bone marrow derived dendritic cells. *FEBS letters* **2007**, 581, (7), 1449-1456.
60. Tu, H.; Rady, P. L.; Juelich, T.; Smith, E. M.; Tying, S. K.; Hughes, T. K., Cytokine regulation of tryptophan metabolism in the hypothalamic-pituitary-adrenal (HPA) axis: implications for protective and toxic consequences in neuroendocrine regulation. *Cellular and molecular neurobiology* **2005**, 25, (3), 673-680.
61. Yadav, M. C.; Burudi, E.; Alirezaei, M.; Flynn, C. C.; Watry, D. D.; Lanigan, C. M.; Fox, H. S., IFN- $\gamma$ -induced IDO and WRS expression in microglia is differentially regulated by IL-4. *Glia* **2007**, 55, (13), 1385-1396.
62. Robinson, C. M.; Shirey, K. A.; Carlin, J. M., Synergistic transcriptional activation of indoleamine dioxygenase by IFN- $\gamma$  and tumor necrosis factor- $\alpha$ . *Journal of interferon & cytokine research* **2003**, 23, (8), 413-421.
63. Alberati-Giani, D.; Ricciardi-Castagnoli, P.; Köhler, C.; Cesura, A., Regulation of the kynurenine pathway by IFN- $\gamma$  in murine cloned macrophages and microglial cells. In *Recent Advances in Tryptophan Research*, Springer: 1996; pp 171-175.

64. Odell, D.; Warren, R. P.; Warren, L.; Burger, R. A.; Maciulis, A., Association of genes within the major histocompatibility complex with attention deficit hyperactivity disorder. *Neuropsychobiology* **1997**, 35, (4), 181-186.
65. Demontis, D. e. a., Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nature Genetics* **2019**, 51, (1), 63-75.
66. Zayats, T.; Athanasiu, L.; Sonderby, I.; Djurovic, S.; Westlye, L. T.; Tamnes, C. K.; Fladby, T.; Aase, H.; Zeiner, P.; Reichborn-Kjennerud, T.; Knappskog, P. M.; Knudsen, G. P.; Andreassen, O. A.; Johansson, S.; Haavik, J., Genome-wide analysis of attention deficit hyperactivity disorder in Norway. *PLoS One* **2015**, 10, (4), e0122501.
67. de Jong, S.; Newhouse, S. J.; Patel, H.; Lee, S.; Dempster, D.; Curtis, C.; Paya-Cano, J.; Murphy, D.; Wilson, C. E.; Horder, J.; Mendez, M. A.; Asherson, P.; Rivera, M.; Costello, H.; Maltezos, S.; Whitwell, S.; Pitts, M.; Tye, C.; Ashwood, K. L.; Bolton, P.; Curran, S.; McGuffin, P.; Dobson, R.; Breen, G., Immune signatures and disorder-specific patterns in a cross-disorder gene expression analysis. *Br J Psychiatry* **2016**, 209, (3), 202-8.
68. Segman, R. H.; Meltzer, A.; Gross-Tsur, V.; Kosov, A.; Frisch, A.; Inbar, E.; Darvasi, A.; Levy, S.; Goltser, T.; Weizman, A.; Galili-Weisstub, E., Preferential transmission of interleukin-1 receptor antagonist alleles in attention deficit hyperactivity disorder. *Mol Psychiatry* **2002**, 7, (1), 72-4.
69. Misener, V. L.; Schachar, R.; Ickowicz, A.; Malone, M.; Roberts, W.; Tannock, R.; Kennedy, J. L.; Pathare, T.; Barr, C. L., Replication test for association of the IL-1 receptor antagonist gene, IL1RN, with attention-deficit/hyperactivity disorder. *Neuropsychobiology* **2004**, 50, (3), 231-4.
70. Irwin, M.; Belendiuk, K.; McCloskey, K.; Freedman, D. X., Tryptophan metabolism in children with attentional deficit disorder. *The American Journal of Psychiatry* **1981**.
71. Oades, R. D.; Dauvermann, M. R.; Schimmelmann, B. G.; Schwarz, M. J.; Myint, A.-M., Attention-deficit hyperactivity disorder (ADHD) and glial integrity: S100B, cytokines and kynurenine metabolism-effects of medication. *Behavioral and Brain Functions* **2010**, 6, (1), 1-14.
72. 2 Biomarkers of Neuroinflammation: Challenges and Potential Opportunities. In *National Academies of Sciences, Engineering, and Medicine. Biomarkers of Neuroinflammation: Proceedings of a Workshop*, 2018.
73. Fond, G.; Micoulaud-Franchi, J. A.; Faugere, M.; Boyer, L.; Faget-Agius, C.; Lançon, C.; Richieri, R.; Cermolacce, M., Abnormal C-reactive protein blood levels as a specific biomarker of major depression and non-remission under antidepressants in schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* **2020**, 97, 109800.
74. Lin, P.; Sun, J.; Lou, X.; Li, D.; Shi, Y.; Li, Z.; Ma, P.; Li, P.; Chen, S.; Jin, W.; Liu, S.; Chen, Q.; Gao, Q.; Zhu, L.; Xu, J.; Zhu, M.; Wang, M.; Liang, K.; Zhao, L.; Xu, H.; Dong, K.; Li, Q.; Cheng, X.; Chen, J.; Guo, X., Consensus on potential biomarkers developed for use in clinical tests for schizophrenia. *General Psychiatry* **2022**, 35, (1), e100685.
75. Chang, J. P.; Su, K. P.; Mondelli, V.; Pariante, C. M., Cortisol and inflammatory biomarker levels in youths with attention deficit hyperactivity disorder (ADHD): evidence from a systematic review with meta-analysis. *Transl Psychiatry* **2021**, 11, (1), 430.
76. Chang, J. P.; Mondelli, V.; Satyanarayanan, S. K.; Chiang, Y. J.; Chen, H. T.; Su, K. P.; Pariante, C. M., Cortisol, inflammatory biomarkers and neurotrophins in children and adolescents with attention deficit hyperactivity disorder (ADHD) in Taiwan. *Brain Behav Immun* **2020**, 88, 105-113.
77. Murata, S.; Castillo, M. F. R.; Murphy, M.; Schwarz, M.; Moll, N.; Martin, B.; Weidinger, E.; Leitner, B.; Mueller, N.; Halaris, A., Effects of inflammation modulation on tryptophan and kynurenine pathway regulation in treatment resistant bipolar depression. *Neurology, Psychiatry and Brain Research* **2019**, 33, 65-72.
78. Bigseth, T. T.; Engh, J. A.; Egeland, J.; Andersen, E.; Andreassen, O. A.; Bang-Kittilsen, G.; Falk, R. S.; Holmen, T. L.; Lindberg, M.; Mordal, J.; Nielsen, J.; Steen, N. E.; Ueland, T.; Vang, T.; Fredriksen, M., Exploring low grade inflammation by soluble urokinase plasminogen activator receptor levels in schizophrenia: a sex-dependent association with depressive symptoms. *BMC Psychiatry* **2021**, 21, (1), 527.

79. Leboyer, M.; Berk, M.; Yolken, R. H.; Tamouza, R.; Kupfer, D.; Groc, L., Immuno-psychiatry: an agenda for clinical practice and innovative research. *BMC Med* **2016**, 14, (1), 173.
80. Dudzińska, E.; Szymona, K.; Bogucki, J.; Koch, W.; Cholewińska, E.; Sitarz, R.; Ognik, K., Increased Markers of Oxidative Stress and Positive Correlation Low-Grade Inflammation with Positive Symptoms in the First Episode of Schizophrenia in Drug-Naïve Patients. *J Clin Med* **2022**, 11, (9).
81. Lee, E. E.; Hong, S.; Martin, A. S.; Eyler, L. T.; Jeste, D. V., Inflammation in Schizophrenia: Cytokine Levels and Their Relationships to Demographic and Clinical Variables. *Am J Geriatr Psychiatry* **2017**, 25, (1), 50-61.
82. Yuan, N.; Chen, Y.; Xia, Y.; Dai, J.; Liu, C., Inflammation-related biomarkers in major psychiatric disorders: a cross-disorder assessment of reproducibility and specificity in 43 meta-analyses. *Translational Psychiatry* **2019**, 9, (1), 233.
83. Morgan, A. R.; Touchard, S.; Leckey, C.; O'Hagan, C.; Nevado-Holgado, A. J.; Consortium, N.; Barkhof, F.; Bertram, L.; Blin, O.; Bos, I.; Dobricic, V.; Engelborghs, S.; Frisoni, G.; Frölich, L.; Gabel, S.; Johannsen, P.; Kettunen, P.; Kłoszewska, I.; Legido-Quigley, C.; Lleó, A.; Martinez-Lage, P.; Mecocci, P.; Meersmans, K.; Molinuevo, J. L.; Peyratout, G.; Popp, J.; Richardson, J.; Sala, I.; Scheltens, P.; Streffer, J.; Soininen, H.; Tainta-Cuezva, M.; Teunissen, C.; Tsolaki, M.; Vandenberghe, R.; Visser, P. J.; Vos, S.; Wahlund, L.-O.; Wallin, A.; Westwood, S.; Zetterberg, H.; Lovestone, S.; Morgan, B. P.; Disorders, A. N. W. T. C. f. N. o. M.; Disease, A. s., Inflammatory biomarkers in Alzheimer's disease plasma. *Alzheimer's & Dementia* **2019**, 15, (6), 776-787.
84. Ceyhun, H. A.; Gürbüz, N., New Hematological Parameters as Inflammatory Biomarkers: Systemic Immune Inflammation Index, Platerethritis, and Platelet Distribution Width in Patients with Adult Attention Deficit Hyperactivity Disorder. *Advances in Neurodevelopmental Disorders* **2022**, 6, (2), 211-223.
85. Lee, E. E.; Sears, D. D.; Liu, J.; Jin, H.; Tu, X. M.; Eyler, L. T.; Jeste, D. V., A novel biomarker of cardiometabolic pathology in schizophrenia? *J Psychiatr Res* **2019**, 117, 31-37.
86. Misiak, B.; Wójta-Kempa, M.; Samochowiec, J.; Schiweck, C.; Aichholzer, M.; Reif, A.; Samochowiec, A.; Stańczykiewicz, B., Peripheral blood inflammatory markers in patients with attention deficit/hyperactivity disorder (ADHD): A systematic review and meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry* **2022**, 118, 110581.
87. Yang, L. L.; Stiernborg, M.; Skott, E.; Söderström, Å.; Giacobini, M.; Lavebratt, C., Proinflammatory mediators and their associations with medication and comorbid traits in children and adults with ADHD. *Eur Neuropsychopharmacol* **2020**, 41, 118-131.
88. Leffa, D. T.; Torres, I. L. S.; Rohde, L. A., A Review on the Role of Inflammation in Attention-Deficit/Hyperactivity Disorder. *Neuroimmunomodulation* **2018**, 25, (5-6), 328-333.
89. Pillinger, T.; Osimo, E. F.; Brugger, S.; Mondelli, V.; McCutcheon, R. A.; Howes, O. D., A meta-analysis of immune parameters, variability, and assessment of modal distribution in psychosis and test of the immune subgroup hypothesis. *Schizophrenia bulletin* **2019**, 45, (5), 1120-1133.
90. Zlotnik, A.; Yoshie, O., The chemokine superfamily revisited. *Immunity* **2012**, 36, (5), 705-16.
91. Bishop, J. R.; Zhang, L.; Lizano, P., Inflammation Subtypes and Translating Inflammation-Related Genetic Findings in Schizophrenia and Related Psychoses: A Perspective on Pathways for Treatment Stratification and Novel Therapies. *Harv Rev Psychiatry* **2022**, 30, (1), 59-70.
92. Corona, J. C., Role of oxidative stress and neuroinflammation in attention-deficit/hyperactivity disorder. *Antioxidants* **2020**, 9, (11), 1039.
93. Instanes, J. T.; Halmøy, A.; Engeland, A.; Haavik, J.; Furu, K.; Klungsøyr, K., Attention-Deficit/Hyperactivity Disorder in Offspring of Mothers With Inflammatory and Immune System Diseases. *Biological Psychiatry* **2017**, 81, (5), 452-459.
94. Gustafsson, H. C.; Sullivan, E. L.; Battison, E. A. J.; Holton, K. F.; Graham, A. M.; Karalunas, S. L.; Fair, D. A.; Loftis, J. M.; Nigg, J. T., Evaluation of maternal inflammation as a marker of future offspring ADHD symptoms: A prospective investigation. *Brain Behav Immun* **2020**, 89, 350-356.

95. Kozłowska, E.; Brzezińska-Błaszczyk, E.; Agier, J.; Wysokiński, A.; Żelechowska, P., Alarmins (IL-33, sST2, HMGB1, and S100B) as potential biomarkers for schizophrenia. *J Psychiatr Res* **2021**, 138, 380-387.
96. Chen, L.; Zheng, W. H.; Du, Y.; Li, X. S.; Yu, Y.; Wang, H.; Cheng, Y., Altered Peripheral Immune Profiles in First-Episode, Drug-Free Patients With Schizophrenia: Response to Antipsychotic Medications. *Front Med (Lausanne)* **2021**, 8, 757655.
97. Elsadek, A.; Al-shokary, A.; Abdelghani, W.; Kamal, N.; Ibrahim, A.; El-shorbagy, H.; Suliman, H.; Barseem, N.; Abdel Maksoud, Y.; Azab, S.; Nour El Din, D., Serum levels of interleukin-6 and tumor necrosis factor alpha in children with attention-deficit hyperactivity disorder. *Journal of Pediatric Neurosciences* **2020**, 15, (4), 402-408.
98. Donfrancesco, R.; Nativio, P.; Borrelli, E.; Giua, E.; Andriola, E.; Villa, M. P., Serum cytokines in paediatric neuropsychiatric syndromes: focus on Attention Deficit Hyperactivity Disorder. *Minerva pediatrica* **2016**.
99. Corominas-Roso, M.; Armario, A.; Palomar, G.; Corrales, M.; Carrasco, J.; Richarte, V.; Ferrer, R.; Casas, M.; Ramos-Quiroga, J. A., IL-6 and TNF- $\alpha$  in unmedicated adults with ADHD: Relationship to cortisol awakening response. *Psychoneuroendocrinology* **2017**, 79, 67-73.
100. Leffa, D. T.; Caye, A.; Santos, I.; Matijasevich, A.; Menezes, A.; Wehrmeister, F. C.; Oliveira, I.; Vitola, E.; Bau, C. H. D.; Grevet, E. H.; Tovo-Rodrigues, L.; Rohde, L. A., Attention-deficit/hyperactivity disorder has a state-dependent association with asthma: The role of systemic inflammation in a population-based birth cohort followed from childhood to adulthood. *Brain, Behavior, and Immunity* **2021**, 97, 239-249.
101. Llorens, M.; Barba, M.; Torralbas, J.; Nadal, R.; Armario, A.; Gagliano, H.; Betriu, M.; Urraca, L.; Pujol, S.; Montalvo, I.; Gracia, R.; Giménez-Palop, O.; Palao, D.; Pàmias, M.; Labad, J., Stress-related biomarkers and cognitive functioning in adolescents with ADHD: Effect of childhood maltreatment. *Journal of Psychiatric Research* **2022**, 149, 217-225.
102. Isaksson, J.; Nilsson, K. W.; Lindblad, F., Early psychosocial adversity and cortisol levels in children with attention-deficit/hyperactivity disorder. *European Child & Adolescent Psychiatry* **2013**, 22, (7), 425-432.
103. Scassellati, C.; Bonvicini, C.; Faraone, S. V.; Gennarelli, M., Biomarkers and Attention-Deficit/Hyperactivity Disorder: A Systematic Review and Meta-Analyses. *Journal of the American Academy of Child & Adolescent Psychiatry* **2012**, 51, (10), 1003-1019.e20.
104. Yimin; Kohanawa, M.; Minagawa, T., Up-regulation of granulomatous inflammation in interleukin-6 knockout mice infected with *Rhodococcus aurantiacus*. *Immunology* **2003**, 110, (4), 501-506.