

## Article

# Analysis of the Spatio-temporal Rainfall Variability in Cameroon over the Period 1950 to 2019

Valentin Brice Ebodé<sup>1,\*</sup><sup>1</sup> Department of Geography, University of Yaounde 1, P.O. Box 755, Yaounde, Cameroon; ebode-brisol@gmail.com

\* Correspondence: ebodebriso@gmail.com (V.B.E.)

**Abstract:** The rainfall study in the long term is essential for climatic change understanding and socioeconomic development. The main goal of this study is to explore the spatial and temporal variations of precipitation in different time scales (seasonal and annual) in Cameroon. The Mann–Kendall and Pettitt tests were applied to analyze the precipitation variability. On temporal plan, the different regions of Cameroon have recorded significant drops in annual rainfall that Pettitt's test generally situates around the 1970s. The decreases observed for the northern part of Cameroon regions are between -5.4% (Adamawa) and -7.4% (Far North). Those of western part regions oscillate between -7.5% (South-West) and -12.5% (West). The southern part of Cameroon regions recorded decreases varying between -4.3% (East) and -5.9% (Center). On spatial plan, the divisions of the northern, western and southern parts of Cameroon respectively recorded after the 1970s (a pivotal period in the evolution of precipitation on temporal plan), a precipitation decrease towards the South, the South-West and the West. This study's findings could be helpful for planning and managing water resources in Cameroon.

**Keywords:** Cameroon; rainfall; long-term variability; trend tests

## 1. Introduction

In the context of current climate change, it is helpful to characterize the Spatio-temporal variability of precipitation to correctly assess its impact on natural resources (water, vegetation, etc.) and certain important activities (agriculture, production of hydro-power, etc.) [1-4]. In some sub-Saharan African countries like Cameroon, the population essentially depends on small-scale agriculture for their livelihood. However, agricultural practices in this region mainly depend on the rainfed farming system. It implies that rainfall plays an essential role in sustaining livelihoods and economic development across this zone of the continent. Thus, any change in the spatial distribution of rainfall and precipitation amounts can have many consequences on agriculture and other socio-economic activities [5].

The majority of studies on the evolution of precipitation in sub-Saharan Africa have been carried out generally at the regional scale [6-10], since most of these studies were conducted where observational rainfall data were available. Let us briefly recall that after the 1990s, long-time series of rainfall became rare in sub-Saharan Africa. This scarcity of data is marked even more in Central Africa [11-12]. The active network now only includes 35 stations in the DRC (i.e. 1 station for 67000 km<sup>2</sup>), 22 in Cameroon and less than 20 for all of Gabon, Congo and Central African Republic [11]. Research dealing directly with the climate and its impact on natural resources (water, vegetation, etc.) is barely emerging in this part of Africa after the 2000s, despite its importance for the sustainable and effective management of these resources.

In sub-Saharan Africa, works devoted to rainfall variability study have been carried out mainly since the 1980s. Many of these works are based on trends and ruptures detection in the rainfall data. The obtained results confirm in West Africa's case a decline from the 1970s [13-16]. In Central Africa, the most perceptible rainfall fluctuations have been observed on a seasonal time scale [17-19]. Ebodé et al. [20] recently highlighted in the case of the Nyong basin that the most significant rainfall variations are those of the dry seasons (summer and winter). Summer rainfall generally increases around the 1970s, while winter rainfall decreases.

In Cameroon, some authors have studied the evolution of rainfall at the national and regional scales [17,21,22]. However, most studies were limited to annual and seasonal series analyses for individual stations or groups of stations. In this study, a much broader vision was adopted and the Spatio-temporal variability of precipitation was approached at the same time steps for the different regions of the country. Beyond that, this study is based on the series that include the two recent decades (2000 and 2010), which is not the case with previous studies. Finally, older studies focus primarily on trend analysis. They have rarely put into perspective the need to examine the variation of the short-term upward and downward sub-trends that characterize the fluctuations of the rainfall series. Separating sub-trends over short periods (e.g. decades) from that of the long period allows physical explanations to be attached to upward and downward trends in the data [23]. In addition, analyses of the correlation between rainfall and possible forcings (such as sea surface temperature) can help predict future rainfall evolution, which is fundamental for planning water resource uses.

This study is devoted to the evolution of rainfall analyses in all the different regions of Cameroon over a recent period, using updated rainfall series. It appears fundamental for this country, whose economy is based essentially on rain-fed agriculture. In addition, the new data and inputs could help in the water resources demand long-term planning as well as the rainfall evolution forecast improving in these regions. Its main objective is to document the trend and amplitude of changes in the precipitation regime in the studied regions.

## 2. Materials and Methods

### 2.1. Study area


Cameroon (475442 km<sup>2</sup>) is a country in the Gulf of Guinea (Central Africa). Very extensive in Latitude (1200 km from North to South), the country has the shape of a triangle whose base runs along the 2nd degree of northern Latitude while the summit, bordering Lake Chad, reaches the 13th parallel. It is limited by Nigeria to the North-North-West, Chad to the North-North-East, the Central African Republic to the East, the Republic of Congo to the South-East, Gabon to the South, Equatorial Guinea and the Gulf of Guinea to the South-West (Figure 1).

The relief of Cameroon is varied. We find the lowlands (Mamfe basin in the South-West, Benoue basin and the northern plain); the plateaus such as the South Cameroonian plateau, with an average altitude of 650 m and the Adamaua plateau, whose average altitude is 1000 m; the highlands, the best known of which are the Mandara mountains (Far North), Alantika (North) and the still-active volcanoes of Oku (North-West) and Mount Cameroon (South-West) which is, at an altitude of 4095 m, the highest point in West Africa.

Climatically, the Adamaua Plateau separates "wet/equatorial" Cameroon from "dry/tropical" Cameroon. The equatorial domain (southern Cameroon plateau, western highlands and coastal lowlands) is characterized by abundant rainfall (Tables 1 and 2). On the South Cameroonian plateau (encompassing the administrative regions of the Center, South and East), there are four distinct seasons (spring, from march to june; summer, july and august; autumn, from september to november; winter, from december to february)

(Table 1). In the Western Highlands (encompassing the North-West, West and South-West Administrative Regions) and the Coastal Lowlands (Littoral), heavy rains fall for nine consecutive months from March to November (Table 2). The tropical domain is distinguished by scanty rainfall. It is characterized by a rainy season from April or May to October.

The vegetation encountered on Cameroonian territory is diversified and can be divided into two large zones: the equatorial part and the tropical part. The vegetation of the equatorial part is composed of the dense humid forest of the South and East formed of very tall trees, of the gallery forests of the West and North-West along the watercourses and in the lowlands funds and mangroves on the Littoral and South-West coasts. The tropical part is covered with savannah. We find there, the wooded savannah of Adamawa rich in shrubs, the grassy savannah of the North and the steppe of the Far North, which is poor in trees and grass [24].



**Figure 1.** Region map of Cameroon. The numbers at the top in each region indicate the area in km<sup>2</sup>, and those below indicate the percentage (%) of occupation of this region in the national territory (Cameroon).

## 2.2. Data sources

The rainfall series used in this work are those of the Climate Research Unit (CRU). CRU data are available since 1901, via the site [https://climexp.knmi.nl/select-field\\_obs2.cgi?id=2833fad3fef1bedc6761d5cba64775f0/](https://climexp.knmi.nl/select-field_obs2.cgi?id=2833fad3fef1bedc6761d5cba64775f0/). They are available in NetCDF format, on a monthly time step and at a spatial resolution of 0.25° x 0.25°. This precipitation product is obtained from gauge interpolations.

## 2.3. Data analysis

The analysis of rainfall (at seasonal and annual time scales) was carried out using Mann-Kendall [25] and Pettitt [26] tests at the 95% significance level.

To analyse rainfall series, the Mann-Kendall test was first used to indicate if there are trends in the lasts and if these trends are significant. This test is based on the test statistic "S" defined as follows:

$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sgn}(x_j - x_i)$$

where the  $x_j$  are the sequential data values;  $n$  is the length of the data set; and  $\text{sgn} = (\theta)$  if  $\theta > 1$ , 0 if  $\theta = 0$  and -1 if  $\theta < 0$ . There is no significant trend in the series analyzed when the calculated p-value is above the chosen significance level.

In addition to the Mann-Kendall test, the Pettitt test was also used to analyze rainfall series. The approach after Pettitt [26] is commonly applied to detect a single change-point in hydrological series or climate series with continuous data. It tests the  $H_0$ : The  $T$  variables follow one or more distributions that have the same location parameter (no change), against the alternative: a change point exists. The Pettitt-test is conducted in such a way:

$$U_t, N = \sum_{i=1}^t \sum_{j=t+1}^N D_{ij}$$

with  $D_{ij} = \text{Sign}(X_i - X_j)$  with:  $\text{sign}(x) = 1$  if  $x > 0$ , 0 if  $x = 0$  and -1 if  $x < 0$ . If the null hypothesis is rejected, an estimate of the date of discontinuity is given by defining the maximum in the absolute value of the variable  $U_t, N$ .

**Table 1.** Statistical properties of annual and seasonal rainfall for the regions of the south Side of the country (Center, East and South). Each of these regions has 4 seasons.

	Annual			Spring			Summer			Autumn			Winter		
	Center	East	South	Center	East	South	Center	East	South	Center	East	South	Center	East	South
Mean rainfall (mm)	1784	1580	1891	664	589	736	390	287	222	673	597	795	57	106	172
Maximum rainfall (mm)	2057	1869	2368	847	746	966	511	422	364	828	750	917	120	180	306
Minimum rainfall (mm)	1479	1348	1356	472	448	526	232	159	85	524	458	544	10.2	38.4	57.9
CV	0.07	0.065	0.09	0.10	0.09	0.12	0.14	0.2	0.27	0.09	0.09	0.10	0.41	0.31	0.3

**Table 2.** Statistical properties of annual and seasonal rainfall for the regions of the North (Adamawa, North and Far North) and West (Littoral, North West, West and South West) Sides of the country. Each of these regions has 2 seasons.

	Adamawa	North	Far North	Littoral	North West	West	South West
Annual							
Mean rainfall (mm)	1540	1174	726	2387	1911	2095	2413
Maximum rainfall (mm)	1799	1356	878	2834	2334	2457	2907
Minimum rainfall (mm)	1305	881	444	1849	1670	1824	2044
CV	0.07	0.08	0.13	0.09	0.07	0.08	0.09
Wet season							
Mean rainfall (mm)	1437	1083	709	2288	1883	2052	2321
Maximum rainfall (mm)	1647	1283	861	2720	2309	2434	2845

Minimum rainfall (mm)	1196	834.2	431	1785	1646	1786	1962
CV	0.07	0.08	0.13	0.09	0.07	0.08	0.09
Dry season							
Mean rainfall (mm)	103	91	17.7	100	28	43	31
Maximum rainfall (mm)	200	164.4	49.8	184.3	86	105	63
Minimum rainfall (mm)	27	22.74	0.6	22.24	6	9.52	7
CV	0.4	0.34	0.64	0.38	0.52	0.45	0.39

### 3. Results and discussion

The precipitation evolution results are presented by zone, including regions with similar climatic conditions from the point of view of the volumes precipitated and the seasonal distribution of the rains. Thus, the ten regions have been grouped into three main zones, namely: the northern part (including Adamaua, the North and the Far North), the western part (including the Littoral, the North-West, the West and the South-West) and the Southern part (including the Centre, the South and the East).

#### 3.1. Interannual evolution of annual and seasonal rainfall

##### 3.1.1. In the northern part


The annual rainfall is decreasing in the regions of the northern part of Cameroon. According to the Mann-Kendall test, these decreases are statistically significant only for the North and Adamaua regions (Figure 2). The Pettitt test revealed breaks in the annual rainfall series in these regions in 1962-63 (Far North) and 1979-80 (North and Adamaua). The deficits recorded following these ruptures are -5.4% (Adamaua), -6.8% (North) and -7.4% (Far North) (Figure 3). Through these deficits, we notice that the decrease in rainfall becomes more significant as one progresses in latitude towards the North. The decadal deviations from the interannual mean analysis shows a general drop in rainfall in the northern part regions of Cameroon from the 1970s (Table 3). However, in the Adamaua and North regions, this decrease was only periodically interrupted during the 1990s (Table 3).

In the Far North region, on the other hand, the increase that began in the 1990s seems to have lasted until the current decade (Table 3). The precipitation evolution in the Adamaua and North regions in recent decades is similar to that observed in equatorial central Africa [19,20,27]. Contemporary rainfall variations in the Far North region are similar to what has been observed in West Africa and near [15,16,28,29].


The rainfall of the rainy season recorded the same evolutions as the annual rainfall. The decreases noted are statistically significant according to the Mann-Kendall test only for the Adamaua and North regions (Figure 2). According to Pettitt's test, the breaks occurred in this season's rainfall series in the same years as the annual rainfall. The deficits recorded following these ruptures are between -4.4% (Adamaua) and -6.4% (Far North) (Figure 3). The decadal deviations of rainfall in the rainy season in these three regions are similar to those of annual rainfall (Table 3). These similar trends observed between the annual rainfall and those of the rainy season are quite logical insofar as, for these three regions, the rainfall of the rainy season represents the bulk of the total rainfall recorded during the year.

The rainfall of the dry season is decreasing in the regions of the northern part of Cameroon. These decreases are all statistically significant according to the Mann-Kendall test (Figure 2). According to the Pettitt test, the breaking point occurred in the Adamaua series in 1964-65 (-28%). The rupture occurred in the North and Far North series in 1996-97. The variations recorded following the latter are -27.7% and -26.8% (Figure 3). The analysis of decadal deviations shows in the case of Adamaua, a general decrease in rainfall since

the 1970s (Table 3). In the case of the other two regions, the decrease in rainfall that occurred during the 1980s was only periodically interrupted during the following decade (Table 3).



**Figure 2.** Results of annual and seasonal rainfall for the regions of the North side of the country (Adamawa (3), North (2) and Far North (1)), using Mann–Kendall's test. There is a significant trend in the series when the calculated p-values are below the significance level ( $\alpha = 0.05$ ). Significant trends are indicated by values in bold.



**Figure 3.** Temporal variation of annual and seasonal rainfall for the regions of the North side of the country (Adamawa, North and Far North), according to the Pettitt test. The rupture years are indicated by the vertical dashed lines. On their right sides, appear the corresponding rates of change. The ruptures in black are those which are identical to several regions.

**Table 3.** Deviations (%) of the decadal annual and seasonal averages of rainfall compared to their interannual means for the regions of the North (Adamawa, North and Far North) and West (Littoral, North West, West and South West) Sides of the country.


Decades	Adamawa	North	Far North	Littoral	North West	West	South West
Annual							
1950	3.9	5.3	11.5	8.3	6.5	7.6	9.1
1960	4.8	6.5	4.5	8.2	5.2	6.7	9.4
1970	0.7	0.6	-2.5	-0.4	0.1	0.6	-1.3
1980	-5.3	-6.8	-12.4	-5.9	-4.2	-4.4	-6.3
1990	1.8	3.3	-0.3	-4.5	-0.7	-3	-3.5
2000	-4.7	-4.8	-1.1	1.1	-1.5	-0.9	-0.7
2010	-2.5	-3.7	1.2	-6.5	-5.4	-6.6	-6.8
Wet season							
1950	1.7	4.4	11.2	7.5	6	7.1	8.2
1960	4.6	4.8	3.3	8.1	5.5	6.9	9.5
1970	1.3	0.2	-3.1	-1	-0.3	0.2	-2
1980	-5	-6.2	-12.4	-5	-3.9	-4.1	-5.4
1990	2.4	3.2	-0.8	-4.1	-0.3	-2.6	-2.8
2000	-2.7	-4.1	-0.2	1.2	-1.5	-0.9	-0.6
2010	-2.4	-2.1	2.1	-6.5	-5.4	-6.6	-6.9
Dry season							
1950	34.8	16.6	19	26.4	39.1	35.4	32.1
1960	8.4	25.8	45	10	-16.5	-4.2	7.4
1970	-8.2	5.3	20.2	13.1	30.9	20	14.8
1980	-8.8	-13.7	-16.5	-26	-22.1	-18	-27.2
1990	-6	4.4	16	-14.4	-31.5	-23	-19.9
2000	-15.1	-13.1	-42.8	-0.1	-1.5	-2.1	-1
2010	-4	-22.3	-40.3	-5.9	-5	-6.9	-5.7

### 3.1.2. In the western part


The annual rainfall in the western part regions of Cameroon decreases significantly according to the Mann-Kendall test (Figure 4). According to the Pettitt test, a rupture was identified in their various series in 1971-72. The deficits recorded following this rupture are between -7.5% (South West) and -12.5% (West) (Figure 5). The analysis of decadal deviations shows that the decline in rainfall that began in the regions of the western part of the country during the 1970s was only interrupted during the 1990s (Table 3). It also emerges from this analysis that the 2010 decade was the driest ever observed in these regions.

As in the annual time step, the rainfall of the rainy season decreases significantly according to the Mann-Kendall test in the different regions of the western part of Cameroon (Figure 4). According to the Pettitt test, they also record a break in 1971-72. The deficits recorded following this rupture vary between -7.3% (South West) and -11.7% (West) (Figure 5). The decadal variations of this season's rainfall are similar to those highlighted for the annual rainfall (Table 3).

According to the Mann-Kendall test, the decreases observed for dry season rainfall are not significant in the case of the North-West and West regions (Figure 4). Pettitt test, for its part, highlights a break in 1975-76 in the different series of rainfall of this season. The deficits generated by this rupture oscillate between -25.8% (Littoral) and -28.2% (West) (Figure 5). The analysis of decadal deviations shows that for this season, rainfall never again reached normal after the 1970s in these four (4) regions (Table 3).



**Figure 4.** Results of annual and seasonal rainfall for the regions of the West side of the country (Littoral (7), West (6), South West (5) and North-West (4)), using Mann-Kendall's test. There is a significant trend in the series when the calculated p-values are below the significance level ( $\alpha = 0.05$ ). Significant trends are indicated by values in bold.



**Figure 5.** Temporal variation of annual and seasonal rainfall for the regions of the West side of the country (Littoral, West, North West, and South West), according to the Pettitt test. The rupture years are indicated by the vertical dashed lines. On their right sides, appear the corresponding rates of change. The ruptures in black are those which are identical to several regions.

### 3.1.3. In the southern part

In the southern part of Cameroon, the annual rainfall in the Center region is the only one for which the observed drop is statistically significant according to the Mann-Kendall test (Figure 6). For its part, the Pettitt test highlights in the case of the Center and East regions a break in 1975-76. The deficits observed following this rupture are respectively -5.9% and -4.3% (Figure 7). For the South region, the break highlighted by the Pettitt test occurred in 1971-72 (-5.6%) (Figure 7). The decadal deviations show a general drop in annual rainfall during the 1970s. However, a slight increase was noted during the 2000s (Center and South) or 2010s (East) (Table 4).

Regarding the rainy seasons, in the case of spring, the East region is the only one to have recorded a significant drop according to the Mann-Kendall test (Figure 6). The Pettitt test identified in the series of spring rainfall of the Center and East regions a break in 1975-75. The deficits caused by these ruptures are -7.9% and -6.2% (Figure 7). According to the Mann-Kendall test, the autumn rainfall statistically decreases in the three (3) regions of the southern part of Cameroon (Figure 6). The Pettitt test identified in the series of the Center and East regions a break in 1978-79. The decreases recorded following the latter are -5.7% and -5.2% (Figure 7). The rupture identified in the series of autumn rainfall in the southern region occurred in 1972-73 (-6.4%) (Figure 7). The decadal deviations of the rainy season's rainfall are similar to those highlighted for the annual rainfall (Table 4).


In the case of the dry seasons (summer and winter), the rainfall evolved in opposite trends. According to the Mann-Kendall test, summer rainfall increases. This increase is only significant for the South region (Figure 6). The Pettitt test puts in the series of summer rainfall of the Center and South regions breaks respectively in 1964-65 (+17.6%) and 1982-

---


83 (+22.1%) (Figure 7). Decadal deviations reveal an increase in summer rainfall during the 1960s (Center) and 1980s (East and South), then a general decrease during the 2010s (Table 4). Overall, winter rainfall in the southern part regions of Cameroon decreases non-significantly according to the Mann-Kendall test (Figure 6). Pettitt's test highlights a break in their series in 1975-76. The deficits recorded are between -20.3% (South) and -30% (Center) (Figure 7). From the 1980s, winter rainfall in these regions experienced a decline that was only interrupted during the 2000s (Table 4).

This study had different results from those obtained by previous studies in the regions of the southern part of Cameroon. The old studies indeed maintain that there is no break in their annual rainfall series [17,19], which is not the case with the present study, which identified downward breaks during the 1970s. The difference in results could arise from the difference in the rainfall sources used. This study is based on CRU data, however, the previous ones use other data sources (stationary, TRMM (Tropical Rainfall Measuring Mission), SIEREM (System of Environmental Information on Water Resources and their Modeling)).

Some studies in Central Africa have highlighted an evolution in opposite trends of the rainfall of the winter and summer dry seasons between the decades 1960/1980 and 1990 [17,19]. This is also the case in this study. However, this study reveals a reversal of these trends once known from the 2000s, which has only been demonstrated so far by very few studies [20,30].



**Figure 6.** Results of annual and seasonal rainfall for the regions of the South side of the country (South (10), East (9) and Center (8)), using Mann–Kendall's test. There is a significant trend in the series when the calculated p-values are below the significance level ( $\alpha = 0.05$ ). Significant trends are indicated by values in bold.



**Figure 7.** Temporal variation of annual and seasonal rainfall for the regions of the South side of the country (South, Center and East), according to the Pettitt test. The rupture years are indicated by the vertical dashed lines. On their right sides, appear the corresponding rates of change. The ruptures in black are those which are identical to several regions.

**Table 4.** Deviations (%) of the decadal annual and seasonal averages of rainfall compared to their interannual means for the regions of the South side of the country (Center, East and South).


Decades	Annual			Spring			Summer			Autumn			Winter		
	Center	East	South												
1950	3.9	2.8	4	7.6	6.4	7	-9.1	-14	-21	6.5	5.4	6.4	19.2	15.3	13.3
1960	4.3	3.4	3.8	3.6	3.8	4.8	3.7	0.4	-3.8	4.3	2.9	1.9	16.3	13.4	17.4
1970	0.9	-1.3	-1.6	-0.9	-1.8	-1.3	4.4	-3.7	-4.8	-0.4	-0.8	-2.2	12.9	5.6	3.4
1980	-3.5	-2.3	-3.6	-5.7	-1.7	-4.9	5.6	5.8	9.3	-4.6	-3.9	-2.4	-26	-18	-20
1990	-2.2	-0.7	-2.6	-4	-3.2	-5.3	4.6	11	13.6	-3	-1.5	-2.7	-19	-13	-12
2000	0.2	-1.8	3.2	-0.4	-4	1.3	0.2	2	5.8	0.8	-2.2	4.5	1.2	3.2	2.5
2010	-3.6	-0.3	-3.3	-0.4	1	-1.2	-8.7	-1	-0.4	-3.6	0	-5.7	-6.6	-6.9	-5.5


### 3.2. Spatial evolution of annual and seasonal rainfall

The fact that the 1970s was identified as a pivotal period in the evolution of rainfall in Cameroon led us to verify whether the spatial distribution of rainfall was also modified after this decade. This is the reason why we chose to study the spatial distribution of rainfall in the regions investigated during the decades 1950-1970 and 1980-2010.

#### 3.2.1. In the northern part

Spatially, the rainfall decreased towards the South in the northern part of Cameroon from the first (decades 1950-1970) to the second defined period (decades 1980-2010). (Figure 8). There is indeed a shift of the isohyets (1000 mm and 1500 mm, for example) towards the South. In addition, we note that the lower limit of the weakest rainfall class (< 1000 mm) progresses towards the South, as does the upper limit of the biggest rainfall class (> 1500 mm) (Figure 8). The seasonal rainfall (wet and dry) follows the same spatial evolution as the annual rainfall. We also note in both cases a progression of the isohyets towards the South (Figure 8).







**Figure 8.** Spatial distribution of mean rainfall for the regions of the North side of the country (Adamawa, North and Far North) at seasonal and annual time scales, between the decades 1950–1970 and 1980–2010.

### 3.2.2. In the western part

In the western part of Cameroon, the annual and seasonal rainfall decreased towards the South-West during the second defined period (decades 1980–2010) (Figure 9). It is perceptible through the displacement of the lower limits of the smallest rainfall classes (< 2000 mm for annual rainfall; < 1950 mm for wet season rainfall, and < 60 mm for dry season rainfall) and the upper limits of the heaviest rain classes in this direction (South-West) (Figure 9).







**Figure 9.** Spatial distribution of mean rainfall for the regions of the West side of the country (Littoral, North West, West and South West) at seasonal and annual time scales, between the decades 1950–1970 and 1980–2010.

### 3.2.3. In the southern part

In the southern part of Cameroon, the annual rainfall and those of the two rainy seasons decreased towards the West after the 1970s (Figure 10). The isohyets and the limits of the rainfall classes progress towards the West (Figure 10). In the case of dry seasons, those of the Summer dry season increase towards the South, while those of its winter counterpart decrease in the same direction, if we stick to the movements of the isohyets and the limits of their rainfall classes (Figure 10). Ebodé et al. [20] obtained similar results in their study in the Nyong basin (South Cameroon).





**Figure 10.** Spatial distribution of mean rainfall for the regions of the South side of the country (Center, East and South) at seasonal and annual time scales, between the decades 1950–1970 and 1980–2010.

#### 4. Conclusion

This study had as objective to analyze over the recent period (1950–2019), the rainfall Spatio-temporal variability in Cameroon. Strong analytical methods (the Pettitt and Mann–Kendall tests) were used. The results show that, temporally, the different regions of Cameroon recorded significant drops in annual rainfall that Pettitt's test generally places around the 1970s. The decreases observed for the northern part of Cameroon regions are between –5.4% (Adamaoua) and –7.4% (Far North). Those of the western part of Cameroon regions oscillate between –7.5% (South-West) and –12.5% (West). The southern part of Cameroon regions recorded decreases varying between –4.3% (East) and –5.9% (Center). On spatial plan, the divisions of the northern, western and southern parts of Cameroon respectively recorded after the 1970s (a pivotal period in the evolution of precipitation on temporal plan), a precipitation decrease towards the South, the South-West and the West. The CRU data used in this study certainly cover the whole country and are available over a long period (1901–2019), but they are only available in monthly time steps and in the form of grids. It does not allow us to go further in the precipitation study by addressing very detailed aspects, such as the evolution of the number of rainy days (Ray and Goel 2021). We recommend regular observation of rainfall in different parts of the country.

**Author Contributions:** Conceptualization: V.B.E.; methodology: V.B.E.; formal analysis: V.B.E.; investigation: V.B.E.; data curation V.B.E.; writing—original draft preparation: V.B.E.

**Funding:** This research received no external funding.

**Data Availability Statement:** All relevant data are included in the paper or its Supplementary Information.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Arati, P.; Riddhidipa, B.; Chowdary V. M.; Dibyendu, D.; Sreedhar, U.; Sankar, R. Trend analysis of time series rainfall data using robust statistics. *Journal of Water and Climate Change* **2017**, *08*, 4.
2. Han, H.; Hou, J.; Jiang, R.; Gong, J.; Bai, G.; Kang, Y.; Qi, W.; Li, Y.; Li, B. Spatial and temporal variation of precipitation characteristics in the semiarid region of Xi'an, northwest China. *Journal of Water and Climate Change* **2021**, *12*, 6.
3. Aher, M.; Yadav, S. M. Assessment of rainfall trend and variability of semi-arid regions of Upper and Middle Godavari basin, India. *Journal of Water and Climate Change* **2021**, *12*, 8. doi: 10.2166/wcc.2021.044.
4. Ray, L.; Goel, N. Spatio-temporal change in rainfall over five different climatic regions of India. *Journal of Water and Climate Change* **2021**, *12*, 7. doi: 10.2166/wcc.2021.052.
5. International Water Management Institute East Africa, 2014 (<http://eastafrika.iwmi.cgiar.org/>). Retrieved 10 Feb 2022.
6. Ogwang, B. A.; Chen, H.; Tan, G.; Ongoma, V.; Ntwali, D. Diagnosis of East African climate and the circulation mechanisms associated with extreme wet and dry events: a study based on RegCM4. *Arab J Geosci* **2015**, *8*, 10255–10265.
7. Camberlin, P. Nile basin climates. In: Dumont HJ (ed) The Nile: origin, environments, limnology and human use, monographiae biologicae, 2009, vol 89. Springer, Dordrecht, pp 307–333.
8. Jury, M. R. Climate trends in southern Africa. *S Afr J Sci* **2013**, *109*, 1/2, 1–11.
9. Diem, J.E.; Hill, T.C.; Milligan, R.A. Diverse multi-decadal changes in streamflow within a rapidly urbanizing region. *J. Hydrol.* **2018**, *556*, 61–71, doi:10.1016/j.jhydrol.2017.10.026.
10. Tierney, J. E.; Smerdon, J. E.; Anchukaitis, K. J.; Seager, R. Multidecadal variability in East African hydroclimate controlled by the Indian Ocean. *Nature* **2013**, *493*, 389–392.
11. Bigot, S.; Moron, V.; Melice, J.L.; Servat, E.; Paturel, J.E. Fluctuations pluviométriques et analyse fréquentielle de la pluviosité en Afrique centrale. In Water Ressources Variability in Africa during the XXth Century, Abidjan, Côte d'ivoire, November 1998; Servat, E., Hughes, D., Fritsch, J.M., Hulme, M., Eds.; IHAS Press, Wallingford, UK, 1998; pp.71–78.
12. Ebodé, V.B. Dzana, J.G.; Nkiaka, E.; Nka, N.B.; Braun, J.J.; Riotte, J. Effects of climate and anthropogenic changes on current and future variability in flows in the So'o River Basin (South of Cameroon). *Hydrology Research* **2022b** (Accepted for publication)
13. Cisse, M.T.; Sambou, S.; Dieme, Y.; Diatta, C.; Bop, M. Analyse des écoulements dans le bassin du fleuve Sénégal de 1960 à 2008. *Revue Sci. Eau* **2014**, *27*, 167–187. <https://doi.org/10.7202/102556ar>.
14. Nka, N.B.; Oudin, L.; Karambiri, H.; Paturel, J.E.; Ribstein, P. Trends in floods in West Africa: Analysis based on 11 catchments in the region. *Hydrol. Earth Syst. Sci.* **2015**, *19*, 4707–4719. <https://doi.org/10.5194/hess-19-4707-2015>.
15. Descroix, L.; Sané, Y.; Thior, M.; Manga, S.-P.; Ba, B.D.; Mingou, J.; Mendy, V.; Coly, S.; Dièye, A.; Badiane, A.; Senghor, M.-J.; Diedhiou, A.-B.; Sow, D.; Bouaita, Y.; Soumaré, S.; Diop, A.; Faty, B.; Sow, B.A.; Machu, E.; Montoroi, J.-P.; Andrieu, J.; Vandervaere, J.-P. Inverse Estuaries in West Africa: Evidence of the Rainfall Recovery? *Water* **2020**, *12*, 647. <https://doi.org/10.3390/w12030647>.
16. Bodian, A.; Diop, L.; Panthou, G.; Dacosta, H.; Deme, A.; Dezetter, A.; Ndiaye, P.M.; Diouf, I.; Vischel, T. Recent Trend in Hydroclimatic Conditions in the Senegal River Basin. *Water* **2020**, *12*, 436. <https://doi.org/10.3390/w12020436>.
17. Liénou, G.; Mahé, G.; Paturel, J.E.; Servat, E.; Sighomnou, D.; Ekodeck, G.E.; Dezetter, A.; Dieulin, C. Evolution des régimes hydrologiques en région équatoriale camerounaise: Un impact de la variabilité climatique en zone équatoriale? *Hydrol. Sci. J.* **2008**, *53*, 789–801. <https://doi.org/10.1623/hysj.53.4.789>.
18. Mahé, G.; Lienou, G.; Descroix, L.; Bamba, F.; Paturel, J.E.; Laraque, A.; Meddi, M.; Habaieb, H.; Adegea, O.; Dieulin, C.; et al. The rivers of Africa: Witness of climate change and the human impact on the environment. *Hydrol. Process.* **2013**, *27*, 2105–2114, doi:10.1002/hyp.9813.
19. Ebodé, V.B.; Mahé, G.; Dzana, J.G.; Amougou, J.A. Anthropization and Climate Change: Impact on the Discharges of Forest Watersheds in Central Africa. *Water* **2020**, *12*, 2718. <https://doi.org/10.3390/w12102718>.
20. Ebodé, V.B.; Braun, J.J.; Nnomo, B.N.; Mahé, G.; Nkiaka, E.; Riotte, J. Impact of Rainfall Variability and Land Use Change on River Discharge in South Cameroon. *Water* **2022a**, *14*, 941. <https://doi.org/10.3390/w14060941>.
21. Sighomnou, D. Analyse Et Redefinition Des Régimes Climatiques Et Hydrologiques Du Cameroun : Perspectives D'évolution Des Ressources En Eau. Ph.D. Thesis, Université de Yaoundé I, Yaoundé, Cameroun, 2004.
22. Amougou, J.A.; Ndam, N.J.R.; Djocgoue, P.F.; Bessoh, B.S. Variabilité climatique et régime hydrologique dans un milieu bioclimatique de transition: Cas du bassin fluvial de la Sanaga. *Afr. Sci.* **2015**, *11*.
23. Onyutha, C. Identification of sub-trends from hydro-meteorological series. *Stoch Environ Res Risk Assess* **2016**, *30*, 189–205. <https://doi.org/10.1007/s00477-015-1070-0>.

---

24. Letouzey, R. Notice de la Carte Phytogéographique du Cameroun au 1/500000; Institut de la carte internationale de la végétation : Toulouse, France, 1985.
25. Yue, S.; Pilon, P.; Cavadias, G. Power of the Man-Kendall and Spearman's rho tests for detecting monotonic trends in hydrological series. *J. Hyrdol.* **2002**, *259*, 254–271. [https://doi.org/10.1016/S0022-1694\(01\)00594-7](https://doi.org/10.1016/S0022-1694(01)00594-7).
26. Pettitt, A.N. A non-parametric approach to the change-point problem. *Appl. Stat.* **1979**, *28*, 126–135.
27. Ebodé, V.B. Impact of rainfall variability and land-use changes on river discharge in Sanaga catchment (forest–savannah transition zone in Central Africa). *Hydrology Research* **2022**, *53*, *7*, 1017–1030. <https://doi.org/10.2166/nh.2022.046>.
28. Descroix, L.; Guichard, F.; Grippa, M.; Lambert, L.A.; Panthou, G.; Mahé, G.; Gal, L.; Dardel, C.; Quantin, G.; Kergoat, L.; Bouaïta, Y.; Hiernaux, P.; Vischel, T.; Pellarin, T.; Faty, B.; Wilcox, C.; Malam Abdou, M.; Mamadou, I.; Vandervaere, J.-P.; Diongue-Niang, A.; Ndiaye, O.; Sané, Y.; Dacosta, H.; Gosset, M.; Cassé, C.; Sultan, B.; Barry, A.; Amogu, O.; Nka Nnomo, B.; Barry, A.; Paturel, J.-E. Evolution of Surface Hydrology in the Sahelo-Sudanian Strip: An Updated Review. *Water* **2018**, *10*, 748. <https://doi.org/10.3390/w10060748>.
29. Ebodé, V. B.; Mahé, G.; Amoussou, E. Impact de la variabilité climatique et de l'anthropisation sur les écoulements de la Bénoué (nord Cameroun), *Proc. IAHS* **2021a**, *384*, 261–267, <https://doi.org/10.5194/piahs-384-261-2021>, 2021.
30. Ebodé, V. B.; Mahé, G.; Amoussou, E. Changement climatique dans le bassin versant de l'Ogooué : évolution récente et impact sur les écoulements. *Proc. IAHS* **2021b**, *384*, 247–253. <https://doi.org/10.5194/piahs-384-247-2021>, 2021.