

Review

Structural determinant of β -amyloid formation: from trans-membrane protein dimerization to β -amyloid aggregates

Nicolas Papadopoulos^{1,2}, Nuria Suelves³, Florian Perrin⁴, Devkee M. Vadukul⁵, Céline Vrancx^{6,7}, Stefan N. Constantinescu^{1,2,8,9}, Pascal Kienlen-Campard^{3*}

¹ de Duve Institute, UCLouvain, Brussels, Belgium.

² Ludwig Institute for Cancer Research Brussels, Brussels, Belgium.

³ Aging and Dementia Research Group, Cellular and Molecular (CEMO) Division, Institute of Neuroscience, UCLouvain, Brussels, Belgium.

⁴ Department of Neurology, Massachusetts General Hospital, Memory Disorders Unit, Harvard Medical School, Charlestown, MA, USA.

⁵ Department of Chemistry, Molecular Sciences Research Hub, Imperial College London, London, United Kingdom.

⁶ Laboratory for Membrane Trafficking, VIB-Center for Brain and Disease Research, KU Leuven, Leuven, Belgium.

⁷ Department of Neurosciences, KU Leuven, Leuven, Belgium.

⁸ Walloon Excellence in Life Sciences and Biotechnology (WELBIO), Brussels, Belgium.

⁹ Ludwig Institute for Cancer Research, Nuffield Department of Medicine, Oxford University, Oxford, UK.

* Correspondence: pascal.kienlen-campard@uclouvain.be

Abstract: Most neurodegenerative diseases have the characteristics of protein folding disorders, i.e. they cause lesions to appear in vulnerable regions of the nervous system, corresponding to protein aggregates that progressively spread through the neuronal network as the symptoms progress. Alzheimer's disease is one of these diseases. It is characterized by two lesions, neurofibrillary tangles (NFTs) and senile plaques, formed essentially of amyloid peptides ($A\beta$). A combination of factors ranging from genetic mutations to age-related changes in the cellular context converge in this disease to accelerate $A\beta$ deposition. Over the last two decades, numerous studies have attempted to elucidate how structural determinants of its precursor (APP) modify $A\beta$ production, and to understand the processes leading to the formation of different $A\beta$ aggregates; e.g. fibrils and oligomers. The synthesis proposed in this review indicates that the same motifs can control APP function and $A\beta$ production essentially by regulating membrane dimerization, and subsequently $A\beta$ aggregation processes. The distinct properties of these motifs and the cellular context regulate the APP conformation to trigger the transition to the amyloid pathology. This concept can be transposed to the study of other protein folding disorders, to better decipher the patterns switching protein conformation from physiological to pathological and improve our understanding of these mechanisms that devastate neuronal functions.

Keywords: Alzheimer's disease; Amyloid Precursor Protein; Amyloid beta; APP-C99; dimerization; Orientations; Aggregation; Oligomerization

1. Introduction

Many neurodegenerative diseases are classified into the family of protein folding (or protein conformation) diseases, which are characterized by the abnormal accumulation of aberrantly folded proteins [1-3]. About 50 diseases are referred to as protein folding diseases, including Alzheimer's disease (AD) and the Tauopathies, Parkinson's disease (PD), Huntington's disease (HD), amyotrophic lateral sclerosis (ALS), and prion diseases such as Creutzfeldt-Jakob disease (CJD). For the vast majority if not all of them, abnormal protein folding and aggregation are considered to play a causative role [1-3]. There is a great variety of disease-causing proteins that can be present alone or in combination, and some of them can be found across several diseases [4].

The concept of protein folding disorder has its origins in the mid-19th century, when, in 1854, Rudolf Virchow coined the term amyloid, from the Latin word “amyrum” (starch), to describe a substance in the starchy brain bodies that exhibited a chemical reaction resembling that of cellulose [5]. Indeed, a great number of protein folding diseases are associated with the formation of highly organized fibrillar aggregates often described as amyloids [4]. This group of protein folding diseases is also known as amyloidoses. However, it is worth noting that some protein folding diseases are characterized by protein aggregates that are not amyloid in nature [6]. In most, if not all protein folding diseases, a change in three-dimensional folding (conformation) increases the tendency of a specific protein to bind to itself [1]. In particular, the disease-causing conformational changes in amyloidoses lead to an enrichment in β -sheet structures, which are associated with an increased propensity to form amyloids. These insoluble protein aggregates, regardless of the source of the amyloid protein/peptide, share a common fibrillar configuration with a unique structural core [7,8]. Each fibril consists of multiple twisted protofilaments which in turn show a so-called cross- β structure, with stacked β -sheet layers that run perpendicular to the long axis of the fibril. Additionally, the formation of amyloid fibrils is usually preceded by the generation of soluble oligomeric intermediates that are believed to be key drivers of cellular toxicity [9].

Proteins can acquire this fibrillogenic potential through pathological conformational changes, the underlying mechanisms of which are not yet fully understood. Several risk factors have been demonstrated to increase the propensity of these vulnerable proteins to self-associate. For example, gene mutations have been linked to the onset of some protein folding diseases (HD, inherited forms of AD and PD, Tauopathies like FTDP17) and it has been proven that some of them affect the conformational dynamics of the disease-causing proteins [10]. Post-translational modifications (e.g., hyperphosphorylation) and changes in environmental conditions, such as temperature, pressure, pH, and concentration of organic solvents, can also affect the aggregation propensity of vulnerable proteins [4]. An imbalance in protein production or clearing mechanisms can also lead to the accumulation of the amyloidogenic protein/peptide, indicating that misfolding processes are protein-concentration dependent [11]. Another key observation emerges from the study of these diseases. Although certain factors such as mutations are present in the individual at the outset, these only develop over time. Ageing is therefore a key factor that contributes to the context leading to protein misfolding. This is illustrated by the progressive build-up of cellular stressors (like reactive oxygen species or toxins) that might perturb the cell proteostasis and therefore increase the accumulation of aberrantly folded proteins [12].

Among neurodegenerative neuropathies, AD is the most prevalent and the most frequent form of amyloidosis. AD produces a progressive loss of memory and other cognitive functions, and it accounts for over 60 percent of dementia diagnoses. The clinical symptoms correlate with the gradual appearance of brain dysfunction and neurodegeneration, and the disease-defining pathological features consisting in extracellular senile plaques formed by the amyloid beta peptide (A β) and neurofibrillary tangles (NFT), which are intraneuronal aggregates primarily composed of hyperphosphorylated tau proteins. The gradual progression of protein aggregates allowed to set a staging that classifies the degree of pathology by correlating the density of lesions and clinical symptoms in specific brain regions [13,14]. From a molecular point of view, our knowledge of the mechanisms underpinning AD pathogeny is mostly rooted in the amyloid cascade hypothesis (ACH), which states that an early alteration in A β production, clearance or deposition is central in the aetiology of AD [15,16]. The debate around ACH has only heated up in recent years [17], mainly because of the failure of compounds targeting amyloid pathology to restore cognitive function when tested in clinical trials. Still, the correlation between A β deposition and the progression of AD is supported by undeniable arguments combining biochemical, anatomopathological and genetic evidence. AD causal mutations (responsible for autosomal-dominant AD or ADAD) are located in the genes encoding the Amyloid Precursor Protein (APP) or Presenilin 1 and 2 (PSEN1 and PSEN2, respectively) [18]. They trigger the accumulation of the A β peptide, which has a high propensity to aggregate,

eventually forming the extracellular senile plaques. Like in other protein folding diseases, these insoluble assemblies gradually accumulate, explaining the characteristic progressive nature of the neurodegenerative process and the commonly late onset of clinical symptoms. In this review, we set to provide a structural perspective on the determinants that drive the generation of pathogenic A β peptides.

2. The Amyloid Precursor Protein Family and A β production

The Amyloid Precursor Protein (APP) belongs to the APP protein family which includes APP, APLP1 and APLP2 in mammals [19-22], APPa and APPb in zebrafish [23] and APPL in drosophila [24]. The evolution of the *APP* gene family (for a review see [25]) indicate that the functions of APP, APLP1 and APLP2 have diverged to contribute distinctly to neuronal processes. After the discovery that A β was the main component of AD amyloid fibrils [26] and subsequent identification of *APP* [19,27] and *APLPs* [22,28,29] genes, extensive studies in knock-out models brought the assumption that *APP*, *APLP1* and *APLP2* are partially functionally redundant (for a review see [30]). The assumption that *APP* and *APLPs* have overlapping functions is also supported by the similarities in their expression pattern, biochemical processing and structure properties. All members of the APP family possess a large ectodomain composed of the E1 and E2 domains. Besides the E2 domain [31-33], the structures of the APP protein family have only partially been solved by conventional techniques. Yet, the recent advances in structure prediction algorithm led by AlphaFold [34] allow a reasonable approximation of the respective structures of APP, APLP1 and APLP2. **Figure 1** depicts the predicted structures of each segment of the APP and APLP proteins. Close analysis reveals that isolated fragments of the extracellular domains are almost identical in structure, with the inter-fragments predicted unstructured. The E1 domain is composed of two separate segments, corresponding to the Growth-Factor Like Domain (GFLD) and copper-binding domain (CuBD). GFLD is made of two β -sheet separated by an α -helix and the CuBD consists of a three-stranded antiparallel β -sheet [35]. The E2 domain is formed by two coiled-coil α -helices which are conserved in all three members of the APP family [33,36] (**Figure 1**). The Kunitz-type protease inhibitor (KPI) domain is specific to APLP2 and long isoforms of APP (APP₇₇₀ and APP₇₅₁) and not present in the neuronal isoform (APP₆₉₅). It is composed of double-stranded β -sheets with a small α -helix [37]. Functionally, the extracellular domain of APP/APLPs binds components of the extracellular matrix such as heparin and collagen via the E1 and E2 domain [38,39]. Interaction with heparin plays a role in neurite outgrowth [40], APP/APLPs dimerization [36,39,41] and cell-cell adhesion [42]. The E1 and E2 domain are also reported to bind zinc and copper [43,44] via the E1 copper/zinc binding domain (Cu/Zn BD) (**Figure 1**) and less well-defined sites of the E2 domain. Binding of zinc and copper regulates APP and APLPs *cis* and *trans*-dimerization, possibly by regulating their interaction with heparin [45-48]. Interestingly, the major neuronal isoform of APP (APP₆₉₅) [49] does not contain the KPI domain whereas longer isoforms that contain the KPI are expressed in the periphery and to a significant level in platelets, in agreement with the reported role of APP KPI as an inhibitor of multiple coagulation factors [50].

The transmembrane (TM) domains of APP and APLPs are formed of a single α -helix and the intracellular domain is mostly unfolded with the exception of the C-terminus which is composed of two small α -helices (**Figure 1**). The similarity between APP/APLPs and Notch processing (see below) led some authors to hypothesize that the APP and APLPs [51] Intracellular C-terminal Domain (AICD) could possess signaling functions (see [52] and [53] for a review). Reporter gene assays based on Gal4-fusion constructs suggested that AICD could act as transcriptional regulator [54,55], but the controversy about the transcriptional activity of AICD and its target genes has only grown since then [56,57].






Figure 1. Structure of the APP proteins family

Structure of the APP and APLPs proteins predicted by AlphaFold 2.0 [34]. The structure of folded domains of each protein are shown as cartoon in red, yellow and blue for APP⁷⁷⁰, APLP1 and APLP2, respectively. Unstructured regions are not shown. The domains corresponding to each structure are detailed with amino acid numbering equivalent to the start and end of each subdomain. The YENPTY motif corresponds to the internalization motif conserved in all members of the APP family. HB: Heparin Binding (orange), CB: Collagen Binding (green), Cu/Zn BD (blue): Copper/zinc Binding Domain, GFLD: Growth factor-like domain.

2.1. Sequential processing of the APP and APLPs and production of β -amyloid peptides

The structural similarities between APP family members goes hand in hand with the overlapping functions of APP, APLP1 and APLP2. Yet, APP is the only member of the family giving rise to the formation of pathogenic A β aggregates. Members of the APP family are processed via two different pathways (Figure 2A). The most common pathway was coined non-amyloidogenic or anti-amyloidogenic due to its inability to produce amyloidogenic A β fragments [58,59]. In the non-amyloidogenic pathway, APP processing begins with the shedding of the ectodomain by a membrane-bound α -secretase of the ADAM family [60]. The main α -secretase for APP is ADAM10 [61] while studies suggest that TACE is the preferred α -secretase of APLP2 [62]. The cleavage by the α -secretase is sequence independent and occurs at a distance of 12-13 residues from the TM domain [63] to release the ectodomain (sAPP α in the case of APP) and the membrane-anchored C83 fragment. In the amyloidogenic pathway, a β -secretase (BACE1) always starts the cleavage a few amino acids upstream of the α -cleavage site to release sAPP β . BACE1 is an aspartyl protease which possesses some level of sequence specificity although large discrepancies do exist between cleavages sites (for a review see [64]). This low level of sequence specificity results in two possible β -cleavage sites of APP. The one after Glu₆₈₂ produces C89 but

only the cleavage after Asp₆₇₂ can generate the C99 that enables production of amyloidogenic A β peptides [65]. Mutations responsible for inherited AD are reported to promote the cleavage at Asp₆₇₂ (i.e. the APP Swedish or APP KM670/671NL mutation) by providing a more favorable environment for β -cleavage at Asp₆₇₂ [66,67]. The two other members of the APP family, APLP1 and APLP2 are also processed by the β -secretase to produce C-terminal fragment (CTF) of 100 and 104 residues, respectively (**Figure 2B**). In both the amyloidogenic and non-amyloidogenic pathways, CTFs resulting from α - or β -cleavage are further processed by a γ -secretase complex. After initial ϵ -cleavage at position 48-49 (A β numbering in APP) which releases AICD and A β ₄₈₋₄₉ [68], the γ -secretase performs intramembrane successive cleavage of three (or four) residues at a time to produce C-terminal A β fragments ranging from 36 to 43 amino acids [69,70], of which A β ₄₂ and A β ₄₃ have strongest oligomerization properties. This highlights the observation that the C-terminus of A β peptides is key for aggregation. Similarly, APLP1 and APLP2 are processed by γ -secretase complex but do not lead to the production of peptides with amyloidogenic properties [71,72], illustrating that A β -like oligomerization is a property specific to the APP sequence (i.e transmembrane (TM) and juxtamembrane (JM) sequences) present in A β peptides. These sequences or motifs provide A β with fibrillogenic properties once it has been released upon γ -cleavage of C99. In addition to a specific C-terminal sequence, the pathologic properties of A β are also associated with the LVFF motif [73] (see below) present in the A β central region, which forms a β -hairpin and also plays a regulatory role in γ -secretase processing [74]. This motif is present only in APP and is not restricted to the C99 produced by β -secretase cleavage (**Figure 2B**), but also present in C83. To date, the role of the N-terminus of A β sequence has been less investigated but it may impact the stabilization of A β aggregates [75]. In this respect, it is important to note that human and murine A β only differ by 3 amino acids residues in the N-terminal regions, and that murine A β is not amyloidogenic. Overall, growing evidence indicates that some peculiar motifs, present in the human A β sequence (and not in related peptides generated by APLP1/APLP2 processing) could play a role in amyloid pathology both (i) by regulating A β production and (ii) by promoting or preventing the assembly of A β building blocks into amyloid fibrils.

A**B****Figure 2. Processing of the APP protein family**

A. Illustration of APP processing. The sites of α , β and β' cleavage are shown with amino acids corresponding to cleavage position. Upon cleavage by α - or β -secretase, fragments called sAPP α , sAPP β and sAPP β' are released and secreted and C83, C89 or C99 remain embedded in the membrane, respectively. C99 contains the complete A β sequence while C89 and C83 have truncated (11-X and 17-X, respectively) A β sequences. The A β 17-X is also called the P3 fragment. The first cleavage by γ -secretase occurs at the ϵ site to release AICD prior to intramembrane proteolytic cleavage of 3 or 4 amino acids to produce amyloid β (A β) peptides from 36 to 43 amino acids. **B.** Sequence alignment of APP C99, C89, C83 and APLP1/2 β -CTF. The alignment illustrates sequence differences between the members of the APP family. The LVFF motif (red) is present only in APP while the C-terminal fragment is similar

between APP and APLP2. The cleavage sites of γ -secretase are mostly conserved between all members of the APP family. The bottom chart illustrates the conservation of amino acid per position based on their properties with values ranging from 0 (no conservation) to 9 with * representing perfect conservation.

3. Transmembrane interactions and amyloidogenic processing

The finding that APP forms dimers by homo- and hetero-interactions with other APP family members [42,76] was instrumental to the hypothesis that APP dimerization was a regulatory mechanism for its processing and function. Dimerization of APP was described early in the field and was first associated to its binding to heparin and collagen via the ectodomain [77]. Later, it was discovered that APP dimerization promoted the production of A β [78]. Homo-dimerization of APP was initially described between different segments of the ectodomain and for its role in cell-cell adhesion [42,79].

Transmembrane (TM) and juxtamembrane (JM) motifs involved in protein-protein interactions have attracted particular interest. Regarding the remarkable conservation of APP and APLPs TM sequences, these motifs might be involved in dimerization between all the members of the family. In addition to their involvement in dimerization, these motifs present in APP are found precisely in the A β sequence and in the vicinity of the APP cleavage sites releasing A β . Such motifs are GxxxG and GxxxG-like motifs, also known as Glycine zippers, and were described as key regulators of APP amyloidogenic processing [80,81]. GxxxG motifs were initially identified in the glycophorin A TM region [82-85]. The groove formed by an amino acid without lateral chain (Gly) in an α -helix context allows close apposition of TM helices and packing into dimers by non-covalent interactions formed by amino acids surrounding the GxxxG sequence. Mutagenesis and structural analyses have revealed numerous examples in which the interaction between TM helices of single-pass membrane proteins is dependent on a GxxxG or (small)xxx(small) motif, where (small) designates amino acid residues with short side chains such as alanine. The interaction strength of motif-containing helices depends strongly on the sequence context and membrane properties. Several GxxxG-containing TM domains can interact via interfaces involving residues (hydrophobic, polar, aromatic) that are not organized in recognizable motifs. Importantly, in multi-pass membrane proteins, GxxxG motifs can be involved in protein folding, and not just oligomerization [86,87]. GxxxG and GxxxG-like motifs are enriched in protein TM regions, and strikingly, APP contains 3 in-register GxxxG motifs in its TM/JM region (Figure 3A), with an ADAD mutation (Flemish or A692G) adding a fourth motif in the JM sequence [88]. Such an occurrence of 3 GxxxG motifs is very rare (Figure 3B) and observed only in 17 human type I TM proteins (22 in *mus musculus*), and only 1 protein contains 4 GxxxG motifs in the region comprising the TM and 6 surrounding residues on each side. Intriguingly, the prion protein involved in prion disease by the misfolding of the endogenously expressed prion protein (PrPC) into an abnormal isoform (PrPSc) that has infectious properties, also contains a string of 3 in-register GxxxG motifs in its hydrophobic region [89]. However, main differences are that PrPC is a membrane-anchored and not a TM protein, and that the hydrophobic regions -known as the hydrophobic tract- lie outside the α -helical C-terminal domain, therefore possibly in another conformation than the APP TM region. In the case of APP, GxxxG motifs are indeed involved in TM interactions once the bulky ectodomain has been removed by α - or β -secretase shedding (Figure 2A) and play key regulatory functions in APP the dimerization and processing of APP CTFs [79,90].

An important advance came from solid state Nuclear Magnetic Resonance (NMR) studies and other biophysical approaches applied to the JM-TM domains of APP. Using peptides resuspended in membrane-like bilayers, like dimyristoyl-phosphocholine (DMPC) : dimyristoylphosphoglycerol (DMPG) bilayers, structural data indicated that the LVFF motif preceding the first GxxxG motif forms a β -sheet structure, that the GxxxG are likely helical and that the helicity of the first GxxxG motif is enhanced by the A21G motif and by cholesterol [91,92]. Earlier NMR studies in

detergent or detergent-lipid mixtures [93,94] showed a different picture for LVFF and GxxxG motifs that did not fit with processing. Careful titration of detergent by FTIR spectroscopy showed that detergent induced the helical structure of LVFF rather than the natural β -sheet. Interestingly, cholesterol appears to bind to the aromatic residues of the LVFF motif and influence the downstream structure [92]. Last but not least, introducing the Flemish mutation (A692G or A21G in A β numbering, see above) and adding cholesterol to the experimental context exerted additive effects on the helicity of the first GxxxG motif and on amyloidogenic processing [95]. Thus, APP processing at γ -sites might be controlled by specific orientations and interactions of its TM domain, these processes (orientation and intercation) being controlled by the cellular context and in particular by the lipid composition of the membrane leaflets.




Figure 3. The role of C99 dimerization in Amyloid β generation

A. Illustration of the role of GxxxG motifs in APP C99 dimerization. The figure illustrates the two dimerization interfaces described in [96]. The conformation shown left contains the $^{25}\text{GXXXG}^{29}$ motif in the interface and was associated with production of SDS-resistant A β hexameric species. The conformation shown right interface contains the $^{33}\text{GXXXG}^{37}$ motif in the interface and was linked to generation of AICD signaling properties and promotion of the A β_{42} processing line [96]. Glycines are shown in red, with the one in the interface shown as bubbles. **B.** Occurrence of GxxxG motif in the TM-JM segments of type I TM proteins in the homo sapiens (black) and mus musculus (blue) proteome. Only one human protein contains 4 GXXXG in the TM-JM segments (not visible in the graph).

3.1. Dimeric conformation of C99 regulates γ -secretase processing

Interest for the dimerization of APP C99 occurred later with the observation that a large proportion of FAD mutations occurred in the TM-JM domain where TM interaction motifs are located. The observation that GxxxG motifs and dimerization of APP regulated A β production provided strong arguments for an influence of TM dimerization in its processing by the γ -secretase. The role of APP/C99 dimerization in the regulation of γ -secretase processing was extensively investigated [81,97-103] but led to two contradicting observations. On one hand, dimerization was found to increase generation of pathogenic A β . For example, Scheuermann and colleagues designed stable APP homodimers by introducing the K699C mutation (K624C in APP₆₉₅ numbering) at the extracellular JM-TM section ahead of γ -secretase cleavage sites and showed that dimerization dramatically increased A β production [78]. In line with these findings, compounds that reduced A β_{42} generation were shown to act by interfering with APP TM dimerization [104]. On the

other hand, different lines of evidence suggested that forced APP TM dimerization was not compatible with γ -secretase processing. Using a system in which dimerization is imposed from the intracellular domain, Eggert and colleagues observed that controlled dimerization resulted in decreased A β generation [105]. But more recently, the acquisition of the cryo-EM structure of γ -secretase in complex with C83 hinted that processing of C-terminal fragments (CTF) could indeed not occur in their dimeric form [106], in line with our own observations using a cell-free γ -secretase processing assay [96]. A first answer to these apparently contradicting observations was provided by our finding that GxxxG motifs were critical in regulating the orientations of TM dimerization and that a specific interface was required for amyloidogenic processing [90]. Using a system of fusion proteins that force dimerization of TM helices in all possible orientations, we demonstrated that precise dimeric orientations of C99 controlled γ -secretase processing by regulating the initial ε -cleavage and therefore influencing AICD-dependent signaling and generation of A β ₄₂ [96]. Noteworthy, the dimeric orientation resulting in enhanced A β ₄₂ production and AICD-dependent signaling contained GxxxG motifs in the interface, but unfolding is mandatory for γ -processing since covalently bound APP CTFs are not further processed [96].

3.2. APP dimeric conformation controls its intracellular localization and A β generation

In addition to its role in the regulation of processing, multiple lines of evidence indicate that dimerization of C99 is both a regulator and a consequence of its subcellular localization. The impact of APP dimerization on its subcellular localization was only recently investigated. A link between dimerization and subcellular localization was first suggested by Ben Khalifa and colleagues who observed that APP dimers were mostly present in the secretory pathway [79]. Using a model of forced dimerization of full-length APP, Eggert and collaborators analyzed in more detail the role of APP dimerization on subcellular localization and observed that APP dimerization resulted in increased localization in the endoplasmic reticulum (ER) and in endosomes, where the generation of pathogenic A β is believed to occur, by modulating its interaction with LRP1 and SorLA [107] (reviewed in [108]). The role of dimerization in intracellular trafficking was extended by Perrin and colleagues who observed that different dimeric orientations of C99 resulted in distinct subcellular localization in primary neurons and that the A β ₄₂ processing line was favored by a dimeric conformation that resulted in decreased cell surface and increased intracellular localization [96].

To better understand the impact of subcellular localization on amyloidogenic processing (see Figure 4), one must keep in mind that γ -secretase is composed of four subunits, namely Nicastrin, Aph1, Pen-2 and Presenilin 1/2 (PS1/2, encoded by the *PSEN1* and *PSEN2* genes, respectively) [109]. The catalytic core of γ -secretase is composed by either PS1 or PS2 which harbor two aspartate catalytic residues (e.g. Asp287 and Asp387 for PS1) [110,111]. The fact that mutations in PSs are the most common cause of FAD highlights their critical role in the pathology (for a review see [112]). The link between subcellular localization of APP and PSs was recently investigated by Sannerud and colleagues [113]. Using a wide variety of approaches, they elegantly demonstrated that PS1 and PS2 have distinct subcellular localizations which impact C99 processing. More specifically, they observed that PS2 was addressed to late endosomes/lysosomes by its N-terminal dileucine sorting motif E₁₆RTSLM₂₁ via interaction with AP-1 while PS1 was more ubiquitously expressed, notably at the plasma membrane. As suggested before [114], secreted A β peptides were increased in the absence of PS2 but intracellular A β and A β ₄₂/A β ₄₀ ratio was higher in the absence of PS1. They further observed that cleavage of C99 in late endosomes was associated with the formation of aggregation-prone A β peptides that were less efficiently secreted. Remarkably, these results correlated with those obtained in two of our recent studies. In Perrin *et al.*, we found that the dimeric orientation of C99 that favored the A β ₄₂ processing line and localized in intracellular compartments was more efficiently processed by PS2 [96]. In Vranckx *et al.*, we reported that specific hexameric-like A β ₄₂ assemblies were majoritily produced by PS2 in vesicular compartments [115]. The fact that PS2 favors A β ₄₂ processing line in late endosomes must also be put in perspective with the observation that BACE1, the limiting enzyme of APP amyloidogenic processing, is predominantly found in endosomal compartments [116-119]. Together, these observations provide strong arguments

for a crosstalk between C99 dimerization, subcellular localization and generation of pathogenic amyloid β . It is also interesting to mention that TM interaction motifs discussed above are not restricted to APP but are also found in PSEN1 and PSEN2 [120], and in the Aph-1 subunit of the γ -secretase [121], and that the hydrophilic loop of PSEN1 together with APP GxxxG TM motif regulate γ -secretase function in generating pathogenic A β peptides [122]. These observations indicate that the role of TM interaction in amyloid pathology might not be restricted to the substrate (APP) but might act all along the processing line controlling the orientation of APP dimers, the docking and fitting in the γ -secretase complex and eventually the formation of pathogenic A β species.




Figure 4. The role of C99 dimerization in A β generation

Illustration of the role of C99 dimerization, cholesterol concentration and PSs localization on APP/C99 processing and A β generation. Initial cleavage by α - or β -secretase occurs at the plasma membrane or in endosomes, respectively. In addition, low cholesterol concentration favors α -cleavage while high cholesterol content is associated with β -cleavage. After initial cleavage, γ -secretase can occur either at the cell surface (preferably by PS1) or in endosomes (preferably by PS). One hypothesis is that medium cholesterol concentration favors dimerization in one of the two dimeric orientations with either $^{25}\text{GxxxG}^{29}$ or $^{33}\text{GxxxG}^{37}$ motif in the interface. Such dimeric concentration targets C99 either to endosomes ($^{33}\text{GxxxG}^{37}$ interface) or to the cell surface ($^{25}\text{GxxxG}^{29}$ interface). When concentration of cholesterol increases (like in lipid rafts), dimerization is destabilized to allow processing by PS1 or PS2-dependent γ -secretase.

3.3. The role of lipids in C99 dimerization and amyloidogenic processing

Cell membranes are a buoyant environment and are largely heterogeneous in their composition. The link between AD and the lipid composition of membranes had already been suggested by Alois Alzheimer who observed “lipoid granules and “adipose inclusion” as a third hallmark of Alzheimer’s disease [123,124]. A large body of evidence has since confirmed that lipid composition was indeed a critical factor influencing AD pathogenesis (reviewed in [125,126]). Cholesterol was early found to be a major risk factor of Late Onset Alzheimer’s Disease (LOAD) [127-129]. This argument is reinforced by the identification of the allele $\epsilon 4$ of ApoE, a major lipoprotein involved in cholesterol transport, as the most important risk factor of non-familial AD [130-132]. Early studies also indicated that low cholesterol promotes α -cleavage and non-amyloidogenic processing [133,134] whereas β -cleavage and amyloidogenic processing is favored by high cholesterol content [135] found in lipid rafts [136]. The role of cholesterol in C99 dimerization remains largely controversial. Different lines of evidence support the observation that C99 directly interacts with cholesterol [93,137,138]. Nuclear Magnetic Resonance (NMR) analyses revealed that interaction between cholesterol and C99 involved essentially the N-terminal loop of C99 and residues Glu693 and Asn698. Interestingly, glycines of the first GxxxG motif also seemed to directly bind cholesterol [138]. In line with the observation that GxxxG motifs interact with cholesterol, Song and colleagues suggested that cholesterol binding to C99 competed with homodimerization [139] and others showed that lowering cholesterol inhibited A β production by promoting APP dimerization [140]. These results are intriguing since a large body of evidence linked C99 dimerization to amyloid β production. Yet, other investigations tend to show that rather than preventing dimerization, cholesterol modulates the conformation of C99 dimers. Studies from Tang and colleagues indicated that the $^{25}\text{GxxxG}^{29}$ motif was in fact stabilized by cholesterol [92] and others suggested that the membrane micro-environment and cholesterol, rather than competing with homodimerization of C99, had a role in the regulation of C99 dimeric conformation [141-143]. These studies differ notably by the composition of the membrane environment used, not only in percentage of cholesterol but also in other lipids that may impact C99 dimerization. One plausible scenario is that a moderately increased level of cholesterol promotes C99 dimerization in a particular conformation related to increase amyloidogenic processing. This conformation may target C99 dimers to lipid raft where the very high cholesterol content induces destabilization of C99 dimers thereby allowing processing by the γ -secretase and generation of A β peptides (Figure 4).

An important question concerns the thickness of the membrane; directly related to lipid composition. A thicker membrane will embed more residues in the juxtamembrane region and change tilt and ionization of the residues bordering the TM domain. The same sequence can also fold as an α -helix, random coil or β -strands in different environments. The use of lipid formulations close to that of the cell membrane allowed the conclusion that the GxxxG motifs are helical [92], yet in less physiological environments they form other structures [94], which is relevant for post-cleavage events and formation of oligomers. Moreover, prediction by replica-exchange molecular dynamics simulations of wild-type (WT) and mutant APP dimer conformations in which Gly residues from GxxxG motifs were changed to Leu illustrated large conformational differences in a membrane context. Dimerization of the WT (GxxxG) is due to two hydrogen bonds between two APP fragments, whereas dimerization of the mutant (LxxxL) is due to hydrophobic interactions. In the mutant, each APP fragment is more tilted, and the γ -cleavage site is shifted toward the center of the membrane. This position produces a mismatch between the active site of γ -secretase and the γ -cleavage site of APP that might impair A β production [144].

3.4. Impact of familial AD mutations on APP TM dimerization: a link with A β production?

The fact that over 20 FAD mutations are located in APP TM domain illustrates the importance of this region for pathogenic A β generation. Yet, their influence on A β production through C99 dimerization remains quite elusive. Using synthetic peptides, Gorman and colleagues found that the V717G, V717F (Indiana) and T714I (Austrian) mutations all

decreased TM dimerization and concluded that TM dimerization was negatively associated with pathogenic A β generation [145]. Likewise, others concluded that FAD mutations in the $^{714}\text{TVIV}^{717}$ TM segments induced lower dimerization [146]. Conversely, So and collaborators analyzed dimerization of full-length APP FAD mutants with techniques including biomolecular fluorescence complementation (BiFC) and live cell crosslinking and found that FAD mutations did not significantly affect the overall level of dimerization [147]. These observations might suggest that APP FAD mutations do not act by increasing C99 dimerization levels but may rather induce modulation of C99 dimeric interface. Consistently, we observed that the APP Flemish mutation (A692G) promotes dimerization in a conformation that puts the GxxxG motif in the interface and increases A β_{42} production [95]. But strikingly, results obtained with split protein assays indicated that mutations of Gly residues present in APP GxxxG motifs did not affect the number of dimers formed, but rather generated specific A β assemblies [99]. This led to a new hypothesis. The TM interaction motifs (i.e. GxxxG or GxxxG-like) might indeed not only control dimerization and processing, but also the processes involved in A β assembly that leads to toxic aggregates. This hypothesis was reinforced by our most recent findings showing that forcing the $^{25}\text{GxxxG}^{29}$ interface in dimeric A β_{42} and A β_{43} assemblies resulted in the generation of SDS-resistant hexameric A β species while forcing dimerization in the $^{33}\text{GxxxG}^{37}$ interface did not allow generation of such oligomers, even in the context of highly amyloidogenic A β_{42} and A β_{43} [96].

3.5. JM/TM determinants drive aggregation steps (nucleation) leading to pathological A β seeds

When the soluble A β (sA β) peptides accumulate they tend to spontaneously self-aggregate, eventually leading to the formation of fibrillar amyloid lesions. Over the past decades, a growing body of evidence indicated that A β intermediates, in particular soluble oligomers, are the primary cause of synaptic dysfunction and A β toxicity observed in AD. Indeed, while A β fibrils are large, insoluble materials aggregating into plaques, A β oligomers are soluble and may easily spread throughout the brain. Many oligomeric structures seem to play an important role in A β assembly and to have deleterious effects that could explain its related toxicity. For instance, dimers and trimers of A β have been consistently linked to long-term potentiation impairments [148-150]. Understanding the process of A β aggregation is crucial for identifying assembly steps that may be targeted by disease-modifying drugs.

Unlike the Tau protein for instance, in which post-translational modifications (and especially phosphorylation) are tightly associated with abnormal folding and aggregation, few post-translational modifications (PTMs) are described on A β and their possible role in A β aggregation remains uncertain. The A β peptide has been demonstrated to undergo several types of posttranslational modification, such as pyroglutamylation, N-terminal truncation, oxidation, glycosylation, nitration, isomerization, racemization and phosphorylation [151,152]. However, whilst modified forms of A β are further investigated as potential markers of AD progression [153], such modifications are known to not be necessary for the induction of A β aggregation and subsequent deleterious effects.

One exception might be the phosphorylation of serine residue at position 8 and 26 (A β numbering) that has been reported to promote the formation of oligomeric A β assemblies or to stabilize them, respectively [154,155]. Thus, intrinsic determinants much more than PTMs are likely to be at hold for aggregation. This might be highly regulated both (i) by the nature of the C-terminal end and (ii) by the presence of peculiar motifs that promote the folding and packaging of A β into oligomers and fibrils. The length of the A β peptide is correlated to its aggregation properties, with the longest forms (A β_{42} and A β_{43}) being more hydrophobic and prone to assemble into oligomers and further elongate into fibrils. A faster rate of aggregation of A β_{42} in comparison to A β_{40} and a stable set of oligomers with a diameter distribution of \sim 7 to 9 nm was prevalently observed uniquely for A β_{42} even after fibril appearance [156]. A β assembly relies on a process called "nucleated polymerization", involving three distinct phases: (i) a nucleation phase -or lag phase- during which several unfolded or partially folded monomers of A β come together to form an oligomeric nucleus, (ii) an elongation phase -or growth phase- in which the nuclei rapidly grow by further addition of monomers, giving rise to prefibrillar

structures then ordered protofibrils, and (iii) a stationary phase -or equilibrium- where the concentration of monomers is low and constant, the assembly process reaches saturation and protofibrils assemble into mature fibrils (Figure 5). The nucleation phase relies on A β monomers undergoing conformational changes and misfolding, which renders it thermodynamically unfavorable, whereas the elongation phase is much more favorable and occurs rapidly. In primary nucleation, the initial formation of amyloidogenic nuclei occurs without contribution of pre-formed oligomers and constitutes the rate-limiting step. Amyloid fibril formation may also be seeded by the presence of preformed aggregates - oligomers, protofibrils or fibrils- [157]. As a consequence, a whole set of A β assemblies intermediate in A β fibril formation are present in the brain parenchyma. Some of them were shown to be readily formed in cells and having seeding properties when monomeric A β_{42} is available [115,158]

Peculiar motifs present in the A β sequence largely contribute to its complex aggregation properties. Strikingly, TM interaction motifs appear to play a key role not only in A β dimerization but also in aggregation. In particular, the GxxxG motifs described above, which acquire aggregation properties after the transition from the α -helical structure found in the TM context to the β -sheet structure present in A β peptides. Glycines from GxxxG allow the turn in the β -helix and form a rigid and groove structure in the β -sheet that allows stacking of amyloid fibrils [159]. It is tempting here to draw a parallel between GxxxG motifs involved in TM dimerization and their role in A β assembly by raising some key questions. Could A β stubs initially folded in an α -helical structure (corresponding to APP TM structure) assemble into dimers that would initiate the nucleation process and promote the conversion to larger A β nuclei (Figure 5)? Some recent work supports this exciting hypothesis. First, specific orientations of APP TM orientations engaging a GxxxG interface lead to the production of specific (hexameric-like) A β assemblies. Remarkably, forcing a specific α -helical orientation of A β_{42} and A β_{43} -like peptide dictates their ability to form oligomeric species [96]. Additionally, it has been recently reported that modular A β C-terminal segment mediates rapid, non-nucleated formation of α -helical oligomers, resulting high local concentration of tethered amyloidogenic segments within these α -oligomers facilitates transition to a β -oligomer population that ultimately generates mature amyloid [160]. Motifs located in the A β N-terminal segment are also involved in A β aggregation. This is partially sound, since the N-terminal region of A β is present in the JM regions of its precursor (APP), that shows poor conservation across the APP family members. It makes therefore sense that intrinsic aggregation properties of A β directly relate to its N-terminal/central region. The LVFF motif (see above) plays a critical role in the A β assembly process. This LVFF sequence was initially proposed as an α -helix region, that acts as a cholesterol sensor regulating APP processing [161,162]. Incorporation of cholesterol into model membranes enhances the structural changes induced by FAD mutations, suggesting a common link between familial mutations and the cellular environment. However, the LVFF motif was more recently shown to be predominantly β -sheet in membrane bilayers that, when disrupted by changing lipid composition, increase A β production. In this β -sheet conformation, it can match hydrophobic motifs in Tau repeats and accelerate their aggregation via polymorphic states, appearing thus as a major interactor in both types of AD protein folding disorders [163].




Figure 5. Steps of amyloid β aggregation

Illustration of putative nucleation steps of A β adapted from [164]. γ -secretase cleaves monomers of C99 (see above) but released A β peptides may rapidly re-form dimers centered on the GxxxG motifs. In the nucleation phase, dimer formation is the initial step leading to the formation of higher-order species (hexamers are represented) whose nucleation is mediated by A β C-terminus. These oligomers are usually forming β -sheets but may go through a transient step to form α -oligomers that rapidly switch to more stable to β -oligomers initiated by the LVFF motif. Elongation occurs by assembly of β -sheet oligomers into fibrils.

4. Conclusion and future directions

Alzheimer's disease is the first cause of dementia and global aging of the population is set to cause a dramatic rise in an already very high number of cases. Despite recent controversy, the amyloid cascade hypothesis remains the most widely accepted in the scientific community given the robustness of its genetic and biochemical arguments. Yet, clinical trials targeting amyloid plaques have all failed to provide the expected benefits. These failures might be rooted in the fact that most trials have focused on targeting amyloid deposits rather than circulating A β oligomers, which are more believed to be the toxic species associated the cognitive decline, although this hypothesis is also fiercely debated. In the end, the involvement of amyloid pathology in AD has been the subject of as much debate -if not more- as recent advances in knowledge.

Still, in recent years, important observations have advanced our understanding of the structural determinants that lead to the formation of toxic oligomers. The discovery that C99 dimerization and its dimeric orientations regulate its processing to generate pathogenic A β species was instrumental to understand how amyloidogenic processing occurs. Recent advances highlight how different factors modulate the conformation of C99 dimers that in turn regulates amyloidogenic processing. Amongst them, membrane composition and in particular cholesterol content seems to play a key role in the development of sporadic AD. The allele $\epsilon 4$ of ApoE, the main apolipoprotein involved in cholesterol homeostasis in the brain, is the principal risk factor for sporadic AD [130,165]. In addition, hypercholesterolemia was found to increase the production of A β oligomers in the brain of mouse models [166] and post-mortem analysis of the brain of AD patients demonstrated increased levels of cholesterol [167]. Despite this observation, the understanding of the role of membrane composition in the modulation of C99 dimeric conformations and generation of toxic amyloid species has been slowed and complicated by the heterogeneity of the methods used to recapitulate biological membranes.

Another critical question is whether the same motifs that regulate C99 processing and localization are also involved in the initial steps of A β oligomerization. Recent studies from two independent groups strongly suggest that early

nucleation of A β could occur in a transient helical conformation and that modulation of the dimeric orientation of helical A β -like peptides dictates the generation of A β oligomers [96,160]. This contrasts with the previous idea that A β oligomers are essentially formed of closely packed β -sheet. If this scenario holds true in future research, this would give additional weight to the role of GxxxG motifs and of membrane composition in the generation of toxic A β oligomers and the onset of Alzheimer's disease. We propose that future studies should elaborate on these recent findings and assess how changes in membrane compositions of the ageing brain affect not only dimerization of C99 but also of A β peptides and how this correlates with generation of toxic A β oligomers. Critically, it would be of major interest to assess how changes in lipid composition associated with ageing [168] affect the structural determinants of C99 processing and amyloid β oligomerization. We believe that understanding how the natural changes in membrane properties of ageing individuals associate with amyloidosis could indeed pave the way for a new array of therapeutic perspectives.

Funding:

NP has received an FSR PhD Fellowship from Université catholique de Louvain and an Aspirant PhD Fellowship from the FRS-FNRS, Belgium. NS is a postdoctoral fellow (Chargé de Recherches) from the FRS-FNRS and received a pilot grant from the Alzheimer Research Foundation (SAO-FRA). Funding to SNC is acknowledged from Ludwig Institute for Cancer Research, Fondation contre le cancer, Salus Sanguinis and Fondation "Les avions de Sébastien", projets Action de recherche concertée (ARC) 16/21-073, research projects from FNRS n°T.0043.21 and WelBio F 44/8/5 - MCF/UIG – 10955. CV is funded by an FWO postdoctoral fellowship 12B7423N and a grant from the consortium Interaction of KULeuven C14/21/095. Funding to PKC acknowledged from SAO-FRA/Stop Alzheimer Research Foundation, Fondation Louvain and Queen Elisabeth Medical Research Foundation, FNRS grant PDRT.0177.18, and UCLouvain Action de Recherche Concertée ARC 21/26-114.

Conflicts of Interest: "The authors declare no conflict of interest."

References

1. Carrell, R.W.; Lomas, D.A. Conformational disease. *The Lancet* **1997**, *350*, 134-138, doi:10.1016/s0140-6736(97)02073-4.
2. Walker, L.C.; LeVine, H. The cerebral proteopathies. *Neurobiology of Aging* **2000**, *21*, 559-561, doi:10.1016/s0197-4580(00)00160-3.
3. Ross, C.A.; Poirier, M.A. Protein aggregation and neurodegenerative disease. *Nat Med* **2004**, *10 Suppl*, S10-17, doi:10.1038/nm1066.
4. Chiti, F.; Dobson, C.M. Protein misfolding, functional amyloid, and human disease. *Annu Rev Biochem* **2006**, *75*, 333-366, doi:10.1146/annurev.biochem.75.101304.123901.
5. Virchow, R. Cellular pathology. As based upon physiological and pathological histology. Lecture XVI—Atheromatous affection of arteries. 1858. *Nutr Rev* **1989**, *47*, 23-25, doi:10.1111/j.1753-4887.1989.tb02747.x.
6. Lomas, D.A.; Carrell, R.W. Serpinopathies and the conformational dementias. *Nat Rev Genet* **2002**, *3*, 759-768, doi:10.1038/nrg907.
7. Sunde, M.; Serpell, L.C.; Bartlam, M.; Fraser, P.E.; Pepys, M.B.; Blake, C.C. Common core structure of amyloid fibrils by synchrotron X-ray diffraction. *J Mol Biol* **1997**, *273*, 729-739, doi:10.1006/jmbi.1997.1348.
8. Serpell, L.C.; Berriman, J.; Jakes, R.; Goedert, M.; Crowther, R.A. Fiber diffraction of synthetic alpha-synuclein filaments shows amyloid-like cross-beta conformation. *Proc Natl Acad Sci U S A* **2000**, *97*, 4897-4902, doi:10.1073/pnas.97.9.4897.
9. Haass, C.; Selkoe, D.J. Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* **2007**, *8*, 101-112, doi:10.1038/nrm2101.
10. Bertram, L.; Tanzi, R.E. The genetic epidemiology of neurodegenerative disease. *J Clin Invest* **2005**, *115*, 1449-1457, doi:10.1172/JCI24761.
11. Goto, Y.; Noji, M.; Nakajima, K.; Yamaguchi, K. Supersaturation-Dependent Formation of Amyloid Fibrils. *Molecules* **2022**, *27*, doi:10.3390/molecules27144588.
12. Morimoto, R.I.; Cuervo, A.M. Proteostasis and the aging proteome in health and disease. *J Gerontol A Biol Sci Med Sci* **2014**, *69 Suppl 1*, S33-38, doi:10.1093/gerona/glu049.
13. Braak, H.; Braak, E. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol. (Berl)* **1991**, *82*, 239-259.
14. Thal, D.R.; Rub, U.; Orantes, M.; Braak, H. Phases of A beta-deposition in the human brain and its relevance for the development of AD. *Neurology* **2002**, *58*, 1791-1800, doi:10.1212/wnl.58.12.1791.
15. Hardy, J. Alzheimer's disease: the amyloid cascade hypothesis: an update and reappraisal. *J. Alzheimers. Dis* **2006**, *9*, 151-153.

16. Hardy, J.; Allsop, D. Amyloid deposition as the central event in the aetiology of Alzheimer's disease. *Trends Pharmacol. Sci* **1991**, *12*, 383-388.

17. Herrup, K. The case for rejecting the amyloid cascade hypothesis. *Nat. Neurosci* **2015**, *18*, 794-799, doi:nn.4017 [pii];10.1038/nn.4017 [doi].

18. Tanzi, R.E. The genetics of Alzheimer disease. *Cold Spring Harb Perspect Med* **2012**, *2*, doi:10.1101/cshperspect.a006296.

19. Kang, J.; Lemaire, H.G.; Unterbeck, A.; Salbaum, J.M.; Masters, C.L.; Grzeschik, K.H.; Multhaup, G.; Beyreuther, K.; Muller-Hill, B. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* **1987**, *325*, 733-736.

20. Slunt, H.H.; Thinakaran, G.; Von Koch, C.; Lo, A.C.; Tanzi, R.E.; Sisodia, S.S. Expression of a ubiquitous, cross-reactive homologue of the mouse beta-amyloid precursor protein (APP). *J Biol Chem* **1994**, *269*, 2637-2644.

21. Tanzi, R.E.; Gusella, J.F.; Watkins, P.C.; Bruns, G.A.; St George-Hyslop, P.; Van Keuren, M.L.; Patterson, D.; Pagan, S.; Kurnit, D.M.; Neve, R.L. Amyloid beta protein gene: cDNA, mRNA distribution, and genetic linkage near the Alzheimer locus. *Science* **1987**, *235*, 880-884, doi:10.1126/science.2949367.

22. Wasco, W.; Bupp, K.; Magendantz, M.; Gusella, J.F.; Tanzi, R.E.; Solomon, F. Identification of a mouse brain cDNA that encodes a protein related to the Alzheimer disease-associated amyloid beta protein precursor. *Proc Natl Acad Sci U S A* **1992**, *89*, 10758-10762, doi:10.1073/pnas.89.22.10758.

23. Musa, A.; Lehrach, H.; Russo, V.A. Distinct expression patterns of two zebrafish homologues of the human APP gene during embryonic development. *Dev Genes Evol* **2001**, *211*, 563-567, doi:10.1007/s00427-001-0189-9.

24. Rosen, D.R.; Martin-Morris, L.; Luo, L.Q.; White, K. A Drosophila gene encoding a protein resembling the human beta-amyloid protein precursor. *Proc Natl Acad Sci U S A* **1989**, *86*, 2478-2482, doi:10.1073/pnas.86.7.2478.

25. Shariati, S.A.; De Strooper, B. Redundancy and divergence in the amyloid precursor protein family. *FEBS Lett* **2013**, *587*, 2036-2045, doi:10.1016/j.febslet.2013.05.026.

26. Glenner, G.G.; Wong, C.W. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun* **1984**, *120*, 885-890, doi:10.1016/s0006-291x(84)80190-4.

27. Tanaka, S.; Nakamura, S.; Ueda, K.; Kameyama, M.; Shiojiri, S.; Takahashi, Y.; Kitaguchi, N.; Ito, H. Three types of amyloid protein precursor mRNA in human brain: Their differential expression in Alzheimer's disease. *Biochemical and Biophysical Research Communications* **1988**, *157*, 472-479, doi:[https://doi.org/10.1016/S0006-291X\(88\)80273-0](https://doi.org/10.1016/S0006-291X(88)80273-0).

28. Sprecher, C.A.; Grant, F.J.; Grimm, G.; O'Hara, P.J.; Norris, F.; Norris, K.; Foster, D.C. Molecular cloning of the cDNA for a human amyloid precursor protein homolog: evidence for a multigene family. *Biochemistry* **1993**, *32*, 4481-4486, doi:10.1021/bi00068a002.

29. Wasco, W.; Gurubhagavatula, S.; Paradis, M.D.; Romano, D.M.; Sisodia, S.S.; Hyman, B.T.; Neve, R.L.; Tanzi, R.E. Isolation and characterization of APLP2 encoding a homologue of the Alzheimer's associated amyloid beta protein precursor. *Nat Genet* **1993**, *5*, 95-100, doi:10.1038/ng0993-95.

30. Müller, U.C.; Zheng, H. Physiological functions of APP family proteins. *Cold Spring Harb Perspect Med* **2012**, *2*, a006288, doi:10.1101/cshperspect.a006288.

31. Wang, Y.; Ha, Y. The X-ray structure of an antiparallel dimer of the human amyloid precursor protein E2 domain. *Mol Cell* **2004**, *15*, 343-353, doi:10.1016/j.molcel.2004.06.037.

32. Lee, S.; Xue, Y.; Hu, J.; Wang, Y.; Liu, X.; Demeler, B.; Ha, Y. The E2 Domains of APP and APLP1 Share a Conserved Mode of Dimerization. *Biochemistry* **2011**, *50*, 5453-5464, doi:10.1021/bi101846x.

33. Roisman, L.C.; Han, S.; Chuei, M.J.; Connor, A.R.; Cappai, R. The crystal structure of amyloid precursor-like protein 2 E2 domain completes the amyloid precursor protein family. *Faseb j* **2019**, *33*, 5076-5081, doi:10.1096/fj.201802315R.

34. Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Žídek, A.; Potapenko, A.; et al. Highly accurate protein structure prediction with AlphaFold. *Nature* **2021**, *596*, 583-589, doi:10.1038/s41586-021-03819-2.

35. Dahms, S.O.; Hoefgen, S.; Roeser, D.; Schlott, B.; Gührs, K.H.; Than, M.E. Structure and biochemical analysis of the heparin-induced E1 dimer of the amyloid precursor protein. *Proc Natl Acad Sci U S A* **2010**, *107*, 5381-5386, doi:10.1073/pnas.0911326107.

36. Lee, S.; Xue, Y.; Hu, J.; Wang, Y.; Liu, X.; Demeler, B.; Ha, Y. The E2 domains of APP and APLP1 share a conserved mode of dimerization. *Biochemistry* **2011**, *50*, 5453-5464, doi:10.1021/bi101846x.

37. Hynes, T.R.; Randal, M.; Kennedy, L.A.; Eigenbrot, C.; Kossiakoff, A.A. X-ray crystal structure of the protease inhibitor domain of Alzheimer's amyloid β -protein precursor. *Biochemistry* **1990**, *29*, 10018-10022, doi:10.1021/bi00495a002.

38. Beher, D.; Hesse, L.; Masters, C.L.; Multhaup, G. Regulation of Amyloid Protein Precursor (APP) Binding to Collagen and Mapping of the Binding Sites on APP and Collagen Type I (*). *Journal of Biological Chemistry* **1996**, *271*, 1613-1620, doi:<https://doi.org/10.1074/jbc.271.3.1613>.

39. Schubert, D.; LaCorbiere, M.; Saitoh, T.; Cole, G. Characterization of an amyloid beta precursor protein that binds heparin and contains tyrosine sulfate. *Proceedings of the National Academy of Sciences* **1989**, *86*, 2066-2069, doi:doi:10.1073/pnas.86.6.2066.

40. Small, D.H.; Nurcombe, V.; Reed, G.; Clarris, H.; Moir, R.; Beyreuther, K.; Masters, C.L. A heparin-binding domain in the amyloid protein precursor of Alzheimer's disease is involved in the regulation of neurite outgrowth. *J Neurosci* **1994**, *14*, 2117-2127, doi:10.1523/jneurosci.14-04-02117.1994.

41. Hoefgen, S.; Coburger, I.; Roeser, D.; Schaub, Y.; Dahms, S.O.; Than, M.E. Heparin induced dimerization of APP is primarily mediated by E1 and regulated by its acidic domain. *Journal of Structural Biology* **2014**, *187*, 30-37, doi:<https://doi.org/10.1016/j.jsb.2014.05.006>.

42. Soba, P.; Eggert, S.; Wagner, K.; Zentgraf, H.; Siehl, K.; Kreger, S.; Löwer, A.; Langer, A.; Merdes, G.; Paro, R.; et al. Homo- and heterodimerization of APP family members promotes intercellular adhesion. *The EMBO Journal* **2005**, *24*, 3624-3634, doi:<https://doi.org/10.1038/sj.emboj.7600824>.

43. Bush, A.I.; Multhaup, G.; Moir, R.D.; Williamson, T.G.; Small, D.H.; Rumble, B.; Pollwein, P.; Beyreuther, K.; Masters, C.L. A novel zinc(II) binding site modulates the function of the beta A4 amyloid protein precursor of Alzheimer's disease. *J Biol Chem* **1993**, *268*, 16109-16112.

44. Multhaup, G.; Schlicksupp, A.; Hesse, L.; Beher, D.; Ruppert, T.; Masters, C.L.; Beyreuther, K. The amyloid precursor protein of Alzheimer's disease in the reduction of copper(II) to copper(I). *Science* **1996**, *271*, 1406-1409, doi:10.1126/science.271.5254.1406.

45. Mayer, M.C.; Kaden, D.; Schauenburg, L.; Hancock, M.A.; Voigt, P.; Roeser, D.; Barucker, C.; Than, M.E.; Schaefer, M.; Multhaup, G. Novel zinc-binding site in the E2 domain regulates amyloid precursor-like protein 1 (APLP1) oligomerization. *J Biol Chem* **2014**, *289*, 19019-19030, doi:10.1074/jbc.M114.570382.

46. Dienemann, C.; Coburger, I.; Mehmedbasic, A.; Andersen, O.M.; Than, M.E. Mutants of metal binding site M1 in APP E2 show metal specific differences in binding of heparin but not of sorLA. *Biochemistry* **2015**, *54*, 2490-2499, doi:10.1021/acs.biochem.5b00111.

47. Dahms, S.O.; König, I.; Roeser, D.; Gührs, K.H.; Mayer, M.C.; Kaden, D.; Multhaup, G.; Than, M.E. Metal binding dictates conformation and function of the amyloid precursor protein (APP) E2 domain. *J Mol Biol* **2012**, *416*, 438-452, doi:10.1016/j.jmb.2011.12.057.

48. Noda, Y.; Asada, M.; Kubota, M.; Maesako, M.; Watanabe, K.; Uemura, M.; Kihara, T.; Shimohama, S.; Takahashi, R.; Kinoshita, A.; et al. Copper enhances APP dimerization and promotes A β production. *Neurosci Lett* **2013**, *547*, 10-15, doi:10.1016/j.neulet.2013.04.057.

49. Rohan de Silva, H.A.; Jen, A.; Wickenden, C.; Jen, L.S.; Wilkinson, S.L.; Patel, A.J. Cell-specific expression of beta-amyloid precursor protein isoform mRNAs and proteins in neurons and astrocytes. *Brain Res Mol Brain Res* **1997**, *47*, 147-156, doi:10.1016/s0169-328x(97)00045-4.

50. Mahdi, F.; Van Nostrand, W.E.; Schmaier, A.H. Protease nexin-2/amyloid beta-protein precursor inhibits factor Xa in the prothrombinase complex. *J Biol Chem* **1995**, *270*, 23468-23474, doi:10.1074/jbc.270.40.23468.

51. Scheinfeld, M.H.; Ghersi, E.; Laky, K.; Fowlkes, B.J.; D'Adamio, L. Processing of beta-amyloid precursor-like protein-1 and -2 by gamma-secretase regulates transcription. *J Biol Chem* **2002**, *277*, 44195-44201, doi:10.1074/jbc.M208110200.

52. Beckett, C.; Nalivaeva, N.N.; Belyaev, N.D.; Turner, A.J. Nuclear signalling by membrane protein intracellular domains: the AICD enigma. *Cell Signal* **2012**, *24*, 402-409, doi:10.1016/j.cellsig.2011.10.007.

53. Pardossi-Piquard, R.; Checler, F. The physiology of the beta-amyloid precursor protein intracellular domain AICD. *J. Neurochem* **2012**, *120 Suppl 1*, 109-124, doi:10.1111/j.1471-4159.2011.07475.x [doi].

54. Cao, X.; Sudhof, T.C. A transcriptively active complex of APP with Fe65 and histone acetyltransferase Tip60. *Science* **2001**, *293*, 115-120.

55. Cao, X.; Sudhof, T.C. Dissection of amyloid-beta precursor protein-dependent transcriptional transactivation. *J. Biol. Chem* **2004**, *279*, 24601-24611.

56. Hebert, S.S.; Serneels, L.; Tolia, A.; Craessaerts, K.; Derkx, C.; Filippov, M.A.; Muller, U.; De Strooper, B. Regulated intramembrane proteolysis of amyloid precursor protein and regulation of expression of putative target genes. *EMBO Rep* **2006**, *7*, 739-745.

57. Opsomer, R.; Contino, S.; Perrin, F.; Gualdani, R.; Tasiaux, B.; Doyen, P.; Vergouts, M.; Vrancx, C.; Doshina, A.; Pierrot, N.; et al. Amyloid Precursor Protein (APP) Controls the Expression of the Transcriptional Activator Neuronal PAS Domain Protein 4 (NPAS4) and Synaptic GABA Release. *eNeuro* **2020**, *7*, doi:ENEURO.0322-19.2020 [pii];10.1523/ENEURO.0322-19.2020 [doi].

58. Sisodia, S.S.; Koo, E.H.; Beyreuther, K.; Unterbeck, A.; Price, D.L. Evidence that beta-amyloid protein in Alzheimer's disease is not derived by normal processing. *Science* **1990**, *248*, 492-495, doi:10.1126/science.1691865.

59. Esch, F.S.; Keim, P.S.; Beattie, E.C.; Blacher, R.W.; Culwell, A.R.; Oltersdorf, T.; McClure, D.; Ward, P.J. Cleavage of amyloid beta peptide during constitutive processing of its precursor. *Science* **1990**, *248*, 1122-1124, doi:10.1126/science.2111583.

60. Eggert, S.; Paliga, K.; Soba, P.; Evin, G.; Masters, C.L.; Weidemann, A.; Beyreuther, K. The proteolytic processing of the amyloid precursor protein gene family members APLP-1 and APLP-2 involves alpha-, beta-, gamma-, and epsilon-like cleavages: modulation of APLP-1 processing by n-glycosylation. *J Biol Chem* **2004**, *279*, 18146-18156, doi:10.1074/jbc.M311601200.

61. Kuhn, P.H.; Wang, H.; Dislich, B.; Colombo, A.; Zeitschel, U.; Ellwart, J.W.; Kremmer, E.; Rossner, S.; Lichtenthaler, S.F. ADAM10 is the physiologically relevant, constitutive alpha-secretase of the amyloid precursor protein in primary neurons. *Embo j* **2010**, *29*, 3020-3032, doi:10.1038/embj.2010.167.

62. Jacobsen, K.T.; Adlerz, L.; Multhaup, G.; Iverfeldt, K. Insulin-like Growth Factor-1 (IGF-1)-induced Processing of Amyloid- β Precursor Protein (APP) and APP-like Protein 2 Is Mediated by Different Metalloproteinases*. *Journal of Biological Chemistry* **2010**, *285*, 10223-10231, doi:<https://doi.org/10.1074/jbc.M109.038224>.

63. Sisodia, S.S. Beta-amyloid precursor protein cleavage by a membrane-bound protease. *Proc Natl Acad Sci U S A* **1992**, *89*, 6075-6079, doi:10.1073/pnas.89.13.6075.

64. Dislich, B.; Lichtenthaler, S.F. The Membrane-Bound Aspartyl Protease BACE1: Molecular and Functional Properties in Alzheimer's Disease and Beyond. *Front Physiol* **2012**, *3*, 8, doi:10.3389/fphys.2012.00008.

65. Deng, Y.; Wang, Z.; Wang, R.; Zhang, X.; Zhang, S.; Wu, Y.; Staufenbiel, M.; Cai, F.; Song, W. Amyloid- β protein (A β) Glu11 is the major β -secretase site of β -site amyloid- β precursor protein-cleaving enzyme 1(BACE1), and shifting the cleavage site to A β Asp1 contributes to Alzheimer pathogenesis. *Eur J Neurosci* **2013**, *37*, 1962-1969, doi:10.1111/ejn.12235.

66. Zhang, S.; Wang, Z.; Cai, F.; Zhang, M.; Wu, Y.; Zhang, J.; Song, W. BACE1 Cleavage Site Selection Critical for Amyloidogenesis and Alzheimer's Pathogenesis. *The Journal of Neuroscience* **2017**, *37*, 6915-6925, doi:10.1523/jneurosci.0340-17.2017.

67. Kimura, A.; Hata, S.; Suzuki, T. Alternative Selection of β -Site APP-Cleaving Enzyme 1 (BACE1) Cleavage Sites in Amyloid β -Protein Precursor (APP) Harboring Protective and Pathogenic Mutations within the A β Sequence. *J Biol Chem* **2016**, *291*, 24041-24053, doi:10.1074/jbc.M116.744722.

68. Weidemann, A.; Eggert, S.; Reinhard, F.B.; Vogel, M.; Paliga, K.; Baier, G.; Masters, C.L.; Beyreuther, K.; Evin, G. A novel epsilon-cleavage within the transmembrane domain of the Alzheimer amyloid precursor protein demonstrates homology with Notch processing. *Biochemistry* **2002**, *41*, 2825-2835, doi:bi015794o [pii].

69. Takami, M.; Nagashima, Y.; Sano, Y.; Ishihara, S.; Morishima-Kawashima, M.; Funamoto, S.; Ihara, Y. γ -Secretase: Successive Tripeptide and Tetrapeptide Release from the Transmembrane Domain of β -Carboxyl Terminal Fragment. *The Journal of Neuroscience* **2009**, *29*, 13042-13052, doi:10.1523/jneurosci.2362-09.2009.

70. Bolduc, D.M.; Montagna, D.R.; Seghers, M.C.; Wolfe, M.S.; Selkoe, D.J. The amyloid-beta forming tripeptide cleavage mechanism of γ -secretase. *Elife* **2016**, *5*, doi:10.7554/elife.17578.

71. Eggert, S.; Paliga, K.; Soba, P.; Evin, G.; Masters, C.L.; Weidemann, A.; Beyreuther, K. The Proteolytic Processing of the Amyloid Precursor Protein Gene Family Members APLP-1 and APLP-2 Involves α -, β -, γ -, and ϵ -Like Cleavages: MODULATION OF APLP-1 PROCESSING BY N-GLYCOSYLATION*. *Journal of Biological Chemistry* **2004**, *279*, 18146-18156, doi:<https://doi.org/10.1074/jbc.M311601200>.

72. Högl, S.; Kuhn, P.H.; Colombo, A.; Lichtenthaler, S.F. Determination of the proteolytic cleavage sites of the amyloid precursor-like protein 2 by the proteases ADAM10, BACE1 and γ -secretase. *PLoS One* **2011**, *6*, e21337, doi:10.1371/journal.pone.0021337.

73. Liu, R.; McAllister, C.; Lyubchenko, Y.; Sierks, M.R. Residues 17-20 and 30-35 of beta-amyloid play critical roles in aggregation. *J Neurosci Res* **2004**, *75*, 162-171, doi:10.1002/jnr.10859.

74. Hu, Y.; Kienlen-Campard, P.; Tang, T.-C.; Perrin, F.; Opsomer, R.; Decock, M.; Pan, X.; Octave, J.-N.; Constantinescu, S.N.; Smith, S.O. β -Sheet Structure within the Extracellular Domain of C99 Regulates Amyloidogenic Processing. *Scientific Reports* **2017**, *7*, 17159, doi:10.1038/s41598-017-17144-0.

75. Söldner, C.A.; Sticht, H.; Horn, A.H.C. Role of the N-terminus for the stability of an amyloid- β fibril with three-fold symmetry. *PLOS ONE* **2017**, *12*, e0186347, doi:10.1371/journal.pone.0186347.

76. Scheuermann, S.; Hambach, B.; Hesse, L.; Stumm, J.; Schmidt, C.; Beher, D.; Bayer, T.A.; Beyreuther, K.; Multhaup, G. Homodimerization of amyloid precursor protein and its implication in the amyloidogenic pathway of Alzheimer's disease. *J. Biol. Chem* **2001**, *276*, 33923-33929.

77. Beher, D.; Hesse, L.; Masters, C.L.; Multhaup, G. Regulation of amyloid protein precursor (APP) binding to collagen and mapping of the binding sites on APP and collagen type I. *J Biol Chem* **1996**, *271*, 1613-1620, doi:10.1074/jbc.271.3.1613.

78. Scheuermann, S.; Hambach, B.; Hesse, L.; Stumm, J.; Schmidt, C.; Beher, D.; Bayer, T.A.; Beyreuther, K.; Multhaup, G. Homodimerization of amyloid precursor protein and its implication in the amyloidogenic pathway of Alzheimer's disease. *J Biol Chem* **2001**, *276*, 33923-33929, doi:10.1074/jbc.M105410200.

79. Ben Khalifa, N.; Tyteca, D.; Marinangeli, C.; Depuydt, M.; Collet, J.F.; Courtoy, P.J.; Renauld, J.C.; Constantinescu, S.; Octave, J.N.; Kienlen-Campard, P. Structural features of the KPI domain control APP dimerization, trafficking, and processing. *Faseb j* **2012**, *26*, 855-867, doi:10.1096/fj.11-190207.

80. Kienlen-Campard, P.; Tasiaux, B.; Van Hees, J.; Li, M.; Huysseune, S.; Sato, T.; Fei, J.Z.; Aimoto, S.; Courtoy, P.J.; Smith, S.O.; et al. Amyloidogenic Processing but Not Amyloid Precursor Protein (APP) Intracellular C-terminal Domain Production Requires a Precisely Oriented APP Dimer Assembled by Transmembrane GXXXG Motifs. *J. Biol. Chem* **2008**, *283*, 7733-7744.

81. Munter, L.M.; Voigt, P.; Harmeyer, A.; Kaden, D.; Gottschalk, K.E.; Weise, C.; Pipkorn, R.; Schaefer, M.; Langosch, D.; Multhaup, G. GxxxG motifs within the amyloid precursor protein transmembrane sequence are critical for the etiology of Abeta42. *EMBO J* **2007**, *26*, 1702-1712.

82. Adams, P.D.; Engelman, D.M.; Brunger, A.T. Improved prediction for the structure of the dimeric transmembrane domain of glycophorin A obtained through global searching. *Proteins* **1996**, *26*, 257-261.

83. Lemmon, M.A.; Treutlein, H.R.; Adams, P.D.; Brunger, A.T.; Engelman, D.M. A dimerization motif for transmembrane alpha-helices. *Nat. Struct. Biol* **1994**, *1*, 157-163.

84. Russ, W.P.; Engelman, D.M. The GxxxG motif: a framework for transmembrane helix-helix association. *J. Mol. Biol* **2000**, *296*, 911-919.

85. Lemmon, M.A.; Flanagan, J.M.; Treutlein, H.R.; Zhang, J.; Engelman, D.M. Sequence specificity in the dimerization of transmembrane alpha-helices. *Biochemistry* **1992**, *31*, 12719-12725, doi:10.1021/bi00166a002.

86. Teese, M.G.; Langosch, D. Role of GxxxG Motifs in Transmembrane Domain Interactions. *Biochemistry* **2015**, *54*, 5125-5135, doi:10.1021/acs.biochem.5b00495.

87. Kirrbach, J.; Krugliak, M.; Ried, C.L.; Pagel, P.; Arkin, I.T.; Langosch, D. Self-interaction of transmembrane helices representing pre-clusters from the human single-span membrane proteins. *Bioinformatics* **2013**, *29*, 1623-1630, doi:10.1093/bioinformatics/btt247.

88. Hendriks, L.; van Duijn, C.M.; Cras, P.; Cruts, M.; Van Hul, W.; van Harskamp, F.; Warren, A.; McInnis, M.G.; Antonarakis, S.E.; Martin, J.J.; et al. Presenile dementia and cerebral haemorrhage linked to a mutation at codon 692 of the beta-amyloid precursor protein gene. *Nat. Genet* **1992**, *1*, 218-221.

89. Harrison, C.F.; Lawson, V.A.; Coleman, B.M.; Kim, Y.S.; Masters, C.L.; Cappai, R.; Barnham, K.J.; Hill, A.F. Conservation of a glycine-rich region in the prion protein is required for uptake of prion infectivity. *J Biol Chem* **2010**, *285*, 20213-20223, doi:10.1074/jbc.M109.093310.

90. Kienlen-Campard, P.; Tasiaux, B.; Van Hees, J.; Li, M.; Huysseune, S.; Sato, T.; Fei, J.Z.; Aimoto, S.; Courtoy, P.J.; Smith, S.O.; et al. Amyloidogenic processing but not amyloid precursor protein (APP) intracellular C-terminal domain production requires a precisely oriented APP dimer assembled by transmembrane GXXXG motifs. *J Biol Chem* **2008**, *283*, 7733-7744, doi:10.1074/jbc.M707142200.

91. Hu, Y.; Kienlen-Campard, P.; Tang, T.C.; Perrin, F.; Opsomer, R.; Decock, M.; Pan, X.; Octave, J.N.; Constantinescu, S.N.; Smith, S.O. beta-Sheet Structure within the Extracellular Domain of C99 Regulates Amyloidogenic Processing. *Sci. Rep* **2017**, *7*, 17159, doi:10.1038/s41598-017-17144-0 [doi];10.1038/s41598-017-17144-0 [pii].

92. Tang, T.C.; Hu, Y.; Kienlen-Campard, P.; El Haylani, L.; Decock, M.; Van Hees, J.; Fu, Z.; Octave, J.N.; Constantinescu, S.N.; Smith, S.O. Conformational changes induced by the A21G Flemish mutation in the amyloid precursor protein lead to increased A β production. *Structure* **2014**, *22*, 387-396, doi:10.1016/j.str.2013.12.012.

93. Beel, A.J.; Molley, C.K.; Kim, H.J.; Tian, F.; Hadziselimovic, A.; Jap, B.; Prestegard, J.H.; Sanders, C.R. Structural studies of the transmembrane C-terminal domain of the amyloid precursor protein (APP): does APP function as a cholesterol sensor? *Biochemistry* **2008**, *47*, 9428-9446, doi:10.1021/bi800993c.

94. Nadezhdin, K.D.; Bocharova, O.V.; Bocharov, E.V.; Arseniev, A.S. Dimeric structure of transmembrane domain of amyloid precursor protein in micellar environment. *FEBS Lett* **2012**, *586*, 1687-1692, doi:S0014-5793(12)00369-9 [pii];10.1016/j.febslet.2012.04.062 [doi].

95. Tang, T.-C.; Hu, Y.; Kienlen-Campard, P.; El Haylani, L.; Decock, M.; Van Hees, J.; Fu, Z.; Octave, J.-N.; Constantinescu, Stefan N.; Smith, Steven O. Conformational Changes Induced by the A21G Flemish Mutation in the Amyloid Precursor Protein Lead to Increased A β Production. *Structure* **2014**, *22*, 387-396, doi:<https://doi.org/10.1016/j.str.2013.12.012>.

96. Perrin, F.; Papadopoulos, N.; Suelves, N.; Opsomer, R.; Vadukul, D.M.; Vrancx, C.; Smith, S.O.; Vertommen, D.; Kienlen-Campard, P.; Constantinescu, S.N. Dimeric Transmembrane Orientations of APP/C99 Regulate γ -Secretase Processing Line Impacting Signaling and Oligomerization. *iScience* **2020**, *23*, 101887, doi:10.1016/j.isci.2020.101887.

97. Higashide, H.; Ishihara, S.; Nobuhara, M.; Ihara, Y.; Funamoto, S. Alanine substitutions in the GXXXG motif alter C99 cleavage by gamma-secretase but not its dimerization. *J Neurochem* **2017**, *140*, 955-962, doi:10.1111/jnc.13942.

98. Decock, M.; El, H.L.; Stanga, S.; Dewachter, I.; Octave, J.N.; Smith, S.O.; Constantinescu, S.N.; Kienlen-Campard, P. Analysis by a highly sensitive split luciferase assay of the regions involved in APP dimerization and its impact on processing. *FEBS Open Bio* **2015**, *5*, 763-773, doi:10.1016/j.fob.2015.09.002 [doi];S2211-5463(15)00087-X [pii].

99. Decock, M.; Stanga, S.; Octave, J.N.; Dewachter, I.; Smith, S.O.; Constantinescu, S.N.; Kienlen-Campard, P. Glycines from the APP GXXXG/GXXXA Transmembrane Motifs Promote Formation of Pathogenic Abeta Oligomers in Cells. *Front Aging Neurosci* **2016**, *8*, 107, doi:10.3389/fnagi.2016.00107 [doi].

100. So, P.P.; Khodr, C.E.; Chen, C.D.; Abraham, C.R. Comparable dimerization found in wildtype and familial Alzheimer's disease amyloid precursor protein mutants. *Am. J. Neurodegener. Dis* **2013**, *2*, 15-28.

101. Sagi, S.A.; Lessard, C.B.; Winden, K.D.; Maruyama, H.; Koo, J.C.; Weggen, S.; Kukar, T.L.; Golde, T.E.; Koo, E.H. Substrate sequence influences gamma-secretase modulator activity, role of the transmembrane domain of the amyloid precursor protein. *J Biol Chem* **2011**, *286*, 39794-39803, doi:10.1074/jbc.M111.277228.

102. Sato, T.; Tang, T.C.; Reubins, G.; Fei, J.Z.; Fujimoto, T.; Kienlen-Campard, P.; Constantinescu, S.N.; Octave, J.N.; Aimoto, S.; Smith, S.O. A helix-to-coil transition at the epsilon-cut site in the transmembrane dimer of the amyloid precursor protein is required for proteolysis. *Proc. Natl. Acad. Sci. U. S. A* **2009**, *106*, 1421-1426.

103. Munter, L.M.; Botev, A.; Richter, L.; Hildebrand, P.W.; Althoff, V.; Weise, C.; Kaden, D.; Multhaup, G. Aberrant amyloid precursor protein (APP) processing in hereditary forms of Alzheimer disease caused by APP familial Alzheimer disease mutations can be rescued by mutations in the APP GxxxG motif. *J. Biol. Chem* **2010**, *285*, 21636-21643, doi:M109.088005 [pii];10.1074/jbc.M109.088005 [doi].

104. Richter, L.; Munter, L.M.; Ness, J.; Hildebrand, P.W.; Dasari, M.; Unterreitmeier, S.; Bulic, B.; Beyermann, M.; Gust, R.; Reif, B.; et al. Amyloid beta 42 peptide (Abeta42)-lowering compounds directly bind to Abeta and interfere with amyloid precursor protein (APP) transmembrane dimerization. *Proc Natl Acad Sci U S A* **2010**, *107*, 14597-14602, doi:10.1073/pnas.1003026107.

105. Eggert, S.; Midthune, B.; Cottrell, B.; Koo, E.H. Induced dimerization of the amyloid precursor protein leads to decreased amyloid-beta protein production. *J Biol Chem* **2009**, *284*, 28943-28952, doi:10.1074/jbc.M109.038646.

106. Zhou, R.; Yang, G.; Guo, X.; Zhou, Q.; Lei, J.; Shi, Y. Recognition of the amyloid precursor protein by human γ -secretase. *Science* **2019**, *363*, doi:10.1126/science.aaw0930.

107. Eggert, S.; Gonzalez, A.C.; Thomas, C.; Schilling, S.; Schwarz, S.M.; Tischer, C.; Adam, V.; Strecker, P.; Schmidt, V.; Willnow, T.E.; et al. Dimerization leads to changes in APP (amyloid precursor protein) trafficking mediated by LRP1 and SorLA. *Cellular and Molecular Life Sciences* **2018**, *75*, 301-322, doi:10.1007/s00018-017-2625-7.

108. Wong, C.-O. Endosomal-Lysosomal Processing of Neurodegeneration-Associated Proteins in Astrocytes. *International Journal of Molecular Sciences* **2020**, *21*, 5149.

109. Edbauer, D.; Winkler, E.; Regula, J.T.; Pesold, B.; Steiner, H.; Haass, C. Reconstitution of gamma-secretase activity. *Nat Cell Biol* **2003**, *5*, 486-488, doi:10.1038/ncb960.

110. Wolfe, M.S.; Xia, W.; Ostaszewski, B.L.; Diehl, T.S.; Kimberly, W.T.; Selkoe, D.J. Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and gamma-secretase activity. *Nature* **1999**, *398*, 513-517, doi:10.1038/19077.

111. Zhang, S.; Zhang, M.; Cai, F.; Song, W. Biological function of Presenilin and its role in AD pathogenesis. *Translational Neurodegeneration* **2013**, *2*, 15, doi:10.1186/2047-9158-2-15.

112. De Strooper, B. Loss-of-function presenilin mutations in Alzheimer disease. Talking Point on the role of presenilin mutations in Alzheimer disease. *EMBO Rep* **2007**, *8*, 141-146, doi:10.1038/sj.embor.7400897.

113. Sannerud, R.; Esselens, C.; Ejsmont, P.; Mattera, R.; Rochin, L.; Tharkeshwar, A.K.; De Baets, G.; De Wever, V.; Habets, R.; Baert, V.; et al. Restricted Location of PSEN2/γ-Secretase Determines Substrate Specificity and Generates an Intracellular Aβ Pool. *Cell* **2016**, *166*, 193-208, doi:10.1016/j.cell.2016.05.020.

114. Bentahir, M.; Nyabi, O.; Verhamme, J.; Tolia, A.; Horré, K.; Wiltfang, J.; Esselmann, H.; De Strooper, B. Presenilin clinical mutations can affect gamma-secretase activity by different mechanisms. *J Neurochem* **2006**, *96*, 732-742, doi:10.1111/j.1471-4159.2005.03578.x.

115. Vrancx, C.; Vadukul, D.M.; Suelves, N.; Contino, S.; D'Auria, L.; Perrin, F.; van Pesch, V.; Hanseeuw, B.; Quinton, L.; Kienlen-Campard, P. Mechanism of Cellular Formation and In Vivo Seeding Effects of Hexameric beta-Amyloid Assemblies. *Mol Neurobiol* **2021**, *58*, 6647-6669, doi:10.1007/s12035-021-02567-8.

116. Ye, X.; Cai, Q. Snapin-mediated BACE1 retrograde transport is essential for its degradation in lysosomes and regulation of APP processing in neurons. *Cell Rep* **2014**, *6*, 24-31, doi:10.1016/j.celrep.2013.12.008.

117. Huse, J.T.; Pijak, D.S.; Leslie, G.J.; Lee, V.M.; Doms, R.W. Maturation and endosomal targeting of beta-site amyloid precursor protein-cleaving enzyme. The Alzheimer's disease beta-secretase. *J Biol Chem* **2000**, *275*, 33729-33737, doi:10.1074/jbc.M004175200.

118. Pastorino, L.; Ikin, A.F.; Nairn, A.C.; Pursnani, A.; Buxbaum, J.D. The carboxyl-terminus of BACE contains a sorting signal that regulates BACE trafficking but not the formation of total A(β). *Mol Cell Neurosci* **2002**, *19*, 175-185, doi:10.1006/mcne.2001.1065.

119. Das, U.; Scott, D.A.; Ganguly, A.; Koo, E.H.; Tang, Y.; Roy, S. Activity-induced convergence of APP and BACE-1 in acidic microdomains via an endocytosis-dependent pathway. *Neuron* **2013**, *79*, 447-460, doi:10.1016/j.neuron.2013.05.035.

120. Marinangeli, C.; Tasiaux, B.; Opsomer, R.; Hage, S.; Sodero, A.O.; Dewachter, I.; Octave, J.N.; Smith, S.O.; Constantinescu, S.N.; Kienlen-Campard, P. Presenilin transmembrane domain 8 conserved AXXXAXXXG motifs are required for the activity of the gamma-secretase complex. *J. Biol. Chem* **2015**, *290*, 7169-7184, doi:M114.601286 [pii];10.1074/jbc.M114.601286 [doi].

121. Lee, S.F.; Shah, S.; Yu, C.; Wigley, W.C.; Li, H.; Lim, M.; Pedersen, K.; Han, W.; Thomas, P.; Lundkvist, J.; et al. A conserved GXXXG motif in APH-1 is critical for assembly and activity of the gamma-secretase complex. *J. Biol. Chem* **2004**, *279*, 4144-4152.

122. Liu, L.; Lauro, B.M.; Wolfe, M.S.; Selkoe, D.J. Hydrophilic loop 1 of Presenilin-1 and the APP GxxxG transmembrane motif regulate gamma-secretase function in generating Alzheimer-causing Abeta peptides. *J Biol Chem* **2021**, *296*, 100393, doi:10.1016/j.jbc.2021.100393.

123. Alzheimer, A. über eigenartige Krankheitsfälle des späteren Alters. *Zeitschrift für die gesamte Neurologie und Psychiatrie* **1911**, *4*, 356, doi:10.1007/BF02866241.

124. Foley, P. Lipids in Alzheimer's disease: A century-old story. *Biochim Biophys Acta* **2010**, *1801*, 750-753, doi:10.1016/j.bbapap.2010.05.004.

125. Morgado, I.; Garvey, M. Lipids in Amyloid-β Processing, Aggregation, and Toxicity. In *Lipids in Protein Misfolding*, Gursky, O., Ed.; Springer International Publishing: Cham, 2015; pp. 67-94.

126. Grimm, M.O.W.; Mett, J.; Grimm, H.S.; Hartmann, T. APP Function and Lipids: A Bidirectional Link. *Frontiers in Molecular Neuroscience* **2017**, *10*, doi:10.3389/fnmol.2017.00063.

127. Kivipelto, M.; Helkala, E.L.; Laakso, M.P.; Hänninen, T.; Hallikainen, M.; Alhainen, K.; Soininen, H.; Tuomilehto, J.; Nissinen, A. Midlife vascular risk factors and Alzheimer's disease in later life: longitudinal, population based study. *Bmj* **2001**, *322*, 1447-1451, doi:10.1136/bmj.322.7300.1447.

128. Kuo, Y.M.; Emmerling, M.R.; Bisgaier, C.L.; Essenburg, A.D.; Lampert, H.C.; Drumm, D.; Roher, A.E. Elevated low-density lipoprotein in Alzheimer's disease correlates with brain abeta 1-42 levels. *Biochem Biophys Res Commun* **1998**, *252*, 711-715, doi:10.1006/bbrc.1998.9652.

129. Sáiz-Vazquez, O.; Puente-Martínez, A.; Ubillos-Landa, S.; Pacheco-Bonrostro, J.; Santabárbara, J. Cholesterol and Alzheimer's Disease Risk: A Meta-Meta-Analysis. *Brain Sci* **2020**, *10*, doi:10.3390/brainsci10060386.

130. Corder, E.H.; Saunders, A.M.; Strittmatter, W.J.; Schmeichel, D.E.; Gaskell, P.C.; Small, G.W.; Roses, A.D.; Haines, J.L.; Pericak-Vance, M.A. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* **1993**, *261*, 921-923, doi:10.1126/science.8346443.

131. Saunders, A.M.; Strittmatter, W.J.; Schmeichel, D.; George-Hyslop, P.H.; Pericak-Vance, M.A.; Joo, S.H.; Rosi, B.L.; Gusella, J.F.; Crapper-MacLachlan, D.R.; Alberts, M.J.; et al. Association of apolipoprotein E allele epsilon 4 with late-onset familial and sporadic Alzheimer's disease. *Neurology* **1993**, *43*, 1467-1472, doi:10.1212/wnl.43.8.1467.

132. Strittmatter, W.J.; Saunders, A.M.; Schmeichel, D.; Pericak-Vance, M.; Enghild, J.; Salvesen, G.S.; Roses, A.D. Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc Natl Acad Sci U S A* **1993**, *90*, 1977-1981, doi:10.1073/pnas.90.5.1977.

133. Bodovitz, S.; Klein, W.L. Cholesterol modulates alpha-secretase cleavage of amyloid precursor protein. *J Biol Chem* **1996**, *271*, 4436-4440, doi:10.1074/jbc.271.8.4436.

134. Simons, M.; Keller, P.; De Strooper, B.; Beyreuther, K.; Dotti, C.G.; Simons, K. Cholesterol depletion inhibits the generation of beta-amyloid in hippocampal neurons. *Proc Natl Acad Sci U S A* **1998**, *95*, 6460-6464, doi:10.1073/pnas.95.11.6460.

135. Ghribi, O.; Larsen, B.; Schrag, M.; Herman, M.M. High cholesterol content in neurons increases BACE, beta-amyloid, and phosphorylated tau levels in rabbit hippocampus. *Exp Neurol* **2006**, *200*, 460-467, doi:10.1016/j.expneurol.2006.03.019.

136. Cordy, J.M.; Hussain, I.; Dingwall, C.; Hooper, N.M.; Turner, A.J. Exclusively targeting α -secretase to lipid rafts by GPI-anchor addition up-regulates α -site processing of the amyloid precursor protein. *Proceedings of the National Academy of Sciences* **2003**, *100*, 11735-11740, doi:doi:10.1073/pnas.1635130100.

137. Song, Y.; Kenworthy, A.K.; Sanders, C.R. Cholesterol as a co-solvent and a ligand for membrane proteins. *Protein Sci* **2014**, *23*, 1-22, doi:10.1002/pro.2385.

138. Barrett, P.J.; Song, Y.; Van Horn, W.D.; Hustedt, E.J.; Schafer, J.M.; Hadziselimovic, A.; Beel, A.J.; Sanders, C.R. The amyloid precursor protein has a flexible transmembrane domain and binds cholesterol. *Science* **2012**, *336*, 1168-1171, doi:10.1126/science.1219988.

139. Song, Y.; Hustedt, E.J.; Brandon, S.; Sanders, C.R. Competition between homodimerization and cholesterol binding to the C99 domain of the amyloid precursor protein. *Biochemistry* **2013**, *52*, 5051-5064, doi:10.1021/bi400735x.

140. Langness, V.F.; van der Kant, R.; Das, U.; Wang, L.; Chaves, R.D.S.; Goldstein, L.S.B. Cholesterol-lowering drugs reduce APP processing to A β by inducing APP dimerization. *Mol Biol Cell* **2021**, *32*, 247-259, doi:10.1091/mbc.E20-05-0345.

141. Dominguez, L.; Foster, L.; Straub, J.E.; Thirumalai, D. Impact of membrane lipid composition on the structure and stability of the transmembrane domain of amyloid precursor protein. *Proceedings of the National Academy of Sciences* **2016**, *113*, E5281-E5287, doi:doi:10.1073/pnas.1606482113.

142. Li, C.-D.; Junaid, M.; Shan, X.; Wang, Y.; Wang, X.; Khan, A.; Wei, D.-Q. Effect of Cholesterol on C99 Dimerization: Revealed by Molecular Dynamics Simulations. *Frontiers in Molecular Biosciences* **2022**, *9*, doi:10.3389/fmolb.2022.872385.

143. Sun, F.; Chen, L.; Wei, P.; Chai, M.; Ding, X.; Xu, L.; Luo, S.-Z. Dimerization and Structural Stability of Amyloid Precursor Proteins Affected by the Membrane Microenvironments. *Journal of Chemical Information and Modeling* **2017**, *57*, 1375-1387, doi:10.1021/acs.jcim.7b00196.

144. Miyashita, N.; Straub, J.E.; Thirumalai, D.; Sugita, Y. Transmembrane structures of amyloid precursor protein dimer predicted by replica-exchange molecular dynamics simulations. *J. Am. Chem. Soc* **2009**, *131*, 3438-3439, doi:10.1021/ja809227c [doi].

145. Gorman, P.M.; Kim, S.; Guo, M.; Melnyk, R.A.; McLaurin, J.; Fraser, P.E.; Bowie, J.U.; Chakrabartty, A. Dimerization of the transmembrane domain of amyloid precursor proteins and familial Alzheimer's disease mutants. *BMC Neurosci* **2008**, *9*, 17, doi:10.1186/1471-2202-9-17.

146. Yan, Y.; Xu, T.H.; Harikumar, K.G.; Miller, L.J.; Melcher, K.; Xu, H.E. Dimerization of the transmembrane domain of amyloid precursor protein is determined by residues around the γ -secretase cleavage sites. *J Biol Chem* **2017**, *292*, 15826-15837, doi:10.1074/jbc.M117.789669.

147. So, P.P.; Khodr, C.E.; Chen, C.D.; Abraham, C.R. Comparable dimerization found in wildtype and familial Alzheimer's disease amyloid precursor protein mutants. *Am J Neurodegener Dis* **2013**, *2*, 15-28.

148. Muller-Schiffmann, A.; Herring, A.; Abdel-Hafiz, L.; Chepkova, A.N.; Schable, S.; Wedel, D.; Horn, A.H.; Sticht, H.; de Souza Silva, M.A.; Gottmann, K.; et al. Amyloid-beta dimers in the absence of plaque pathology impair learning and synaptic plasticity. *Brain* **2016**, *139*, 509-525, doi:10.1093/brain/awv355.

149. Shankar, G.M.; Bloodgood, B.L.; Townsend, M.; Walsh, D.M.; Selkoe, D.J.; Sabatini, B.L. Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J Neurosci* **2007**, *27*, 2866-2875, doi:10.1523/JNEUROSCI.4970-06.2007.

150. Townsend, M.; Shankar, G.M.; Mehta, T.; Walsh, D.M.; Selkoe, D.J. Effects of secreted oligomers of amyloid beta-protein on hippocampal synaptic plasticity: a potent role for trimers. *J Physiol* **2006**, *572*, 477-492, doi:10.1113/jphysiol.2005.103754.

151. Barykin, E.P.; Mitkevich, V.A.; Kozin, S.A.; Makarov, A.A. Amyloid beta Modification: A Key to the Sporadic Alzheimer's Disease? *Front Genet* **2017**, *8*, 58, doi:10.3389/fgene.2017.00058.

152. Kummer, M.P.; Heneka, M.T. Truncated and modified amyloid-beta species. *Alzheimers Res Ther* **2014**, *6*, 28, doi:10.1186/alzrt258.

153. Domingo, G.; Benussi, L.; Saraceno, C.; Bertuzzi, M.; Nicsanu, R.; Longobardi, A.; Bellini, S.; Cagnotto, A.; Salmona, M.; Binetti, G.; et al. N-Terminally Truncated and Pyroglutamate-Modified Abeta Forms Are Measurable in Human Cerebrospinal Fluid and Are Potential Markers of Disease Progression in Alzheimer's Disease. *Front Neurosci* **2021**, *15*, 708119, doi:10.3389/fnins.2021.708119.

154. Kumar, S.; Rezaei-Ghaleh, N.; Terwel, D.; Thal, D.R.; Richard, M.; Hoch, M.; Mc Donald, J.M.; Wullner, U.; Glebov, K.; Heneka, M.T.; et al. Extracellular phosphorylation of the amyloid beta-peptide promotes formation of toxic aggregates during the pathogenesis of Alzheimer's disease. *EMBO J* **2011**, *30*, 2255-2265, doi:10.1038/emboj.2011.138.

155. Kumar, S.; Wirths, O.; Stuber, K.; Wunderlich, P.; Koch, P.; Theil, S.; Rezaei-Ghaleh, N.; Zweckstetter, M.; Bayer, T.A.; Brustle, O.; et al. Phosphorylation of the amyloid beta-peptide at Ser26 stabilizes oligomeric assembly and increases neurotoxicity. *Acta Neuropathol* **2016**, *131*, 525-537, doi:10.1007/s00401-016-1546-0 [doi];10.1007/s00401-016-1546-0 [pii].

156. Nirmalraj, P.N.; List, J.; Battacharya, S.; Howe, G.; Xu, L.; Thompson, D.; Mayer, M. Complete aggregation pathway of amyloid beta (1-40) and (1-42) resolved on an atomically clean interface. *Sci Adv* **2020**, *6*, eaaz6014, doi:10.1126/sciadv.aaz6014.

157. Almeida, Z.L.; Brito, R.M.M. Structure and Aggregation Mechanisms in Amyloids. *Molecules* **2020**, *25*, 1195.

158. Vadukul, D.M.; Vrancx, C.; Burguet, P.; Contino, S.; Suelves, N.; Serpell, L.C.; Quinton, L.; Kienlen-Campard, P. An evaluation of the self-assembly enhancing properties of cell-derived hexameric amyloid-beta. *Sci. Rep* **2021**, *11*, 11570, doi:10.1038/s41598-021-90680-y [doi];10.1038/s41598-021-90680-y [pii].

159. Sato, T.; Kienlen-Campard, P.; Ahmed, M.; Liu, W.; Li, H.; Elliott, J.I.; Aimoto, S.; Constantinescu, S.N.; Octave, J.N.; Smith, S.O. Inhibitors of amyloid toxicity based on beta-sheet packing of Abeta40 and Abeta42. *Biochemistry* **2006**, *45*, 5503-5516.

160. Misra, P.; Kodali, R.; Chemuru, S.; Kar, K.; Wetzel, R. Rapid alpha-oligomer formation mediated by the Abeta C terminus initiates an amyloid assembly pathway. *Nat Commun* **2016**, *7*, 12419, doi:10.1038/ncomms12419.

161. Barrett, P.J.; Song, Y.; Van Horn, W.D.; Hustedt, E.J.; Schafer, J.M.; Hadziselimovic, A.; Beel, A.J.; Sanders, C.R. The amyloid precursor protein has a flexible transmembrane domain and binds cholesterol. *Science* **2012**, *336*, 1168-1171, doi:336/6085/1168 [pii];10.1126/science.1219988 [doi].

162. Beel, A.J.; Sakakura, M.; Barrett, P.J.; Sanders, C.R. Direct binding of cholesterol to the amyloid precursor protein: An important interaction in lipid-Alzheimer's disease relationships? *Biochim. Biophys. Acta* **2010**, *1801*, 975-982, doi:S1388-1981(10)00064-8 [pii];10.1016/j.bbapap.2010.03.008 [doi].

163. Miller, Y.; Ma, B.; Nussinov, R. Synergistic interactions between repeats in tau protein and Abeta amyloids may be responsible for accelerated aggregation via polymorphic states. *Biochemistry* **2011**, *50*, 5172-5181, doi:10.1021/bi200400u.

164. Misra, P.; Kodali, R.; Chemuru, S.; Kar, K.; Wetzel, R. Rapid α -oligomer formation mediated by the A β C terminus initiates an amyloid assembly pathway. *Nat Commun* **2016**, *7*, 12419, doi:10.1038/ncomms12419.

165. Liu, C.C.; Liu, C.C.; Kanekiyo, T.; Xu, H.; Bu, G. Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. *Nat Rev Neurol* **2013**, *9*, 106-118, doi:10.1038/nrneurol.2012.263.

166. Umeda, T.; Tomiyama, T.; Kitajima, E.; Idomoto, T.; Nomura, S.; Lambert, M.P.; Klein, W.L.; Mori, H. Hypercholesterolemia accelerates intraneuronal accumulation of A β oligomers resulting in memory impairment in Alzheimer's disease model mice. *Life Sci* **2012**, *91*, 1169-1176, doi:10.1016/j.lfs.2011.12.022.

167. Lazar, A.N.; Bich, C.; Panchal, M.; Desbenoit, N.; Petit, V.W.; Touboul, D.; Dauphinot, L.; Marquer, C.; Laprévote, O.; Brunelle, A.; et al. Time-of-flight secondary ion mass spectrometry (TOF-SIMS) imaging reveals cholesterol overload in the cerebral cortex of Alzheimer disease patients. *Acta Neuropathol* **2013**, *125*, 133-144, doi:10.1007/s00401-012-1041-1.

168. Gille, B.; Galuska, C.E.; Fuchs, B.; Peleg, S. Recent Advances in Studying Age-Associated Lipids Alterations and Dietary Interventions in Mammals. *Front Aging* **2021**, *2*, 773795, doi:10.3389/fragi.2021.773795.