

Article

In Vitro Multiplication of *Lophostemon suaveolens* (Sol.ex Gaertn.) Peter G.Wilson & J.T. Waterh): Peatland Tree Species for Rehabilitation

Asri Insiana Putri *, Noor Khomsah Kartikawati, Arif Nirsatmanto, Sri Sunarti, Liliek Haryjanto, Toni Herawan, Purwanto Budi Santosa, Reni Setyo Wahyuningtyas, Fajar Lestari and Anto Rimbawanto

¹ National Research and Innovation Agency (BRIN) Republic of Indonesia, Jl. Ir. H. Juanda No. 18 Paledang, Kota Bogor, Jawa Barat, Indonesia; asri008@brin.go.id (AIP); noor018@brin.go.id (N.K.K.); arif050@brin.go.id (A.N.); sris017@brin.go.id (S.S.); lili019@brin.go.id (L.H.); toni005@brin.go.id (T.H.); purw021@brin.go.id (P.B.S.); reni009@brin.go.id (R.S.W.); fajar.lestari@brin.go.id (F.L.); anto022@brin.go.id (A.R.)

* Correspondence author: asri008@brin.go.id, +62 81559728514

Abstract: Peatlands in Indonesia are one of the world's largest carbon sinks, helping to regulate greenhouse gas emissions and global climate change. *Lophostemon suaveolens* is a relatively unexplored plant found in Papua's endemic peat ecosystem that grows well in wet areas with low fertility. It is geographically dispersed and has the potential for peatland rehabilitation. Seed is one of materials for the reproduction of *L. suaveolens*. However, the difficulty in seed collection and the limitation in seed production become a current problem for its cultivation. Seed multiplication by using *in vitro* method would be one of the mechanisms to overcome the problem. We present an efficient and reproducible protocol for *in vitro* multiplication of plantlets using nodal segments and shoot apices collected from plantlets. After 3 months of the culture initiation stage, the elongated axillary shoots were separated from the clumps and further multiplied using Murashige and Skoog (MS) media supplemented with (1) BAP (0.5 ml/l) as single PGR, (2) NAA (0.1 ml/l) as a single PGR and (3) a combination of two types of PGR BAP (0.5 ml/l) and NAA (0.1 ml/l). Up to an incubation period of 6 months, the efficiency of leaf axillary shoot propagation was determined by counting the number of nodule multiplication coefficient (NMC), shoot length, root length, and number of leaves (6 consecutive subcultures). The higher the NMC, the higher the plantlets that will be obtained, increasing shoot regeneration from nodules physiologically increasing evapotranspiration *in vitro*. The highest of NMC (8.4) was observed in MS medium with a combination of 0.5 ml/l BAP and 0.1 ml/l NAA (double PGRs), with the longest shoots (5.91 cm), the longest root length (8.83 cm), and the most leaves (32). When combination of BAP and NAA were used simultaneously, the plantlets during acclimatization was the highest survived. It was concluded that MS in combination with 0.5 ml/l BAP and 0.1 ml/l NAA has been the most appropriate protocol for the success of *in vitro* multiplication of *L. suaveolens*. This is the first report of *L. suaveolens in vitro* multiplication, and the protocol could be used to propagate this peatland species on a large scale. The authors acknowledge the limitations of the experimental work and recommend further work to increase the sample size and complete the field-testing phase to help verify the initial findings presented in this paper

Keywords: acclimatization; axillary shoots; BAP; NAA; propagation

1. Introduction

Indonesia has the world's largest tropical peatland [1]. It is estimated that tropical peat land retains approximately 40% of terrestrial carbon [2], Traditional slash and burn practices for agricultural and plantation purposes have resulted in large peat fires that released a large amount of carbon to the atmosphere [3], potentially affecting atmospheric

composition and the climate system. In 2015, the worst peat fires since 1997 ravaged Sumatera, Kalimantan, and Papua. The sensitivity of peatland degradation varies depending on geographic location. Zhang [4] reported on the sensitivity of peat degradation via remote sensing observation of trench ecology in the Zoige peatland, East Tibetan plateau. Because peat soils decompose more strongly at higher elevations, Alpine peat is more sensitive to disturbances associated with the groundwater table. Valjarevic et al, [5] used remote sensing, topographic maps, geographic information system (GIS) analysis, and official data from cadastral and census books to reconstruct forest conditions, including degraded land, in Toplica, Republic of Serbia over the last 60 years. The combination of remote sensing techniques and micro-techniques can be a mutually reinforcing option for land rehabilitation. The combination of remote sensing techniques and micro-techniques can be a mutually reinforcing alternative for the rehabilitation of degraded land.

Peatland rehabilitation aims to restore former peatland ecosystem processes, productivity, and services, but it does not imply restoring pre-existing biotic integrity in terms of species composition, community structure, and ecosystem functions [6]. Peatland restoration, also known as the process of restoring peatlands, refers to the process of restoring peatland ecosystems.

Peat restoration, also known as peatland restoration process, refers to the process of restoring peatland ecosystems. Peatland restoration, or the process of returning degraded or degraded peatlands to their natural state, takes this a step further [7]. The global policy for the rehabilitation of marginal lands is focused on the concept of tropical forest and landscape restoration. Intensification techniques are the focus of ecology to improve crop quality in degraded landscape areas [14]. Unlike well-established studies on ecological restoration of isolated peatlands [7], tropical peatland restoration is still in its early stages [3][8],[9]. This research supports the vegetative propagation of quality plants for land rehabilitation. The basic operational model for the purpose of reforestation using *in vitro* propagation techniques has been carried out in Brazil, mainly for 15 tree species, ten woody trees (*Couroupita guianensis* Aubl., *Tabebuia heptaphylla* (Vell.) Toledo, *Tabebuia impetiginosa* (Mart. ex DC.) Standl., *Tabebuia roseoalba* (Ridl.) Sandwith, *Vochysia haenkeana* (Spreng.) Mart., *Vitex montevidensis* Cham., *Copaifera coriacea* Mart., *Spondias tuberosa* Arruda, *Schinus terebinthifolia* Raddi, and *Talisia esculenta* (A.) Radlk.-Hil) and five palm trees (*Syagrus coronate* (Mart.) Becc., *Attalea oleifera* Barb. Rodr., *Elaeis guineensis* Jacq., *Colubrina glandulosa* Perk., and *Astrocaryum vulgare* Mart.) [14].

According to the ministerial regulation, one of the plant criteria for peat ecosystem rehabilitation was to prioritize native species and land suitability. *Lophostemon suaveolens*, a plant in the Myrtaceae family, was found in the peat ecosystem. *L. suaveolens* is a tree or shrub with terminal buds covered in scales, stems and petioles that exude a milky resin when cut. On mature branches, the leaves alternate, and the inflorescences are axillary and cymose. White 5-merous flowers sepals are lobed and sometimes persistent in fruit. Fruit a loculicidal capsule not exceeding, or barely exceeding, the fruiting hypanthium, 3-valved; seeds linear [9]. *L. suaveolens* is the dominant tree species in the Merauke area, and it is widely used by the community for building materials, firewood, and a variety of other purposes [10]. Forest rehabilitation is the process of restoring degraded forest structures in order to generate forest products and productivity, but it does not always include all plant and animal species that are expected to exist in a given location. A large number of long-lasting seeds are required for large-scale restoration. Seedling propagation must be efficient in order to meet demand on a large scale. Seed is one of materials for the reproduction of *L. suaveolens*. However, the difficulty in seed collection and the limitation in seed production become a current problem for its cultivation. Seedlings of *Lophostemon* sp. cannot be propagated naturally in sufficient numbers. Micropropagation is an alternative method of vegetative reproduction to traditional methods. In forestry, *in vitro* propagation techniques are useful for mass reproduction of superior trees found in nature, and

axillary shoot proliferation has been found to be the most successful in achieving plantlet regeneration from mature forest trees [12].

One technique for vegetative propagation is multiplication *in vitro* culture. *In vitro* culture is a propagation technique that isolates certain plant parts and then stimulates them to multiplication and rooting form new plants by modifying the growing media and environment suitable for plants. *In vitro* culture techniques are able to produce large amounts of planting material in a relatively short time, are uniform, free of pathogens, and do not depend on the season. Multiplication *in vitro* culture, growth and development of explants are strongly influenced by the type of base media and plant growth regulators. To increase explant propagation, growth regulators in the form of auxin and cytokinin groups are generally added to the growing media. This growth regulator stimulates the process of growth and development of plant tissue [13]. The auxin-cytokinin ratio determines the type of culture established or regeneration type. 1-Naphthaleneacetic acid (NAA) is classified as an auxin and 6-Benzylaminopurine (BAP) is classified as a cytokinin based on their primary function *in vitro* and *ex vitro* propagation [15]. NAA, which is synthesized in the shoot apex of young leaves and transported basipetally to the roots [16], is essential for maintaining apical dominance and cell elongation. Their main effects include rooting, stimulation, and inhibition of axillary bud outgrowth [17]. Cytokinin regulate meristem activity [18], plant shape determination, side condition adaptation, and environmental responses. NAA and BAP act either antagonistically or synergistically to control developmental processes such as meristem formation and maintenance [19]. BAP and NAA are a type of PGRs the most often used in shoot propagation because these are stable, resistant to degradation and readily available. Murashige and Skoog (MS) media are the basic media that are generally used for the propagation of various types of plants. This basic medium is rich in minerals that stimulate organogenesis. To increase the propagation of explants, growth regulators in the form of auxin and cytokinin groups are usually added to the growing media. This growth regulator stimulates the process of growth and development of plant tissue [20].

In vitro multiplication has emerged as a successful and effective technique for mass propagation of species suitable for peatland rehabilitation. The lack of efficient propagation methods, combined with the high growth potential in these marginal peatlands, highlights the importance of germplasm and the development of *in vitro* propagation protocols. The purpose of this study was to observe the addition of exogenous BAP and NAA which were thought to increase the multiplication coefficient, increase shoot elongation, root elongation, and number of leaves on *L. suaveolens* to maintain the availability of stronger vegetative propagation seeds for rehabilitation on marginal land. Stable nursery architecture will promote optimal growth in nursery acclimatization. The effect of NAA and BAP on the *in vitro* multiplication of *L. suaveolens* has not been studied in depth.

This research has a strategic value to obtain techniques for providing plantlets or seedlings on marginal lands that are difficult to manage. *In vitro* enrichment of PGRs has the potential to strengthen seedlings resulting from more efficient vegetative multiplication on a large scale. The weakness of this study is that it does not fully display the entire micropropagation process, starting from the preparation of explant material from trees in nature to field trials of planting seedlings resulting from *in vitro* multiplication. However, it can actually focus more on displaying research results on the most important part of *in vitro* multiplication for *L. suaveolens* which has not been reported by other studies

2. Materials and Methods

The research was carried out at the tissue culture laboratory, greenhouse, and nursery at the Center for Forest Biotechnology and Tree Improvement in Yogyakarta, Indonesia (7°40'20" S and 110°23'30"), 600 meters above sea level, with the highest temperature being 32 degrees Celsius and the lowest temperature being 18 degrees Celsius, and a mean humidity of 95%. Researchers observed for 13 months in the laboratory and 14 months in the greenhouse from January 2018 to March 2021. Genetic material source and

newly formed shoots of plantlet were taken from PT. Arara Abadi tissue culture laboratory at Riau, Sumatera, Indonesia. The equipment used in this study met tissue culture laboratory standards.

3. Multiplication of *L. suaveolens*

The goal of this phase is to produce more *L. suaveolens* propagules from shoots. Micro cutting technique and repeated subcultures multiply the number of propagules until the desired number of plants is obtained. The treatment composition was transferred into several new culture media after the nodule of the axenic plantlets was separated as an explant. This study's in vitro multiplication of *L. suaveolens* is a follow-up to preliminary observations on the initiation of axillary buds on various basic media and PGRs [21]. Several modifications of explants disinfection method by [22] were used in this study to obtain aseptic explants. Disinfection of explants was continued in the LAF by soaking and shaking the explants for 40 minutes in 15% (v/v) antimicrobial compound biocide isothiazolone (BI) and a few drops of Tween⁸⁰ for 40 minutes. The final step of disinfection is to soak and shake the explants for one minute in 70% ethanol. After every disinfection, explants rinsed with sterile distilled water.

Aseptic explants were cultured in glass tubes (25 mm 150 mm) with 25 ml of MS medium supplemented with single PGR treatment (one kind of BAP 0.5 ml/l and one kind of NAA 0.1 ml/l) and double PGRs treatment (combination of 0.5 ml/l BAP and 0.1 ml/l NAA). Before being gelled in 8 g/l of Phyto agar (King of Fiber, Indonesia) for multiplication and 6 g/l of Phyto agar for rooting, each basal medium contained vitamin and was supplemented with 30 g/l sugar (Sugar Group Companies, Indonesia). Before autoclaving at 121°C, the pH of the medium was adjusted to 5.8 by adding a few drops of 1 N hydrochloric acid 32% (Merck 100313) or sodium hydroxide (Merck 1310-73-2), and all steps were repeated three times to ensure reproducibility in media preparation. The experimental setup was kept in a laboratory at 25°C and 60% relative humidity under white fluorescent light at 40 mol/m²/s intensity with a 16 h light and 8 h dark cycle.

Each PGR and PGRs treatment at this multiplication stage were tested in 20 culture replicates, each culture containing one explant consisting of one nodule. The multiplication potential of explants was expressed as nodule multiplication coefficient (NMC), shoot length, root length, and number of leaves of *L. suaveolens* in vitro which were determined after 6 months of incubation. The following is how NMC is defined:

$$\text{NMC} = \frac{(\% \text{ viable explants}) \times (\text{average of nodules/explant})}{\text{number of plantlets transplanted}} \times 100 \quad (1)$$

3.1. Rooting and acclimatization of *L. suaveolens*

After 6 months of incubation and rooting regeneration, another 4 months without subculture was performed to ensure that the plantlets were healthy with well-developed roots *in vitro* prior to acclimatization in the green house. The plantlets were washed under running tap water, soaked in fungicide (Mankoze 80%) for several minutes, and transplanted into a polybag containing a mixture of regosol topsoil (sandy soil) and compost from bamboo leaf litter in a 1:1 ratio. Acclimatization of plantlets was done in a greenhouse with 30-40% natural light and watered once a day for 6 months. Total 50 plantlets were individually transplanted in a poly bag with a plastic cover in this experiment. No control in acclimatization, just make sure the best rooting in vitro can survive in soil medium and green house condition.

Acclimatization was carried out carefully by removing the plantlets from the glass tube, the roots were washed with running tap water to remove the adhering agar and soaked in fungicide (Mankoze 80%) for several minutes. Plantlets were transferred to polybags containing a mixture of regosol topsoil and compost from bamboo leaf litter in

1:1 ratio. Acclimatization was carried out in a greenhouse with 30-40% natural light and watered once a day for 6 months. In this experiment, 50 plantlets were transplanted one by one into polybags with plastic covers without control. The growth of plantlets was observed in the nursery after 8 months. The following formula is used to calculate the percentage of plantlets that survive after planting

Plantlets growth was observed at the nursery after 8 months. The following formula was used to calculate the percentage of plantlets that survived after being planted:

$$\frac{\text{Number of plantlets survival}}{\text{Number of plantlets transplanted}} \times 100 \quad (2)$$

A completely randomized design (CRD) with three replicates and at least twenty explants per replicate was used in the experiment. The data was analyzed using ANOVA, and the mean values were separated using Duncan's multiple range test (DMRT) at a significance level of $\alpha = 0.05$. The statistical package SPSS (Version 24) was used for analysis.

4. Results

4.1. Multiplication of *L. suaveolens*

Continuous subculture on the same medium resulted in multiple shoot regeneration in explant nodules. Figure 1 depicts the multiplication of *L. suaveolens* in vitro, these plantlets were separated and transferred to new media for rooting stage. The separation of plantlets into several new cultures will then elongate the shoots again and produce multiple nodules again. This multiplication capability is an advantage for the purpose of mass seed production because it does not occur in every species.




Figure 1. *In vitro* multiplication of *L. suaveolens*, plantlets as the source of explants from axenic culture (a), separation of the axenic plantlets into several new culture media (b), explants of multiple nodules from shoot elongation (c), shoot elongation of explants after 6 months sub-culture (d), (Credit: Putri 2018).

Figure 2 shows the sprouting ability in multiplication of *L. suaveolens* in vitro, regeneration in media with single PGRs BAP, regeneration in media with single PGRs NAA, and regeneration in media with BAP and NAA after 36 subcultures.



(a) (b) (c)

Figure 2. Sprouting ability in *in vitro* multiplication of *L. suaveolens*, regeneration in media with BAP (a), regeneration in media with NAA (b), regeneration in media with combination of BAP and NAA (c) (Credit: Putri 2018).

Table 1 shows the effect of single and multiple PGRs on the multiplication parameters of *L. suaveolens* after 6 months of *in vitro* incubation. The number of nodule multiplication coefficients (NMC) significantly different in each treatment, as well as for shoot elongation. Root elongation was not significantly different between single hormone treatments, but significantly different between single hormone and double hormone combinations. Combination of double PGRs addition were higher than single PGRs in NMC (8.4 ± 0.13), shoot length (5.91 ± 0.17 cm) and root length (8.83 ± 0.69 cm). The multiplication stage in this study obtained 100% axenic cultures, thus all cultures could be used to calculate the multiplication coefficient. According to the observations, all shoots are from axillary, not adventitious shoots. The ability to germinate after being transferred to a new medium is indeed different, some are single, and some can form multiple shoots. This ability is very advantageous at the multiplication stage. During the observation that callus growth did not occur, there may have to be additional exogenous hormones to grow callus.

Table 1. Effect of single and double PGRs on *L. suaveolens* multiplication parameter after 6 months incubation *in vitro*. Each value is a mean of twenty replicates with standard error (Mean \pm S.E). Mean with different letters (a,b,c) in each column are significantly different at 0.05 probability level by Duncan multiple range test. The results of ANOVA showed significant differences ($p \leq 0.05$) between the concentration of NAA, BAP, NAA & BAP tested for NMC, shoot length and root length.

PGRs	Nodule multiplication coefficients (NMC)	Shoot length (cm)	Root length (cm)	Leave number
BAP	6.0 \pm 0.07 ^b	4.72 \pm 0.13 ^b	5.03 \pm 0.11 ^a	30 \pm 0.01 ^a
NAA	5.1 \pm 0.09 ^a	2.21 \pm 0.09 ^a	5.10 \pm 0.09 ^a	29 \pm 0.12 ^a
BAP & NAA	8.4 \pm 0.13 ^c	5.91 \pm 0.17 ^c	8.83 \pm 0.69 ^b	32 \pm 0.07 ^a

Nodule multiplication efficiency of *L. suaveolens* *in vitro* for 6 months incubation is the initial multiplication time of the average of several plantlets. On a mass scale, this multiplication is carried out on thousands of plantlets continuously. During the first 6 months the highest yield was 8.4 nodules/explant and each nodule produced 1 to 3 shoots, so that approximately 24 new shoots were obtained per explant to become 24 new plantlets. As a comparison, the vegetative propagation of *L. suaveolens* by cuttings conducted by Kartikaningtyas et al., [23] resulted in 3 new shoots from one stem cuttings without incubation time restrictions and root formation has not occurred. Not only macro and micro vegetative propagation of *L. suaveolens* but also for *Lophostemon* sp. has not been published previously. Seedlings material is obtained from seeds, which are very limited in availability for large areas of land rehabilitation. So that the information in this study is very important to be able to continue research on a mass scale

4.2. Rooting and acclimatization of *L. suaveolens*

Figure 3 depicts root elongation after a 10-month subculture (6 months for multiplication and rooting regeneration, and another 4 months to ensure that the plantlets were healthy with well-developed roots *in vitro* prior to acclimatization), with exogenous BAP, exogenous NAA, and double PGRs.





Figure 3. Root lengthening of *L. suaveolens* after 10 months incubation (one subculture per month) with BAP (a), NAA (b) and double PGRs (c), (Credit: Putri 2018).

Figure 4 depicts the transition of a healthy *L. suaveolens* from *in vitro* regeneration to *ex vitro* development in a green house. After hardening, the rooted shoots were transferred to paper cups (Fig. 4d). The leaf color changed to dark green two weeks after the plastic cover was removed, and a wax coat and villi were observed on the leaf surface; the stems were strong (Fig. 4e). Plantlets were transplanted to pots after 6 months (Fig. 4f), with survival rates of 97%, 92%, and 94% for double PGRs, BAP, and NAA treatments, respectively (Fig. 5).






Figure 4. Acclimatization of *L. suaveolens* in greenhouse, healthy plantlet from *in vitro* regeneration (Figure 4.a), in fungicide solution (Figure 4.b) and after transplanted to the media with individual plastic cover (Figure 4.c and 4.d), tissue culture seedlings grow well after opening the plastic cover at 4 weeks of incubation (Figure 4.e and 4.f), (Credit: Putri 2018).

The percentage of plantlets that survived after single and double PGR treatments were measured 8 weeks after acclimatization (Figure 5). The plants were then kept in a greenhouse for eight months. There were no deaths in any of the treatments, and the combination of BAP and NAA treatment was the best. Plantlets care is currently limited to watering and no fertilizing.




Figure 5. The percentage of *L. suaveolens* plantlets survival with PGRs treatments 8 weeks after transplanting. Each value is a mean of twenty replicates, significantly different from each other at 0.05 probability level by Duncan multiple range test. The results of ANOVA showed a significant difference ($p \leq 0.05$) between NAA, BAP, NAA & BAP treatment for plantlets survival.

Figure 6 depicts the post-acclimatization shoot height of *L. suaveolens* seedlings treated with PGRs. The plants were then kept in a greenhouse and grew normally, with no noticeable morphological differences from stock plants. The regenerated plants showed 100% survival after 8 months of growing in the green house, with an average height of 43 ± 0.34 cm for double PGRs, 37 ± 0.21 cm for BAP, and 38 ± 0.53 cm for NAA treatments.




Figure 6. The shoots height of *L. suaveolens* plantlets in the green house. Each value is a mean of twenty replicates with standard error (Mean \pm S.E). Mean with different letters (a,b) are significantly different at 0.05 probability level by Duncan multiple range test. The results of ANOVA showed a significant difference ($p \leq 0.05$) between NAA, BAP, NAA & BAP treatment for shoot height.

5. Discussion

5.1. Disinfection culture condition

The success of the in vitro multiplication study of *L. suaveolens* is highly dependent on the effectiveness of the disinfection stage to obtain axenic culture and can maintain axenic conditions continuously until new plantlets are obtained so that the subsequent multiplication regeneration process is not disturbed by contamination. The results of this study showed that the disinfection conditions of *L. suaveolens* culture could be maintained until the plantlets were obtained in each treatment in vitro up to 100%.

In this study, we used Biocide Isothiazolones (BI) as disinfectant. BI utilizes a two-step mechanism involving rapid inhibition (minutes) of growth and metabolism, followed by irreversible cell damage resulting in loss of viability (hours). Cells are inhibited by disruption of the metabolic pathways involving dehydrogenase enzymes. Critical physiological functions are rapidly inhibited in microbes, including growth, respiration (oxygen consumption), and energy generation (ATP synthesis). In this study, without microscopic observations of contaminants were carried out, so that contamination by endophytes in the explant tissue was still possible. However, axenic culture under these conditions was able to maintain explant regeneration in an in vitro environment. The BI unique mechanism produces a broad spectrum of activity, which is expected to increase the endophytic contaminant disinfection ability, low usage rate, and difficulty in achieving resistance [26].

The disinfectant efficiency can be achieved through optimized disinfectant concentrations as well as exposure periods [27], [28] [29]. Although better disinfection can be obtained with high disinfectants concentrations with longer exposure, explant viability can be negatively affected by disinfectants under these conditions, resulting in yellow dehydrated explants along with low viability [28]. The desirable disinfection procedures should be proposed in a cheap, simple, efficient, and environmentally friendly way for eliminating the endogenous and surface contaminations [30].

5.2. Multiplication of *L. suaveolens*

Our preliminary research revealed that MS media produced the best response to the initiation of *L. suaveolens* axillary buds when compared to WPM [21]. At this early stage, no PGRs were added to test the ability of endogenous PGRs to regenerate in vitro. After 8 weeks of incubation, there was a decrease in growth, preventing the multiplication stage. Exogenous PGRs were used as an explant source in this study, and the results were as expected. Growth regulators are organic substances that plants require in small amounts to regulate plant growth [31].

The results showed that the axillary shoots of *L. suaveolens* were successfully used as a source of explants for in vitro multiplication. The increase in the number of nodules formed as a result of shoot elongation indicates the effectiveness of the hormone treatment

in this study. Moreover, the formation of multiple shoots per nodule will further increase the effectiveness and efficiency of *L. suaveolens* multiplication. In a previous study, [16] used species that can only form multiple shoots to calculate the shoot multiplication coefficient; however, *L. suaveolens* can form multiple nodules from elongation in one shoot as well as multiple shoots. The explants were placed in new culture media and incubated in a sterile environment (Figure 1.d).

At the multiplication stage, the ability to form explant shoots on single PGRs media was lower than on double PGRs media subcultures. Figure 2 depicts the high sprouting ability of *L. suaveolens* at the multiplication stage. The concentrations and combinations of PGRs and endogenous hormones in the culture medium influenced the growth qualities of micro-propagules [17],[18]. Although each species responds differently to the combination of endogenous and exogenous PGRs, BAP and NAA supplementation may cause hormonal imbalance in plants [19]. Our findings suggest that the response of *L. suaveolens* plantlets to BAP and NAA may differ between species. The ability of *L. suaveolens* to form multiple shoots (Figs. 2.a, 2b, and 2c) demonstrates that this study was successful in revealing this species' high in vitro multiplication potential to provide plants for peat rehabilitation. This is the first time this technique for *L. suaveolens* has been studied.

According to George [32][33], hormones can be found naturally in plant metabolic systems (endogenous) or synthetically (exogenously). They help to promote cell differentiation, growth, and division. BAP and NAA are the two most common classes of PGRs used in in vitro growth techniques. The type of culture specified, or regeneration type is determined by a comparison of BAP and NAA [34]. In terms of plant sensitivity to PGRs, the ratio varies by genus and species. The comparison of BAP and NAA is a determinant in meristem formation, and the balance of PGRs has been shown to be capable of stimulating the type of shoot formation. Endogenous PGR levels in plants, which influence exogenous PGR intake and also depend on the plant species.

The auxin-cytokine ratio in vitro multiplication varies depending on the type of plant used; this is critical to the success of *in vitro* plant regeneration. Single shoots can be propagated from the shoot lengthening nodules, while multiple shoots can be separated. More research is needed to determine the ability of *L. suaveolens* to respond to a combination of PGR types and concentrations on plantlet growth. PGRs influence the genetic mechanism and regulate the differentiation and multiplication processes through differential gene action, cellular events such as replication, transcription, and translation [35]. Plantlet regeneration in a variety of tree species, including *Eucalyptus nitens* [36], *Paulownia tomentosa* [37] and *Himalayan poplar* [38], has been reported using various explants and modulating auxin-cytokinin balances. BA (2.2 M) in combination with NAA (0.1 M) increased axillary budding and shoot multiplication in *Acacia mangium* and *Acacia mangium* x *Acacia auriculiformis* [39].

The addition of single BAP, single NAA, and double PGRs all resulted in significant differences in nodule multiplication coefficients and average shoot length, but with single PGRs being lower than double PGRs (Table 1). On average root length, double PGRs have a significant difference compared to single PGRs, but there is no significant difference between the three treatments on average leave number. For large-scale propagation, the multiplication coefficient number of shoots regenerated per explant/nodule of *L. suaveolens* is the most important factor to consider. The cumulative effect of PGR combinations on the percentage of viable explants, average nodule per explant, NMC, average shoot elongation, and number of leaves was observed and measured. The combination of BAP and NAA has the potential to provide a better effect than either single PGR, as shown in Table 1. BAP is the most effective growth hormone for shoot development in general. It works by inhibiting apical dominance of shoot induction and shoot bud formation.

According to several studies, media supplemented with BAP and NAA as double PGR is beneficial for shoot multiplication. One study found that MS media supplemented with 2 mg/l BA and 2 mg/l NAA yielded the best results. Another study discovered that

2 mg/l BAP resulted in the most shoot multiplication [40]. Using 2.22-M BAP, the median number of shoots produced Myrtaceae was determined. In some clones, this dose resulted in extremely high shoot numbers; explants frequently produced extensive callus and multiple short shoots. PGRs play an important role in cytodifferentiation and morphogenesis, regulating the complex processes of cell proliferation, xylogenetic differentiation, and root-shoot differentiation [41].

5.3. Rooting and acclimatization of *L. suaveolens*

The addition of exogenous PGR resulted in greater root elongation after 6 months of incubation (Figure 3). There was no discernible difference in the number of leaves among treatments. After 6 months of *in vitro* incubation, exogenous PGR had no effect on leaf regeneration. To facilitate acclimatization, the ability to successfully root regenerated shoots *in vitro* is required. This study looked at the role of different PGR compositions in root induction in sub cultured plants for four months. Full strength solid MS medium containing 0.5 ml/l BAP and 0.1 ml/l NAA (double PGRs) achieved the best *in vitro* rooting of individual shoots among the PGRs concentration treatments (Table 1). The average root length was greater in the double PGRs treatment than in the single PGR treatments (8.83 ± 0.69 cm). Healthy plantlets with well-developed roots were transferred from the culture tube after 10 months of *in vitro* rooting regeneration for *ex vitro* acclimatization in a greenhouse. In this study, 100% of the micro-cutting explants in the multiple PGRs treatment were healthy, compared to 72% in the single BAP treatment and 86% in the single NAA treatment. The high mortality during or after transfer from *in vitro* to *ex vitro* conditions is a limitation in the large-scale application of tissue culture techniques.

The results of this study show that this species is highly resistant to abiotic stress (changes in temperature, light intensity, and humidity) and biotic stress (changes in the axenic culture environment to an abundant microbial environment in soil media) during acclimatization. Because plants lack sufficient resistance to the soil microflora, their sudden exposure (particularly the root system) to microbial communities present in the soil was the leading cause of plant mortality during acclimatization [42]. Acclimatization is essential for improving plantlet survival and obtaining ready-to-plant seedlings. Direct planting of seedlings resulting from plant tissue culture into the field was not possible due to the high mortality rate of plants. *In vitro* control of humidity, temperature, light, and nutrient availability necessitates *ex vivo* adjustments. Acclimatization to the various biotic environments in soil media requires special care as well.

In terms of seedling survival of *L. suaveolens*, the combination of BAP and NAA produced the best results (Fig. 5). All seedlings can live healthy lives until the fifth week, but there is a slight decrease due to the death of some seedlings. Previous research has discovered that a variety of factors can cause seedling death during acclimatization. Because the cultured plants have nonfunctional stomata, a weak root system, and a poorly developed cuticle, transferring micro shoots to *ex vitro* conditions results in death [43]. Plantlets that received an excessive amount of PGRs developed morphological and anatomical abnormalities and were dubbed vitrified plants or hyperhydrate plants. Micro-propagated plantlets, due to their physiology and anatomy, must gradually adapt to the greenhouse or field environment [44]. Plant mortality occurred in all treatments after the eighth week, however. At this point, plantlets' care is limited to watering and no fertilization; this condition also demonstrates that a plant that can survive in waterlogged areas can also thrive in nutrient-deficient soils.

L. suaveolens has the potential to be one of the species used for revegetation and restore peatland ecosystems that adapted to wetlands and have economic value in cultivation functions. Given the importance of peatland rehabilitation, the scarcity of seedlings for millions of hectares of degraded peatlands, and the lack of protocols for *L. suaveolens* regeneration *in vitro*, this study attempted to develop an *in vitro* propagation protocol that was evaluated for faster propagation. The contribution of this research to science in-

cludes the refinement or improvement of protocol-specific in vitro multiplication techniques for the addition of PGRs NAA and BAP for *L. suaveolens* species and contribute to research originality or novelty the species which has never been done before.

In vitro multiplication is the most important step in tissue culture propagation; the technique has never been used before for *L. suaveolens*. Revegetation will ensure the sustainability of the ecosystem, protect the peat from erosion, and provide economic value to local communities living near the peatland ecosystem (Indonesian Water Portal, 2021) [45]. Myrtacea is an alternative species that has been widely developed for land rehabilitation, within one genus it is possible to have close identification physiologically and morphologically, so it is possible to do in vitro multiplication as is the case with *L. suaveolens*. The technique can be used as a support for environmental conditions that are degraded in a sustainable manner.

In general, in vitro plant multiplication techniques have been applied to the agricultural and forestry industries on a large scale, but for the purposes of planting production, they are not widely used for conservation purposes. The cost of maintaining the environment can be emphasized by mastering effective and efficient biotechnology propagation as well as mastering the multiplication technique in this study.

6. Implication

A sufficient supply of long-lasting seedlings is required for large-scale restoration. Effective seedling propagation is required to meet demand on a large scale. When wild plants are scarce due to the time lapse since the last mass flowering of many forest plant species, tissue culture of *L. suaveolens* can provide an ongoing supply of plant seedlings. Tissue culture techniques have been developed to enable large-scale production of trees while also making these techniques more cost effective. The cost of producing wildlings grows exponentially over time, whereas the cost of producing cuttings grows more slowly.

The highest NMC value is 8.4, this indicates that this species has the potential to produce 8 nodules per shoot. Multiple shoots will produce a minimum of 10 shoots per plantlet, then a plantlet will produce 80 plants. If a 10% failure occurs, about 70 plants ready for planting will be obtained. In plantation with 2×3 m of spacing, then only requires about 17 plantlets per hectare. Thus, the *in vitro* multiplication technique of *L. suaveolens* propagation will be more efficient for plantation.

7. Conclusions

This research has a strategic value to obtain techniques for providing plantlets or seedlings on marginal lands that are difficult to manage. As an alternative to mass propagation of seedlings for peatland rehabilitation, we developed an in vitro multiplication technique of *L. suaveolens*, a water-resistant tree species. This study is a new high-efficiency innovation technique with a mutually reinforcing effect on seedling growth. Increasing the multiplication coefficient in vitro by combining the PGRs BAP and NAA. The availability of balanced nutrients and PGRs during in vitro multiplication resulted in improved plantlets growth after acclimatization and ready for field planting. This protocol can be used to spread *L. suaveolens* on a large scale. This protocol will need to be tested on a larger scale of cultivation.

Acknowledgments The authors gratefully acknowledge the Research and Development of PT. Arara Abadi in Perawang, Riau for providing the plant materials. We wish to thank to Ministry of Environment and Forestry of Indonesia for fully supporting this study.

Author Contributions: Conceptualization, A.I.P., L.H and T.H.; methodology, A.I.P.; software, A.I.P.; validation, A.I.P., T.H., A.N., S.S., and L.H.; formal analysis, A.I.P., A.R.; investigation, A.I.P., P.B.S., R.S.W., and F.L.; resources, A.I.P.; data curation, L.H.; writing—original draft preparation, A.I.P., N.K.K.; writing—review and editing, A.I.P., N.K.K.; visualization, A.I.P.; supervision, A.I.P., N.K.K., P.B.S.; project administration, A.I.P.; funding acquisition. A.I.P., N.K.K., A.N., S.S., L.H.,

T.H., P.B.S., R.S.W., F.L., and A.R. had an equal role as main contributors in discussing the conceptual ideas and the outline, providing critical feedback for each section and writing the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: No Funding

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest the funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Reference

- [1] Agus, F.; Anda, M.; Jamil, A., *LAHAN GAMBUT INDONESIA Pembentukan, Karakteristik, dan Potensi Mendukung Ketahanan Pangan*, Edisi rev. IAARD Press, 2016.
- [2] M. A. Cochrane, *Fire in the tropics*. In: *Tropical Fire Ecology*. Springer, Berlin, Heidelberg, 2009.
- [3] J. Page, S; Siegert, F; Rieley, "The Amount of Carbon Released from Peat and Forest Fires in Indonesia During 1997," *Nature*, vol. 420, no. 7 November 2002, pp. 61–65, 2002.
- [4] H. Zhang, W., Lu, Q., Song, K., Qu, G., Wang, Y., Wang, X., Li, H., Li, J., Liy, G., Li, "Remotely sensing the ecological influences of ditches in Zoige Peatland, eastern Tibetan Plateau," *Int. J. Remote Sens.*, vol. 35, no. 13, 2014.
- [5] Valjarević, A., Djekić, T., Stevanović, V., Ivanović, R., Jandziković, B., "GIS numerical and remote sensing analyses of forest changes in the Toplica region for the period of 1953–2013," *Appl. Geogr.*, vol. 92, pp. 131–139, 2018.
- [6] D. M. . Clark, J.S.; Bell, M. C. . Hersh, M. H.; Kwit, E. Moran, and Salk, C.; Stine, A; Valle, D.; Zhu, K., "Individual-scale variation, species-scale differences: inference needed to understand diversity," *Ecol. Lett.*, vol. 14, pp. 1273–1287, 2011.
- [7] H. Anzaldúa, G. and Gerdes, *Restoring Peatlands and Applying Concepts for Sustainable Management in Belarus-An analysis of project implementation and cost-effectiveness*. Gerardo Anzaldúa and Holger Gerdes (Ecologic Institute, 2011).
- [8] H. Jauhiainen, J; Silvennoinen, H; Mari, Könönen, M; Limin, S; & Vasander, "Management driven changes in carbon mineralization dynamics of tropical peat," *Biogeochemistry*, vol. 129, pp. 115–132, 2016.
- [9] F. Jaenice, J; Lohberger, S; Siegert, "Monitoring the effect of restoration measures in Indonesian peatlands by radar satellite imagery," *J. Environ. Manage.*, vol. 92, no. 3, pp. 630–638, 2010.
- [10] M. Wilson, PG; and Heslewood, "Phylogenetic position of *Meteoromyrtus* (Myrtaceae)," *Telopea, J. Plant Syst.*, vol. 19, pp. 45–55, 2016.
- [11] Y. Mangera, "ANALISIS VEGETASI JENIS POHON DI KAWASAN HUTAN KAMPUNG WASUR PADA TAMAN NASIONAL WASUR DISTRIK MERAUKE KABUPATEN MERAUKE," *J. Agric.*, vol. 1 (1), no. Mei, pp. 18–36, 2008.
- [12] S. Von Arnold and T. Eriksson, "In vitro studies of adventitious shoot formation in *Pinus contorta*," *Can. J. Bot.*, vol. 59, no. 5, pp. 870–874., 1981.
- [13] T. Khan, M. A. Khan, K. Karam, N. Ullah, Z. U. R. Mashwani, and A. Nadhman, "Plant in vitro Culture Technologies; A Promise Into Factories of Secondary Metabolites Against COVID-19," *Frontiers in Plant Science*, vol. 12. Frontiers Media S.A., Mar-2021.
- [14] P. D.-F. . T. R. R. Souza, R. A.; Dantas, P. V. P.; Cavalcante, "Basic procedure for the in vitro propagation of Brazilian trees for reforestation purposes," *J. Environ. Anal. Prog.*, vol. 2, no. 2, pp. 1–8, 2017.
- [15] Henry G. B; Ross F. B; & Andrew VB;, "The impact of phytohormone concentration in *Moringa oleifera* leaf extract on wheat yield and components of yield," *Publ. Cover 2019 Impact Factor 1.132 J. Plant Nutr.*, vol. 43, no. 3, 2020.
- [16] Müller, D., Leyser, O., "Auxin, cytokinin and the control of shoot branching," *Ann. Bot.*, vol. 107, no. 7, pp. 1203–1212, 2012.
- [17] Sauer, M., Robert, S., Kleine-Vehn, J., "Auxin: simply complicated," *J. Exp. Bot.*, vol. 64, no. 9, pp. 2565–2577, 2013.
- [18] Z. Ying-Hua, S; Yu-Bo, L; Xian-Sheng, "Auxin–Cytokinin Interaction Regulates Meristem Development," *Mol. Plant*, vol. 4, no. 4, 2011.
- [19] Lee, Y.C., Chang, J.C., "Development of an Improved Micropropagation Protocol for Red-Fleshed Pitaya 'Da Hong' with and without Activated Charcoal and Plant Growth Regulator Combinations," *Horticulturae*, vol. 8, p. 104, 2022.
- [20] Bhojwani, S. S., Dantu, P. K., "Micropropagation," in *Plant Tissue Culture: An Introductory Text*, Bhojwani, S.S., Dantu, P.K., Eds., Ed. New Delhi, India: Springer, 2013, pp. 245–274.
- [21] A. I. P. Putri, "Not published," 2018.
- [22] Hesami, M., Naderi, R., Tohidfar, M., "Modeling and Optimizing in vitro Sterilization of Chrysanthemum via Multilayer Perceptron-Non-dominated Sorting Genetic Algorithm-II (MLP-NSGAI)," *Front. Plant Sci. Plant Sci.*, vol. 10, pp. 1–13, 2019.
- [23] B. G. Tabachnick and L. S. Fidell, *Experimental Designs Using ANOVA*, vol. 15, no. 2. Belmont-California, USA: Duxbury Press, Belmont, CA., 2020.
- [24] A. E. Permasari, D. R. A. Rambli, and P. D. D. Dominic, "Forecasting method selection using ANOVA and Duncan multiple range tests on time series dataset," in *Proceedings 2010 International Symposium on Information Technology - Engineering Technology, ITSIM'10*, 2010, vol. 2, pp. 941–945.

- [25] Artikaningtyas, D., Setiyaji, T., Surip, N. F. N., "The Effectivity of Planting Media and Fertilizer in Sprouting Ability of Kess (Lophostemon suaveolens (Sol.ex Gaertn.) Peter G.Wilson & J.T. Waterh) Stool Plant," *J. Perbenihan Tanam. Hutan*, vol. 7, pp. 67–76, 2019.
- [26] Williams, T. M., "The Mechanism of Action of Isothiazolone Biocide," CORROSION 2006, San Diego, California, NACE-06090, 2006.
- [27] Altan, F., Burun, B., Sahin, N., "Fungal contaminants observed during micropropagation of Lilium candidum L. and the effect of chemotherapeutic substances applied after sterilization," *African J. Biotechnol.*, vol. 7, 2010.
- [28] Valente, T. A. M., Silva, D. M., Gomes, P. S., Fernandes, M. H., Santos, J. D., Sencadas, V., "Effect of Sterilization Methods on Electrospun Poly(lactic acid) (PLA) Fiber Alignment for Biomedical Applications," *ACS Appl. Mater. Interfaces*, vol. 8, no. 5, pp. 3241–3249, 2014.
- [29] Hesami, M., Naderi, R., Yoosefzadeh-N. M., "Optimizing sterilization conditions and growth regulator effects on in vitro shoot regeneration through direct organogenesis in Chenopodium quinoa," *J. Biotechnol. Comput. Biol. Bionanotechnol.*, vol. 99, no. 1, pp. 49–57, 2018.
- [30] Blin, T., Purohit, V., Leprince, J., Jouenne, T., Glinel, K., "Bactericidal Microparticles Decorated by an Antimicrobial Peptide for the Easy Disinfection of Sensitive Aqueous Solutions," *Biomacromolecules*, vol. 12, no. 4, pp. 1259–1264, 2011.
- [31] Shen, X., Kane, M.E., Chen, J., "Effects of genotype, explant source, and plant growth regulators on indirect shoot organogenesis in Dieffenbachia cultivars," *Vitr. Cellul. Dev. Biol. - Plant*, vol. 44, pp. 282–288, 2008.
- [32] George, E.F., Hall, M.A., Klerk, G-J De., "Plant Growth Regulators I: Introduction; Auxins, their Analogues and Inhibitors," in *Plant Propagation by Tissue Culture*, 2008, pp. 175–204.
- [33] G.-J. D. George, E.F., Hall, M.A., Klerk, "Plant Growth Regulators II: Cytokinins, their Analogues and Antagonists," in *Plant Propagation by Tissue Culture*, 2008, pp. 205–226.
- [34] E. B. Herman, *Recent Advances in Plant Tissue Culture*. Agritech Consultants, Inc., Shrub Oak, 2006.
- [35] T. A. Thorpe, "History of Plant Tissue Culture," *Mol. Biotechnol.*, vol. 37, no. 2, pp. 169–180, 2007.
- [36] J. D. Bandyopadhyay, S. Hamil, "Ultrastructural Studies of Somatic Embryos of Eucalyptus nitens and Comparisons with Zygotic Embryos Found in Mature Seeds," *Ann. Bot.*, vol. 86, no. 2, 2000.
- [37] Carmen, S.J., Jose, C.M., Corredoira, E., "Histology of the regeneration of Paulownia tomentosa (Paulowniaceae) by organogenesis," *Rev. Biol. Trop.*, vol. 62, no. 2, pp. 809–818, 2014.
- [38] Anggarwal, G., Gaur, A., Srivastava, D.K., "Establishment of high frequency shoot regeneration system in Himalayan poplar (Populus ciliata Wall. ex Royle) from petiole explants using Thidiazuron cytokinin as plant growth regulator," *J. For. Res.*, vol. 26, no. 3, 2015.
- [39] Moteuuis, O., Galiana, A., Goh, D., "In Vitro Propagation of Acacia mangium and A. mangium × A. auriculiformis," <https://www.researchgate.net/journal/Methods-in-molecular-biology-Clifton-NJ-1064-3745>, vol. 11013, pp. 199–211, 2013.
- [40] F. Ibáñez, CM; Camus, PA; Rocha, "Diversity and distribution of cephalopod species off the coast of Chile," *Mar. Biol. Res.*, vol. 5, no. 4, p. 374, 2019.
- [41] Darby, D., Bai, S.H., Wallace, H. M., Truemen, S.J., "Micropropagation of the therapeutic-honey plants Leptospermum polygalifolium and L. scoparium (Myrtaceae)," *Aust. J. Bot.*, vol. 69, no. 5, pp. 310–317, 2021.
- [42] K. Lebedev, V; Schestibratov, "Effect of natural and synthetic growth stimulators on in vitro rooting and acclimatization of common ash (Fraxinus excelsior L.) microplants," *Nat. Sci.*, vol. 5, no. 10, 2013.
- [43] R. F. B. A. V. Brockman, H. G; Ross F. Brennan, "The impact of phytohormone concentration in Moringa oleifera leaf extract on wheat yield and components of yield," *J. Plant Nutr.*, vol. 43, no. 3, 2020.
- [44] S. Mathur, A; Mathur, A; Verma, P; Yadaf, "Biological hardening and genetic fidelity testing of micro-cloned progeny of Chlorophytum borivilianum Sant. et Fernand.," *African J. Biotechnol.*, vol. 7, no. 8, pp. 1046–1053, 2008.
- [45] Indonesia Water Portal, Sustainability Times., "A peatland restoration project in Indonesia could set a global example," 2021.