

Article

Wildlife Hosts of *Leishmania Infantum* in a Re-Emerging Focus of Human Leishmaniasis, in Emilia-Romagna, Northeast Italy

Roberta Taddei^{1,†}, Arianna Bregoli^{1,†}, Giorgio Galletti¹, Elena Carra¹, Laura Fiorentini¹, Maria Cristina Fontana¹, Matteo Frasnelli¹, Carmela Musto², Giovanni Pupillo¹, Alessandro Reggiani¹, Annalisa Santi¹, Arianna Rossi¹, Marco Tamba¹, Mattia Calzolari^{1,‡}, Gianluca Rugna^{1,‡,*}

¹ Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna, via A. Bianchi 9, 25124 Brescia, Italy

² Department of Veterinary Medical Sciences – University of Bologna, Via Tolara di Sopra 50, 40064 Ozzano Emilia, Italy

† These authors contributed equally to this work.

‡ These authors also contributed equally to this work.

* Correspondence: gianluca.rugna@izsler.it

Abstract: In the last decade, an upsurge of human leishmaniasis has been reported in the Emilia-Romagna region, Northeast Italy. Epidemiologic data raised doubts about the role of dogs as the main reservoirs for *Leishmania infantum*. In the present study, a total of 1,077 wild animals were screened for *L. infantum* DNA in earlobe and spleen samples from 2019 to 2022. The lymph nodes were tested only in the 23 animals already positive in the earlobe and/or spleen. A total of 71 (6.6%) animals resulted positive in at least one of the sampled tissues, including 3/18 (16.7%) wolves, 6/39 (15.4%) European hares, 38/309 (12.3%) roe deer, 1/11 (9.1%) red deer, 8 (4.9%) wild boars, 13/319 (4.1%) red foxes, 1/54 (1.9%) porcupine, and 1/59 (1.7%) European badger. Most of the infected animals (62/71) tested positive only in the earlobe tissue, just 4 animals (2 roe deer and 2 wild boars) tested positive only in the spleen, and 5 animals (3 roe deer and 2 red foxes) resulted positive for both tissues. *L. infantum* DNA was detected in the lymph nodes of 6/23 animals. *L. infantum* detection occurred in all seasons associated with low real-time PCR Ct values. Further research is needed in order to clarify the role of wildlife in the re-emerging focus of leishmaniasis in Northeast Italy.

Keywords: *Leishmania infantum*; reservoir; wildlife; artiodactyls; roe deer; European hare; red fox; wild boar

1. Introduction

Leishmaniasis caused by the protozoan *Leishmania (L.) infantum* is a neglected vector-borne zoonotic disease that is endemic in the Mediterranean basin [1,2]. *L. infantum* is the causative agent of zoonotic visceral (VL), cutaneous (CL) and mucosal (ML) leishmaniasis in humans [3,4]. It is also the cause of canine leishmaniasis (CanL) in dogs, which are considered the major domestic reservoir of infection for people [5]. In the Old World, transmission occurs via the bite of female sand flies of the genus *Phlebotomus* [1].

Italy is endemic to *L. infantum*, representing the country with the highest CanL seroprevalence in Europe [6]. The classical endemic areas are the rural and hilly peri-urban zones along the Tyrrhenian littoral, the southern peninsular regions, and the islands. Since the 1990s, the incidence of leishmaniasis in both dogs and humans has been on the rise in northern areas, previously regarded as non-endemic [7]. Among the northern Italian regions, the Emilia-Romagna region (RER) is characterized by a distinctive epidemiological situation: since 1934, CL has been frequently reported -with an incidence exceeding 2,000 cases in the 1950s- before declining, most likely as a result of pesticide use in agriculture [8]. In contrast, until the early 1970s, there were just 4 recorded autochthonous cases of VL. In 1971–1972, a severe VL outbreak with 60 cases occurred in the foothill areas close to Bologna municipality, which is located in the central RER [9]. This outbreak was

regarded as atypical, as only human cases were diagnosed but a canine reservoir was not clearly evidenced [9]. Furthermore, follow-up surveillance of this VL focus failed to reveal human and canine cases over the next 15 years [10].

Since 2012, an upsurge of VL cases has been described in RER [11,12]. Similar to the VL outbreak in the 1970s, no rise in *L. infantum* infection among dogs was reported, which motivated research to better understand the regional *L. infantum* epidemiology. Molecular typing studies showed that *L. infantum* strains circulating in dogs belonged to a distinct population in comparison with strains circulating in VL cases and sand flies [13,14]. Moreover, the blood-meal analysis of *Phlebotomus* (*Ph.*) *perfiliewi* – the strongly suspected vector of *L. infantum* in RER [15] – showed a biting preference for wild animals and humans [16]. These findings led to hypothesize a role for an unknown wildlife species as a reservoir for the parasite, also according to the increasing number of studies reporting infection of *Leishmania* spp. in wild animals [17].

The aim of the present study was to evaluate the role of different wild mammals as natural hosts of *L. infantum* in RER by testing them for *L. infantum* DNA over a 3-year period.

2. Materials and Methods

2.1. Study area

RER is located in the northeastern part of the Italian peninsula. It covers an area of approximately 22,500 km² and is inhabited by a population of 4.5 million people. The region is geographically divided into two homogeneous areas: the northern half of the region is entirely occupied by the Po Valley, while in the southern half, the flat area gives way to the hilly and then to the mountainous part of the Apennine chain. On its eastern side, it borders the Adriatic Sea. The climate is sub-continental in the inner part of the region and becomes Mediterranean near the coasts. The average temperature from 1991 to 2015 was 12,8 °C. The rainfall ranges from 650 to 1,200 mm per year [18], depending on the altitude and the distance from the sea: the minimum value is recorded in the plains and increases towards the hills and the mountains. The rains are concentrated mainly in autumn and spring; the summer is often characterized by severe dryness.

RER is divided into 9 administrative provinces; this study included the warm-blooded fauna of the 6 provinces most affected by human leishmaniasis in the last decade: Reggio Emilia, Modena, Bologna, Ravenna, Forlì-Cesena, and Rimini [19]. This territory comprises the central-eastern part of the region, characterized by flat, hilly, and mountainous areas as well as the Adriatic coast.

2.2. Sampling

From February 2019 to March 2022, a total of 1,077 wild mammal carcasses were collected as part of the wildlife health surveillance program set up in RER [20].

They consisted of: 508 artiodactyls, including 309 roe deer (*Capreolus capreolus*), 164 wild boars (*Sus scrofa*), 24 fallow deer (*Dama dama*), 11 red deer (*Cervus elaphus*); 401 carnivores, including 319 red foxes (*Vulpes vulpes*), 59 European badgers (*Meles meles*); 18 wolves (*Canis lupus italicus*), 2 stone martens (*Martes foina*), 1 pine marten (*Martes martes*), 1 raccoon (*Procyon lotor*), 1 Western polecat (*Mustela putorius*); 67 rodents, including 54 porcupines (*Hystrix cristata*), 7 squirrels (*Sciurus vulgaris*), 5 dormice (*Glis glis*), 1 common dormouse (*Muscardinus avellanarius*); 62 hedgehogs (*Erinaceus europaeus*) belonging to the Eulipotyphla order; 39 lagomorphs European hares (*Lepus europaeus*).

The animals were legally hunted, found dead by local authorities, accidentally road-killed, or conferred by wildlife rescue and rehabilitation centres.

Post-mortem examination and tissue sampling were conducted at the Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia-Romagna. The presence of pathological alterations in the organs was recorded. Spleen and earlobe samples were collected from all animals included in the study. Superficial lymph nodes of the head were

sampled in a limited number of carcasses and tested only if the spleen and/or earlobe resulted positive.

All the sampled tissues were stored frozen (-20°C) until molecular analysis.

2.3. Molecular analysis

Genomic DNA was extracted from 25 mg sample of the spleen, lymph node, and the central hairless part of the earlobe. Tissue samples were homogenized and incubated overnight at 56° C in 200 µL buffered lysis solution containing proteinase K (10 µg/mL). DNA was extracted using the NucleoSpin Tissue Kit (Macherey Nagel, Duren, Germany) according to the manufacturer’s instructions and eluted in 200 µL of elution buffer. Each sample was co-extracted and amplified with a commercial internal control DNA template (QuantiFast Pathogen PCR +IC Kit, Qiagen GmbH, Hilden, Germany). The detection of *Leishmania* DNA was carried out by a TaqMan MGB probe real-time PCR targeting a highly repetitive kinetoplast minicircle DNA sequence [21]. Each amplification was performed in 25 µL of reaction mixture with 5 µL of sample DNA on BioRad CFX96 (Bio-Rad Laboratories, Hercules, CA, USA). The PCR was considered positive when a threshold cycle (Ct) value lower than 40 was detected.

2.4. Data analysis

Data were summarized with the proportion of positive samples, with exact confidence interval [22]. Analysis was performed with R 4.0.2 [23]. The spatial distribution of samples was provided with QGis (version 3.14.15-Pi).

3. Results

A total of 2,177 tissue samples were analysed for the presence of *L. infantum* DNA: spleen and earlobe samples from each of the 1,077 animals included in the study, and the lymph nodes of 23 animals found positive in the earlobe and/or spleen.

The overall species and tissue-specific PCR detection is summarized in Table 1. A total of 71 (6.6%) animals resulted positive in at least one of the sampled tissues, including 3/18 (16.7%) wolves, 6/39 (15.4%) European hares, 38/309 (12.3%) roe deer, 1/11 (9.1%) red deer, 8/164 (4.9%) wild boars, 13/319 (4.1%) red foxes, 1/54 (1.9%) porcupine, and 1/59 (1.7%) European badger.

Table 1. *L. infantum* real-time PCR positive wildlife species in the Emilia-Romagna region in the period 2019-2022. Detection rates (including 95% confidence interval) for the target species and tissues are reported.

Order	Species	Earlobe		Spleen		Total	
		positive (tested)	prevalence (IC95%)	positive (tested)	prevalence (CI 95%)	positive (tested)	prevalence (CI 95%)
Artiodactyla	Roe deer	36 (309)	11.7 (8.3 – 15.8)	5 (309)	1.6 (0.5 – 3.7)	38 (309)	12.3 (8.9 – 16.5)
	Red deer	1 (11)	9.1 (0.2 – 41.3)	0 (11)	0.0 (0.0 – 28.5)	1 (11)	9.1 (0.2 – 41.3)
	Wild boar	6 (164)	3.7 (1.4 – 7.8)	2 (164)	1.2 (0.2 – 4.3)	8 (164)	4.9 (2.1 – 9.4)
Carnivora	Wolf	3 (18)	16.7 (3.6 – 41.4)	0 (18)	0.0 (0.0 – 18.5)	3 (18)	16.7 (3.6 – 41.4)
	Red fox	13 (319)	4.1 (2.2 – 6.9)	2 (319)	0.6 (0.1 – 2.3)	13 (319)	4.1 (2.2 – 6.9)

	European badger	1 (59)	1.7 (0.0 – 9.1)	0 (59)	0.0 (0.0 – 6.1)	1 (59)	1.7 (0.0 – 9.1)
Lagomorpha	European hare	6 (39)	15.4 (5.9 – 30.5)	0 (39)	0.0 (0.0 – 9.0)	6 (39)	15.4 (5.9 – 30.6)
Rodentia	Porcupine	1 (54)	1.9 (0.1 – 9.9)	0 (54)	0.0 (0.0 – 6.6)	1 (54)	1.9 (0.1 – 9.9)
Other species ¹		0 (104)	--	0 (104)	--	0 (104)	--
<i>All Species</i>		67 (1077)	6.2 (4.9 – 7.8)	9 (1077)	0.8 (0.4 – 1.6)	71 (1077)	6.6 (5.2 – 8.2)

¹species with no positive individuals, namely 62 hedgehogs, 24 fallow deer, 2 stone martens, 1 pine marten, 1 raccoon, 1 Western polecat, 7 squirrels, 5 dormice, 1 common dormouse.

In relation to the tissue location of the parasite, five animals (3 roe deer and 2 foxes) tested positive in both the spleen and earlobe, 4 animals (2 roe deer and 2 wild boars) tested positive only in the spleen, while the large majority of the positive animals (62/71) tested positive only in the earlobe. Earlobe samples showed the highest *L. infantum* DNA detection rate (6.2%; CI95%: 4.9-7.8) compared to spleen samples (0.8%; CI95%: 0.4-1.6). The superficial lymph nodes were tested on a subgroup of 23 animals already positive for the ear (19), spleen (1), or both (3). Among them, 6 animals tested positive and 1 roe deer resulted positive in all the three organs (earlobe, spleen and lymph nodes) (Tab. 2).

Table 2. Lymph node analysis on 23 animals already tested positive for *L. infantum* DNA in the earlobe and/or spleen.

Order	Species	No. of lymph node positive animals (No. tested)			
		earlobe	spleen	both	total
Artiodactyla	Roe deer	2 (10)	0 (1)	1 (2)	3 (13)
	Red deer	1 (1)			1 (1)
	Wild boar	0 (1)			0 (1)
Carnivora	Wolf	1 (3)			1 (3)
	Red fox	1 (3)		0 (1)	1 (4)
	European badger	0 (1)			0 (1)
	total	5 (19)	0 (1)	1 (3)	6 (23)

None of the positive animals showed macroscopic dermal and/or internal lesions suggestive of *Leishmania* infection, with the exception of one roe deer with a localized ulcerative lesion located on the positive earlobe (further investigations are underway).

The PCR Ct values ranged from 13.9 to 39.1; 15.9% of the samples had Ct values lower than 25 and 17.1% of the samples had Ct values greater than 35 (Fig. 1). Overall, median Ct values were lower in the earlobe than in spleen and lymph node samples, with results of 29.0, 33.9, and 34.8, respectively. In particular, the median Ct values were less than 35 only in the earlobes of roe deer, red deer, red foxes, European hares, and

porcupines. In addition, Ct values resulted less than 20 in 4 earlobe samples (2 from roe deer and 2 from red foxes), always exceeding 30 in all spleen samples.




Figure 1. Distribution of *L. infantum* real-time PCR cycle threshold (Ct) values for the target species and tissues.

The spatial distribution of both the sampled and positive animals is shown in Figure 2. The highest detection rates were concentrated in the provinces of Forlì-Cesena (11.7%) and Bologna (8.7%) (Fig. 2 and Tab. 3). At a province level, the highest number of positive species was also recorded in the Bologna and Forlì-Cesena provinces, with 5 and 4 species, respectively (Tab. S1).




Figure 2. Sampled and *L. infantum* real-time PCR positive areas, Emilia-Romagna region, Italy.

Table 3. Overall prevalence (including 95% confidence interval) for *L. infantum* DNA in wildlife in the provinces of the Emilia-Romagna region.

Province	No. of positive (tested animals)	Prevalence (%)	CI 95%
Bologna	27 (309)	8.7	(5.8 - 12.5)
Forlì-Cesena	22 (188)	11.7	(7.5 - 17.2)
Modena	11 (250)	4.4	(2.2 - 7.7)
Ravenna	3 (40)	7.5	(1.6 - 20.4)
Reggio Emilia	1 (162)	0.6	(0.0 - 3.4)
Rimini	7 (128)	5.5	(2.2 - 10.9)

CI: confidence interval.

The yearly temporal distribution of *Leishmania* DNA detection showed that the animals tested positive in all seasons and Ct values lower than 25 (from 13.9 to 21.6) were recorded throughout the whole year (Fig. 3). Roe deer tested positive in all seasons (Tab. 4), with minimum associated Ct values ranging from 13.9 to 26.3. Overall, one roe deer, positive in both the earlobe and spleen, one badger, and one wild boar tested positive for *L. infantum* DNA during spring (Tab. 4).




Figure 3. Distribution of *L. infantum* real-time PCR cycle threshold (Ct) values among the weather seasons.

Table 4. Prevalence (including 95% confidence interval) of *L. infantum* real-time PCR positive species along the weather seasons.

		Season											
Order	Species	spring			summer			autumn			winter		
		N° positive	Prevalence	95% C.I.	N° positive	Prevalence	95% C.I.	N° positive	Prevalence	95% C.I.	N° positive	Prevalence	95% C.I.

		(n° teste d)			(n° teste d)			(n° teste d)			(n° teste d)		
Artio- dactyla	Red deer							1(6)	16.7	(0.4- 64.1)	0(5)	0	(0- 52.2)
	Roe deer	1(75)	1.3	(0-7.2)	6(53)	11.3	(4.3- 23)	13(6 4)	20.3	(11.3- 32.2)	18(1 17)	15.4	(9.4- 23.2)
	Wild boar	1(10)	10	(0.3- 44.5)	0(21)	0	(0- 16.1)	3(77)	3.9	(0.8- 11)	4(56)	7.1	(2- 17.3)
Car- nivora	European badger	1(6)	16.7	(0.4- 64.1)	0(10)	0	(0- 30.8)	0(21)	0	(0- 16.1)	0(22)	0	(0- 15.4)
	Red fox	0(4)	0	(0- 60.2)	5(15 7)	3.2	(1- 7.3)	4(64)	6.3	(1.7- 15.2)	4(94)	4.3	(1.2- 10.5)
	Wolf				0(1)	0	(0- 97.5)	0(6)	0	(0- 45.9)	3(11)	27.3	(6-61)
Lago- morpha	European hare	0(7)	0	(0-41)	0(4)	0	(0- 60.2)	1(16)	6.3	(0.2- 30.2)	5(12)	41.7	(15.2- 72.3)
	Porcupine	0(8)	0	(0- 36.9)	0(13)	0	(0- 24.7)	0(17)	0	(0- 19.5)	1(16)	6.3	(0.2- 30.2)
<i>All species</i>		3(11 3)	2.7	(0.6- 7.6)	11(2 88)	3.8	(1.9- 6.7)	22(3 29)	6.7	(4.2- 10.0)	35(3 47)	10.1	(7.1- 13.8)

C.I.: confidence interval.

4. Discussion

L. infantum is a zoonotic parasite steadily endemic at a low prevalence (around 2%) in the canine population in RER [19,24]. In contrast, an increasing number of VL, CL, and ML cases have been reported in the region in the last decade [3,12,25,26]. Molecular typing of strains from canine and human hosts and from sand flies, raised doubts about the dogs' role as main reservoirs to human infection in this region [13,14]. Moreover, studies on the blood-meals of *Ph. perfiliewi* – the most abundant sand fly species in the study area – showed a feeding preference for wild mammals (especially roe deer and European hares) and humans, while no trace of canine or rodents' blood was observed [16]. In addition, high infection rates in *L. infantum* naturally infected sand flies were found, suggesting the existence of a dog independent sylvatic cycle [15,16].

These findings led to the investigation of the potential role of wild mammals as natural reservoirs of *L. infantum* in the region. Different species of mammals have been found positive for *L. infantum* DNA in the study area, with an overall prevalence of 6.6% (CI95%: 5.2-8.2), ranging among positives from 1.7% to 16.7% according to species. The overall highest positivity rates in wildlife were observed in the foothill areas of the Forlì-Cesena and Bologna provinces (Fig. 1), which are the areas more affected by human leishmaniasis [19] and where a high density of sand flies has been historically reported [27,28]. Even though the results could be skewed by the non-homogeneous sample across provinces, the largest number of infected species was also detected in these provinces, perhaps as a result of a higher infection pressure.

Among the wild carnivores sampled in the present study, the highest *Leishmania* DNA detection rate was observed in wolves (16.7%; CI 95%: 3.6-41.4). This value was lower than those reported from other European countries, including other northern Italian regions, where prevalence rates of up to 50% were observed [29-31]. Also, the prevalence

found in foxes (4.1%; CI 95%: 2.2-6.9), ranging from 3.6% to 7.2% according to the sampling areas [Tab. S1], was lower than that reported previously in northwestern (12.3%), central (52.2%), and southern Italy (28.6%-40%) [31-34], and in Spain (14.1%) [35,36]. Similarly, a low *L. infantum* detection rate was found in European badgers (1.7%), when compared with the high values observed in northern Spain (26%) and northwestern Italy (53.3%) [31,37].

Among rodents, porcupines were the unique species found positive for *L. infantum* DNA with a detection rate of 1.9% (CI 95%: 0.1-9.9). To our best knowledge, there is only a single paper reporting porcupines of the genus *Hystrix* as possible wild reservoir species for visceral leishmaniasis [38]. The small sample size of other species prevented a real assessment of the current prevalence in rodents in the study area. Recently, anthroponophilic rodents have been found to be infected at a prevalence of 11% in plain areas of RER [39], which are historically less involved in human leishmaniasis than the foothill areas. However, the increase in seasonal abundance and the distribution of sand flies due to climate change, together with the recent detection of *Leishmania* DNA from lowland sand flies [40], indicate that the role of rodents in the epidemiology of leishmaniasis in RER needs to be monitored and further investigated.

Lagomorphs such as European hares were found infected in the study area with a detection rate of 15.4% (CI 95%: 5.9-30.6). Even though this value was lower than those reported from other Mediterranean countries, such as Spain (43.6%) and northern Greece (23.5%) [41,42], it is worth noting that all the positive subjects (6/13, 46%) were sampled in the Bologna province (Tab. S1), which is the most ancient and re-emerging focus of VL in RER [11]. In addition to black rats (*Rattus rattus*) in Italy [43], infectiousness to sand flies has been demonstrated in the Iberian hare (*Lepus granatensis*), which was suggested to be the main sylvatic reservoir in the human leishmaniasis outbreak occurred in Fuenlabrada, Spain [41,44].

Although rodents, carnivores, and lagomorphs are the orders more widely explored in leishmaniasis foci, being the most suitable wild reservoirs for the parasite [45,46], over the last decade, the list of wild animals infected with *L. infantum* has been enlarged with bats, primates, and different species of hedgehog [17,47-49]. Interestingly, *L. infantum* has been recently detected or isolated from domestic species belonging to the families Equidae and Bovidae, respectively [50,51]. However, wild artiodactyls have been only rarely investigated.

The present study provided convincing evidence for *L. infantum* DNA detection in wild boar (*Sus scrofa*) and in other two artiodactyls belonging to the family Cervidae, namely roe deer (*Capreolus capreolus*) and red deer (*Cervus elaphus*).

Moraes-Silva and colleagues [52], by combining serologic and molecular methods as well as experimental infection, concluded that domestic swine (*Sus scrofa*) is resistant to infection by *L. infantum*, excluding the possible involvement of this species in the epidemiology of visceral leishmaniasis. However, the same authors commented on a previous study [53] showing the presence of numerous amastigotes of an uncharacterized *Leishmania* species, probably *L. braziliensis*, in one pig. In the present study, wild boars showed a median Ct value of 35.9, with the lowest value above 30, probably indicating a natural resistance to the disease and a transient infection. Nevertheless, the detection of an infected animal in the spring indicates that additional research is required to clarify the role of wild boar.

Data generated in the present study suggest the need for an in-depth analysis of the role of cervids. The detection rate of *Leishmania* DNA in roe deer over the entire study area was 12.3% (95% CI: 8.9-16.5). Considering only the areas most affected by human leishmaniasis, it was found to be 9.4% in the province of Bologna, while the highest infection rate was detected in the province of Forlì-Cesena with 31.8% of infected roe deer. In addition, the roe deer has been shown to be the preferred host species by *Ph. perfiliewi* in selected endemic sites in the RER region, according to blood-meal identification. In the same study, a *Ph. perfiliewi* specimen was found engorged with a double meal, i.e., roe deer and man, and it was infected by a high load of *L. infantum* as well [16].

Absence of pathological alterations has been a common finding among wild animals infected by *Leishmania* spp., but in a study carried out in Spain, almost half the animals tested exhibited at least one lesion compatible with leishmaniasis in their earlobe [54]. In the present study, the earlobe tissue of a single roe deer showed an ulcerated area with an inflammatory histologic pattern typical of cutaneous leishmaniasis (data not shown). Further investigation is ongoing at the time of writing this paper.

In this study, three different tissues were targeted to detect the parasite in wildlife. It is worth noting that earlobe samples showed a higher level of *Leishmania* DNA detection than the spleen samples. Results of lymph node examination suggested a low tendency for *L. infantum* visceralization in wildlife from RER. Other authors have also compared different tissues, with results varying between studies as well as between animal species. Abbate and colleagues [34] showed that in most foxes and wild rabbits, *L. infantum* DNA was found mainly in the spleen. In other lagomorph surveys, *L. infantum* was more frequently detected in the skin than in other tissues [30,55].

The mere finding of *L. infantum* DNA in a given animal species does not necessarily imply that this species is involved in *Leishmania* epidemiology, acting as a reservoir. Indeed, xenodiagnoses should be carried out to demonstrate host infectiousness to sand flies. However, it is likely that the success of host to sand fly transmission depends on the extent of skin parasite load, which has been suggested as the best marker to identify potential reservoir animal species [37,56,57]. Although the molecular analysis performed did not allow for exact quantification, the real-time PCR threshold cycle (Ct) of positivity can serve as an estimate of parasite load [15]. A low-to-intermediate number of parasites (median Ct between 25 and 35) were found in the earlobes of red deer, wolves, European hares, porcupines, roe deer, and red foxes. In the latter two species, we also found high parasite loads (Ct values <20) in several earlobe samples. Amastigotes in skin are directly accessible to sand flies, which are known to prefer safe, hairless parts of hosts, such as the ear pinnae, and feed copiously on them [58,59].

The results of the present study showed detection of *Leishmania* DNA also in winter and spring, seasons free for adult sand flies [28]. Roe deer tested positive in all seasons, but the limited number of sampled animals does not allow a definitive assessment to be drawn on all other species. Furthermore, the occurrence of Ct values below 25 in earlobe samples independently of the season suggests that some species could not clear the *L. infantum* infection, thus fitting the most important of the five criteria to consider a species as a reservoir of *Leishmania* parasites, i.e., the long course of infection [60,61]. In light of this, roe deer need to be further investigated, as they are among the species with low Ct values in spring, thus representing a possible host for *L. infantum* overwintering.

The present study confirmed previous findings on the existence of a sylvatic cycle for *L. infantum* in the study area. We cannot exclude that differences observed in wild hosts and parasite's tissue tropism in RER could be linked to: i) the peculiar parasitic population circulating in humans and sand flies in the region [14], being phylogenetically distant from the *L. infantum* MON-1 strains commonly affecting dogs and humans [62-65]; ii) the predominant sand fly species, i.e., *Ph. perfiliewi*, in comparison with other Mediterranean endemic areas, where *Ph. perniciosus* is the main vector of *L. infantum*. Finally, the exhibited multi-host infection pattern and preferential tropism for skin tissues would allow the parasite to widen its reservoir spectrum, allowing greater exploitation of the trophic (host) activity of the vector, determined by the vector host preference and the host abundance. The involvement of wild animals in the parasite's epidemiology will be determined by ongoing molecular research on the *L. infantum* DNA detected in wildlife in RER.

5. Conclusions

A deep knowledge of the transmission dynamics of *Leishmania* parasites is essential in order to apply control measures or monitoring programmes. This study provides evidence that *L. infantum* is widespread in wildlife in RER and gives preliminary data on tissue tropism, geographic and seasonal pattern of the parasite. Additionally, the study

expands the number of mammalian hosts for *L. infantum* in Italy. Further investigation is needed in order to clarify whether wildlife species should be considered maintenance hosts, sources of infection, or sentinels for *L. infantum* circulation in the re-emerging focus of leishmaniasis in Northeast Italy.

Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1, Table S1: Prevalence (including 95% confidence interval) of *L. infantum* real-time PCR positive species among the provinces of the study area.

Author Contributions: Conceptualization, G.R., R.T. and M.C.; methodology, R.T. and A.B.; validation, G.R., M.C. and R.T.; formal analysis, G.G., A.R. (Arianna Rossi) and A.S.; investigation, R.T., A.B. and G.R.; resources, G.P., M.F., L.F., M.C.F. and C.M. and A.R. (Alessandro Reggiani); data curation, R.T. and G.G.; writing—original draft preparation, R.T. and G.R.; writing—review and editing, A.S., M.C., E.C., C.M. and M.T.; visualization, G.G. and A.R. (Arianna Rossi); supervision, G.R. and R.T.; project administration, G.R.; funding acquisition, G.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Ministero della Salute, Italy (Ricerca Corrente 2019, IZSLER PRC2019016)

Institutional Review Board Statement: Ethical review and approval were waived for this study that did not involve killing of animals. The samples did not originate from experimental trial but took advantage just from diagnostic activity with Regional Surveillance Plans for wildlife diseases (DGR, 13 November 2017, no. 1703). Therefore, since the sampling was not specifically programmed as experimental study, but originating from diagnostic activity, we believed that it does not fall in the provisions of the National Law (i.e., DLGS 4/3 2014, n. 26. Application at national level of the EU Directive 2010/63/UE) and no ethical approval or permit for animal experimentation was required

Informed Consent Statement: Not applicable.

Data Availability Statement: Data generated or analyzed during this study are included in the published article.

Acknowledgments: The authors are grateful to all the local hunters, forest guards, and staff from the Wildlife Rescue and Rehabilitation Centres for their help in collecting samples. We would also like to express our gratitude to Dr. Elio Licata and Dr. Stefano Palminteri of the Veterinary Public Health Services for the implementation and coordination of the wildlife surveillance program in the Emilia-Romagna region.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Moriconi, M.; Rugna, G.; Calzolari, M.; Bellini, R.; Albieri, A.; Angelini, P.; Cagarelli, R.; Landini, M.P.; Charrel, R.N.; Varani, S. Phlebotomine sand fly-borne pathogens in the Mediterranean Basin: Human leishmaniasis and phlebovirus infections. *PLoS Negl Trop Dis* 2017, 11, e0005660, DOI 10.1371/journal.pntd.0005660.
2. Berriatua, E.; Maia, C.; Conceicao, C.; Ozbek, Y.; Toz, S.; Baneth, G.; Perez-Cutillas, P.; Ortuno, M.; Munoz, C.; Jumakanova, Z.; Pereira, A.; Rocha, R.; Monge-Maillo, B.; Gasimov, E.; Van der Stede, Y.; Torres, G.; Gossner, C.M. Leishmaniasis in the European Union and Neighboring Countries. *Emerg Infect Dis* 2021, 27, 10.3201/eid2706.210239, DOI 10.3201/eid2706.210239.
3. Gaspari, V.; Zaghi, I.; Macri, G.; Patrizi, A.; Salfi, N.; Locatelli, F.; Carra, E.; Re, M.C.; Varani, S. Autochthonous Cases of Mucosal Leishmaniasis in Northeastern Italy: Clinical Management and Novel Treatment Approaches. *Microorganisms* 2020, 8, 10.3390/microorganisms8040588, DOI E588.
4. Gradoni, L.; López-Vélez, R.; Mokni, M. Manual on case management and surveillance of the leishmaniasis in the WHO European Region. 2017 Available online: <https://apps.who.int/iris/bitstream/handle/10665/344118/9789289052511-eng.pdf?sequence=1&isAllowed=y>. (Accessed on: 25/10/2022).
5. Solano-Gallego, L.; Koutinas, A.; Miro, G.; Cardoso, L.; Pennisi, M.G.; Ferrer, L.; Bourdeau, P.; Oliva, G.; Baneth, G. Directions for the diagnosis, clinical staging, treatment and prevention of canine leishmaniosis. *Vet Parasitol* 2009, 165, 1-18, DOI 10.1016/j.vet-par.2009.05.022.
6. Maia, C.; Cardoso, L. Spread of *Leishmania infantum* in Europe with dog travelling. *Vet Parasitol* 2015, 213, 2-11, DOI 10.1016/j.vet-par.2015.05.003.
7. Maroli, M.; Rossi, L.; Baldelli, R.; Capelli, G.; Ferroglio, E.; Genchi, C.; Gramiccia, M.; Mortarino, M.; Pietrobelli, M.; Gradoni, L. The northward spread of leishmaniasis in Italy: evidence from retrospective and ongoing studies on the canine reservoir and phlebotomine vectors. *Trop Med Int Health* 2008, 13, 256-264, DOI 10.1111/j.1365-3156.2007.01998.x.
8. Pampiglione, S. La Leishmaniosi Viscerale in Emilia-Romagna. *Ann. Sanità Pubblica* 1974, 35, 1021-1028.

9. Pampiglione, S.; La Placa, M.; Schlick, G. Studies on mediterranean Leishmaniasis. I. An outbreak of visceral leishmaniasis in Northern Italy. *Trans R Soc Trop Med Hyg* 1974, 68, 349-359, DOI 10.1016/0035-9203(74)90148-5.
10. Baldelli, R.; Di Francesco, A. Leishmaniosi in Italia: risultati di indagini sierologiche condotte su cani di diversa provenienza geografica, 1992; , pp. 1395– 1399.
11. Varani, S.; Cagarelli, R.; Melchionda, F.; Attard, L.; Salvadori, C.; Finarelli, A.C.; Gentilomi, G.A.; Tigani, R.; Rangoni, R.; Todeschini, R.; Scalone, A.; Di Muccio, T.; Gramiccia, M.; Gradoni, L.; Viale, P.; Landini, M.P. Ongoing outbreak of visceral leishmaniasis in Bologna Province, Italy, November 2012 to May 2013. *Euro Surveill* 2013, 18, 20530, DOI 20530.
12. Franceschini, E.; Puzzolante, C.; Menozzi, M.; Rossi, L.; Bedini, A.; Orlando, G.; Gennari, W.; Meacci, M.; Rugna, G.; Carra, E.; Codeluppi, M.; Mussini, C. Clinical and Microbiological Characteristics of Visceral Leishmaniasis Outbreak in a Northern Italian Nonendemic Area: A Retrospective Observational Study. *Biomed Res Int* 2016, 2016, 6481028, DOI 10.1155/2016/6481028.
13. Rugna, G.; Carra, E.; Corpus, F.; Calzolari, M.; Salvatore, D.; Bellini, R.; Di Francesco, A.; Franceschini, E.; Bruno, A.; Poglayen, G.; Varani, S.; Vitale, F.; Merialdi, G. Distinct Leishmania infantum Strains Circulate in Humans and Dogs in the Emilia-Romagna Region, Northeastern Italy. *Vector Borne Zoonotic Dis* 2017, 17, 409-415, DOI 10.1089/vbz.2016.2052.
14. Rugna, G.; Carra, E.; Bergamini, F.; Calzolari, M.; Salvatore, D.; Corpus, F.; Gennari, W.; Baldelli, R.; Fabbi, M.; Natalini, S.; Vitale, F.; Varani, S.; Merialdi, G. Multilocus microsatellite typing (MLMT) reveals host-related population structure in Leishmania infantum from northeastern Italy. *PLoS Negl Trop Dis* 2018, 12, e0006595, DOI 10.1371/journal.pntd.0006595.
15. Calzolari, M.; Carra, E.; Rugna, G.; Bonilauri, P.; Bergamini, F.; Bellini, R.; Varani, S.; Dottori, M. Isolation and Molecular Typing of Leishmania infantum from Phlebotomus perfiliewi in a Re-Emerging Focus of Leishmaniasis, Northeastern Italy. *Microorganisms* 2019, 7, 10.3390/microorganisms7120644, DOI E644.
16. Calzolari, M.; Romeo, G.; Bergamini, F.; Dottori, M.; Rugna, G.; Carra, E. Host preference and Leishmania infantum natural infection of the sand fly Phlebotomus perfiliewi in northern Italy. *Acta Trop* 2022, 226, 106246, DOI S0001-706X(21)00424-1 [pii].
17. Cardoso, L.; Schallig, H.; Persichetti, M.F.; Pennisi, M.G. New Epidemiological Aspects of Animal Leishmaniosis in Europe: The Role of Vertebrate Hosts Other Than Dogs. *Pathogens* 2021, 10, 10.3390/pathogens10030307, DOI 307.
18. Antolini, G.; Pavan, V.; Tomozeiu, R.; Marletto, V. Atlante climatico dell'Emilia-Romagna 1961-2015. ARPAE Emilia-Romagna 2017.
19. Regione Emilia-Romagna. Piano regionale di sorveglianza e controllo della leishmaniosi canina. Available online: <https://www.anagrafecaninarer.it/acrer/Default.aspx?tabid=160#>. (Accessed on: 25/10/2022).
20. Regione Emilia Romagna Sorveglianza e monitoraggio della fauna selvatica. Available online: <https://www.alimenti-salute.it/documentazione-regionale/90355>. (Accessed on: 25/10/2022).
21. Galletti, E.; Bonilauri, P.; Bardasi, L.; Fontana, M.C.; Ramini, M.; Renzi, M.; Dosa, G.; Merialdi, G. Development of a minor groove binding probe based real-time PCR for the diagnosis and quantification of Leishmania infantum in dog specimens. *Res Vet Sci* 2011, 91, 243-245, DOI 10.1016/j.rvsc.2011.01.004.
22. Clopper, C.J.; Pearson, E.S. The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial. *Biometrika* 1934, 26, 404-413, DOI 10.2307/2331986.
23. R Core Team R: A language and environment for statistical computing. Available online: R Core Team (2020). R Foundation for Statistical Computing, Vienna, Austria. Available on: <https://www.R-project.org>.
24. Santi, A.; Renzi, M.; Baldelli, R.; Calzolari, M.; Caminiti, A.; Dell'Anna, S.; Galletti, G.; Lombardini, A.; Paternoster, G.; Tamba, M. A surveillance program on canine leishmaniasis in the public kennels of Emilia-Romagna Region, Northern Italy. *Vector Borne Zoonotic Dis* 2014, 14, 206-211, DOI 10.1089/vbz.2013.1362.
25. Cesinaro, A.M.; Nosseir, S.; Mataka, E.; Mengoli, M.C.; Cavatorta, C.; Gennari, W. An outbreak of cutaneous leishmaniasis in Modena province (Northern Italy): report of 35 cases. *Pathologica* 2017, 109, 363-367.
26. Varani, S.; Cagarelli, R.; Melchionda, F.; Attard, L.; Salvadori, C.; Finarelli, A.C.; Gentilomi, G.A.; Tigani, R.; Rangoni, R.; Todeschini, R.; Scalone, A.; Di Muccio, T.; Gramiccia, M.; Gradoni, L.; Viale, P.; Landini, M.P. Ongoing outbreak of visceral leishmaniasis in Bologna Province, Italy, November 2012 to May 2013. *Euro Surveill* 2013, 18, 20530, DOI 20530.
27. Corradetti, A. Phlebotomus and leishmaniasis in North-Central Italy (Apennine Region), 1962; , pp. 103-109.
28. Calzolari, M.; Romeo, G.; Callegari, E.; Bonilauri, P.; Chiapponi, C.; Carra, E.; Rugna, G.; Taddei, R.; Lelli, D.; Dottori, M. Co-Circulation of Phlebotomus and Leishmania Parasites in Sand Flies from a Single Site in Italy Monitored between 2017 and 2020. *Viruses* 2021, 13, 10.3390/v13081660, DOI 1660.
29. Oleaga, A.; Zanet, S.; Espi, A.; Pegoraro de Macedo, M R; Gortazar, C.; Ferroglio, E. Leishmania in wolves in northern Spain: A spreading zoonosis evidenced by wildlife sanitary surveillance. *Vet Parasitol* 2018, 255, 26-31, DOI S0304-4017(18)30120-1 [pii].
30. Risueno, J.; Ortuno, M.; Perez-Cutillas, P.; Goyena, E.; Maia, C.; Cortes, S.; Campino, L.; Bernal, L.J.; Munoz, C.; Arcenillas, I.; Martinez-Rondan, F.J.; Gonzalez, M.; Collantes, F.; Ortiz, J.; Martinez-Carrasco, C.; Berriatua, E. Epidemiological and genetic studies suggest a common Leishmania infantum transmission cycle in wildlife, dogs and humans associated to vector abundance in Southeast Spain. *Vet Parasitol* 2018, 259, 61-67, DOI S0304-4017(18)30199-7.
31. Battisti, E.; Zanet, S.; Khalili, S.; Trisciuglio, A.; Hertel, B.; Ferroglio, E. Molecular Survey on Vector-Borne Pathogens in Alpine Wild Carnivorans. *Front Vet Sci* 2020, 7, 1, DOI 10.3389/fvets.2020.00001.
32. Dipineto, L.; Manna, L.; Baiano, A.; Gala, M.; Fioretti, A.; Gravino, A.E.; Menna, L.F. Presence of Leishmania infantum in red foxes (Vulpes vulpes) in southern Italy. *J Wildl Dis* 2007, 43, 518-520, DOI 43/3/518.
33. Ranieri, V.; Poli, A.; Ariti, G.; Nardoni, S.; Bertuccelli, M.; Fanucchi Manciant Detection of Leishmania infantum DNA in tissues of free-ranging red foxes (Vulpes vulpes) in Central Italy. *Eur J Wildl Res* 2010, 56, 689-692.

34. Abbate, J.M.; Arfuso, F.; Napoli, E.; Gaglio, G.; Giannetto, S.; Latrofa, M.S.; Otranto, D.; Brianti, E. *Leishmania infantum* in wild animals in endemic areas of southern Italy. *Comp Immunol Microbiol Infect Dis* 2019, 67, 101374, DOI S0147-9571(19)30171-7.
35. Criado-Fornelio, A.; Gutierrez-Garcia, L.; Rodriguez-Cabeiro, F.; Reus-Garcia, E.; Roldan-Soriano, M.A.; Diaz-Sanchez, M.A. A parasitological survey of wild red foxes (*Vulpes vulpes*) from the province of Guadalajara, Spain. *Vet Parasitol* 2000, 92, 245-251, DOI S0304-4017(00)00329-0.
36. Sobrino, R.; Ferroglio, E.; Oleaga, A.; Romano, A.; Millan, J.; Revilla, M.; Arnal, M.C.; Trisciuglio, A.; Gortazar, C. Characterization of widespread canine leishmaniasis among wild carnivores from Spain. *Vet Parasitol* 2008, 155, 198-203, DOI 10.1016/j.vetpar.2008.05.003.
37. Del Rio, L.; Chitimia, L.; Cubas, A.; Victoriano, I.; De la Rua, P.; Gerrikagoitia, X.; Barral, M.; Munoz-Garcia, C.I.; Goyena, E.; Garcia-Martinez, D.; Fisa, R.; Riera, C.; Murcia, L.; Segovia, M.; Berriatua, E. Evidence for widespread *Leishmania infantum* infection among wild carnivores in *L. infantum* periendemic northern Spain. *Prev Vet Med* 2014, 113, 430-435, DOI 10.1016/j.prevetmed.2013.12.001.
38. Petrisceva, P.A. The natural focalities of leishmaniasis in the USSR. *Bull World Health Organ* 1971, 44, 567-576.
39. Magri, A.; Galuppi, R.; Fioravanti, M.; Caffara, M. Survey on the presence of *Leishmania* sp. in peridomestic rodents from the Emilia-Romagna Region (North-Eastern Italy). *Vet Res Commun* 2022, DOI 10.1007/s11259-022-09925-4.
40. Calzolari, M.; Romeo, G.; Munari, M.; Bonilauri, P.; Taddei, R.; Sampieri, M.; Bariselli, S.; Rugna, G.; Dottori, M. Sand Flies and Pathogens in the Lowlands of Emilia-Romagna (Northern Italy). *Viruses* 2022, 14, DOI 10.3390/v14102209.
41. Ruiz-Fons, F.; Ferroglio, E.; Gortazar, C. *Leishmania infantum* in free-ranging hares, Spain, 2004-2010. *Euro Surveill* 2013, 18, 20541, DOI 20541.
42. Tsokana, C.N.; Sokos, C.; Giannakopoulos, A.; Mamuris, Z.; Birtsas, P.; Papaspyropoulos, K.; Valiakos, G.; Spyrou, V.; Lefkaditis, M.; Chatzopoulos, D.C.; Kantere, M.; Manolakou, K.; Touloudi, A.; Burriel, A.R.; Ferroglio, E.; Hadjichristodoulou, C.; Billinis, C. First evidence of *Leishmania* infection in European brown hare (*Lepus europaeus*) in Greece: GIS analysis and phylogenetic position within the *Leishmania* spp. *Parasitol Res* 2016, 115, 313-321, DOI 10.1007/s00436-015-4749-8 [doi].
43. Gradoni, L.; Pozio, E.; Gramiccia, M.; Maroli, M.; Bettini, S. Leishmaniasis in Tuscany (Italy): VII. Studies on the role of the black rat, *Rattus rattus*, in the epidemiology of visceral leishmaniasis. *Trans R Soc Trop Med Hyg* 1983, 77, 427-431, DOI 10.1016/0035-9203(83)90102-5.
44. Molina, R.; Jimenez, M.I.; Cruz, I.; Iriso, A.; Martin-Martin, I.; Sevillano, O.; Melero, S.; Bernal, J. The hare (*Lepus granatensis*) as potential sylvatic reservoir of *Leishmania infantum* in Spain. *Vet Parasitol* 2012, 190, 268-271, DOI 10.1016/j.vetpar.2012.05.006.
45. Millan, J.; Ferroglio, E.; Solano-Gallego, L. Role of wildlife in the epidemiology of *Leishmania infantum* infection in Europe. *Parasitol Res* 2014, 113, 2005-2014, DOI 10.1007/s00436-014-3929-2.
46. Alcover, M.M.; Riera, M.C.; Fisa, R. Leishmaniasis in Rodents Caused by *Leishmania infantum*: A Review of Studies in the Mediterranean Area. *Front Vet Sci* 2021, 8, 702687, DOI 10.3389/fvets.2021.702687.
47. Munoz-Madrid, R.; Belinchon-Lorenzo, S.; Iniesta, V.; Fernandez-Cotrino, J.; Parejo, J.C.; Serrano, F.J.; Monroy, I.; Baz, V.; Gomez-Luque, A.; Gomez-Nieto, L.C. First detection of *Leishmania infantum* kinetoplast DNA in hair of wild mammals: application of qPCR method to determine potential parasite reservoirs. *Acta Trop* 2013, 128, 706-709, DOI 10.1016/j.actatropica.2013.08.009.
48. Miro, G.; Troyano, A.; Montoya, A.; Farinas, F.; Fermin, M.L.; Flores, L.; Rojo, C.; Checa, R.; Galvez, R.; Marino, V.; Frago, C.; Martinez-Navado, E. First report of *Leishmania infantum* infection in the endangered orangutan (*Pongo pygmaeus pygmaeus*) in Madrid, Spain. *Parasit Vectors* 2018, 11, 185-1, DOI 10.1186/s13071-018-2772-1.
49. Azami-Conesa, I.; Gomez-Munoz, M.T.; Martinez-Diaz, R.A. A Systematic Review (1990-2021) of Wild Animals Infected with Zoonotic *Leishmania*. *Microorganisms* 2021, 9, 10.3390/microorganisms9051101, DOI 1101.
50. Mhadhbi, M.; Sassi, A. Infection of the equine population by *Leishmania* parasites. *Equine Vet J* 2020, 52, 28-33, DOI 10.1111/evj.13178.
51. Paixao-Marques, M.D.S.; Alves-Martin, M.F.; Guiraldi, L.M.; Dos Santos, W.J.; de Lemos, F.A.; Sanchez, G.P.; Richini-Pereira, V.B.; Lucheis, S.B. First isolation of *Leishmania infantum* by blood culture in bovines from endemic area for canine visceral leishmaniasis. *Parasitology* 2019, 146, 911-913, DOI 10.1017/S0031182019000088.
52. Moraes-Silva, E.; Antunes, F.R.; Rodrigues, M.S.; da Silva Juliao, F.; Dias-Lima, A.G.; Lemos-de-Sousa, V.; de Alcantara, A.C.; Reis, E.A.; Nakatani, M.; Badaro, R.; Reis, M.G.; Pontes-de-Carvalho, L.; Franke, C.R. Domestic swine in a visceral leishmaniasis endemic area produce antibodies against multiple *Leishmania infantum* antigens but apparently resist to *L. infantum* infection. *Acta Trop* 2006, 98, 176-182, DOI S0001-706X(06)00073-8.
53. Brazil, R.P.; Desterro, M.D.; Nascimento, S.B.; Macau, R.P. Natural infection of a pig (*Sus scrofa*) by *Leishmania* in a recent focus of cutaneous leishmaniasis on the Island of Sao Luis, Maranhao. *Mem Inst Oswaldo Cruz* 1987, 82, 145, DOI S0074-02761987000100025.
54. Martin-Sanchez, J.; Torres-Medina, N.; Morillas-Marquez, F.; Corpas-Lopez, V.; Diaz-Saez, V. Role of wild rabbits as reservoirs of leishmaniasis in a non-epidemic Mediterranean hot spot in Spain. *Acta Trop* 2021, 222, 106036, DOI S0001-706X(21)00215-1.
55. Ortega-Garcia, M.V.; Salguero, F.J.; Rodriguez-Bertos, A.; Moreno, I.; Garcia, N.; Garcia-Seco, T.; Luz Torre, G.; Dominguez, L.; Dominguez, M. A pathological study of *Leishmania infantum* natural infection in European rabbits (*Oryctolagus cuniculus*) and Iberian hares (*Lepus granatensis*). *Transbound Emerg Dis* 2019, 66, 2474-2481, DOI 10.1111/tbed.13305.
56. Aslan, H.; Oliveira, F.; Meneses, C.; Castrovinci, P.; Gomes, R.; Teixeira, C.; Derenge, C.A.; Orandle, M.; Gradoni, L.; Oliva, G.; Fischer, L.; Valenzuela, J.G.; Kamhawi, S. New Insights Into the Transmissibility of *Leishmania infantum* From Dogs to Sand Flies: Experimental Vector-Transmission Reveals Persistent Parasite Depots at Bite Sites. *J Infect Dis* 2016, 213, 1752-1761, DOI 10.1093/infdis/jiw022.

-
57. Arumugam, S.; Scorza, B.M.; Petersen, C. Visceral Leishmaniasis and the Skin: Dermal Parasite Transmission to Sand Flies. *Pathogens* 2022, 11, 10.3390/pathogens11060610, DOI 610.
 58. Svobodova, M.; Votypka, J. Experimental transmission of *Leishmania tropica* to hamsters and mice by the bite of *Phlebotomus sergenti*. *Microbes Infect* 2003, 5, 471-474, DOI S1286457903000662.
 59. Ready, P.D. Biology of phlebotomine sand flies as vectors of disease agents. *Annu Rev Entomol* 2013, 58, 227-250, DOI 10.1146/annurev-ento-120811-153557.
 60. WHO Expert Committee on the Control of the Leishmaniases. Meeting; World Health Organization *Control of the Leishmaniases: Report of a Meeting of the WHO Expert Committee on the Control of Leishmaniases, Geneva, 22-26 March 2010*, World Health Organization: 2010.
 61. Chaves, L.F.; Hernandez, M.; Dobson, A.P.; Pascual, M. Sources and sinks: revisiting the criteria for identifying reservoirs for American cutaneous leishmaniasis. *Trends Parasitol* 2007, 23, 311-316, DOI <https://doi.org/10.1016/j.pt.2007.05.003>.
 62. Chicharro, C.; Llanes-Acevedo, I.P.; Garcia, E.; Nieto, J.; Moreno, J.; Cruz, I. Molecular typing of *Leishmania infantum* isolates from a leishmaniasis outbreak in Madrid, Spain, 2009 to 2012. *Euro Surveill* 2013, 18, 20545, DOI 20545.
 63. Cortes, S.; Mauricio, I.L.; Kuhls, K.; Nunes, M.; Lopes, C.; Marcos, M.; Cardoso, L.; Schonian, G.; Campino, L. Genetic diversity evaluation on Portuguese *Leishmania infantum* strains by multilocus microsatellite typing. *Infect Genet Evol* 2014, 26, 20-31, DOI 10.1016/j.meegid.2014.04.023.
 64. Pomares, C.; Marty, P.; Banuls, A.L.; Lemichez, E.; Pratlong, F.; Faucher, B.; Jeddi, F.; Moore, S.; Michel, G.; Aluru, S.; Piarroux, R.; Hide, M. Genetic Diversity and Population Structure of *Leishmania infantum* from Southeastern France: Evaluation Using Multi-Locus Microsatellite Typing. *PLoS Negl Trop Dis* 2016, 10, e0004303, DOI 10.1371/journal.pntd.0004303.
 65. El Hamouchi, A.; El Kacem, S.; Ejghal, R.; Lemrani, M. Genetic polymorphism in *Leishmania infantum* isolates from human and animals determined by nagt PCR-RFLP. *Infect Dis Poverty* 2018, 7, 54-y, DOI 10.1186/s40249-018-0439-y.