

Review

Use of antiandrogens as therapeutic agents in COVID-19 patients

Efstathios S. Giotis ^{1,2,*}, Emine Cil ² and Greg N. Brooke ²

¹ Department of Infectious Diseases, Imperial College London, London, W2 1PG, UK

² School of Life Sciences, University of Essex, Colchester, ESSEX, CO4 3SQ, UK

* Correspondence: author

Abstract: COVID-19, caused by the severe acute respiratory syndrome coronavirus 2 (SARS CoV-2), is estimated to have caused over 6.5 million deaths to date worldwide. The emergence of fast-evolving SARS-CoV-2 variants of concern alongside increased transmissibility and/or virulence as well as immune and vaccine escape capabilities highlight the urgent need for more effective antivirals to combat the disease in the long run along with regularly updated vaccine boosters. One of the early risk factors identified during the COVID-19 pandemic was that men are more likely to get infected by the virus, more likely to develop severe disease and exhibit higher likelihood of hospitalisation and mortality rates compared to women. An association exists between SARS-CoV-2 infectiveness and disease severity with sex steroid hormones and in particular androgens. Several studies underlined the importance of the androgen-mediated regulation of the host protease TMPRSS2 and the cell entry protein ACE2 as well as the key role of these factors in the entry of the virus into target cells. In this context, modulating androgen signalling is a promising strategy to block viral infection and antiandrogens could be used as a preventative measure at the pre- or early-hospitalisation stage of COVID-19 disease. Different antiandrogens, including commercial drugs used to treat metastatic castration-sensitive prostate cancer and other conditions, have been tested as antivirals with varying success. In this review, we summarise the most recent updates concerning the use of antiandrogens as prophylactic and therapeutic options for COVID-19.

Keywords: coronavirus; TMPRSS2; ACE2; SARS-CoV-2; androgens; sex hormones; antiandrogens

1. Emergence of COVID-19 and variants of concern

Coronaviruses (CoV) are a large family of positive-sense, enveloped, single-stranded RNA viruses that are common in people and many animal species. Human coronaviruses (hCoV) include the common cold CoV 229E, OC43, NL63 and HKU1 which predominantly cause mild respiratory illness, and three highly pathogenic coronaviruses which cause more severe and fatal disease in people; severe acute respiratory syndrome coronavirus (SARS-CoV, 2003), Middle East respiratory syndrome coronavirus (MERS-CoV, 2012), and more recently SARS-CoV-2 (2019) the causative agent of COVID-19 [1]. At the end of September 2022, over 600 million cases of COVID-19 have been confirmed globally, with a 4-5% crude mortality risk (deaths per 100 patients hospitalized primarily for COVID-19) [2]. COVID-19 may present in patients with mild, moderate or severe symptomatology. Mild symptoms include fever, dry cough, dyspnoea, fatigue, while COVID-19 associated mortality is attributed to a severe respiratory failure resembling acute respiratory distress syndrome (ARDS) [3-5]. Severe pneumonia COVID-19 patients exhibit exacerbated inflammatory response resulting in excessive release of pro-inflammatory cytokines, known as the “cytokine storm”, which leads to alveolar damage, fibrosis, and progressive respiratory failure [5-8]. It is becoming increasingly clear that COVID-19 is a multi-system disease and can also lead to heart failure, acute kidney and neurovascular injuries [5, 9, 10]. In addition, growing patient testimony and scientific evidence demonstrate that a large number of patients (approximately 2% of the UK population alone) also

suffer from post-COVID syndrome (long COVID or postacute sequelae of COVID-19 (PASC)) which cause debilitating physical or mental symptoms for long periods after infection [5, 11-13].

The vaccination programmes have been key to reducing COVID-19 hospital admissions and mortality rates, but for some countries and groups of people the uptake is still low due to various socioeconomic factors and accessibility inequalities. In addition, the vaccination-acquired immunity wanes substantially with time and boosters are required to restore the efficiency of vaccines against severe disease [14, 15]. Despite progress in countries with high vaccination rates, the pandemic is still actively unfolding with emerging virus variants that can evade vaccine or natural immunity triggering new public health challenges. Virus variants are characterised by the WHO as Variants of Concern (VOC) if they demonstrate increased transmissibility, virulence, or cause reduced effectiveness of vaccine induced protection, diagnostic tests and management measures [16, 17]. Important variants include B.1.1.7 (Alpha), B.1.351 (Beta), P.1 (Gamma) and B.1.617.2 (Delta) [17-19]. Two new sub-variants (B.1.1.529; Omicron) designated as BA.4 and BA.5 emerged in December 2021 and January 2022 respectively, which replaced the other circulating Omicron lineages and have driven the current wave of infections in Africa, the United States, and Europe [20]. These two subvariants appear more contagious and seem to circumvent immunity from previous infections or vaccination [21, 22]. As a result of the emergence of Omicron variants, a number of updated COVID-19 vaccines have been developed and approved for use in United States and United Kingdom and elsewhere. However, a recent study suggested that the direct protective benefit of the Omicron-updated bivalent vaccines is only marginal compared to ancestral-based vaccines [23].

COVID-19 hospitalised patient management in high-income countries is largely focused on highly-efficient albeit high-priced monoclonal antibodies [24], anti-inflammatory drugs such as dexamethasone, which reduce cytokine levels and immune-mediated pathogenesis in the hyper-inflammatory phase of the disease [25], and/or drugs which directly interfere with viral replication. The latter act either by inhibition of the viral 3CL^{pro} protease (paxlovid), by inhibition of the RNA-dependent RNA polymerase (remdesivir)[26], or by introducing copying errors in the viral genome (molnupiravir) [27]. Recent clinical studies reported that remdesivir has no significant effect on patients who are already being ventilated [28] and that some paxlovid- or molnupiravir-treated patients experience rebound infections after completing a course of treatment [29, 30]. Taken together, more efficient and less expensive antivirals drugs for pre-emptive protection against emerging variants, reducing reliance on boosters, are needed especially in low- and middle-income countries.

2. Risk factors and gender disparity in COVID-19 patients

Most COVID-19 patients are expected to have a favourable prognosis. However, some groups are at higher risk to develop severe or fatal illness due to underlying health conditions. These include demographic factors (i.e. older age), gender, occupational exposure, ethnicity, and more importantly comorbidities such as diabetes, hypertension, obesity, neurological disorders, cancer, heart, lung, liver and kidney disease [31-35]. Researchers are in the early stages of understanding the risk factors of viral persistence and long COVID. Accumulating evidence showed that people who experienced multisystem inflammatory syndrome (MIS) with underlying health conditions and who were not vaccinated were more likely to develop long-term post-COVID sequelae [36-38].

Since the start of the coronavirus pandemic, emerging gender-disaggregated data from multiple countries suggested that men were disproportionately affected by COVID-19. Initially, it was hypothesised that the higher occupational exposure and comorbidities of males compared to females were contributory to this discrepancy. Follow-up studies confirmed that men globally exhibit higher infection, hospitalisation and mortality rates compared to women [39-41]. Female COVID-19 patients were also found to clear the virus and recover substantially earlier compared to male patients [42, 43]. Sex differences in prevalence, severity and outcome of viral infection have been previously reported for

other viruses such as influenza viruses, HIV, herpes simplex viruses, hepatitis B, measles, and West Nile virus [44-46]. The reasons accounting for the higher vulnerability of males to viruses are not clear but several immunological as well as genetic, hormonal, and socio-behavioural explanations have been put forward.

The innate immune system serves as the first line of host defence against invading viruses and involves sensing of viruses through pattern recognition receptors and activating inflammatory pathways, which eventually result in viral clearance. Innate immune cells i.e. monocytes, dendritic cells, and macrophages, are functionally more active in females [47-49]. Females are typically able to launch more vigorous innate and adaptive immune responses, have higher resistance to most viral infections and exhibit higher antibody responses following infection and vaccination compared to males [47, 50-52]. The bi-allelic expression of immune-related genes present on the X chromosome such as the RNA receptors TLR7 and TLR8 may be responsible for sexually dimorphic immunophenotypes [51-53]. Hormonal regulation may also be responsible for the gender disparity in the disease progression and outcome following viral challenges. Steroid hormones, such as androgens estrogens, progesterone and glucocorticoids control the activity of innate immune cells and the survival and differentiation of T and B lymphocytes in both men and women [47, 53-55]. Many studies have been published on the effects of hormones to virus susceptibility. The consensus is that androgens (a primarily male hormone) and progestones are immunosuppressive and increase susceptibility to viruses while estrogens (the primary female hormones) are immunoprotective and enhance antiviral responses [48, 49, 56]. Besides the effect on antiviral responses, androgens can also directly influence virus activity i.e. HBV, HCV [57]. Conversely, viruses are also able to manipulate sex steroid receptor signalling mechanisms to serve their own survival and enhance their replication rate [58]. The extent to which male and female immune responses differ during SARS-CoV-2 infection and the possibility of a bi-directional interaction between the virus and androgens have been overlooked so far and deserve greater attention.

3. Cell entry of SARS-CoV-2

Viral attachment, fusion and entry of SARS-CoV-2 into target cells is mediated by the virus' envelope spike (S) glycoprotein [59-61]. The S protein is the main virus' antigenic site for inducing antibody responses and is therefore a key target for the development of antivirals and vaccines. S is synthesized as a precursor protein that is cleaved into two subunits: the N-terminal S1 subunit, which contains the receptor binding domain, and the C-terminal S2 subunit, which harbors the viral fusion machinery [59, 60, 62]. SARS-CoV-2 initiates cell entry by binding its S to the transmembrane glycoprotein Angiotensin-Converting Enzyme 2 (ACE2), a negative controller of the Renin Angiotensin system [63, 64]. Following binding, the virus gains access into the cell by two alternative routes either via direct fusion with the cell membrane or via clathrin-mediated endocytosis [60-62]. Both entry mechanisms require priming of the S protein by proteolytic cleavage at two sites in a manner reminiscent to that of the proteolytic cleavage and maturation of influenza A viruses' entry protein haemagglutinin [65, 66]. The first cleavage site is a polybasic insertion (PRRAR) at the S1-S2 boundary, and the second cleavage occurs at the S2' site in the S2 subunit, a crucial step for triggering the fusion of viral and host cell membranes [60, 61, 67, 68]. The S1-S2 site is cleaved by the serine protease furin, which is ubiquitously expressed in the respiratory tract, while the S2' site can be cleaved by either the endosomal cathepsins (B and L) in the endolysosome or by the cell-surface transmembrane protease serine 2 (TMPRSS2) at the plasma membrane [59, 69, 70]. TMPRSS2 is part of the type 2 transmembrane serine protease (TTSP) family and has been extensively studied in the context of prostate cancer as its expression is regulated in response to androgens through direct transcriptional regulation by the androgen receptor (AR) [71, 72]. Other TTSPs such as TMPRSS4 may also act as SARS-CoV-2 cell entry mediators [73]. The aftermath of SARS-CoV-2 cell-entry is governed by a complicated network of host-pathogen interactions, where a range of viral and host factors can demarcate the final outcome.




Figure 1. Alternative entry routes of SARS-CoV-2 into target cells. Elements of the schematic are from Motifolio.

4. ACE2 and TMPRSS2 as candidate targets for antiviral therapy

A large number of viral and host factors that play key roles in the virus pathogenesis has been discovered. Among them, ACE2 and TMPRSS2 have been proposed as the most attractive targets for the development of antivirals that could inhibit or delay virus infection, thus possibly reducing viral transmission and symptom severity. Both factors are expressed in the nasal and bronchial secretory cells including the alveolar epithelial type II cells (AT2), which is the main target cell subpopulation of SARS-CoV-2, as well as gastrointestinal and other virus-targeted tissues [74-78]. ACE2 regulates key processes in the human body such as blood pressure, cardiovascular function, wound healing and inflammation and acts as the main cellular receptor for SARS-CoV and the hCoV NL63 [79, 80]. TMPRSS2 is also essential for the cell entry of other viruses i.e. influenza A and B, HCV as well as other coronaviruses [81-85].

At the onset of the COVID-19 pandemic, there were studies suggesting that higher ACE2 and TMPRSS2 levels may be associated with higher susceptibility to SARS-CoV-2 infection, and this spurred extensive research that unveiled novel insights into the function and biology of ACE2/TMPRSS2. Several reports revealed an association between the variable expression of ACE2 and TMPRSS2 in different tissues across individuals with COVID-19 severity/fatality variations [86-90]. The lung airway expression of both ACE2 and TMPRSS2 has been found to be lower in children which are less susceptible to infection compared with adults, and significantly higher in smokers compared to non-smokers, and in patients with chronic obstructive pulmonary disease (COPD) compared to healthy individuals [91, 92].

Targeting ACE2 was considered the obvious first-choice target for prophylactic and therapy interventions aiming to block virus accessibility to respiratory cells early on in infection. Several strategies have been evaluated *in vitro* and *in vivo* with varied success, including decoy or soluble ACE2 molecules, pseudoligands with a high ACE2 affinity, and blocking antibodies [63, 93, 94]. However, a disadvantage of these approaches is the potential dysregulation of ACE2-mediated vasodilation, amino acid transport and pancreatic insulin secretion [94-97]. TMPRSS2-inhibitors are arguably better candidates for COVID-19 antivirals than those for ACE2 as knockout of TMPRSS2 protein causes no overt detrimental phenotype [98, 99]. Numerous TMPRSS2 inhibitors have been tested in

in vitro studies that showed they inhibit entry of SARS-CoV-2 in lung cells including notably camostat, nafamostat, and several peptidomimetic inhibitors [63, 100-103]. These inhibitors were either 'repurposed' commercially available drugs, designed *de novo* or identified by virtual screenings and have been reviewed extensively elsewhere [104-109]. Several of these compounds have already or are currently been investigated in randomized, controlled trials to assess their use as monotherapies against COVID-19. Clinical trials testing efficacy of camostat and nafamostat have shown contradictory results. Some studies demonstrated a clear, beneficial effect for COVID-19 patients [110-112] while others showed only modest clinical improvements [101]. A possible caveat for these inhibitors is that the currently prevailing Omicron subvariants are largely entering target cells via the endosomal pathway [113] and it is possible that inhibitors targeting the TMPRSS2 (non-endosomal) entry route might not have a significant effect. Separate studies have targeted the expression of TMPRSS2 by using either androgen receptor (AR) antagonists or drugs that lower circulating androgen levels. The relationship of androgens and COVID-19 is not entirely clear and is even considered controversial for some researchers. Conflicting studies have been published reporting that both subdued or excessive testosterone levels can lead to severe COVID-19 disease e.g. in hypogonadism in elder men and in testosterone-treated postmenopausal women respectively, which suggests that the androgen-mediated effects to COVID-19 patients are likely multidimensional and interdependent on a range of confounding factors [114].

5. Androgen receptor structure and signalling

The Androgen Receptor (AR), a member of the Steroid Receptor Family, is a ligand dependent transcription factor with a modular structure [115]. The N-terminal Activation Function (AF-1) is a region that mediates protein-protein interactions and is important for transcriptional activity. Adjacent to this region, is the DNA binding domain (DBD), consisting of two zinc-finger like modules, which modulate the interaction of the receptors with specific DNA response elements present in the regulatory regions of target genes. The C-terminus houses the ligand binding domain (LBD)/Activation Function 2 (AF2). The AR LBD is formed of 12 α -helices which fold to form a pocket into which androgen fits. Ligand binding promotes a conformational change, resulting in receptor activation [116, 117].

In the absence of ligand, the AR is located in the cytoplasm and is held in a ligand binding competent state by a heat shock protein complex (Figure 2). Testosterone can diffuse into the cell and is converted to the more potent androgen dihydrotestosterone (DHT) by 5 α -reductase. Upon ligand binding, the AR undergoes a conformational change which promotes dissociation of the heat shock protein complex, dimerisation and nuclear localisation. Next, the AR binds to androgen response elements (AREs) in the regulatory regions of target genes and via the recruitment of coactivators (proteins that enhance the receptors transcriptional activity) and the basal transcriptional machinery, enhances or represses gene expression [115-118].




Figure 2. Androgen receptor signalling and how androgen deprivation therapies might block viral uptake. In the absence of androgen, the AR is located in the cytoplasm bound to a heat shock protein (HSP) complex. Ligand binding promotes dissociation of this complex, nuclear localisation and dimerisation. The AR binds to androgen response elements in the regulatory regions of target genes (e.g. TMPRSS2 and ACE2) and through the recruitment of coactivators and the basal transcriptional machinery, regulates transcription. LHRH antagonists/analogs reduce androgen production to inhibit AR signalling. Antiandrogens bind to and inhibit the AR. The down-regulation of AR signalling reduces TMPRSS2 and ACE2 expression, reducing COVID-19 entry. Elements of the schematic are from Motifolio.

6. Androgen receptor activity in the lung and regulation of TMPRSS2 and ACE2

Our understanding of AR action has predominantly come from studies that have investigated the receptor's role in the prostate, a small secretory gland at the base of the bladder, and prostate cancer [118]. In the prostate, the AR regulates various genes important in the development and function of the gland. In prostate cancer, the AR has been shown to regulate a transcriptional profile that promotes tumour growth. However, it is well documented that the AR is expressed in, and plays an important role in, other organs [117]. For example, analysis of an AR reporter mouse demonstrated that the AR is expressed and transcriptionally active in multiple organs in male and female mice, including the testes, prostate, ovaries, uterus, salivary glands and spleen [119]. Importantly, the study also demonstrated that AR is transcriptionally active in the lung, albeit weakly. Further, microarray analysis of the AR transcriptome in the lung adenocarcinoma cell line A459, demonstrated that the AR regulates pathways involved in e.g. oxygen transport, DNA repair and DNA recombination [120].

Several studies have investigated AR expression in the lung, with single cell analysis demonstrating that the receptor is expressed in multiple cell types e.g. club cells [121-123]. Importantly, the AR was shown to be co-expressed with TMPRSS2 and ACE2 in AT2 cells [122], the cell type predominantly targeted by SARS-CoV-2 [124]. The AR has been shown to regulate the expression of TMPRSS2 in multiple cell types, including prostate, breast and lung cells [71, 120, 122, 125]. Analysis of the regions up-stream of TMPRSS2 identified multiple potential AREs, with a binding site approximately 13 kb upstream from the start site being crucial for optimal androgen regulation of the gene [125]. Interestingly, chromatin immunoprecipitation analysis of AR binding up-stream regions of TMPRSS2

demonstrated that this regulation may be tissue specific [122, 126]. In the prostate cancer cell line LNCaP, the AR was found to bind to response elements present in approximately the first 80 kb upstream of TMPRSS2. However, in the immortalised lung cells A549 and H1944, the AR is bound to more distal regions; binding sites present approximately 100-200 kb upstream of the transcriptional start site [122]. ACE2 expression has also been shown to be regulated by the AR. For example, Baratchian *et al.* demonstrated that ACE2 is regulated at the RNA and protein level by the AR in LNCaP cells [121]. Similar to TMPRSS2, AR binding sites were also identified upstream of the ACE2 gene, suggesting direct gene regulation by androgens.

7. Androgen receptor modulators as treatments for COVID

AR regulation of TMPRSS2 and ACE2 expression led to speculation that androgens could be a contributing factor to SARS-CoV-2 infection and disease severity (e.g. [127]). Subsequently, it was hypothesised that inhibition of AR signalling could be a method to reduce SARS-CoV-2 infection/treat the disease [128]. Multiple inhibitors of the AR signalling pathway have been developed and these either block androgen synthesis, also known as androgen deprivation therapies (ADT, e.g. LHRH antagonists/agonists), or bind directly to the receptor and inhibit its activity (antiandrogens). These androgen signalling inhibitors are widely used in men for the treatment of prostate cancer as well as for other diseases such as benign prostatic hyperplasia and male pattern baldness/androgenetic alopecia [129, 130]. They have also been trialled/used in women for the treatment of diseases such as breast cancer, ovarian cancer and polycystic ovarian syndrome [131-134].

Antiandrogens (e.g. enzalutamide) have been shown to significantly reduce ACE2 and TMPRSS2 expression in prostate and lung cell lines [122, 126]. Castration of mice (removal of testicular androgen production) and treatment of mice with enzalutamide, has also been shown to reduce ACE2 and/or TMPRSS2 expression in the mouse lung [122, 135]. Importantly, inhibition of AR signalling has been shown to block cellular entry of pseudotyped virus expressing the SARS-CoV-2 spike protein and also the authentic virus in lung and prostate cancer cell lines [122]. However, Li *et al.* found that enzalutamide was unable to block SARS-CoV-2 infection in human lung organoids [126].

8. Retrospective/observational studies to assess the potential efficacy of ADT in relation to COVID-19

A number of observational studies, to assess the potential benefits of ADT in relation to COVID-19, have been undertaken. The first clinical data to support the hypothesis that ADT could be useful for the management of COVID-19 came from Montopoli *et al.* who analysed infection rates in prostate cancer patients in the Veneto region of Italy [136]. Comparison of 5,273 patients receiving ADT with 37,161 patients not receiving ADT found that although cancer patients were at higher risk of COVID-19 infection, ADT partially protected men from the disease. Soon after, a second but smaller study (22 prostate cancer patients receiving ADT and 36 that were not) performed at Mount Sinai Health System in New York City also found that ADT had potential as a treatment option for COVID-19 [137]. The study found that those receiving ADT were significantly less likely to require hospitalisation or oxygen support. However, ADT was not found to significantly reduce morbidity or intubation [137]. A more recent observational study of veterans in the US also demonstrated that men receiving ADT had reduced incidence of COVID-19 and were less likely to suffer from severe symptoms [138].

In addition to the analysis of prostate cancer patients receiving ADT, studies have also looked at the therapeutic potential of AR signalling inhibitors as a treatment option for COVID in other cohorts of men. For example, McCoy *et al.* investigated the potential benefit of 5-alpha-reductase inhibitors, molecules that block the conversion of testosterone into the more potent dihydrotestosterone, in men being treated for androgenetic alopecia such as spironolactone [139]. In agreement with the ADT studies, the authors found that there was a significant reduction in the frequency of COVID-19-related symptoms in the men receiving the 5-alpha-reductase inhibitor (n=48) compared to those not receiving the

therapy (n=65). However, not all epidemiological studies have supported the use of ADT as a therapy for COVID-19. For example, Welén *et al.* investigated COVID-19 severity in prostate cancer patients in the Swedish national registers, who had received different types of ADT [140]. After adjustment for age and comorbidities, the authors found no evidence that ADT protected the patients from infection, nor did they reduce the severity of the disease.

9. Clinical trials to assess the efficacy of ADT as a treatment option for COVID-19

Due to the experimental and observational clinical data investigating ADT as a therapeutic approach for COVID-19, multiple clinical trials were initiated. One study, using the novel and experimental antiandrogen Proxalutamide, showed significant promise in clinical trials. The trial demonstrated a significant improvement in survival and a reduction in hospitalisations in the group receiving the antiandrogen. However, concerns were raised about the study design and the publication has since been retracted [138]. A second larger follow-up multicentre clinical trial was performed for Proxalutamide. The trial included a total of 778 subjects who had been hospitalised with COVID-19, 423 of whom received Proxalutamide in addition to standard care. Patients who received Proxalutamide were found to have a significantly better recovery rate, reduced mortality rate and on average spent less time in hospital [139].

In contrast to the proxalutamide results, several studies investigating alternative androgen signalling inhibitors have shown limited/no benefit to patients. For example, Nickols *et al.* undertook a clinical trial to assess the therapeutic value of degarelix, which blocks androgen production through inhibition of the hypothalamus-pituitary signalling axis. The study was terminated after an interim analysis demonstrated that there was no significant difference in outcomes (e.g. mortality, ongoing need for hospitalization, or requirement for mechanical ventilation) between the deglarelix (plus standard care) and placebo groups [141]. Similarly, the COVIDENZA trial also found that inhibition of AR signalling had no therapeutic value for COVID-19. COVIDENZA was a randomized phase 2 clinical trial that investigated the efficacy of the antiandrogen enzalutamide. The trial enrolled 42 hospitalised COVID-19 patients and following a safety evaluation, the trial was halted as it was found that enzalutamide-treated patients required longer stays in hospital. A number of other clinical trials that aimed to investigate antiandrogens as a therapeutic approach for COVID-19 were subsequently withdrawn. However, at the time of writing, two proxalutamide clinical trials are ongoing (NCT05009732 and NCT04869228) and hopefully these will provide some clarity on the efficacy of targeting the AR as a treatment option for COVID-19.

10. Conclusions and further perspectives

Considering the prohibiting cost of current COVID-19 drug regimens for low- and middle-income countries, the emerging SARS-CoV-2 variants and the COVID-19 vaccine rollout and efficacy challenges, the need for cost effective, orally available and broad-spectrum antivirals remains urgent. Despite the promising antiviral effect that a range of antiandrogens display *in vitro* against SARS-CoV-2, the results of finalized clinical trials on the efficacy of ADT or antiandrogens in COVID-19 patients have not been conclusive enough to inform clinical practice. Various next generation antiandrogens have been formulated and the development of a lot more is underway including apalutamide, darolutamide, orteronel and galeterone. These new drugs should be explored for their antiviral effects and clinical outcomes as they might be more effective against SARS-CoV-2 and perhaps more amenable for widespread use in COVID-19 patients.

A disadvantage of antiandrogens as standalone therapeutic agents is that alternative TMPRSS2-independent virus entry pathways can counteract the inhibitory effects of antiandrogens to TMPRSS2-dependent viral entry. This might explain the mixed or inconclusive results of clinical trials to date which evaluated the monotherapy of ADT/antiandrogens in COVID-19 patients. The use of ADT/antiandrogens in combination therapy has not been evaluated so far. Combinations of antiviral drugs are more likely to function

synergistically if they have distinct mechanisms of action and target different stages of the virus lifecycle. Therefore, while we wait for the outcomes of the remaining ADT clinical trials and notably proxalutamide, the next wave of investigations should be focused on the combination of antiandrogen therapy with other treatments such as viral replication inhibitors.

In addition, it is surprising that experimental approaches to target TMPRSS2-independent virus entry pathways have been comparably limited and the crosstalk between androgen signalling and virus pathogenicity has not been investigated thoroughly. The concept that androgens may serve dual roles in SARS-CoV-2 infection is intriguing and remains understudied. Future efforts should be focused on simultaneous targeting alternative viral entry mechanisms and defining better the mechanistic roles of androgens in the respiratory tract. In conclusion, antiandrogens have more exploratory potential as antivirals. Building on results from former clinical trials, future trials should focus on antiandrogens or ADT in combinatorial therapeutic modalities against COVID-19.

Author Contributions: Conceptualization, data curation, writing-original draft preparation: E.S.G and G.N.B.; Data curation: E. C.. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References:

1. Cui, J.; Li, F.; Shi, Z. L., Origin and evolution of pathogenic coronaviruses. *Nat Rev Microbiol* **2019**, *17*, (3), 181-192.
2. Adjei, S.; Hong, K.; Molinari, N. M.; Bull-Otterton, L.; Ajani, U. A.; Gundlapalli, A. V.; Harris, A. M.; Hsu, J.; Kadri, S. S.; Starnes, J.; Yeoman, K.; Boehmer, T. K., Mortality Risk Among Patients Hospitalized Primarily for COVID-19 During the Omicron and Delta Variant Pandemic Periods - United States, April 2020-June 2022. *MMWR Morb Mortal Wkly Rep* **2022**, *71*, (37), 1182-1189.
3. Zhu, J.; Ji, P.; Pang, J.; Zhong, Z.; Li, H.; He, C.; Zhang, J.; Zhao, C., Clinical characteristics of 3,062 COVID-19 patients: a meta-analysis. *J Med Virol* **2020**.
4. Whitaker, M.; Elliott, J.; Chadeau-Hyam, M.; Riley, S.; Darzi, A.; Cooke, G.; Ward, H.; Elliott, P., Persistent COVID-19 symptoms in a community study of 606,434 people in England. *Nat Commun* **2022**, *13*, (1), 1957.
5. Evans, R. A.; McAuley, H.; Harrison, E. M.; Shikotra, A.; Singapuri, A.; Sereno, M.; Elneima, O.; Docherty, A. B.; Lone, N. I.; Leavy, O. C.; Daines, L.; Baillie, J. K.; Brown, J. S.; Chalder, T.; De Soyza, A.; Diar Bakerly, N.; Easom, N.; Geddes, J. R.; Greening, N. J.; Hart, N.; Heaney, L. G.; Heller, S.; Howard, L.; Hurst, J. R.; Jacob, J.; Jenkins, R. G.; Jolley, C.; Kerr, S.; Kon, O. M.; Lewis, K.; Lord, J. M.; McCann, G. P.; Neubauer, S.; Openshaw, P. J. M.; Parekh, D.; Pfeffer, P.; Rahman, N. M.; Raman, B.; Richardson, M.; Rowland, M.; Semple, M. G.; Shah, A. M.; Singh, S. J.; Sheikh, A.; Thomas, D.; Toshner, M.; Chalmers, J. D.; Ho, L. P.; Horsley, A.; Marks, M.; Poinasamy, K.; Wain, L. V.; Brightling, C. E.; Group, P.-C. C., Physical, cognitive, and mental health impacts of COVID-19 after hospitalisation (PHOSP-COVID): a UK multicentre, prospective cohort study. *Lancet Respir Med* **2021**, *9*, (11), 1275-1287.
6. Zhang, Z.; Penn, R.; Barclay, W. S.; Giotis, E. S., Naive Human Macrophages Are Refractory to SARS-CoV-2 Infection and Exhibit a Modest Inflammatory Response Early in Infection. *Viruses* **2022**, *14*, (2).
7. Montazersaheb, S.; Hosseiniyan Khatibi, S. M.; Hejazi, M. S.; Tarhriz, V.; Farjami, A.; Ghasemian Sorbeni, F.; Farahzadi, R.; Ghasemnejad, T., COVID-19 infection: an overview on cytokine storm and related interventions. *Virol J* **2022**, *19*, (1), 92.
8. Henderson, L. A.; Canna, S. W.; Schulert, G. S.; Volpi, S.; Lee, P. Y.; Kernan, K. F.; Caricchio, R.; Mahmud, S.; Hazen, M. M.; Halyabar, O.; Hoyt, K. J.; Han, J.; Grom, A. A.; Gattorno, M.; Ravelli, A.; De Benedetti, F.; Behrens, E. M.; Cron, R. Q.; Nigrovic, P. A., On the Alert for Cytokine Storm: Immunopathology in COVID-19. *Arthritis Rheumatol* **2020**, *72*, (7), 1059-1063.
9. Elrobaa, I. H.; New, K. J., COVID-19: Pulmonary and Extra Pulmonary Manifestations. *Front Public Health* **2021**, *9*.
10. Lee, M. H.; Perl, D. P.; Steiner, J.; Pasternack, N.; Li, W.; Maric, D.; Safavi, F.; Horkayne-Szakaly, I.; Jones, R.; Stram, M. N.; Moncur, J. T.; Hefti, M.; Folkerth, R. D.; Nath, A., Neurovascular injury with complement activation and inflammation in COVID-19. *Brain* **2022**, *145*, (7), 2555-2568.
11. Greenhalgh, T.; Sivan, M.; Delaney, B.; Evans, R.; Milne, R., Long covid-an update for primary care. *BMJ* **2022**, *378*, e072117.
12. Brightling, C. E.; Evans, R. A., Long COVID: which symptoms can be attributed to SARS-CoV-2 infection? *Lancet* **2022**, *400*, (10350), 411-413.
13. Huang, L.; Li, X.; Gu, X.; Zhang, H.; Ren, L.; Guo, L.; Liu, M.; Wang, Y.; Cui, D.; Wang, Y.; Zhang, X.; Shang, L.; Zhong, J.; Wang, X.; Wang, J.; Cao, B., Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study. *Lancet Respir Med* **2022**, *10*, (9), 863-876.

14. Feikin, D. R.; Abu-Raddad, L. J.; Andrews, N.; Davies, M. A.; Higdon, M. M.; Orenstein, W. A.; Patel, M. K., Assessing vaccine effectiveness against severe COVID-19 disease caused by omicron variant. Report from a meeting of the World Health Organization. *Vaccine* **2022**, *40*, (26), 3516-3527.

15. Andrews, N.; Tessier, E.; Stowe, J.; Gower, C.; Kirsebom, F.; Simmons, R.; Gallagher, E.; Thelwall, S.; Groves, N.; Dabrera, G.; Myers, R.; Campbell, C. N. J.; Amirthalingam, G.; Edmunds, M.; Zambon, M.; Brown, K.; Hopkins, S.; Chand, M.; Ladhani, S. N.; Ramsay, M.; Lopez Bernal, J., Duration of Protection against Mild and Severe Disease by Covid-19 Vaccines. *N Engl J Med* **2022**, *386*, (4), 340-350.

16. Shrestha, L. B.; Foster, C.; Rawlinson, W.; Tedla, N.; Bull, R. A., Evolution of the SARS-CoV-2 omicron variants BA.1 to BA.5: Implications for immune escape and transmission. *Rev Med Virol* **2022**, *32*, (5).

17. Joshi, N.; Tyagi, A.; Nigam, S., Molecular Level Dissection of Critical Spike Mutations in SARS-CoV-2 Variants of Concern (VOCs): A Simplified Review. *ChemistrySelect* **2021**, *6*, (31), 7981-7998.

18. Tao, K.; Tzou, P. L.; Nouhin, J.; Gupta, R. K.; de Oliveira, T.; Kosakovsky Pond, S. L.; Fera, D.; Shafer, R. W., The biological and clinical significance of emerging SARS-CoV-2 variants. *Nat Rev Genet* **2021**, *22*, (12), 757-773.

19. Plante, J. A.; Mitchell, B. M.; Plante, K. S.; Debbink, K.; Weaver, S. C.; Menachery, V. D., The variant gambit: COVID-19's next move. *Cell Host Microbe* **2021**, *29*, (4), 508-515.

20. Tegally, H.; Moir, M.; Everatt, J.; Giovanetti, M.; Scheepers, C.; Wilkinson, E.; Subramoney, K.; Makatini, Z.; Moyo, S.; Amoako, D. G.; Baxter, C.; Althaus, C. L.; Anyaneji, U. J.; Kekana, D.; Viana, R.; Giandhari, J.; Lessells, R. J.; Maponga, T.; Maruapula, D.; Choga, W.; Matshaba, M.; Mbulawa, M. B.; Msomi, N.; Naidoo, Y.; Pillay, S.; Sanko, T. J.; San, J. E.; Scott, L.; Singh, L.; Magini, N. A.; Smith-Lawrence, P.; Stevens, W.; Dor, G.; Tshiabuila, D.; Wolter, N.; Preiser, W.; Treurnicht, F. K.; Venter, M.; Chiloane, G.; McIntyre, C.; O'Toole, A.; Ruis, C.; Peacock, T. P.; Roemer, C.; Pond, S. L. K.; Williamson, C.; Pybus, O. G.; Bhiman, J. N.; Glass, A.; Martin, D. P.; Jackson, B.; Rambaut, A.; Laguda-Akingba, O.; Gaseitsiwe, S.; von Gottberg, A.; de Oliveira, T.; Bester, A. P.; Claassen, M.; Doolabh, D.; Mudau, I.; Mbhele, N.; Engelbrecht, S.; Goedhals, D.; Hardie, D.; Hsiao, N. Y.; Iranzadeh, A.; Ismail, A.; Joseph, R.; Maharaj, A.; Mahlangu, B.; Mahlakwane, K.; Davis, A.; Marais, G.; Mlisana, K.; Mnguni, A.; Mohale, T.; Motsatsi, G.; Mwangi, P.; Ntuli, N.; Nyaga, M.; Olubayo, L.; Radibe, B.; Ramphal, Y.; Ramphal, U.; Strasheim, W.; Tebeila, N.; van Wyk, S.; Wilson, S.; Lucaci, A. G.; Weaver, S.; Maharaj, A.; Pillay, Y.; Davids, M.; Mendes, A.; Mayaphi, S.; Consortium, N. S., Emergence of SARS-CoV-2 Omicron lineages BA.4 and BA.5 in South Africa. *Nat Med* **2022**.

21. Tallei, T. E.; Alhumaid, S.; AlMusa, Z.; Fatimawali; Kusumawaty, D.; Alynbawi, A.; Alshukairi, A. N.; Rabaan, A. A., Update on the omicron sub-variants BA.4 and BA.5. *Rev Med Virol* **2022**.

22. Thompson, M. G.; Natarajan, K.; Irving, S. A.; Rowley, E. A.; Griggs, E. P.; Gaglani, M.; Klein, N. P.; Grannis, S. J.; DeSilva, M. B.; Stenehjem, E.; Reese, S. E.; Dickerson, M.; Naleway, A. L.; Han, J.; Konatham, D.; McEvoy, C.; Rao, S.; Dixon, B. E.; Dascomb, K.; Lewis, N.; Levy, M. E.; Patel, P.; Liao, I. C.; Kharbanda, A. B.; Barron, M. A.; Fadel, W. F.; Grisel, N.; Goddard, K.; Yang, D. H.; Wondimu, M. H.; Murthy, K.; Valvi, N. R.; Arndorfer, J.; Fireman, B.; Dunne, M. M.; Embi, P.; Azziz-Baumgartner, E.; Zerbo, O.; Bozio, C. H.; Reynolds, S.; Ferdinand, J.; Williams, J.; Link-Gelles, R.; Schrag, S. J.; Verani, J. R.; Ball, S.; Ong, T. C., Effectiveness of a Third Dose of mRNA Vaccines Against COVID-19-Associated Emergency Department and Urgent Care Encounters and Hospitalizations Among Adults During Periods of Delta and Omicron Variant Predominance - VISION Network, 10 States, August 2021-January 2022. *MMWR Morb Mortal Wkly Rep* **2022**, *71*, (4), 139-145.

23. Khoury, D. S.; Docken, S. S.; Subbarao, K.; Kent, S. J.; Davenport, M. P.; Cromer, D., Predicting the efficacy of variant-modified COVID-19 vaccine boosters. *bioRxiv* **2022**.

24. Weinreich, D. M.; Sivapalasingam, S.; Norton, T.; Ali, S.; Gao, H.; Bhore, R.; Musser, B. J.; Soo, Y.; Rofail, D.; Im, J.; Perry, C.; Pan, C.; Hosain, R.; Mahmood, A.; Davis, J. D.; Turner, K. C.; Hooper, A. T.; Hamilton, J. D.; Baum, A.; Kyrtos, C. A.; Kim, Y.; Cook, A.; Kampman, W.; Kohli, A.; Sachdeva, Y.; Gruber, X.; Kowal, B.; DiCioccio, T.; Stahl, N.; Lipsich, L.; Braunstein, N.; Herman, G.; Yancopoulos, G. D.; Trial, I., REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with Covid-19. *N Engl J Med* **2021**, *384*, (3), 238-251.

25. Group, R. C.; Horby, P.; Lim, W. S.; Emberson, J. R.; Mafham, M.; Bell, J. L.; Linsell, L.; Staplin, N.; Brightling, C.; Ustianowski, A.; Elmahi, E.; Prudon, B.; Green, C.; Felton, T.; Chadwick, D.; Rege, K.; Fegan, C.; Chappell, L. C.; Faust, S. N.; Jaki, T.; Jeffery, K.; Montgomery, A.; Rowan, K.; Juszczak, E.; Baillie, J. K.; Haynes, R.; Landray, M. J., Dexamethasone in Hospitalized Patients with Covid-19. *N Engl J Med* **2021**, *384*, (8), 693-704.

26. Beigel, J. H.; Tomashek, K. M.; Dodd, L. E.; Mehta, A. K.; Zingman, B. S.; Kalil, A. C.; Hohmann, E.; Chu, H. Y.; Luetkemeyer, A.; Kline, S.; Lopez de Castilla, D.; Finberg, R. W.; Dierberg, K.; Tapson, V.; Hsieh, L.; Patterson, T. F.; Paredes, R.; Sweeney, D. A.; Short, W. R.; Touloumi, G.; Lye, D. C.; Ohmagari, N.; Oh, M. D.; Ruiz-Palacios, G. M.; Benfield, T.; Fatkenheuer, G.; Kortepeter, M. G.; Atmar, R. L.; Creech, C. B.; Lundgren, J.; Babiker, A. G.; Pett, S.; Neaton, J. D.; Burgess, T. H.; Bonnett, T.; Green, M.; Makowski, M.; Osinusi, A.; Nayak, S.; Lane, H. C.; Members, A.-S. G., Remdesivir for the Treatment of Covid-19 - Final Report. *N Engl J Med* **2020**, *383*, (19), 1813-1826.

27. Vangeel, L.; Chiu, W.; De Jonghe, S.; Maes, P.; Slechten, B.; Raymenants, J.; Andre, E.; Leyssen, P.; Neyts, J.; Jochmans, D., Remdesivir, Molnupiravir and Nirmatrelvir remain active against SARS-CoV-2 Omicron and other variants of concern. *Antiviral Res* **2022**, *198*, 105252.

28. Consortium, W. H. O. S. T., Remdesivir and three other drugs for hospitalised patients with COVID-19: final results of the WHO Solidarity randomised trial and updated meta-analyses. *Lancet* **2022**, *399*, (10339), 1941-1953.

29. Wang, L.; Berger, N. A.; Davis, P. B.; Kaelber, D. C.; Volkow, N. D.; Xu, R., COVID-19 rebound after Paxlovid and Molnupiravir during January-June 2022. *medRxiv* **2022**.

30. Rubin, R., From Positive to Negative to Positive Again-The Mystery of Why COVID-19 Rebounds in Some Patients Who Take Paxlovid. *JAMA* **2022**, 327, (24), 2380-2382.

31. Li, L. Q.; Huang, T.; Wang, Y. Q.; Wang, Z. P.; Liang, Y.; Huang, T. B.; Zhang, H. Y.; Sun, W.; Wang, Y., COVID-19 patients' clinical characteristics, discharge rate, and fatality rate of meta-analysis. *J Med Virol* **2020**.

32. Ioannidis, J. P. A.; Axfors, C.; Contopoulos-Ioannidis, D. G., Population-level COVID-19 mortality risk for non-elderly individuals overall and for non-elderly individuals without underlying diseases in pandemic epicenters. *Environ Res* **2020**, 188, 109890.

33. Williamson, E. J.; Walker, A. J.; Bhaskaran, K.; Bacon, S.; Bates, C.; Morton, C. E.; Curtis, H. J.; Mehrkar, A.; Evans, D.; Inglesby, P.; Cockburn, J.; McDonald, H. I.; MacKenna, B.; Tomlinson, L.; Douglas, I. J.; Rentsch, C. T.; Mathur, R.; Wong, A. Y. S.; Grieve, R.; Harrison, D.; Forbes, H.; Schultze, A.; Croker, R.; Parry, J.; Hester, F.; Harper, S.; Perera, R.; Evans, S. J. W.; Smeeth, L.; Goldacre, B., Factors associated with COVID-19-related death using OpenSAFELY. *Nature* **2020**, 584, (7821), 430-436.

34. Fauci, A. S.; Lane, H. C.; Redfield, R. R., Covid-19 - Navigating the Uncharted. *N Engl J Med* **2020**, 382, (13), 1268-1269.

35. Zhang, J. J.; Dong, X.; Liu, G. H.; Gao, Y. D., Risk and Protective Factors for COVID-19 Morbidity, Severity, and Mortality. *Clin Rev Allergy Immunol* **2022**.

36. Antonelli, M.; Pujol, J. C.; Spector, T. D.; Ourselin, S.; Steves, C. J., Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2. *Lancet* **2022**, 399, (10343), 2263-2264.

37. Antonelli, M.; Penfold, R. S.; Merino, J.; Sudre, C. H.; Molteni, E.; Berry, S.; Canas, L. S.; Graham, M. S.; Klaser, K.; Modat, M.; Murray, B.; Kerfoot, E.; Chen, L.; Deng, J.; Osterdahl, M. F.; Cheetham, N. J.; Drew, D. A.; Nguyen, L. H.; Pujol, J. C.; Hu, C.; Selvachandran, S.; Polidori, L.; May, A.; Wolf, J.; Chan, A. T.; Hammers, A.; Duncan, E. L.; Spector, T. D.; Ourselin, S.; Steves, C. J., Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study. *Lancet Infect Dis* **2022**, 22, (1), 43-55.

38. Fernandez-de-Las-Penas, C.; de-la-Llave-Rincon, A. I.; Ortega-Santiago, R.; Ambite-Quesada, S.; Gomez-Mayordomo, V.; Cuadrado, M. L.; Arias-Navalon, J. A.; Hernandez-Barrera, V.; Martin-Guerrero, J. D.; Pellicer-Valero, O. J.; Arendt-Nielsen, L., Prevalence and risk factors of musculoskeletal pain symptoms as long-term post-COVID sequelae in hospitalized COVID-19 survivors: a multicenter study. *Pain* **2022**, 163, (9), e989-e996.

39. Wenham, C.; Smith, J.; Morgan, R.; Gender; Group, C.-W., COVID-19: the gendered impacts of the outbreak. *Lancet* **2020**, 395, (10227), 846-848.

40. Stasi, V. D.; Rastrelli, G., The Role of Sex Hormones in the Disparity of COVID-19 Outcomes Based on Gender. *J Sex Med* **2021**, 18, (12), 1950-1954.

41. Jin, J. M.; Bai, P.; He, W.; Wu, F.; Liu, X. F.; Han, D. M.; Liu, S.; Yang, J. K., Gender Differences in Patients With COVID-19: Focus on Severity and Mortality. *Front Public Health* **2020**, 8, 152.

42. Xu, K.; Chen, Y.; Yuan, J.; Yi, P.; Ding, C.; Wu, W.; Li, Y.; Ni, Q.; Zou, R.; Li, X.; Xu, M.; Zhang, Y.; Zhao, H.; Zhang, X.; Yu, L.; Su, J.; Lang, G.; Liu, J.; Wu, X.; Guo, Y.; Tao, J.; Shi, D.; Yu, L.; Cao, Q.; Ruan, B.; Liu, L.; Wang, Z.; Xu, Y.; Liu, Y.; Sheng, J.; Li, L., Factors Associated With Prolonged Viral RNA Shedding in Patients with Coronavirus Disease 2019 (COVID-19). *Clin Infect Dis* **2020**, 71, (15), 799-806.

43. Chakravarty, D.; Nair, S. S.; Hammouda, N.; Ratnani, P.; Gharib, Y.; Wagaskar, V.; Mohamed, N.; Landon, D.; Dovey, Z.; Kyriyanou, N.; Tewari, A. K., Sex differences in SARS-CoV-2 infection rates and the potential link to prostate cancer. *Commun Biol* **2020**, 3, (1), 374.

44. Klein, S. L.; Huber, S., Sex differences in susceptibility to viral infection. In *Sex Hormones and Immunity to Infection*, Klein S. L., R. C. W., Ed. Springer-Verlag: Berlin, Germany, 2010; pp 93-122.

45. Klein, S. L.; Hodgson, A.; Robinson, D. P., Mechanisms of sex disparities in influenza pathogenesis. *J Leukoc Biol* **2012**, 92, (1), 67-73.

46. Klein, S. L., Sex differences in prophylaxis and therapeutic treatments for viral diseases. *Handb Exp Pharmacol* **2012**, (214), 499-522.

47. Klein, S. L.; Flanagan, K. L., Sex differences in immune responses. *Nat Rev Immunol* **2016**, 16, (10), 626-38.

48. Liao, Z. H.; Huang, T.; Xiao, J. W.; Gu, R. C.; Ouyang, J.; Wu, G.; Liao, H., Estrogen signaling effects on muscle-specific immune responses through controlling the recruitment and function of macrophages and T cells. *Skelet Muscle* **2019**, 9, (1), 20.

49. Al-Attar, A.; Presnell, S. R.; Peterson, C. A.; Thomas, D. T.; Lutz, C. T., The effect of sex on immune cells in healthy aging: Elderly women have more robust natural killer lymphocytes than do elderly men. *Mech Ageing Dev* **2016**, 156, 25-33.

50. Fish, E. N., The X-files in immunity: sex-based differences predispose immune responses. *Nat Rev Immunol* **2008**, 8, (9), 737-44.

51. Libert, C.; Dejager, L.; Pinheiro, I., The X chromosome in immune functions: when a chromosome makes the difference. *Nat Rev Immunol* **2010**, 10, (8), 594-604.

52. vom Steeg, L. G.; Klein, S. L., SeXX Matters in Infectious Disease Pathogenesis. *PLoS Pathog* **2016**, 12, (2), e1005374.

53. Oertelt-Prigione, S., The influence of sex and gender on the immune response. *Autoimmun Rev* **2012**, 11, (6-7), A479-85.

54. Straub, R. H., The complex role of estrogens in inflammation. *Endocr Rev* **2007**, *28*, (5), 521-74.

55. Gilliver, S. C., Sex steroids as inflammatory regulators. *J Steroid Biochem Mol Biol* **2010**, *120*, (2-3), 105-15.

56. Khan, D.; Ansar Ahmed, S., The Immune System Is a Natural Target for Estrogen Action: Opposing Effects of Estrogen in Two Prototypical Autoimmune Diseases. *Front Immunol* **2015**, *6*, 635.

57. Ruggieri, A.; Gagliardi, M. C.; Anticoli, S., Sex-Dependent Outcome of Hepatitis B and C Viruses Infections: Synergy of Sex Hormones and Immune Responses? *Front Immunol* **2018**, *9*, 2302.

58. Tian, Y.; Kuo, C. F.; Chen, W. L.; Ou, J. H., Enhancement of hepatitis B virus replication by androgen and its receptor in mice. *J Virol* **2012**, *86*, (4), 1904-10.

59. Jackson, C. B.; Farzan, M.; Chen, B.; Choe, H., Mechanisms of SARS-CoV-2 entry into cells. *Nat Rev Mol Cell Biol* **2022**, *23*, (1), 3-20.

60. Yadav, R.; Chaudhary, J. K.; Jain, N.; Chaudhary, P. K.; Khanra, S.; Dhamija, P.; Sharma, A.; Kumar, A.; Handu, S., Role of Structural and Non-Structural Proteins and Therapeutic Targets of SARS-CoV-2 for COVID-19. *Cells* **2021**, *10*, (4).

61. Bayati, A.; Kumar, R.; Francis, V.; McPherson, P. S., SARS-CoV-2 infects cells after viral entry via clathrin-mediated endocytosis. *J Biol Chem* **2021**, *296*, 100306.

62. Shang, J.; Wan, Y.; Luo, C.; Ye, G.; Geng, Q.; Auerbach, A.; Li, F., Cell entry mechanisms of SARS-CoV-2. *Proc Natl Acad Sci U S A* **2020**, *117*, (21), 11727-11734.

63. Hoffmann, M.; Kleine-Weber, H.; Schroeder, S.; Kruger, N.; Herrler, T.; Erichsen, S.; Schiergens, T. S.; Herrler, G.; Wu, N. H.; Nitsche, A.; Muller, M. A.; Drosten, C.; Pohlmann, S., SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **2020**, *181*, (2), 271-280.

64. Jia, H., Pulmonary Angiotensin-Converting Enzyme 2 (ACE2) and Inflammatory Lung Disease. *Shock* **2016**, *46*, (3), 239-48.

65. Giotis, E. S.; Carnell, G.; Young, E. F.; Ghanny, S.; Soteropoulos, P.; Wang, L. F.; Barclay, W. S.; Skinner, M. A.; Temperton, N., Entry of the bat influenza H17N10 virus into mammalian cells is enabled by the MHC class II HLA-DR receptor. *Nat Microbiol* **2019**, *4*, (12), 2035-2038.

66. Long, J. S.; Mistry, B.; Haslam, S. M.; Barclay, W. S., Host and viral determinants of influenza A virus species specificity. *Nat Rev Microbiol* **2019**, *17*, (2), 67-81.

67. Matsuyama, S.; Nagata, N.; Shirato, K.; Kawase, M.; Takeda, M.; Taguchi, F., Efficient activation of the severe acute respiratory syndrome coronavirus spike protein by the transmembrane protease TMPRSS2. *J Virol* **2010**, *84*, (24), 12658-64.

68. Shulla, A.; Heald-Sargent, T.; Subramanya, G.; Zhao, J.; Perlman, S.; Gallagher, T., A transmembrane serine protease is linked to the severe acute respiratory syndrome coronavirus receptor and activates virus entry. *J Virol* **2011**, *85*, (2), 873-82.

69. Belouzard, S.; Millet, J. K.; Licitra, B. N.; Whittaker, G. R., Mechanisms of coronavirus cell entry mediated by the viral spike protein. *Viruses* **2012**, *4*, (6), 1011-33.

70. Ibrahim, I. M.; Abdelmalek, D. H.; Elshahat, M. E.; Elfiky, A. A., COVID-19 spike-host cell receptor GRP78 binding site prediction. *J Infect* **2020**, *80*, (5), 554-562.

71. Lin, B.; Ferguson, C.; White, J. T.; Wang, S.; Vessella, R.; True, L. D.; Hood, L.; Nelson, P. S., Prostate-localized and androgen-regulated expression of the membrane-bound serine protease TMPRSS2. *Cancer Res* **1999**, *59*, (17), 4180-4.

72. Lucas, J. M.; Heinlein, C.; Kim, T.; Hernandez, S. A.; Malik, M. S.; True, L. D.; Morrissey, C.; Corey, E.; Montgomery, B.; Mostaghel, E.; Clegg, N.; Coleman, I.; Brown, C. M.; Schneider, E. L.; Craik, C.; Simon, J. A.; Bedalov, A.; Nelson, P. S., The androgen-regulated protease TMPRSS2 activates a proteolytic cascade involving components of the tumor microenvironment and promotes prostate cancer metastasis. *Cancer Discov* **2014**, *4*, (11), 1310-25.

73. Katopodis, P.; Kerslake, R.; Davies, J.; Randeva, H. S.; Chatha, K.; Hall, M.; Spandidos, D. A.; Anikin, V.; Polychronis, A.; Robertus, J. L.; Kyrou, I.; Karteris, E., COVID19 and SARSCoV2 host cell entry mediators: Expression profiling of TMRSS4 in health and disease. *Int J Mol Med* **2021**, *47*, (4).

74. Lukassen, S.; Chua, R. L.; Trefzer, T.; Kahn, N. C.; Schneider, M. A.; Muley, T.; Winter, H.; Meister, M.; Veith, C.; Boots, A. W.; Hennig, B. P.; Kreuter, M.; Conrad, C.; Eils, R., SARS-CoV-2 receptor ACE2 and TMPRSS2 are primarily expressed in bronchial transient secretory cells. *EMBO J* **2020**, *39*, (10), e105114.

75. Kumar, A.; Faiq, M. A.; Pareek, V.; Raza, K.; Narayan, R. K.; Prasoon, P.; Kumar, P.; Kulandhasamy, M.; Kumari, C.; Kant, K.; Singh, H. N.; Qadri, R.; Pandey, S. N.; Kumar, S., Relevance of SARS-CoV-2 related factors ACE2 and TMPRSS2 expressions in gastrointestinal tissue with pathogenesis of digestive symptoms, diabetes-associated mortality, and disease recurrence in COVID-19 patients. *Med Hypotheses* **2020**, *144*, 110271.

76. Collin, J.; Queen, R.; Zerti, D.; Dorgau, B.; Georgiou, M.; Djidrovski, I.; Hussain, R.; Coxhead, J. M.; Joseph, A.; Rooney, P.; Liso, S.; Figueiredo, F.; Armstrong, L.; Lako, M., Co-expression of SARS-CoV-2 entry genes in the superficial adult human conjunctival, limbal and corneal epithelium suggests an additional route of entry via the ocular surface. *Ocul Surf* **2021**, *19*, 190-200.

77. Zou, X.; Chen, K.; Zou, J.; Han, P.; Hao, J.; Han, Z., Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med* **2020**, *14*, (2), 185-192.

78. Vaarala, M. H.; Porvari, K. S.; Kellokumpu, S.; Kyllonen, A. P.; Vihko, P. T., Expression of transmembrane serine protease TMPRSS2 in mouse and human tissues. *J Pathol* **2001**, *193*, (1), 134-40.

79. de Kloet, A. D.; Krause, E. G.; Woods, S. C., The renin angiotensin system and the metabolic syndrome. *Physiol Behav* **2010**, *100*, (5), 525-34.

80. Samavati, L.; Uhal, B. D., ACE2, Much More Than Just a Receptor for SARS-CoV-2. *Front Cell Infect Microbiol* **2020**, *10*, 317.

81. Shen, L. W.; Qian, M. Q.; Yu, K.; Narva, S.; Yu, F.; Wu, Y. L.; Zhang, W., Inhibition of Influenza A virus propagation by benzoelenoxanthenes stabilizing TMPRSS2 Gene G-quadruplex and hence down-regulating TMPRSS2 expression. *Sci Rep* **2020**, *10*, (1), 7635.

82. Esumi, M.; Ishibashi, M.; Yamaguchi, H.; Nakajima, S.; Tai, Y.; Kikuta, S.; Sugitani, M.; Takayama, T.; Tahara, M.; Takeda, M.; Wakita, T., Transmembrane serine protease TMPRSS2 activates hepatitis C virus infection. *Hepatology* **2015**, *61*, (2), 437-46.

83. Shirato, K.; Kawase, M.; Matsuyama, S., Middle East respiratory syndrome coronavirus infection mediated by the transmembrane serine protease TMPRSS2. *J Virol* **2013**, *87*, (23), 12552-61.

84. Gierer, S.; Bertram, S.; Kaup, F.; Wrensch, F.; Heurich, A.; Kramer-Kuhl, A.; Welsch, K.; Winkler, M.; Meyer, B.; Drost, C.; Dittmer, U.; von Hahn, T.; Simmons, G.; Hofmann, H.; Pohlmann, S., The spike protein of the emerging betacoronavirus EMC uses a novel coronavirus receptor for entry, can be activated by TMPRSS2, and is targeted by neutralizing antibodies. *J Virol* **2013**, *87*, (10), 5502-11.

85. Limburg, H.; Harbig, A.; Bestle, D.; Stein, D. A.; Moulton, H. M.; Jaeger, J.; Janga, H.; Hardes, K.; Koepke, J.; Schulte, L.; Koczulla, A. R.; Schmeck, B.; Klenk, H. D.; Bottcher-Frieberthaus, E., TMPRSS2 Is the Major Activating Protease of Influenza A Virus in Primary Human Airway Cells and Influenza B Virus in Human Type II Pneumocytes. *J Virol* **2019**, *93*, (21).

86. Iwata-Yoshikawa, N.; Okamura, T.; Shimizu, Y.; Hasegawa, H.; Takeda, M.; Nagata, N., TMPRSS2 Contributes to Virus Spread and Immunopathology in the Airways of Murine Models after Coronavirus Infection. *J Virol* **2019**, *93*, (6).

87. Chen, J.; Jiang, Q.; Xia, X.; Liu, K.; Yu, Z.; Tao, W.; Gong, W.; Han, J. J., Individual variation of the SARS-CoV-2 receptor ACE2 gene expression and regulation. *Aging Cell* **2020**, *19*, (7).

88. Singh, H.; Choudhari, R.; Nema, V.; Khan, A. A., ACE2 and TMPRSS2 polymorphisms in various diseases with special reference to its impact on COVID-19 disease. *Microb Pathog* **2021**, *150*, 104621.

89. Calcagnile, M.; Forgez, P.; Iannelli, A.; Bucci, C.; Alifano, M.; Alifano, P., Molecular docking simulation reveals ACE2 polymorphisms that may increase the affinity of ACE2 with the SARS-CoV-2 Spike protein. *Biochimie* **2021**, *180*, 143-148.

90. Irham, L. M.; Chou, W. H.; Calkins, M. J.; Adikusuma, W.; Hsieh, S. L.; Chang, W. C., Genetic variants that influence SARS-CoV-2 receptor TMPRSS2 expression among population cohorts from multiple continents. *Biochem Biophys Res Commun* **2020**, *529*, (2), 263-269.

91. Saheb Sharif-Askari, N.; Saheb Sharif-Askari, F.; Alabed, M.; Temsah, M. H.; Al Hejaly, S.; Hamid, Q.; Halwani, R., Airways Expression of SARS-CoV-2 Receptor, ACE2, and TMPRSS2 Is Lower in Children Than Adults and Increases with Smoking and COPD. *Mol Ther Methods Clin Dev* **2020**, *18*, 1-6.

92. Wang, A.; Chiou, J.; Poirion, O. B.; Buchanan, J.; Valdez, M. J.; Verheyden, J. M.; Hou, X.; Kudtarkar, P.; Narendra, S.; Newsome, J. M.; Guo, M.; Faddah, D. A.; Zhang, K.; Young, R. E.; Barr, J.; Sajti, E.; Misra, R.; Huyck, H.; Rogers, L.; Poole, C.; Whitsett, J. A.; Pryhuber, G.; Xu, Y.; Gaulton, K. J.; Preissl, S.; Sun, X.; Consortium, N. L., Single-cell multiomic profiling of human lungs reveals cell-type-specific and age-dynamic control of SARS-CoV2 host genes. *Elife* **2020**, *9*.

93. Monteil, V.; Dyczynski, M.; Lauschke, V. M.; Kwon, H.; Wirnsberger, G.; Youhanna, S.; Zhang, H.; Slutsky, A. S.; Hurtado Del Pozo, C.; Horn, M.; Montserrat, N.; Penninger, J. M.; Mirazimi, A., Human soluble ACE2 improves the effect of remdesivir in SARS-CoV-2 infection. *EMBO Mol Med* **2021**, *13*, (1), e13426.

94. Jia, H.; Neptune, E.; Cui, H., Targeting ACE2 for COVID-19 Therapy: Opportunities and Challenges. *Am J Respir Cell Mol Biol* **2021**, *64*, (4), 416-425.

95. Lelis, D. F.; Freitas, D. F.; Machado, A. S.; Crespo, T. S.; Santos, S. H. S., Angiotensin-(1-7), Adipokines and Inflammation. *Metabolism* **2019**, *95*, 36-45.

96. Zhang, H.; Penninger, J. M.; Li, Y.; Zhong, N.; Slutsky, A. S., Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med* **2020**, *46*, (4), 586-590.

97. Zhang, J.; Xie, B.; Hashimoto, K., Current status of potential therapeutic candidates for the COVID-19 crisis. *Brain Behav Immun* **2020**, *87*, 59-73.

98. Kim, T. S.; Heinlein, C.; Hackman, R. C.; Nelson, P. S., Phenotypic analysis of mice lacking the Tmprss2-encoded protease. *Mol Cell Biol* **2006**, *26*, (3), 965-75.

99. Stopsack, K. H.; Mucci, L. A.; Antonarakis, E. S.; Nelson, P. S.; Kantoff, P. W., TMPRSS2 and COVID-19: Serendipity or Opportunity for Intervention? *Cancer Discov* **2020**, *10*, (6), 779-782.

100. Bestle, D.; Heindl, M. R.; Limburg, H.; Van Lam van, T.; Pilgram, O.; Moulton, H.; Stein, D. A.; Hardes, K.; Eickmann, M.; Dolnik, O.; Rohde, C.; Klenk, H. D.; Garten, W.; Steinmetzer, T.; Bottcher-Frieberthaus, E., TMPRSS2 and furin are both essential for proteolytic activation of SARS-CoV-2 in human airway cells. *Life Sci Alliance* **2020**, *3*, (9).

101. Gunst, J. D.; Staerke, N. B.; Pahus, M. H.; Kristensen, L. H.; Bodilsen, J.; Lohse, N.; Dalgaard, L. S.; Bronnum, D.; Frobert, O.; Honge, B.; Johansen, I. S.; Monrad, I.; Erikstrup, C.; Rosendal, R.; Vilstrup, E.; Mariager, T.; Bove, D. G.; Offersen, R.; Shakar, S.; Cajander, S.; Jorgensen, N. P.; Sritharan, S. S.; Breining, P.; Jespersen, S.; Mortensen, K. L.; Jensen, M. L.; Kolte, L.; Frattari, G. S.; Larsen, C. S.; Storgaard, M.; Nielsen, L. P.; Tolstrup, M.; Saedder, E. A.; Ostergaard, L. J.; Ngo, H. T. T.; Jensen, M. H.; Hojen, J. F.; Kjolby, M.; Sogaard, O. S., Efficacy of the TMPRSS2 inhibitor camostat mesilate in patients hospitalized with Covid-19-a double-blind randomized controlled trial. *EClinicalMedicine* **2021**, *35*, 100849.

102. Shapira, T.; Monreal, I. A.; Dion, S. P.; Buchholz, D. W.; Imbiakha, B.; Olmstead, A. D.; Jager, M.; Desilets, A.; Gao, G.; Martins, M.; Vandal, T.; Thompson, C. A. H.; Chin, A.; Rees, W. D.; Steiner, T.; Nabi, I. R.; Marsault, E.; Sahler, J.; Diel, D. G.; Van de Walle, G. R.; August, A.; Whittaker, G. R.; Boudreault, P. L.; Leduc, R.; Aguilar, H. C.; Jean, F., A TMPRSS2 inhibitor acts as a pan-SARS-CoV-2 prophylactic and therapeutic. *Nature* **2022**, *605*, (7909), 340-348.

103. Mahoney, M.; Damalanka, V. C.; Tartell, M. A.; Chung, D. H.; Lourenco, A. L.; Pwee, D.; Mayer Bridwell, A. E.; Hoffmann, M.; Voss, J.; Karmakar, P.; Azouz, N. P.; Klingler, A. M.; Rothlauf, P. W.; Thompson, C. E.; Lee, M.; Klampfer, L.; Stallings, C. L.; Rothenberg, M. E.; Pohlmann, S.; Whelan, S. P. J.; O'Donoghue, A. J.; Craik, C. S.; Janetka, J. W., A novel class of TMPRSS2 inhibitors potently block SARS-CoV-2 and MERS-CoV viral entry and protect human epithelial lung cells. *Proc Natl Acad Sci U S A* **2021**, *118*, (43).

104. Rahman, N.; Basharat, Z.; Yousuf, M.; Castaldo, G.; Rastrelli, L.; Khan, H., Virtual Screening of Natural Products against Type II Transmembrane Serine Protease (TMPRSS2), the Priming Agent of Coronavirus 2 (SARS-CoV-2). *Molecules* **2020**, *25*, (10).

105. Rahbar Saadat, Y.; Hosseiniyan Khatibi, S. M.; Zununi Vahed, S.; Ardalan, M., Host Serine Proteases: A Potential Targeted Therapy for COVID-19 and Influenza. *Front Mol Biosci* **2021**, *8*, 725528.

106. Hu, X.; Shrimp, J. H.; Guo, H.; Xu, M.; Chen, C. Z.; Zhu, W.; Zakharov, A. V.; Jain, S.; Shinn, P.; Simeonov, A.; Hall, M. D.; Shen, M., Discovery of TMPRSS2 Inhibitors from Virtual Screening as a Potential Treatment of COVID-19. *ACS Pharmacol Transl Sci* **2021**, *4*, (3), 1124-1135.

107. Alzain, A. A.; Elbadwi, F. A.; Alsamani, F. O., Discovery of novel TMPRSS2 inhibitors for COVID-19 using in silico fragment-based drug design, molecular docking, molecular dynamics, and quantum mechanics studies. *Inform Med Unlocked* **2022**, *29*, 100870.

108. Mahgoub, M. A.; Alnaem, A.; Fadlelmola, M.; Abo-Idris, M.; Makki, A. A.; Abdelgadir, A. A.; Alzain, A. A., Discovery of novel potential inhibitors of TMPRSS2 and Mpro of SARS-CoV-2 using E-pharmacophore and docking-based virtual screening combined with molecular dynamic and quantum mechanics. *J Biomol Struct Dyn* **2022**, *1*-14.

109. Manandhar, S.; Pai, K. S. R.; Krishnamurthy, P. T.; Kiran, A.; Kumari, G. K., Identification, virtual screening and molecular dynamic analysis of novel TMPRSS2 inhibitors from natural compound database as potential entry-blocking agents in SARS-CoV-2 therapy. *Struct Chem* **2022**, *33*, (5), 1609-1617.

110. Hofmann-Winkler, H.; Moerer, O.; Alt-Epping, S.; Brauer, A.; Buttner, B.; Muller, M.; Fricke, T.; Grundmann, J.; Harnisch, L. O.; Heise, D.; Kernchen, A.; Pressler, M.; Stephani, C.; Tampe, B.; Kaul, A.; Gartner, S.; Kramer, S.; Pohlmann, S.; Winkler, M. S., Camostat Mesylate May Reduce Severity of Coronavirus Disease 2019 Sepsis: A First Observation. *Crit Care Explor* **2020**, *2*, (11), e0284.

111. Doi, K.; Ikeda, M.; Hayase, N.; Moriya, K.; Morimura, N.; Group, C.-U. S., Nafamostat mesylate treatment in combination with favipiravir for patients critically ill with Covid-19: a case series. *Crit Care* **2020**, *24*, (1), 392.

112. Jang, S.; Rhee, J. Y., Three cases of treatment with nafamostat in elderly patients with COVID-19 pneumonia who need oxygen therapy. *Int J Infect Dis* **2020**, *96*, 500-502.

113. Breining, P.; Frolund, A. L.; Hojen, J. F.; Gunst, J. D.; Staerke, N. B.; Saedder, E.; Cases-Thomas, M.; Little, P.; Nielsen, L. P.; Sogaard, O. S.; Kjolby, M., Camostat mesylate against SARS-CoV-2 and COVID-19-Rationale, dosing and safety. *Basic Clin Pharmacol Toxicol* **2021**, *128*, (2), 204-212.

114. Punjani, N.; Flannigan, R., Androgens and COVID-19: exploring the role of testosterone replacement therapy. *Int J Impot Res* **2022**.

115. Brooke, G. N., Waxman, J. Bevan, C. L., The Role of Androgen Receptor Mutations in Progression of Prostate Cancer. In *Advances in Genome Science, Changing Views on Living Organisms*, Neri, C., Ed. Bentham Science Publishers: United Arab Emirates, 2013; Vol. 1, pp 45-65.

116. Brooke, G. N.; Bevan, C. L., The role of androgen receptor mutations in prostate cancer progression. *Current genomics* **2009**, *10*, (1), 18-25.

117. Bleach, R.; McIlroy, M., The Divergent Function of Androgen Receptor in Breast Cancer; Analysis of Steroid Mediators and Tumor Introcrinology. *Front Endocrinol (Lausanne)* **2018**, *9*, 594.

118. Feng, Q.; He, B., Androgen Receptor Signaling in the Development of Castration-Resistant Prostate Cancer. *Front Oncol* **2019**, *9*, 858.

119. Dart, D. A.; Waxman, J.; Aboagye, E. O.; Bevan, C. L., Visualising androgen receptor activity in male and female mice. *PLoS One* **2013**, *8*, (8), e71694.

120. Mikkonen, L.; Pihlajamaa, P.; Sahu, B.; Zhang, F. P.; Janne, O. A., Androgen receptor and androgen-dependent gene expression in lung. *Mol Cell Endocrinol* **2010**, *317*, (1-2), 14-24.

121. Baratchian, M.; McManus, J. M.; Berk, M. P.; Nakamura, F.; Mukhopadhyay, S.; Xu, W.; Erzurum, S.; Drazba, J.; Peterson, J.; Klein, E. A.; Gaston, B.; Sharifi, N., Androgen regulation of pulmonary AR, TMPRSS2 and ACE2 with implications for sex-discordant COVID-19 outcomes. *Sci Rep* **2021**, *11*, (1), 11130.

122. Leach, D. A.; Mohr, A.; Giotis, E. S.; Cil, E.; Isac, A. M.; Yates, L. L.; Barclay, W. S.; Zwacka, R. M.; Bevan, C. L.; Brooke, G. N., The antiandrogen enzalutamide downregulates TMPRSS2 and reduces cellular entry of SARS-CoV-2 in human lung cells. *Nat Commun* **2021**, *12*, (1), 4068.

123. McManus, J. M.; Gaston, B.; Zein, J.; Sharifi, N., Association Between Asthma and Reduced Androgen Receptor Expression in Airways. *J Endocr Soc* **2022**, *6*, (5), bvac047.

124. Rockx, B.; Kuiken, T.; Herfst, S.; Bestebroer, T.; Lamers, M. M.; Oude Munnink, B. B.; de Meulder, D.; van Amerongen, G.; van den Brand, J.; Okba, N. M. A.; Schipper, D.; van Run, P.; Leijten, L.; Sikkema, R.; Verschoor, E.; Verstrepen, B.; Bogers, W.; Langermans, J.; Drosten, C.; Fentener van Vlissingen, M.; Fouchier, R.; de Swart, R.; Koopmans, M.; Haagmans, B. L., Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate model. *Science* **2020**, *368*, (6494), 1012-1015.

125. Clinckemalie, L.; Spans, L.; Dubois, V.; Laurent, M.; Helsen, C.; Joniau, S.; Claessens, F., Androgen regulation of the TMPRSS2 gene and the effect of a SNP in an androgen response element. *Mol Endocrinol* **2013**, *27*, (12), 2028-40.

126. Li, F.; Han, M.; Dai, P.; Xu, W.; He, J.; Tao, X.; Wu, Y.; Tong, X.; Xia, X.; Guo, W.; Zhou, Y.; Li, Y.; Zhu, Y.; Zhang, X.; Liu, Z.; Aji, R.; Cai, X.; Li, Y.; Qu, D.; Chen, Y.; Jiang, S.; Wang, Q.; Ji, H.; Xie, Y.; Sun, Y.; Lu, L.; Gao, D., Distinct mechanisms for TMPRSS2 expression explain organ-specific inhibition of SARS-CoV-2 infection by enzalutamide. *Nat Commun* **2021**, *12*, (1), 866.

127. Wadman, M. Why coronavirus hits men harder: sex hormones offer clues. <https://www.science.org/content/article/why-coronavirus-hits-men-harder-sex-hormones-offer-clues>

128. Mauvais-Jarvis, F., Do Anti-androgens Have Potential as Therapeutics for COVID-19? *Endocrinology* **2021**, *162*, (8).

129. Yip, L.; Sinclair, R. D., Antiandrogen therapy for androgenetic alopecia. *Expert Review of Dermatology* **2014**, *1*, (2), 261-269.

130. Lanz, C.; Bennamoun, M.; Macek, P.; Cathelineau, X.; Sanchez-Salas, R., The importance of antiandrogen in prostate cancer treatment. *Ann Transl Med* **2019**, *7*, (Suppl 8), S362.

131. Papadatos-Pastos, D.; Dedes, K. J.; de Bono, J. S.; Kaye, S. B., Revisiting the role of antiandrogen strategies in ovarian cancer. *Oncologist* **2011**, *16*, (10), 1413-21.

132. Fioretti, F. M.; Sita-Lumsden, A.; Bevan, C. L.; Brooke, G. N., Revising the role of the androgen receptor in breast cancer. *Journal of molecular endocrinology* **2014**, *52*, (3), R257-65.

133. Anestis, A.; Zoi, I.; Papavassiliou, A. G.; Karamouzis, M. V., Androgen Receptor in Breast Cancer-Clinical and Preclinical Research Insights. *Molecules* **2020**, *25*, (2), 358.

134. Alpanes, M.; Fernandez-Duran, E.; Escobar-Morreale, H. F., Androgens and polycystic ovary syndrome. *Expert Rev Endocrinol Metab* **2012**, *7*, (1), 91-102.

135. Deng, Q.; Rasool, R. U.; Russell, R. M.; Natesan, R.; Asangani, I. A., Targeting androgen regulation of TMPRSS2 and ACE2 as a therapeutic strategy to combat COVID-19. *iScience* **2021**, *24*, (3), 102254.

136. Montopoli, M.; Zumerle, S.; Vettor, R.; Rugge, M.; Zorzi, M.; Catapano, C. V.; Carbone, G. M.; Cavalli, A.; Pagano, F.; Raga, E.; Prayer-Galetti, T.; Alimonti, A., Androgen-deprivation therapies for prostate cancer and risk of infection by SARS-CoV-2: a population-based study (N = 4532). *Ann Oncol* **2020**, *31*, (8), 1040-1045.

137. Patel, V. G.; Zhong, X.; Liaw, B.; Tremblay, D.; Tsao, C. K.; Galsky, M. D.; Oh, W. K., Does androgen deprivation therapy protect against severe complications from COVID-19? *Ann Oncol* **2020**, *31*, (10), 1419-1420.

138. Lee, K. M.; Heberer, K.; Gao, A.; Becker, D. J.; Loeb, S.; Makarov, D. V.; Gulanski, B.; DuVall, S. L.; Aslan, M.; Lee, J.; Shih, M. C.; Lynch, J. A.; Hauger, R. L.; Rettig, M., A Population-Level Analysis of the Protective Effects of Androgen Deprivation Therapy Against COVID-19 Disease Incidence and Severity. *Front Med (Lausanne)* **2022**, *9*, 774773.

139. McCoy, J.; Cadegiani, F. A.; Wambier, C. G.; Herrera, S.; Vano-Galvan, S.; Mesinkovska, N. A.; Ramos, P. M.; Shapiro, J.; Sinclair, R.; Tosti, A.; Goren, A., 5-alpha-reductase inhibitors are associated with reduced frequency of COVID-19 symptoms in males with androgenetic alopecia. *J Eur Acad Dermatol Venereol* **2021**, *35*, (4), e243-e246.

140. Welen, K.; Rosendal, E.; Gisslen, M.; Lenman, A.; Freyhult, E.; Fonseca-Rodriguez, O.; Bremell, D.; Stranne, J.; Balkhed, A. O.; Niward, K.; Repo, J.; Robinsson, D.; Henningsson, A. J.; Styrke, J.; Angelin, M.; Lindquist, E.; Allard, A.; Becker, M.; Rudolfsson, S.; Buckland, R.; Carlsson, C. T.; Bjartell, A.; Nilsson, A. C.; Ahlm, C.; Connolly, A. F.; Overby, A. K.; Josefsson, A., A Phase 2 Trial of the Effect of Antiandrogen Therapy on COVID-19 Outcome: No Evidence of Benefit, Supported by Epidemiology and In Vitro Data. *Eur Urol* **2022**, *81*, (3), 285-293.

141. Nickols, N. G.; Mi, Z.; DeMatt, E.; Biswas, K.; Clise, C. E.; Huggins, J. T.; Maraka, S.; Ambrogini, E.; Mirsaeidi, M. S.; Levin, E. R.; Becker, D. J.; Makarov, D. V.; Adorno Febles, V.; Belligund, P. M.; Al-Ajam, M.; Muthiah, M. P.; Montgomery, R. B.; Robinson, K. W.; Wong, Y. N.; Bedimo, R. J.; Villareal, R. C.; Aguayo, S. M.; Schoen, M. W.; Goetz, M. B.; Gruber, C. J.; Bhattacharya, D.; Soo Hoo, G.; Orshansky, G.; Norman, L. E.; Tran, S.; Ghayouri, L.; Tsai, S.; Geelhoed, M.; Rettig, M. B., Effect of Androgen Suppression on Clinical Outcomes in Hospitalized Men With COVID-19: The HITCH Randomized Clinical Trial. *JAMA Netw Open* **2022**, *5*, (4), e227852.