

Article

Completeness of Bethe ansatz for Gaudin models with $\mathfrak{gl}(1|1)$ symmetry and diagonal twists

Kang Lu * 

* Department of Mathematics, University of Virginia, 141 Cabell Dr, Charlottesville, VA 22903, United States; Kang.Lu@Virginia.edu;

Abstract: We study the Gaudin models with $\mathfrak{gl}(1|1)$ symmetry that are twisted by a diagonal matrix and defined on tensor products of polynomial evaluation $\mathfrak{gl}(1|1)[t]$ -modules, extending all the results of [11] to the twisted case. Namely, we give an explicit description of the algebra of Hamiltonians (Gaudin Hamiltonians) acting on tensor products of polynomial evaluation $\mathfrak{gl}(1|1)[t]$ -modules and show that there exists a bijection between common eigenvectors (up to proportionality) of the algebra of Hamiltonians and monic divisors of an explicit polynomial written in terms of the highest weights and evaluation parameters. In particular, our result implies that each common eigenspace of the algebra of Hamiltonians has dimension one. We also give dimensions of the generalized eigenspaces.

Keywords: Gaudin models; Lie superalgebras; Bethe ansatz; Pseudo-differential operators; Berezinian

1. Introduction

In the last half of a century, Gaudin models for simple Lie algebras have been intensively studied by many mathematicians and physicists using various methods which produced numerous spectacular results, see e.g. [2,3,20]. In recent years, the Gaudin models for Lie superalgebras have steadily gained attention within the mathematical community, see e.g. [5–7,11,15,17,21,24]. In this paper, we study the completeness of Bethe ansatz for $\mathfrak{gl}(1|1)$ Gaudin models with diagonal twists (quasi-periodic $\mathfrak{gl}(1|1)$ Gaudin models).

The results of this paper are quite similar to those of [11,13] with suitable modifications, following the strategy of [19,20]. Surprisingly, to the best of my knowledge, most of the previous work on Gaudin models for Lie superalgebras were done in the periodic case, except e.g. [5]. Therefore, we also need to establish the results on algebraic Bethe ansatz for $\mathfrak{gl}(1|1)$ Gaudin models in the quasi-periodic case, see Section 2.4. In particular, we show that Bethe ansatz is complete for generic evaluation parameters, see Theorem 2.5. Using the completeness of Bethe ansatz for generic parameters, we are able to describe the image of algebra of Hamiltonians (Bethe algebra) explicitly and show that the quasi-periodic $\mathfrak{gl}(1|1)$ Gaudin models are perfectly integrable, cf. [9]. Consequently, we obtain the completeness of Bethe ansatz for quasi-periodic $\mathfrak{gl}(1|1)$ Gaudin models with pairwise distinct evaluation parameters.

Note that the perfect integrability for the quasi-periodic $\mathfrak{gl}(m|n)$ Gaudin models defined on tensor products of symmetric powers of the vector representations was established in [5, Corollary 5.3] by studying duality between $\mathfrak{gl}(m|n)$ and $\mathfrak{gl}(k)$ Gaudin models and using the known results from [19]. In particular, it gives rise to the perfect integrability for the quasi-periodic $\mathfrak{gl}(1|1)$ Gaudin models defined on tensor products of polynomial modules. However, an explicit description of the image of Bethe algebra and the complete spectrum of Bethe algebra were not discussed in [5].

The paper is organized as follows. In Section 2, we fix notations and discuss basic facts of the algebraic Bethe ansatz for quasi-periodic $\mathfrak{gl}(1|1)$ Gaudin models. Then we recall the space $\mathcal{V}^{\mathfrak{G}}$ and Weyl modules, and their properties in Section 3. Section 4 contains the main theorems where we also discuss the higher Gaudin transfer matrices and the relations

Citation: Lastname, E.; Lastname, E.; Lastname, F. Title. *Journal Not Specified* 2022, 1, 0. <https://doi.org/>

Received:

Accepted:

Published:

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2022 by the authors. Submitted to *Journal Not Specified* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

between higher Gaudin transfer matrices and the first two Gaudin transfer matrices. Section 5 is dedicated to the proofs of main theorems.

2. Preliminaries

2.1. Lie superalgebra $\mathfrak{gl}(1|1)$ and its representations

A vector superspace $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is a \mathbb{Z}_2 -graded vector space. Elements of $V_{\bar{0}}$ are called *even*; elements of $V_{\bar{1}}$ are called *odd*. We write $|v| \in \{\bar{0}, \bar{1}\}$ for the parity of a homogeneous element $v \in V$. Set $(-1)^{\bar{0}} = 1$ and $(-1)^{\bar{1}} = -1$.

Consider the vector superspace $\mathbb{C}^{1|1}$, where $\dim(\mathbb{C}_{\bar{0}}^{1|1}) = 1$ and $\dim(\mathbb{C}_{\bar{1}}^{1|1}) = 1$. We choose a homogeneous basis v_1, v_2 of $\mathbb{C}^{1|1}$ such that $|v_1| = \bar{0}$ and $|v_2| = \bar{1}$. For brevity we shall write their parities as $|v_i| = |i|$. Denote by $E_{ij} \in \text{End}(\mathbb{C}^{1|1})$ the linear operator of parity $|i| + |j|$ such that $E_{ij}v_r = \delta_{jr}v_i$ for $i, j, r = 1, 2$.

The Lie superalgebra $\mathfrak{gl}(1|1)$ is spanned by elements e_{ij} , $i, j = 1, 2$, with parities $e_{ij} = |i| + |j|$ and the supercommutator relations are given by

$$[e_{ij}, e_{rs}] = \delta_{jr}e_{is} - (-1)^{(|i|+|j|)(|r|+|s|)}\delta_{is}e_{rj}.$$

Let \mathfrak{h} be the commutative Lie subalgebra of $\mathfrak{gl}(1|1)$ spanned by e_{11}, e_{22} . Denote the universal enveloping algebras of $\mathfrak{gl}(1|1)$ and \mathfrak{h} by $U(\mathfrak{gl}(1|1))$ and $U(\mathfrak{h})$, respectively.

We call a pair $\lambda = (\lambda_1, \lambda_2)$ of complex numbers a $\mathfrak{gl}(1|1)$ -weight. Set $|\lambda| = \lambda_1 + \lambda_2$. A $\mathfrak{gl}(1|1)$ -weight λ is *non-degenerate* if $\lambda_1 + \lambda_2 \neq 0$.

Let M be a $\mathfrak{gl}(1|1)$ -module. A non-zero vector $v \in M$ is called *singular* if $e_{12}v = 0$. Denote the subspace of all singular vectors of M by $(M)^{\text{sing}}$. A non-zero vector $v \in M$ is called of *weight* $\lambda = (\lambda_1, \lambda_2)$ if $e_{11}v = \lambda_1v$ and $e_{22}v = \lambda_2v$. Denote by $(M)_\lambda$ the subspace of M spanned by vectors of weight λ .

Let $\Lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$ be a sequence of $\mathfrak{gl}(1|1)$ -weights. Set $|\Lambda| = \sum_{s=1}^k |\lambda^{(s)}|$.

Denote by L_λ the irreducible $\mathfrak{gl}(1|1)$ -module generated by an even singular vector v_λ of weight λ . Then L_λ is two-dimensional if λ is non-degenerate and one-dimensional otherwise. Clearly, $\mathbb{C}^{1|1} \cong L_{\omega_1}$, where $\omega_1 = (1, 0)$, if we identify the action of e_{ij} on $\mathbb{C}^{1|1}$ with the operator E_{ij} .

A $\mathfrak{gl}(1|1)$ -module M is called a *polynomial module* if M is a submodule of $(\mathbb{C}^{1|1})^{\otimes n}$ for some $n \in \mathbb{Z}_{\geq 0}$. We say that λ is a *polynomial weight* if L_λ is a polynomial module. Weight $\lambda = (\lambda_1, \lambda_2)$ is a polynomial weight if and only if $\lambda_1, \lambda_2 \in \mathbb{Z}_{\geq 0}$ and either $\lambda_1 > 0$ or $\lambda_1 = \lambda_2 = 0$. We also write $L_{(\lambda_1, \lambda_2)}$ for L_λ .

For non-degenerate polynomial weights $\lambda = (\lambda_1, \lambda_2)$ and $\mu = (\mu_1, \mu_2)$, we have

$$L_{(\lambda_1, \lambda_2)} \otimes L_{(\mu_1, \mu_2)} = L_{(\lambda_1 + \mu_1, \lambda_2 + \mu_2)} \oplus L_{(\lambda_1 + \mu_1 - 1, \lambda_2 + \mu_2 + 1)}.$$

2.2. Current superalgebra $\mathfrak{gl}(1|1)[t]$

Denote by $\mathfrak{gl}(1|1)[t]$ the Lie superalgebra $\mathfrak{gl}(1|1) \otimes \mathbb{C}[t]$ of $\mathfrak{gl}(1|1)$ -valued polynomials with the point-wise supercommutator. Call $\mathfrak{gl}(1|1)[t]$ the *current superalgebra* of $\mathfrak{gl}(1|1)$. We identify $\mathfrak{gl}(1|1)$ with the subalgebra $\mathfrak{gl}(1|1) \otimes 1$ of constant polynomials in $\mathfrak{gl}(1|1)[t]$.

We write $e_{ij}[r]$ for $e_{ij} \otimes t^r$, $r \in \mathbb{Z}_{\geq 0}$. A basis of $\mathfrak{gl}(1|1)[t]$ is given by $e_{ij}[r]$, $i, j = 1, 2$ and $r \in \mathbb{Z}_{\geq 0}$. They satisfy the supercommutator relations

$$[e_{ij}[r], e_{kl}[s]] = \delta_{jk}e_{il}[r+s] - (-1)^{(|i|+|j|)(|k|+|l|)}\delta_{il}e_{kj}[r+s].$$

In particular, one has

$$(e_{12}[r])^2 = (e_{21}[r])^2 = 0, \quad e_{21}[r]e_{21}[s] = -e_{21}[s]e_{21}[r] \quad (2.1)$$

in the universal enveloping superalgebra $U(\mathfrak{gl}(1|1)[t])$. The universal enveloping superalgebra $U(\mathfrak{gl}(1|1)[t])$ is a Hopf superalgebra with the coproduct given by

$$\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \text{for } X \in \mathfrak{gl}(1|1)[t].$$

Let $e_{ij}(x) = \sum_{r=0}^{\infty} e_{ij}[r]x^{-r-1}$, where x is a formal variable. Then we have

$$(u - v)[e_{ij}(u), e_{rs}(v)] = -[e_{ij}, e_{rs}](u) + [e_{ij}, e_{rs}](v). \quad (2.2)$$

In particular,

$$[e_{ij}(x), e_{rs}(x)] = -\partial_x [e_{ij}, e_{rs}](x). \quad (2.3)$$

For each $a \in \mathbb{C}$, there exists an automorphism of $U(\mathfrak{gl}(1|1)[t])$, $\rho_a : e_{ij}(x) \rightarrow e_{ij}(x - a)$. Given a $\mathfrak{gl}(1|1)[t]$ -module M , denote by $M(a)$ the pull-back of M through the automorphism ρ_a .

For each $a \in \mathbb{C}$, we have the evaluation map

$$\text{ev}_a : U(\mathfrak{gl}(1|1)[t]) \rightarrow U(\mathfrak{gl}(1|1)), \quad e_{ij}(x) \mapsto e_{ij}/(x - a).$$

For a $\mathfrak{gl}(1|1)$ -module L , denote by $L(a)$ the $\mathfrak{gl}(1|1)[t]$ -module obtained by pulling back L through the evaluation map ev_a . We call $L(a)$ an *evaluation module* at a .

Given any series $\zeta(x) \in x^{-1}\mathbb{C}[x^{-1}]$, we have the one-dimensional $\mathfrak{gl}(1|1)[t]$ -module generated by an even vector v satisfying $e_{ij}(x)v = \delta_{ij}(-1)^{|j|}\zeta(x)v$. We denote this module by \mathbb{C}_{ζ} .

If b_1, \dots, b_n are pairwise distinct complex numbers and L_1, \dots, L_n are finite-dimensional irreducible $\mathfrak{gl}(1|1)$ -modules, then the $\mathfrak{gl}(1|1)[t]$ -module $\otimes_{s=1}^n L_s(b_s)$ is irreducible.

There is a natural $\mathbb{Z}_{\geq 0}$ -gradation on $U(\mathfrak{gl}(1|1)[t])$ such that $\deg(e_{ij}[r]) = r$ which induces the filtration $\mathcal{F}_0 U(\mathfrak{gl}(1|1)[t]) \subset \mathcal{F}_1 U(\mathfrak{gl}(1|1)[t]) \subset \dots \subset U(\mathfrak{gl}(1|1)[t])$, where $\mathcal{F}_s U(\mathfrak{gl}(1|1)[t])$ be the subspace of $U(\mathfrak{gl}(1|1)[t])$ spanned by all elements of degree $\leq s$.

Let M be a $\mathbb{Z}_{\geq 0}$ -graded space with finite-dimensional homogeneous components. Let $M_j \subset M$ be the homogeneous component of degree j . We call the formal power series in variable q ,

$$\text{ch}(M) = \sum_{j=0}^{\infty} \dim(M_j) q^j, \quad (2.4)$$

the *graded character* of M .

2.3. Gaudin Hamiltonians

In this section, we discuss the inhomogeneous Gaudin Hamiltonians. Throughout the paper, we shall fix two complex numbers $\mathbf{q} = (q_1, q_2)$. Moreover, we assume that $q_1 \neq q_2$, see the end of this section.

Let $\Lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$ be a sequence of polynomial $\mathfrak{gl}(1|1)$ -weights and $\mathbf{b} = (b_1, \dots, b_k)$ a sequence of distinct complex numbers, where $\lambda^{(s)} = (\alpha_s, \beta_s)$. Set $n = |\Lambda| = \sum_{s=1}^k (\alpha_s + \beta_s)$ and $L_{\Lambda} = \otimes_{s=1}^k L_{\lambda^{(s)}}$. The *quadratic Gaudin Hamiltonians* are the linear maps $\mathcal{H}_r \in \text{End}(L_{\Lambda})$ given by

$$\mathcal{H}_r := q_1 e_{11}^{(r)} + q_2 e_{22}^{(r)} + \sum_{s=1, s \neq r}^k \frac{e_{11}^{(r)} e_{11}^{(s)} - e_{12}^{(r)} e_{21}^{(s)} + e_{21}^{(r)} e_{12}^{(s)} - e_{22}^{(r)} e_{22}^{(s)}}{b_r - b_s}, \quad 1 \leq r \leq k. \quad (2.5)$$

where $e_{ab}^{(r)} = 1^{\otimes(r-1)} \otimes e_{ab} \otimes 1^{\otimes(k-r)}$.

Lemma 2.1. *The Gaudin Hamiltonians \mathcal{H}_r*

1. *are mutually commuting:* $[\mathcal{H}_r, \mathcal{H}_s] = 0$ for all r, s ;
2. *commute with the action of \mathfrak{h} :* $[\mathcal{H}_r, X] = 0$ for all r and $X \in \mathfrak{h}$.

Proof. This follows immediately from [21, Proposition 3.1] for non-twisted (i.e. $q_1 = q_2 = 0$) Gaudin Hamiltonians. \square

Instead of working on Gaudin Hamiltonians \mathcal{H}_s , we work on the generating function of Gaudin Hamiltonians,

$$\mathcal{H}(x) := \sum_{r=1}^{\infty} \mathcal{H}_r x^{-r} = q_1 e_{11}(x) + q_2 e_{22}(x) + \frac{1}{2} \sum_{a,b=1}^2 e_{ab}(x) e_{ba}(x) (-1)^{|b|}. \quad (2.6)$$

The operator $\mathcal{H}(x)$ acts on the tensor product of the evaluation $\mathfrak{gl}(1|1)[t]$ -modules

$$L_{\Lambda}(\mathbf{b}) := \bigotimes_{s=1}^k L_{\Lambda^{(s)}}(b_s).$$

Note that $L_{\Lambda}(\mathbf{b})$ and L_{Λ} are isomorphic as $\mathfrak{gl}(1|1)$ -modules via the identity map, then we have

$$\mathcal{H}(x) = \frac{1}{2} \sum_{s=1}^k \frac{\alpha_s(\alpha_s - 1) - \beta_s(\beta_s + 1)}{(x - b_s)^2} \text{Id} + \sum_{s=1}^k \frac{1}{x - b_s} \mathcal{H}_s, \quad (2.7)$$

as operators in $\text{End}(L_{\Lambda}) = \text{End}(L_{\Lambda}(\mathbf{b}))$. We call $\mathcal{H}(x)$ the *Gaudin transfer matrix*.

We are interested in finding the eigenvalues and eigenvectors of the Gaudin transfer matrix in $L_{\Lambda}(\mathbf{b})$. To be more precise, we call

$$\xi(x) = \sum_{r=1}^{\infty} \xi_r x^{-r}, \quad \xi_r \in \mathbb{C}, \quad (2.8)$$

an *eigenvalue* of $\mathcal{H}(x)$ if there exists a non-zero vector $v \in L_{\Lambda}(\mathbf{b})$ such that $\mathcal{H}_r v = \xi_r v$ for all $r \in \mathbb{Z}_{\geq 1}$. If $\xi(x)$ is a rational function, we consider it as a power series in x^{-1} as in (2.8). The vector v is called an *eigenvector* of $\mathcal{H}(x)$ corresponding to eigenvalue $\xi(x)$. We also define the *eigenspace* of $\mathcal{H}(x)$ in $L_{\Lambda}(\mathbf{b})$ corresponding to eigenvalue $\xi(x)$ as $\bigcap_{r=1}^{\infty} \ker(\mathcal{H}_r|_{L_{\Lambda}(\mathbf{b})} - \xi_r)$.

It is sufficient to consider L_{Λ} with $\beta_s = 0$ for all s . Indeed, if $L_{\Lambda}(\mathbf{b})$ is an arbitrary tensor product and

$$\xi(x) = \sum_{s=1}^k \frac{\beta_s}{x - b_s},$$

then

$$L_{\Lambda}(\mathbf{b}) \otimes \mathbb{C}_{\xi} \cong L_{\tilde{\Lambda}}(\mathbf{b}), \quad \tilde{\lambda}^{(s)} = (\alpha_s + \beta_s, 0).$$

Identify $L_{\Lambda}(\mathbf{b}) \otimes \mathbb{C}_{\xi}$ with $L_{\tilde{\Lambda}}(\mathbf{b})$ as vector spaces. Then $\mathcal{H}(x)$ acting on $L_{\Lambda}(\mathbf{b}) \otimes \mathbb{C}_{\xi}$ coincides with $\mathcal{H}(x) + \xi(x)(e_{11}(x) + e_{22}(x)) + (q_1 - q_2)\xi(x)$ acting on $L_{\tilde{\Lambda}}(\mathbf{b})$. Note that the coefficients of $e_{11}(x) + e_{22}(x)$ are central in $U(\mathfrak{gl}(1|1)[t])$ and hence $e_{11}(x) + e_{22}(x)$ acts on $L_{\tilde{\Lambda}}(\mathbf{b})$ by the scalar series

$$\sum_{s=1}^k \frac{\alpha_s + \beta_s}{x - b_s},$$

therefore the problem of diagonalization of the Gaudin transfer matrix in $L_{\Lambda}(\mathbf{b})$ is reduced to diagonalization of the Gaudin transfer matrix in $L_{\tilde{\Lambda}}(\mathbf{b})$.

Again by the fact that the coefficients of $e_{11}(x) + e_{22}(x)$ are central, if $q_1 = q_2$, then the diagonalization problem of $\mathcal{H}(x)$ is the same as the one for homogeneous case $q_1 = q_2 = 0$ which has been discussed in [11]. Thus, for the rest of the paper, we shall assume that $q_1 \neq q_2$.

Since L_{λ} is one-dimensional if λ is degenerate, similarly, it suffices to consider the case that all participant $\mathfrak{gl}(1|1)$ -weights are non-degenerate. Hence, we shall always assume throughout the paper that $\lambda^{(s)}$ are non-degenerate for all $1 \leq s \leq k$.

2.4. Bethe ansatz

The main method to find eigenvalues and eigenvectors of the Gaudin transfer matrix in L_Λ is the algebraic Bethe ansatz. We give the results for algebraic Bethe ansatz of quasi-periodic $\mathfrak{gl}(1|1)$ Gaudin models in this section, following e.g. [21, Section VI].

Fix a non-negative integer l . Let $\mathbf{t} = (t_1, \dots, t_l)$ be a sequence of complex numbers. Define the polynomial $y_{\mathbf{t}} = \prod_{i=1}^l (x - t_i)$. We say that polynomial $y_{\mathbf{t}}$ represents \mathbf{t} .

Set

$$\zeta_{\Lambda, \mathbf{b}}(x) := q_1 - q_2 + \sum_{s=1}^k \frac{\alpha_s + \beta_s}{x - b_s}. \quad (2.9)$$

A sequence of complex numbers \mathbf{t} is called a *solution to the Bethe ansatz equation associated to Λ, \mathbf{b}, l* if

$$y_{\mathbf{t}}(x) \text{ divides the polynomial } \varphi_{\Lambda, \mathbf{b}}(x) := \zeta_{\Lambda, \mathbf{b}}(x) \prod_{s=1}^k (x - b_s). \quad (2.10)$$

We do not distinguish solutions which differ by a permutation of coordinates (that is represented by the same polynomial).

Let v_s be the highest weight vector of $L_{\lambda^{(s)}}$, and set $|0\rangle = v_1 \otimes \dots \otimes v_k$. We call $|0\rangle$ the *vacuum vector*.

Define the *off-shell Bethe vector* $\mathbb{B}_l(\mathbf{t}) \in (L_\Lambda)_{(n-l, l)}$ by

$$\mathbb{B}_l(\mathbf{t}) = e_{21}(t_1) \cdots e_{21}(t_l) |0\rangle. \quad (2.11)$$

Since $e_{21}(x)e_{21}(u) = -e_{21}(u)e_{21}(x)$, the order of t_i is not important. Moreover, the off-shell Bethe vector is zero if $t_i = t_j$ for some $1 \leq i \neq j \leq l$.

If \mathbf{t} is a solution of the Bethe ansatz equation (2.10), we call $\mathbb{B}_l(\mathbf{t})$ an *on-shell Bethe vector*.

Let \mathbf{t} be a solution of the Bethe ansatz equation associated to Λ, \mathbf{b}, l .

Theorem 2.2. *If the on-shell Bethe vector $\mathbb{B}_l(\mathbf{t})$ is non-zero, then $\mathbb{B}_l(\mathbf{t})$ is an eigenvector of the Gaudin transfer matrix $\mathcal{H}(x)$ with the corresponding eigenvalue*

$$\mathcal{E}_{y_{\mathbf{t}}, \Lambda, \mathbf{b}}(x) = \frac{1}{2} \zeta'_{\Lambda, \mathbf{b}}(x) - \zeta_{\Lambda, \mathbf{b}}(x) \frac{y'_{\mathbf{t}}(x)}{y_{\mathbf{t}}(x)} + \sum_{r,s=1}^k \frac{\alpha_r \alpha_s - \beta_r \beta_s}{2(x - b_r)(x - b_s)} + \sum_{s=1}^k \frac{q_1 \alpha_s + q_2 \beta_s}{x - b_s}. \quad (2.12)$$

where $\zeta_{\Lambda, \mathbf{b}}(x)$ is given by (2.9).

Proof. By (2.2) and the fact that coefficients of $e_{11}(x) + e_{22}(x)$ are central in $U(\mathfrak{gl}(1|1)[t])$, we have

$$[\mathcal{H}(x), e_{21}(t)] = -\frac{1}{u-t} (\zeta_{\Lambda, \mathbf{b}}(x)e_{21}(t) - \zeta_{\Lambda, \mathbf{b}}(t)e_{21}(x)),$$

as operators on $L_\Lambda(\mathbf{b})$. Note that if t is a coordinate of a solution of the Bethe ansatz equation, then $\zeta_{\Lambda, \mathbf{b}}(t) = 0$. Therefore, we have

$$[\mathcal{H}(x), e_{21}(t_i)] = -\frac{1}{u-t} \zeta_{\Lambda, \mathbf{b}}(x) e_{21}(t_i)$$

for $1 \leq i \leq l$. Hence, we conclude that

$$\mathcal{H}(x) \mathbb{B}_l(\mathbf{t}) = -\zeta_{\Lambda, \mathbf{b}}(x) \sum_{j=1}^l \frac{1}{x - t_j} \mathbb{B}_l(\mathbf{t}) + e_{21}(t_1) \cdots e_{21}(t_l) \mathcal{H}(x) |0\rangle.$$

The theorem now follows from the straightforward computation of the eigenvalue of $\mathcal{H}(x)$ corresponding to the vector $|0\rangle$. \square

Consider another Gaudin transfer matrix

$$\mathcal{T}(x) = \frac{1}{2}(\dot{e}_{11}(x) + \dot{e}_{22}(x)) + \frac{1}{2}(e_{11}(x) + e_{22}(x))^2 + q_1(e_{11}(x) + e_{22}(x)) - \mathcal{H}(x), \quad (2.13)$$

where $\dot{e}_{ii}(x) = \partial_x(e_{ii}(x))$, $i = 1, 2$. Then the eigenvalue of $\mathcal{T}(x)$ acting on the on-shell Bethe vector $\mathbb{B}_l(\mathbf{t})$ is

$$\begin{aligned} \mathcal{E}_{y_t, \Lambda, \mathbf{b}}(x) &= \zeta_{\Lambda, \mathbf{b}}(x) \frac{y'_t(x)}{y_t(x)} + \sum_{r,s=1}^k \frac{\alpha_r \beta_s + \alpha_s \beta_r + 2\beta_r \beta_s}{2(x-b_r)(x-b_s)} + \sum_{s=1}^k \frac{(q_1 - q_2)\beta_s}{x-b_s} \\ &= \zeta_{\Lambda, \mathbf{b}}(x) \left(\frac{y'_t(x)}{y_t(x)} + \sum_{s=1}^k \frac{\beta_s}{x-b_s} \right). \end{aligned} \quad (2.14)$$

It is important to know if the on-shell Bethe vectors are non-zero. 128

Proposition 2.3. *Suppose the polynomial $\varphi_{\Lambda, \mathbf{b}}(x)$ only has simple roots, then the on-shell Bethe vector $\mathbb{B}_l(\mathbf{t})$ is nonzero.* 129
130

Proof. Since $\varphi_{\Lambda, \mathbf{b}}(x)$ only has simple roots, we have $t_i \neq t_j$ for $i \neq j$. Note that b_s are distinct and $\alpha_s + \beta_s > 0$ (since the weights are nondegenerate by our assumption), we have $b_s \neq t_i$. Hence $\zeta_{\Lambda, \mathbf{b}}(t_i) = 0$. Moreover, we have

$$0 \neq \varphi'_{\Lambda, \mathbf{b}}(t_i) = \zeta'_{\Lambda, \mathbf{b}}(t_i) \prod_{s=1}^k (t_i - b_s) + \zeta_{\Lambda, \mathbf{b}}(t_i) \left(\prod_{s=1}^k (x - b_s) \right)' \Big|_{x=t_i} = \zeta'_{\Lambda, \mathbf{b}}(t_i) \prod_{s=1}^k (t_i - b_s).$$

Therefore $\zeta'_{\Lambda, \mathbf{b}}(t_i) \neq 0$. 131

By (2.2) and the fact that coefficients of $e_{11}(x) + e_{22}(x)$ are central in $U(\mathfrak{gl}(1|1)[t])$, we have

$$[e_{12}(t), e_{21}(\tilde{t})] = -\frac{1}{t-\tilde{t}} (\zeta_{\Lambda, \mathbf{b}}(t) - \zeta_{\Lambda, \mathbf{b}}(\tilde{t}))$$

as operators on $L_{\Lambda}(\mathbf{b})$. Therefore, we have $[e_{12}(t), e_{21}(\tilde{t})] = 0$ if t and \tilde{t} are distinct coordinates of \mathbf{t} while $[e_{12}(t), e_{21}(t)] = -\zeta'_{\Lambda, \mathbf{b}}(t)$. One finds that

$$e_{12}(t_1) \cdots e_{12}(t_1) e_{21}(t_1) \cdots e_{21}(t_1) |0\rangle = (-1)^l \prod_{i=1}^l \zeta'_{\Lambda, \mathbf{b}}(t_i) |0\rangle \neq 0,$$

completing the proof. \square 132

The conjecture of completeness of Bethe ansatz for Gaudin models associated with $\mathfrak{gl}(1|1)$ was formulated as follows, cf. [6, Conjecture 8.2]. 133
134

Conjecture 2.4. *Suppose all weights $\lambda^{(s)}$, $1 \leq s \leq k$, are polynomial $\mathfrak{gl}(1|1)$ -weights. Then the Gaudin transfer matrix $\mathcal{H}(x)$ has a simple spectrum in $L_{\Lambda}(\mathbf{b})$. There exists a bijective correspondence between the monic divisors y of the polynomial $\varphi_{\Lambda, \mathbf{b}}$ and the eigenvectors v of the Gaudin transfer matrices (up to multiplication by a non-zero constant). Moreover, this bijection is such that $\mathcal{H}(x)v = \mathcal{E}_{y, \Lambda, \mathbf{b}}(x)v$, where $\mathcal{E}_{y, \Lambda, \mathbf{b}}(x)$ is given by (2.12). \square* 135
136
137
138
139

By simple spectrum we mean that if v_1, v_2 are eigenvectors of $\mathcal{H}(x)$ and $v_1 \neq cv_2$, $c \in \mathbb{C}^\times$, then the eigenvalues of $\mathcal{H}(x)$ on v_1 and v_2 are different. 140
141

The conjecture follows from Theorem 4.8 proved in Section 5.3. 142

The conjecture is clear for the case when $\varphi_{\Lambda, \mathbf{b}}$ only has simple roots. Note that $\dim L_{\Lambda}(\mathbf{b}) = 2^k$. If the polynomial $\varphi_{\Lambda, \mathbf{b}}$ has no multiple roots, then $\varphi_{\Lambda, \mathbf{b}}$ has the desired number of distinct monic divisors. Therefore, we have desired the number of on-shell Bethe vectors which are also nonzero by Proposition 2.3. By Theorem 2.2, it implies that we do have an eigenbasis of the Gaudin transfer matrix consisting of on-shell Bethe vectors 143
144
145
146
147

in $L_\Lambda(\mathbf{b})$ with different eigenvalues. Thus the algebraic Bethe ansatz works well for this situation. 148
149

Theorem 2.5. *Suppose all weights $\lambda^{(s)}$, $1 \leq s \leq k$, are polynomial $\mathfrak{gl}(1|1)$ -weights. If the polynomial $\varphi_{\Lambda, \mathbf{b}}$ has no multiple roots, then the Gaudin transfer matrix $\mathcal{H}(x)$ is diagonalizable and the Bethe ansatz is complete. In particular, for any given Λ and generic \mathbf{b} , the Gaudin transfer matrix $\mathcal{H}(x)$ is diagonalizable and the Bethe ansatz is complete. \square* 150
151
152
153

3. Space $\mathcal{V}^\mathfrak{S}$ and Weyl modules 154

In this section, we discuss the super-analog of \mathcal{V}^S in [20, Section 2.5], cf. [13, Section 3]. 155

The symmetric group \mathfrak{S}_n acts naturally on $\mathbb{C}[z_1, \dots, z_n]$ by permuting variables. Denote by $\sigma_i(z)$ the i -th elementary symmetric polynomial in z_1, \dots, z_n . The algebra of symmetric polynomials $\mathbb{C}[z_1, \dots, z_n]^\mathfrak{S}$ is freely generated by $\sigma_1(z), \dots, \sigma_n(z)$. 156
157
158

Fix $\ell \in \{0, 1, \dots, n\}$. We have a subgroup $\mathfrak{S}_\ell \times \mathfrak{S}_{n-\ell} \subset \mathfrak{S}_n$. Then \mathfrak{S}_ℓ permutes the first ℓ variables while $\mathfrak{S}_{n-\ell}$ permutes the last $n - \ell$ variables. Denote by $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}_\ell \times \mathfrak{S}_{n-\ell}}$ the subalgebra of $\mathbb{C}[z_1, \dots, z_n]$ consisting of $\mathfrak{S}_\ell \times \mathfrak{S}_{n-\ell}$ -invariant polynomials. It is known that $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}_\ell \times \mathfrak{S}_{n-\ell}}$ is a free $\mathbb{C}[z_1, \dots, z_n]^\mathfrak{S}$ -module of rank $\binom{n}{\ell}$. 159
160
161
162

3.1. Definition of $\mathcal{V}^\mathfrak{S}$ 163

Let $V = (\mathbb{C}^{1|1})^{\otimes n}$ be the tensor power of the vector representation of $\mathfrak{gl}(1|1)$. The $\mathfrak{gl}(1|1)$ -module V has weight decomposition

$$V = \bigoplus_{\ell=0}^n (V)_{(n-\ell, \ell)}.$$

Let \mathcal{V} be the space of polynomials in variables $\mathbf{z} = (z_1, z_2, \dots, z_n)$ with coefficients in V ,

$$\mathcal{V} = V \otimes \mathbb{C}[z_1, z_2, \dots, z_n].$$

The space V is identified with the subspace $V \otimes 1$ of constant polynomials in \mathcal{V} . The space \mathcal{V} has a natural grading induced from the grading on $\mathbb{C}[z_1, \dots, z_n]$ with $\deg(z_i) = 1$. Namely, the degree of an element $v \otimes p$ in \mathcal{V} is given by the degree of the polynomial p , $\deg(v \otimes p) = \deg p$. Clearly, the space $\text{End}(\mathcal{V})$ has a gradation structure induced from that on \mathcal{V} . 164
165
166
167
168

Let $P^{(i,j)}$ be the graded flip operator which acts on the i -th and j -th factors of V . Let s_1, s_2, \dots, s_{n-1} be the simple permutations of the symmetric group \mathfrak{S}_n . Define the \mathfrak{S}_n -action on \mathcal{V} by the rule:

$$s_i : f(z_1, \dots, z_n) \mapsto P^{(i,i+1)} f(z_1, \dots, z_{i+1}, z_i, \dots, z_n),$$

for $f(z_1, \dots, z_n) \in \mathcal{V}$. Note that the \mathfrak{S}_n -action respects the gradation on \mathcal{V} . Denote the subspace of all vectors in \mathcal{V} invariant with respect to the \mathfrak{S}_n -action by $\mathcal{V}^\mathfrak{S}$. 169
170

Clearly, the $\mathfrak{gl}(1|1)$ -action on \mathcal{V} commutes with the \mathfrak{S}_n -action on \mathcal{V} and preserves the grading. Therefore, $\mathcal{V}^\mathfrak{S}$ is a graded $\mathfrak{gl}(1|1)$ -module. Hence we have the weight decomposition for both $\mathcal{V}^\mathfrak{S}$ and $(\mathcal{V}^\mathfrak{S})^{\text{sing}}$,

$$\mathcal{V}^\mathfrak{S} = \bigoplus_{\ell=0}^n (\mathcal{V}^\mathfrak{S})_{(n-\ell, \ell)}, \quad (\mathcal{V}^\mathfrak{S})^{\text{sing}} = \bigoplus_{\ell=0}^n (\mathcal{V}^\mathfrak{S})_{(n-\ell, \ell)}^{\text{sing}}.$$

Note that $(\mathcal{V}^\mathfrak{S})_{(n-\ell, \ell)}$ and $(\mathcal{V}^\mathfrak{S})_{(n-\ell, \ell)}^{\text{sing}}$ are also graded $\mathbb{C}[z_1, \dots, z_n]^\mathfrak{S}$ -modules. 171

The space \mathcal{V} is a $\mathfrak{gl}(1|1)[t]$ -module where $e_{ij}[r]$ acts by

$$\begin{aligned} e_{ij}[r](p(z_1, \dots, z_n)w_1 \otimes \cdots \otimes w_n) \\ = p(z_1, \dots, z_n) \sum_{s=1}^n (-1)^{(|w_1| + \cdots + |w_{s-1}|)(|i| + |j|)} z_s^r w_1 \otimes \cdots \otimes e_{ij} w_s \otimes \cdots \otimes w_n, \end{aligned} \quad (3.1)$$

for $p(z_1, \dots, z_n) \in \mathbb{C}[z_1, \dots, z_n]$ and $w_s \in \mathbb{C}^{1|1}$. 172

Lemma 3.1. *The $\mathfrak{gl}(1|1)[t]$ -action on \mathcal{V} commutes with the \mathfrak{S}_n -action on \mathcal{V} . Both \mathcal{V} and $\mathcal{V}^{\mathfrak{S}}$ are graded $\mathfrak{gl}(1|1)[t]$ -modules. \square* 173
174

3.2. Properties of $\mathcal{V}^{\mathfrak{S}}$ and $(\mathcal{V}^{\mathfrak{S}})^{\text{sing}}$

175

In this section, we recall properties of $\mathcal{V}^{\mathfrak{S}}$ and $(\mathcal{V}^{\mathfrak{S}})^{\text{sing}}$ from [13, Section 3]. 176

Lemma 3.2. *The space $(\mathcal{V}^{\mathfrak{S}})_{(n-\ell, \ell)}$ is a free $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$ -module of rank $\binom{n}{\ell}$. In particular, the space $\mathcal{V}^{\mathfrak{S}}$ is a free $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$ -module of rank 2^n . \square* 177
178

$$\text{Set } v^+ = v_1^{\otimes n} = v_1 \otimes \cdots \otimes v_1. \quad 179$$

Lemma 3.3. *The $\mathfrak{gl}(1|1)[t]$ -module $\mathcal{V}^{\mathfrak{S}}$ is a cyclic module generated by v^+ . \square* 180

Lemma 3.4. *The set*

$$\{e_{21}[r_1]e_{21}[r_2] \cdots e_{21}[r_\ell]v^+ \mid 0 \leq r_1 < r_2 < \cdots < r_\ell \leq n-1\} \quad (3.2)$$

is a free generating set of $(\mathcal{V}^{\mathfrak{S}})_{(n-\ell, \ell)}$ over $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$. \square 181

Lemma 3.5. *The space $(\mathcal{V}^{\mathfrak{S}})^{\text{sing}}_{(n-\ell, \ell)}$ is a free $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$ -module of rank $\binom{n-1}{\ell}$ with a free generating set given by*

$$\{e_{12}[0]e_{21}[0]e_{21}[r_1] \cdots e_{21}[r_\ell]v^+, \quad 1 \leq r_1 < r_2 < \cdots < r_\ell \leq n-1\}. \quad (3.3)$$

In particular, the space $(\mathcal{V}^{\mathfrak{S}})^{\text{sing}}$ is a free $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$ -module of rank 2^{n-1} . \square 182

$$\text{Set } (q)_r = \prod_{i=1}^r (1 - q^i). \quad 183$$

Proposition 3.6. *We have*

$$\text{ch}((\mathcal{V}^{\mathfrak{S}})_{(n-\ell, \ell)}) = \frac{q^{\ell(\ell-1)/2}}{(q)_\ell (q)_{n-\ell}}, \quad \text{ch}((\mathcal{V}^{\mathfrak{S}})^{\text{sing}}_{(n-\ell, \ell)}) = \frac{q^{\ell(\ell+1)/2}}{(q)_\ell (q)_{n-1-\ell} (1 - q^n)}.$$

Given $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{C}^n$, let $I_{\mathbf{a}}$ be the ideal of $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$ generated by $\sigma_i(\mathbf{z}) - a_i$, $i = 1, \dots, n$. Then for any \mathbf{a} , by Lemmas 3.1 and 3.2, the quotient space $\mathcal{V}^{\mathfrak{S}}/I_{\mathbf{a}}\mathcal{V}^{\mathfrak{S}}$ is a $\mathfrak{gl}(1|1)[t]$ -module of dimension 2^n over \mathbb{C} . Denote by \bar{v}^+ be the image of v^+ under this quotient. 184
185
186
187

3.3. Weyl modules

188

In this section, we recall a special family of Weyl modules for $\mathfrak{gl}(1|1)[t]$ and their properties from [11, Section 3.3]. 189
190

Let $\eta(x)$ be a monic polynomial of degree m with complex coefficients, where $m \in \mathbb{Z}_{\geq 0}$,

$$\eta(x) = \sum_{i=0}^m \gamma_i x^i, \quad \gamma_m = 1.$$

Denote by W_η the $\mathfrak{gl}(1|1)[t]$ -module generated by an even vector w subject to the relations:

$$e_{11}(x)w = \eta'(x)/\eta(x)w, \quad e_{22}(x)w = e_{12}(x)w = 0, \quad (3.4)$$

$$\sum_{i=0}^m \gamma_i e_{21}[i]w = 0. \quad (3.5)$$

It is convenient to write (3.5) as $(e_{21} \otimes \eta(t))w = 0$.

Clearly, we have $\dim W_\eta \leq 2^m$ by PBW theorem and (2.1), (3.5). The module W_η is the universal $\mathfrak{gl}(1|1)[t]$ -module satisfying (3.4), (3.5) which we call a *Weyl module*.

If $\eta(x) = (x - b)^m$, we write W_η as $W_m(b)$.

Lemma 3.7. Let $\mathbf{a} = (0, \dots, 0) \in \mathbb{C}^n$. Then $\mathcal{V}^\mathfrak{S} / I_{\mathbf{a}}\mathcal{V}^\mathfrak{S}$ is isomorphic to $W_n(0)$ as $\mathfrak{gl}(1|1)[t]$ -modules. \square

In particular, we have $\dim W_m(b) = 2^m$.

Lemma 3.8. Let $\eta(x) = \prod_{s=1}^k (x - b_s)^{n_s}$, where $b_s \neq b_r$ for $1 \leq s \neq r \leq k$. Then W_η is isomorphic to $\otimes_{s=1}^k W_{n_s}(b_s)$ as $\mathfrak{gl}(1|1)[t]$ -modules. \square

Given sequences $\mathbf{n} = (n_1, \dots, n_k)$ of nonnegative integers and $\mathbf{b} = (b_1, \dots, b_s)$ of distinct complex numbers, by Lemma 3.8, we call $\otimes_{s=1}^k W_{n_s}(b_s)$ the *Weyl module associated with \mathbf{n} and \mathbf{b}* .

Given $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{C}^n$, define $k \in \mathbb{Z}_{>0}$, $b_s \in \mathbb{C}$, and $n_s \in \mathbb{Z}_{>0}$ for $1 \leq s \leq k$ by

$$x^n + \sum_{i=1}^n (-1)^i a_i x^{n-i} = \prod_{s=1}^k (x - b_s)^{n_s}, \quad (3.6)$$

where b_1, \dots, b_k are distinct. Note that $n = \sum_{s=1}^k n_s$.

Lemma 3.9. The $\mathfrak{gl}(1|1)[t]$ -module $\mathcal{V}^\mathfrak{S} / I_{\mathbf{a}}\mathcal{V}^\mathfrak{S}$ is isomorphic to $\otimes_{s=1}^k W_{n_s}(b_s)$. \square

We also need the following statements.

Lemma 3.10. Let $b \in \mathbb{C}$. We have the following properties for $W_m(b)$.

1. As a $\mathfrak{gl}(1|1)$ -module, $W_m(b)$ is isomorphic to $(\mathbb{C}^{1|1})^{\otimes m}$.
2. A $\mathfrak{gl}(1|1)[t]$ -module M is an irreducible subquotient of $W_m(b)$ if and only if M has the form $L_\lambda(b)$, where λ is a polynomial weight such that $|\lambda| = m$. \square

Corollary 3.11. A $\mathfrak{gl}(1|1)[t]$ -module M is an irreducible subquotient of $\otimes_{s=1}^k W_{n_s}(b_s)$ if and only if M has the form $\otimes_{s=1}^k L_{\lambda^{(s)}}(b_s)$, where $\lambda^{(s)}$ is a polynomial weight such that $|\lambda^{(s)}| = n_s$ for each $1 \leq s \leq k$. \square

4. Main theorems

4.1. The algebra \mathcal{O}_l

Let Ω_l be the n -dimensional affine space with coordinates $f_1, \dots, f_l, g_1, \dots, g_{n-l}$. Introduce two polynomials

$$f(x) = x^l + \sum_{i=1}^l f_i x^{l-i}, \quad g(x) = x^{n-l} + \sum_{i=1}^{n-l} g_i x^{n-l-i}. \quad (4.1)$$

Denote by \mathcal{O}_l the algebra of regular functions on Ω_l , namely

$$\mathcal{O}_l = \mathbb{C}[f_1, \dots, f_l, g_1, \dots, g_{n-l-1}, g_{n-l}].$$

Define the degree function by

$$\deg f_i = i, \quad \deg g_j = j,$$

for all $i = 1, \dots, l$ and $j = 1, \dots, n - l$. The algebra \mathcal{O}_l is graded with the graded character given by

$$\text{ch}(\mathcal{O}_l) = \frac{1}{(q)_l(q)_{n-l}}. \quad (4.2)$$

Let $\mathcal{F}_0\mathcal{O}_l \subset \mathcal{F}_1\mathcal{O}_l \subset \dots \subset \mathcal{O}_l$ be the increasing filtration corresponding to this grading, where $\mathcal{F}_s\mathcal{O}_l$ consists of elements of degree at most s . 215

Let $\Sigma_1, \dots, \Sigma_n$ be the elements of \mathcal{O}_l such that 216

$$(q_1 - q_2)f(x)g(x) = (q_1 - q_2)x^n + \sum_{i=1}^n (-1)^i ((q_1 - q_2)\Sigma_i - (n + 1 - i)\Sigma_{i-1})x^{n-i}, \quad (4.3)$$

where $\Sigma_0 = 1$. The homomorphism

$$\pi_l : \mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}} \rightarrow \mathcal{O}_l, \quad \sigma_i(z) \mapsto \Sigma_i, \quad i = 1, \dots, n, \quad (4.4)$$

is injective and induces a $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$ -module structure on \mathcal{O}_l . 217

Express $f'(x)g(x)$ as follows,

$$(q_1 - q_2)f'(x)g(x) = (q_1 - q_2)lx^{n-1} + \sum_{i=1}^{n-1} G_i x^{n-1-i}, \quad (4.5)$$

where $G_i \in \mathcal{O}_l$. 218

Lemma 4.1. *The elements G_i and Σ_j , $i = 1, \dots, n - 1$, $j = 1, \dots, n$, generate the algebra \mathcal{O}_l . \square* 219

Lemma 4.2. *We have $G_i \in \mathcal{F}_i\mathcal{O}_l \setminus \mathcal{F}_{i-1}\mathcal{O}_l$ and $\Sigma_j \in \mathcal{F}_j\mathcal{O}_l \setminus \mathcal{F}_{j-1}\mathcal{O}_l$, $i = 1, \dots, n - 1$, $j = 1, \dots, n$. \square* 220

4.2. Bethe algebra 222

We call the unital subalgebra of $U(\mathfrak{gl}(1|1)[t])$ generated by the coefficients of

$$e_{11}(x) + e_{22}(x), \quad \mathcal{H}(x) = q_1 e_{11}(x) + q_2 e_{22}(x) + \frac{1}{2} \sum_{i,j=1}^2 e_{ij}(x)e_{ji}(x)(-1)^{|j|}$$

the *Bethe algebra*. We denote the Bethe algebra by \mathcal{B} . Note that the coefficients of $e_{11}(x) + e_{22}(x)$ generate the center of $U(\mathfrak{gl}(1|1)[t])$. 223

Lemma 4.3 ([17]). *The Bethe algebra \mathcal{B} is commutative. The Bethe algebra \mathcal{B} commutes with the subalgebra $U(\mathfrak{h}) \subset U(\mathfrak{gl}(1|1)[t])$. \square* 225

Being a subalgebra of $U(\mathfrak{gl}(1|1)[t])$, the Bethe algebra \mathcal{B} acts on any $\mathfrak{gl}(1|1)[t]$ -module M . Since \mathcal{B} commutes with $U(\mathfrak{h})$, the Bethe algebra preserves the subspace $(M)_\lambda$ for any weight λ . If $K \subset M$ is a \mathcal{B} -invariant subspace, then we call the image of \mathcal{B} in $\text{End}(K)$ the *Bethe algebra associated with K* . 227

Let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{C}^n$. Define $k \in \mathbb{Z}_{>0}$, a sequence of positive integers $\mathbf{n} = (n_1, \dots, n_k)$, and a sequence of distinct complex numbers $\mathbf{b} = (b_1, \dots, b_k)$ by (3.6). Let $\Lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$ be a sequence of polynomial $\mathfrak{gl}(1|1)$ -weights such that $|\lambda^{(s)}| = n_s$. 231

We study the action of the Bethe algebra \mathcal{B} on the following \mathcal{B} -modules: 232

$$\mathcal{M}_l = (\mathcal{V}^{\mathfrak{S}})_{(n-l,l)}, \quad \mathcal{M}_{l,\mathbf{a}} = \left(\bigotimes_{s=1}^k W_{n_s}(b_s) \right)_{(n-l,l)}, \quad \mathcal{M}_{l,\Lambda,\mathbf{b}} = \left(\bigotimes_{s=1}^k L_{\lambda^{(s)}}(b_s) \right)_{(n-l,l)}.$$

Denote the Bethe algebras associated with $\mathcal{M}_l, \mathcal{M}_{l,a}, \mathcal{M}_{l,\Lambda,b}$ by $\mathcal{B}_l, \mathcal{B}_{l,a}, \mathcal{B}_{l,\Lambda,b}$, respectively. For any element $X \in \mathcal{B}$, we denote by $X(z), X(a), X(\Lambda, b)$ the respective linear operators.

Since by Lemma 3.3 the $\mathfrak{gl}(1|1)[t]$ -module $\mathcal{V}^{\mathfrak{S}}$ is generated by $v_1^{\otimes n} = v_1 \otimes \cdots \otimes v_1$, the series $e_{11}(x) + e_{22}(x)$ acts on $\mathcal{V}^{\mathfrak{S}}$ by multiplication by the series

$$\sum_{i=1}^n \frac{1}{x - z_i} = \sum_{i=1}^n \sum_{j=0}^{\infty} z_i^j x^{-j-1}.$$

Therefore there exist unique central elements C_1, \dots, C_n of $U(\mathfrak{gl}(1|1)[t])$ of minimal degrees such that each C_i acts on $\mathcal{V}^{\mathfrak{S}}$ by multiplication by $\sigma_i(z)$.

Define $B_i \in \mathcal{B}$ by

$$\left(x^n + \sum_{i=1}^n (-1)^i C_i x^{n-i}\right) \mathcal{T}(x) = x^n \sum_{i=1}^{\infty} B_i x^{-i}, \quad (4.6)$$

where $\mathcal{T}(x)$ is defined in (2.13).

Lemma 4.4. *We have $B_i(z) = 0$ for $i > n$ and $B_1(z) = (q_1 - q_2)l$.*

Proof. Let $V(c) = \otimes_{i=1}^n \mathbb{C}^{1|1}(c_i)$, where $c_i \in \mathbb{C}$. Note that $B_i(z)$ is a polynomial in z with values in $\text{End}((V)_{(n-l,l)})$. For any sequence of complex numbers $c = (c_1, \dots, c_n)$, we can evaluate $B_i(z)$ at $z = c$ to an operator on $(V(c))_{(n-l,l)}$. By Theorem 2.5, the Gaudin transfer matrix $\mathcal{H}(x)$ is diagonalizable and the Bethe ansatz is complete for $(V(c))_{(n-l,l)}$ when $c \in \mathbb{C}^n$ is generic. Hence by (2.14) and (4.6) that $(x^n + \sum_{i=1}^n (-1)^i C_i x^{n-i}) \mathcal{T}(x)$ acts on $(V(c))_{(n-l,l)}$ as a polynomial in x for generic c . In particular, it implies that $B_i, i > n$, acts on $(V(c))_{(n-l,l)}$ by zero for generic c . Therefore $B_i(z), i > n$, is identically zero.

By the same reasoning, one shows that $B_1(z) = (q_1 - q_2)l$. Alternatively, it also follows from $B_1 = (q_1 - q_2)e_{22}$. \square

Lemma 4.5. *The elements $B_i(z)$ and $C_j(z)$, for $1 < i \leq n$ and $1 \leq j \leq n$, generate the algebra \mathcal{B}_l .*

Proof. It follows from the definition of \mathcal{B} , (4.6), and Lemma 4.4. \square

One can restrict the filtration on $U(\mathfrak{gl}(1|1)[t])$ to the Bethe algebra, $\mathcal{F}_0 \mathcal{B} \subset \mathcal{F}_1 \mathcal{B} \subset \cdots \subset \mathcal{B}$.

Lemma 4.6. *We have $B_i \in \mathcal{F}_{i-1} \mathcal{B} / \mathcal{F}_{i-2} \mathcal{B}$ and $C_j \in \mathcal{F}_j \mathcal{B} / \mathcal{F}_{j-1} \mathcal{B}$ for $1 < i \leq n$ and $1 \leq j \leq n$. \square*

4.3. Main theorems

Recall from Proposition 3.6 that there exists a unique vector (up to proportionality) of degree $l(l-1)/2$ in \mathcal{M}_l explicitly given by

$$u_l := e_{21}[0]e_{21}[1] \cdots e_{21}[l-1]v^+,$$

see Lemma 3.4.

Any commutative algebra \mathcal{A} is a module over itself induced by left multiplication. We call it the *regular representation* of \mathcal{A} . The dual space \mathcal{A}^* is naturally an \mathcal{A} -module which is called the *coregular representation*. A bilinear form $(\cdot | \cdot) : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathbb{C}$ is called *invariant* if $(ab|c) = (a|bc)$ for all $a, b, c \in \mathcal{A}$. A finite-dimensional commutative algebra \mathcal{A} admitting an invariant non-degenerate symmetric bilinear form $(\cdot | \cdot) : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathbb{C}$ is called a *Frobenius algebra*. The regular and coregular representations of a Frobenius algebra are isomorphic.

Let M be an \mathcal{A} -module and $\mathcal{E} : \mathcal{A} \rightarrow \mathbb{C}$ a character, then the *\mathcal{A} -eigenspace associated to \mathcal{E}* in M is defined by $\bigcap_{a \in \mathcal{A}} \ker(a|_M - \mathcal{E}(a))$. The *generalized \mathcal{A} -eigenspace associated to \mathcal{E}* in M is defined by $\bigcap_{a \in \mathcal{A}} (\bigcup_{m=1}^{\infty} \ker(a|_M - \mathcal{E}(a))^m)$.

Theorem 4.7. *The action of the Bethe algebra \mathcal{B}_1 on \mathcal{M}_1 has the following properties.* 266

1. *The map $\eta_l : G_i \mapsto B_{i+1}(z), \Sigma_j \mapsto C_j(z), i = 1, \dots, n - 1, j = 1, \dots, n$, extends uniquely to an isomorphism $\eta_l : \mathcal{O}_1 \rightarrow \mathcal{B}_1$ of filtered algebras. Moreover, the isomorphism η_l is an isomorphism of $\mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}}$ -modules.* 267
268
269
2. *The map $\rho_l : \mathcal{O}_1 \mapsto \mathcal{M}_1, F \mapsto \eta_l(F)u_l$, is an isomorphism of filtered vector spaces identifying the \mathcal{B}_1 -module \mathcal{M}_1 with the regular representation of \mathcal{O}_1 .* 270
271

Theorem 4.7 is proved in Section 5. 272

Let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{C}^n$. Define $k \in \mathbb{Z}_{>0}$, a sequence of positive integers $\mathbf{n} = (n_1, \dots, n_k)$, and a sequence of distinct complex numbers $\mathbf{b} = (b_1, \dots, b_k)$ by (3.6). Let $\Lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$ be a sequence of non-degenerate polynomial weights such that $|\lambda^{(s)}| = n_s$ for each $1 \leq s \leq k$. 273
274
275
276

Theorem 4.8. *The action of the Bethe algebra $\mathcal{B}_{l,\Lambda,\mathbf{b}}$ on $\mathcal{M}_{l,\Lambda,\mathbf{b}}$ has the following properties.* 277

1. *The Bethe algebra $\mathcal{B}_{l,\Lambda,\mathbf{b}}$ is isomorphic to*

$$\mathbb{C}[w_1, \dots, w_k]^{\mathfrak{S}_1 \times \dots \times \mathfrak{S}_{k-1}} / \langle \sigma_i(\mathbf{w}) - \varepsilon_i \rangle_{i=1, \dots, k},$$

where ε_i are given by

$$\varphi_{\Lambda,\mathbf{b}}(x) := \prod_{s=1}^k (x - b_s) \left(q_1 - q_2 + \sum_{s=1}^k \frac{n_s}{x - b_s} \right) = (q_1 - q_2) \left(x^k + \sum_{i=1}^k (-1)^i \varepsilon_i x^{k-i} \right)$$

and $\sigma_i(\mathbf{w})$ are elementary symmetric functions in w_1, \dots, w_k . 278

2. *The Bethe algebra $\mathcal{B}_{l,\Lambda,\mathbf{b}}$ is a Frobenius algebra. Moreover, the $\mathcal{B}_{l,\Lambda,\mathbf{b}}$ -module $\mathcal{M}_{l,\Lambda,\mathbf{b}}$ is isomorphic to the regular representation of $\mathcal{B}_{l,\Lambda,\mathbf{b}}$.* 279
280
3. *The Bethe algebra $\mathcal{B}_{l,\Lambda,\mathbf{b}}$ is a maximal commutative subalgebra in $\mathcal{M}_{l,\Lambda,\mathbf{b}}$ of dimension $\binom{k}{l}$.* 281
4. *Every \mathcal{B} -eigenspace in $\mathcal{M}_{l,\Lambda,\mathbf{b}}$ has dimension one.* 282
5. *The \mathcal{B} -eigenspaces in $\mathcal{M}_{l,\Lambda,\mathbf{b}}$ bijectively correspond to the monic degree l divisors $y(x)$ of the polynomial $\varphi_{\Lambda,\mathbf{b}}(x)$. Moreover, the eigenvalue of $\mathcal{H}(x)$ corresponding to the monic divisor y is described by $\mathcal{E}_{y,\Lambda,\mathbf{b}}(x)$, see (2.12).* 283
284
285
6. *Every generalized \mathcal{B} -eigenspace in $\mathcal{M}_{l,\Lambda,\mathbf{b}}$ is a cyclic \mathcal{B} -module.* 286
7. *The dimension of the generalized \mathcal{B} -eigenspace associated to $\mathcal{E}_{y,\Lambda,\mathbf{b}}(x)$ is*

$$\prod_{a \in \mathbb{C}} \binom{\text{Mult}_a(\varphi_{\Lambda,\mathbf{b}})}{\text{Mult}_a(y)},$$

where $\text{Mult}_a(p)$ is the multiplicity of a as a root of the polynomial p . 287

Theorem 4.8 is proved in Section 5. 288

Note that the results of is quite parallel to that of XXX spin chains, see [13, Theorem 4.11]. 289
290

4.4. Higher Gaudin transfer matrices 291

To define higher Gaudin transfer matrices, we first recall basics about pseudo-differential operators. Let \mathcal{A} be a differential superalgebra with an even derivation $\partial : \mathcal{A} \rightarrow \mathcal{A}$. For $r \in \mathbb{Z}_{>0}$, denote the r -th derivative of $a \in \mathcal{A}$ by $a_{[r]}$. Define the superalgebra of pseudo-differential operators $\mathcal{A}((\partial^{-1}))$ as follows. Elements of $\mathcal{A}((\partial^{-1}))$ are Laurent series in ∂^{-1} with coefficients in \mathcal{A} , and the product is given by

$$\partial \partial^{-1} = \partial^{-1} \partial = 1, \quad \partial^r a = \sum_{s=0}^{\infty} \binom{r}{s} a_{[s]} \partial^{r-s}, \quad r \in \mathbb{Z}, \quad a \in \mathcal{A},$$

where

$$\binom{r}{s} = \frac{r(r-1)\cdots(r-s+1)}{s!}.$$

Let

$$\mathcal{A}_x^{m|n} = \mathbf{U}(\mathfrak{gl}(1|1)[t])((x^{-1})) = \left\{ \sum_{r=-\infty}^s g_r x^r, r \in \mathbb{Z}, g_r \in \mathbf{U}(\mathfrak{gl}(1|1)[t]) \right\}$$

Consider the operator in $\text{End}(\mathbb{C}^{1|1}) \otimes \mathcal{A}_x^{m|n}((\partial_x^{-1}))$,

$$\mathfrak{Z}(x, \partial_x) := \sum_{a,b=1}^2 E_{ab} \otimes \left(\delta_{ab}(\partial_x - q_a) - e_{ab}(x)(-1)^{|a|} \right).$$

which is a Manin matrix, see [17, Lemma 3.1] and [5, Lemma 4.2]. Define the *Berezinian*, see [22], of $\mathfrak{Z}(x, \partial_x)$ by

$$\begin{aligned} & \text{Ber}(\mathfrak{Z}(x, \partial_x)) \\ &= (\partial_x - q_1 - e_{11}(x)) \left(\partial_x - q_2 + e_{22}(x) + e_{21}(x) (\partial_x - q_1 - e_{11}(x))^{-1} e_{12}(x) \right)^{-1}. \end{aligned} \quad (4.7)$$

Denote the Berezinian by $\mathfrak{D}(x, \partial_x)$ and expand it as an element in $\mathcal{A}_x^{m|n}((\partial_x^{-1}))$,

$$\mathfrak{D}(x, \partial_x) = \sum_{r=0}^{\infty} (-1)^r \mathcal{G}_r(x) \partial_x^{-r}. \quad (4.8)$$

We call the series $\mathcal{G}_r(x) \in \mathcal{A}_x^{m|n}$, $r \in \mathbb{Z}_{\geq 0}$, the *higher Gaudin transfer matrices*. In particular, we call $\mathcal{G}_1(x)$ and $\mathcal{G}_2(x)$ the *first and second Gaudin transfer matrices*, respectively. 292
293

Example 4.9. We have $\mathcal{G}_0(x) = 1$,

$$\mathcal{G}_1(x) = q_1 - q_2 + e_{11}(x) + e_{22}(x),$$

$$\mathcal{G}_2(x) = (q_1 - q_2 + e_{11}(x) + e_{22}(x))(-q_2 + e_{22}(x)) - e_{21}(x)e_{12}(x). \quad \square$$

Remark 4.10. In principle, the Bethe algebra should be the unital subalgebra of $\mathbf{U}(\mathfrak{gl}(1|1)[t])$ generated by coefficients $\mathcal{G}_r(x)$, $r \in \mathbb{Z}_{>0}$, cf. [16]. However, it turns out that the first two transfer matrices already give (almost) complete information about the Bethe algebra, see the discussion below. 294
295
296
297 \square

Now we describe the eigenvalues of higher Gaudin transfer matrices acting on the on-shell Bethe vector. 298
299

Let $\Lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$ be a sequence of $\mathfrak{gl}(1|1)$ -weights and $\mathbf{b} = (b_1, \dots, b_k)$ a sequence of distinct complex numbers, where $\lambda^{(s)} = (\alpha_s, \beta_s)$. Let $\mathbf{t} = (t_1, \dots, t_l)$, where $0 \leq l < k$. Suppose that $y_{\mathbf{t}}$ divides the polynomial $\varphi_{\Lambda, \mathbf{b}}$ (namely \mathbf{t} satisfies the Bethe ansatz equation), see (2.10). 300
301
302
303

Theorem 4.11 ([12, Theorem 5.2]). *If $t_i \neq t_j$ for $1 \leq i < j \leq l$, then*

$$\mathfrak{D}(x, \partial_x) \mathbb{B}_l(\mathbf{t}) = \mathbb{B}_l(\mathbf{t}) \left(\partial_x - q_1 - \sum_{s=1}^k \frac{\alpha_s}{x - b_s} + \frac{y'_{\mathbf{t}}}{y_{\mathbf{t}}} \right) \left(\partial_x - q_2 + \sum_{s=1}^k \frac{\beta_s}{x - b_s} + \frac{y'_{\mathbf{t}}}{y_{\mathbf{t}}} \right)^{-1}. \quad (4.9)$$

The theorem is a differential analog of [13, Theorem 6.4]. Note that the pseudo-differential operator in the right hand side of (4.9), denoted by $\mathfrak{D}_{y, \Lambda, \mathbf{b}}$, was introduced [6, Section 5.3]. This theorem is generalized to the $\mathfrak{gl}(m|n)$ case in [12, Theorem 5.2] where on the right hand side the pseudo-differential operator describing the eigenvalues of higher Gaudin transfer matrices should be replaced by the pseudo-differential operator in [6, 304
305
306
307
308

Equation (6.5)]. This generalization is a classical limit of [14, Conjecture 5.15] and [12, Corollary 3.6] which connects the rational difference operator introduced in [4, Equation (5.6)] with the eigenvalues of higher transfer matrices on the on-shell Bethe vector for XXX spin chains associated with $\mathfrak{gl}(m|n)$.

Remark 4.12. As shown in [6, Lemma 5.7], the odd reflection of $\mathfrak{D}_{y,\Lambda,b}$, cf. [6, equation (3.1)], which comes from the study of the fermionic reproduction procedure of the Bethe ansatz equation, are compatible with the odd reflection of Lie superalgebras. The difference analog of this fact has been used in [10] to investigate the relations between the odd reflections of super Yangian of type A and the fermionic reproduction procedure of the Bethe ansatz equation for XXX spin chains. \square

We conclude this section by discussing the connections between $\mathcal{G}_i(x)$, $i \geq 3$, and $\mathcal{G}_1(x)$, $\mathcal{G}_2(x)$.

Let

$$\mu(x) = q_1 + \sum_{s=1}^k \frac{\alpha_s}{x - b_s} - \frac{y'_t}{y_t}, \quad \nu(x) = -q_2 + \sum_{s=1}^k \frac{\beta_s}{x - b_s} + \frac{y'_t}{y_t}.$$

For simplicity, we do not write the dependence of $\mu(x)$ and $\nu(x)$ on $\Lambda, \mathbf{b}, \mathbf{t}$ explicitly. Then the eigenvalue of $\mathfrak{D}(x, \partial_x)$ acting on $\mathbb{B}_l(\mathbf{t})$ is given by

$$(\partial_x - \mu(x))(\partial_x + \nu(x))^{-1} = 1 - (\mu(x) + \nu(x))(\partial_x + \nu(x))^{-1}. \quad (4.10)$$

Hence the eigenvalues of $\mathcal{G}_i(x)$ are essentially determined only by $\mu(x) + \nu(x)$ and $\nu(x)$. Comparing (4.8) and the expansion of (4.10), we have

$$\mathcal{G}_1(x)\mathbb{B}_l(\mathbf{t}) = (\mu(x) + \nu(x))\mathbb{B}_l(\mathbf{t}), \quad \mathcal{G}_2(x)\mathbb{B}_l(\mathbf{t}) = (\mu(x) + \nu(x))\nu(x)\mathbb{B}_l(\mathbf{t}), \quad (4.11)$$

see also (2.14). Therefore, the spectrum of all higher transfer matrices are determined simply by that of the first two transfer matrices which justifies our definition of Bethe algebra.

Lemma 4.13. *Let the complex parameters c_1, \dots, c_m and the positive integer m vary. Then the kernels of the representations $\otimes_{i=1}^m \mathbb{C}^{1|1}(c_i)$ of $U(\mathfrak{gl}(1|1)[t])$ have the zero intersection.*

Proof. The proof is contained in the proof of [23, Proposition 1.7]. \square

Corollary 4.14. *We have*

$$\mathfrak{D}(x, \partial_x) = \left(\partial_x - \mathcal{G}_1(x) + \frac{\mathcal{G}_2(x)}{\mathcal{G}_1(x)} \right) \left(\partial_x + \frac{\mathcal{G}_2(x)}{\mathcal{G}_1(x)} \right)^{-1}. \quad (4.12)$$

Proof. By Lemma 4.13, it suffices to check that the left hand side and the right hand side of (4.12) act identically on a basis of $\otimes_{i=1}^m \mathbb{C}^{1|1}(c_i)$ for all $m \in \mathbb{Z}_{>0}$ and generic $c = (c_1, \dots, c_m)$.

By Theorem 2.5, there is a basis of $\otimes_{i=1}^m \mathbb{C}^{1|1}(c_i)$ consisting of on-shell Bethe vectors for generic c . Therefore, the statement follows from Theorem 4.11, (4.10), (4.11). \square

5. Proof of main theorems

In this section, we prove the main theorems. For completeness, we provide all details even they are parallel to those in [11, Section 5].

5.1. The first isomorphism

Proof of Theorem 4.7. We first show the homomorphism defined by η_l is well-defined.

Consider the tensor product $V(c) = \otimes_{i=1}^n \mathbb{C}^{1|1}(c_i)$, where $c_i \in \mathbb{C}$, and the corresponding Bethe ansatz equation associated to weight $(n - l, l)$. Let \mathbf{t} be a solution with distinct coordinates and $\mathbb{B}_l(\mathbf{t})$ the corresponding on-shell Bethe vector. Denote $\mathcal{E}_{i,t}$ the eigenvalues of B_i acting on $\mathbb{B}_l(\mathbf{t})$, see Theorem 2.2 and (2.14).

Define a character $\pi : \mathcal{O}_l \rightarrow \mathbb{C}$ by sending

$$f(x) \mapsto y_t(x), \quad \Sigma_n \mapsto \prod_{i=1}^n c_i,$$

$$g(x) \mapsto \frac{1}{(q_1 - q_2)y_t(x)} \prod_{i=1}^n (x - c_i) \left(q_1 - q_2 + \sum_{i=1}^n \frac{1}{x - c_i} \right).$$

Then

$$\pi(\Sigma_i) = \sigma_i(c), \quad \pi(G_i) = \mathcal{E}_{i,t}, \quad (5.1)$$

by (4.3) and by (2.12), (2.14), (4.5), respectively. 340

Let now $P(G_i, \Sigma_j)$ be a polynomial in G_i, Σ_j such that $P(G_i, \Sigma_j)$ is equal to zero in \mathcal{O}_l . 341
It suffices to show $P(B_i(z), C_j(z))$ is equal to zero in \mathcal{B}_l . 342

Note that $P(B_i(z), C_j(z))$ is a polynomial in z_1, \dots, z_n with values in $\text{End}((V)_{(n-l,l)})$. 343
For any sequence c of complex numbers, we can evaluate $P(B_i(z), C_j(z))$ at $z = c$ to an 344
operator on $(V(c))_{(n-l,l)}$. By Theorem 2.5, the transfer matrix $\mathcal{T}(x)$ is diagonalizable and 345
the Bethe ansatz is complete for $(V(c))_{(n-l,l)}$ when $c \in \mathbb{C}^n$ is generic. Hence by (5.1) the 346
value of $P(B_i(z), C_j(z))$ at $z = c$ is also equal to zero for generic c . Therefore $P(B_i(z), C_j(z))$ 347
is identically zero and the map η_l is well-defined. 348

Let us now show that the map η_l is injective. Let $P(G_i, \Sigma_j)$ be a polynomial in G_i, Σ_j 349
such that $P(G_i, \Sigma_j)$ is non-zero in \mathcal{O}_l . Then the value at a generic point of Ω_l (e.g. the 350
non-vanishing points of $P(G_i, \Sigma_j)$ such that f and g are relatively prime and have only 351
simple zeros) is not equal to zero. Moreover, at those points the transfer matrix $\mathcal{T}(x)$ is 352
diagonalizable and the Bethe ansatz is complete again by Theorem 2.5. Therefore, again 353
by (5.1), the polynomial $P(B_i(z), C_j(z))$ is a non-zero element in \mathcal{B}_l . Thus the map η_l is 354
injective. 355

The surjectivity of η_l follows from Lemma 4.5. Hence η_l is an isomorphism of algebras. 356

The fact that η_l is an isomorphism of graded algebra respecting the gradation follows 357
from Lemmas 4.2 and 4.6. This completes the proof of part (i). 358

The kernel of ρ_l is an ideal of \mathcal{O}_l . If we identify $\sigma_i(z)$ with Σ_i , then the algebra 359
 \mathcal{O}_l contains the algebra $\mathbb{C}[z_1, \dots, z_n]^{\otimes}$, see (4.4). The kernel of ρ_l intersects $\mathbb{C}[z_1, \dots, z_n]^{\otimes}$ 360
trivially. Therefore the kernel of ρ_l is trivial as well. Hence ρ_l is an injective map. Comparing 361
(4.2) and Proposition 3.6, we have $\text{ch}(\mathcal{M}_l) = q^{l(l-1)/2} \text{ch}(\mathcal{O}_l)$. Thus ρ_l is an isomorphism 362
of graded vector spaces which shifts the degree by $l(l-1)/2$, completing the proof of part 363
(ii). \square 364

5.2. The second isomorphism 365

Let $\mathbf{a} = (a_1, \dots, a_n)$ be a sequence of complex numbers. Define $k \in \mathbb{Z}_{>0}$, a sequence 366
of positive integers $\mathbf{n} = (n_1, \dots, n_k)$, and a sequence of distinct complex numbers $\mathbf{b} =$
 (b_1, \dots, b_k) by (3.6). Let $I_{l,\mathbf{a}}^{\mathcal{O}}$ be the ideal of \mathcal{O}_l generated by the elements $\Sigma_i - a_i$, $i = 1, \dots, n$,
where $\Sigma_1, \dots, \Sigma_{n-1}$ are defined in (4.3). Let $\mathcal{O}_{l,\mathbf{a}}$ be the quotient algebra

$$\mathcal{O}_{l,\mathbf{a}} = \mathcal{O}_l / I_{l,\mathbf{a}}^{\mathcal{O}}.$$

Let $I_{l,\mathbf{a}}^{\mathcal{B}}$ be the ideal of \mathcal{B}_l generated by $C_i(z) - a_i$, $i = 1, \dots, n$. Consider the subspace

$$I_{l,\mathbf{a}}^{\mathcal{M}} = I_{l,\mathbf{a}}^{\mathcal{B}} \mathcal{M}_l = (I_{\mathbf{a}} \mathcal{V}^{\otimes})_{(n-l,l)},$$

where $I_{\mathbf{a}}$ as before is the ideal of $\mathbb{C}[z_1, \dots, z_n]^{\otimes}$ generated by $\sigma_i(z) - a_i$, $i = 1, \dots, n$. 366

Lemma 5.1. *We have*

$$\eta_l(I_{l,\mathbf{a}}^{\mathcal{O}}) = I_{l,\mathbf{a}}^{\mathcal{B}}, \quad \rho_l(I_{l,\mathbf{a}}^{\mathcal{O}}) = I_{l,\mathbf{a}}^{\mathcal{M}}, \quad \mathcal{B}_{l,\mathbf{a}} = \mathcal{B}_l / I_{l,\mathbf{a}}^{\mathcal{B}}, \quad \mathcal{M}_{l,\mathbf{a}} = (\mathcal{V}^{\otimes})_{(n-l,l)}^{\text{sing}} / I_{l,\mathbf{a}}^{\mathcal{M}}.$$

Proof. The lemma follows from Theorem 4.7 and Lemma 3.9. \square 367

By Lemma 5.1, the maps η_l and ρ_l induce the maps

$$\eta_{l,a} : \mathcal{O}_{l,a} \rightarrow \mathcal{B}_{l,a}, \quad \rho_{l,a} : \mathcal{O}_{l,a} \rightarrow \mathcal{M}_{l,a}.$$

The map $\eta_{l,a}$ is an isomorphism of algebras. Since $\mathcal{B}_{l,a}$ is finite-dimensional, by e.g. [20, Lemma 3.9], $\mathcal{O}_{l,a}$ is a Frobenius algebra, so is $\mathcal{B}_{l,a}$. The map $\rho_{l,a}$ is an isomorphism of vector spaces. Moreover, it follows from Theorem 4.7 and Lemma 5.1 that $\rho_{l,a}$ identifies the regular representation of $\mathcal{O}_{l,a}$ with the $\mathcal{B}_{l,a}$ -module $\mathcal{M}_{l,a}$. 368
369
370
371

The statement of this section implies, by e.g. [9, Lemma 1.3], the following. Set

$$\zeta_{n,b}(x) = q_1 - q_2 + \sum_{s=1}^k \frac{n_s}{x - b_s}, \quad \psi_{n,b}(x) := \zeta_{n,b}(x) \prod_{r=1}^k (x - b_r)^{n_s}.$$

Theorem 5.2. Suppose $\mathbf{b} = (b_1, \dots, b_k)$ is a sequence of distinct complex numbers. Then the Gaudin transfer matrix $\mathcal{H}(x)$ has a simple spectrum in $(\otimes_{s=1}^k W_{n_s}(b_s))^{\text{sing}}$. There exists a bijective correspondence between the monic divisors y of the polynomial $\psi_{n,b}$ and the eigenvectors v_y of the Gaudin transfer matrix $\mathcal{H}(x)$ (up to multiplication by a non-zero constant). Moreover, this bijection is such that

$$\mathcal{H}(x)v_y = \left(\frac{1}{2}\zeta'_{n,b}(x) - \zeta_{n,b}(x) \frac{y'(x)}{y(x)} + \frac{1}{2} \left(\sum_{s=1}^k \frac{n_s}{x - b_s} \right) \left(\sum_{s=1}^k \frac{n_s}{x - b_s} + 2q_1 \right) \right) v_y. \quad \square$$

Remark 5.3. Fix $l \in \mathbb{Z}_{\geq 0}$ and set $\mathbf{t} = (t_1, \dots, t_l)$. Let \mathbf{y}_t represent \mathbf{t} . Then the Bethe ansatz equation for $(\otimes_{s=1}^k W_{n_s}(b_s))$ is

$$y_t(x) \quad \text{divides the polynomial} \quad \psi_{n,b}(x).$$

Note that in this case, y_t may have multiple roots. If there are multiple roots in y_t , then the corresponding on-shell Bethe vector is zero. Therefore an actual eigenvector should be obtained via an appropriate derivative as pointed out in [6, Section 8.2]. \square 372
373
374

5.3. The third isomorphism 375

Recall from Section 2.3, that without loss of generality, we can assume that $\beta_s = 0$, $1 \leq s \leq k$. In this case, $\alpha_s = n_s$, $1 \leq s \leq k$. 376
377

Lemma 5.4. There exists a surjective $\mathfrak{gl}(1|1)[t]$ -module homomorphism from $\otimes_{s=1}^k W_{n_s}(b_k)$ to $\otimes_{s=1}^k L_{\lambda^{(s)}}(b_k)$ which maps vacuum vector to vacuum vector. 378
379

Proof. It follows from Lemma 3.10 and our assumption that $\beta_s = 0$ for all $1 \leq s \leq k$. \square 380

By Lemma 3.9, the surjective $\mathfrak{gl}(1|1)[t]$ -module homomorphism

$$\otimes_{s=1}^k W_{n_s}(b_k) \rightarrow \otimes_{s=1}^k L_{\lambda^{(s)}}(b_k)$$

induces a surjective $\mathfrak{gl}(1|1)[t]$ -module homomorphism

$$\mathcal{V}^{\mathfrak{S}} \rightarrow \otimes_{s=1}^k L_{\lambda^{(s)}}(b_k).$$

The second map then induces a projection of the Bethe algebras $\mathcal{B}_l \rightarrow \mathcal{B}_{l,\Lambda,b}$. We describe the kernel of this projection. We consider the corresponding ideal in the algebra \mathcal{O}_l . 381
382

Suppose $l \leq k$. Define the polynomial $h(x)$ by

$$h(x) = \prod_{s=1}^k (x - b_s)^{n_s - 1}.$$

Divide the polynomial $g(x)$ in (4.1) by $h(x)$ and let

$$p(x) = x^{k-l} + p_1 x^{k-l-1} + \cdots + p_{k-l-1} x + p_{k-l}, \quad (5.2)$$

$$r(x) = r_1 x^{n-k-1} + r_2 x^{n-k-2} + \cdots + r_{n-k-1} x + r_{n-k} \quad (5.3)$$

be the quotient and the remainder, respectively. Clearly, $p_i, r_j \in \mathcal{O}_l$. 383

Denote by $I_{l,\Lambda,b}^{\mathcal{O}}$ the ideal of \mathcal{O}_l generated by r_1, \dots, r_{n-k} , and the coefficients of polynomial

$$\varphi_{\Lambda,b}(x) - (q_1 - q_2)p(x)f(x) = \prod_{s=1}^k (x - b_s) \left(q_1 - q_2 + \sum_{s=1}^k \frac{n_s}{x - b_s} \right) - (q_1 - q_2)p(x)f(x).$$

Let $\mathcal{O}_{l,\Lambda,b}$ be the quotient algebra

$$\mathcal{O}_{l,\Lambda,b} = \mathcal{O}_l / I_{l,\Lambda,b}^{\mathcal{O}}.$$

Clearly, if $\mathcal{O}_{l,\Lambda,b}$ is finite-dimensional, then it is a Frobenius algebra. 384

Let $I_{l,\Lambda,b}^{\mathcal{B}}$ be the image of $I_{l,\Lambda,b}^{\mathcal{O}}$ under the isomorphism η_l . 385

Lemma 5.5. *The ideal $I_{l,\Lambda,b}^{\mathcal{B}}$ is contained in the kernel of the projection $\mathcal{B}_l \rightarrow \mathcal{B}_{l,\Lambda,b}$.* 386

Proof. We treat $\mathbf{b} = (b_1, \dots, b_k)$ as variables. Note that the elements of $I_{l,\Lambda,b}^{\mathcal{B}}$ act on $\mathcal{M}_{l,\Lambda,b}$ as polynomials in \mathbf{b} with values in $\text{End}((L_{\Lambda})_{(n-l)})$. Therefore it suffices to show it for generic \mathbf{b} . Let $\mathfrak{f}(x)$ be the image of $f(x)$ under η_l . The condition that $I_{l,\Lambda,b}^{\mathcal{B}}$ vanishes is equivalent to the condition that $\varphi_{\Lambda,b}(x)$ is divisible by $\mathfrak{f}(x)$. 387
388
389
390

By Theorem 2.5, there exists an eigenbasis of the operator $\mathcal{T}(x)$ in $\mathcal{M}_{l,\Lambda,b}$ for generic \mathbf{b} . Clearly, a solution of Bethe ansatz equation associated to Λ, \mathbf{b}, l is also a solution to Bethe ansatz equation for $\mathcal{M}_{l,\Lambda}$, see Theorem 5.2 and Remark 5.3. Moreover, the expressions of corresponding on-shell Bethe vectors coincide (with different vacuum vectors). By Lemma 5.4 and Theorems 2.2, 5.2, $\varphi_{\Lambda,b}(x)$ is divisible by $\mathfrak{f}(x)$ for generic \mathbf{b} since the eigenvalue of $\mathfrak{f}(x)$ corresponds to $y_t(x)$ in (2.12). Therefore $I_{l,\Lambda,b}^{\mathcal{B}}$ vanishes for generic \mathbf{b} , thus completing the proof. \square 391
392
393
394
395
396
397

Therefore, we have the epimorphism

$$\mathcal{O}_{l,\Lambda,b} \cong \mathcal{B}_l / I_{l,\Lambda,b}^{\mathcal{B}} \twoheadrightarrow \mathcal{B}_{l,\Lambda,b}. \quad (5.4)$$

We claim that the surjection in (5.4) is an isomorphism by checking $\dim \mathcal{O}_{l,\Lambda,b} = \dim \mathcal{B}_{l,\Lambda,b}$. 398

Lemma 5.6. *We have $\dim \mathcal{O}_{l,\Lambda,b} = \binom{k}{l}$.* 399

Proof. Note that $\mathbb{C}[p_1, \dots, p_{k-l}, r_1, \dots, r_{n-k}] \cong \mathbb{C}[g_1, \dots, g_{n-l}]$, where p_i and r_j are defined in (5.2) and (5.3). It is not hard to check that

$$\mathcal{O}_{l,\Lambda,b} \cong \mathbb{C}[f_1, \dots, f_l, p_1, \dots, p_{k-l}] / \tilde{I}_{l,\Lambda,b}^{\mathcal{O}}, \quad (5.5)$$

where $\tilde{I}_{l,\Lambda,b}^{\mathcal{O}}$ is the ideal of $\mathbb{C}[f_1, \dots, f_l, p_1, \dots, p_{k-l}]$ generated by the coefficients of the polynomial $\varphi_{\Lambda,b}(x) - (q_1 - q_2)p(x)f(x)$. 400
401

Introduce new variables $w = (w_1, \dots, w_k)$ such that

$$f(x) = \prod_{i=1}^l (x - w_i), \quad p(x) = \prod_{i=1}^{k-l} (x - w_{l+i}).$$

Let $\varepsilon = (\varepsilon_1, \dots, \varepsilon_k)$ be complex numbers such that

$$\varphi_{\Lambda, b}(x) = \prod_{s=1}^k (x - b_s) \left(q_1 - q_2 + \sum_{s=1}^k \frac{n_s}{x - b_s} \right) = (q_1 - q_2) \left(x^k + \sum_{i=1}^k (-1)^i \varepsilon_i x^{k-i} \right).$$

Then

$$\mathbb{C}[f_1, \dots, f_l, p_1, \dots, p_{k-l}] / \tilde{\mathcal{I}}_{l, \Lambda, b}^{\mathcal{O}} \cong \mathbb{C}[w_1, \dots, w_{k-1}]^{\mathfrak{S}_l \times \mathfrak{S}_{k-l}} / \langle \sigma_i(w) - \varepsilon_i \rangle_{i=1, \dots, k}. \quad (5.6)$$

The lemma now follows from the fact that $\mathbb{C}[w_1, \dots, w_{k-1}]^{\mathfrak{S}_l \times \mathfrak{S}_{k-l}}$ is a free $\mathbb{C}[w_1, \dots, w_{k-1}]^{\mathfrak{S}_l}$ -module of rank $\binom{k}{l}$. \square

Note that we have the projection $(\mathcal{V}^{\mathfrak{S}})_{(n-l, l)} \rightarrow \mathcal{M}_{l, \Lambda, b}$. Since by Theorem 4.7 the Bethe algebra \mathcal{B}_l acts on $(\mathcal{V}^{\mathfrak{S}})_{(n-l, l)}$ cyclically, the Bethe algebra $\mathcal{B}_{l, \Lambda, b}$ acts on $\mathcal{M}_{l, \Lambda, b}$ cyclically as well. Therefore we have

$$\dim \mathcal{B}_{l, \Lambda, b} = \dim \mathcal{M}_{l, \Lambda, b} = \binom{k}{l}. \quad (5.7)$$

Proof of Theorem 4.8. Part (i) follows from Lemma 5.6 and (5.4), (5.5), (5.6), (5.7). Clearly, we have $\mathcal{B}_{l, \Lambda, b} \cong \mathcal{O}_{l, \Lambda, b}$ is a Frobenius algebra. Moreover, the map ρ_l from Theorem 4.7 induces a map

$$\rho_{l, \Lambda, b} : \mathcal{O}_{l, \Lambda, b} \rightarrow \mathcal{M}_{l, \Lambda, b}$$

which identifies the regular representation of $\mathcal{O}_{l, \Lambda, b}$ with the $\mathcal{B}_{l, \Lambda, b}$ -module $\mathcal{M}_{l, \Lambda, b}$. Therefore part (ii) is proved.

Since $\mathcal{B}_{l, \Lambda, b}$ is a Frobenius algebra, the regular and coregular representations of the algebra $\mathcal{B}_{l, \Lambda, b}$ are isomorphic to each other. Parts (iii)–(vi) follow from the general facts about the coregular representations, see e.g. [20, Section 3.3] or [9, Lemma 1.3].

Due to part (iv), it suffices to consider the algebraic multiplicity of every eigenvalue. It is well known that roots of a polynomial depend continuously on its coefficients. Hence the eigenvalues of $\mathcal{T}(x)$ depend continuously on b . Part (vii) follows from the deformation argument and Theorem 2.5. \square

6. Conclusions

In this paper, we investigated the $\mathfrak{gl}(1|1)$ Gaudin models that are twisted by a diagonal matrix G and defined on tensor products of polynomial evaluation $\mathfrak{gl}(1|1)[t]$ -modules. Our results generalize all the results of [11] to the twisted case. In the meanwhile, we gave an explicit description of the algebra of Hamiltonians acting on tensor products of polynomial evaluation $\mathfrak{gl}(1|1)[t]$ -modules by generators and relations. Moreover, we showed that there exists a bijection between common eigenvectors (up to proportionality) of the algebra of Hamiltonians and monic divisors of an explicit polynomial written in terms of the highest weights and evaluation parameters. In particular, our result implies that each common eigenspace of the algebra of Hamiltonians has dimension one. We also gave dimensions of the generalized eigenspaces. Our results give a confirmed answer to the completeness of Bethe ansatz in the case of $\mathfrak{gl}(1|1)$ Gaudin models. We expect our results will be an essential step towards understanding the more general $\mathfrak{gl}(m|n)$ Gaudin models.

References

1. S. Belliard, E. Ragoucy, *The nested Bethe ansatz for 'all' closed spin chains*. J. Phys A: Math. and Theor. **41** (2008), 295202.
2. B. Feigin, E. Frenkel, N. Reshetikhin, *Gaudin model, Bethe ansatz and critical level*. Comm. Math. Phys., **166** (1994), no. 1, 27–62.

3. I. Halacheva, J. Kamnitzer, L. Rybnikov, A. Weekes, *Crystals and monodromy of Bethe vectors*. *Duke Math. J.* **169** (2020), no. 12, 2337–2419. 429
430
4. C.-L. Huang, K. Lu, E. Mukhin, *Solutions of $\mathfrak{gl}(m|n)$ XXX Bethe ansatz equation and rational difference operators*. *J. Phys. A* **52** (2019), 375204, 31pp. 431
432
5. C.-L. Huang, E. Mukhin, *The duality of $\mathfrak{gl}_{m|n}$ and \mathfrak{gl}_k Gaudin models*. *J. Algebra* **548** (2020), 1–24. 433
6. C.-L. Huang, E. Mukhin, B. Vicedo, C. Young, *The solutions of $\mathfrak{gl}(m|n)$ Bethe ansatz equation and rational pseudodifferential operators*. *Sel. Math. New Ser.* (2019) **25**:52. 434
435
7. P. Kulish, N. Manojlovic, *Bethe vectors of the $\mathfrak{osp}(1|2)$ Gaudin model*. *Lett. Math. Phys.* **55** (2001), no. 1, 77–95. 436
8. P. Kulish, N. Yu. Reshetikhin, *Diagonalization of $GL(N)$ invariant transfer-matrices and quantum N -wave system (Lee model)*. *J. Phys A: Math. Theor.* **15** (1983), L591–L596. 437
438
9. K. Lu, *Perfect integrability and Gaudin models*. *SIGMA* **16** (2020), 132, 10 pages. 439
10. K. Lu, *A note on odd reflections of super Yangian and Bethe ansatz*. *Lett. Math. Phys.* **112** (2022), Article no.: 29. 440
11. K. Lu, *Completeness of Bethe ansatz for Gaudin models associated with $\mathfrak{gl}(1|1)$* . *Nuclear Phys. B* **980** (2022), Paper No. 115790, 23 pp. 441
12. K. Lu, *On Bethe eigenvectors and higher transfer matrices for supersymmetric spin chains*. preprint, [arXiv:2209.14416](https://arxiv.org/abs/2209.14416). 442
13. K. Lu, E. Mukhin, *On the supersymmetric XXX spin chains associated with $\mathfrak{gl}_{1|1}$* . *Commun. Math. Phys.* **386** (2021), 711–747. 443
14. K. Lu, E. Mukhin, *Jacobi-Trudi identity and Drinfeld functor for super Yangian*. *Int. Math. Res. Not. IMRN*, **2021** (2021), no. 21, 16749–16808. 444
445
15. K. Lu, E. Mukhin, *Bethe ansatz equations for orthosymplectic Lie superalgebra and self-dual superspaces*. *Ann. Henri Poincaré* **22** (2021), 4087–4130. 446
447
16. A. Molev, E. Mukhin, *Invariants of the vacuum module associated with the Lie superalgebra $\mathfrak{gl}(1|1)$* . *J. Phys A: Math. Theor.* **48** (2015), no. 31, 314001, 20 pp. 448
449
17. A. Molev, E. Ragoucy, *The MacMahon Master Theorem for right quantum superalgebras and higher Sugawara operators for $\widehat{\mathfrak{gl}}(m|n)$* . *Moscow Math. J.*, **14** (2014), no. 1, 83–119. 450
451
18. E. Mukhin, V. Tarasov, A. Varchenko, *Bethe eigenvectors of higher transfer matrices*. *J. Stat. Mech. Theor. Exp.* (2006) P08002. 452
19. E. Mukhin, V. Tarasov, A. Varchenko, *Spaces of quasi-exponentials and representations of \mathfrak{gl}_N* . *J. Phys A: Math. Theor.* **41** (2008), no. 19, 194017, 28 pp. 453
454
20. E. Mukhin, V. Tarasov, A. Varchenko, *Schubert calculus and representations of general linear group*. *J. Amer. Math. Soc.* **22** (2009), no. 4, 909–940. 455
456
21. E. Mukhin, B. Vicedo, C. Young, *Gaudin models for $\mathfrak{gl}(m|n)$* . *J. Math. Phys.* **56** (2015), no. 5, 051704, 30 pp. 457
22. M. Nazarov, *Quantum Berezinian and the classical Capelli identity*. *Lett. Math. Phys.* **21** (1991), 123–131. 458
23. M. Nazarov, *Yangian of the General Linear Lie Superalgebra*. *SIGMA Symmetry Integrability Geom. Methods Appl.* **16** (2020), Paper No. 112, 24 pages. 459
460
24. A. Zeitlin, *Superopers on Supercurves*. *Lett. Math. Phys.* **105** (2015), 149–167. 461