

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

Influencing Factors for the Implementation of School-based Interventions Promoting Obesity Prevention Behaviors in Children with Low Socioeconomic Status: A Systematic Review

Friederike Butscher^{1*}, Jan Ellinger¹, Monika Singer¹ and Christoph Mall¹

¹ Department of Sport and Health Sciences, Associate Professorship of Didactics in Sport and Health, Technical University of Munich, Georg-Brauchle-Ring 60/62, 80992, Munich, Germany

*Corresponding author: friedericke.butscher@tum.de

Abstract: Background: Health inequity (HI) remains a major challenge in public health. Improving the health of children with low socioeconomic status (SES) can help to reduce overall HI in children. Childhood obesity is a global problem, entailing several adverse health effects. It is crucial to assess the influencing factors for adoption, implementation and sustainment of interventions. This review aims to identify articles reporting about influencing factors for the implementation of school-based interventions promoting obesity prevention behaviors in children with low SES. It aims to critically appraise the articles' quality, assess influencing factors, categorize and evaluate them, and to discuss possible implications. Methods: A systematic search was conducted in 7 databases with the following main inclusion criteria: 1) school-based interventions and 2) target group aged 5–14 years. The Consolidated Framework for Implementation Research, its five domains (intervention characteristics, inner setting, outer setting, characteristics of individuals, process) along with 39 constructs within these domains were used to deductively analyze the data. We grouped the articles with regard to the characteristics of the interventions in *simple* and *complex* interventions. For each domain, and for the groups of *simple* and *complex* interventions, the most commonly reported influencing factors are identified. Results: 6452 articles were screened, and 16 met all eligibility criteria. Included articles applied mixed methods (n=10), qualitative (n=5) and quantitative design (n=1). Of these, five were considered to report *simple* interventions and eleven were considered to report *complex* interventions. In total, 295 influencing factors were assessed. Aspects of the *inner setting* were reported in every study, aspects of the *outer setting* were the least reported domain, and in the group of simple interventions not reported at all. In the *inner setting*, most reported influencing factors were *time* (n=7), *scheduling* (n=6) and *communication* (n=6). Conclusion: This review found a wide range of influencing factors for implementation. Most important influencing factors need to be assessed for every setting. Including all stakeholders involved in the implementation process enhances the prioritization of the most important influencing factors for the specific setting. More empirical research and practical guidance are needed to promote obesity prevention behaviors among children with low SES.

Keywords: health equity; implementation; obesity; systematic review; CFIR; qualitative review; school-based interventions; children; facilitators

1. Background

“Implementation Science could, quite literally, put health equity back on the fast track.” Beryne

Odeny (1)

Health inequalities exist within and between populations (2). Life expectancy differs between countries as well as within them, such as, for example, between men and women (2). The social determinants of health (SDH) are a powerful driver for health inequalities

(3-5). Under certain conditions, we no longer speak of health inequalities but of health inequity (HI) (6, 7): If health inequalities arise due to the unequal distribution of the SDH, such as income, wealth, and access to health care (5, 8), and if health inequalities are therefore avoidable and unfair (9). HI is social injustice in health (7).

On an individual level, SDH can be represented by socioeconomic status (SES) (10). SES is a multidimensional concept and incorporates several socioeconomic factors. It can be described by past or current income, family wealth, educational level, occupation and social standing within the community (11). HI follows a social gradient, as groups with a low SES have poorer health (e.g., higher mortality and morbidity) than groups with high SES (5, 12).

It is especially important to protect health of children, as they have less control over their health and the circumstances influencing their health than to adults (13). Negative health influences in childhood can lead to health consequences throughout life (14, 15). Being overweight in childhood, for example, is associated with also being overweight as an adult (16), and diverse adverse health effects such as, cardiovascular diseases or mental disorders, can result from overweight and obesity in childhood (17, 18). The prevalence of obesity in children and youths is increasing globally (19, 20) and therefore, it is important to develop and implement interventions addressing childhood obesity.

Obesity in childhood lead to adverse health effects in adulthood, and in industrialized countries, childhood obesity exhibits HI: low SES is associated with higher rates of obesity among children (21).

Furthermore, due to societal processes, low SES and poor health implicate and maintain each other (7, 10, 22, 23). In their model of child health inequalities, Pearce et al. (13) show that low SES and low child health status are in a mutually reinforcing cycle, conditioning and maintaining each other. To break the cycle health-promoting interventions that improve the health status and therefore help to reduce HI in children may be a suitable entry point (13).

Interventions to prevent obesity among children generally take place in schools (24), and moderate evidence has been found for school-based combined diet and physical activity (PA) interventions (25-27). School-based interventions are a proposed approach for preventing obesity, because in the school setting almost all children in society can be reached (28). Furthermore, it is important to implement those interventions in real-world settings, as the implementation of an intervention influences its effectiveness (29, 30). Improving the reach and the adoption, delivery, and sustainment of effective interventions is the aim of implementation science (31). Because several factors influencing the speed and extent of the adoption, uptake, and use of an intervention (32), a suggested first step in the implementation process is the identification of those influencing factors in order to address them (33).

The influencing factors for the implementation of interventions have been assessed in the school setting, both for PA promoting interventions (30) and for interventions to promote PA and reducing sedentary behavior (34). Barriers and facilitators were assessed for the sustainment of health behavior interventions in schools and childcare settings (35), for PA during school lessons (36), and for the provision of fruit and vegetable in kindergartens and schools (37).

Those reviews present important results, but none of those reviews distinguished between different SES, although this factor is an important differentiator every study should take into account to approach health equity (38). Furthermore, none of the existing reviews assessed the implementation of interventions addressing the combination of the two leading behaviors in obesity development, namely, PA and nutrition (39), in the school setting. From these considerations, it is essential that factors influencing the implementation of school-based interventions be systematically assessed to promote obesity prevention behaviors for children with low SES. These findings can help improving the understanding of specific needs, to guide practice, to improve implementation, and therefore, to enhance the sustainment of effective interventions. This can further contribute to reducing HI in children. This reviews aimed to identify articles reporting about

influencing factors for the implementation of school-based interventions promoting obesity prevention behaviors for children with low SES. Furthermore, to assess the methodological quality of the identified articles, to categorize and evaluate reported influencing factors, to analyze differences of reported influencing factors regarding *simple* and *complex* interventions, and to discuss possible implications.

2. Methods

To identify, critically appraise, and summarize evidence from the articles conducted, we conducted a systematic review in accordance with established guidelines for such systematic reviews (40), the PRISMA Checklist can be found in the Additional file 1. This review was previously registered at PROSPERO (ID: CRD42021281209).

Information sources and searches

The databases Scopus, PubMed, ERIC, SportDiscus, PsychArticles, Education Source and SocINDEX were searched for relevant articles. The terms shown in Table 1 were used to construct the search term, following database specifications (see Additional file 2). There were no limitations with respect to the publication date of the articles, as no systematic review with the same aim had previously been conducted. The database search was completed on July 2, 2021.

Table 1. Search term.

	AND →					
Population	kid	Child	student	pupil		
Setting	school	education				
Intervention	intervention	program				
Topic of intervention	“health promotion” obesity	“over weight” overweight	adiposity	“physical activity”	“physical inactivity”	
OR	“physical active”	sport	sitting	sedentary	“media use”	“screen time” nutrition
	food	Diet				
Implementation phases	adoption	dissemination	adaptation	implementation	sustainability	“process evaluation”
Socioeconomic status	socioeconomic	socio-economic	income			

Eligibility criteria

We adopted the following eligibility criteria:

- Regarding the design, the article had to be an implementation evaluation or process evaluation study, or a hybrid process-effectiveness study as described by Curran et al. (41).
- The article had to investigate an intervention promoting obesity prevention behaviors (e.g., promotion of PA, promotion of health nutrition).
- The intervention reported, had to address children aged 5–14 years exclusively. The youngest age for beginning primary school is five years (42) and 14 years is the last year of childhood, before entering the youth category (43).
- The intervention reported, had to be conducted in a school setting
- The intervention reported, had to be conducted in an area with population of low SES or address children with low SES in particular.
- The article had to report influencing factors for the implementation of the intervention in their results section regarding children with low SES.

Operationalization of SES

Parental income, parental education and parental occupation are often used to measure children's SES (44). These measures are correlated but not interchangeable (44). Aggregated measures are also used to establish SES for school population or for regions or districts. Aggregated measures are drawn from administrative data and therefore depend on the institutional understanding of SES (44) and the availability of data. In this review, the mention of low SES was considered sufficient, and the specific criteria used to measure SES were not questioned, although this information was extracted from the articles and is shown in the results.

Screening process

After deduplication, two reviewer (FB and JE) independently screened the articles on title and abstract level and in a second step on full text level using the software Rayyan to determine inclusion (45). Conflicts were discussed and resolved between the reviewers. Additionally, all articles included in the review by Cassar et al. (34) were screened at full text level (n=26), as that review had a very similar aim, with the exception of the focus on SES.

Data extraction and synthesis

Article title, year of publication, country, main outcome of the article, means of data collection, criteria for low SES, and description of the intervention were extracted into an Word file (see Additional file 3) by MS and FB from each article. After the screening process, two reviewers (FB and JE) extracted barriers and facilitators for implementation from the results section of the articles into an Excel file. FB and JE extracted data from two articles independently and then matched their results through discussion. The remaining articles were split between FB and JE for data extraction. Any uncertainties about what details to extract, were additionally extracted by the second reviewer, matched and resolved through discussion.

The extracted data was coded using MAXQDA (46). The data coding was guided by the *Consolidated Framework of Implementation Research* (CFIR) (47). This comprises the five domains *intervention characteristics, inner setting, outer setting, characteristics of individuals* and *process*, along with 39 constructs within these domains (47). These 39 constructs from CFIR were used as categories for deductively coding the data. Each sense unit was coded into only one category.

FB and JE coded 25% of the data independently, then the codes were reconciled, and the rest of the data was coded by FB. In the next step, all coded segments for each category were reviewed by FB, and if necessary, the coding was adjusted, and additional specifications were developed for the categories for this review to have a clear differentiation between categories (see Additional file 4). To test the specified category descriptions, two reviewers (JE and a student assistant) coded 50% in total of the data again. The double-coded data were compared, and differences were discussed and resolved. All categories that caused more than one disagreement on the segment level were reviewed again for all data by FB. In the last step, all categories were reviewed, and reasonable inductive sub-categories were developed. These sub-categories then were coded again by a student assistant, and any disagreements were discussed and resolved.

For each of the five domains, the average number of articles that reported the (sub-)categories was calculated. (Sub-)Categories reported more than on average are presented in the results section.

We grouped and then compared the articles with regard to the characteristics of the interventions, following the definition for complex interventions by Craig et al. (48). If the intervention met two out of the three following aspects, it was considered to be *complex*, and otherwise it was considered the be *simple*: the intervention 1) addressed more than one obesity prevention behavior, 2) consisted of more than one component (e.g., classroom activities and teacher training), and 3) included parental involvement. We

analyzed the most frequently reported influencing factors within those groups of *simple* and *complex* interventions.

Methodological Quality assessment

We assessed the methodological quality of the articles using the Mixed Methods Appraisal Tool (MMAT) Version 2018 (49, 50). The MMAT allows the rating of quantitative and qualitative articles in the two separate corresponding categories, and mixed methods articles are rated in both, as well as an additional third mixed methods category.

Two reviewers (FB and JE) individually assessed the methodological quality of three articles, and the results were discussed with a third reviewer (CM). All of the remaining articles were split between two reviewers (FB and JE), and the methodological quality was individually assessed. If any uncertainties arose regarding the methodological quality of any particular, it was assessed and evaluated by the reviewers individually, resulted were discussed and the uncertainties resolved. No overall score was calculated, as recommended by the authors of MMAT (49).

3. Results

Study selection

From 6446 articles screened, 15 articles were identified as meeting all eligibility criteria. Screening the articles included by Cassar et al. (34), one additional article met all eligibility criteria, resulting in 16 articles for inclusion in this review (51-66) (see Figure 1).




Figure 1. Flow chart of screening.

Study characteristics

Additional file 3 presents detailed information on the included articles, such as their aim and information on the intervention reported. The aims of ten articles was to assess the implementation of the intervention and additional influencing factors for implementation (51, 53, 54, 56, 57, 59-61, 64, 66). Five articles only assessed influencing factors for implementation (52, 58, 62, 63, 65), and one assessed the influencing factors for implementation, as well as the effectiveness of the intervention (55). The articles applied mixed methods (n=10), qualitative (n=5) and quantitative design (n=1). The interventions reported by the articles promoted PA (54, 55, 59, 64, 66), healthy nutrition (56, 60, 61), PA and healthy nutrition (51, 52, 58, 62, 63, 65), and PA, healthy nutrition and reducing screen time (57) and PA, healthy nutrition, healthy sleep, and reduce screen time (53). In total, 14 independent interventions were reported, and three articles reported on the same intervention (58, 62, 63). Five articles were considered reporting on *simple* interventions (55, 59, 61, 64, 66), eleven articles were considered reporting on *complex* interventions (51, 52, 54, 56-58, 60, 62, 63, 65) (see Additional file 3).

Quality assessment results

Additional file 5 presents the ratings of the methodological quality assessment. Four (52, 58, 62, 63) of the five qualitative articles received a “yes” for all criteria, one articles received a “no” for the criterion “Is the interpretation of results sufficiently substantiated by data?” (65). The only quantitative article received a “can’t tell” for the criterion “Is the sampling strategy relevant to address the research question?” (61). Of the ten mixed methods articles, five (51, 55, 57, 59, 64) received a “yes” for all qualitative criteria, whereas only one of them received all quantitative criteria rated with “yes” (64). Five (51, 53, 54, 56, 57) of the 10 mixed methods articles received a rating of “yes” for all mixed methods criteria, one article (59) received a “no” for the criterion “Is there an adequate rationale for using a mixed methods design to address the research question?”. None of the mixed methods articles received only “yes” ratings for all criteria. In the mixed methods articles, the qualitative items rated lower than the qualitative articles.

Influencing factors for implementation

In the following, selected results are presented to answer the research question what the influencing factors for the implementation of school-based interventions promoting obesity prevention behaviors for children with low SES are.

Table 2 presents all included articles and their reporting of influencing factors in the five domains of CFIR, as well as the assignment to the groups of *simple* or *complex* interventions. The *inner setting* was reported in all articles (n=16), and the least reported domain was the *outer setting* (n=7). The *outer setting* was only reported in the group of *complex* interventions. Every article reported influencing factors in at least three different domains. In the 16 articles, 295 influencing factors were found across 93 (sub-)categories. Additional file 6 presents all influencing factors, categorized in barriers and facilitators, extracted from the articles for each domain and (sub-)category. Of the 39 original CFIR categories, four categories were not reported at all. Additional file 7 presents the most commonly reported (sub-)categories in the group of *simple* and *complex* interventions for each domain.

Table 2. Reported CFIR domains in included articles.

CFIR domain (number of (sub-)categories in domain)	article	Bartelink (51)*	Bauer (52)*	Blaine (53)*	de Meij (54)*	Gadais (55)°	Lepe (56)*	Lubans (57)*	Malek(58)*	Marchant(59)°	Martens (60)*	Muckelbauer (61)°	Norman (a) (62)*	Norman (b) (63)*	Steward (64)°	Verjans-Janssen (65)*	Wright(66)°	
Intervention characteristics (12)	x		x	x	x	x	x	x	x	x	x	x	x	x	x	x	15	
Inner setting (32)	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	16	
Outer setting (11)	x	x	x	x									x	x	x	x	7	
Characteristics of the individual (17)	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	14	
Process (21)	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	15	
	4	4	5	5	4	4	4	4	4	4	4	3	5	5	4	5	3	

°group of *simple* interventions*group of *complex* interventions

The results for each domain are presented below. Figure 2–8 show selected (sub-)categories for each domain, as well as the reported barriers and facilitators in those (sub-)categories. Furthermore, the most reported barrier(s) or facilitator(s) for each group of interventions is marked. Additional file 8 presents all (sub-)categories for each domain, and all the number of articles reporting the relevant barriers and facilitators.

Intervention characteristics

Intervention characteristics were reported by 14 articles, with 3.3 articles on average for each (sub)category. Figure 2 displays the four (sub-)categories that were reported more than averagely. Most articles reported *evidence strengths* (n=7) and *preparation* (n=7) influencing the implementation. Within the group of *simple* interventions, *adaptability* (n=3) was the most reported influencing factor, and within the group of *complex* interventions, *evidence strengths* (n=7) was the most reported influencing factor.




Figure 2. Intervention characteristics and most reported (sub-)categories.

Evidence strengths (n=7) was reported as a barrier, because no short- or long-term effects of the intervention (n=2) were seen. As facilitating for the implementation of the interventions, noticeable improvements in health (e.g., in the fitness level or self-confidence of children) (n=3) and successful connection of intervention to everyday life (n=3) were reported.

The preparation (n=7) of the intervention in the sub-category of design quality & packaging was reported as facilitator as introduction of the intervention (n=1) and intervention components with real life relevance (n=4) e.g., hands-on sessions, real life relevance of intervention components. In n=3 articles, barriers as inadequate intervention materials (e.g., wordiness of lessons) were reported.

Inner setting

All 16 articles reported (sub-)categories of the *inner setting*, with 2.8 articles on average for each (sub-)category. Figure 3 and Figure 4 display the 14 (sub-)categories that were reported more than the average. Most articles reported that *time* (n=7), *scheduling* (n=6) and *communication* (n=6) influenced the implementation. Within the group of *simple* interventions, *communication* and *scheduling* (n=2) were the most reported influencing factors, and within the group of *complex* interventions, *time* (n=6) was the most commonly reported influencing factor.




Figure 3. Inner setting (part 1) and most reported (sub-)categories.




Figure 4. Inner setting (part 2) and most reported (sub-)categories.

The most often reported aspect related to time (n=7), with sufficient time (n=2) facilitating and insufficient time hindering (n=5) for meetings, for training, for the children, for implementation as well as insufficient time due to the evaluation timeline (n=1) was also reported as hindering implementation.

Scheduling (n=6) was reported as a barrier (n=6), due to conflicts with scheduling (n=4), insufficient fine-tuned organizational procedures (n=1) and the school year was already planned, when the intervention was introduced (n=1). As a facilitator, scheduling was reported (n=2) in terms of a good fit of intervention in the work tasks (n=1) and scheduling the intervention activity before school was successful (n=1).

In the sub-category of *communication* (n=6), *good communication between stakeholders within school and with externals* was reported as a facilitator (n=5) and *miscommunication between school stakeholders* (n=2) was considered a barrier for implementation.

Outer setting

The *outer setting* were reported by seven articles, with 2.8 articles on average for each (sub-)category. Figure 5 displays the four (sub-)categories that were reported more than the average. *Abilities* (n=4), as a sub-category of *needs and resources of those served by the organization* and *existing policy* (n=4), as sub-category of *external policy and incentives* were the most reported. None of the articles within the group of *simple interventions* reported any influencing factors in the *outer setting*, and within the group of *complex interventions*, *abilities* and *existing policies* (n=4) were the most reported influencing factors.




Figure 5. Outer setting and most reported (sub-)categories.

Abilities (n=4) was reported as a barrier, because of lack of sufficient abilities among parents to conduct intervention / support children (n=3), whereas recognized and considered parents' abilities (n=1) facilitated implementation.

Existing policy (n=4), where lack of policy/expectation for intervention implementation (n=2) and lack of control (e.g., over administrative changes, food in cafeteria) (n=2) were reported as barriers. Financial support for the intervention (n=2) and fit between policies and intervention topics (n=2) were reported as facilitators for implementation.

Characteristics of individuals

Characteristics of individuals were reported by 14 articles, with 3.4 articles on average for each (sub-)category. Figure 6 presents the five (sub-)categories reported more than average. The most reported were *intervention strategy* (n=6) as sub-category of *individual stage of change* and *interest in intervention* (n=6) as sub-category of *other personal attributes*. Within the group of *simple* interventions, *effect of stage* (n=2) was the most reported influencing factor, and within the group of *complex* interventions, *character* and *interest in intervention* (n=5) were reported as sub-categories of *other personal attributes* as the most reported influencing factor.




Figure 6. characteristics of individuals and most reported (sub-)categories.

Intervention strategy (n=6) was reported as a barrier due to competitive elements of the intervention (n=1) and difficulty/ease of the intervention (n=1). As Competitive, playful and applied intervention components (n=5), and fitting the intervention to children's abilities and leading to gradual improvement (in fitness) (n=2) were reported as facilitators.

Interest in intervention (n=6) was reported as a barrier, due to a lack of interest by parents and children (n=2) and teachers' wish for additional training on topics other than the intervention topics (n=1). Interest in, enthusiasm for and commitment to the intervention from children (n=1) and from teachers (n=3) facilitated implementation.

Process

(Sub-)Categories in the *process* domain were reported by 15 articles, with 3.75 articles on average for each (sub-)category. Figure 7 and Figure 8 display the seven (sub-)categories reported more than average. The most reported (sub-)category that influences the implementation is *influence on executing* (n=8) as sub-category of *executing*. Within the group of *simple* interventions, *influence on executing* (n=3) was the most reported influencing factor. Within the group of *complex* interventions, *outcome* as a sub-category of *engaging parents*, *strategy* as a sub-category of *engaging key stakeholders* and *influence on executing* (n=5) were the most reported influencing factor.




Figure 7. process domain (part 1) with most reported (sub-)categories.




Figure 8. process domain (part 2) with most reported (sub-)categories.

Influence on executing (n=8) was reported as lack of support for children to finish intervention activities (n=1), in-class lessons that were too long or too diverse (n=2), parents lacking structure regarding the intervention (n=1), teachers conducting home activities in school due to lack of parental ability (n=1) and maintaining intervention guidelines leading to lack of enthusiasm (n=1). Practice-oriented thinking (n=1), children receiving support to finish intervention activities (n=1) and following the intervention guidelines (n=1) were reported as facilitators.

4. Discussion

The review identified, categorized and evaluated influencing factors for the implementation of school-based interventions promoting obesity prevention behaviors in children with low SES. We identified 295 influencing factors reported in 16 articles across 93 (sub-)categories in the five domains of CFIR. The articles examined were grouped in a set of five *simple* and eleven *complex* interventions.

Aspects of the *inner setting* (also referred to as *organizational*) were reported in every article, and aspects of the *outer setting* (also referred to as *context*) constituted the least reported domain. These findings are consistent with the results of comparable reviews that assessed influencing factors on interventions promoting PA (30, 36) and reducing sedentary behavior in school settings (34), as well as the sustainment of health behavior interventions in school settings and childcare services (35). It is likely that the eligibility criteria for 'school-based' interventions lead to the accumulation of factors in the *inner setting*. Furthermore, the *inner setting* is the most comprehensive domain of the CFIR. Although the *outer setting* is of great importance for implementation (29, 67-69), it is the least reported domain and in the group of *simple* interventions it is not reported at all.

On the (sub-)category level, some results are also consistent with the results of comparable reviews (30, 34-36). Yet, a comparison with other such reviews is only limited reasonable. E.g., the sub-category *insufficient time: insufficient time* was also found as barrier for implementation by Naylor et al. (30). However, if we consider the aspects reported in this review in the sub-category *insufficient time*, the five articles that reported this barrier indicated four different aspects where *insufficient time* was felt. For practical application, this means that even though there are consistent results on an aggregated level, the underlying aspects can be very divers.

Influencing factors—for intervention developers

We grouped the articles in a set of *simple* and *complex* interventions, because *complex* interventions might entail a wider range of influencing factors than *simple* interventions. Furthermore, there is moderate evidence that complex interventions are more effective than simple interventions (25-27). Comparing the groups of *simple* and *complex* interventions, one of the most reported influencing factors in the *process* domain was *executing* in both groups. *Executing* (also referred to as *fidelity*) defined as "carrying out or accomplishing the implementation according to plan" (47) and *adaptability* are highly connected. *Adaptability* was indeed the most reported influencing factor within the group of *simple* interventions in the domain *intervention characteristics*. Modifying *executing* is highly related to *adaptations* made, as adaptations mostly decrease the *executing* of an intervention. Adaptations are quite relevant for implementation (70) and *executing* is often used as outcome (71). To analyze the influence of *adaptability* and *executing* on health outcome, it is important to document and consider both (70).

We grouped the articles with regard to the intervention characteristics, following the criteria for *complex* interventions by Craig (48). Still, it seems like every intervention itself can be considered a complex intervention, following different criteria (e.g., synergies between intervention components, degree of flexibility, and multiplicity of mediators or moderators) (72). If the intervention itself is not complex, the school setting, with its context and stakeholders, and the interactions between them certainly can be considered complex (regardless of how simple or complicated the intervention is) (73, 74).

In addition, we noticed that the results of this review and those of other reviews varied. We assume that in school-based interventions, many different influencing factors appear, and every article differs in the degree of detail they use in assessing and reporting them.

Comparable reviews (30, 34-36) did not specifically address the target population of children with low SES. We found different influencing factors for the implementation of interventions for the population of children with low SES than such reviews without the focus on children with a low SES. We cannot evaluate whether these differences due to

the different target populations. Articles were included in our review, if they reported that their population has a low SES. All included articles measured the SES on an aggregated school or area level, so their comparability is limited. There are children with low SES living in areas where the population is predominantly of a mid to high SES. Those children might have received an intervention, but due to eligibility criteria, these articles are not included in this review. On the other hand, in an area where the population is predominantly of a low SES, not all children necessarily will have a low SES. (75)

The questions now arises, which categories or aspects should be considered when developing new or adapting existing interventions for new settings and scaling them up. Every setting and organization has different needs and resources. The group that receive interventions, such as those of a low SES, must be involved in the uptake of effective health-promoting interventions (76). Participatory approaches can increase the likelihood of successful implementation, and improving the sustainability of interventions and can help balance top-down and bottom-up approaches (77, 78). The intervention mapping approach (79), implementation mapping (33), and the closely related method of co-creation (80) offer guidance for a participatory intervention development and the implementation process.

School setting—for school staff

All organizations require resources to conduct health-promoting activities. A team for implementation, a health supporting culture, and a head teacher, who supports the intervention are likely to be important factors for successful implementation (81).

In the following, we would like to give some suggestions for school (head) teachers, school health-promoter, and social workers: Networks and collaboration can be developed with those who are willing within the school and with other schools and institutions. It is important to be open and willing to try different things, and to be ready to adjust aspects of the intervention, as each institution has different preconditions, needs, and resources. Working with the community is a promising opportunity, as those collaborations can improve the community networks and benefit school and students (82).

For children outside the school, the family is a very important setting. With regard to school-based nutrition and PA interventions, there is no clear conclusion whether parental involvement has a positive influence on the effectiveness of school-based PA and nutrition interventions (83). Regarding school-based nutrition and PA interventions with direct parental involvement (e.g., completing a questionnaire would not count as direct involvement), Verjans-Janssen et al. (84) found mainly positive effects. This may indicate the influence and importance of direct parental involvement in school-based interventions (84), especially in obesity prevention interventions (39) and for children with low SES (75, 85).

Guidance has been created addressed to schools on how to implement health-promoting activities (86, 87). Evidence-informed guidance is of great importance, however those guidance tend to exhibit a quite theoretical perspective. Building on this foundation, there is still a need for empirical tested and actionable strategies the theory practice translation.

Methodological considerations—for researcher

Because a wide variety of implementation frameworks exist (88, 89), this review also indented to help standardize and increases comparability of results in implementation research in general (90) and for school-based/obesity interventions specifically, using CFIR.

CFIR offers several advantages, due to its constant development (91) and its method of rating determinants (92). Furthermore, the CFIR outcome addendum (93), and the CFIR-ERIC (Expert Recommendations for Implementation Change: a summary of 73 implementation strategies) (94) matching tool (95). Identifying and reporting influencing

factors using CFIR, very detailed information can be presented, using the categories, as well as more generally by using the domains. This offers comparability on different levels.

In this review and in many other instances, the application of CFIR is descriptive and linear. This review focused on identifying and evaluating influencing factors for implementation of school-based interventions preventing obesity prevention behaviors for children with low SES. The search term, and the eligibility criteria were chosen accordingly. There are issues this review could not and would not answer, but future research should address. It is important to quantify the influencing factors (92, 96), and to analyze which influencing factors are interconnected. It is essential to analyze the factors that have an influence on implementation outcomes (97-99) and on health outcomes (100). Choosing appropriate implementation strategies (101) and organizing the whole process with an overall evaluation plan (33) should be the standard in implementation evaluation in general and in children with low SES in particular.

Limitations

There are several limitations to the results of this review. Any intervention automatically does the implementation by conducting the intervention in real-world settings, and the corresponding articles might also report about implementation aspects. Articles in which reported aspects are not labeled as implementation evaluation or process evaluation results though, were not found with the search term and therefore not included in this review, although they might have contained important findings. Due to lack of transparency how the influencing factors were identified, influencing factors must have been reported in the results section. Articles reporting influencing factors only in the discussion were excluded. In addition, some clustering of influencing factors may not have occurred, because the number of 16 articles in total, five in the group of *simple* interventions, and eleven in the group of *complex* interventions, was too small.

5. Conclusion

This review is the first assessing influencing factors for the implementation of interventions promoting obesity prevention behaviors in children with low SES specifically, and contributes to existing literature regarding health equity. We identified influencing factors for implementation and presented them on a very detailed level. This enhances the presentation of results at the most applicable level, and contributes to the translation between theory and practice. This review highlights the need of empirical research investigating the processes and dynamics during the adoption, implementation and sustainment of an intervention as a whole.

List of abbreviations: CFIR: Consolidated Framework of Implementation Research; HI: health inequity; MMAT: Mixed Methods Appraisal Tool; PA: physical activity; SES: socioeconomic status

Author Contributions: FB directed the planning of this review. FB developed and refined the search strategy. FB conducted the database search, the screening, methodological quality assessment, data extraction and data analysis. JE contributed to the screening, methodological quality assessment, data extraction and data analysis. MS contributed to the data extraction and preparation of supplementary files. CM contributed to developing and refining the search strategy and to the methodological quality assessment. FB wrote the manuscript, CM and JE commented on the manuscript. CM advised throughout the review process, and contributed to and commented on the manuscript. All authors read and approved the final manuscript.

Funding: This review was funded by the German Federal Ministry of Health in the Project "Familie+ – living healthy together in family and school" (ZMVI1-2519KIG006). Open Access funding by Technical University of Munich enabled and organized by Project DEAL. English lecture service funding by the Graduate School of the Technical University of Munich.

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments: We would like to thank Elaine Fischer and Julia Frank for supporting the qualitative analysis for this article as well as Antonia Frank for supporting the preparation of the graphics. Furthermore, we would like to thank Doris Gebhard for providing important advice in the course of shaping this article.

Conflicts of Interest: The authors declare that they have no competing interests.

References

1. Odeny B. Closing the health equity gap: A role for implementation science? *PLoS Med.* 2021;18(9):e1003762.
2. World Health Organization. *World health statistics 2021: monitoring health for the SDGs, sustainable development goals.* World Health Organization; 2021.
3. World Health Organization. *Action in the social determinants of health: Learning from previous experiences.* World Health Organization; 2010.
4. World Health Organization. *Report of the International Conference on Primary Health Care, Alma-Ata, USSR, 6-12 September 1978.* Alma-Ata: World Health Organization; 1978.
5. Marmot M, Friel S, Bell R, Houweling TA, Taylor S, Health CoSDo. Closing the gap in a generation: health equity through action on the social determinants of health. *The lancet.* 2008;372(9650):1661-9.
6. Whitehead M, Dahlgren G. *Concepts and principles for tackling social inequities in health: Levelling up Part 1.* Copenhagen: World Health Organization; 2006.
7. Braveman P. What is health equity: and how does a life-course approach take us further toward it? *Matern Child Health J.* 2014;18(2):366-72.
8. Nutbeam D. *Health Promotion Glossary.* Genf: World Health Organization; 1998. Report No.: 0957-4824.
9. Whitehead M. The concepts and principles of equity and health. *Int J Health Serv.* 1992;22(3):429-45.
10. Diderichsen F, Evans T, Whitehead M. The Social Basis of Disparities in Health. In: Evans T, Whitehead M, Diderichsen F, Bhuiya A, Wirth M, editors. *Challenging Inequities in Health: From Ethics to Action.* New York: Oxford University Press; 2001. p. 12-23.
11. Braveman PA, Cubbin C, Egerter S, Chideya S, Marchi KS, Metzler M, et al. Socioeconomic status in health research: one size does not fit all. *Jama.* 2005;294(22):2879-88.
12. World Health Organization. *Report on social determinants of health and the health divide in the WHO European Region.* Copenhagen: World Health Organization; 2012.
13. Pearce A, Dundas R, Whitehead M, Taylor-Robinson D. Pathways to inequalities in child health. *Arch Dis Child.* 2019;104(10):998-1003.
14. Guyer B, Ma S, Grason H, Frick KD, Perry DF, Sharkey A, et al. Early childhood health promotion and its life course health consequences. *Acad Pediatr.* 2009;9(3):142-9. e71.
15. Braveman P, Barclay C. Health disparities beginning in childhood: a life-course perspective. *Pediatrics.* 2009;124 Suppl 3:163-75.
16. World Health Organization. *Report of the Commission on Ending Childhood Obesity. Implementation plan: executive summary.* Genf: World Health Organization; 2017.
17. Reilly JJ, Kelly J. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: systematic review. *Int J Obes.* 2011;35(7):891-8.
18. Barton M. Childhood obesity: a life-long health risk. *Acta Pharmacol Sin.* 2012;33(2):189-93.
19. Abarca-Gómez L, Abdeen ZA, Hamid ZA, Abu-Rmeileh NM, Acosta-Cazares B, Acuin C, et al. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *The Lancet.* 2017;390(10113):2627-42.
20. United Nations Children's Fund (UNICEF), World Health Organization, Bank IBfRaDTW. *Levels and trends in child malnutrition: key findings of the 2021 edition of the joint child malnutrition estimates.* Geneva: World Health Organization; 2021.
21. Vazquez CE, Cubbin C. Socioeconomic Status and Childhood Obesity: a Review of Literature from the Past Decade to Inform Intervention Research. *Current Obesity Reports.* 2020;1-9.
22. Dahlgren G, Whitehead M. *European strategies for tackling social inequities in health: Levelling up Part 2.* Copenhagen: World Health Organization; 2006.
23. Evans T, Whitehead M, Bhuiya A, Diderichsen F, Wirth M. *Challenging inequities in health: from ethics to action.* New York: Oxford University Press; 2001.
24. Brown T, Moore TH, Hooper L, Gao Y, Zayegh A, Ijaz S, et al. Interventions for preventing obesity in children. *Cochrane Database Syst Rev.* 2019;7:CD001871.
25. Bleich SN, Vercammen KA, Zatz LY, Frelier JM, Ebbeling CB, Peeters A. Interventions to prevent global childhood overweight and obesity: a systematic review. *The Lancet Diabetes & Endocrinology.* 2018;6(4):332-46.

26. Khambalia AZ, Dickinson S, Hardy LL, Gill T, Baur LA. A synthesis of existing systematic reviews and meta-analyses of school-based behavioural interventions for controlling and preventing obesity. *Obes Rev.* 2012;13(3):214-33.

27. Sobol-Goldberg S, Rabinowitz J, Gross R. School-based obesity prevention programs: a meta-analysis of randomized controlled trials. *Obesity (Silver Spring).* 2013;21(12):2422-8.

28. World Health Organization. School policy framework: implementation of the WHO global strategy on diet, physical activity and health. . World Health Organization; 2008.

29. Durlak JA, DuPre EP. Implementation matters: a review of research on the influence of implementation on program outcomes and the factors affecting implementation. *Am J Community Psychol.* 2008;41(3-4):327-50.

30. Naylor PJ, Nettlefold L, Race D, Hoy C, Ashe MC, Wharf Higgins J, et al. Implementation of school based physical activity interventions: a systematic review. *Prev Med.* 2015;72:95-115.

31. Brown CH, Curran G, Palinkas LA, Aarons GA, Wells KB, Jones L, et al. An Overview of Research and Evaluation Designs for Dissemination and Implementation. *Annu Rev Public Health.* 2017;38:1-22.

32. Rabin B, Brownson R. Terminology for Dissemination and Implementation Research. In: Brownson R, Colditz G, Proctor E, editors. *Dissemination and Implementation Research in Health: Translating Science to Practice.* 2nd edn. New York: Oxford University Press; 2017.

33. Fernandez ME, Ten Hoor GA, van Lieshout S, Rodriguez SA, Beidas RS, Parcel G, et al. Implementation Mapping: Using Intervention Mapping to Develop Implementation Strategies. *Front Public Health.* 2019;7:158.

34. Cassar S, Salmon J, Timperio A, Naylor PJ, van Nassau F, Contardo Ayala AM, et al. Adoption, implementation and sustainability of school-based physical activity and sedentary behaviour interventions in real-world settings: a systematic review. *Int J Behav Nutr Phys Act.* 2019;16(1):120.

35. Shoesmith A, Hall A, Wolfenden L, Shelton RC, Powell BJ, Brown H, et al. Barriers and facilitators influencing the sustainment of health behaviour interventions in schools and childcare services: a systematic review. *Implement Sci.* 2021;16(1):62.

36. Michael RD, Webster CA, Egan CA, Nilges L, Brian A, Johnson R, et al. Facilitators and Barriers to Movement Integration in Elementary Classrooms: A Systematic Review. *Res Q Exerc Sport.* 2019;90(2):151-62.

37. Meshkovska B, Scheller DA, Wendt J, Jilani H, Scheidmeir M, Stratil JM, et al. Barriers and facilitators to implementation of direct fruit and vegetables provision interventions in kindergartens and schools: a qualitative systematic review applying the consolidated framework for implementation research (CFIR). *Int J Behav Nutr Phys Act.* 2022;19(1).

38. Brownson RC, Kumanyika SK, Kreuter MW, Haire-Joshu D. Implementation science should give higher priority to health equity. *Implement Sci.* 2021;16(1):28.

39. Rennie KL, Johnson L, Jebb SA. Behavioural determinants of obesity. *Best Pract Res Clin Endocrinol Metab.* 2005;19(3):343-58.

40. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71.

41. Landes SJ, McBain SA, Curran GM. Reprint of: An introduction to effectiveness-implementation hybrid designs. *Psychiatry Res.* 2020;283:112630.

42. UNESCO Institute for Statistics. Official Entrance Age to Primary Education (years). 2021.

43. United Nations GA. International Youth Year: Participation, Development, Peace. 1981.

44. Sirin SR. Socioeconomic Status and Academic Achievement: A Meta-Analytic Review of Research. *Review of Educational Research.* 2005;75(3):417-53.

45. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev.* 2016;5(1):210.

46. VERBI Software. MAXQDA 2020. Berlin, Germany: VERBI Software; 2020.

47. Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci.* 2009;4:50.

48. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M, et al. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ.* 2008;337:a1655.

49. Hong Q, Pluye P, Fàbregues S, Bartlett G, Boardman F, Cargo M, et al. Mixed Methods Appraisal Tool (MMAT). Canadian Intellectual Property Office. 2018;Industry Canada.

50. Pace R, Pluye P, Bartlett G, Macaulay AC, Salsberg J, Jagosh J, et al. Testing the reliability and efficiency of the pilot Mixed Methods Appraisal Tool (MMAT) for systematic mixed studies review. *Int J Nurs Stud.* 2012;49(1):47-53.

51. Bartelink NHM, van Assema P, Jansen MWJ, Savelberg H, Moore GF, Hawkins J, et al. Process evaluation of the healthy primary School of the Future: the key learning points. *BMC Public Health.* 2019;19(1):698.

52. Bauer KW, Patel A, Prokop LA, Austin SB. Swimming upstream: faculty and staff members from urban middle schools in low-income communities describe their experience implementing nutrition and physical activity initiatives. *Prev Chronic Dis.* 2006;3(2):A37.

53. Blaine RE, Franckle RL, Ganter C, Falbe J, Giles C, Criss S, et al. Using School Staff Members to Implement a Childhood Obesity Prevention Intervention in Low-Income School Districts: the Massachusetts Childhood Obesity Research Demonstration (MA-CORD Project), 2012-2014. *Prev Chronic Dis.* 2017;14:E03.

54. De Meij JS, van der Wal MF, van Mechelen W, Chinapaw MJ. A mixed methods process evaluation of the implementation of JUMP-in, a multilevel school-based intervention aimed at physical activity promotion. *Health Promot Pract.* 2013;14(5):777-90.

55. Gadais T, Caron T, Ayoub MB, Karelis A, Nadeau L. The Role of the Teacher in the Implementation of a School-Based Intervention on the Physical Activity Practice of Children. *Int J Environ Res Public Health.* 2020;17(19):7344.

56. Lepe S, Goodwin J, Mulligan KT, Balestracci K, Sebelia L, Greene G. Process Evaluation of a Policy, Systems, and Environmental Change Intervention in an Urban School District. *J Nutr Educ Behav.* 2019;51(3):307-17.

57. Lubans DR, Smith JJ, Skinner G, Morgan PJ. Development and implementation of a smartphone application to promote physical activity and reduce screen-time in adolescent boys. *Front Public Health.* 2014;2:42.

58. Malek ME, Nyberg G, Elinder LS, Patterson E, Norman Å. Children's experiences of participating in a school-based health promotion parental support programme - a qualitative study. *BMC Pediatr.* 2021;21(1):228.

59. Marchant E, Todd C, Stratton G, Brophy S. The Daily Mile: Whole-school recommendations for implementation and sustainability. A mixed-methods study. *PLoS One.* 2020;15(2):e0228149.

60. Martens M, van Assema P, Paulussen T, Schaalma H, Brug J. Krachtvoer: process evaluation of a Dutch programme for lower vocational schools to promote healthful diet. *Health Educ Res.* 2006;21(5):695-704.

61. Muckelbauer R, Libuda L, Clausen K, Kersting M. Long-term process evaluation of a school-based programme for overweight prevention. *Child Care Health Dev.* 2009;35(6):851-7.

62. Norman Å, Nyberg G, Elinder LS, Berlin A. One size does not fit all-qualitative process evaluation of the Healthy School Start parental support programme to prevent overweight and obesity among children in disadvantaged areas in Sweden. *BMC Public Health.* 2016;16:37.

63. Norman Å, Nyberg G, Berlin A. School-based obesity prevention for busy low-income families-Organisational and personal barriers and facilitators to implementation. *PLoS One.* 2019;14(11):e0224512.

64. Stewart G, Webster CA, Weaver RG, Stodden DF, Brian A, Egan CA, et al. Evaluation of a classroom movement integration training delivered in a low socioeconomic school district. *Eval Program Plann.* 2019;73:187-94.

65. Verjans-Janssen SRB, Gerards S, Verhees AH, Kremers SPJ, Vos SB, Jansen MWJ, et al. Implementation of KEIGAAF in Primary Schools: A Mutual Adaptation Physical Activity and Nutrition Intervention. *Int J Environ Res Public Health.* 2020;17(3):751.

66. Wright CM, Chomitz VR, Duquesnay PJ, Amin SA, Economos CD, Sacheck JM. The FLEX study school-based physical activity programs - measurement and evaluation of implementation. *BMC Public Health.* 2019;19(1):73.

67. Pfadenhauer LM, Gerhardus A, Mozygemba K, Lysdahl KB, Booth A, Hofmann B, et al. Making sense of complexity in context and implementation: the Context and Implementation of Complex Interventions (CICI) framework. *Implement Sci.* 2017;12(1):21.

68. Nilsen P, Bernhardsson S. Context matters in implementation science: a scoping review of determinant frameworks that describe contextual determinants for implementation outcomes. *BMC Health Serv Res.* 2019;19(1):189.

69. Edwards N, Barker PM. The importance of context in implementation research. *J Acquir Immune Defic Syndr.* 2014;67 Suppl 2:157-62.

70. Chambers DA, Norton WE. The Adaptome: Advancing the Science of Intervention Adaptation. *Am J Prev Med.* 2016;51(4 Suppl 2):124-31.

71. Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Administration and Policy in Mental Health and Mental Health Services Research.* 2011;38(2):65-76.

72. Petticrew M, Anderson L, Elder R, Grimshaw J, Hopkins D, Hahn R, et al. Complex interventions and their implications for systematic reviews: a pragmatic approach. *J Clin Epidemiol.* 2013;66(11):1209-14.

73. Rosas SR. Systems thinking and complexity: considerations for health promoting schools. *Health Promot Int.* 2017;32(2):301-11.

74. Dryden-Palmer KD, Parshuram CS, Berta WB. Context, complexity and process in the implementation of evidence-based innovation: a realist informed review. *BMC Health Serv Res.* 2020;20(1):81.

75. Lynch JW, Law C, Brinkman S, Chittleborough C, Sawyer M. Inequalities in child healthy development: some challenges for effective implementation. *Soc Sci Med.* 2010;71(7):1244-8.

76. Yancey A, Glenn BA, Ford CL, Bell-Lewis L. Dissemination and Implementation Research among Racial/Ethnic Minority and Other Vulnerable Populations. *Dissemination and Implementation Research in Health: Translating Science to Practice:* Oxford University Press; 2017.

77. Ramanadhan S, Davis MM, Armstrong R, Baquero B, Ko LK, Leng JC, et al. Participatory implementation science to increase the impact of evidence-based cancer prevention and control. *Cancer Causes Control.* 2018;29(3):363-9.

78. Minkler M, Salvatore AL, Chang C. Participatory Approaches for Study Design and Analysis in Dissemination and Implementation Research. In: Brownson RC, Colditz GA, Proctor EK, editors. *Dissemination and Implementation Research in Health: Translating Science to Practice:* Oxford University Press; 2017.

79. Bartholomew Eldredge LK, Markham CM, Ruiter RA, Fernandez ME, Kok G, Parcel GS. Planning health promotion programs: an intervention mapping approach. 4 ed: John Wiley & Sons; 2016.

80. Leask CF, Sandlund M, Skelton DA, Altenburg TM, Cardon G, Chinapaw MJM, et al. Framework, principles and recommendations for utilising participatory methodologies in the co-creation and evaluation of public health interventions. *Res Involv Engagem.* 2019;5:2.

81. Aarons GA, Moullin JC, Ehrhart MG. The Role of Organizational Processes in Dissemination and Implementation Research. In: Brownson RC, Colditz GA, Proctor EK, editors. *Dissemination and Implementation Research in Health: Translating Science to Practice:* Oxford University Press; 2017.

82. Sanders M. School-Community Partnerships The Little Extra That Makes a Big Difference. In: Epstein JL SM, Sheldon SB, Simon BS, Clark Salinas K, Rodriguez Jansorn N, Van Voorhis FL, Martin CS, Thomas BG, Greenfeld MD, Hutchins DJ, Williams KJ., editor. *School, Family, and Community Partnerships: your handbook for action.* 4. ed. California: Corwin; 2019.

83. Van Lippevelde W, Verloigne M, De Bourdeaudhuij I, Brug J, Bjelland M, Lien N, et al. Does parental involvement make a difference in school-based nutrition and physical activity interventions? A systematic review of randomized controlled trials. *Int J Public Health*. 2012;57(4):673-8.

84. Verjans-Janssen SRB, van de Kolk I, Van Kann DHH, Kremers SPJ, Gerards S. Effectiveness of school-based physical activity and nutrition interventions with direct parental involvement on children's BMI and energy balance-related behaviors - A systematic review. *PLoS One*. 2018;13(9):e0204560.

85. Kader M, Sundblom E, Elinder LS. Effectiveness of universal parental support interventions addressing children's dietary habits, physical activity and bodyweight: A systematic review. *Prev Med*. 2015;77:52-67.

86. Sharples J, Albers B, Fraser S, Kime S. Putting Evidence to Work: A School's Guide to Implementation. London: Education Endowment Foundation (EEF); 2019.

87. Organization WH. Making every school a health-promoting school-Implementation Guidance. World Health Organization; 2021

88. Nilsen P. Making sense of implementation theories, models and frameworks. *Implement Sci*. 2015;10:53.

89. Shelton RC, Lee M, Brotzman LE, Wolfenden L, Nathan N, Wainberg ML. What Is Dissemination and Implementation Science?: An Introduction and Opportunities to Advance Behavioral Medicine and Public Health Globally. *Int J Behav Med*. 2020;27(1):3-20.

90. Kirk MA, Kelley C, Yankey N, Birken SA, Abadie B, Damschroder L. A systematic review of the use of the Consolidated Framework for Implementation Research. *Implement Sci*. 2016;11:72.

91. Damschroder L, Reardon CM, Widerquist MAO, Lowery JC. The Updated Consolidated Framework for Implementation Research: CFIR 2.0. Research Square. 2022.

92. Damschroder L, Lowery J. Evaluation of a large-scale weight management program using the consolidated framework for implementation research (CFIR). *Implement Sci*. 2013;8.

93. Damschroder LJ, Reardon CM, Opra Widerquist MA, Lowery J. Conceptualizing outcomes for use with the Consolidated Framework for Implementation Research (CFIR): the CFIR Outcomes Addendum. *Implement Sci*. 2022;17(1):7.

94. Powell BJ, Waltz TJ, Chinman MJ, Damschroder LJ, Smith JL, Matthieu MM, et al. A refined compilation of implementation strategies: results from the Expert Recommendations for Implementing Change (ERIC) project. *Implement Sci*. 2015;10:21.

95. CFIR Research Team-Center for Clinical Management Research. CFIR-ERIC Implementation Strategy Matching Tool 2022 [Available from: <https://cfirguide.org/choosing-strategies/>].

96. Lobczowska K, Banik A, Brukalo K, Forberger S, Kubiak T, Romaniuk P, et al. Meta-review of implementation determinants for policies promoting healthy diet and physically active lifestyle: application of the Consolidated Framework for Implementation Research. *Implement Sci*. 2022;17(1):2.

97. Jong ST, Croxson CHD, Foubister C, Brown HE, Guell C, Lawlor ER, et al. Reach, Recruitment, Dose, and Intervention Fidelity of the GoActive School-Based Physical Activity Intervention in the UK: A Mixed-Methods Process Evaluation. *Children (Basel)*. 2020;7(11):231.

98. Trigwell J, McGee CE, Murphy RC, Porcellato LA, Ussher M, Garnham-Lee K, et al. Process evaluation of a sport-for-health intervention to prevent smoking amongst primary school children: SmokeFree Sports. *BMC Public Health*. 2015;15:347.

99. Griffin TL, Clarke JL, Lancashire ER, Pallan MJ, Adab P, investigators Wst. Process evaluation results of a cluster randomised controlled childhood obesity prevention trial: the WAVES study. *BMC Public Health*. 2017;17(1):681.

100. Springer AE, Kelder SH, Ranjit N, Hochberg-Garrett H, Crow S, Delk J. Promoting physical activity and fruit and vegetable consumption through a community-school partnership: the effects of Marathon Kids® on low-income elementary school children in Texas. *J Phys Act Health*. 2012;9(5):739-53.

101. Cook CR, Lyon AR, Locke J, Waltz T, Powell BJ. Adapting a Compilation of Implementation Strategies to Advance School-Based Implementation Research and Practice. *Prev Sci*. 2019;20(6):914-35.

List of additional files

Additional file 1

- Word document .docx
- PRISMA guidelines for reporting
- Checklist of PRISMA guidelines for reporting

Additional file 2

- Word document .docx
- Search strategy
- Search strategy and serach term used in the different data bases

Additional file 3

- Word document .docx
- List of included articles
- Description of included articles, with e.g. information about article design, data collection and intervention's description

Additional file 4

- Excel file .xlsx
- Consolidated Framework for Implementation Research Constructs with specifications
- Consolidated Framework for Implementation Research Constructs with specifications made for this review

Additional file 5

- Excel file .xlsx
- Mixed Methods Appraisal Tool ratings
- Methodological quality assessment ratings for each article using the Mixed Methods Appraisal Tool items

Additional file 6

- Excel file .xlsx
- Extracted and categorized influencing factors
- Extracted and categorized influencing factors for all domains and categories of the Consolidated Framework for Implementation Research Constructs

Additional file 7

- Excel file .xlsx
- Most commonly reported (sub-)categories by group of simple and complex interventions
- The most commonly reported (sub-)categories in the group of simple and complex interventions for each CFIR domain.

Additional file 8

- PowerPoint file .pptx
- Number of reported barriers/facilitators and (sub-)categories for each CFIR domain
- All (sub-)categories for each of the five CFIR domains, and all the number of articles reporting the relevant barriers and facilitators

Figure 2-8 editable

- PowerPoint file .pptx
- Figure 2-8 are provided as PDF as part of the manuscript and they are provided in PowerPoint as editable files.