

Review

1

Improving Abiotic Stress Tolerance of Forage Grasses with Ge-Nome Editing Tools

2

3

Ferenz Sustek-Sánchez ¹, Odd-Arne Rognli ², Nils Rostoks ³, Merike Sõmera ¹, Kristina Jaškunė ⁴, Mallikarjuna Rao Kovi ², Gražina Statkevičiūtė ⁴ and Cecilia Sarmiento ^{1,*}

4

5

¹ Department of Chemistry and Biotechnology, Tallinn University of Technology, Tallinn, Estonia

6

² Department of Plant Sciences, Norwegian University of Life Sciences (NMBU), Ås, Norway.

7

³ Faculty of Biology, University of Latvia, Riga, Latvia.

8

⁴ Institute of Agriculture, Lithuanian Research Centre for Agriculture and Forestry, Kedainiai, Lithuania.

9

* Correspondence: cecilia.sarmiento@taltech.ee

10

Abstract: Due to an increase in the consumption of food, feed, and fuel and to meet global food security needs for the rapidly growing human population, there is a necessity to obtain high-yielding crops that can adapt to future climate changes. Currently, the main feed source used for ruminant livestock production is forage grasses. In temperate climate zones, perennial grasses grown for feed are widely distributed and tend to suffer under unfavorable environmental conditions. Gene editing has been shown to be an effective tool for the development of abiotic stress-resistant plants. The highly versatile CRISPR-Cas system enables increasingly complex modifications in genomes while maintaining precision and low off-target frequency mutations. In this review, we provide an overview of forage grass species that have been subjected to gene editing. We offer a perspective view on the generation of plants resilient to abiotic stresses. Due to the broad factors contributing to these stresses the review focuses on drought, salt, heat, and cold stresses. The application of new genomic techniques (e.g., CRISPR-Cas) allows addressing several challenges caused by climate change and abiotic stresses for developing forage grass cultivars with improved adaptation to the future climatic conditions. Gene editing will contribute towards developing safe and sustainable food systems.

Keywords: CRISPR; genome editing; gene editing; forage grass; abiotic stress; plant

26

27

1. Introduction

28

Grasses belong to the family of Poaceae, which constitutes the most economically important plant family [1,2]. Grasslands and meadows extend over vast portions of the planet, on land, and even under the sea [3,4]. Their importance in Earth's ecosystems goes beyond their use in fields and pastures. Grassy biomes comprise more than one-quarter of the planet's land area. Grasses not only provide food, shelter, and building materials for animals and humans, but they also generate oxygen and store carbon [5]. This storage, mainly subterranean, contributes towards the fertilization of soils and makes grasslands valuable sinks of CO₂ [6,7]. Furthermore, grasses are considered more resilient to dryer and warmer conditions than trees. These facts suggest that in the climatic conditions predicted for the future, grasslands could be a better and more robust carbon sink than forests [8].

29

30

31

32

33

34

35

36

37

38

39

Grass crops provide the most essential dietary food sources globally. From these, forage grasses are the main component used to feed ruminant livestock [9,10]. Grasses can be cultivated in less fertile lands compared to other crops. In these zones, normally associated with developing countries [11,12], animal husbandry and its derivatives e.g., dairy products, remain essential [13,14]. To cope with the predicted population growth and the consequential increase in food needs, high-yielding crops must be further developed [15]. To reach food security, the strategies used must avoid causing negative environmental

40

41

42

43

44

45

46



impacts. Synthetic nitrogen-based fertilizers have been important for reaching high yields, nevertheless, their production and usage are a source of massive generation and emission of greenhouse gases (GHGs) [16]. It is well known that the high concentration of atmospheric GHGs is closely related to climate change. Therefore, the challenge is to increase farming efficiency while reducing the impact of agricultural activity on climate change [17]. Importantly, climate change not only directly affects crop productivity but also has indirect and socio-economic impacts, for instance soil fertility, need for irrigation, food demand, policy, rising costs (reviewed in [15]).

Grasses usage as forage and as reliable sinks of carbon emissions, call for an improvement in their biomass yield, and their resistance towards the new abiotic and biotic stresses caused by climate change [18]. Especially, plants will have to cope with variations in temperature, water availability, and soil composition [19]. Said variations will generate stresses due to heat, cold, drought, and salinity conditions. A promising approach to provide grasses with stress resistance is using gene editing techniques [20,21]. The first attempts have been performed to use gene editing in forage grasses [22–29]. This is not an easy task due to their reproductive and genetic characteristics which are difficult to work with. The inability of forage grasses to self-pollinate hinders inbreeding. Additionally, forage grasses have high variability between the genetic background of different individuals. This provides them with a considerable gene pool, responsible for their adaptability and resilience towards environmental changes. Conversely, it creates difficulties for studies focused on identifying the genetic cause of traits or phenotypes of interest [30,31]. There are diverse ways of classifying grasses beyond their taxonomy. For instance, forage grasses can be divided into different types depending on their life cycle and ecotype. In the first case, according to the survival of the plant after going through its reproductive phase, grasses can be considered annual, biannual, or perennial. In terms of their ecotype, grasses can be separated into warm- or cool-season plants if their optimal growth happens during winter or summer, respectively. Importantly, warm-season grasses are C_4 plants, while cool-season grasses are C_3 plants [32,33].

In this review, we provide an overview of the main metabolic and molecular changes that plants suffer to cope with the effects of abiotic stress derived from climate change. Additionally, we summarize the actual state of gene editing applications in forage grasses. We propose how gene editing could be used to generate grass plants able to resist these abiotic stresses. Finally, we hypothesize how the new genetic resources and tools can be used to improve forage grass breeding that will help achieve food security in a sustainable way.

2. Cellular and molecular responses to cope with the main abiotic stresses

Extreme temperatures, uncommon precipitation patterns, and deterioration of soils are being observed due to climate change. These environmental consequences have a great impact on agriculture since plants are of sessile nature. The responses used by plants when encountering a stressor aim firstly to achieve acclimation to the new environment and later adaptation to it. Acclimation includes adjusting the physiology and metabolism of a plant to achieve a new state of homeostasis, while adaptation involves both phenotypic and genotypic alterations. Acclimation mediates quick responses to ensure the survival of a plant, whereas adaptation is considered an evolutionary and lengthy process whose goal is to preserve a population. Plants must cope with new and more extreme conditions, which lead to different abiotic and biotic stresses than those commonly present in their biomes [34]. Abiotic stresses are those derived from the physical and chemical factors of an environment and are independent of living organisms [35]. As a response to these environmental alterations, plants undergo morphological, metabolic, and physiological changes. In this review, we will focus on drought, salinity, cold, and heat stress responses at the cellular and molecular levels. These are not the only abiotic conditions that will vary due to climate change, but they represent some of the major alterations that will result from it [35,36]. The stresses discussed in this review have a significant impact

on the growth and development of plants, which is directly connected to crops' yield and profitability [37,38].

Even though the abiotic stresses will be described separately, in nature they tend to interact producing greater effects than individually. Therefore, plants normally must acclimate to a combination of stresses. This should not be ignored when designing strategies to improve crops' tolerance to stress [39].

2.1. Temperature conditions

One of climate change's main effects is the alteration of temperature conditions [40]. Temperature affects and limits plant growth and development directly [41]. Therefore, it has a great impact on crop yield which is associated with food security [9,10,40]. It is considered that there are two abiotic stresses derived from temperature variations: heat and cold stress.

2.1.1. Heat stress

As a direct consequence of climate change, global warming has led to steady and yearly temperature rises. It has become common to experience warmer seasons with particularly extreme temperatures during summer. Hence, heat waves have increased worldwide causing heat stress for plants [42]. Heat stress appears with sudden increases in temperature, 10 or 15 °C above usual conditions [43], and its consequence depends on the plant genotype and ecotype, on the level of incremented temperature, and on the length of the stress [44,45]. Plants may survive heat stress through heat-avoidance or heat-tolerance mechanisms [46]. The avoidance processes intend to ensure the survival of a plant, for example altering its leaf orientation or regulating its stomatal conductance, while heat-tolerance mechanisms are related to the plant's ability to maintain its growth under heat stress. These processes involve the synthesis and regulation of different enzymes and proteins [44]. Plants primary sensing mechanism towards heat stress is located in the plasma membrane of cells. These membranes become more fluid and permeable under heat stress, which activates heat sensor proteins. It is believed that these heat sensors are, or interact, with calcium channels [47]. Calcium is known to be a key molecule involved in the activation of diverse stress responses mechanisms [48]. Different transmembrane proteins related to calcium transport have been proposed to act as heat sensors. Members of the Annexin gene family, the protein Synaptotagmin A (SYTA) in *Arabidopsis thaliana* (L.) Heynh. and the Cyclic Nucleotide-Gated Channels (CNGCs) are examples of heat sensor proteins from plants [49–51]. The CNGCs are cation channels that regulate the entrance of ions, e.g., Ca^{2+} , into the cytosol from the apoplast and have a calmodulin-binding domain in their cytosolic region. This suggests that increased levels of cytosolic Ca^{2+} trigger an unknown signaling cascade that mediates the accumulation of heat-shock proteins (HSPs) [47]. In rice, the induced loss of function of two of these CNGCs proteins, OsCNGC14 and OsCNGC16, showed that mutant plants exhibited reduced survival when exposed to both heat and cold stresses. This concurs with the observed role of CNGCs in heat stress signaling and shows that temperature stresses have overlapping signaling mechanisms [52]. The abrupt changes derived from heat stress can degrade cellular components, altering the composition of membranes and denaturing proteins. Moreover, oxidative stress is also a common result of abiotic stresses. In consequence, the production of reactive oxygen species (ROS) increases. ROS can be generated in different cellular compartments, such as peroxisomes, mitochondria, and chloroplasts [53]. These molecules are very toxic and can end up inducing cell death due to damage to proteins, cell membranes, and even DNA [54]. To avoid drastic consequences, cells induce the synthesis of HSPs and heat-shock transcription factors (HSFs). In response to heat stress, these transcription factors bind the heat-shock elements (HSEs) that are conserved regions of the HSPs genes. This leads to increased levels of HSPs in the cells, which aims to preserve the integrity of cell proteins by preventing their misfolding and aggregation thanks to the chaperoning role of HSPs [55]. The overexpression of *Lolium arundinaceum* (Schreb.) Darbysh. HsfA2c produced

plants tolerant to heat stress [56]. In addition, to prevent damage from oxidative stress plants can use different antioxidant enzymes like peroxidase and catalase. The plant species and ecotype determine which enzymes will be responsible for coping with oxidative stress [40]. Importantly, metabolic changes, like alterations in enzymes' activity, also occur due to heat stress. In plants, for example, the oxygenase activity of rubisco rises, leading to more photorespiration and therefore reduced carbon fixation and photosynthesis. Furthermore, heat stress alters the degradation and synthesis of carotenoids and chlorophyll that causes a more pronounced decrease in photosynthetic activity [41].

2.1.2 Low temperature tolerance and winter hardiness

Winter survival of forage grasses is a very complex trait determined by the interaction of abiotic stresses like low temperature, frost, desiccation, water logging, ice-encasement and snow cover, which also can cause biotic stress by low-temperature fungi [57]. Winter hardiness, persistency and stable high yields are limiting factors for forage grass production in temperate regions. Short growing seasons with long days, the long winter with short days and low light intensity cause stressful conditions for perennial plants. Cold acclimation, tolerance to freezing and ice-encasement are crucial components of winter survival. Plant species from temperate climates, which are frequently exposed to sub-zero temperatures have developed advanced mechanisms to cope with extended periods of cold during winters. These plant species, when exposed to low but non-lethal temperatures, increase their freezing tolerance through a process called cold acclimation [58,59]. Most forage grass species and winter-types of cereals need vernalization, i.e., the induction of flowering when exposed to low temperatures [60]. During autumn the plants produce only leaves until the vernalization requirement is met and the tillers switch from vegetative to generative growth. However, stem elongation and flowering need long days and normal growth temperatures and will not happen until spring [61].

Long duration of ice cover (ice-encasement) is the major cause of winter damage [62]. Warm spells in winter cause snowmelt, which then form non-permeable ice layers when the temperature returns to below zero, causing anoxic conditions for plants [63]. Though freezing tolerance gives a good estimate for winter hardiness, the correlation between freezing tolerance and tolerance to ice-encasement is relatively less known [64]. Studies by Gudleifsson et al [65] showed a weak correlation ($r=0.36$) between freezing tolerance and ice-encasement.

Freezing tolerance is a complex dynamic trait which requires a fine-tuned coordinated response at the physiological and sub-cellular level in relation to environmental cues to induce physiological, biochemical, and metabolic changes [66,67]. Many of these resulting cold-associated changes are mainly due to changes in gene expression [66,68,69]. Temperature, light, and a complex interaction of these two variables are key factors driving the process of cold acclimation and determining the extent of freezing tolerance acquired [70–73].

With the increase in autumn temperatures, cold acclimation will occur during late autumn or early winter under different irradiance levels than normal conditions [73,74]. Water logging conditions as a result of the heavy precipitation in autumn during cold acclimation may also negatively affect cold acclimation and freezing tolerance [75]. Winter survival under novel climate conditions is likely to be determined by the ability to cold acclimate at low non-freezing temperatures, resist deacclimation during short warm spells in mid-winters and re-acclimation when the temperatures drop again after the warm spells [76–78].

The ICE-CBF-COR signaling cascade is known to play a key role in freezing tolerance and remains the best-characterized pathway to date [69,79]. CBF regulon consisting of genes *CBF1*, *CBF2* and *CBF3* amongst others, which contributes to acclimation to cold temperatures [80]. These genes were first studied in *Arabidopsis* and encode transcription factors that bind to dehydration responsive genes, as well as those with an early response to cold and dehydration [81]. Other important proteins contributing to winter survival are

dehydrins (DHNs) or group 2 Late Embryogenesis Abundant (LEA) proteins. Many grass species are tolerant to freezing by upregulating DHN genes [82]. Dehydrins are often regulated by CBF cold-responsive pathways. CBF/DREB transcription factors recognize and bind to the DRE/CRT elements in the promoter of cold-responsive genes (COR) [83]. The transcriptome analysis in *Elymus nutans* showed that the genes encoding LEA14-A, cold-regulated plasma membrane protein COR413PM, cold-responsive protein COR14a and dehydrin COR410 (Table 2) had higher transcriptional abundance in a genotype with higher tolerance to cold [84]. Further, quantitative trait loci (QTLs) for winter survival, frost and drought tolerance have been mapped in meadow fescue (*Lolium pratense* (Huds.) Darbysh.). Several of the QTLs were located in the same chromosomal regions as QTLs and genes in Triticeae species, notably DHNs, CBFs and vernalization response genes. The major frost tolerance/winter survival QTL co-located with the position of the CBF6 gene. Some of the winter survival QTLs co-located with frost tolerance QTLs, others with drought QTLs, while some were unique and most likely this was due to segregation for genes affecting seasonal adaptation, e.g., photoperiodic sensitivity [85].

In addition, perennial grass species produce water soluble carbohydrates, such as fructans and raffinose family oligosaccharides during cold acclimation [86]. Fructans are an important energy source found in temperate forage grasses. They are synthesized from sucrose and can be defined as storage carbohydrates that are non-structural [87]. Fructans are stored in vacuoles and will either have linear or branched fructose polymers with glycosidic bonds to sucrose [88]. The linear polyfructose molecules tend to accumulate in plants either as an addition to or instead of starch [89]. The levels of fructan in wintering plants are involved in freezing tolerance and they are important for survival during winter and regeneration or sprouting of tissues in spring, being an important sugar supply [90]. Accumulation of fructans involves fructosyltransferases, invertases and fructan exohydrolases, which are regulated tightly and moreover, their genes have been characterized and isolated [89,91].

2.2. Drought

Drought is one of the main environmental factors limiting crop productivity and predicted climate change shifts in the future will result in temperature increases and changes in precipitation patterns [40]. In the semiarid regions, plants have evolved defense mechanisms allowing them to cope with stressful environments and survive prolonged desiccation. These mechanisms include an elaborated antioxidant defense system and complex gene expression programs, ensuring transcription and translation of LEA proteins, heat shock proteins, and other stress-responsive genes, as well as metabolic modulations consisting of various phytohormones and phytochemicals [92–95]. Annual crops escape the limited water conditions by completing their reproductive cycle producing seeds. While annuals can ensure the survival of species via seeds, perennial crops must cope with water shortage using drought tolerance and avoidance strategies [41,96]. Plants avoid drought by reducing transpiration and maintaining or even increasing water uptake resulting in postponed tissue dehydration. In contrast, drought tolerant perennial crops experiencing stress survive by suspending shoot growth leading to leaf desiccation. However, the crowns of the plants stay vigorous and recover under adequate rainfall. The latter two strategies are of particular importance in forage crops because they are expected to be high yielding under mild stress and to quickly recover after it. Recent studies on vegetative desiccation tolerance have linked this mechanism to seed-development processes, by showing increased expression of seed-related genes in vegetative tissues during drying [97]. The finding suggests that desiccation and water-deficit tolerance mechanisms in grasses derive from an alternative use or “rewiring” of seed-development pathways. Unraveling the key players involved in this mechanism could be a significant step towards engineering the resurrection trait into drought tolerant forage crops.

Compared to semiarid regions, the typical mild summer drought of temperate zones does not threaten crop survival but causes a significant yield penalty [98,99]. The strategies result in reduction of aboveground biomass growth and accumulation, which is one of the most agronomically important traits to achieve. Genotypes adapted to water deficit might maintain growth, and under temporary drought scenario they might be considered as competitive in terms of stable biomass accumulation [100]. The limited water availability triggers responses at the whole-plant, tissue, cellular and molecular levels [35,101]. The perceived stress signal is converted to increased levels of abscisic acid (ABA) production and accumulation in stomata guard cells which regulate transpiration through stomata closure and thus conserve water in tissues [102,103]. However, this type of water loss prevention negatively affects the photosynthetic activity and this in turn results in a slowdown of growth and, under prolonged water shortage, growth halt [101]. Though ABA negatively impacts the aboveground biomass accumulation, at the same time it has an opposite effect on growth and development of roots that largely help to overcome stress [104–106]. Nevertheless, improving forage crops for superior yield through ABA-induced drought adaptation remains a great challenge because of ABA mediated stomatal closure leading to reduced carbon gain and ABA-induced senescence [107]. Another consequence of drought stress in plants is overproduction of ROS causing an oxidative stress which in turn results in cellular membrane damage, imbalance of ions and oxidation of bioactive molecules [108,109].

ABA also plays an important role in inducing the protective role of DHNs. Dehydrins are a subfamily of group 2 LEA proteins that accumulate during late stages of seed development, when plant water content often decreases. In addition, DHNs accumulate in vegetative tissues that are exposed to various stress factors related to dehydration (drought, high salinity, low temperatures, wounding) [110]. Hundreds of DHN genes have been sequenced in both dicotyledonous and monocotyledonous plant species [111]. The regulation of these genes involves Ca^{2+} signaling pathways as well as ABA and mitogen-activated protein kinase (MAPK) cascades. Dehydrins help to detoxify ROS binding to metal ions and scavenging ROS through oxidative modification. Importantly, the characteristic lysine-rich K-segment of dehydrins displays high membrane affinity. DHNs are known to bind and to protect membranes and even DNA from potential damaging caused by adverse environment. It has been shown that DHNs interact with plasma membrane intrinsic proteins that are important members of the aquaporin family [112,113]. The coordination of intracellular functions, including stress response, depends on the flow of information from the nucleus to cell organelles and back. The expression of many nuclear stress response genes is regulated by 3'-phosphoadenosine 5'-phosphate (PAP), known as a key player in chloroplast stress retrograde signaling, which accumulates during drought, salinity and intensive light stress [114]. The concentrations of PAP are regulated by phosphatase SAL1, which dephosphorylates PAP to Adenosine monophosphate (AMP) and thus reduces PAP levels [115]. The studies on *TaSal1* knockout wheat mutants obtained using CRISPR-Cas9 confirmed PAP accumulation, resulting in enhanced stress signaling and induced stomatal closure. Consequently, mutant plants had bent stem and rolled-leaf phenotype with better regulation of stomatal closure and seed germination [116].

Plants' early responses to a deficit of water and to a high content of salt in the soil are very similar. The initial water stress is followed by alterations in photosynthesis and cell growth [117,118].

2.3. Salinity

Salt stress is considered one of the most devastating environmental stresses that limits the productivity and quality of agricultural crops worldwide. Nowadays, over 20% of the world's cultivable lands are affected by salinity stress and these areas are in a continuous increase [119].

During the process of soil salinization, an excessive increase in water-soluble salts occurs. The most common cations found in saline soils are Na^+ , Ca^{2+} , and Mg^{2+} , whereas chloride, sulfates, and carbonates are the main source of anions. The high concentration of dissolved salts in the root zone reduces the osmotic potential difference between the soil and roots, which limits water uptake in plants, causing physiological water deficiency and malabsorption of essential elements [120]. The toxic effect of a high concentration of Na^+ is the most prominent one – Na^+ is not needed for plant metabolism, whereas it competes for binding sites with K^+ that is essential for many cellular functions [121].

In cells, exposition to salt stress primarily induces osmotic stress and ionic stress. Sensing salt ions and hyperosmolality triggers Ca^{2+} accumulation in the cytosol, activation of ROS signaling, and alteration of membrane phospholipid composition. These signals change phytohormone signaling, cytoskeleton dynamics, and the cell wall structure. Moreover, various physiological and molecular changes inhibit photosynthesis and alter sugar signaling, which may lead to plant growth retention [122].

Several Na^+ -binding molecules have been demonstrated to act as sensors able to respond and signal an excess of Na^+ [123]. The best-studied of them is the hyperosmolality-gated calcium-permeable channel family OSCA that has been identified in many species, including important cereals [124,125].

The environment-triggered Ca^{2+} influx signal in the cytoplasm is received by Ca^{2+} -sensing proteins. Among those, calcineurin B-like proteins (CBLs) are responsible for maintaining the ion transport and homeostasis through interactions with the serine/threonine protein kinases (CIPKs) which activate Na^+ , K^+ , H^+ , NO_3^- , NH_4^+ and Mg^{2+} transporters located in different cellular membranes. In addition, regulation of ROS and ABA signaling is also modulated by CBL-CIPK complexes [118]. Regulation of Na^+ transport from cytosol to the apoplast is mediated by the salt overly sensitive (SOS) pathway where the specific complexes of CBLs-CIPKs interact with Na^+/H^+ antiporter SOS1 that removes excessive Na^+ . Another CBL-CIPK complex activates Na^+/H^+ exchange transporter 1 located in the vacuole tonoplast to transport the excess of Na^+ to that organelle [118]. The CBL and CIPK encoding genes seem to be conserved among dicots and monocots [126,127]. Sequestering of the ions into vacuoles helps to avoid stress but needs the osmotic potential adjustment in the cytosol by the accumulation of osmotically active substances such as polyols, amides and amino acids, soluble carbohydrates, and quaternary ammonium compounds. The toxic and osmotic effects of salt ions in the cytoplasm are usually reached by scavenging ROS by antioxidant enzymes that also help to tolerate the toxic effects of salt ions [128].

Other early events in salt stress response include rise of cyclic nucleotides (e.g., cGMP) and ROS. The cGMP inhibits Na^+ influx via non-selective ion channel. In addition, rise in cGMP and ROS induces transcriptional regulation that can activate MAPK cascades. Rise in expression of MAPKs leads to increased osmolyte synthesis to alleviate salt-induced osmotic stress. Osmolytes are also a signal for production of ABA, regulating stomatal closure and therefore osmotic homeostasis and water balance [122]. Salt stress-induced accumulation of ABA activates the sucrose non-fermenting-1 related protein kinases 2 (SnRK2s). In turn, activated MAPKs and SnRK2s transduce signals to downstream transcription factors to induce the expression of stress-responsive genes [129].

The ability to resist saline environments differs remarkably among plants. Non-halophytic plants (i.e., glycophytes) are sensitive to salinity stress, and their growth and development are hampered by a salinized environment. However, glycophytes exhibit natural variation in their salinity tolerance. Such variation often relies on an allelic variation of genes involved in salinity stress response [130]. For example, it has been noticed that under salt treatment to reduce sodium influx in response to osmotic stress, an aquaporin, a cation antiporter, and a calcium-transporting ATPase were downregulated, while a manganese transporter and a vacuolar-type proton ATPase subunit were upregulated in the roots of a salt-tolerant accession of *Poa pratensis* L. when compared to a susceptible accession of *P. pratensis* [131].

Halophytic plants have adapted to salinized environments and they show stimulation of growth enhancement and productivity at moderate salinity (50–250 mM NaCl) [128]. These plant species have developed specific mechanisms that regulate internal salt load, e.g., many have developed specialized salt glands which excrete ions on the leaf surface. Such structures are characteristic of C4 grasses belonging to the tribes Chlorideae, Sporoboleae and Aelropodeae. Other halophytes, including as well C4 grasses (e.g., *Paspalum vaginatum* Sw.), use bladder-like protrusions from epidermal cells into which ions are sequestered and accumulated until these cells senesce and die [132,133]. The number and density of salt glands or salt bladders depends on salt concentration in the soil during plant growth indicating the dynamic adaptation to environmental conditions [128].

Identification of genetic components and their variance underlying salinity tolerance is a useful source for plant breeders [134]. The overexpression of several halophytic genes in glycophytic recipients has been demonstrated to enhance abiotic stress tolerance [135]. An increasing number of transcriptomic studies from salt-tolerant non-halophytic and halophytic grasses grown under different salinity conditions will help to elucidate the gene networking process behind the effective salinity response [136–138].

3. Gene editing: a tool for developing stress resistant forage grasses

The biggest challenge for agriculture nowadays is to obtain plants that are resilient to adverse environmental conditions that at the same time provide enough yield to fulfill food and feed security in a sustainable way. In the case of perennial forage grasses, yield is determined by repeated harvesting of herbage over as many years as possible. Therefore, forage grass genotypes with improved survival and growth under abiotic stress conditions are needed.

Genome editing tools have proven to be useful for achieving such aims, especially the Nobel prize-winning discovery of application of RNA-directed Cas9 nuclease for genome editing [139,140] abbreviated as CRISPR-Cas9. Although this editing strategy was immediately applied in model and crop plants, almost ten years ago [141–145], not much has been achieved in the forage grasses landscape. The European GMO database EU-GENIUS lists only green foxtail (*Setaria viridis* (L.) P. Beauv.) line 193-31 that has been modified using CRISPR-Cas9 mediated mutagenesis. The expressed CRISPR-Cas9 system targeted the coding region of the *S. viridis* homolog of the *Zea mays* L. *ID1* gene, which promotes flowering in maize. The deactivation of the homolog in *S. viridis* led to delayed flowering. In the knockout line 193-31, the CRISPR-Cas9 DNA construct was segregated away [146].

To find out how many publications have been released showing edited genes in forage grasses, a search was carried out in the following databases: Scopus, Web of Science, Google scholar and PubMed. The search included the scientific or the common names of 47 grass species (Table S1) or the name of each of the 12 subfamilies of Poaceae and in addition, one of the following terms: "CRISPR", "gene editing", "genome editing". The outcome of the search is shown in Table 1. The genome of only six species, three annual grasses and three perennial ones, all growing in temperate regions, has been targeted with CRISPR-Cas tools. Genome editing in *S. viridis*, a model plant for C4 grasses, has been reported three times. Most of the work has been done knocking out one gene using the easiest genome editing approach, i.e., CRISPR-Cas9.

362
363
364
365
366
367
368
369
370
371

372
373
374
375
376
377

378
379
380
381
382
383
384

385
386
387
388
389
390
391
392

393
394
395
396
397
398
399
400
401
402
403
404

405
406
407
408
409
410
411
412
413

Table 1. Gene editing in forage grasses.

414

Species	Common name	Biome	Life cycle	Editing system	Publication
<i>Lolium multiflorum</i>	Italian ryegrass	Temperate	Annual	CRISPR-Cas9	[24]
<i>Lolium perenne</i>	Perennial ryegrass	Temperate	Perennial	CRISPR-Cas9	[24,28]
<i>Panicum virgatum</i>	Switchgrass	Temperate	Perennial	CRISPR-Cas9	[22]
<i>Lolium arundinaceum</i> *	Tall fescue	Temperate	Perennial	CRISPR-Cas9/Cas12a	[147]
<i>Setaria italica</i>	Foxtail millet	Temperate	Annual	CRISPR-Cas9	[25,29,148]
<i>Setaria viridis</i>	Green foxtail	Temperate	Annual	CRISPR-Cas9_Trex2	[23]
				CRISPR-Cas9	[27,149]

* *Festuca arundinacea*.

415

CRISPR-Cas9 as a system for carrying out simple mutations (indels: insertions/deletions) that change the reading frame of a coding region and therefore generate knockouts, is straightforward and still mainly used for functional genomics. It consists of two main components: the Cas9 nuclease from *Streptococcus pyogenes* and the short guide RNA (gRNA) that targets the DNA sequence of interest. Designing the gRNA with precision enables the simultaneous mutations of all alleles of a gene in a polyploid plant, as it was the case for *Panicum virgatum* L. (tetraploid) and *Lolium arundinaceum* (allohexaploid, Table 1). Specific genes that have been knocked-out in forage grasses are related to flowering (*PHYC* of *Setaria italica* (L.) P.Beauv. and *FON2* of *S. viridis*), tillering and branching (*tb1a* and *tb1b* of *Panicum virgatum*), meiosis (*DMC1* of *Lolium multiflorum* Lam.), haploid induction (*MTL* of *S. italica*) and heat stress response (*HSP17.9* of *L. arundinaceum*), apart from the *PDS* gene used as endogenous marker (Table 1 and references therein). In most of the cases the cited publications discuss the targeted mutagenesis method and results obtained, but the phenotypic characterization of the mutants is limited and far away from field trials. Interestingly, not only classical CRISPR-Cas9 system has been used, but also CRISPR-Cas12a in the case of *L. arundinaceum* [26] and CRISPR-Cas9_Trex2 in the case of *S. viridis* [23].

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

The toolkit of CRISPR-Cas applications has expanded to around twenty different techniques that allow diverse targeted modifications in the genome [36,150]. On one hand, Cas enzymes from different bacteria have been characterized and adopted for use. That is the case for Cas12a (former Cpf1), an enzyme from the *Lachnospiraceae bacterium* ND2006 that cuts DNA strands distal from the sequence recognized by the nuclease (the PAM site), generating 4-5 nucleotide overhangs that enable an easy insertion of donor DNA sequences [151,152]. Other modifications of the CRISPR-Cas9 system imply the co-expression or the fusion of different proteins to the Cas9 nuclease, in its original or mutated versions. CRISPR-Cas9_Trex2, for example, has the Trex2 exonuclease co-expressed with Cas9 for increasing the mutation efficiency [23,153]. Importantly, an enzymatically inactive variant of Cas9, called "dead Cas9" (dCas9) that maintains its specific DNA binding ability, can be fused to transcription activators or repressors to regulate transcriptional levels of endogenous genes [154]. Therefore, CRISPR-Cas tools are not only meant to inactivate genes and create loss-of-function mutants, but also gain-of-function mutants can be obtained. In addition, thanks to the Super Nova Tag (SunTag) system, the transcriptional regulation can be potentiated. The SunTag contains peptide repeats that bind several transcription factors for cooperatively activating a target gene [155]. Moreover, a gene of interest may also be up- or downregulated epigenetically. For instance, CRISPR-dCas9 linked to DRM methyltransferase catalytic domain targets methylation to specific loci and thereby inactivates the target gene [156].

453

454

455

456

457

An alternative way of inducing a change in the levels of expression of a gene is altering its promoter sequence. In fact, the promoter can be even swapped by another one that ensures e.g., higher levels of expression in a ubiquitous manner. Using CRISPR-Cas9 such a substitution is possible, as shown for the *ARGOS8* gene in maize, whose overexpression was associated with improved grain yield under field drought stress conditions [157].

It should be pointed out that yield and stress resistances are among the most difficult polygenic traits to improve through genetic engineering, but examples as the former one give hope that it can be achieved by CRISPR-Cas. Another example is the knockout via CRISPR-Cas9 of the main effect gene *OsRR22* that controls salt tolerance in rice. Obtained plants showed salt tolerance in growth chambers and no difference in agronomic traits compared to wild type plants in field trials under normal growth conditions [158,159].

As explained in section 2, abiotic stress responses are complex, linked to different metabolic pathways and the genes involved in those mechanisms are mainly pleiotropic. Fishing out a specific key player, a master gene to be mutated, could be possible in some cases and it is worth trying. Since gene editing in grasses is in its early stages (Table 1), we selected specific genes related to the four abiotic stresses discussed in this review and figured out if those target genes would need to be overexpressed or downregulated to gain tolerance to specific stresses. If a candidate gene was found in forage grasses or at least in a Poaceae species, that species was selected, but this was not possible in all cases (Table 2). As shown in the table, there are genes that are related to more than one stress response. For simplicity, it is not shown that, e.g., *DHN11* seems to be also involved in cold and drought stresses and *COR410* appears to be related to drought stress as well.

Table 2. Target genes for improvement of abiotic stress tolerance.

Abiotic stress	Target gene	Species	Role	Proposed Strategy	Publication
Heat	<i>OspsbA</i>	<i>Oryza sativa</i>	Stress response	Upregulate	[160]
	<i>LaHsfA2c</i>	<i>Lolium arundinaceum</i> *	Stress response	Upregulate	[56]
	<i>OsCNGC14/CNGC16</i>	<i>Oryza sativa</i>	Stress sensing	Upregulate	[52]
	<i>SiMAPK3</i>	<i>Solanum lycopersicum</i>	Stress response	Downregulate	[161]
	<i>OsPYL1/4/6</i>	<i>Oryza sativa</i>	Stress response	Downregulate	[162]
	<i>SIPHYA</i> and <i>SIPHYB1B2</i>	<i>Solanum lycopersicum</i>	Stress response	Downregulate	[163]
Cold	<i>EnCOR410</i>	<i>Elymus nutans</i>	Stress response	Upregulate	[84]
	<i>AcSnRK2.11</i>	<i>Agropyron cristatum</i>	Stress response	Upregulate	[164]
	<i>OsCOLD1</i>	<i>Oryza sativa</i>	Stress sensing	Upregulate	[165]
	<i>OsMYB30</i>	<i>Oryza sativa</i>	Stress response	Downregulate	[166]
	<i>AtEGR2</i>	<i>Arabidopsis thaliana</i>	Stress response	Downregulate	[167]
	<i>AtCRPK1</i>	<i>Arabidopsis thaliana</i>	Stress response	Downregulate	[168]
Drought	<i>CdDHN4</i>	<i>Cynodon dactylon</i>	Stress response	Upregulate	[169]
	<i>OsSYT-5</i>	<i>Oryza sativa</i>	Stress sensing	Upregulate	[170]
	<i>AcSnRK2.11</i>	<i>Agropyron cristatum</i>	Stress response	Upregulate	[164]
	<i>OsDST</i>	<i>Oryza sativa</i>	Stress response	Downregulate	[171]
	<i>TaSal1</i>	<i>Triticum aestivum</i>	Stress response	Downregulate	[116]
	<i>HvCBP20</i>	<i>Hordeum vulgare</i>	Stress response	Downregulate	[172]
Salinity	<i>ZmDHN11</i>	<i>Zea mays</i>	Stress response	Upregulate	[173]
	<i>AcSnRK2.11</i>	<i>Agropyron cristatum</i>	Stress response	Upregulate	[164]
	<i>OsOSCA1.4</i>	<i>Oryza sativa</i>	Stress sensing	Upregulate	[134]
	<i>OsbHLH024</i>	<i>Oryza sativa</i>	Stress response	Downregulate	[174]
	<i>HvITPK1</i>	<i>Hordeum vulgare</i>	Stress response	Downregulate	[175]
	<i>OsRR22</i>	<i>Oryza sativa</i>	Stress response	Downregulate	[158]

* *Festuca arundinacea*.

Section 2 mentioned that plants detect an increase in temperature (in the soil or air) when the structure and fluidity of their cell membranes change. Heat stress tends to make membranes more fluid [176], which activates pathways through heat sensors like the CNGCs. In theory, an increased expression of stress receptors can lead to an improved response to stress. Consequently, the genes involved in the heat stress response signalling pathway can be upregulated by overexpressing a heat sensor coding gene. In *A. thaliana*,

458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474

475
476
477
478
479
480
481
482

an overexpression of the SYTA gene resulted in higher germination and seedlings survival rates than in wild-type and knockout lines after heat stress exposition. Moreover, the overexpression plants presented higher expression of both HSPs and HSFs, together with lower levels of membrane lipid peroxidation than in non-overexpression lines [50]. All these changes provide evidence that upregulating a heat stress sensor can improve the stress tolerance of a plant. Therefore, overexpressing a similar gene in grasses, like a homologous of rice *OsCNGC14* or *OsCNGC16* gene, could result in forage species with higher tolerance to heat stress. A similar approach can be followed by upregulating proteins present in plants in a basal state that are involved in the responses to abiotic pressures (Figure 1). Kinase proteins are suitable for this goal since they are involved in most stress response pathways, regulating posttranslational modifications of other proteins as a response to both abiotic and biotic stress [177]. Therefore, overexpressing a gene from the SnRK2 family, a group of kinases specific to plants that have been shown to play important roles in abiotic stress regulation is an adequate approach [178]. The heterologous overexpression of the gene *TaSnRK2.3* from wheat in *Arabidopsis* produced plants that had higher tolerance to drought conditions [179]. Similarly, another study was able to overexpress the *AcSnRK2.11* gene from *Agropyron cristatum* (L.) Gaertn., a forage grass species, in *Nicotiana tabacum* L. The overexpression plants had significantly higher survival rates than the wild-type ones after recovery periods from cold stress and presented significantly upregulated patterns of abiotic stress-related genes like dehydrins. Possibly, upregulation of these protein kinases could provide drought, cold and salinity stress tolerance to forage grasses plants.

On the other hand, negative regulators of abiotic stress responses are also suitable targets for abiotic stress tolerance improvement by downregulating them via genome editing (Figure 1). Possible candidates for downregulation could be enzymes that degrade signaling molecules involved in stress response, like for example the inositol phosphatases [180]. As previously mentioned in this review, the phosphatase SAL1 negatively regulates plants' response to drought [181]. Using the CRISPR-Cas9 system, scientists have already generated *Tasal1* knockout mutant wheat with fewer and smaller stomata, that germinate and grow better under drought conditions [116]. Likewise, modifying the expression of transcription factors related to abiotic stress is another alternative for producing tolerant plants. The transcription factors of the bHLH family have been shown to participate in abiotic stress regulation in different plant species [182]. In rice, the *OsbHLH024* gene seems to negatively regulate salinity tolerance. This was demonstrated by generating knockout plants using the CRISPR-Cas9 system. The mutated plants had an increased salinity tolerance when compared to the wild-type ones. Additionally, the knockout lines presented a reduced accumulation of sodium ions and ROS, but higher concentrations of potassium ions than the control plants. Finally, the expression of genes encoding ion transporter was upregulated in the knockout plants in comparison to the wild-type ones [174]. All these variations suggest that the downregulation of homologues of the *OsbHLH024* gene in grasses could provide them with salinity stress tolerance.

During the last years, innovative ways of inserting specific targeted mutations based on CRISPR-Cas have been developed, e.g., base- and prime editing and for now, some technical problems need to be overcome when applied to plants [183–185]. In these cases, the Cas9 nuclease is mutated in such a way that it acts as nickase, cutting only one strand of the targeted DNA. These strategies and the activation of homology- directed repair (HDR) instead of Non-Homologous End Joining (NHEJ), makes it possible to replace, insert or delete large sequences and even to generate chromosomal rearrangements [186,187].

The possibilities to induce targeted changes with CRISPR-Cas in the genome of crops, and specifically in forage grasses, are immense, not to mention the speed of obtaining the desired traits compared to conventional breeding techniques. In addition, gene editing can be easily multiplexed for targeting different sequences at one shot. Depending on the specific trait and species, there can be bottlenecks to be removed like specific ways of


transforming a plant or quality in the annotation of the genome. These obstacles are thought to be solved with technical advances, however in the case of grasses, important biological features need to be taken into consideration when aiming to combine gene editing with a breeding program. These challenges are elaborated in section 5. Here we briefly mention that also biological characteristics of grasses related to the way they reproduce could be changed with genome editing.

Forage grasses have a strong gametophytic self-incompatibility (SI) system that makes inbreeding almost impossible. The two multi-allelic *S* and *Z* genes have since long been known to govern SI in grasses [188,189], and recently it was shown that two *DUF247* genes are behind the *S* and *Z* loci [190,191]. With the sequences and molecular function of these genes known, they would be an obvious target for generating self-fertile knockout lines by gene editing. A similar approach has been used to develop self-compatibility in potato [192].

To obtain male sterile lines is also of importance in the case of forage grasses. The way has been paved by research in maize, where genes *Ms1* and *Ms45* have been targeted by CRISPR-Cas9 and male-sterile wheat lines for hybrid seed production have been obtained [193,194].

Also, double haploids can be generated artificially inducing haploids with a knock-out of *MTL*, as it has been done already in *S. italica* (Table 1) [25].

Finally, apomixis is present in several grass species, e.g., *Poa pratensis*, a species used both in lawns, pastures, and leys. Inducing apomixis in other forage grasses would be of importance for fixing hybrid vigor. Some steps towards achieving this aim have been taken already in rice. Mutations using CRISPR-Cas of several genes related to the abolishment of meiotic steps produced clonal diploid gametes. Then, parthenogenesis was induced by ectopic expression in the egg cell of *BABY BOOM1* and clonal progeny was obtained [184,195].



537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562

Figure 1. Proposed strategy for the improvement of abiotic stress tolerance in forage grasses using genome editing. Four abiotic stresses (heat, low temperature, drought, salinity) hinder the overall wellbeing of a non-tolerant grass (plant shown in yellow). Using the CRISPR-Cas system, different genes can be targeted. Agrobacterium-mediated transformation or biolistics are suitable delivery methods of the CRISPR-Cas+gRNAs complex for in-vitro culture modifications that lead towards the generation of abiotic stress tolerant plant (blue rectangle). Once tolerant parental plants are obtained (GE: gene editing), these can be crossed to produce a population able to overcome the effects of abiotic stress (green rectangle). The green plant on the bottom left represents a tolerant grass. 564
565
566
567
568
569
570
571

4. Gene editing versus traditional genetic modifications

 572

Genetic variation is fundamental to crop improvement. Modern plant breeding started in the late 19th century with the advent of cross-breeding which still is the backbone of most plant breeding efforts [196,197]. After the discovery that physical and chemical factors can lead to heritable changes in genetic material, random mutagenesis became a valuable tool for plant breeding to increase genetic diversity and to develop specific traits. With the discovery of recombinant DNA technology in the 1970s, the development of new combinations of genetic elements by splicing genes and regulatory elements from different species became possible. The discovery of *Agrobacterium*-mediated transformation enabled scientists to introduce these novel combinations of genes into plant genomes to produce new traits [197]. While the introduction of transgenes into plant genomes has contributed enormously to the understanding of gene functions in plants, the commercial applications have been limited to mostly herbicide tolerance and insect resistance, which provide obvious advantages for farmers, but little direct, tangible benefits for consumers in developed countries. Only a few commercial applications of transgenic plants with improved yield and abiotic stress resistance are known. Wheat expressing the sunflower transcription factor HaHB4 has been shown to provide improved water use efficiency resulting in higher grain production [198]. Wheat HB4 marketed by the company Bioceres Crop Solutions has been authorized for food and feed uses in a number of countries, such as Argentina, Australia, Brazil and United States, but its cultivation is approved only in Argentina [199,200]. Maize MON87403 contains the *ATHB17* gene from *A. thaliana* encoding a transcription factor of the HD-Zip II family with reported increase in ear biomass at the early reproductive phase [201], which may provide an opportunity for increased grain yield under field conditions [202]. Maize MON87460 expresses the *Bacillus subtilis* cold shock protein B (CspB) resulting in increased grain yield under drought conditions [203]. Both GMO events have been assessed by the European Food Safety Authority (EFSA) [204,205], and MON87460 was authorized for food and feed uses in the EU. Transformation techniques have been developed for most of the economically important forage and turf grass [206], however, very few transgenic forage grasses have been registered for commercial cultivation. The ISAAA GMO approval database lists only one transgenic event in creeping bentgrass (*Agrostis stolonifera* L.) with tolerance to glyphosate (ASR368) [207]. 573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603

Even though commercial cultivation of GM crops has brought clear benefits to farmers and more indirect benefits to environment through reduced land and pesticides [208,209], cultivation and use of transgenic plants for food and feed have been controversial in many regions of the world, and especially in Europe. Agronomic, environmental, human health, social and economic effects of transgenic crops have been comprehensively reviewed by the US National Academies of Sciences in 2016 [210]. 604
605
606
607
608
609

Genome editing became possible with advances in protein engineering which allowed production of site-directed nucleases (SDNs), such as zinc finger nucleases (ZFNs) and transcription activator-like effector nucleases (TALENs) [211,212]. As outlined in section 3, genome editing has several advantages over the transgenic techniques including precision, lower number of off-target effects, more streamlined production, multiplex possibility, as well as potential for modification of many more different traits. A few examples include lower gluten content in wheat through simultaneous editing of alpha-gliadin 610
611
612
613
614
615
616

genes [213], increased production of gamma-aminobutyric acid in tomato [214] or increased accumulation of provitamin D3 in tomato [215]. The maize with an increased expression of *ARGOS8* gene, as detailed in section 3, contained no exogenous DNA sequences, thus, theoretically, it could be exempt from GMO regulation depending on country-specific policies.

The increased precision, low off-target potential and the absence of exogenous DNA in some of the genome-edited plants suggested that genome editing would not be regulated similarly to GMOs. It certainly has worked out that way in Japan, where Sanatech Seed has commercialized high gamma-aminobutyric acid tomato [216]. In the EU, however, the Court of Justice of the European Union (CJEU, case C-528/16) ruled that organisms resulting from mutagenesis techniques in legal aspects are GMOs and are subject to the regulations laid down by the Directive 2001/18/EC. This applies to mutagenesis techniques introduced since 2001, when the GMO Directive was adopted. Site-directed nucleases can modify plant genomes according to three scenarios, SDN-1, SDN-2 and SDN-3 [217], where only SDN-3 scenario results in transgenic plants, while under SDN-1 and SDN-2 scenarios no exogenous DNA is inserted into the genome. Under the CJEU ruling, also the SDN-1 and SDN-2 techniques, including CRISPR-Cas fall under the GMO Directive, while chemical and radiation random mutagenesis remains exempt according to Annex IB of the Directive 2001/18/EC. The ruling provoked a public outcry from both academia and biotech industry, which stressed that from a scientific point of view the application of GMO Directive to products created by a much more precise technique than random mutagenesis and transgenesis results in a disproportionate regulatory burden [218–222]. It was also noted that this ruling leads to a situation when two identical products with the same mutation resulting in, e.g., herbicide tolerance trait could be regulated in totally different ways. In addition, it would create an unsustainable situation with detection, since no technology can determine the origin of simple mutations, such as single nucleotide polymorphisms. Consequently, reliable detection methods for SDN-1 and SDN-2 products are problematic [223]. This legal uncertainty makes genome-editing research in the EU less appealing, as seeking regulatory approval for gene-edited products would involve the same cumbersome procedure as for GMOs. So far there are no applications for regulatory approval involving gene-editing, although a few applications for authorization of products obtained with CRISPR-Cas9 in SDN-3 scenario, e.g., maize DP-915635-4 have been submitted to member states and are currently under review by EFSA [224].

According to the EU (Council Decision (EU) 2019/1904), the European Commission (EC) conducted a study involving input from the Member States and different stakeholders regarding the status of new genomic techniques (NGTs) including genome editing. Within this framework, the EC mandated EFSA to issue a scientific opinion on the risk assessment of plants produced by the SDN-1, SDN-2, and oligonucleotide-directed mutagenesis techniques. EFSA has assessed the safety of plants developed using SDN-1 and SDN-2 techniques and did not identify new hazards specifically linked to these techniques compared to both SDN-3 and conventional breeding. In addition, EFSA concluded that the existing Guidance for risk assessment of food and feed from GM plants and the Guidance on the environmental risk assessment of genetically modified plants are sufficient, but only partially applicable, to plants generated via SDN-1 and SDN-2 [225,226]. As part of the ongoing effort to update the EU GMO legislation upon EC request, EFSA recently produced an updated scientific opinion on cisgenesis and intragenesis [227]. The EFSA scientific opinion concluded that no new risks were identified in cisgenic and intragenic plants obtained with NGTs, as compared with those already considered for plants obtained with conventional breeding and established genomic techniques, although only limited information on such plants was available. EFSA determined that the use of NGTs reduces the risks associated with potential unintended modifications of the host genome resulting in fewer requirements for the assessment of cisgenic and intragenic plants, due

to site-specific integration of the added genetic material. However, there was no legal necessity to overhaul the GMO legislation, since the EFSA concluded that the current guidelines were partially applicable and sufficient. Importantly, the data requirements could be reduced on a case-by-case basis for the risk assessment of cisgenic or intragenic plants obtained through NGTs. While cisgenesis and intragenesis is just one of the possible approaches for forage grass breeding, EFSA also recently issued a statement on criteria for risk assessment of plants produced by targeted mutagenesis, cisgenesis and intragenesis [228]. These criteria could be used by policy makers to design a more flexible and proportionate risk assessment framework for gene edited plants. Recently, several regulatory options have been proposed [229–231]. They range from maintaining the *status quo* (full risk assessment of genome edited organisms as GMOs) to product-based regulation or regulation based on the presence of foreign DNA in the genome. These two options would be preferable for commercial deployment of genome edited crops, but they would require substantial reexamination of GMO Directive and authorization procedure. The EC is expected to present a new policy and/or legal proposal by the second quarter of 2023. Meanwhile, other jurisdictions around the world have already developed legal framework for genome edited plants, e.g., under Argentina NBT Resolution N° 21/2021, if a product (plant, animal or microorganism) does not have a new combination of genetic material, the product is non-GM and considered as conventional product [232]. Different regulatory approaches are summarized in a recent review [233].

Interestingly, the “EU GMO database of Deliberate Release into the environment of plants GMOs for any other purposes than placing on the market (experimental releases)” lists over 900 applications for field trials registered by the Member States since 2002 [149]. Among those there is only one application for field trial of high fructan transgenic ryegrass in 2006, and there are no applications for field trials of genome edited forage grasses, although at least 14 field trials of plants edited with CRISPR-Cas9 have been authorized [234].

In conclusion, while there are a few basic studies on gene function in forage grasses using genome editing technique as described in section 3 of this review, these are yet to see commercial application. The main limiting factor for the investment in research and development of genome edited forage grasses is probably the regulatory uncertainty, especially in the EU. Although edited plants without foreign DNA in the genome are expected to receive the least amount of regulatory scrutiny, they are also less prone to show major changes in relevant traits. This is because gene knockouts or simple gene edits are unlikely to result in complex phenotypes, such as enhanced abiotic stress tolerance, higher yield or improved nutritional composition, especially considering the genetic complexity that has hindered progress in characterization of the genes underlying such traits in forage grasses. Nevertheless, as recent years have witnessed a dynamic development of gene editing tools and genotype-independent transformation approaches along with increasing genomic resources, the manipulation of plant responses may become possible to overcome abiotic stresses when combining modern techniques and good breeding management strategies.

5. Breeding grasses in the gene editing era

Forage grasses are outbreeding species and highly heterozygous due to the strong gametophytic self-incompatibility (SI) system. Inbred line development is thus very difficult with strong inbreeding depression as a result. Therefore, cultivars of forage grasses are usually synthetic populations [235]. Forage grass breeders usually start by phenotypic selection of superior candidate genotypes for traits with high heritability, e.g., heading date and disease resistance, among a large number of spaced plants. However, forage grasses are sown in swards and because yield and other traits will be affected by competition in the swards, such traits cannot be selected on single spaced plants. The candidate genotypes are therefore put in some form of progeny testing system, e.g., polycross to produce half-sib (HS) families or bi-parental crosses producing full-sib (FS) families, and

selection for yield and forage quality traits are based the performance of such families in swards (genotypic selection). Synthetic populations/cultivars are constructed by crossing the best genotypes based on their performance in the progeny test or by mixing HS or FS families. The synthetic populations are further multiplied to obtain enough seed for establishing sward plots for testing in multi-location-year trials before the best candidate cultivars are being submitted to official variety testing. A typical breeding cycle will take 10-15 years before synthetic cultivars are available for farmers. With the advent of high-throughput molecular markers, whole-genome sequences, and genomic selection methods, the breeding cycle can be shortened [236,237].

The success of a breeding program is very much dependent on the genetic variation present in the initial breeding material. Many agronomically important traits, like yield and adaptability to biotic and abiotic stresses, have been partly fixed within elite germplasm, however, they still exhibit large genetic variation and are thus of primary importance in breeding programs [238,239]. This variation might be employed for future improvements of crop productivity and tolerance to stress; however, landraces, closely related species and wild relatives can offer much wider and unexploited germplasm resources [240,241]. Extensive studies of perennial ryegrass diversity among modern European cultivars revealed that modern cultivars are mostly related to ecotypes from north-western Europe [242], while most of the natural genetic variation remains unexploited. Later studies on the genetic structure of geographically diverse perennial ryegrass collection supported these findings and in addition showed that latitude was a prominent force shaping the diversity of wild-growing perennial ryegrass populations [100]. Furthermore, the ecotypes exhibit biomass and seed yielding potential similar to cultivars [243,244], suggesting that ecotypes could serve as valuable trait donors in breeding programs. Field testing of many *L. perenne* ecotypes and cultivars at several Nordic and Baltic locations identified tetraploid Baltic breeding lines and diploid ecotypes from Eastern Europe as being most winter hardy with stable performances across environments [245]. None of the cultivars were among the most stable entries, and diploid ecotypes displayed a larger variation in heading date, regrowth, and winter survival than the cultivars. Thus, there is ample genetic variation still to be exploited within the genetic resources of perennial ryegrass. Induced polyploidization is also widely exploited in forage crop breeding as one of unconventional techniques to develop new superior yielding and abiotic stress tolerant breeding material [246–248].

To utilize transgenes or gene-edits in grass breeding, first, efficient methods for introduction and regeneration *in vitro* need to be available in a range of independent genotypes. In principle, introgression of new genes can either be introduced into the parental clones of already existing varieties (variety-parent approach) or transferred into a new base population (population approach) [235]. Repeated backcrossing and an efficient selection system is needed to bring transgenes/gene-edits to homozygosity in the parental clones. A side-effect of this could be increased inbreeding depression due to linkage drags creating longer homozygous chromosomal segments. Traditional random insertion of transgenes in several genotypes that are intercrossed to construct synthetic cultivars is problematic due to the presence of multiple insertion sites, silencing and variable expression levels. The availability of complete genome sequences also of forage grass species, notably *L. perenne* [249], and gene editing technologies, makes it possible to do induce precise genome alterations. This will make it easier to develop synthetic cultivars of outbreeding crops like forage grasses with stable expression of genetic modifications.

Integration of transgenic traits in perennial grasses and the challenges associated with deployment and management of transgenic cultivars has been discussed by Smith et al. [250] and Badenhorst et al. [251]. Using gene-drive technologies [252] would in principle be an efficient method for spreading gene-edits through breeding populations of grasses. However, the risk of gene flow between cultivars and to feral populations is high and would probably preclude practical use of such technologies.

A pertinent question is what the most important targets for genetic engineering in forage grasses would be. Genetic gain for yield has been modest due to the long breeding cycles and extensive field testing [253,254]. The potential heterosis is only partially exploited in synthetic cultivars, and it is expected that great yield increased could be achieved if F1 hybrids, which has been very successfully exploited in maize, could be developed [255]. Self-incompatibility, inbreeding depression, and the lack of male-sterile lines for making hybrids are major obstacles for developing F1 hybrids. Inbreeding depression needs to be tackled to implement self-fertile lines in forage breeding programs. By generating a large number of self-fertile plants with diverse genetic backgrounds by gene-editing, and selecting genotypes with good seed set, the prospects of developing inbred lines in forage grasses have never been better. These lines could be used for F1 hybrid production and would also be very useful for functional studies. Other methods for capturing heterosis would be the development of facultative apomixis. The evolution of apomixis in natural populations and the challenges of utilizing apomixis in breeding has been reviewed recently by Hojsgaard and Hörndl [256].

6. Conclusions

In the current review, we focus on possible improvements of abiotic stress tolerance in forage grasses using new genome editing tools. The potential impact of climate change is described in relation to forage grass tolerance to four important abiotic stresses, heat, low temperature, drought and salinity. We propose approaches for editing the genome of grasses to regulate stress responses. Furthermore, we discuss the latest developments in the regulatory framework for genome editing, especially with regard to the EU, and identify factors affecting the application of genome editing techniques for the improvement of grasses. Finally, we address breeding strategies specific to the reproductive biology of forage grasses and identify how genome editing could be used to facilitate breeding and achieve food security in a sustainable way. In conclusion, we describe pathways for developing abiotic stress tolerance in forage grasses under climate change using genome editing technologies, provided that an appropriate legal framework is developed.

Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1, Table S1: Forage grasses.

Author Contributions: Conceptualization, F. S., C. S.; writing—manuscript preparation, F. S., N. R., K. J., M. S., O.A. R., M. R., G. S. and C. S.; writing—review and editing, F. S. and C.S.; funding acquisition, K. J., N. R., O.A. R. and C.S. All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Funding: Project "Improving adaptability and resilience of perennial ryegrass for safe and sustainable food systems through CRISPR-Cas9 technology – EditGrass4Food", ID No. EEA-RESEARCH-64, Contract No. EEZ/BPP/VIAA/2021/4 is financially supported by the European Economic Area (EEA) grants.

Acknowledgments: The authors are thankful to students and technicians who have contributed to the EditGrass4Food project. Figure 1 was created with BioRender.com.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the writing of the manuscript.

References

776
777
778
779
780
781
782
783
784
785
786
787
788
789
790

791
792
793
794
795
796
797
798
799
800
801
802
803

804
805
806
807
808
809

810
811
812
813
814
815
816

817
818
819
820

821

1. Huang, W.; Zhang, L.; Columbus, J.T.; Hu, Y.; Zhao, Y.; Tang, L.; Guo, Z.; Chen, W.; McKain, M.; Bartlett, M.; et al. A Well-Supported Nuclear Phylogeny of Poaceae and Implications for the Evolution of C4 Photosynthesis. *Mol. Plant* **2022**, *15*, 755–777, doi:10.1016/j.molp.2022.01.015. 822
2. Lee, S.; Choi, S.; Jeon, D.; Kang, Y.; Kim, C. Evolutionary Impact of Whole Genome Duplication in Poaceae Family. *J. Crop Sci. Biotechnol.* **2020**, *23*, 413–425, doi:10.1007/s12892-020-00049-2. 825
3. McSteen, P.; Kellogg, E.A. Molecular, Cellular, and Developmental Foundations of Grass Diversity. *Science* **2022**, *377*, 599–602, doi:10.1126/science.abo5035. 827
4. Lopez, B.; Hines, P.J.; Ash, C. The Unrecognized Value of Grass. *Science* **2022**, *377*, 590–591, doi:10.1126/science.add6362. 829
5. Strömborg, C.A.E.; Staver, A.C. The History and Challenge of Grassy Biomes. *Science* **2022**, *377*, 592–593, doi:10.1126/science.add1347. 831
6. Bengtsson, J.; Bullock, J.M.; Egoh, B.; Everson, C.; Everson, T.; O'Connor, T.; O'Farrell, P.J.; Smith, H.G.; Lindborg, R. Grasslands—More Important for Ecosystem Services than You Might Think. *Ecosphere* **2019**, *10*, e02582, doi:10.1002/ecs2.2582. 833
7. Terrer, C.; Phillips, R.P.; Hungate, B.A.; Rosende, J.; Pett-Ridge, J.; Craig, M.E.; van Groenigen, K.J.; Keenan, T.F.; Sulman, B.N.; Stocker, B.D.; et al. A Trade-off between Plant and Soil Carbon Storage under Elevated CO₂. *Nature* **2021**, *591*, 599–603, doi:10.1038/s41586-021-03306-8. 838
8. Dass, P.; Houlton, B.Z.; Wang, Y.; Warlind, D. Grasslands May Be More Reliable Carbon Sinks than Forests in California. *Environ. Res. Lett.* **2018**, *13*, 074027, doi:10.1088/1748-9326/aacb39. 839
9. *Building Climate Resilience for Food Security and Nutrition*; FAO, Ed.; The state of food security and nutrition in the world; FAO: Rome, 2018; ISBN 978-92-5-130571-3. 841
10. *Moving Forward on Food Loss and Waste Reduction*; FAO, Ed.; The state of food and agriculture; Food and Agriculture Organization of the United Nations: Rome, 2019; ISBN 978-92-5-131789-1. 843
11. Kwadzo, M.; Quayson, E. Factors Influencing Adoption of Integrated Soil Fertility Management Technologies by Smallholder Farmers in Ghana. *Helijon* **2021**, *7*, e07589, doi:10.1016/j.helijon.2021.e07589. 845
12. Feller, C.; Blanchart, E.; Bernoux, M.; Lal, R.; Manlay, R. Soil Fertility Concepts over the Past Two Centuries: The Importance Attributed to Soil Organic Matter in Developed and Developing Countries. *Arch. Agron. Soil Sci.* **2012**, *58*, S3–S21, doi:10.1080/03650340.2012.693598. 847
13. Moorby, J.M.; Fraser, M.D. Review: New Feeds and New Feeding Systems in Intensive and Semi-Intensive Forage-Fed Ruminant Livestock Systems. *Animal* **2021**, *100297*, doi:10.1016/j.animal.2021.100297. 850
14. Capstaff, N.M.; Miller, A.J. Improving the Yield and Nutritional Quality of Forage Crops. *Front. Plant Sci.* **2018**, *9*, doi:https://doi.org/10.3389/fpls.2018.00535. 852
15. Raza, A.; Razzaq, A.; Mehmood, S.S.; Zou, X.; Zhang, X.; Zhang, X.; Lv, Y.; Jinsong Xu; Xu, J. Impact of Climate Change on Crops Adaptation and Strategies to Tackle Its Outcome: A Review. *2019*, *8*, 34, doi:10.3390/plants8020034. 854
16. Chai, R.; Ye, X.; Ma, C.; Wang, Q.; Tu, R.; Zhang, L.; Gao, H. Greenhouse Gas Emissions from Synthetic Nitrogen Manufacture and Fertilization for Main Upland Crops in China. *Carbon Balance Manag.* **2019**, *14*, 20, doi:10.1186/s13021-019-0133-9. 856
17. Rivero, R.M.; Mittler, R.; Blumwald, E.; Zandalinas, S.I. Developing Climate-Resilient Crops: Improving Plant Tolerance to Stress Combination. *Plant J.* **2021**, doi:10.1111/tpj.15483. 859
18. Giridhar, K.; Samireddypalle, A. Impact of Climate Change on Forage Availability for Livestock. In *Climate Change Impact on Livestock: Adaptation and Mitigation*; Sejian, V., Gaughan, J., Baumgard, L., Prasad, C., Eds.; Springer India: New Delhi, 2015; pp. 97–112 ISBN 978-81-322-2265-1. 861

19. Cushman, J.C.; Denby, K.J.; Mittler, R. Plant Responses and Adaptations to a Changing Climate. *Plant J.* **2022**, *109*, 319–322, doi:10.1111/tpj.15641. 864

20. Bailey-Serres, J.; Parker, J.E.; Ainsworth, E.A.; Oldroyd, G.E.D.; Schroeder, J.I. Genetic Strategies for Improving Crop Yields. *Nature* **2019**, *575*, 109–118, doi:10.1038/s41586-019-1679-0. 866

21. P. Dhakate; D. Sehgal; S. Vaishnavi; A. Chandra; Apekshita Singh; S. Raina; Vijay Rani Rajpal Comprehending the Evolution of Gene Editing Platforms for Crop Trait Improvement. *Front. Genet.* **2022**, doi:10.3389/fgene.2022.876987. 868

22. Liu, Y.; Wang, W.; Yang, B.; Currey, C.; Fei, S. Functional Analysis of the Teosinte Branched 1 Gene in the Tetraploid Switchgrass (Panicum Virgatum L.) by CRISPR/Cas9-Directed Mutagenesis. *Front. Plant Sci.* **2020**, *11*, doi:https://doi.org/10.3389/fpls.2020.572193. 870

23. Weiss, T.; Wang, C.; Kang, X.; Zhao, H.; Elena Gamo, M.; Starker, C.G.; Crisp, P.A.; Zhou, P.; Springer, N.M.; Voytas, D.F.; et al. Optimization of Multiplexed CRISPR/Cas9 System for Highly Efficient Genome Editing in Setaria Viridis. *Plant J.* **2020**, *104*, 828–838, doi:10.1111/tpj.14949. 873

24. Zhang, Y.; Ran, Y.; Nagy, I.; Lenk, I.; Qiu, J.-L.; Asp, T.; Jensen, C.S.; Gao, C. Targeted Mutagenesis in Ryegrass (Lolium Spp.) Using the CRISPR/Cas9 System. *Plant Biotechnol. J.* **2020**, *18*, 1854–1856, doi:10.1111/pbi.13359. 876

25. Cheng, Z.; Sun, Y.; Yang, S.; Zhi, H.; Yin, T.; Ma, X.; Zhang, H.; Diao, X.; Guo, Y.; Li, X.; et al. Establishing in *Planta* Haploid Inducer Line by Edited SiMTL in Foxtail Millet (Setaria Italica). *Plant Biotechnol. J.* **2021**, *19*, 1089–1091, doi:10.1111/pbi.13584. 878

26. Zhang, L.; Wang, T.; Wang, G.; Bi, A.; Wassie, M.; Xie, Y.; Cao, L.; Xu, H.; Fu, J.; Chen, L.; et al. Simultaneous Gene Editing of Three Homoeoalleles in Self-Incompatible Allohexaploid Grasses. *J. Integr. Plant Biol.* **2021**, *63*, 1410–1415, doi:10.1111/jipb.13101. 881

27. Zhu, C.; Liu, L.; Crowell, O.; Zhao, H.; Brutnell, T.P.; Jackson, D.; Kellogg, E.A. The CLV3 Homolog in Setaria Viridis Selectively Controls Inflorescence Meristem Size. *Front. Plant Sci.* **2021**, *12*, doi:https://doi.org/10.3389/fpls.2021.636749. 885

28. Kumar, R.; Kamuda, T.; Budhathoki, R.; Tang, D.; Yer, H.; Zhao, Y.; Li, Y. Agrobacterium- and a Single Cas9-SgRNA Transcript System-Mediated High Efficiency Gene Editing in Perennial Ryegrass. *Front. Genome Ed.* **2022**, *4*, doi:https://doi.org/10.3389/fgeed.2022.960414. 887

29. Wang, H.; Jia, G.; Zhang, N.; Zhi, H.; Xing, L.; Zhang, H.; Sui, Y.; Tang, S.; Li, M.; Zhang, H.; et al. Domestication-Associated PHYTOCHROME C Is a Flowering Time Repressor and a Key Factor Determining Setaria as a Short-Day Plant. *New Phytol.* **2022**, *236*, 1809–1823, doi:10.1111/nph.18493. 890

30. Muguerza, M.B.; Gondo, T.; Ishigaki, G.; Shimamoto, Y.; Umami, N.; Nitthaisong, P.; Rahman, M.M.; Akashi, R. Tissue Culture and Somatic Embryogenesis in Warm-Season Grasses—Current Status and Its Applications: A Review. *Plants* **2022**, *11*, 1263, doi:10.3390/plants11091263. 893

31. Cropano, C.; Manzanares, C.; Yates, S.; Copetti, D.; Do Canto, J.; Lübbertedt, T.; Koch, M.; Studer, B. Identification of Candidate Genes for Self-Compatibility in Perennial Ryegrass (Lolium Perenne L.). *Front. Plant Sci.* **2021**, *12*, doi:https://doi.org/10.3389/fpls.2021.707901. 896

32. Moser, L.E.; Hoveland, C.S. Cool-Season Grass Overview. In *Cool-Season Forage Grasses*; John Wiley & Sons, Ltd, 1996; pp. 1–14 ISBN 978-0-89118-226-9. 899

33. Moser, L.E.; Burson, B.L.; Sollenberger, L.E. Warm-Season (C4) Grass Overview. In *Warm-Season (C4) Grasses*; John Wiley & Sons, Ltd, 2004; pp. 1–14 ISBN 978-0-89118-237-5. 901

34. Sugimoto, K.; Nowack, M.K. Plant Development Meets Climate Emergency – It's Time to Plant an Apple Tree. *Curr. Opin. Plant Biol.* **2022**, *65*, 102175, doi:10.1016/j.pbi.2022.102175. 903

904

35. He, M.; He, C.-Q.; Ding, N.-Z. Abiotic Stresses: General Defenses of Land Plants and Chances for Engineering Multistress Tolerance. *Front. Plant Sci.* **2018**, *9*, doi:<https://doi.org/10.3389/fpls.2018.01771>. 905

36. Villalobos-López, M.A.; Arroyo-Becerra, A.; Quintero-Jiménez, A.; Iturriaga, G. Biotechnological Advances to Improve Abiotic Stress Tolerance in Crops. *Int. J. Mol. Sci.* **2022**, *23*, 12053, doi:10.3390/ijms231912053. 906

37. Bita, C.; Gerats, T. Plant Tolerance to High Temperature in a Changing Environment: Scientific Fundamentals and Production of Heat Stress-Tolerant Crops. *Front. Plant Sci.* **2013**, *4*, doi:<https://doi.org/10.3389/fpls.2013.00273>. 907

38. Bulgari, R.; Franzoni, G.; Ferrante, A. Biostimulants Application in Horticultural Crops under Abiotic Stress Conditions. *Agronomy* **2019**, *9*, 306, doi:10.3390/agronomy9060306. 908

39. Pascual, L.S.; Segarra-Medina, C.; Gómez-Cadenas, A.; López-Climent, M.F.; Vives-Peris, V.; Zandalinas, S.I. Climate Change-Associated Multifactorial Stress Combination: A Present Challenge for Our Ecosystems. *J. Plant Physiol.* **2022**, doi:10.1016/j.jplph.2022.153764. 909

40. Rama, H.-O.; Roberts, D.; Tignor, M.; Poloczanska, E.S.; Mintenbeck, K.; Alegría, A.; Craig, M.; Langsdorf, S.; Löschke, S.; Möller, V.; et al. *Climate Change 2022: Impacts, Adaptation and Vulnerability Working Group II Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*; 2022; 910

41. Loka, D.; Harper, J.; Humphreys, M.; Gasior, D.; Wootton-Beard, P.; Gwynn-Jones, D.; Scullion, J.; Doonan, J.; Kingston-Smith, A.; Dodd, R.; et al. Impacts of Abiotic Stresses on the Physiology and Metabolism of Cool-Season Grasses: A Review. *Food Energy Secur.* **2019**, *8*, e00152, doi:10.1002/fes3.152. 911

42. Jagadish, S.V.K.; Way, D.A.; Sharkey, T.D. Plant Heat Stress: Concepts Directing Future Research. *Plant Cell Environ.* **2021**, *44*, 1992–2005, doi:10.1111/pce.14050. 912

43. Liu, X.; Meng, P.; Yang, G.; Zhang, M.; Peng, S.; Zhai, M.Z. Genome-Wide Identification and Transcript Profiles of Walnut Heat Stress Transcription Factor Involved in Abiotic Stress. *BMC Genomics* **2020**, *21*, 474, doi:10.1186/s12864-020-06879-2. 913

44. Hasanuzzaman, M.; Nahar, K.; Alam, Md.M.; Roychowdhury, R.; Fujita, M. Physiological, Biochemical, and Molecular Mechanisms of Heat Stress Tolerance in Plants. *Int. J. Mol. Sci.* **2013**, *14*, 9643–9684, doi:10.3390/ijms14059643. 914

45. Wang, J.; Vanga, S.K.; Saxena, R.; Orsat, V.; Raghavan, V. Effect of Climate Change on the Yield of Cereal Crops: A Review. *Climate* **2018**, *6*, 41, doi:10.3390/cli6020041. 915

46. Aleem, S.; Sharif, I.; Amin, E.; Tahir, M.; Parveen, N.; Aslam, R.; Najeebulah, M.; Shahid, M.T.H. Heat Tolerance in Vegetables in the Current Genomic Era: An Overview. *Plant Growth Regul.* **2020**, *92*, 497–516, doi:10.1007/s10725-020-00658-5. 916

47. Bourgine, B.; Guihur, A. Heat Shock Signaling in Land Plants: From Plasma Membrane Sensing to the Transcription of Small Heat Shock Proteins. *Front. Plant Sci.* **2021**, *12*, 710801, doi:10.3389/fpls.2021.710801. 917

48. Xu, T.; Niu, J.; Jiang, Z. Sensing Mechanisms: Calcium Signaling Mediated Abiotic Stress in Plants. *Front. Plant Sci.* **2022**, *13*, doi:<https://doi.org/10.3389/fpls.2022.925863>. 918

49. DeFalco, T.A.; Moeder, W.; Yoshioka, K. Opening the Gates: Insights into Cyclic Nucleotide-Gated Channel-Mediated Signaling. *Trends Plant Sci.* **2016**, *21*, 903–906, doi:10.1016/j.tplants.2016.08.011. 919

50. Yan, Q.; Huang, Q.; Chen, J.; Li, J.; Liu, Z.; Yang, Y.; Li, X.; Wang, J. SYTA Has Positive Effects on the Heat Resistance of Arabidopsis. *Plant Growth Regul.* **2017**, *81*, 467–476, doi:10.1007/s10725-016-0224-5. 920

51. Wu, X.; Wang, Y.; Bian, Y.; Ren, Y.; Xu, X.; Zhou, F.; Ding, H. A Critical Review on Plant Annexin: Structure, Function, and Mechanism. *Plant Physiol. Biochem.* **2022**, *190*, 81–89, doi:10.1016/j.plaphy.2022.08.019. 921

52. Cui, Y.; Lu, S.; Li, Z.; Cheng, J.; Hu, P.; Zhu, T.; Wang, X.; Jin, M.; Wang, X.; Li, L.; et al. CYCLIC NUCLEOTIDE-GATED ION CHANNELS 14 and 16 Promote Tolerance to Heat and Chilling in Rice. *Plant Physiol.* **2020**, *183*, 1794–1808, doi:10.1104/pp.20.00591. 922

53. Hasanuzzaman, M.; Bhuyan, M.H.M.B.; Zulfiqar, F.; Raza, A.; Mohsin, S.M.; Mahmud, J.A.; Fujita, M.; Fotopoulos, V. Reactive Oxygen Species and Antioxidant Defense in Plants under Abiotic Stress: Revisiting the Crucial Role of a Universal Defense Regulator. *Antioxidants* **2020**, *9*, 681, doi:10.3390/antiox9080681. 947
948
949

54. Singh, A.; Kumar, A.; Yadav, S.; Singh, I.K. Reactive Oxygen Species-Mediated Signaling during Abiotic Stress. *Plant Gene* **2019**, *18*, 100173, doi:10.1016/j.plgene.2019.100173. 950
951

55. Krishna, P. Plant Responses to Heat Stress. In *Plant Responses to Abiotic Stress*; Hirt, H., Shinozaki, K., Eds.; Topics in Current Genetics; Springer: Berlin, Heidelberg, 2004; pp. 73–101 ISBN 978-3-540-39402-0. 952
953

56. Wang, X.; Huang, W.; Liu, J.; Yang, Z.; Huang, B. Molecular Regulation and Physiological Functions of a Novel FaHsfA2c Cloned from Tall Fescue Conferring Plant Tolerance to Heat Stress. *Plant Biotechnol. J.* **2017**, *15*, 237–248, doi:10.1111/pbi.12609. 954
955
956

57. Rognli, O.A. Breeding for Improved Winter Survival in Forage Grasses. In *Proceedings of the Plant and Microbe Adaptations to Cold in a Changing World*; Imai, R., Yoshida, M., Matsumoto, N., Eds.; Springer: New York, NY, 2013; pp. 197–208. 957
958
959

58. Thomashow, M.F. PLANT COLD ACCLIMATION: Freezing Tolerance Genes and Regulatory Mechanisms. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **1999**, *50*, 571–599, doi:10.1146/annurev.arplant.50.1.571. 960
961

59. Chinnusamy, V.; Zhu, J.; Zhu, J.-K. Cold Stress Regulation of Gene Expression in Plants. *Trends Plant Sci.* **2007**, *12*, 444–451, doi:10.1016/j.tplants.2007.07.002. 962
963

60. Fjellheim, S.; Boden, S.; Trevaskis, B. The Role of Seasonal Flowering Responses in Adaptation of Grasses to Temperate Climates. *Front. Plant Sci.* **2014**, *5*, doi:https://doi.org/10.3389/fpls.2014.00431. 964
965

61. Heide, O.M. Control of Flowering and Reproduction in Temperate Grasses. *New Phytol.* **1994**, *128*, 347–362, doi:10.1111/j.1469-8137.1994.tb04019.x. 966
967

62. Gudleifsson, B.E. Ice Encasement Damage on Grass Crops and Alpine Plants in Iceland - Impact of Climate Change. In *Plant cold hardiness: from the laboratory to the field*; CABI, 2009; pp. 163–172 ISBN 978-1-84593-513-9. 968
969

63. Larsen, A. Breeding Winter Hardy Grasses. In *Breeding Fodder Crops for Marginal Conditions: Proceedings of the 18th Eucarpia Fodder Crops Section Meeting, Loen, Norway, 25–28 August 1993*; Rognli, O.A., Solberg, E., Schjelderup, I., Eds.; *Developments in Plant Breeding*; Springer Netherlands: Dordrecht, 1994; pp. 149–158 ISBN 978-94-011-0966-6. 970
971
972
973

64. Andrews, C.J.; Gudleifsson, B.E. A Comparison of Cold Hardiness and Ice Encasement Tolerance of Timothy Grass and Winter Wheat. *Can. J. Plant Sci.* **1983**, *63*, 429–435, doi:10.4141/cjps83-049. 974
975

65. Gudleifsson, B.E.; Andrews, C.J.; Bjornsson, H. Cold Hardiness and Ice Tolerance of Pasture Grasses Grown and Tested in Controlled Environments. *Can. J. Plant Sci.* **1986**, *66*, 601–608, doi:10.4141/cjps86-080. 976
977

66. Maruyama, K.; Urano, K.; Yoshiwara, K.; Morishita, Y.; Sakurai, N.; Suzuki, H.; Kojima, M.; Sakakibara, H.; Shibata, D.; Saito, K.; et al. Integrated Analysis of the Effects of Cold and Dehydration on Rice Metabolites, Phytohormones, and Gene Transcripts. *Plant Physiol.* **2014**, *164*, 1759–1771, doi:10.1104/pp.113.231720. 978
979
980

67. Nakaminami, K.; Matsui, A.; Nakagami, H.; Minami, A.; Nomura, Y.; Tanaka, M.; Morosawa, T.; Ishida, J.; Takahashi, S.; Uemura, M.; et al. Analysis of Differential Expression Patterns of mRNA and Protein During Cold-Acclimation and De-Acclimation in *Arabidopsis*. *Mol. Cell. Proteomics* **2014**, *13*, 3602–3611, doi:10.1074/mcp.M114.039081. 981
982
983
984

68. Yamaguchi-Shinozaki, K.; Shinozaki, K. Transcriptional Regulatory Networks in Cellular Responses and Tolerance to Dehydration and Cold Stresses. *Annu. Rev. Plant Biol.* **2006**, *57*, 781–803, doi:10.1146/annurev.arplant.57.032905.105444. 985
986
987

69. Thomashow, M.F. Molecular Basis of Plant Cold Acclimation: Insights Gained from Studying the CBF Cold Response Pathway. *Plant Physiol.* **2010**, *154*, 571–577, doi:10.1104/pp.110.161794. 988

70. Gray, G.R.; Chauvin, L.P.; Sarhan, F.; Huner, N.P.A. Cold Acclimation and Freezing Tolerance (A Complex Interaction of Light and Temperature). *Plant Physiol.* **1997**, *114*, 467–474, doi:10.1104/pp.114.2.467. 990

71. Janda, T.; Majláth, I.; Szalai, G. Interaction of Temperature and Light in the Development of Freezing Tolerance in Plants. *J. Plant Growth Regul.* **2014**, *33*, 460–469, doi:10.1007/s00344-013-9381-1. 992

72. Rapacz, M.; Ergon, Å.; Höglind, M.; Jørgensen, M.; Jurczyk, B.; Østrem, L.; Rognli, O.A.; Tronsmo, A.M. Overwintering of Herbaceous Plants in a Changing Climate. Still More Questions than Answers. *Plant Sci.* **2014**, *225*, 34–44, doi:10.1016/j.plantsci.2014.05.009. 994

73. Dalmannsdottir, S.; Jørgensen, M.; Rapacz, M.; Østrem, L.; Larsen, A.; Rødven, R.; Rognli, O.A. Cold Acclimation in Warmer Extended Autumns Impairs Freezing Tolerance of Perennial Ryegrass (*Lolium Perenne*) and Timothy (*Phleum Pratense*). *Physiol. Plant.* **2017**, *160*, 266–281, doi:10.1111/ppl.12548. 998

74. Dalmannsdottir, S.; Rapacz, M.; Jørgensen, M.; Østrem, L.; Larsen, A.; Rødven, R.; Rognli, O.A. Temperature Before Cold Acclimation Affects Cold Tolerance and Photoacclimation in Timothy (*Phleum Pratense* L.), Perennial Ryegrass (*Lolium Perenne* L.) and Red Clover (*Trifolium Pratense* L.). *J. Agron. Crop Sci.* **2016**, *202*, 320–330, doi:10.1111/jac.12149. 1000

75. Jørgensen, M.; Torp, T.; Mølmann, J.A.B. Impact of Waterlogging and Temperature on Autumn Growth, Hardening and Freezing Tolerance of Timothy (*Phleum Pratense*). *J. Agron. Crop Sci.* **2020**, *206*, 242–251, doi:10.1111/jac.12385. 1004

76. Kovi, M.R.; Ergon, Å.; Rognli, O.A. Freezing Tolerance Revisited—Effects of Variable Temperatures on Gene Regulation in Temperate Grasses and Legumes. *Curr. Opin. Plant Biol.* **2016**, *33*, 140–146, doi:10.1016/j.pbi.2016.07.006. 1006

77. Rapacz, M.; Jurczyk, B.; Sasal, M. Deacclimation May Be Crucial for Winter Survival of Cereals under Warming Climate. *Plant Sci.* **2017**, *256*, 5–15, doi:10.1016/j.plantsci.2016.11.007. 1008

78. Jaškunė, K.; Armonienė, R.; Liatukas, Ž.; Statkevičiūtė, G.; Cesevičienė, J.; Brazauskas, G. Relationship between Freezing Tolerance and Leaf Growth during Acclimation in Winter Wheat. *Agronomy* **2022**, *12*, 859, doi:10.3390/agronomy12040859. 1010

79. Ding, Y.; Shi, Y.; Yang, S. Advances and Challenges in Uncovering Cold Tolerance Regulatory Mechanisms in Plants. *New Phytol.* **2019**, *222*, 1690–1704, doi:10.1111/nph.15696. 1013

80. Park, S.; Gilmour, S.J.; Grumet, R.; Thomashow, M.F. CBF-Dependent and CBF-Independent Regulatory Pathways Contribute to the Differences in Freezing Tolerance and Cold-Regulated Gene Expression of Two *Arabidopsis* Ecotypes Locally Adapted to Sites in Sweden and Italy. *PLOS ONE* **2018**, *13*, e0207723, doi:10.1371/journal.pone.0207723. 1015

81. Galiba, G.; Vágújfalvi, A.; Li, C.; Soltész, A.; Dubcovsky, J. Regulatory Genes Involved in the Determination of Frost Tolerance in Temperate Cereals. *Plant Sci.* **2009**, *176*, 12–19, doi:10.1016/j.plantsci.2008.09.016. 1019

82. Liu, Y.; Song, Q.; Li, D.; Yang, X.; Li, D. Multifunctional Roles of Plant Dehydrins in Response to Environmental Stresses. *Front. Plant Sci.* **2017**, *8*, doi:https://doi.org/10.3389/fpls.2017.01018. 1021

83. Vazquez-Hernandez, M.; Romero, I.; Escribano, M.I.; Merodio, C.; Sanchez-Ballesta, M.T. Deciphering the Role of CBF/DREB Transcription Factors and Dehydrins in Maintaining the Quality of Table Grapes Cv. Autumn Royal Treated with High CO₂ Levels and Stored at 0°C. *Front. Plant Sci.* **2017**, *8*, doi:https://doi.org/10.3389/fpls.2017.01591. 1023

84. Fu, J.; Miao, Y.; Shao, L.; Hu, T.; Yang, P. De Novo Transcriptome Sequencing and Gene Expression Profiling of *Elymus Nutans* under Cold Stress. *BMC Genomics* **2016**, *17*, 870, doi:10.1186/s12864-016-3222-0. 1026

85. 1027

85. Alm, V.; Busso, C.S.; Ergon, Å.; Rudi, H.; Larsen, A.; Humphreys, M.W.; Rognli, O.A. QTL Analyses and Comparative Genetic Mapping of Frost Tolerance, Winter Survival and Drought Tolerance in Meadow Fescue (*Festuca pratensis* Huds.). *Theor. Appl. Genet.* **2011**, *123*, 369–382, doi:10.1007/s00122-011-1590-z. 1028
1029
1030

86. Bhowmik, P.K.; Tamura, K.; Sanada, Y.; Tase, K.; Yamada, T. Sucrose Metabolism of Perennial Ryegrass in Relation to Cold Acclimation. *Z. Für Naturforschung C* **2006**, *61*, 99–104, doi:10.1515/znc-2006-1-218. 1031
1032

87. Waterhouse, A.L.; Chatterton, N.J. Glossary of Fructan Terms. In *Science and technology of fructans*; CRC Press Boca Raton, FL, 1993; pp. 1–7 ISBN 0-8493-5111-1. 1033
1034

88. Valluru, R.; Van den Ende, W. Plant Fructans in Stress Environments: Emerging Concepts and Future Prospects. *J. Exp. Bot.* **2008**, *59*, 2905–2916, doi:10.1093/jxb/ern164. 1035
1036

89. Chalmers, J.; Lidgett, A.; Cummings, N.; Cao, Y.; Forster, J.; Spangenberg, G. Molecular Genetics of Fructan Metabolism in Perennial Ryegrass. *Plant Biotechnol. J.* **2005**, *3*, 459–474, doi:10.1111/j.1467-7652.2005.00148.x. 1037
1038

90. Yoshida, M. Fructan Structure and Metabolism in Overwintering Plants. *Plants* **2021**, *10*, 933, doi:10.3390/plants10050933. 1039
1040

91. Wu, S.; Greiner, S.; Ma, C.; Zhong, J.; Huang, X.; Rausch, T.; Zhao, H. A Fructan Exohydrolase from Maize Degrades Both Inulin and Levan and Co-Exists with 1-Kestotriose in Maize. *Int. J. Mol. Sci.* **2021**, *22*, 5149, doi:10.3390/ijms22105149. 1041
1042
1043

92. Farrant, J.M.; Cooper, K.; Hilgart, A.; Abdalla, K.O.; Bentley, J.; Thomson, J.A.; Dace, H.J.W.; Peton, N.; Mundree, S.G.; Rafudeen, M.S. A Molecular Physiological Review of Vegetative Desiccation Tolerance in the Resurrection Plant *Xerophyta Viscosa* (Baker). *Planta* **2015**, *242*, 407–426, doi:10.1007/s00425-015-2320-6. 1044
1045
1046

93. VanBuren, R.; Wai, C.M.; Zhang, Q.; Song, X.; Edger, P.P.; Bryant, D.; Michael, T.P.; Mockler, T.C.; Bartels, D. Seed Desiccation Mechanisms Co-Opted for Vegetative Desiccation in the Resurrection Grass *Oropetium Thomaeum*. *Plant Cell Environ.* **2017**, *40*, 2292–2306, doi:10.1111/pce.13027. 1047
1048
1049

94. Hilhorst, H.W.M.; Costa, M.-C.D.; Farrant, J.M. A Footprint of Plant Desiccation Tolerance. Does It Exist? *Mol. Plant* **2018**, *11*, 1003–1005, doi:10.1016/j.molp.2018.07.001. 1050
1051

95. Oliver, M.J.; Farrant, J.M.; Hilhorst, H.W.M.; Mundree, S.; Williams, B.; Bewley, J.D. Desiccation Tolerance: Avoiding Cellular Damage During Drying and Rehydration. *Annu. Rev. Plant Biol.* **2020**, *71*, 435–460, doi:10.1146/annurev-plant-071219-105542. 1052
1053
1054

96. Kooyers, N.J. The Evolution of Drought Escape and Avoidance in Natural Herbaceous Populations. *Plant Sci. Int. J. Exp. Plant Biol.* **2015**, *234*, 155–162, doi:10.1016/j.plantsci.2015.02.012. 1055
1056

97. Pardo, J.; Man Wai, C.; Chay, H.; Madden, C.F.; Hilhorst, H.W.M.; Farrant, J.M.; VanBuren, R. Intertwined Signatures of Desiccation and Drought Tolerance in Grasses. *Proc. Natl. Acad. Sci.* **2020**, *117*, 10079–10088, doi:10.1073/pnas.2001928117. 1057
1058
1059

98. Moore, F.C.; Lobell, D.B. The Fingerprint of Climate Trends on European Crop Yields. *Proc. Natl. Acad. Sci.* **2015**, *112*, 2670–2675, doi:10.1073/pnas.1409606112. 1060
1061

99. Ergon, Å.; Seddaiu, G.; Korhonen, P.; Virkajarvi, P.; Bellocchi, G.; Jørgensen, M.; Østrem, L.; Reheul, D.; Volaire, F. How Can Forage Production in Nordic and Mediterranean Europe Adapt to the Challenges and Opportunities Arising from Climate Change? *Eur. J. Agron.* **2018**, *92*, 97–106, doi:10.1016/j.eja.2017.09.016. 1062
1063
1064

100. Jaškūnė, K.; Aleliūnas, A.; Statkevičiūtė, G.; Kemešytė, V.; Studer, B.; Yates, S. Genome-Wide Association Study to Identify Candidate Loci for Biomass Formation Under Water Deficit in Perennial Ryegrass. *Front. Plant Sci.* **2020**, *11*, doi:https://doi.org/10.3389/fpls.2020.570204. 1065
1066
1067

101. Farooq, M.; Wahid, A.; Kobayashi, N.; Fujita, D.; Basra, S.M.A. Plant Drought Stress: Effects, Mechanisms and Management. In *Sustainable Agriculture*; Lichtfouse, E., Navarrete, M., Debaeke, P., Véronique, S., Alberola, C., Eds.; Springer Netherlands: Dordrecht, 2009; pp. 153–188 ISBN 978-90-481-2666-8. 1068
1069
1070

102. Wilkinson, S.; Davies, W.J. Drought, Ozone, ABA and Ethylene: New Insights from Cell to Plant to Community. *Plant Cell Environ.* **2010**, *33*, 510–525, doi:10.1111/j.1365-3040.2009.02052.x. 1071
1072

103. Lee, S.C.; Luan, S. ABA Signal Transduction at the Crossroad of Biotic and Abiotic Stress Responses. *Plant Cell Environ.* **2012**, *35*, 53–60, doi:10.1111/j.1365-3040.2011.02426.x. 1073
1074

104. Saab, I.N.; Sharp, R.E.; Pritchard, J.; Voetberg, G.S. Increased Endogenous Abscisic Acid Maintains Primary Root Growth and Inhibits Shoot Growth of Maize Seedlings at Low Water Potentials 1. *Plant Physiol.* **1990**, *93*, 1329–1336, doi:10.1104/pp.93.4.1329. 1075
1076
1077

105. Li, X.; Wilkinson, S.; Shen, J.; Forde, B.G.; Davies, W.J. Stomatal and Growth Responses to Hydraulic and Chemical Changes Induced by Progressive Soil Drying. *J. Exp. Bot.* **2017**, *68*, 5883–5894, doi:10.1093/jxb/erx381. 1078
1079

106. Khadka, V.S.; Vaughn, K.; Xie, J.; Swaminathan, P.; Ma, Q.; Cramer, G.R.; Fennell, A.Y. Transcriptomic Response Is More Sensitive to Water Deficit in Shoots than Roots of *Vitis Riparia* (Michx.). *BMC Plant Biol.* **2019**, *19*, 72, doi:10.1186/s12870-019-1664-7. 1080
1081
1082

107. Sah, S.K.; Reddy, K.R.; Li, J. Abscisic Acid and Abiotic Stress Tolerance in Crop Plants. *Front. Plant Sci.* **2016**, *7*, doi:https://doi.org/10.3389/fpls.2016.00571. 1083
1084

108. Hussain, S.; Khan, F.; Cao, W.; Wu, L.; Geng, M. Seed Priming Alters the Production and Detoxification of Reactive Oxygen Intermediates in Rice Seedlings Grown under Sub-Optimal Temperature and Nutrient Supply. *Front. Plant Sci.* **2016**, *7*, doi:https://doi.org/10.3389/fpls.2016.00439. 1085
1086
1087

109. Hussain, M.; Farooq, S.; Hasan, W.; Ul-Allah, S.; Tanveer, M.; Farooq, M.; Nawaz, A. Drought Stress in Sunflower: Physiological Effects and Its Management through Breeding and Agronomic Alternatives. *Agric. Water Manag.* **2018**, *201*, 152–166, doi:10.1016/j.agwat.2018.01.028. 1088
1089
1090

110. Svensson, J.; Ismail, A.M.; Tapio Palva, E.; Close, T.J. Chapter 11 - Dehydrins. In *Cell and Molecular Response to Stress*; Storey, K.B., Storey, J.M., Eds.; Sensing, Signaling and Cell Adaptation; Elsevier, 2002; Vol. 3, pp. 155–171. 1091
1092

111. Kosová, K.; Prášil, I.T.; Vítámvás, P. Role of Dehydrins in Plant Stress Response. In *Handbook of Plant and Crop Stress, Fourth Edition*; CRC Press, 2019 ISBN 978-1-351-10460-9. 1093
1094

112. Liu, Y.; Song, Q.; Li, D.; Yang, X.; Li, D. Multifunctional Roles of Plant Dehydrins in Response to Environmental Stresses. *Front. Plant Sci.* **2017**, *8*, doi:https://doi.org/10.3389/fpls.2017.01018. 1095
1096

113. Sun, Z.; Li, S.; Chen, W.; Zhang, J.; Zhang, L.; Sun, W.; Wang, Z. Plant Dehydrins: Expression, Regulatory Networks, and Protective Roles in Plants Challenged by Abiotic Stress. *Int. J. Mol. Sci.* **2021**, *22*, 12619, doi:10.3390/ijms222312619. 1097
1098
1099

114. Pornsiriwong, W.; Estavillo, G.M.; Chan, K.X.; Tee, E.E.; Ganguly, D.; Crisp, P.A.; Phua, S.Y.; Zhao, C.; Qiu, J.; Park, J.; et al. A Chloroplast Retrograde Signal, 3'-Phosphoadenosine 5'-Phosphate, Acts as a Secondary Messenger in Abscisic Acid Signaling in Stomatal Closure and Germination. *eLife* **2017**, *6*, e23361, doi:10.7554/eLife.23361. 1100
1101
1102

115. Estavillo, G.M.; Crisp, P.A.; Pornsiriwong, W.; Wirtz, M.; Collinge, D.; Carrie, C.; Giraud, E.; Whelan, J.; David, P.; Javot, H.; et al. Evidence for a SAL1-PAP Chloroplast Retrograde Pathway That Functions in Drought and High Light Signaling in Arabidopsis. *Plant Cell* **2011**, *23*, 3992–4012, doi:10.1105/tpc.111.091033. 1103
1104
1105

116. Abdallah, N.A.; Elsharawy, H.; Abulela, H.A.; Thilmony, R.; Abdelhadi, A.A.; Elarabi, N.I. Multiplex CRISPR/Cas9-Mediated Genome Editing to Address Drought Tolerance in Wheat. *GM Crops Food* **2022**, *0*, 1–17, doi:10.1080/21645698.2022.2120313. 1106
1107
1108

117. Chaves, M.M.; Flexas, J.; Pinheiro, C. Photosynthesis under Drought and Salt Stress: Regulation Mechanisms from Whole Plant to Cell. *Ann. Bot.* **2009**, *103*, 551–560, doi:10.1093/aob/mcn125. 1109
1110

118. Ma, Y.; Dias, M.C.; Freitas, H. Drought and Salinity Stress Responses and Microbe-Induced Tolerance in Plants. *Front. Plant Sci.* **2020**, *11*, doi:https://doi.org/10.3389/fpls.2020.591911. 1111
1112

119. Qadir, M.; Quillérou, E.; Nangia, V.; Murtaza, G.; Singh, M.; Thomas, R. j.; Drechsel, P.; Noble, A. d. Economics of Salt-Induced Land Degradation and Restoration. *Nat. Resour. Forum* **2014**, *38*, 282–295, doi:10.1111/1477-8947.12054. 1113
1114

120. Farooq, M.; Asif, S.; Jang, Y.-H.; Park, J.-R.; Zhao, D.-D.; Kim, E.-G.; Kim, K.-M. Effect of Different Salts on Nutrients Uptake, Gene Expression, Antioxidant, and Growth Pattern of Selected Rice Genotypes. *Front. Plant Sci.* **2022**, *13*, doi:https://doi.org/10.3389/fpls.2022.895282. 1115
1116
1117

121. Tester, M.; Davenport, R. Na⁺ Tolerance and Na⁺ Transport in Higher Plants. *Ann. Bot.* **2003**, *91*, 503–527, doi:10.1093/aob/mcg058. 1118
1119

122. Zhao, S.; Zhang, Q.; Liu, M.; Zhou, H.; Ma, C.; Wang, P. Regulation of Plant Responses to Salt Stress. *Int. J. Mol. Sci.* **2021**, *22*, 4609, doi:10.3390/ijms22094609. 1120
1121

123. Shabala, S.; Wu, H.; Bose, J. Salt Stress Sensing and Early Signalling Events in Plant Roots: Current Knowledge and Hypothesis. *Plant Sci.* **2015**, *241*, 109–119, doi:10.1016/j.plantsci.2015.10.003. 1122
1123

124. Han, Y.; Wang, Y.; Zhai, Y.; Wen, Z.; Liu, J.; Xi, C.; Zhao, H.; Wang, Y.; Han, S. OsOSCA1.1 Mediates Hyperosmolality and Salt Stress Sensing in *Oryza Sativa*. *Biology* **2022**, *11*, 678, doi:10.3390/biology11050678. 1124
1125

125. She, K.; Pan, W.; Yan, Y.; Shi, T.; Chu, Y.; Cheng, Y.; Ma, B.; Song, W. Genome-Wide Identification, Evolution and Expressional Analysis of OSCA Gene Family in Barley (*Hordeum Vulgare L.*). *Int. J. Mol. Sci.* **2022**, *23*, 13027, doi:10.3390/ijms232113027. 1126
1127
1128

126. Martínez-Atienza, J.; Jiang, X.; Garciadeblas, B.; Mendoza, I.; Zhu, J.-K.; Pardo, J.M.; Quintero, F.J. Conservation of the Salt Overly Sensitive Pathway in Rice. *Plant Physiol.* **2007**, *143*, 1001–1012, doi:10.1104/pp.106.092635. 1129
1130

127. Kanwar, P.; Sanyal, S.K.; Tokas, I.; Yadav, A.K.; Pandey, A.; Kapoor, S.; Pandey, G.K. Comprehensive Structural, Interaction and Expression Analysis of CBL and CIPK Complement during Abiotic Stresses and Development in Rice. *Cell Calcium* **2014**, *56*, 81–95, doi:10.1016/j.ceca.2014.05.003. 1131
1132
1133

128. Flowers, T.J.; Colmer, T.D. Salinity Tolerance in Halophytes. *New Phytol.* **2008**, *179*, 945–963, doi:10.1111/j.1469-8137.2008.02531.x. 1134
1135

129. Zhao, C.; Zhang, H.; Song, C.; Zhu, J.-K.; Shabala, S. Mechanisms of Plant Responses and Adaptation to Soil Salinity. *The Innovation* **2020**, *1*, 100017, doi:10.1016/j.xinn.2020.100017. 1136
1137

130. Jamil, A.; Riaz, S.; Ashraf, M.; Foolad, M.R. Gene Expression Profiling of Plants under Salt Stress. *Crit. Rev. Plant Sci.* **2011**, *30*, 435–458, doi:10.1080/07352689.2011.605739. 1138
1139

131. Bushman, B.S.; Amundsen, K.L.; Warnke, S.E.; Robins, J.G.; Johnson, P.G. Transcriptome Profiling of Kentucky Bluegrass (*Poa Pratensis L.*) Accessions in Response to Salt Stress. *BMC Genomics* **2016**, *17*, 48, doi:10.1186/s12864-016-2379-x. 1140
1141
1142

132. Chavarria, M.R.; Wherley, B.; Jessup, R.; Chandra, A. Leaf Anatomical Responses and Chemical Composition of Warm-Season Turfgrasses to Increasing Salinity. *Curr. Plant Biol.* **2020**, *22*, 100147, doi:10.1016/j.cpb.2020.100147. 1143
1144

133. Spiekerman, J.J.; Devos, K.M. The Halophyte Seashore Paspalum Uses Adaxial Leaf Papillae for Sodium Sequestration. *Plant Physiol.* **2020**, *184*, 2107–2119, doi:10.1104/pp.20.00796. 1145
1146

134. Zhai, Y.; Wen, Z.; Han, Y.; Zhuo, W.; Wang, F.; Xi, C.; Liu, J.; Gao, P.; Zhao, H.; Wang, Y.; et al. Heterogeneous Expression of Plasma-Membrane-Localised OsOSCA1.4 Complements Osmotic Sensing Based on Hyperosmolality and Salt Stress in *Arabidopsis* Osca1 Mutant. *Cell Calcium* **2020**, *91*, 102261, doi:10.1016/j.ceca.2020.102261. 1147
1148
1149

135. Mishra, A.; Tanna, B. Halophytes: Potential Resources for Salt Stress Tolerance Genes and Promoters. *Front. Plant Sci.* **2017**, *8*, doi:<https://doi.org/10.3389/fpls.2017.00829>. 1150
1151

136. Xu, H.-S.; Guo, S.-M.; Zhu, L.; Xing, J.-C. Growth, Physiological and Transcriptomic Analysis of the Perennial Ryegrass *Lolium Perenne* in Response to Saline Stress. *R. Soc. Open Sci.* **2020**, *7*, 200637, doi:[10.1098/rsos.200637](https://doi.org/10.1098/rsos.200637). 1152
1153

137. Mann, A.; Kumar, N.; Kumar, A.; Lata, C.; Kumar, A.; Meena, B.L.; Mishra, D.; Grover, M.; Gaba, S.; Parameswaran, C.; et al. De Novo Transcriptomic Profiling of Differentially Expressed Genes in Grass Halophyte *Urochondra Setulosa* under High Salinity. *Sci. Rep.* **2021**, *11*, 5548, doi:[10.1038/s41598-021-85220-7](https://doi.org/10.1038/s41598-021-85220-7). 1154
1155
1156

138. Vaziriyaneganeh, M.; Khan, S.; Zwiazek, J.J. Transcriptome and Metabolome Analyses Reveal Potential Salt Tolerance Mechanisms Contributing to Maintenance of Water Balance by the Halophytic Grass *Puccinellia Nuttalliana*. *Front. Plant Sci.* **2021**, *12*. 1157
1158
1159

139. Gasiunas, G.; Barrangou, R.; Horvath, P.; Siksnys, V. Cas9–CrRNA Ribonucleoprotein Complex Mediates Specific DNA Cleavage for Adaptive Immunity in Bacteria. *Proc. Natl. Acad. Sci.* **2012**, *109*, E2579–E2586, doi:[10.1073/pnas.1208507109](https://doi.org/10.1073/pnas.1208507109). 1160
1161
1162

140. Jinek, M.; Chylinski, K.; Fonfara, I.; Hauer, M.; Doudna, J.A.; Charpentier, E. A Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity. *Science* **2012**, *337*, 816–821, doi:[10.1126/science.1225829](https://doi.org/10.1126/science.1225829). 1163
1164

141. Feng, Z.; Zhang, B.; Ding, W.; Liu, X.; Yang, D.-L.; Wei, P.; Cao, F.; Zhu, S.; Zhang, F.; Mao, Y.; et al. Efficient Genome Editing in Plants Using a CRISPR/Cas System. *Cell Res.* **2013**, *23*, 1229–1232, doi:[10.1038/cr.2013.114](https://doi.org/10.1038/cr.2013.114). 1165
1166

142. Jiang, W.; Zhou, H.; Bi, H.; Fromm, M.; Yang, B.; Weeks, D.P. Demonstration of CRISPR/Cas9/SgRNA-Mediated Targeted Gene Modification in *Arabidopsis*, Tobacco, Sorghum and Rice. *Nucleic Acids Res.* **2013**, *41*, e188, doi:[10.1093/nar/gkt780](https://doi.org/10.1093/nar/gkt780). 1167
1168
1169

143. Li, J.-F.; Norville, J.E.; Aach, J.; McCormack, M.; Zhang, D.; Bush, J.; Church, G.M.; Sheen, J. Multiplex and Homologous Recombination–Mediated Genome Editing in *Arabidopsis* and *Nicotiana Benthamiana* Using Guide RNA and Cas9. *Nat. Biotechnol.* **2013**, *31*, 688–691, doi:[10.1038/nbt.2654](https://doi.org/10.1038/nbt.2654). 1170
1171
1172

144. Nekrasov, V.; Staskawicz, B.; Weigel, D.; Jones, J.D.G.; Kamoun, S. Targeted Mutagenesis in the Model Plant *Nicotiana Benthamiana* Using Cas9 RNA-Guided Endonuclease. *Nat. Biotechnol.* **2013**, *31*, 691–693, doi:[10.1038/nbt.2655](https://doi.org/10.1038/nbt.2655). 1173
1174
1175

145. Shan, Q.; Wang, Y.; Li, J.; Zhang, Y.; Chen, K.; Liang, Z.; Zhang, K.; Liu, J.; Xi, J.J.; Qiu, J.-L.; et al. Targeted Genome Modification of Crop Plants Using a CRISPR-Cas System. *Nat. Biotechnol.* **2013**, *31*, 686–688, doi:[10.1038/nbt.2650](https://doi.org/10.1038/nbt.2650). 1176
1177

146. GE *Setaria Viridis* Molecular Characterization Details Available online: https://www.euginius.eu/euginius/pages/gmo_detail.jsf?gmoname=GE-193-31 (accessed on 24 November 2022). 1178
1179

147. Zhang, L.; Wang, T.; Wang, G.; Bi, A.; Wassie, M.; Xie, Y.; Cao, L.; Xu, H.; Fu, J.; Chen, L.; et al. Simultaneous Gene Editing of Three Homoeoalleles in Self-Incompatible Allohexaploid Grasses. *J. Integr. Plant Biol.* **2021**, *63*, 1410–1415, doi:[10.1111/jipb.13101](https://doi.org/10.1111/jipb.13101). 1180
1181
1182

148. Zhang, W.; Zhi, H.; Tang, S.; Zhang, H.; Sui, Y.; Jia, G.; Wu, C.; Diao, X. Identification of No Pollen 1 Provides a Candidate Gene for Heterosis Utilization in Foxtail Millet (*Setaria Italica* L.). *Crop J.* **2021**, *9*, 1309–1319, doi:[10.1016/j.cj.2021.03.018](https://doi.org/10.1016/j.cj.2021.03.018). 1183
1184
1185

149. Part B Notifications (Experimental Releases) – GM Plants Available online: https://webgate.ec.europa.eu/fip/GMO_Registers/GMO_Part_B_Plants.php (accessed on 24 November 2022). 1186
1187

150. Capdeville, N.; Schindele, P.; Puchta, H. Getting Better All the Time – Recent Progress in the Development of CRISPR/Cas-Based Tools for Plant Genome Engineering. *Curr. Opin. Biotechnol.* **2023**, *79*, 102854, doi:[10.1016/j.copbio.2022.102854](https://doi.org/10.1016/j.copbio.2022.102854). 1188
1189
1190

151. Zetsche, B.; Gootenberg, J.S.; Abudayyeh, O.O.; Slaymaker, I.M.; Makarova, K.S.; Essletzbichler, P.; Volz, S.E.; Joung, J.; van der Oost, J.; Regev, A.; et al. Cpf1 Is a Single RNA-Guided Endonuclease of a Class 2 CRISPR-Cas System. *Cell* **2015**, *163*, 759–771, doi:10.1016/j.cell.2015.09.038. 1191
1192
1193

152. Moreno-Mateos, M.A.; Fernandez, J.P.; Rouet, R.; Vejnar, C.E.; Lane, M.A.; Mis, E.; Khokha, M.K.; Doudna, J.A.; Giraldez, A.J. CRISPR-Cpf1 Mediates Efficient Homology-Directed Repair and Temperature-Controlled Genome Editing. *Nat. Commun.* **2017**, *8*, 2024, doi:10.1038/s41467-017-01836-2. 1194
1195
1196

153. Čermák, T.; Curtin, S.J.; Gil-Humanes, J.; Čegan, R.; Kono, T.J.Y.; Konečná, E.; Belanto, J.J.; Starker, C.G.; Mathre, J.W.; Greenstein, R.L.; et al. A Multipurpose Toolkit to Enable Advanced Genome Engineering in Plants. *Plant Cell* **2017**, *29*, 1196–1217, doi:10.1105/tpc.16.00922. 1197
1198
1199

154. Ding, X.; Yu, L.; Chen, L.; Li, Y.; Zhang, J.; Sheng, H.; Ren, Z.; Li, Y.; Yu, X.; Jin, S.; et al. Recent Progress and Future Prospect of CRISPR/Cas-Derived Transcription Activation (CRISPRa) System in Plants. *Cells* **2022**, *11*, 3045, doi:10.3390/cells11193045. 1200
1201
1202

155. Tanenbaum, M.E.; Gilbert, L.A.; Qi, L.S.; Weissman, J.S.; Vale, R.D. A Protein-Tagging System for Signal Amplification in Gene Expression and Fluorescence Imaging. *Cell* **2014**, *159*, 635–646, doi:10.1016/j.cell.2014.09.039. 1203
1204

156. Papikian, A.; Liu, W.; Gallego-Bartolomé, J.; Jacobsen, S.E. Site-Specific Manipulation of *Arabidopsis* Loci Using CRISPR-Cas9 SunTag Systems. *Nat. Commun.* **2019**, *10*, 729, doi:10.1038/s41467-019-08736-7. 1205
1206

157. Shi, J.; Gao, H.; Wang, H.; Lafitte, H.R.; Archibald, R.L.; Yang, M.; Hakimi, S.M.; Mo, H.; Habben, J.E. ARGOS8 Variants Generated by CRISPR-Cas9 Improve Maize Grain Yield under Field Drought Stress Conditions. *Plant Biotechnol. J.* **2017**, *15*, 207–216, doi:10.1111/pbi.12603. 1207
1208
1209

158. Zhang, A.; Liu, Y.; Wang, F.; Li, T.; Chen, Z.; Kong, D.; Bi, J.; Zhang, F.; Luo, X.; Wang, J.; et al. Enhanced Rice Salinity Tolerance via CRISPR/Cas9-Targeted Mutagenesis of the OsRR22 Gene. *Mol. Breed.* **2019**, *39*, 47, doi:10.1007/s11032-019-0954-y. 1210
1211
1212

159. Han, X.; Chen, Z.; Li, P.; Xu, H.; Liu, K.; Zha, W.; Li, S.; Chen, J.; Yang, G.; Huang, J.; et al. Development of Novel Rice Germplasm for Salt-Tolerance at Seedling Stage Using CRISPR-Cas9. *Sustainability* **2022**, *14*, 2621, doi:10.3390/su14052621. 1213
1214
1215

160. Chen, J.-H.; Chen, S.-T.; He, N.-Y.; Wang, Q.-L.; Zhao, Y.; Gao, W.; Guo, F.-Q. Nuclear-Encoded Synthesis of the D1 Subunit of Photosystem II Increases Photosynthetic Efficiency and Crop Yield. *Nat. Plants* **2020**, *6*, 570–580, doi:10.1038/s41477-020-0629-z. 1216
1217
1218

161. Yu, W.; Wang, L.; Zhao, R.; Sheng, J.; Zhang, S.; Li, R.; Shen, L. Knockout of SlMAPK3 Enhances Tolerance to Heat Stress Involving ROS Homeostasis in Tomato Plants. *BMC Plant Biol.* **2019**, *19*, 354, doi:10.1186/s12870-019-1939-z. 1219
1220

162. Miao, C.; Xiao, L.; Hua, K.; Zou, C.; Zhao, Y.; Bressan, R.A.; Zhu, J.-K. Mutations in a Subfamily of Abscisic Acid Receptor Genes Promote Rice Growth and Productivity. *Proc. Natl. Acad. Sci.* **2018**, *115*, 6058–6063, doi:10.1073/pnas.1804774115. 1221
1222
1223

163. Abdellatif, I.M.Y.; Yuan, S.; Na, R.; Yoshihara, S.; Hamada, H.; Suzuki, T.; Ezura, H.; Miura, K. Functional Characterization of Tomato Phytochrome A and B1B2 Mutants in Response to Heat Stress. *Int. J. Mol. Sci.* **2022**, *23*, 1681, doi:10.3390/ijms23031681. 1224
1225
1226

164. Xiang, D.-J.; Man, L.-L.; Cao, S.; Liu, P.; Li, Z.-G.; Wang, X.-D. Heterologous Expression of an *Agropyron Cristatum* SnRK2 Protein Kinase Gene (AcSnRK2.11) Increases Freezing Tolerance in Transgenic Yeast and Tobacco. *3 Biotech* **2020**, *10*, 209, doi:10.1007/s13205-020-02203-7. 1227
1228
1229

165. Ma, Y.; Dai, X.; Xu, Y.; Luo, W.; Zheng, X.; Zeng, D.; Pan, Y.; Lin, X.; Liu, H.; Zhang, D.; et al. COLD1 Confers Chilling Tolerance in Rice. *Cell* **2015**, *160*, 1209–1221, doi:10.1016/j.cell.2015.01.046. 1230
1231

166. Zeng, Y.; Wen, J.; Zhao, W.; Wang, Q.; Huang, W. Rational Improvement of Rice Yield and Cold Tolerance by 1232
Editing the Three Genes OsPIN5b, GS3, and OsMYB30 With the CRISPR–Cas9 System. *Front. Plant Sci.* **2020**, *10*, 1233
doi:<https://doi.org/10.3389/fpls.2019.01663>. 1234

167. Ding, Y.; Lv, J.; Shi, Y.; Gao, J.; Hua, J.; Song, C.; Gong, Z.; Yang, S. EGR2 Phosphatase Regulates OST1 Kinase 1235
Activity and Freezing Tolerance in Arabidopsis. *EMBO J.* **2019**, *38*, e99819, doi:10.15252/embj.201899819. 1236

168. Liu, Z.; Jia, Y.; Ding, Y.; Shi, Y.; Li, Z.; Guo, Y.; Gong, Z.; Yang, S. Plasma Membrane CRPK1-Mediated Phosphor- 1237
ylation of 14-3-3 Proteins Induces Their Nuclear Import to Fine-Tune CBF Signaling during Cold Response. *Mol. 1238
Cell* **2017**, *66*, 117–128.e5, doi:10.1016/j.molcel.2017.02.016. 1239

169. Lv, A.; Fan, N.; Xie, J.; Yuan, S.; An, Y.; Zhou, P. Expression of CdDHN4, a Novel YSK2-Type Dehydrin Gene from 1240
Bermudagrass, Responses to Drought Stress through the ABA-Dependent Signal Pathway. *Front. Plant Sci.* **2017**, *8*, 1241
doi:<https://doi.org/10.3389/fpls.2017.00748>. 1242

170. Shanmugam, S.; Boyett, V.A.; Khodakovskaya, M. Enhancement of Drought Tolerance in Rice by Silencing of the 1243
OsSYT-5 Gene. *PLOS ONE* **2021**, *16*, e0258171, doi:10.1371/journal.pone.0258171. 1244

171. Santosh Kumar, V.V.; Verma, R.K.; Yadav, S.K.; Yadav, P.; Watts, A.; Rao, M.V.; Chinnusamy, V. CRISPR-Cas9 1245
Mediated Genome Editing of Drought and Salt Tolerance (OsDST) Gene in Indica Mega Rice Cultivar MTU1010. 1246
Physiol. Mol. Biol. Plants **2020**, *26*, 1099–1110, doi:10.1007/s12298-020-00819-w. 1247

172. Daszkowska-Golec, A.; Karcz, J.; Plociniczak, T.; Sitko, K.; Szarejko, I. Cuticular Waxes—A Shield of Barley Mu- 1248
tant in CBP20 (Cap-Binding Protein 20) Gene When Struggling with Drought Stress. *Plant Sci.* **2020**, *300*, 110593, 1249
doi:10.1016/j.plantsci.2020.110593. 1250

173. Ju, H.; Li, D.; Li, D.; Yang, X.; Liu, Y. Overexpression of ZmDHN11 Could Enhance Transgenic Yeast and Tobacco 1251
Tolerance to Osmotic Stress. *Plant Cell Rep.* **2021**, *40*, 1723–1733, doi:10.1007/s00299-021-02734-0. 1252

174. Alam, M.S.; Kong, J.; Tao, R.; Ahmed, T.; Alamin, M.; Alotaibi, S.S.; Abdelsalam, N.R.; Xu, J.-H. CRISPR/Cas9 1253
Mediated Knockout of the OsbHLH024 Transcription Factor Improves Salt Stress Resistance in Rice (*Oryza Sativa* 1254
L.). *Plants* **2022**, *11*, 1184, doi:10.3390/plants11091184. 1255

175. Vlčko, T.; Ohnoutková, L. Allelic Variants of CRISPR/Cas9 Induced Mutation in an Inositol Trisphosphate 5/6 1256
Kinase Gene Manifest Different Phenotypes in Barley. *Plants* **2020**, *9*, 195, doi:10.3390/plants9020195. 1257

176. Niu, Y.; Xiang, Y. An Overview of Biomembrane Functions in Plant Responses to High-Temperature Stress. *Front. 1258
Plant Sci.* **2018**, *9*, doi:<https://doi.org/10.3389/fpls.2018.00915>. 1259

177. Damaris, R.N.; Yang, P. Protein Phosphorylation Response to Abiotic Stress in Plants. In *Plant Phosphoproteomics: 1260
Methods and Protocols*; Wu, X.N., Ed.; Methods in Molecular Biology; Springer US: New York, NY, 2021; pp. 17–43 1261
ISBN 978-1-07-161625-3. 1262

178. Zhang, H.; Li, W.; Mao, X.; Jing, R.; Jia, H. Differential Activation of the Wheat SnRK2 Family by Abiotic Stresses. 1263
Front. Plant Sci. **2016**, *7*, doi:<https://doi.org/10.3389/fpls.2016.00420>. 1264

179. Tian, S.; Mao, X.; Zhang, H.; Chen, S.; Zhai, C.; Yang, S.; Jing, R. Cloning and Characterization of TaSnRK2.3, a 1265
Novel SnRK2 Gene in Common Wheat. *J. Exp. Bot.* **2013**, *64*, 2063–2080, doi:10.1093/jxb/ert072. 1266

180. Jia, Q.; Kong, D.; Li, Q.; Sun, S.; Song, J.; Zhu, Y.; Liang, K.; Ke, Q.; Lin, W.; Huang, J. The Function of Inositol 1267
Phosphatases in Plant Tolerance to Abiotic Stress. *Int. J. Mol. Sci.* **2019**, *20*, 3999, doi:10.3390/ijms20163999. 1268

181. Chan, K.X.; Mabbitt, P.D.; Phua, S.Y.; Mueller, J.W.; Nisar, N.; Gigolashvili, T.; Stroehner, E.; Grassl, J.; Arlt, W.; 1269
Estavillo, G.M.; et al. Sensing and Signaling of Oxidative Stress in Chloroplasts by Inactivation of the SAL1 Phos- 1270
phoadenosine Phosphatase. *Proc. Natl. Acad. Sci.* **2016**, *113*, E4567–E4576, doi:10.1073/pnas.1604936113. 1271

182. Guo, J.; Sun, B.; He, H.; Zhang, Y.; Tian, H.; Wang, B. Current Understanding of BHLH Transcription Factors in 1272
Plant Abiotic Stress Tolerance. *Int. J. Mol. Sci.* **2021**, *22*, 4921, doi:10.3390/ijms22094921. 1273

183. Anzalone, A.V.; Koblan, L.W.; Liu, D.R. Genome Editing with CRISPR–Cas Nucleases, Base Editors, Transposases and Prime Editors. *Nat. Biotechnol.* **2020**, *38*, 824–844, doi:10.1038/s41587-020-0561-9. 1274
1275

184. Zhu, H.; Li, C.; Gao, C. Applications of CRISPR–Cas in Agriculture and Plant Biotechnology. *Nat. Rev. Mol. Cell Biol.* **2020**, *21*, 661–677, doi:10.1038/s41580-020-00288-9. 1276
1277

185. Hua, K.; Han, P.; Zhu, J.-K. Improvement of Base Editors and Prime Editors Advances Precision Genome Engineering in Plants. *Plant Physiol.* **2022**, *188*, 1795–1810, doi:10.1093/plphys/kiab591. 1278
1279

186. Puchta, H.; Jiang, J.; Wang, K.; Zhao, Y. Updates on Gene Editing and Its Applications. *Plant Physiol.* **2022**, *188*, 1725–1730, doi:10.1093/plphys/kiac032. 1280
1281

187. Villalobos-López, M.A.; Arroyo-Becerra, A.; Quintero-Jiménez, A.; Iturriaga, G. Biotechnological Advances to Improve Abiotic Stress Tolerance in Crops. *Int. J. Mol. Sci.* **2022**, *23*, 12053, doi:10.3390/ijms231912053. 1282
1283

188. Lundqvist, A. Genetics of Self-Incompatibility in *Festuca Pratensis* Huds. *Hereditas* **1955**, *41*, 518–520, doi:10.1111/j.1601-5223.1955.tb03007.x. 1284
1285

189. Cornish, M.A.; Hayward, M.D.; Lawrence, M.J. Self-Incompatibility in Ryegrass. *Heredity* **1979**, *43*, 95–106, doi:10.1038/hdy.1979.63. 1286
1287

190. Manzanares, C.; Barth, S.; Thorogood, D.; Byrne, S.L.; Yates, S.; Czaban, A.; Asp, T.; Yang, B.; Studer, B. A Gene Encoding a DUF247 Domain Protein Cosegregates with the S Self-Incompatibility Locus in Perennial Ryegrass. *Mol. Biol. Evol.* **2016**, *33*, 870–884, doi:10.1093/molbev/msv335. 1288
1289

191. Herridge, R.; McCourt, T.; Jacobs, J.M.E.; Mace, P.; Brownfield, L.; Macknight, R. Identification of the Genes at S and Z Reveals the Molecular Basis and Evolution of Grass Self-Incompatibility. *Front. Plant Sci.* **2022**, *13*, doi:https://doi.org/10.3389/fpls.2022.1011299. 1291
1292

192. Ye, M.; Peng, Z.; Tang, D.; Yang, Z.; Li, D.; Xu, Y.; Zhang, C.; Huang, S. Generation of Self-Compatible Diploid Potato by Knockout of S-RNase. *Nat. Plants* **2018**, *4*, 651–654, doi:10.1038/s41477-018-0218-6. 1294
1295

193. Singh, M.; Kumar, M.; Albertsen, M.C.; Young, J.K.; Cigan, A.M. Concurrent Modifications in the Three Homeologs of Ms45 Gene with CRISPR-Cas9 Lead to Rapid Generation of Male Sterile Bread Wheat (*Triticum Aestivum* L.). *Plant Mol. Biol.* **2018**, *97*, 371–383, doi:10.1007/s11103-018-0749-2. 1296
1297

194. Okada, A.; Arndell, T.; Borisjuk, N.; Sharma, N.; Watson-Haigh, N.S.; Tucker, E.J.; Baumann, U.; Langridge, P.; Whitford, R. CRISPR/Cas9-Mediated Knockout of Ms1 Enables the Rapid Generation of Male-Sterile Hexaploid Wheat Lines for Use in Hybrid Seed Production. *Plant Biotechnol. J.* **2019**, *17*, 1905–1913, doi:10.1111/pbi.13106. 1299
1300

195. Khanday, I.; Skinner, D.; Yang, B.; Mercier, R.; Sundaresan, V. A Male-Expressed Rice Embryogenic Trigger Redirected for Asexual Propagation through Seeds. *Nature* **2019**, *565*, 91–95, doi:10.1038/s41586-018-0785-8. 1302
1303

196. Hickey, L.T.; N. Hafeez, A.; Robinson, H.; Jackson, S.A.; Leal-Bertioli, S.C.M.; Tester, M.; Gao, C.; Godwin, I.D.; Hayes, B.J.; Wulff, B.B.H. Breeding Crops to Feed 10 Billion. *Nat. Biotechnol.* **2019**, *37*, 744–754, doi:10.1038/s41587-019-0152-9. 1304
1305

197. Gao, C. Genome Engineering for Crop Improvement and Future Agriculture. *Cell* **2021**, *184*, 1621–1635, doi:10.1016/j.cell.2021.01.005. 1307
1308

198. González, F.G.; Capella, M.; Ribichich, K.F.; Curín, F.; Giacomelli, J.I.; Ayala, F.; Watson, G.; Otegui, M.E.; Chan, R.L. Field-Grown Transgenic Wheat Expressing the Sunflower Gene HaHB4 Significantly Outyields the Wild Type. *J. Exp. Bot.* **2019**, *70*, 1669–1681, doi:10.1093/jxb/erz037. 1309
1310

199. Argentina First to Market with Drought-Resistant GM Wheat. *Nat. Biotechnol.* **2021**, *39*, 652–652, doi:10.1038/s41587-021-00963-y. 1312
1313

200. HB4 Wheat | GM Approval Database- ISAAA.Org Available online: <https://www.isaaa.org/gmapprovaldatabase/event/default.asp?EventID=574> (accessed on 30 November 2022). 1314
1315

201. Rice, E.A.; Khandelwal, A.; Creelman, R.A.; Griffith, C.; Ahrens, J.E.; Taylor, J.P.; Murphy, L.R.; Manjunath, S.; Thompson, R.L.; Lingard, M.J.; et al. Expression of a Truncated ATHB17 Protein in Maize Increases Ear Weight at Silking. *PLOS ONE* **2014**, *9*, e94238, doi:10.1371/journal.pone.0094238. 1316
1317
1318

202. Leibman, M.; Shryock, J.J.; Clements, M.J.; Hall, M.A.; Loida, P.J.; McClerren, A.L.; McKiness, Z.P.; Phillips, J.R.; Rice, E.A.; Stark, S.B. Comparative Analysis of Maize (*Zea Mays*) Crop Performance: Natural Variation, Incremental Improvements and Economic Impacts. *Plant Biotechnol. J.* **2014**, *12*, 941–950, doi:10.1111/pbi.12202. 1319
1320
1321

203. Nemali, K.S.; Bonin, C.; Dohleman, F.G.; Stephens, M.; Reeves, W.R.; Nelson, D.E.; Castiglioni, P.; Whitsel, J.E.; Sammons, B.; Silady, R.A.; et al. Physiological Responses Related to Increased Grain Yield under Drought in the First Biotechnology-Derived Drought-Tolerant Maize. *Plant Cell Environ.* **2015**, *38*, 1866–1880, doi:10.1111/pce.12446. 1322
1323
1324

204. EFSA Panel on Genetically Modified Organisms (GMO); Arpaia, S.; Birch, A.N.; Chesson, A.; du Jardin, P.; Gathmann, A.; Gropp, J.; Herman, L.; Hoen-Sorteberg, H.-G.; Jones, H.; et al. Scientific Opinion on an Application (EFSA-GMO-NL-2009-70) for the Placing on the Market of Genetically Modified Drought Tolerant Maize MON 87460 for Food and Feed Uses, Import and Processing under Regulation (EC) No 1829/2003 from Monsanto. *EFSA J.* **2012**, *10*, 1325
1326
1327
1328
1329

205. EFSA Panel on Genetically Modified Organisms (GMO); Naegeli, H.; Birch, A.N.; Casacuberta, J.; De Schrijver, A.; Gralak, M.A.; Guerche, P.; Jones, H.; Manachini, B.; Messéan, A.; et al. Assessment of Genetically Modified Maize MON 87411 for Food and Feed Uses, Import and Processing, under Regulation (EC) No 1829/2003 (Application EFSA-GMO-NL-2015-124). *EFSA J.* **2018**, *16*, e05310, doi:10.2903/j.efsa.2018.5310. 1330
1331
1332
1333

206. Wang, Z.-Y.; Ge, Y. Recent Advances in Genetic Transformation of Forage and Turf Grasses. *Vitro Cell. Dev. Biol. - Plant* **2006**, *42*, 1–18, doi:10.1079/IVP2005726. 1334
1335

207. Creeping Bentgrass (*Agrostis Stolonifera*) GM Events | GM Approval Database - ISAAA.Org Available online: <https://www.isaaa.org/gmapprovaldatabase/crop/default.asp?CropID=8&Crop=Creeping%20Bentgrass> (accessed on 2 December 2022). 1336
1337
1338

208. Brookes, G. Twenty-One Years of Using Insect Resistant (GM) Maize in Spain and Portugal: Farm-Level Economic and Environmental Contributions. *GM Crops Food* **2019**, *10*, 90–101, doi:10.1080/21645698.2019.1614393. 1339
1340

209. Brookes, G. Farm Income and Production Impacts from the Use of Genetically Modified (GM) Crop Technology 1996–2020. *GM Crops Food* **2022**, *13*, 171–195, doi:10.1080/21645698.2022.2105626. 1341
1342

210. National Academies of Sciences, Engineering, and Medicine; Division on Earth and Life Studies; Board on Agriculture and Natural Resources; Committee on Genetically Engineered Crops: Past Experience and Future Prospects *Genetically Engineered Crops: Experiences and Prospects*; National Academies Press (US): Washington (DC), 2016; ISBN 978-0-309-43738-7. 1343
1344
1345
1346

211. Shukla, V.K.; Doyon, Y.; Miller, J.C.; DeKelver, R.C.; Moehle, E.A.; Worden, S.E.; Mitchell, J.C.; Arnold, N.L.; Gopal, S.; Meng, X.; et al. Precise Genome Modification in the Crop Species *Zea Mays* Using Zinc-Finger Nucleases. *Nature* **2009**, *459*, 437–441, doi:10.1038/nature07992. 1347
1348
1349

212. Urnov, F.D.; Rebar, E.J.; Holmes, M.C.; Zhang, H.S.; Gregory, P.D. Genome Editing with Engineered Zinc Finger Nucleases. *Nat. Rev. Genet.* **2010**, *11*, 636–646, doi:10.1038/nrg2842. 1350
1351

213. Sánchez-León, S.; Gil-Humanes, J.; Ozuna, C.V.; Giménez, M.J.; Sousa, C.; Voytas, D.F.; Barro, F. Low-Gluten, Nontransgenic Wheat Engineered with CRISPR/Cas9. *Plant Biotechnol. J.* **2018**, *16*, 902–910, doi:10.1111/pbi.12837. 1352
1353

214. Li, R.; Li, R.; Li, X.; Fu, D.; Zhu, B.; Tian, H.; Luo, Y.; Zhu, H. Multiplexed CRISPR/Cas9-Mediated Metabolic Engineering of γ -Aminobutyric Acid Levels in *Solanum Lycopersicum*. *Plant Biotechnol. J.* **2018**, *16*, 415–427, doi:10.1111/pbi.12781. 1354
1355
1356

215. Li, J.; Scarano, A.; Gonzalez, N.M.; D'Orso, F.; Yue, Y.; Nemeth, K.; Saalbach, G.; Hill, L.; de Oliveira Martins, C.; Moran, R.; et al. Biofortified Tomatoes Provide a New Route to Vitamin D Sufficiency. *Nat. Plants* **2022**, *8*, 611–616, doi:10.1038/s41477-022-01154-6. 1357
1358
1359

216. Waltz, E. GABA-Enriched Tomato Is First CRISPR-Edited Food to Enter Market. *Nat. Biotechnol.* **2021**, *40*, 9–11, doi:10.1038/d41587-021-00026-2. 1360
1361

217. EFSA Panel on Genetically Modified Organisms (GMO); Arpaia, S.; Birch, A.N.; Chesson, A.; du Jardin, P.; Gathmann, A.; Gropp, J.; Herman, L.; Hoen-Sorteberg, H.-G.; Jones, H.; et al. Scientific Opinion Addressing the Safety Assessment of Plants Developed Using Zinc Finger Nuclease 3 and Other Site-Directed Nucleases with Similar Function. *EFSA J.* **2012**, *10*, 2943, doi:10.2903/j.efsa.2012.2943. 1362
1363
1364
1365

218. Purnhagen, K.P.; Kok, E.; Kleter, G.; Schebesta, H.; Visser, R.G.F.; Wesseler, J. EU Court Casts New Plant Breeding Techniques into Regulatory Limbo. *Nat. Biotechnol.* **2018**, *36*, 799–800, doi:10.1038/nbt.4251. 1366
1367

219. Urnov, F.D.; Ronald, P.C.; Carroll, D. A Call for Science-Based Review of the European Court's Decision on Gene-Edited Crops. *Nat. Biotechnol.* **2018**, *36*, 800–802, doi:10.1038/nbt.4252. 1368
1369

220. Christiansen, A.T.; Andersen, M.M.; Kappel, K. Are Current EU Policies on GMOs Justified? *Transgenic Res.* **2019**, *28*, 267–286, doi:10.1007/s11248-019-00120-x. 1370
1371

221. Wasmer, M. Roads Forward for European GMO Policy—Uncertainties in Wake of ECJ Judgment Have to Be Mitigated by Regulatory Reform. *Front. Bioeng. Biotechnol.* **2019**, *7*, doi:https://doi.org/10.3389/fbioe.2019.00132. 1372
1373

222. Schulman, A.H.; Oksman-Caldentey, K.; Teeri, T.H. European Court of Justice Delivers No Justice to Europe on Genome-edited Crops. *Plant Biotechnol. J.* **2020**, *18*, 8–10, doi:10.1111/pbi.13200. 1374
1375

223. European Network of GMO Laboratories (ENG) *Detection of Food and Feed Plant Products Obtained by New Mutagenesis Techniques*; 2019; 1376
1377

224. Maize DP-915635-4 Available online: <https://open.efsa.europa.eu> (accessed on 5 December 2022). 1378

225. EFSA Panel on Genetically Modified Organisms (GMO); Naegeli, H.; Bresson, J.-L.; Dalmay, T.; Dewhurst, I.C.; Epstein, M.M.; Firbank, L.G.; Guerche, P.; Hejatko, J.; Moreno, F.J.; et al. Applicability of the EFSA Opinion on Site-Directed Nucleases Type 3 for the Safety Assessment of Plants Developed Using Site-Directed Nucleases Type 1 and 2 and Oligonucleotide-Directed Mutagenesis. *EFSA J.* **2020**, *18*, e06299, doi:10.2903/j.efsa.2020.6299. 1379
1380
1381
1382

226. Rostoks, N. Implications of the EFSA Scientific Opinion on Site Directed Nucleases 1 and 2 for Risk Assessment of Genome-Edited Plants in the EU. *Agronomy* **2021**, *11*, 572, doi:10.3390/agronomy11030572. 1383
1384

227. EFSA Panel on Genetically Modified Organisms (GMO); Mullins, E.; Bresson, J.-L.; Dalmay, T.; Dewhurst, I.C.; Epstein, M.M.; Firbank, L.G.; Guerche, P.; Hejatko, J.; Moreno, F.J.; et al. Updated Scientific Opinion on Plants Developed through Cisgenesis and Intragenesis. *EFSA J.* **2022**, *20*, e07621, doi:10.2903/j.efsa.2022.7621. 1385
1386
1387

228. EFSA Panel on Genetically Modified Organisms (GMO); Mullins, E.; Bresson, J.-L.; Dalmay, T.; Dewhurst, I.C.; Epstein, M.M.; Firbank, L.G.; Guerche, P.; Hejatko, J.; Moreno, F.J.; et al. Criteria for Risk Assessment of Plants Produced by Targeted Mutagenesis, Cisgenesis and Intragenesis. *EFSA J.* **2022**, *20*, e07618, doi:10.2903/j.efsa.2022.7618. 1388
1389
1390
1391

229. Bratlie, S.; Halvorsen, K.; Myskja, B.K.; Mellegård, H.; Bjorvatn, C.; Frost, P.; Heiene, G.; Hofmann, B.; Holst-Jensen, A.; Holst-Larsen, T.; et al. A Novel Governance Framework for GMO. *EMBO Rep.* **2019**, *20*, e47812, doi:10.15252/embr.201947812. 1392
1393
1394

230. Kearns, P.W.E.; Kleter, G.A.; Bergmans, H.E.N.; Kuiper, H.A. Biotechnology and Biosafety Policy at OECD: Future Trends. *Trends Biotechnol.* **2021**, *39*, 965–969, doi:10.1016/j.tibtech.2021.03.001. 1395
1396

231. Gould, F.; Amasino, R.M.; Brossard, D.; Buell, C.R.; Dixon, R.A.; Falck-Zepeda, J.B.; Gallo, M.A.; Giller, K.E.; Glenna, L.L.; Griffin, T.; et al. Toward Product-Based Regulation of Crops. *Science* **2022**, *377*, 1051–1053, doi:10.1126/science.abo3034. 1397
1398
1399

232. Goberna, M.F.; Whelan, A.I.; Godoy, P.; Lewi, D.M. Genomic Editing: The Evolution in Regulatory Management Accompanying Scientific Progress. *Front. Bioeng. Biotechnol.* **2022**, *10*, doi:https://doi.org/10.3389/fbioe.2022.835378. 1400
1401

233. Entine, J.; Felipe, M.S.S.; Groenewald, J.-H.; Kershen, D.L.; Lema, M.; McHughen, A.; Nepomuceno, A.L.; Ohsawa, R.; Ordonio, R.L.; Parrott, W.A.; et al. Regulatory Approaches for Genome Edited Agricultural Plants in Select Countries and Jurisdictions around the World. *Transgenic Res.* **2021**, *30*, 551–584, doi:10.1007/s11248-021-00257-8. 1402
1403
1404

234. Rostoks, N. Unpublished. 1405

235. Posselt, U.K. Breeding Methods in Cross-Pollinated Species. In *Fodder Crops and Amenity Grasses*; Boller, B., Posselt, U.K., Veronesi, F., Eds.; Handbook of Plant Breeding; Springer: New York, NY, 2010; pp. 39–87 ISBN 978-1-4419-0760-8. 1406
1407
1408

236. Rognli, O.A.; Pecetti, L.; Kovi, M.R.; Annicchiarico, P. Grass and Legume Breeding Matching the Future Needs of European Grassland Farming. *Grass Forage Sci.* **2021**, *76*, 175–185, doi:10.1111/gfs.12535. 1409
1410

237. Barre, P.; Asp, T.; Byrne, S.; Casler, M.; Faville, M.; Rognli, O.A.; Roldan-Ruiz, I.; Skøt, L.; Ghesquière, M. Genomic Prediction of Complex TraitsComplex Traits in Forage Plants Species: Perennial Grasses Case. In *Genomic Prediction of Complex Traits: Methods and Protocols*; Ahmadi, N., Bartholomé, J., Eds.; Methods in Molecular Biology; Springer US: New York, NY, 2022; pp. 521–541 ISBN 978-1-07-162205-6. 1411
1412
1413
1414

238. Meyer, R.S.; DuVal, A.E.; Jensen, H.R. Patterns and Processes in Crop Domestication: An Historical Review and Quantitative Analysis of 203 Global Food Crops. *New Phytol.* **2012**, *196*, 29–48, doi:10.1111/j.1469-8137.2012.04253.x. 1415
1416

239. Swinnen, G.; Goossens, A.; Pauwels, L. Lessons from Domestication: Targeting Cis-Regulatory Elements for Crop Improvement. *Trends Plant Sci.* **2016**, *21*, 506–515, doi:10.1016/j.tplants.2016.01.014. 1417
1418

240. Jonavičienė, K.; Paplauskienė, V.; Brazauskas, G. Isozymes and ISSR Markers as a Tool for the Assessment of Genetic Diversity in *Phleum* Spp. *Žemdirb. Agric.* **2009**, *96*, 47–57. 1419
1420

241. Brozynska, M.; Furtado, A.; Henry, R.J. Genomics of Crop Wild Relatives: Expanding the Gene Pool for Crop Improvement. *Plant Biotechnol. J.* **2016**, *14*, 1070–1085, doi:10.1111/pbi.12454. 1421
1422

242. Blanco-Pastor, J.L.; Manel, S.; Barre, P.; Roschanski, A.M.; Willner, E.; Dehmer, K.J.; Hegarty, M.; Muylle, H.; Rutting, T.; Roldán-Ruiz, I.; et al. Pleistocene Climate Changes, and Not Agricultural Spread, Accounts for Range Expansion and Admixture in the Dominant Grassland Species *Lolium Perenne* L. *J. Biogeogr.* **2019**, *46*, 1451–1465, doi:10.1111/jbi.13587. 1423
1424
1425
1426

243. Bachmann-Pfabe, S.; Willner, E.; Dehmer, K.J. Ex-Situ Evaluations of *Lolium Perenne* L. Ecotypes Collected in Bulgaria, Croatia, Spain and Ireland Reveal Valuable Breeding Material. *Genet. Resour. Crop Evol.* **2018**, *65*, 1423–1439, doi:10.1007/s10722-018-0623-8. 1427
1428
1429

244. Jaškūnė, K.; Kemešytė, V.; Aleliūnas, A.; Statkevičiūtė, G. Genome-Wide Markers for Seed Yield and Disease Resistance in Perennial Ryegrass. *Crop J.* **2022**, *10*, 508–514, doi:10.1016/j.cj.2021.07.005. 1430
1431

245. Gylstrøm, K.H. Phenotypic Characterization and Ploidy Level Determination of Perennial Ryegrass (*Lolium Perenne* L.) Gene Bank Accessions and Cultivars. Master thesis, Norwegian University of Life Sciences, Ås, 2020. 1432
1433

246. Akinroluyo, O.K.; Urbanavičiūtė, I.; Jaškūnė, K.; Kemešytė, V.; Statkevičiūtė, G. Differences in Salt Tolerance between Diploid and Autotetraploid Lines of *Lolium Multiflorum* at the Germination and Vegetative Stages. *Žemdirb. Agric.* **2019**, *106*, 329–336, doi:10.13080/z-a.2019.106.042. 1434
1435
1436

247. Akinroluyo, O.K.; Jaškūnė, K.; Kemešytė, V.; Statkevičiutė, G. Drought Stress Response of Westerwolths Ryegrass (*Lolium Multiflorum* Ssp. *Multiflorum*) Cultivars Differing in Their Ploidy Level. *Žemdirb. Agric.* **2020**, *107*, 161–170, doi:10.13080/z-a.2020.107.021. 1437
1438
1439

248. Rauf, S.; Ortiz, R.; Malinowski, D.P.; Clarindo, W.R.; Kainat, W.; Shehzad, M.; Waheed, U.; Hassan, S.W. Induced Polyploidy: A Tool for Forage Species Improvement. *Agriculture* **2021**, *11*, 210, doi:10.3390/agriculture11030210. 1440
1441

249. Nagy, I.; Veeckman, E.; Liu, C.; Bel, M.V.; Vandepoele, K.; Jensen, C.S.; Ruttink, T.; Asp, T. Chromosome-Scale Assembly and Annotation of the Perennial Ryegrass Genome. *BMC Genomics* **2022**, *23*, 505, doi:10.1186/s12864-022-08697-0. 1442
1443
1444

250. Smith, K.F.; Spangenberg, G. Considerations for Managing Agricultural Co-Existence between Transgenic and Non-Transgenic Cultivars of Outcrossing Perennial Forage Plants in Dairy Pastures. *Agronomy* **2016**, *6*, 59, doi:10.3390/agronomy6040059. 1445
1446
1447

251. Badenhorst, P.E.; Smith, K.F.; Spangenberg, G. Development of a Molecular Breeding Strategy for the Integration of Transgenic Traits in Outcrossing Perennial Grasses. *Agronomy* **2016**, *6*, 56, doi:10.3390/agronomy6040056. 1448
1449

252. Bier, E. Gene Drives Gaining Speed. *Nat. Rev. Genet.* **2022**, *23*, 5–22, doi:10.1038/s41576-021-00386-0. 1450

253. Sampoux, J.-P.; Baudouin, P.; Bayle, B.; Béguier, V.; Bourdon, P.; Chosson, J.-F.; Deneufbourg, F.; Galbrun, C.; Ghesquière, M.; Noël, D.; et al. Breeding Perennial Grasses for Forage Usage: An Experimental Assessment of Trait Changes in Diploid Perennial Ryegrass (*Lolium Perenne* L.) Cultivars Released in the Last Four Decades. *Field Crops Res.* **2011**, *123*, 117–129, doi:10.1016/j.fcr.2011.05.007. 1451
1452
1453
1454

254. McDonagh, J.; O'Donovan, M.; McEvoy, M.; Gilliland, T.J. Genetic Gain in Perennial Ryegrass (*Lolium Perenne*) Varieties 1973 to 2013. *Euphytica* **2016**, *212*, 187–199, doi:10.1007/s10681-016-1754-7. 1455
1456

255. Herridge, R.P.; Macknight, R.C.; Brownfield, L.R. Prospects for F1 Hybrid Production in Ryegrass. *N. Z. J. Agric. Res.* **2020**, *63*, 405–415, doi:10.1080/00288233.2018.1559867. 1457
1458

256. Hojsgaard, D.; Hörandl, E. The Rise of Apomixis in Natural Plant Populations. *Front. Plant Sci.* **2019**, *10*. 1459
1460