



Article

On the significance of parameters in the Choice and Collection schemata in the 2nd order Peano arithmetic

Vladimir Kanovei ^{*,†}  and Vassily Lyubetsky ^{*,†} 

Institute for Information Transmission Problems of the Russian Academy of Sciences (Kharkevich Institute),
127051 Moscow, Russia

* Correspondence: kanovei@iitp.ru (V.K.); lyubetsk@iitp.ru (V.L.)

† These authors contributed equally to this work.

Abstract: We make use of generalized iterations of the Sacks forcing to define cardinal-preserving generic extensions of the constructible universe \mathbf{L} in which the axioms of \mathbf{ZF} hold and in addition either 1) the parameter-free countable axiom of choice \mathbf{AC}_ω^* fails, or 2) \mathbf{AC}_ω^* holds but the full countable axiom of choice \mathbf{AC}_ω fails in the domain of reals. In another generic extension of \mathbf{L} , we define a set $X \subseteq \mathcal{P}(\omega)$, which is a model of the parameter-free part \mathbf{PA}_2^* of the 2nd order Peano arithmetic \mathbf{PA}_2 , in which $\mathbf{CA}(\Sigma_2^1)$ (Comprehension for Σ_2^1 formulas with parameters) holds, yet an instance of Comprehension \mathbf{CA} for a more complex formula fails.

Treating the iterated Sacks forcing as a class forcing over \mathbf{L}_{ω_1} , we infer the following consistency results as corollaries. If the 2nd order Peano arithmetic \mathbf{PA}_2 is formally consistent then so are the theories: 1) $\mathbf{PA}_2 + \neg\mathbf{AC}_\omega^*$, 2) $\mathbf{PA}_2 + \mathbf{AC}_\omega^* + \neg\mathbf{AC}_\omega$, 3) $\mathbf{PA}_2^* + \mathbf{CA}(\Sigma_2^1) + \neg\mathbf{CA}$.

Keywords: forcing; projective well-orderings; projective classes; Jensen's forcing

MSC: 03E15; 03E35

Contents (added for the convenience of the refereeing process)

1. Introduction	3
2. Preliminaries	4
3. A cardinal-collapse model where the parameter-free AC_ω^* fails	5
4. A cardinal-collapse model where the parameter-free AC_ω^* holds but the full AC_ω fails	5
5. Iterated perfect sets	6
6. The forcing notion and the basic extension	8
7. Structure of the basic extension	9
8. A model in which the parameter-free AC_ω^* fails	10
9. The non-definability lemma	11
10. A model in which the parameter-free AC_ω^* holds but the full AC_ω fails	12
11. Models in which the parameter-free CA^* holds but the full CA fails	13
12. Working on the basis of the consistency of PA_2	14
13. Conclusions, remarks, and problems	16

1. Introduction

Let \mathbf{PA}_2 be the second-order Peano arithmetic **without** the schema of (countable) Choice in this paper. Discussing the structure and deductive properties of \mathbf{PA}_2 , Kreisel [?, § III, page 366] wrote that the selection of subsystems “is a central problem”. In particular, Kreisel notes, that

[...] if one is convinced of the significance of something like a given axiom schema, it is natural to study details, such as the effect of parameters.

Recall that *parameters* in this context are free variables in various axiom schemata in \mathbf{PA} , \mathbf{PA}_2 , \mathbf{ZFC} , and other similar theories. Thus the most obvious way to study “the effect of parameters” is to compare the strength of a given axiom schema S with its parameter-free subschema S^* . (The asterisk will mean the parameter-free subschema in this paper.)

Some work in this direction was done in the early years of modern set theory. In particular Levy [?] proved that the generic collapse of cardinals below \aleph_ω (called the Levy collapse, see Solovay [?]) results in a generic extension of \mathbf{L} in which \mathbf{AC}_ω^* fails, where \mathbf{AC}_ω^* is the parameter-free subschema of the (countable) choice schema \mathbf{AC}_ω in the language of \mathbf{PA}_2 . Later Guzicki [?] established that the Levy-style generic collapse below \aleph_{ω_1} results in a generic extension of \mathbf{L} in which \mathbf{AC}_ω (in the language of \mathbf{PA}_2) fails, but the parameter-free subschema \mathbf{AC}_ω^* holds, so that \mathbf{AC}_ω^* is strictly weaker than \mathbf{AC}_ω . This can be compared with an opposite result for the *dependent choice* schema \mathbf{DC} , in the language of \mathbf{PA}_2 , which happens to be equivalent to its parameter-free subschema \mathbf{DC}^* by a simple argument given for instance in [?].

Some results related to parameter-free versions of the Separation and Replacement axiom schemata in \mathbf{ZFC} also are known from [? ? ?].

This paper is devoted to further clarification of the role of parameters in the Choice schema \mathbf{AC}_ω and *comprehension schema* \mathbf{CA} in \mathbf{PA}_2 . Special attention will be paid to the evaluation of those proof theoretic tools used in the arguments. That is, we show that the formal consistency of \mathbf{PA}_2 suffices. This is a crucial advantage comparably to some earlier results, like e.g. the abovementioned results by Levy [?] and Guzicki [?] which definitely cannot be obtained on the base of the consistency of \mathbf{PA}_2 .

The following theorems 1, 2, 3 are the main results of this paper.

Theorem 1. *In \mathbf{ZF} , let \mathbf{L} be the constructible universe. Then:*

- (i) *There is a cardinal-preserving generic extension of \mathbf{L} in which $\mathbf{AC}_\omega(\text{OD})$ (that is, \mathbf{AC}_ω for ordinal-definable relations) holds, but the full \mathbf{AC}_ω fails in the domain of reals.*
- (ii) *If \mathbf{PA}_2 is consistent then $\mathbf{PA}_2 + \mathbf{AC}_\omega^*$ does not prove \mathbf{AC}_ω .*

Theorem 1 is entirely new. Part (i) greatly surpasses the abovementioned result of Guzicki [?] by the requirement of cardinal-preservation. This is a *condicio sine qua non* for Claim (ii) to be derived as a consequence, because involvement of uncountable cardinals in the arguments, as in [?], is definitely beyond the formal consistency of \mathbf{PA}_2 .

In the next theorem, \mathbf{PA}_2^* is the subtheory of \mathbf{PA}_2 in which the full schema \mathbf{CA} is replaced by its parameter-free version \mathbf{CA}^* , and the Induction principle is formulated as a schema rather than one sentence.

Theorem 2. *In \mathbf{ZF} , let \mathbf{L} be the constructible universe. Then:*

- (i) *There is a cardinal-preserving generic extension of \mathbf{L} , and a set $M \subseteq \mathcal{P}(\omega)$ in this extension, such that $\mathcal{P}(\omega) \cap \mathbf{L} \subseteq M$ and M models $\mathbf{PA}_2^* + \mathbf{CA}(\Sigma_1^1) + \neg \mathbf{CA}$.*
- (ii) *If \mathbf{PA}_2 is consistent then $\mathbf{PA}_2^* + \mathbf{CA}(\Sigma_1^1)$ does not prove \mathbf{CA} .*

This is a new result as well, appeared in our recent ArXiv preprint [?].

The next theorem, albeit not entirely new, is added in for good measure as its proof involves basically the same type of generic extensions.

Theorem 3. In \mathbf{ZF} , let \mathbf{L} be the constructible universe. Then:

- (i) There is a cardinal-preserving generic extension of \mathbf{L} in which \mathbf{AC}_ω^* fails.
- (ii) If \mathbf{PA}_2 is consistent then \mathbf{PA}_2 does not prove \mathbf{AC}_ω^* .

Part (i) of this theorem was essentially established by Enayat [?], where it is shown that using the finite-support infinite product of Jensen's minimal- Δ_3^1 -real forcing [?] results in a permutation model of \mathbf{ZF} with an infinite Dedekind-finite Π_2^1 set of reals, which easily yields the refutation of \mathbf{AC}_ω^* .

The paper is organized as follows. After a short review of \mathbf{PA}_2 preliminaries in Section 2, we take some space to briefly describe the aforementioned cardinal-collapse models by Levy [?] and Guzik [?] in Sections 3 and 4.

The first claims of all three theorems will be established by means of a complex iteration of the Sacks forcing which resembles the generalized iteration by Groszek and Jech [?], but is carried out in a pure geometric way that avoids any machinery of forcing iterations. We call this technique *arboreal Sacks iterations*. The associated coding by degrees of constructibility is also involved, approximately along the lines discussed in [?, page 143].

Our basic forcing notion $\mathbf{Perf} = \mathbb{P}$ is introduced in Section 5; it consists of *iterated perfect sets*. The structure of \mathbb{P} -generic extensions $\mathbf{L}[G]$ of \mathbf{L} is studied in Sections 6 and 7. In particular, Theorem 5 contains several important results on the degrees of constructibility of reals and the relation of true $\leq_{\mathbf{L}}$ -successor in the generic extensions considered.

The proof of Theorem 3(i) is carried out in Section 8 modulo an important lemma established in Section 9. Basically, a generic extension that proves Theorem 3(i) will be obtained as a certain subextension of a \mathbb{P} -generic extension $\mathbf{L}[G]$.

Claims (i) of Theorems 1 and 2 will be established in Sections resp. 10, 11, also via different subextensions of a \mathbb{P} -generic extension.

Finally Section 12 contains the proof of claims (ii) of all three theorems. To do this, we will redo proofs of claims (i) in some uniform manner.

The paper ends with a usual conclusion-style material.

It remains to note that topics in subsystems of second order arithmetic remain of big interest in modern studies, see e.g. [?], and our paper contributes to this research line.

2. Preliminaries

Following [?] we define the second order Peano arithmetic \mathbf{PA}_2 as a theory in the language $\mathcal{L}(\mathbf{PA}_2)$ with two sorts of variables – for natural numbers and for sets of them. We use j, k, m, n for variables over ω and x, y, z for variables over $\mathcal{P}(\omega)$, reserving capital letters for subsets of $\mathcal{P}(\omega)$ and other sets. The axioms are as follows in (1), (2), (3), (4):

- (1) **Peano's axioms** for numbers.
- (2) The **Induction** schema: $\Phi(0) \wedge \forall k (\Phi(k) \implies \Phi(k+1)) \implies \forall k \Phi(k)$, for every formula $\Phi(k)$ in $\mathcal{L}(\mathbf{PA}_2)$, and in $\Phi(k)$ we allow parameters, i.e., free variables other than k . (We do not formulate Induction as one sentence here because the Comprehension schema **CA** will not be assumed in full generality in Section 11.)
- (3) **Extensionality** for sets of natural numbers.
- (4) The **Comprehension** schema **CA**: $\exists x \forall k (k \in x \iff \Phi(k))$, for every formula Φ in which x does not occur, and in Φ we allow parameters.

\mathbf{PA}_2 is also known as A_2^- (see e.g. an early survey [?]), as Z_2 (see e.g. Simpson [?] and Friedman [?]), as Z_2^- (in [?] or elsewhere). Note that the schema of Choice (see below) is not included in \mathbf{PA}_2 .

The following schemata are not assumed to be parts of \mathbf{PA}_2 , yet they are often considered in the context of and in the connection with \mathbf{PA}_2 .

The Schema of Choice \mathbf{AC}_ω : $\forall k \exists x \Phi(k, x) \implies \exists x \forall k \Phi(k, (x)_k)$, for every formula Φ , where we allow parameters in Φ , and $(x)_k = \{j : 2^k(2j+1) - 1 \in x\}$, as usual.

We use \mathbf{AC}_ω instead of \mathbf{AC} , more common in \mathbf{PA}_2 studies, because \mathbf{AC} is the general axiom of choice in the \mathbf{ZFC} context.

Dependent Choices DC: $\forall x \exists y \Phi(x, y) \implies \exists x \forall k \Phi((x)_k, (x)_{k+1})$, for every formula Φ , and in Φ we allow parameters.

We let \mathbf{CA}^* be the parameter-free sub-schema of \mathbf{CA} (that is, $\Phi(k)$ contains no free variables other than k). We define the parameter-free sub-schema $\mathbf{AC}_\omega^* \subseteq \mathbf{AC}_\omega$ the same way. The parameter-free sub-schema $\mathbf{DC}^* \subseteq \mathbf{DC}$ can be defined as well, but this does not make much sense because \mathbf{DC}^* is known to be equivalent to \mathbf{DC} by a simple argument, see e.g. [?].

In set-theoretic setting, \mathbf{AC}_ω and \mathbf{DC} can be considered in the assumption that Φ is a set-theoretic binary relation on $\omega \times \mathcal{P}(\omega)$, whose type can be restricted in this or another way depending on the context. In particular, $\mathbf{AC}_\omega(\text{OD})$ assumes that Φ is an OD (ordinal-definable) relation. (See [?] on ordinal definability.) In addition, say $\mathbf{AC}_\omega^*(\Pi_3^1)$ or $\mathbf{AC}_\omega(\Pi_3^1)$ means the restriction to the type of lightface Π_3^1 (parameter-free) or resp. boldface Π_3^1 (with parameters in $\mathcal{P}(\omega)$ allowed) formulas.

3. A cardinal-collapse model where the parameter-free \mathbf{AC}_ω^* fails

Here we recall an old model by Levy [?] in which the parameter-free \mathbf{AC}_ω^* fails for a certain (lightface) Π_2^1 relation. This is basically any model of $\mathbf{ZF} + (\aleph_1 = \aleph_\omega^{\mathbf{L}})$. To get this model, Levy makes use of the collapse below \aleph_ω , i.e., a Cohen-style generic sequence $f = \langle f_n \rangle_{n < \omega}$ of (generic) collapse maps $f_n : \omega \xrightarrow{\text{onto}} \aleph_n^{\mathbf{L}}$ is adjoined to the Gödel constructible universe \mathbf{L} . Consider the set $F = \{f_n : n < \omega\}$ and the class $N = \text{HOD}(F)$ of all sets hereditarily F -ordinal-definable in $\mathbf{L}[f]$. Then N is a model of $\mathbf{ZF} + (\aleph_1 = \aleph_\omega^{\mathbf{L}})$.

We may note that the set $\mathcal{P}(\omega) \cap N$ of all reals in N is equal to the set $\mathcal{P}(\omega) \cap \bigcup_{n < \omega} \mathbf{L}[f_0, f_1, \dots, f_n]$.

To prove that \mathbf{AC}_ω fails under $\aleph_1 = \aleph_\omega^{\mathbf{L}}$, Levy considers the relation

$$R(n, f) := n < \omega, f \in \omega^\omega, \text{ and } f \text{ codes a well-ordering of length } \geq \aleph_n^{\mathbf{L}}.$$

Then, first, \mathbf{AC}_ω fails for R under $\aleph_1 = \aleph_\omega^{\mathbf{L}}$ by obvious reasons, and second, R can be presented as a lightface Π_2^1 relation.

To prove the second claim, we may note, following Levy, that $R(n, f)$ is equivalent to the following relation:

$$R'(n, f) := n < \omega, f \in \omega^\omega, f \text{ codes a well-ordering, whose length we denote by } \alpha, \text{ and, for every countable transitive set } X \text{ which models } \mathbf{ZF} \text{ minus the Power Set axiom, if } \alpha \in X \text{ then it is true in } \langle X; \in \rangle \text{ that "there are at least } n+1 \text{ infinite cardinals } \leq \alpha".$$

To see that R' is a Π_2^1 relation, Levy uses well-founded relations on ω as a substitution for countable transitive sets. Since the well-foundedness is a Π_1^1 property, the definition of R' can be converted to a Π_2^1 form.

From a more modern perspective, we may note that R' is a Π_1^{HC} relation, where $\text{HC} = H_{\omega_1}$ is the transitive set of all *hereditarily countable* sets, and then make use of the conversion theorem (see e.g. Theorem 25.25 in [?]) saying that Π_1^{HC} relations on the reals are the same as Π_2^1 relations.

4. A cardinal-collapse model where the parameter-free \mathbf{AC}_ω^* holds but the full \mathbf{AC}_ω fails

The Guzikki model with such an effect appeared in [?]. It is similar to Levy's model of [?], yet it makes use of the Levy collapse below \aleph_{ω_1} . To get such a model, we adjoin, to the Gödel constructible universe \mathbf{L} , a Cohen-style (finite-support) generic sequence $f = \langle f_\xi \rangle_{\xi < \omega_1^{\mathbf{L}}}$ of (generic) collapsing maps $f_\xi : \omega \xrightarrow{\text{onto}} \aleph_\xi^{\mathbf{L}}$. Consider the set $F = \{f \restriction \beta : \beta < \omega_1^{\mathbf{L}}\}$ and the class N of all sets hereditarily F -real-ordinal definable in $\mathbf{L}[f]$. Then N is a model of $\mathbf{ZF} + (\aleph_1 = \aleph_{\omega_1}^{\mathbf{L}})$.

The set $\mathcal{P}(\omega) \cap N$ of all reals in N is equal to $\mathcal{P}(\omega) \cap \bigcup_{\beta < \omega_1^{\mathbf{L}}} \mathbf{L}[f \restriction \beta]$.

To check that \mathbf{AC}_ω fails in N for a Π_2^1 relation, let $p \in N, p \subseteq \omega$ code a strictly increasing map $g = g_p : \omega \rightarrow \omega_1^L$ whose range is cofinal in ω_1^L . Accordingly the sequence of cardinals $\aleph_{g(n)}^L \in N$ is cofinal in $\aleph_{\omega_1}^L$. This allows to accomodate the arguments in Section 3, with minor changes *mutatis mutandis*, and prove that \mathbf{AC}_ω fails in N for a Π_2^1 relation similar to R but defined with p as a parameter.

To see that the parameter-free \mathbf{AC}_ω^* , and even $\mathbf{AC}_\omega(\text{OD})$ for all ordinal-definable relations holds in N , let $\varphi(k, x, \gamma)$ be an \in -formula with an ordinal γ as the only parameter. Assume that $\forall k \exists x \subseteq \omega \varphi(k, x, \gamma)$ holds in N . Then for every k there exist ordinals $\beta < \omega_1^L$ such that a set $x \subseteq \omega$ satisfying $\varphi(k, x, \gamma)$ in N exists in $\mathbf{L}[f \restriction \beta]$. Let β_k be the least such an ordinal. The sequence $\langle \beta_n \rangle_{n < \omega}$ immediately belongs to $\mathbf{L}[f]$. Yet using the homogeneous character of the product collapse forcing that yields f , one can prove that in fact the sequence $\langle \beta_n \rangle_{n < \omega}$ in fact belongs to \mathbf{L} . Therefore $\beta = \sup_n \beta_n < \omega_1^L$, and accordingly for any k there is a set $x \subseteq \omega, x \in \mathbf{L}[f \restriction \beta]$ satisfying $\varphi(k, x, \gamma)$ in N . It remains to note that $\mathbf{L}[f \restriction \beta] \subseteq N$.

5. Iterated perfect sets

Here we begin the proof of Theorems 1, 2, 3. The proof involves the engine of generalized iterated Sacks forcing developed in [??] on the base of earlier papers [??] and others. We consider the constructible universe \mathbf{L} as the ground model.

Arguing in \mathbf{L} in this section, we define, in \mathbf{L} , the set

$$I = \omega_1^{<\omega} \setminus \{\Lambda\}; \quad I \in \mathbf{L};$$

of all non-empty tuples $i = \langle \xi_0, \dots, \xi_n \rangle, n < \omega$, of ordinals $\xi_k < \omega_1$, partially ordered by the extension \subset of tuples. I is a tree without the minimal node Λ (the empty tuple), which we exclude.

Our plan is to define a generic extension $\mathbf{L}[a]$ of \mathbf{L} by an array $a = \langle a_i \rangle_{i \in I}$ of reals $a_i \subseteq \omega$, in which the structure of “sacksness” is determined by this set I , so that in particular each a_i is Sacks-generic over the submodel $\mathbf{L}[\langle a_j \rangle_{j \subset i}]$. Then Theorems 1, 2, 3 will be obtained via submodels of the basic model $\mathbf{L}[a]$.

Let Ξ be the set of all countable and finite initial segments (in the sense of \subset) $\zeta \subseteq I$. If $\zeta \in \Xi$ then IS_ζ is the set of all initial segments of ζ .

Greek letters $\xi, \eta, \zeta, \vartheta$ will denote sets in Ξ .

Characters i, j are used to denote *elements* of I .

For any $i \in \zeta \in \Xi$, we consider initial segments $\zeta[\subset i] = \{j \in \zeta : j \subset i\}$ and $\zeta[\not\subset i] = \{j \in \zeta : j \not\subset i\}$, and $\zeta[\subseteq i], \zeta[\not\subseteq i]$ defined analogously.

We consider $\mathcal{P}(\omega)$ as identic to 2^ω , so that both $\mathcal{P}(\omega)$ and $\mathcal{P}(\omega)^\xi$ for $\xi \in \Xi$ are homeomorphic Polish compact spaces. Points of $\mathcal{P}(\omega)$ will be called *reals*.

Assume that $\eta \subseteq \zeta \in \Xi$. If $x \in \mathcal{P}(\omega)^\xi$ then let $x \restriction \eta \in \mathcal{P}(\omega)^\eta$ denote the usual restriction. If $X \subseteq \mathcal{P}(\omega)^\xi$ then let $X \restriction \eta = \{x \restriction \eta : x \in X\}$. To save space, let $X \restriction_{\subset i}$ mean $X \restriction \zeta[\subset i]$, $X \restriction_{\not\subset i}$ mean $X \restriction \zeta[\not\subset i]$, etc.

But if $Y \subseteq \mathcal{P}(\omega)^\eta$ then we put $Y \restriction^{-1} \zeta = \{x \in \mathcal{P}(\omega)^\xi : x \restriction \eta \in Y\}$.

To describe the idea behind the definition of iterated perfect sets, recall that the Sacks forcing consists of perfect subsets of $\mathcal{P}(\omega)$, that is, sets of the form $H''\mathcal{P}(\omega) = \{H(a) : a \in \mathcal{P}(\omega)\}$, where $H : \mathcal{P}(\omega) \xrightarrow{\text{onto}} X$ is a homeomorphism.

To get a product Sacks model, with two factors (the case of a two-element unordered set as the length of iteration), we have to consider sets $X \subseteq \mathcal{P}(\omega)^2$ of the form $X = H''\mathcal{P}(\omega)^2$ where H is any homeomorphism defined on $\mathcal{P}(\omega)^2$ so that it splits in obvious way into a pair of one-dimensional homeomorphisms.

To get an iterated Sacks model, with two stages of iteration (the case of a two-element ordered set as the length of iteration), we have to consider sets $X \subseteq \mathcal{P}(\omega)^2$ of the form $X = H''\mathcal{P}(\omega)^2$, where H is any homeomorphism defined on $\mathcal{P}(\omega)^2$ such that if $H(a_1, a_2) = \langle x_1, x_2 \rangle$ and $H(a'_1, a'_2) = \langle x'_1, x'_2 \rangle$ then $a_1 = a'_1 \iff x_1 = x'_1$.

The combined product/iteration case results in the following definition.

Definition 1 (iterated perfect sets, [? ?]). For any $\zeta \in \Xi$, \mathbf{Perf}_ζ is the collection of all sets $X \subseteq \mathcal{P}(\omega)^\zeta$ such that there is a homeomorphism $H : \mathcal{P}(\omega)^\zeta \xrightarrow{\text{onto}} X$ satisfying

$$x_0 \upharpoonright \zeta = x_1 \upharpoonright \zeta \iff H(x_0) \upharpoonright \zeta = H(x_1) \upharpoonright \zeta$$

for all $x_0, x_1 \in \text{dom } H$ and $\zeta \in \Xi$, $\xi \subseteq \zeta$. Homeomorphisms H satisfying this requirement will be called *projection-keeping*. In other words, sets in \mathbf{Perf}_ζ are images of $\mathcal{P}(\omega)^\zeta$ via projection-keeping homeomorphisms.

We put $\mathbf{Perf} = \bigcup_{\zeta \in \Xi} \mathbf{Perf}_\zeta$. \square

Remark 1. Note that \emptyset , the empty set, formally belongs to Ξ , and then $\mathcal{P}(\omega)^\emptyset = \{\emptyset\}$, and we easily see that $\mathbb{1} = \{\emptyset\}$ is the only set in \mathbf{Perf}_\emptyset . \square

For the convenience of the reader, we now present five lemmas on sets in \mathbf{Perf}_ζ established in [? ?].

Lemma 1 (Proposition 4 in [? ?]). Let $\zeta \in \Xi$. Every set $X \in \mathbf{Perf}_\zeta$ is closed and satisfies the following properties:

1. If $\mathbf{i} \in \zeta$ and $z \in X \upharpoonright_{\zeta \setminus \mathbf{i}}$ then $D_{Xz}(\mathbf{i}) = \{x(\mathbf{i}) : x \in X \wedge x \upharpoonright_{\zeta \setminus \mathbf{i}} = z\}$ is a perfect set in $\mathcal{P}(\omega)$.
2. If $\xi \in \text{IS}_\zeta$, and a set $X' \subseteq X$ is open in X (in the relative topology) then the projection $X' \upharpoonright \xi$ is open in $X \upharpoonright \xi$. In other words, the projection from X to $X \upharpoonright \xi$ is an open map.
3. If $\xi, \eta \in \text{IS}_\zeta$, $x \in X \upharpoonright \xi$, $y \in X \upharpoonright \eta$, and $x \upharpoonright (\xi \cap \eta) = y \upharpoonright (\xi \cap \eta)$, then $x \cup y \in X \upharpoonright (\xi \cup \eta)$.

Proof (sketch). Clearly $\mathcal{P}(\omega)^\zeta$ satisfies P-1, P-2, P-3, and one easily shows that projection-keeping homeomorphisms preserve the requirements. \square

Lemma 2 (Lemma 5 in [? ?]). Suppose that $\xi, \zeta, \theta \in \Xi$, $\xi \cup \zeta \subseteq \theta$, $W \in \mathbf{Perf}_\theta$, $C \subseteq W \upharpoonright \zeta$ is any set, and $U = W \cap (C \upharpoonright^{-1} \theta)$. Then $U \upharpoonright \xi = (W \upharpoonright \xi) \cap (C \upharpoonright (\xi \cap \zeta) \upharpoonright^{-1} \xi)$.

Lemma 3 (Lemma 6 in [? ?]). If $\zeta \in \Xi$, $X \in \mathbf{Perf}_\zeta$, $\xi \in \text{IS}_\zeta$, then $X \upharpoonright \xi \in \mathbf{Perf}_\xi$.

Lemma 4 (Lemma 8 in [? ?]). If $\zeta \in \Xi$, $X \in \mathbf{Perf}_\zeta$, a set $U \subseteq X$ is open in X , and $x_0 \in U$, then there is a set $X' \in \mathbf{Perf}_\zeta$, $X' \subseteq U$, clopen in X and containing x_0 .

Lemma 5 (Lemma 9 in [? ?]). Suppose that $\zeta \in \Xi$, $\eta \in \text{IS}_\zeta$, $X \in \mathbf{Perf}_\zeta$, $Y \in \mathbf{Perf}_\eta$, and $Y \subseteq X \upharpoonright \eta$. Then $Z = X \cap (Y \upharpoonright^{-1} \zeta)$ belongs to \mathbf{Perf}_ζ .

In particular $Y \upharpoonright^{-1} \zeta \in \mathbf{Perf}_\zeta$, since obviously $\mathcal{P}(\omega)^\zeta \in \mathbf{Perf}_\zeta$.

Corollary 1. Assume that $\xi, \eta \in \Xi$, $\theta = \xi \cup \eta$, $X \in \mathbf{Perf}_\xi$, $Y \in \mathbf{Perf}_\eta$, and $X \upharpoonright (\xi \cap \eta) = Y \upharpoonright (\xi \cap \eta)$. Then $Z = (X \upharpoonright^{-1} \theta) \cap (Y \upharpoonright^{-1} \theta) \in \mathbf{Perf}_\theta$.

Proof. The bigger set $X' = X \upharpoonright^{-1} \theta$ belongs to \mathbf{Perf}_θ by Lemma 5. In addition, $X' \upharpoonright \eta = X \upharpoonright (\xi \cap \eta) \upharpoonright^{-1} \eta$ by Lemma 2 (with $C = X$, $W = \mathcal{P}(\omega)^\theta$). It follows that $Y \subseteq X' \upharpoonright \eta$, because $Y \upharpoonright (\xi \cap \eta) = X \upharpoonright (\xi \cap \eta)$. We conclude that $X' \cap (Y \upharpoonright^{-1} \theta) \in \mathbf{Perf}_\theta$ by Lemma 5. Finally, we have $X' \cap (Y \upharpoonright^{-1} \theta) = Z$ by construction. \square

Corollary 2. Assume that $\xi_0, \xi_1, \xi_2, \dots \in \Xi$ are pairwise disjoint, $\theta = \bigcup_k \xi_k$, and $X_k \in \mathbf{Perf}_{\xi_k}$ for each k . Then the set $Z = \bigcap_k (X_k \upharpoonright^{-1} \theta)$ belongs to \mathbf{Perf}_θ , $Z \upharpoonright \xi_k = X_k$ and $Z \leq X_k$ for all k .

Proof. For each k , there exists a projection-keeping homeomorphism $H_k : \mathcal{P}(\omega)^{\xi_k} \xrightarrow{\text{onto}} X_k$. Define $H : \mathcal{P}(\omega)^\theta \rightarrow \mathcal{P}(\omega)^\theta$ by $H(x) \upharpoonright \xi_k = H_k(x \upharpoonright \xi_k)$ for all k . Then H is projection-keeping and $H : \mathcal{P}(\omega)^\theta \xrightarrow{\text{onto}} Z$. \square

Still **arguing in \mathbf{L}** , we let Π be the group of all **permutations** π of the index set I , i.e. all bijections $\pi : I \xrightarrow{\text{onto}} I$ such that $i \subset j \iff \pi(i) \subset \pi(j)$. Any such a permutation $\pi \in \Pi$ induces a transformation acting on several types of objects as follows.

- If $\zeta \in \mathfrak{E}$, or generally $\zeta \subseteq I$, then $\pi\zeta = \pi''\zeta = \{\pi(i) : i \in \zeta\}$.
- If $\zeta \subseteq I$ and $x \in \mathcal{P}(\omega)^\zeta$ then $\pi x \in \mathcal{P}(\omega)^{\pi\zeta}$ is defined by $\pi x(\pi(i)) = x(i)$ for all $i \in \zeta$. That is, formally $\pi x = x \circ \pi^{-1}$, the superposition.
- If $\zeta \subseteq I$ and $X \subseteq \mathcal{P}(\omega)^\zeta$ then $\pi X = \{\pi x : x \in X\}$.
- If $G \subseteq \mathbf{Perf}$ then $\pi G = \{\pi X : X \in G\}$.

The following lemma is obvious.

Lemma 6. *If $X \in \mathbf{Perf}_\zeta$ then $\pi X \in \mathbf{Perf}_{\pi\zeta}$.*

Moreover π is an order preserving automorphism of \mathbf{Perf} . \square

6. The forcing notion and the basic extension

This section introduces the forcing notion we consider and the according generic extension called the basic extension.

We continue to argue in \mathbf{L} . Recall that a partially ordered set $I \in \mathbf{L}$ is defined in Section 5, and \mathfrak{E} is the set of all at most countable initial segments $\zeta \subseteq I$ in \mathbf{L} . For any $\zeta \in \mathfrak{E}$, let $\mathbb{P}_\zeta = (\mathbf{Perf}_\zeta)^\mathbf{L}$.

The set $\mathbb{P} = \mathbb{P}_I = \bigcup_{\zeta \in \mathfrak{E}} \mathbb{P}_\zeta \in \mathbf{L}$ will be the *forcing notion*.

To define the order, we put $\|X\| = \zeta$ whenever $X \in \mathbb{P}_\zeta$. Now we set $X \leq Y$ (i.e. X is *stronger* than Y) iff $\zeta = \|Y\| \subseteq \|X\|$ and $X \restriction \zeta \subseteq Y$.

Remark 2. We may note that the set $\mathbb{1} = \{\emptyset\}$ as in Remark 1 belongs to \mathbb{P} and is the \leq -largest (i.e., the weakest) element of \mathbb{P} . \square

Now let $G \subseteq \mathbb{P}$ be a \mathbb{P} -generic set (filter) over \mathbf{L} .

Remark 3. If $X \in \mathbb{P}_\zeta$ in \mathbf{L} then X is not even a closed set in $\mathcal{P}(\omega)^\zeta$ in $\mathbf{L}[G]$. However we can transform it to a perfect set in $\mathbf{L}[G]$ by the closure operation. Indeed the topological closure $X^\#$ of such a set X in $\mathcal{P}(\omega)^\zeta$ taken in $\mathbf{L}[G]$ belongs to \mathbf{Perf}_ζ from the point of view of $\mathbf{L}[G]$. \square

It easily follows from Lemma 4 that there exists a unique array $\mathbf{a}[G] = \langle \mathbf{a}_i[G] \rangle_{i \in I}$, all $\mathbf{a}_i[G]$ being elements of $\mathcal{P}(\omega)$, such that $\mathbf{a}[G] \restriction \zeta \in X^\#$ whenever $X \in G$ and $\|X\| = \zeta \in \mathfrak{E}$. Then $\mathbf{L}[G] = \mathbf{L}[\langle \mathbf{a}_i[G] \rangle_{i \in I}] = \mathbf{L}[\mathbf{a}[G]]$ is a \mathbb{P} -generic extension of \mathbf{L} , which we call *the basic extension*.

For the sake of convenience, let $\mathbf{a}_\Lambda[G] = \emptyset$.

Theorem 4 (Thm 24 in both [?] and [?]). *Every cardinal in \mathbf{L} remains a cardinal in $\mathbf{L}[G]$. Every $\mathbf{a}_i[G]$ is Sacks generic over the model $\mathbf{L}[\mathbf{a}[G] \restriction_{\subset i}]$.*

Proof (idea). The forcing \mathbf{Perf} has the following property in \mathbf{L} , common with the ordinary one-step Sacks forcing:

- (*) if sets $D_n \subseteq \mathbf{Perf}$ are open dense in \mathbf{Perf} , and $X \in \mathbf{Perf}$, then there is a stronger condition $Y \in \mathbf{Perf}$, $Y \leq X$, and finite sets $D'_n \subseteq D_n$ *pre-dense* in \mathbf{Perf} below Y , in the sense that any stronger $Z \in \mathbf{Perf}$, $Z \leq Y$, is compatible with some $Z' \in D'_n$.

This property, established in [?] , [?] by means of a splitting/fusion technique, easily implies the preservation of all \mathbf{L} -cardinals in \mathbb{P} -generic extensions of \mathbf{L} . \square

Here follow several lemmas on reals in \mathbb{P} -generic models $\mathbf{L}[G]$, established in [?]. In the lemmas, we let $G \subseteq \mathbb{P}$ be a set \mathbb{P} -generic over \mathbf{L} .

Lemma 7 (Lemma 22 in [?]). Suppose that sets $\eta, \xi \in \Xi$ satisfy $\forall j \in \eta \exists i \in \xi (j \subseteq i)$. Then $\mathbf{a}[G] \upharpoonright \eta \in \mathbf{L}[\mathbf{a}[G] \upharpoonright \xi]$. \square

Lemma 8 (Lemma 26 in [?]). Suppose that $K \in \mathbf{L}$ is an initial segment in I , and $i \in I \setminus K$. Then $\mathbf{a}_i[G] \notin \mathbf{L}[\mathbf{a}[G] \upharpoonright K]$. \square

Lemma 9 (Corollary 27 in [?]). If $i \neq j$ then $\mathbf{a}_i[G] \neq \mathbf{a}_j[G]$ and even $\mathbf{L}[\mathbf{a}_i[G]] \neq \mathbf{L}[\mathbf{a}_j[G]]$. \square

Lemma 10 (Lemma 29 in [?]). If $K \in \mathbf{L}$ is an initial segment of I , and $r \in \mathcal{P}(\omega) \cap \mathbf{L}[G]$, then either $r \in \mathbf{L}[\mathbf{a}[G] \upharpoonright K]$ or $\mathbf{a}_i[G] \in \mathbf{L}[r]$ for some $i \in I \setminus K$. \square

7. Structure of the basic extension

We apply the lemmas above in the proof of the next theorem. Let $\leq_{\mathbf{L}}$ denote the Gödel wellordering on $\mathcal{P}(\omega)$, so that $x \leq_{\mathbf{L}} y$ iff $x \in \mathbf{L}[y]$. Let $x <_{\mathbf{L}} y$ mean that $x \leq_{\mathbf{L}} y$ but $y \not\leq_{\mathbf{L}} x$, and $x \equiv_{\mathbf{L}} y$ mean that $x \leq_{\mathbf{L}} y$ and $y \leq_{\mathbf{L}} x$.

Say that y is a **true** $\leq_{\mathbf{L}}$ -successor of x (where $x, y \in \mathcal{P}(\omega)$) iff $x <_{\mathbf{L}} y$ and any real $z \in \mathcal{P}(\omega)$ satisfies $z <_{\mathbf{L}} y \implies z \leq_{\mathbf{L}} x$.

Theorem 5. Let $G \subseteq \mathbb{P}$ be a set \mathbb{P} -generic over \mathbf{L} , and $i \in I$. Then we have the following:

- (i) if $j \in I$ and $j \subseteq i$ then $\mathbf{a}_j[G] \leq_{\mathbf{L}} \mathbf{a}_i[G]$;
- (ii) if $j \in I$ and $j \not\subseteq i$ then $\mathbf{a}_j[G] \not\leq_{\mathbf{L}} \mathbf{a}_i[G]$;
- (iii) if $r \in \mathbf{L}[G] \cap \mathcal{P}(\omega)$ and $r \leq_{\mathbf{L}} \mathbf{a}_i[G]$ then $r \in \mathbf{L}$ or $r \equiv_{\mathbf{L}} \mathbf{a}_j[G]$ for some $j \in I, j \subseteq i$;
- (iv) if $i \in I, \gamma < \omega_1^{\mathbf{L}}$, then $\mathbf{a}_{i \cap \gamma}[G]$ is a true $\leq_{\mathbf{L}}$ -successor of $\mathbf{a}_i[G]$;
- (v) if $i \in I$, and $y \in \mathcal{P}(\omega) \cap \mathbf{L}[G]$ is a true $\leq_{\mathbf{L}}$ -successor of $\mathbf{a}_i[G]$, then there is $\gamma < \omega_1^{\mathbf{L}}$ such that $y \equiv_{\mathbf{L}} \mathbf{a}_{i \cap \gamma}[G]$;
- (vi) if $\gamma < \omega_1^{\mathbf{L}}$, then $\mathbf{a}_{\langle \gamma \rangle}[G]$ is a true $\leq_{\mathbf{L}}$ -successor of $\mathbf{a}_{\Lambda}[G]$;
- (vii) if $y \in \mathcal{P}(\omega) \cap \mathbf{L}[G]$ is a true $\leq_{\mathbf{L}}$ -successor of $\mathbf{a}_{\Lambda}[G]$, then there is $\gamma < \omega_1^{\mathbf{L}}$ such that $x \equiv_{\mathbf{L}} \mathbf{a}_{\langle \gamma \rangle}[G]$.

Proof. (i) Apply Lemma 7 with $\eta = \{j\}$ and $\xi = \{i\}$.

(ii) Apply Lemma 8 with $K = [\subseteq i]$.

(iii) If there are elements $j \in I, j \subseteq i$, such that $\mathbf{a}_j[G] \in \mathbf{L}[r]$, then let j be the largest such one. Let $\xi = [\subseteq j]$ (a finite initial segment of I). By Lemma 10, either $r \in \mathbf{L}[\mathbf{a}[G] \upharpoonright \xi]$, or there is $i' \notin \xi$ such that $\mathbf{a}_{i'}[G] \in \mathbf{L}[r]$. In the “either” case, we have $r \in \mathbf{L}[\mathbf{a}_j[G]]$ by (i), so that $\mathbf{L}[r] = \mathbf{L}[\mathbf{a}_j[G]]$ by the choice of j . In the “or” case we have $\mathbf{a}_{i'}[G] \in \mathbf{L}[\mathbf{a}_j[G]]$, hence $i' \subseteq i$ by (ii). But this contradicts the choice of j and i' .

Finally if there is no $j \in I, j \subseteq i$, such that $\mathbf{a}_j[G] \in \mathbf{L}[r]$, then the same argument with $\xi = \emptyset$ gives $r \in \mathbf{L}$.

(iv) The relation $\mathbf{a}_i[G] <_{\mathbf{L}} \mathbf{a}_{i \cap \gamma}[G]$ is implied by Lemmas 7 and 8. If now $z <_{\mathbf{L}} \mathbf{a}_{i \cap \gamma}[G]$ then $z \in \mathbf{L}$ or $z \equiv_{\mathbf{L}} \mathbf{a}_j[G]$ for some $j \subseteq i \cap \gamma$ by (iii), and in the latter case in fact $j \subset i \cap \gamma$, hence $j \subseteq i$, and then $z \leq_{\mathbf{L}} \mathbf{a}_i[G]$.

(v) As $y \not\leq_{\mathbf{L}} \mathbf{a}_i[G]$, by Lemma 10 there is $j \in I$ such that $j \not\subseteq i$ and $\mathbf{a}_j[G] \leq_{\mathbf{L}} y$. If $\mathbf{a}_j[G] <_{\mathbf{L}} y$ strictly then $\mathbf{a}_j[G] \leq_{\mathbf{L}} \mathbf{a}_i[G]$ by the true $\leq_{\mathbf{L}}$ -successor property, hence $j \subseteq i$ by (ii), contrary to the choice of j . Therefore in fact $\mathbf{a}_j[G] \equiv_{\mathbf{L}} y$. Then we have $i \subset j$ still by the true $\leq_{\mathbf{L}}$ -successor property and (i), (ii). This implies $j = i \cap \gamma$ for some $\gamma < \omega_1^{\mathbf{L}}$, because if say $j = i \cap \gamma \cap \delta$ then $z = \mathbf{a}_{i \cap \gamma}[G]$ is strictly between $\mathbf{a}_i[G]$ and $\mathbf{a}_j[G]$, contrary to the true $\leq_{\mathbf{L}}$ -successor property.

(vi) Similar to (iv). Recall that $\mathbf{a}_{\Lambda}[G] = \emptyset \in \mathbf{L}$. This implies $\mathbf{a}_{\Lambda}[G] \leq_{\mathbf{L}} \mathbf{a}_{\langle \gamma \rangle}[G]$. On the other hand, $\mathbf{a}_{\langle \gamma \rangle}[G] \not\leq_{\mathbf{L}} \mathbf{a}_{\Lambda}[G]$ holds by Lemma 8 with $K = \emptyset$. If now $z <_{\mathbf{L}} \mathbf{a}_{\langle \gamma \rangle}[G]$ then $z \in \mathbf{L}$ or $z \equiv_{\mathbf{L}} \mathbf{a}_j[G]$ for some $j \subseteq \langle \gamma \rangle$ by (iii), and in the latter case in fact $j = \langle \gamma \rangle$, hence then $z \equiv_{\mathbf{L}} \mathbf{a}_{\langle \gamma \rangle}[G]$, contrary to the choice of z .

(viii) As $y \not\leq_L \mathbf{a}_\Lambda[G] \in \mathbf{L}$, by Lemma 10 (with $K = \emptyset$) there is $j \in I$ such that $\mathbf{a}_j[G] \leq_L y$. If $\mathbf{a}_j[G] <_L y$ strictly then $\mathbf{a}_j[G] \leq_L \mathbf{a}_\Lambda[G]$ by the true \leq_L -successor property, hence $\mathbf{a}_j[G] \in \mathbf{L}$, contrary to Lemma 8 with $K = \emptyset$. Therefore in fact $\mathbf{a}_j[G] \equiv_L y$. This implies $j = \langle \gamma \rangle$ for some $\gamma < \omega_1^L$, because if say $j = \langle \gamma, \delta \rangle$ then $y = \mathbf{a}_{\langle \gamma \rangle}[G]$ is strictly between $\mathbf{a}_\Lambda[G]$ and $y \equiv_L \mathbf{a}_j[G]$, contrary to the true \leq_L -successor property. \square

Now consider the following formula:

$\mathfrak{A}(n, \vec{x}) := \vec{x} = \langle x_0, x_1, \dots, x_n \rangle$ is a tuple of reals $x_k \subseteq \omega$ such that $x_0 = \emptyset$ and each x_k ($0 < k \leq n$) is a true \leq_L -successor of x_{k-1} .

Thus $\mathfrak{A}(n, \vec{x})$ separates tuples of true successor iterations, of length n .

Remark 4. $\mathfrak{A}(n, \vec{x})$ is a Π_3^1 relation, absolute for any transitive model of **ZF** containing the true ω_1 , and componentwise \equiv_L -invariant in the argument $\vec{x} = \langle x_0, x_1, \dots, x_n \rangle$. Indeed to see that \mathfrak{A} is Π_3^1 note that ‘being a true \leq_L -successor’ is Π_3^1 by direct estimation. To see the absoluteness note that both ‘being a true \leq_L -successor’ and \mathfrak{A} are relativized to the lower \leq_L -cone of the arguments. The invariance is obvious. \square

Corollary 3 (of Theorem 5). Let $G \subseteq \mathbb{P}$ be a set \mathbb{P} -generic over \mathbf{L} .

(i) If $i = \langle \gamma_1, \gamma_2, \dots, \gamma_n \rangle \in I$, $\text{dom } i = n \geq 1$, and

$$\mathbf{a}_{\subseteq i}[G] = \langle \mathbf{a}_\Lambda[G], \mathbf{a}_{\langle \gamma_1 \rangle}[G], \mathbf{a}_{\langle \gamma_1, \gamma_2 \rangle}[G], \dots, \mathbf{a}_{\langle \gamma_1, \gamma_2, \dots, \gamma_n \rangle}[G] \rangle, \quad (1)$$

then $\mathfrak{A}(n, \mathbf{a}_{\subseteq i}[G])$ holds in $\mathbf{L}[G]$.

(ii) Conversely if $\vec{x} = \langle x_0, x_1, \dots, x_n \rangle \in \mathbf{L}[G]$ and $\mathfrak{A}(n, \vec{x})$ holds in $\mathbf{L}[G]$ then there is $i = \langle \gamma_1, \gamma_2, \dots, \gamma_n \rangle \in I$ such that $\vec{x} \equiv_L \mathbf{a}_{\subseteq i}[G]$ componentwise, that is, $x_0 \equiv_L \mathbf{a}_\Lambda[G]$, $x_1 \equiv_L \mathbf{a}_{\langle \gamma_1 \rangle}[G]$, $x_2 \equiv_L \mathbf{a}_{\langle \gamma_1, \gamma_2 \rangle}[G]$, \dots , $x_n \equiv_L \mathbf{a}_{\langle \gamma_1, \gamma_2, \dots, \gamma_n \rangle}[G]$. \square

8. A model in which the parameter-free \mathbf{AC}_ω^* fails

Here we prove Theorem 3(i). Let us fix a set $G \subseteq \mathbb{P}$, \mathbb{P} -generic over \mathbf{L} and consider the according \mathbb{P} -generic array $\mathbf{a}[G] = \langle \mathbf{a}_i[G] \rangle_{i \in I}$ and the \mathbb{P} -generic extension $\mathbf{L}[G] = \mathbf{L}[\mathbf{a}[G]]$. The goal is to define a sub-extension of $\mathbf{L}[G]$ in which the parameter-free \mathbf{AC}_ω^* fails.

- Let $\Omega \in \mathbf{L}$ be the set of all finite or \mathbf{L} -countable initial segments $\zeta \subseteq I$ such that there is a number $n < \omega$ satisfying $\text{dom } i < n$ for all $i \in \zeta$.
- Let $W[G] \in \mathbf{L}[G]$ be the set of all restrictions of the form $\mathbf{a}[G] \upharpoonright \zeta$, $\zeta \in \Omega$, of the generic array $\mathbf{a}[G]$.
- Let $\text{OD}(W[G])^{\mathbf{L}[G]}$ be the class of all sets $W[G]$ -ordinal-definable in $\mathbf{L}[G]$. Thus $x \in \text{OD}(W[G])^{\mathbf{L}[G]}$ iff x is definable in $\mathbf{L}[G]$ by a set-theoretic formula with parameters in $W[G] \cup \mathbf{Ord}$.

Here \mathbf{Ord} is the class of all ordinals, as usual. See [?] , [?] on ordinal definability.

- Let $\mathfrak{M}_G = \text{HOD}(W[G])^{\mathbf{L}[G]}$ be the class of all sets $x \in \mathbf{L}[G]$, hereditarily $W[G]$ -ordinal-definable in $\mathbf{L}[G]$, i.e., it is required that x itself, all elements of x , all elements of elements of x , etc., belong to the above defined class $\text{OD}(W[G])^{\mathbf{L}[G]}$ in $\mathbf{L}[G]$.

Theorem 6. If a set $G \subseteq \mathbb{P}$ is \mathbb{P} -generic over \mathbf{L} then \mathfrak{M}_G is a model of **ZF** in which the parameter-free $\mathbf{AC}_\omega^*(\Pi_3^1)$ fails.

It follows that $\mathfrak{M}_G \cap \mathcal{P}(\omega)$ is a model of $\mathbf{PA}_2 + \neg \mathbf{AC}_\omega^*(\Pi_3^1)$.

Proof. That classes of the form $\text{HOD}(X)$ model **ZF** see [?] , Chapter 13.

Note that if $i \in I$ then $\mathbf{a}_i[G] \in \mathfrak{M}_G = \text{HOD}(W[G])^{\mathbf{L}[G]}$ via the initial segment $\zeta = [\subseteq a] = \{j \in I : j \subseteq i\} \in \Omega$, and hence $\mathbf{a}_{\subseteq i}[G] \in \mathfrak{M}_G$ as well. It follows by Corollary 3(i) that

$\exists x \mathfrak{A}(m, x)$ is true in \mathfrak{M}_G , where $m = \text{dom } i$. Our goal will be to show that the parameter-free formula $\exists x \forall m \mathfrak{A}(m, (x)_m)$, the right-hand side of \mathbf{AC}_ω , fails in \mathfrak{M}_G , meaning that \mathbf{AC}_ω^* fails in \mathfrak{M}_G for the formula \mathfrak{A} .

Suppose to the contrary that there is $x \in \mathfrak{M}_G$ satisfying $\forall m \mathfrak{A}(m, (x)_m)$. This obviously results in a sequence $\langle \vec{y}_m \rangle_{m < \omega} \in \mathfrak{M}_G$ of tuples $\vec{y}_m = \langle y_0^m, y_1^m, \dots, y_m^m \rangle \in \mathfrak{M}_G$ of reals $y_k^m \subseteq \omega$ satisfying $\mathfrak{A}(k, \vec{y}_k)$, that is, $y_0^m = \emptyset$ and each y_k ($0 < k \leq m$) is a true \leq_L -successor of y_{k-1} .

By definition there is an \in -formula $\varphi(m, k, y, \mathbf{a}[G] \upharpoonright \xi)$ with free variables m, k, y , a parameter of the form $\mathbf{a}[G] \upharpoonright \xi$, where $\xi \in \Omega$, and some ordinals as parameters — such that if $k \leq m < \omega$ and $y \in \mathfrak{M}_G \cap \mathcal{P}(\omega)$ then $\varphi(m, k, y, \mathbf{a}[G] \upharpoonright \xi)$ is true in $\mathbf{L}[G]$ iff $y = y_k^m$. (The case of several parameters of the form $\mathbf{a}[G] \upharpoonright \xi$, $\xi \in \Omega$, can be easily reduced to the case of one parameter.)

As $\xi \in \Omega$, there is a number $1 \leq m < \omega$ such that $\text{dom } i < m$ for all $i \in \xi$. Fix this m and consider the tuple $\vec{y}_m = \langle y_0^m, y_1^m, \dots, y_m^m \rangle \in \mathfrak{M}_G = \text{HOD}(W[G])^{\mathbf{L}[G]}$. By Corollary 3(ii), there is a tuple $j = \langle \gamma_1, \gamma_2, \dots, \gamma_m \rangle \in I$, such that $\vec{y}_m \equiv_{\mathbf{L}} \mathbf{a}_{\subseteq j}[G]$ componentwise, that is, $y_k^m \equiv_{\mathbf{L}} \mathbf{a}_j[G] = \mathbf{a}_{\langle \gamma_1, \gamma_2, \dots, \gamma_k \rangle}[G]$ for all $k \leq m$.

Note that $j \notin \xi$ by the choice of m . There is a number $n \leq m$ such that still $i_0 = \langle \gamma_1, \gamma_2, \dots, \gamma_{n-1}, \gamma_n \rangle \notin \xi$ but the shorter tuple $i = \langle \gamma_1, \gamma_2, \dots, \gamma_{n-1} \rangle$ belongs to ξ , and hence $\mathbf{a}_{\subseteq i}[G] \in \text{HOD}(W[G])^{\mathbf{L}[G]}$. Then by Corollary 3 the \mathbf{L} -degree $[\mathbf{a}_{i_0}[G]]_{\mathbf{L}} = \{a \subseteq \omega : a \equiv_{\mathbf{L}} \mathbf{a}_{i_0}[G]\}$ is definable in $\mathbf{L}[G]$ by the next formula, in which $(\mathbf{a}[G] \upharpoonright \xi)(i) = \mathbf{a}_i[G]$.

$$\psi(a, \mathbf{a}[G] \upharpoonright \xi) := a \subseteq \omega \text{ is a true } \leq_L\text{-successor of } (\mathbf{a}[G] \upharpoonright \xi)(i).$$

To conclude, $i_0 \notin \xi \in \Omega$ and the \mathbf{L} -degree $[\mathbf{a}_{i_0}[G]]_{\mathbf{L}}$ is definable in $\mathbf{L}[G]$ by an \in -formula with $\mathbf{a}[G] \upharpoonright \xi$ and ordinals as parameters. But this contradicts Lemma 11 that follows in the next Section. The contradiction refutes the contrary assumption above.

We finally note that \mathfrak{A} is a Π_3^1 formula by Remark 4. \square

9. The non-definability lemma

Here we prove the following lemma.

Lemma 11. *If a set $G \subseteq \mathbb{P}$ is \mathbb{P} -generic over \mathbf{L} , $\xi \in \Xi$, and $i_0 \in I \setminus \xi$ then the \mathbf{L} -degree $[\mathbf{a}_{i_0}[G]]_{\mathbf{L}} = \{a \subseteq \omega : a \equiv_{\mathbf{L}} \mathbf{a}_{i_0}[G]\}$ cannot be defined in $\mathbf{L}[G]$ by an \in -formula with $\mathbf{a}[G] \upharpoonright \xi$ and ordinals as parameters.*

Proof. Suppose to the contrary that $\psi(x, \mathbf{a}[G] \upharpoonright \xi)$ is a formula as indicated, and it holds in $\mathbf{L}[G]$ that $[\mathbf{a}_{i_0}[G]]_{\mathbf{L}} = \{x \subseteq \omega : \psi(x, \mathbf{a}[G] \upharpoonright \xi)\}$. Then there is a “condition” $X_0 \in G$ such that

$$X_0 \Vdash [\mathbf{a}_{i_0}[G]]_{\mathbf{L}} = \{x \subseteq \omega : \psi(x, \mathbf{a}[\underline{G}] \upharpoonright \xi)\}, \quad (2)$$

where \Vdash is the \mathbb{P} -forcing relation over \mathbf{L} , and \underline{G} is the canonical \mathbb{P} -name for the generic filter G . Let $\zeta = \|X_0\|$, so that $X_0 \in \mathbb{P}_\zeta$.

We argue in \mathbf{L} . Thus $X \in \text{Perf}_\zeta$. See Section 5 on permutations of I .

As ξ, ζ are countable initial segments of I , it does not take much effort to define, in \mathbf{L} , a permutation $\pi \in \Pi$ satisfying the following:

- (A) $\pi \upharpoonright \xi$ is the identity;
- (B) $\pi(i_0) \neq i_0$, and if $i \in (\zeta \setminus \xi)$ then $\pi(i) \notin \zeta \setminus \xi$.

Coming back to (2) above, we put $Y_0 = \pi X_0$, $j_0 = \pi(i_0)$. Note that $Y_0 \in \mathbb{P}_{\zeta'}$ by Lemma 6, where $\zeta' = \pi\zeta = \pi''\zeta$. We claim that

$$Y_0 \Vdash [\mathbf{a}_{j_0}[G]]_{\mathbf{L}} = \{x \subseteq \omega : \psi(x, \mathbf{a}[\underline{G}] \upharpoonright \xi)\} \quad (3)$$

To prove the claim, let $H' \subseteq \mathbb{P}$ be \mathbb{P} -generic over \mathbf{L} , and $Y_0 \in H'$. We have to check that, in $\mathbf{L}[H']$, $[\mathbf{a}_{j_0}[H']]_{\mathbf{L}} = \{x \subseteq \omega : \psi(x, \mathbf{a}[H'] \upharpoonright \xi)\}$.

The set $H = \pi^{-1}H'$ is \mathbb{P} -generic over \mathbf{L} and obviously $X_0 \in H$. It follows from (2) that $[\mathbf{a}_{i_0}[H]]_{\mathbf{L}} = \{x \subseteq \omega : \psi(x, \mathbf{a}[H] \upharpoonright \xi)\}$ in $\mathbf{L}[H]$. Yet $\mathbf{L}[H] = \mathbf{L}[H']$ (since $\pi \in \mathbf{L}$), $\mathbf{a}[H'] \upharpoonright \xi = \mathbf{a}[H] \upharpoonright \xi$ by (A), and finally $\mathbf{a}_{j_0}[H'] = \mathbf{a}_{i_0}[H]$ by construction. Thus, indeed $[\mathbf{a}_{j_0}[H']]_{\mathbf{L}} = \{x \subseteq \omega : \psi(x, \mathbf{a}[H'] \upharpoonright \xi)\}$ in $\mathbf{L}[H']$, as required. This completes the proof of (3).

The next step is to establish

(C) X_0 and Y_0 are compatible in \mathbb{P} .

We check this claim *arguing in* \mathbf{L} , so that $X_0 \in \mathbf{Perf}_{\zeta}$ and $Y_0 \in \mathbf{Perf}_{\zeta'}$, where $\zeta' = \pi\zeta = \pi''\zeta$. It follows from (A), (B) that the set $\eta = \zeta \cap \zeta' \in \Xi$ satisfies $\eta = \zeta' \cap \zeta = \zeta' \cap \zeta$, and in addition $X_0 \upharpoonright \eta = Y_0 \upharpoonright \eta$. Let $\vartheta = \zeta \cup \zeta'$. Then $Z = (X_0 \upharpoonright^{-1} \vartheta) \cup (Y_0 \upharpoonright^{-1} \vartheta)$ belongs to $\mathbf{Perf}_{\vartheta}$ by Corollary 1. Thus $Z \in \mathbb{P}$, hence (C) holds. This implies (3) since $Z \leq X_0, Y_0$ is obvious.

But it follows from (2) and (3) that X_0 and Y_0 force contradictory statements (because $i_0 \neq j_0$, and hence $[\mathbf{a}_{i_0}[G]]_{\mathbf{L}} \neq [\mathbf{a}_{j_0}[G]]_{\mathbf{L}}$). The contradiction obtained completes the proof of the lemma. This accomplishes the proof of Theorem 6 as well. \square

10. A model in which the parameter-free \mathbf{AC}_{ω}^* holds but the full \mathbf{AC}_{ω} fails

Here we prove Theorem 1(i). The model will be a modification of the model studied in Section 8. We still fix a set $G \subseteq \mathbb{P}$, \mathbb{P} -generic over \mathbf{L} and consider the \mathbb{P} -generic array $\mathbf{a}[G] = \langle \mathbf{a}_i[G] \rangle_{i \in I}$ and the \mathbb{P} -generic extension $\mathbf{L}[G] = \mathbf{L}[\mathbf{a}[G]]$. We are going to define a sub-extension of $\mathbf{L}[G]$ in which the parameter-free \mathbf{AC}_{ω}^* holds but the full \mathbf{AC}_{ω} fails.

- Let $\Omega' \in \mathbf{L}$ be the set of all finite or \mathbf{L} -countable initial segments $\xi \subseteq I$ such that for any $\gamma < \omega_1$ there is a number $n = n_{\gamma} < \omega$ satisfying $\text{dom } i < n$ for all $i \in \xi$ satisfying $i(0) = \gamma$.
- Let $W'[G] \in \mathbf{L}[G]$ be the set of all restrictions of the form $\mathbf{a}[G] \upharpoonright \xi$, $\xi \in \Omega'$, of the generic array $\mathbf{a}[G]$.
- Let $\text{OD}(W'[G])^{\mathbf{L}[G]}$ be the class of all sets $W'[G]$ -ordinal-definable in $\mathbf{L}[G]$. Thus $x \in \text{OD}(W'[G])^{\mathbf{L}[G]}$ iff x is definable in $\mathbf{L}[G]$ by a set-theoretic formula with sets in $W'[G] \cup \mathbf{Ord}$ as parameters.
- Let $\mathfrak{M}'_G = \text{HOD}(W'[G])^{\mathbf{L}[G]}$ be the class of all sets $x \in \mathbf{L}[G]$, hereditarily $W'[G]$ -ordinal-definable in $\mathbf{L}[G]$.

Theorem 7. *If a set $G \subseteq \mathbb{P}$ is \mathbb{P} -generic over \mathbf{L} then \mathfrak{M}'_G is a model of \mathbf{ZF} in which the parameter-free \mathbf{AC}_{ω}^* holds, even $\mathbf{AC}_{\omega}(\text{OD})$ (with ordinals as parameters) holds, but the full $\mathbf{AC}_{\omega}(\Pi_3^1)$ fails. It follows that $\mathfrak{M}'_G \cap \mathcal{P}(\omega)$ is a model of $\mathbf{PA}_2 + \mathbf{AC}_{\omega}^* + \neg \mathbf{AC}_{\omega}(\Pi_3^1)$.*

Proof. Let $\mathfrak{A}'(n, \vec{x})$ be the formula ' $\mathfrak{A}(n, \vec{x}) \wedge x_0 = \mathbf{a}_{\langle 0 \rangle}[G]$ '. (See the definition of \mathfrak{A} in Section 7.) Note the parameter $\mathbf{a}_{\langle 0 \rangle}[G]$ in this formula. Similarly to the proof of Theorem 6, if $i \in I$ then $\mathbf{a}_i[G] \in \mathfrak{M}'_G$ and $\mathbf{a}_{\subseteq i}[G] \in \mathfrak{M}'_G$. It still follows by Corollary 3(i) that $\exists x \mathfrak{A}'(n, x)$ is true in \mathfrak{M}'_G , where $n = \text{dom } i$. Moreover, arguments pretty similar to the proof of Theorem 6, which we leave for the reader, show that the formula $\exists x \forall m \mathfrak{A}(k, (x)_m)$, the right-hand side of \mathbf{AC}_{ω} , fails in \mathfrak{M}'_G . Thus $\mathbf{AC}_{\omega}(\Pi_3^1)$ (with real parameters) fails in \mathfrak{M}'_G .

It remains to prove that $\mathbf{AC}_{\omega}(\text{OD})$ (with ordinals as parameters) holds in \mathfrak{M}'_G . Suppose towards the contrary that $\varphi(k, x)$ is an \in -formula with ordinals as parameters, such that \mathbf{AC}_{ω} fails for φ in \mathfrak{M}'_G . Thus there exists a condition $X^* \in G$ satisfying

(+) $X^* \Vdash$ "it holds in $\mathfrak{M}'_G = \text{HOD}(W'[G])^{\mathbf{L}[G]}$ that $\forall k \exists x \varphi(k, x)$ but $\neg \exists x \forall k \varphi(k, (x)_k)$ ".

Here \Vdash is the \mathbb{P} -forcing relation over \mathbf{L} , and \underline{G} is the canonical \mathbb{P} -name for the generic filter G , as above.

As $\forall k \exists x \varphi(k, x)$ holds in \mathfrak{M}'_G , there is a sequence $\langle x_k \rangle_{k < \omega} \in \mathbf{L}[G]$ of reals $x_k \in \mathfrak{M}'_G$, $x_k \subseteq \omega$, satisfying $\varphi(k, x_k)$, $\forall k$. By definition, for any k there is a set $\delta_k \in \Omega'$ such that $x_k \in \text{HOD}[\mathbf{a}[G] \upharpoonright \delta_k]^{\mathbf{L}[G]}$ (meaning that only $\mathbf{a}[G] \upharpoonright \delta_k$ and ordinals are admitted as parameters), and the sequence $\langle \delta_k \rangle_{k < \omega}$ belongs to $\mathbf{L}[G]$ as well. Furthermore, as the forcing

relation is definable in \mathbf{L} , there exist sequences $\langle X_k \rangle_{k < \omega} \in \mathbf{L}$ of conditions $X_k \in \mathbb{P}$ (possibly $X_k \notin G$), and $\langle \tau_k \rangle_{k < \omega} \in \mathbf{L}$ of sets $\tau_k \in \Omega'$, such that

$$X_k \Vdash \exists x \in \text{HOD}[\mathbf{a}[\underline{G}] \upharpoonright \tau_k] (\mathfrak{M}'_{\underline{G}} \models \varphi(k, x)). \quad (4)$$

Now, **arguing in \mathbf{L}** , we let $\zeta_k = \|X_k\|$, $\eta_k = \zeta_k \cup \tau_k$, and $\zeta^* = \|X^*\|$. Thus ζ^* and all τ_k, ζ_k, η_k belong to Ξ . Clearly there exists a sequence of permutations $\pi_k \in \Pi$ (see Section 5), $k < \omega$, such that the sets $\eta'_k = \pi_k''\eta_k = \{\pi_k(i) : i \in \eta_k\} \in \Xi$ are pairwise disjoint and disjoint with ζ^* .

Let $X'_k = \pi_k X_k$, so that $X'_k \in \text{Perf}_{\zeta'_k}$ in \mathbf{L} by Lemma 6, where $\zeta'_k = \pi_k''\zeta_k = \{\pi_k(i) : i \in \zeta_k\} \subseteq \eta'_k$. Define $\zeta = \zeta^* \cup \bigcup_k \zeta'_k$; $\zeta \in \Xi$. It follows by Corollary 2 that the set $X' = (X^* \upharpoonright^{-1} \zeta) \cap \bigcap_k (X'_k \upharpoonright^{-1} \zeta)$ belongs to Perf_{ζ} and $X' \leq X^*$, $X' \leq X'_k$ for all k .

On the other hand, the sets $\tau'_k = \pi_k''\tau_k$ belong to Ω' (because so do τ_k) and are pairwise disjoint (because so are the sets $\eta'_k = \zeta'_k \cup \tau'_k$). However Ω' is closed in \mathbf{L} under countable disjoint union, hence $\tau' = \bigcup_k \tau'_k \in \Omega'$.

We still **work in \mathbf{L}** . Starting with (4) and arguing as in the proof of Lemma 11 (the proof of 3 on page 11), we deduce that, for all k ,

$$X'_k \Vdash \exists x \in \text{HOD}[\mathbf{a}[\underline{G}] \upharpoonright \tau'_k] (\mathfrak{M}'_{\underline{G}} \models \varphi(k, x)),$$

and hence

$$X' \Vdash \forall k \exists x \in \text{HOD}[\mathbf{a}[\underline{G}] \upharpoonright \tau'] (\mathfrak{M}'_{\underline{G}} \models \varphi(k, x)), \quad (5)$$

because $X' \leq X'_k$ and $\tau'_k \subseteq \tau'$.

Finally, if H is \mathbb{P} -generic then the class $\text{HOD}[\mathbf{a}[H] \upharpoonright \tau']$ has a well-ordering, say \preceq_H , also $\{\mathbf{a}[H] \upharpoonright \tau'\}$ -ordinal-definable in $\text{HOD}[\mathbf{a}[H] \upharpoonright \tau']$. See e.g. [?], Section 13, the class $\text{HOD}[\mathbf{a}[H] \upharpoonright \tau']$ is identic to $\text{HOD}[\mathbf{a}[H] \upharpoonright \tau']$ as in [?]. Therefore, if H is any \mathbb{P} -generic set over \mathbf{L} containing X' , then, arguing on the basis of (5), we can define $y \subseteq \omega$ in \mathfrak{M}'_H such that, for each k , $(y)_k$ is equal to the \preceq_H -least set $x \subseteq \omega$ in $\text{HOD}[\mathbf{a}[H] \upharpoonright \tau']$, satisfying $\varphi(k, x)$. This proves that $\mathfrak{M}'_H \models \exists y \forall k \varphi(k, (y)_k)$ for any such H , and hence

$$X' \Vdash (\mathfrak{M}'_{\underline{G}} \models \exists y \forall k \varphi(k, (y)_k)).$$

But this contradicts (†) above since $X' \leq X^*$. \square

11. Models in which the parameter-free \mathbf{CA}^* holds but the full \mathbf{CA} fails

Here we sketch a proof of Theorem 2(i). See a full proof in our recent ArXiv preprint [?]. Thus the goal is to define a set $X \subseteq \mathcal{P}(\omega)$ in a cardinal-preserving generic extension of \mathbf{L} , which is a model of \mathbf{PA}_2^* (with the parameter-free Comprehension \mathbf{CA}^*) in which the full \mathbf{CA} fails.

Following the arguments above, assume that $G \subseteq \mathbb{P}$ is a set \mathbb{P} -generic over \mathbf{L} , define $\mathbf{a}_i[G] \subseteq \omega$ ($i \in I$) and the array $\mathbf{a}[G] = \langle \mathbf{a}_i[G] \rangle_{i \in I}$ as above, and consider the set

$$J[G] = \{\gamma \wedge 0^n : \gamma < \omega_1 \wedge n < \omega\} \cup \{\gamma \wedge 0^n \wedge 1 : \gamma < \omega_1 \wedge n \in \mathbf{a}_{\gamma \wedge 1}[G]\}.$$

$$\text{Here } \gamma \wedge 0^n = \langle \gamma, \underbrace{0, \dots, 0}_{n \text{ 0s}} \rangle, \gamma \wedge 0^n \wedge 1 = \langle \gamma, \underbrace{0, \dots, 0}_{n \text{ 0s}}, 1 \rangle, \gamma \wedge 1 = \langle \gamma, 1 \rangle.$$

Thus $J[G] \subseteq I$ and $J[G] \in \mathbf{L}[G]$. (Not necessarily $J[G] \in \mathbf{L}$.) We put

$$M_G = \mathcal{P}(\omega) \cap \bigcup_{i_1, \dots, i_n \in J[G]} \mathbf{L}[a_{i_1}[G], \dots, a_{i_n}[G]]; \quad M_G \subseteq \mathcal{P}(\omega).$$

Theorem 8. *If a set $G \subseteq \mathbb{P}$ is \mathbb{P} -generic over \mathbf{L} then M_G is a model of \mathbf{PA}_2^* (with the parameter-free Comprehension \mathbf{CA}^*) in which the full $\mathbf{CA}(\Sigma_2^1)$ holds but the full $\mathbf{CA}(\Sigma_4^1)$ fails.*

Proof (sketch, see [?] for a full proof). That M_G is a model of $\mathbf{CA}(\Sigma_2^1)$ (with parameters) follows by the Shoenfield absoluteness theorem, because M_G is Gödel-closed downwards

by construction. That the parameter-free \mathbf{AC}_ω^* holds in M_G follows by the ordinary permutation technique by a method rather similar to the verification of \mathbf{AC}_ω^* in the proof of Theorem 7 above.

Finally, M_G fails to satisfy the full **CA**. Indeed the reals $\mathbf{a}_{\gamma 1}[G]$ ($\gamma < \omega_1$) do not belong to M_G , since $\gamma \cap 1 \notin J[G]$ by construction. On the other hand, each $\mathbf{a}_{\gamma \cap 1}[G]$ is analytically definable in M_G as the set containing the numbers $n \geq 1$ such that the structure of true \leq_L -successors above $\mathbf{a}_{\langle \gamma \rangle}[G]$ has a split at n -th level, and possibly containing or not containing 0. Note the role of $\mathbf{a}_{\langle \gamma \rangle}[G] \in M_G$ as a parameter in this definition of $\mathbf{a}_{\gamma \cap 1}[G]$ in M_G . The ensuing definability formula for $\mathbf{a}_{\gamma \cap 1}[G]$ is Σ_4^1 by direct estimation, because it is based on the Π_3^1 definability of the relation of ‘being a true \leq_L -successor’. \square

Another model of \mathbf{PA}_2^* , in which **CA** fails even in the most elementary form of the nonexistence of complements of some its members, is also presented in [?]. It has the form $M = (\mathcal{P}(\omega) \cap \mathbf{L}) \cup \{y_n : n < \omega\}$, where $\langle y_n \rangle_{n < \omega}$ is a Cohen-generic sequence over \mathbf{L} . Note that the complements $y'_n = \omega \setminus y_n$ are **not** adjoined to M , so that **CA** is violated in M even in the form $\exists x \forall k (k \in x \iff k \notin y_n)$, with y_n as a parameter. On the other hand, the parameter-free \mathbf{CA}^* holds in M by ordinary permutation arguments.

12. Working on the basis of the consistency of \mathbf{PA}_2

This section is devoted to claims (ii) of our main Theorems 1, 2, 3. We recall that the consistency of \mathbf{PA}_2 is a common assumption in claims (ii). As the proofs of claims (i) of the theorems, given above, contain a heavy dose of the forcing technique, first of all we have to adequately replace \mathbf{PA}_2 with a more **ZFC**-like, forcing-friendly theory. This will be \mathbf{ZFC}^- , a subtheory of **ZFC** obtained as follows:

- (a) the Power Set axiom **PS** is excluded;
- (b) the Axiom of Choice **AC** is replaced with the wellorderability axiom **WA** saying that every set can be wellordered;
- (c) the Replacement schema, which is not sufficiently strong in the absence of **PS**, is replaced with the Collection schema;

See, e.g., [?] for a comprehensive account of main features of \mathbf{ZFC}^- .

Two more principles are considered in the context of \mathbf{ZFC}^- , namely

HC: every set is finite or countable,

V = L: every set is Gödel-constructible, i.e., the axiom of constructibility.

Theorem 9. *Theories \mathbf{PA}_2 and $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$ are equiconsistent. In fact they are interpretable in each other.*

Proof. This has been a well-known fact since while ago, see e.g. Theorem 5.25 in [?]. A more natural way of proof is as follows.

Firstly the theory \mathbf{Z}^- (i.e., \mathbf{ZFC}^- without **WA** and Collection) is interpreted in \mathbf{PA}_2 by the *tree interpretation* described e.g. in [?], §5, especially Theorem 5.11, or in [?], Definition VII.3.10 ff. Kreisel [?], VI(a)(ii), attributed this interpretation to the category of “crude” results. Secondly the whole theory $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$ is interpreted in \mathbf{Z}^- by means of the same tree interpretation, but restricted to only those trees that define sets constructible below the first *gap ordinal*, see a rather self-contained proof in [?]. This second part belongs to the category of “delicate” results of Kreisel [?], VI(b)(ii) \square

Theorem 9 allows us to replace the consistency of \mathbf{PA}_2 in claims (ii) of our Theorems 1, 2, 3 by the equivalent consistency of \mathbf{ZFC}^- , which is a much more forcing-friendly theory.

This makes it possible to argue in the frameworks of \mathbf{ZFC}^- in the following proof of Theorem 3(ii). The proof is an adaptation of the proof of the statement (i) of the same Theorem 3, on the basis of $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$.

Proof of Claims (ii) of Theorems 1, 2, 3. We argue on the basis of $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$. In other words, all sets are countable and constructible, so that the ground universe behaves like \mathbf{L}_{ω_1} in many ways. Yet, to avoid unnecessary misunderstanding, we accept the following.

Definition 2. The ground universe of $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$ is denoted by \mathbf{L}^- . Accordingly ω_1^- will be the collection (a proper \mathbf{L}^- -class) of all ordinals in \mathbf{L}^- . \square

Emulating the construction in Section 5, we define proper classes $\mathbf{I} = (\omega_1^-)^{<\omega} \setminus \{\Lambda\}$ and $\mathbf{\Xi}$, and sets \mathbf{IS}_ξ , $\xi[\subset \mathbf{i}]$, $\xi[\not\subset \mathbf{i}]$, etc., similar to Section 5. But coming to Definition 1, we face a problem. Indeed, each space $\mathcal{P}(\omega)^\xi$ and any homeomorphism $H : \mathcal{P}(\omega)^\xi \rightarrow \mathcal{P}(\omega)^\xi$ is now a proper class, hence \mathbf{Perf}_ξ as by Definition 1 is a class of proper classes, which cannot be considered. Therefore we have to *parametrize* homeomorphisms by sets.

Definition 3 (\mathbf{ZFC}^- form of Definition 1). Arguing in \mathbf{L}^- , let $\xi \in \mathbf{\Xi}$. Define

$$Q_\xi = \{x \in \mathcal{P}(\omega)^\xi : \text{the set } \{\langle i, k \rangle : x(i)(k) = 1\} \text{ is finite}\};$$

this is a countable dense subset of $\mathcal{P}(\omega)^\xi$ in \mathbf{ZFC}^- .

Let $h : Q_\xi \rightarrow \mathcal{P}(\omega)^\xi$ be any map (a set in \mathbf{L}^-). Let $[h]$ be its extension defined on $\mathcal{P}(\omega)^\xi$ by $[h](x) = \lim_{y \rightarrow x} h(y)$ whenever the limit exists, so $[h] : \text{dom}[h] \rightarrow \mathcal{P}(\omega)^\xi$ is a continuous map defined on $\text{dom}[h]$, a topologically closed “subset” or rather subclass of $\mathcal{P}(\omega)^\xi$ (also a proper class).

We define \mathcal{H}_ξ to be the class of all maps $h : Q_\xi \rightarrow \mathcal{P}(\omega)^\xi$ such that $\text{dom}[h] = \mathcal{P}(\omega)^\xi$, $[h]$ is 1 – 1 and $[h]$ is a projection-keeping homeomorphism.

Finally if $h \in \mathcal{H}_\xi$ then let $X_h = [h]''\mathcal{P}(\omega)^\xi = \{[h](x) : x \in \mathcal{P}(\omega)^\xi\}$.

Then $\mathbf{Perf}_\xi^- = \mathcal{H}_\xi$ and $\mathbf{Perf}^- = \bigcup_{\xi \in \mathbf{\Xi}} \mathbf{Perf}_\xi^-$ are proper classes, of course. \square

It is quite obvious that in the \mathbf{ZFC} setting \mathbf{Perf}_ξ coincides with the collection of all sets X_h , $h \in \mathcal{H}_\xi$. This allows us to use the map $h \rightarrow X_h$ as a *parametrization* of \mathbf{Perf} in \mathbf{L}^- , so that \mathbf{Perf}^- is the set of codes for the \mathbf{Perf} and each particular $\mathbf{Perf}_\xi^- = \mathcal{H}_\xi$ is the set of codes for \mathbf{Perf}_ξ . We will use \mathbf{Perf}^- as a **forcing notion**, that is, put $\mathbb{P}^- = \mathbf{Perf}^-$, with the order $g \leq h$ iff $X_g \leq X_h$ in the sense of Section 5.

Note that both \mathbb{P}^- and the order are definable proper classes in \mathbf{L}^- .

Conditions $h \in \mathbb{P}^-$ should be informally identified with corresponding objects (parametrically defined proper classes) X_g .

The property $(*)$ in the proof of Theorem 4 transforms to the following property of the forcing \mathbf{Perf}^- has a property in \mathbf{L}^- :

$(*)^-$ if a parametrized sequence of classes $D_n \subseteq \mathbf{Perf}^-$ is such that each D_n is open dense in \mathbf{Perf}^- , and $X \in \mathbf{Perf}$, then there is a stronger condition $Y \in \mathbf{Perf}$, $Y \leq X$, and finite sets $D'_n \subseteq D_n$ *pre-dense* in \mathbf{Perf}^- below Y .

In other words, \mathbf{Perf}^- is a *pretame* forcing notion in \mathbf{L}^- in the sense of [?] or [?].

It follows (see e.g. [?]) that any \mathbf{Perf}^- -generic extension of \mathbf{L}^- is still a model of \mathbf{ZFC}^- , and the forcing and definability theorems hold similar to the case of usual set-size forcing. Furthermore all constructions and arguments involved in the proofs of Theorems 6, 7, 8 above (i.e., claims (i) of Theorems resp. 3, 1, 2), as well as the results of [?] cited in the course of the proofs, can be reproduced *mutatis mutandis* on the basis of the theory $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$. In particular, Theorem 6 takes the form asserting that the $\mathcal{P}(\omega)$ -part of a certain subextension of any \mathbb{P}^- -generic extension of \mathbf{L}^- satisfies $\mathbf{PA}_2 + \neg \mathbf{AC}_\omega^*(\Pi_3^1)$.

Metamathematically, this means that the formal consistency of $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$ implies the consistency of $\mathbf{PA}_2 + \neg \mathbf{AC}_\omega^*(\Pi_3^1)$. However the consistency of $\mathbf{ZFC}^- + \mathbf{HC} + (\mathbf{V} = \mathbf{L})$ is equivalent to the consistency of \mathbf{PA}_2 by Theorem 9. This concludes the proof of Claim (ii) of Theorem 3.

Pretty similarly, Theorems 7 and 8 take appropriate forms sufficient to infer the consistency of resp.

$$\mathbf{PA}_2 + \mathbf{AC}_\omega^* + \neg \mathbf{AC}_\omega(\Pi_3^1), \quad \mathbf{PA}_2^* + \mathbf{CA}(\Sigma_2^1) + \neg \mathbf{CA}(\Pi_4^1),$$

from the consistency of \mathbf{PA}_2 , as required.

(Claims (ii) of Theorems 1, 2, 3) \square

13. Conclusions, remarks, and problems

In this study, the method of generalized arboreal iterations of the Sacks forcing is employed to the problem of obtaining cardinal-preserving models of \mathbf{ZFC} , and models of \mathbf{ZFC}^- and the second-order Peano arithmetic \mathbf{PA}_2 , in which the parameter-free version of the Comprehension or Choice schema holds but the full schema fails. These results (Theorems 1, 2, 3 above) contribute to the ongoing study of both subsystems and extensions of \mathbf{PA}_2 as in [?], [?], [?] among many others, as well as to modern studies of forcing extensions in class theories and \mathbf{ZFC}^- -like theories as in [?], [?], [?], [?].

From our study, it is concluded that the technique of generalized arboreal iterations of the Sacks forcing succeeds to solve important problems in descriptive set theory and second-order Peano arithmetic related to parameter-free versions of such crucial axiom schemata as Comprehension and Choice, by our Theorems 1, 2, 3.

From the results of this paper, the following remarks and problems arise.

Remark 5. Identifying the theories with their deductive closures, we may present the concluding statements of Theorems 1, 2, 3 as resp.

$$\mathbf{PA}_2 + \mathbf{AC}_\omega^* \subsetneq \mathbf{PA}_2 + \mathbf{AC}_\omega, \quad \mathbf{PA}_2^* + \mathbf{CA}(\Sigma_2^1) \subsetneq \mathbf{PA}_2, \quad \mathbf{PA}_2 \subsetneq \mathbf{PA}_2 + \mathbf{AC}_\omega^*. \quad (6)$$

Studies on subsystems of \mathbf{PA}_2 have discovered many cases in which $S \subsetneq S'$ holds for a given pair of subsystems S, S' , see e.g. [?]. And it is a rather typical case that such a strict extension is established by demonstrating that S' proves the consistency of S . One may ask whether this is the case for the results in (6). The answer is in the negative: namely

the theories \mathbf{PA}_2^ , $\mathbf{PA}_2^* + \mathbf{CA}(\Sigma_2^1)$, and the full \mathbf{PA}_2 are equiconsistent*

by a result in [?], also mentioned in [?]. This equiconsistency result also follows from a somewhat sharper theorem in [?], 1.5. \square

Remark 6. There is another meaningful submodel of the basic model $\mathbf{L}[G] = \mathbf{L}[\mathbf{a}[G]]$. Namely, consider the set W'' of all finite or countable **well-founded** initial segments $\zeta \in \mathbf{L}$, $\zeta \subseteq I$, instead of the sets W (as in Section 8) and W' (as in Section 10). Define a corresponding submodel \mathfrak{M}_G'' accordingly. Then \mathbf{AC}_ω holds in \mathfrak{M}_G'' but $\mathbf{DC}(\Pi_3^1)$ fails. Yet a better model is defined in [?], in which \mathbf{AC}_ω holds but even $\mathbf{DC}(\Pi_2^1)$ (the best possible in this case) fails. \square

We proceed with a list of open problems.

Problem 1. Is the parameter-free countable choice schema \mathbf{AC}_ω^* in the language $\mathcal{L}(\mathbf{PA}_2)$ true in the models defined in Section 11?

Problem 2. Can we sharpen the result of Theorem 8 by specifying that $\mathbf{CA}(\Sigma_3^1)$, rather than Σ_4^1 , is violated? The combination $\mathbf{CA}(\Sigma_2^1)$ plus $\neg \mathbf{CA}(\Sigma_3^1)$ over \mathbf{PA}_2^* would be optimal for Theorem 2. Can we similarly sharpen the result of Theorems 6 and 7 by specifying that $\mathbf{AC}_\omega^*(\Sigma_2^1)$, resp., $\mathbf{AC}_\omega(\Sigma_2^1)$ are violated? As suggested by V. Gitman, Jensen's iterated forcing introduced in [?] may lead to a solution.

Problem 3. As a generalization of Problem 2, prove that, for any $n \geq 2$, $\mathbf{PA}_2^* + \mathbf{CA}(\Sigma_n^1)$ does not imply $\mathbf{CA}(\Sigma_{n+1}^1)$. In this case, it would be possible to conclude that the full

schema **CA** is not finitely axiomatizable over PA_2^* . There are similar questions related to Theorems 6 and 7, of course. Compare to Problem 9 in [? , §11]. We expect that methods of inductive construction of forcing notions in **L** that carry *hidden automorphisms*, recently developed in our papers [? ? ? ? ?], may lead to solutions.

Problem 4 (Communicated by Ali Enayat). A natural question is whether the results of this note also hold for second order set theory (the Kelley-Morse theory of classes). This may involve a generalization of the Sacks forcing to uncountable cardinals, as in Kanamori [?], and new models of set theory recently defined by Fuchs [?].

Author Contributions: Conceptualization, V.K. and V.L.; methodology, V.K. and V.L.; validation, V.K.; formal analysis, V.K. and V.L.; investigation, V.K. and V.L.; writing original draft preparation, V.K.; writing review and editing, V.K. and V.L.; project administration, V.L.; funding acquisition, V.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable. The study did not report any data.

Acknowledgments: We thank Ali Enayat, Gunter Fuchs, Victoria Gitman, and Kameryn Williams, for their enlightening comments that made it possible to accomplish this research, and separately Ali Enayat for references to [? ? ?] in matters of Theorem 3.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kreisel, G. A survey of proof theory. *J. Symb. Log.* **1968**, *33*, 321–388. <https://doi.org/10.2307/2270324>.
2. Levy, A. Definability in axiomatic set theory II. In *Proceedings of the Math. Logic Found. Set Theory, Proc. Int. Colloqu., Jerusalem 1968*; Bar-Hillel, Y., Ed.; North-Holland: Amsterdam-London, 1970; pp. 129–145. [https://doi.org/10.1016/S0049-237X\(08\)71935-9](https://doi.org/10.1016/S0049-237X(08)71935-9).
3. Solovay, R.M. A model of set-theory in which every set of reals is Lebesgue measurable. *Ann. Math. (2)* **1970**, *92*, 1–56. <https://doi.org/10.2307/1970696>.
4. Guzicki, W. On weaker forms of choice in second order arithmetic. *Fundam. Math.* **1976**, *93*, 131–144. <https://doi.org/10.4064/fm-93-2-131-144>.
5. Corrada, M. Parameters in theories of classes. *Mathematical logic in Latin America, Proc. Symp., Santiago 1978*, 121–132 (1980)., 1980. [https://doi.org/10.1016/S0049-237X\(09\)70483-5](https://doi.org/10.1016/S0049-237X(09)70483-5).
6. Levy, A. Parameters in comprehension axiom schemes of set theory. *Proc. Tarski Symp., internat. Symp. Honor Alfred Tarski, Berkeley 1971, Proc. Symp. Pure Math.* **25**, 309–324 (1974)., 1974.
7. Ralf Schindler and Philipp Schlicht. ZFC without parameters (A note on a question of Kai Wehmeier). https://ivv5hpp.uni-muenster.de/u/rds/ZFC_without_parameters.pdf. Accessed: 2022-09-06.
8. Kanovei, V.; Lyubetsky, V. The parameterfree Comprehension does not imply the full Comprehension in the 2nd order Peano arithmetic. *arXiv e-prints* **2022**, p. arXiv:2209.07599, [arXiv:math.LO/2209.07599]. <https://doi.org/https://arxiv.org/abs/2209.07599>.
9. Enayat, A. On the Leibniz – Mycielski axiom in set theory. *Fundam. Math.* **2004**, *181*, 215–231. <https://doi.org/10.4064/fm181-3-2>.
10. Jensen, R. Definable sets of minimal degree. In *Math. Logic Found. Set Theory, Proc. Int. Colloqu., Jerusalem 1968*; Bar-Hillel, Y., Ed.; North-Holland: Amsterdam-London, 1970; Vol. 59, *Studies in logic and the foundations of mathematics*, pp. 122–128. [https://doi.org/10.1016/S0049-237X\(08\)71934-7](https://doi.org/10.1016/S0049-237X(08)71934-7).
11. Groszek, M.; Jech, T. Generalized iteration of forcing. *Trans. Amer. Math. Soc.* **1991**, *324*, 1–26. <https://doi.org/10.1090/S0002-9947-1991-0946221-X>.
12. Mathias, A.R.D. Surrealist landscape with figures (a survey of recent results in set theory). *Period. Math. Hung.* **1979**, *10*, 109–175. <https://doi.org/10.1007/BF02025889>.
13. Frittaion, E. A note on fragments of uniform reflection in second order arithmetic. *The Bulletin of Symbolic Logic* **2022**, pp. 1–16. <https://doi.org/10.1017/bsl.2022.23>.
14. Apt, K.R.; Marek, W. Second order arithmetic and related topics. *Ann. Math. Logic* **1974**, *6*, 177–229. [https://doi.org/10.1016/0003-4843\(74\)90001-1](https://doi.org/10.1016/0003-4843(74)90001-1).

- Simpson, S.G. *Subsystems of second order arithmetic*, 2nd ed.; Perspectives in Logic, Cambridge: Cambridge University Press; Urbana, IL: ASL, 2009. Pages xvi + 444.
- Friedman, H. On the necessary use of abstract set theory. *Advances in Mathematics* **1981**, *41*, 209–280. [https://doi.org/https://doi.org/10.1016/0001-8708\(81\)90021-9](https://doi.org/https://doi.org/10.1016/0001-8708(81)90021-9).
- Schindler, T. A disquotational theory of truth as strong as Z_2^- . *J. Philos. Log.* **2015**, *44*, 395–410. <https://doi.org/10.1007/s10992-014-9327-5>.
- Jech, T. *Set theory*, The third millennium revised and expanded ed.; Springer-Verlag: Berlin-Heidelberg-New York, 2003. Pages xiii + 769, <https://doi.org/10.1007/3-540-44761-X>.
- Kanovei, V. Non-Glimm-Effros equivalence relations at second projective level. *Fund. Math.* **1997**, *154*, 1–35. <https://doi.org/10.4064/fm-154-1-1-35>.
- Kanovei, V. On non-wellfounded iterations of the perfect set forcing. *J. Symb. Log.* **1999**, *64*, 551–574. <https://doi.org/10.2307/2586484>.
- Baumgartner, J.E.; Laver, R. Iterated perfect-set forcing. *Ann. Math. Logic* **1979**, *17*, 271–288. [https://doi.org/10.1016/0003-4843\(79\)90010-X](https://doi.org/10.1016/0003-4843(79)90010-X).
- Groszek, M.J. Applications of iterated perfect set forcing. *Ann. Pure Appl. Logic* **1988**, *39*, 19–53. [https://doi.org/10.1016/0168-0072\(88\)90044-9](https://doi.org/10.1016/0168-0072(88)90044-9).
- Myhill, J.; Scott, D. Ordinal definability. *Axiomatic Set Theory*, Proc. Sympos. Pure Math. 13, Part I, 271–278 (1971)., 1971.
- Gitman, V.; Hamkins, J.D.; Johnstone, T.A. What is the theory ZFC without power set? *Math. Log. Q.* **2016**, *62*, 391–406.
- Kanovei, V.G. Theory of Zermelo without power set axiom and the theory of Zermelo- Fraenkel without power set axiom are relatively consistent. *Math. Notes* **1981**, *30*, 695–702. <https://doi.org/10.1007/BF01141627>.
- Friedman, S.D. *Fine structure and class forcing*; Vol. 3, *De Gruyter Series in Logic and Its Applications*, de Gruyter: Berlin, 2000; pp. x + 221.
- Antos, C.; Gitman, V. Modern Class Forcing. In *Research Trends in Contemporary Logic*; Daghighi, A.; Rezus, A.; Pourmahdian, M.; Gabbay, D.; Fitting, M., Eds.; College Publications, forthcoming. [LINK](#), accessed: 2022-12-06.
- Enayat, A.; Schmerl, J.H. The Barwise-Schlipf theorem. *Proc. Am. Math. Soc.* **2021**, *149*, 413–416. <https://doi.org/10.1090/proc/15216>.
- Friedman, S.D.; Gitman, V.; Kanovei, V. A model of second-order arithmetic satisfying AC but not DC. *J. Math. Log.* **2019**, *19*, 1–39. Article No 1850013, <https://doi.org/10.1142/S0219061318500137>.
- Antos, C.; Friedman, S.D.; Gitman, V. Boolean-valued class forcing. *Fundam. Math.* **2021**, *255*, 231–254. <https://doi.org/10.4064/fm20-7-2021>.
- Gitman, V.; Hamkins, J.D.; Holy, P.; Schlicht, P.; Williams, K.J. The exact strength of the class forcing theorem. *J. Symb. Log.* **2020**, *85*, 869–905. <https://doi.org/10.1017/jsl.2019.89>.
- Holy, P.; Krapf, R.; Lücke, P.; Njegomir, A.; Schlicht, P. Class forcing, the forcing theorem and Boolean completions. *J. Symb. Log.* **2016**, *81*, 1500–1530. <https://doi.org/10.1017/jsl.2016.4>.
- Schmerl, J.H. Peano arithmetic and hyper-Ramsey logic. *Trans. Am. Math. Soc.* **1986**, *296*, 481–505. <https://doi.org/10.2307/2000376>.
- Kanovei, V.; Lyubetsky, V. Models of set theory in which nonconstructible reals first appear at a given projective level. *Mathematics* **2020**, *8*. Article No 910, <https://doi.org/10.3390/math8060910>.
- Kanovei, V.; Lyubetsky, V. On the Δ_n^1 problem of Harvey Friedman. *Mathematics* **2020**, *8*. Article No 1477, <https://doi.org/10.3390/math8091477>.
- Kanovei, V.; Lyubetsky, V. Models of set theory in which separation theorem fails. *Izvestiya: Mathematics* **2021**, *85*, 1181–1219. <https://doi.org/10.1070/IM8521>.
- Kanovei, V.; Lyubetsky, V. The full basis theorem does not imply analytic wellordering. *Ann. Pure Appl. Logic* **2021**, *172*, 46. Id/No 102929, <https://doi.org/10.1016/j.apal.2020.102929>.
- Kanovei, V.; Lyubetsky, V. A model in which the Separation principle holds for a given effective projective Sigma-class. *Axioms* **2022**, *11*. Article No 122, <https://doi.org/10.3390/axioms11030122>.
- Kanamori, A. Perfect-set forcing for uncountable cardinals. *Ann. Math. Logic* **1980**, *19*, 97–114. [https://doi.org/10.1016/0003-4843\(80\)90021-2](https://doi.org/10.1016/0003-4843(80)90021-2).
- Fuchs, G. Blurry Definability. *Mathematics* **2022**, *10* (3), Article No 452. <https://doi.org/10.3390/math10030452>.