

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Proof-of-concept of a novel multiepitope recombinant protein for the serodiagnosis of patients with Chagas Disease

Juliana Martins Machado^{1*}, Isabela Amorim Gonçalves Pereira^{2*}, Ana Clara Gontijo Maia^{1*}, Mariana Ferraz Chaves Francisco¹, Lais Moreira Nogueira¹, Isadora Braga Gandra¹, Anna Julia Ribeiro¹, Kamila Alves Silva¹, Carlos Ananias Aparecido Resende¹, Jonatas Oliveira da Silva¹, Michelli dos Santos¹, Ana Alice Maia Gonçalves², Grasiele de Sousa Vieira Tavares², Miguel Angel Chávez-Fumagalli³, Mariana Campos da Paz⁴, Rodolfo Cordeiro Giunchetti⁵, Manoel Otávio C. Rocha², Ana Thereza Chaves², Eduardo Antônio Ferraz Coelho², Alexsandro S. Galdino^{1*}

1 *Laboratório de Biotecnologia de Microrganismos, Universidade Federal de São João Del-Rei, Divinópolis, MG, Brazil.*

2 *Programa de Pós-Graduação em Ciências da Saúde: Infectologia e Medicina Tropical, Faculdade de Medicina, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil*

3 *Computational Biology and Chemistry Research Group, Vicerrectorado de Investigación, Universidad Católica de Santa María, Arequipa 04000, Peru*

4 *Laboratório de NanoBiotecnologia & Bioativos, Universidade Federal de São João Del-Rei, Sebastião Gonçalves Coelho, 400, 355901-296, Divinópolis, MG, Brazil*

5 *Laboratório de Biologia das Interações Celulares, Departamento de Morfologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, 31270-901, Belo Horizonte, MG, Brazil*

* Correspondence: A.S.G - Email address: asgaldino@ujs.edu.br - Laboratório de Biotecnologia de Microrganismos, Universidade Federal de São João Del-Rei, Divinópolis – 35501-296, Brazil.

Abstract: **Background:** Chagas disease is still a neglected disease considered a public health problem. Early diagnosis of cases is important to improve the prognosis of infected patients and prevent transmission. Serological tests are the method of choice for diagnosis. However, two serological tests are currently recommended to confirm positive cases. In this sense, more sensitive and specific serological tests need to be developed to overcome the problems currently faced. This study aimed to develop a new recombinant multiepitope protein for the diagnosis of Chagas disease, hereafter named rTC. **Methods:** The rTC was constructed based on amino acid sequences from different combinations of *Trypanosoma cruzi* antigens in the same polypeptide and tested in ELISA for detecting different Chagas disease. **Results:** rTC1 was able to discriminate between indeterminate (IND) and cardiac (CARD) cases and cross-reactive diseases, and healthy samples, with 96.6% sensitivity and 97.96% specificity, respectively. **Conclusion:** These data suggest a preliminary study that rTC1 has the potential to be tested in future studies against a larger serological panel for the diagnosis of Chagas disease.

Keywords: multiepitope; recombinant protein; immunodiagnosis; Chagas disease

1. Introduction

Chagas disease (CD) is a neglected tropical disease caused by the hemoflagellate *Trypanosoma cruzi* (*T. cruzi*) [1,2]. The infection occurs mainly by the vectorial route, transmitted to humans by hematophagous insects of the subfamily *Triatominae* [3]. However, the parasite can also be transmitted congenitally, by blood transfusions, organ transplantation, or the ingestion of contaminated food and beverages [3]. It is considered a public health problem and there has been an increase in its global spread in recent years. Currently, the disease is endemic in 21 regions, and it is occurring in non-endemic countries, such as the United States of America, Canada, Spain, France, Switzerland, Italy, Japan, and other continents such as Asia and Oceania [1,4].

According to the Pan American Health Organization (PAHO) about six to eight million people are infected with *T. cruzi* in the Americas, while about 70% of these people are living with the disease without knowing they are infected due to the absence of an appropriate diagnosis [5]. More than 10,000 people die every year from clinical complications of Chagas disease and 75 million people are at risk of infection [4,6]. In Brazil, new cases

were reported in most states, with 95% concentrated in the northern region, with 83% of the cases reported in Pará state [7].

T. cruzi infection in humans results in an acute illness lasting six to eight weeks, which, if untreated, may evolve into a chronic phase [8–10]. In the acute phase, a local inflammatory lesion appears at the site of infection where the parasite undergoes the first rounds of replication [11,12]. Afterward, the parasites spread through the body, circulating trypomastigotes being easily found in the blood. During the chronic phase, circulating parasites cannot be detected by inspection of blood but progressive tissue damage ensues, often with serious cardiac manifestations or gastrointestinal involvement. The chronic phase is characterized by the high production of specific immunoglobulin G (IgG) antibodies, and diagnosis at this stage relies mainly on serological techniques [13,14].

Parasitological and molecular diagnostic tests are the most appropriate in the acute phase. However, direct parasitological tests are not recommended in the chronic phase due to the absence of parasitemia [15–17]. Direct parasitological tests, such as thick, fresh blood, and smears, are based on an analysis of the parasite in the blood of infected patients. But these tests have low sensitivity and can generate false-negative results [18]. In addition, indirect parasitological tests, such as xenodiagnoses or blood culture, have low sensitivity varying between 20% and 50% and require assistance from other methods, such as the concentration test, to confirm the result. The ELISA and Polymerase Chain Reaction (PCR) tests are indicated for the chronic phase because they are more sensitive and specific than parasitological tests [19]. Despite its high sensitivity and specificity, PCR is a costly test and difficult to standardize [20].

Immunoglobulin G antibody detection is the method of choice for diagnosis in the chronic phase of the disease because parasitemia is scarce. However, two different serological methods are recommended for diagnosis confirmation. Usually, the enzyme-linked immunosorbent assay (ELISA), indirect immunofluorescence, or indirect hemagglutination are employed [21]. ELISA techniques are generally based on crude extracts or antigens purified of epimastigote or trypomastigote forms. In addition, recombinant antigens, either alone or combined, and synthetic peptides are also widely used [22,23]. This assay is a widely used method for screening donors at blood banks in Brazil [24]. However, a gold standard for serological diagnosis has not yet been established. According to the PAHO, at least two different serological tests or two different *T. cruzi* antigens are required as serological discrepancies are common [25–27].

The chronic phases of CD can be divided into four clinical forms: i) indeterminate form, ii) cardiac form, iii) digestive form, and iv) mixed form when cardiac and digestive manifestations coexist simultaneously [28]. Individuals in the indeterminate form have a significant risk of developing chronic cardiomyopathy [29], and about 40% of individuals with chronic diseases may develop heart damage, digestive diseases, or both manifestations [30]. Therefore, the development of new markers capable of recording the recognition of different clinical forms and assessing the potential for progression and monitoring of the most severe form may contribute to the early adoption of therapeutic therapies [31,32].

The use of assays that employ a combination of recombinant antigens and the incorporation of different detection systems increase the sensitivity and specificity of the serological techniques [33]. The major advantage of recombinant antigen-based tests is that they minimize the extent of specificity problems [34,35]. The present study designed a recombinant multiepitope protein for diagnosing Chagas disease (rTC) and evaluated discrimination of clinical forms and cross-reactivity. The findings demonstrated that rTC can diagnose *T. cruzi*-infected individuals with low cross-reactivity and could be a valuable tool for producing immunodiagnostic kits for Chagas disease.

2. Materials and Methods

2.1 Strains, reagents, and samples

Escherichia coli (E. coli) BL21 (λ DE3) and expression vector pET21a were purchased from Novagen. HIS-Select® Nickel Affinity Gel resin was used to purify rTC. Antibody-enzyme conjugates [monoclonal anti-poly histidine-Horseradish Peroxidase (HRP) antibody and monoclonal anti-human IgG-HRP] were purchased from Sigma. Other reagents of analytical grade were obtained from standard commercial sources.

2.2 Serum samples

The study was approved by the Human Research Ethics Committee of UFMG, with protocol numbers: CAAE-32343114.9.0000.5149 / CAAE-48354315.8.0000.5149. CD-positive serum samples (n=58) were classified according to the clinical form of the disease as described by Rocha et al. (2003) [36]: indeterminate form (IND, n=30) for asymptomatic individuals who have sinus rhythm, without significant changes in the electrocardiogram, chest radiograph, and echocardiogram; and cardiac form 5 (CARD, n= 28) for individuals with echocardiographic signs of dilated left ventricle with systolic function impaired ventricular function, with or without manifestations of heart failure. Healthy individuals (n=30) from a non-endemic area for the CD with negative characterization tests and individuals infected with Leishmaniasis, visceral (n=30), and cutaneous (n=30), for cross-reactivity assessment, were included as the control group (NI). All sera were collected and characterized by two or more tests (indirect immune fluorescence, ELISA, or indirect hemagglutination).

2.3 Design of the synthetic gene, cloning, and expression

The synthetic gene was custom synthesized by Genone with codon usage for *Escherichia coli* (*E. coli*) and cloned as a *NdeI*/*XhoI* fragment into pET21a in-frame with a C-terminal histidine tag to allow protein purification by affinity chromatography. The resulting plasmid was used to transform *E. coli* BL21 (DE3) *plysS* competent cells and selection was performed on LB agar plates containing 100 μ g/mL ampicillin. The DNA and amino acid sequences for the entire synthetic gene construct are proprietary (under Brazilian patent No. BR1020220183139) and are not available to be shared at this stage. An individual colony was inoculated in 5 mL LB (10 g/L Casein Peptone, 5 g/L Yeast extract, 10 g/L NaCl, pH 7.2) containing 100 g/mL ampicillin and allowed to grow overnight at 37°C under agitation (200 rpm). At one point twenty-five mL of the pre-culture was transferred to 25 mL LB in a 250 mL Erlenmeyer. The culture was grown in the conditions until an OD600 of 0.6 at which point 1 mM IPTG was added. Aliquots were withdrawn at 0.5, 1.5, and 2.5 h after induction. The induced culture was harvested by centrifugation at 6000 \times g for 15 min at 4°C and the pellet was stored at -80°C.

2.4 Purification of the rTC

The purification strategy involved the addition of a lysis buffer (8 M urea, 50 mM NaH₂PO₄, 300 mM NaCl, 10 mM imidazole, and pH 8.0) to the pellet following incubation at 4°C overnight. The system was then sonicated, and 500 μ L Ni-NTA resin (Sigma) was added to the supernatant for batch purification. The resin was washed (8 M urea, 50 mM NaH₂PO₄, 300 mM NaCl, 5 mM imidazole, and pH 8.0) and proteins eluted with 8 M urea, 50 mM NaH₂PO₄, 300 mM NaCl, and 100 mM imidazole (pH 8.0).

2.5 Electrophoresis and Western Blotting

Protein integrity and molecular mass calculation were evaluated by running samples on 12% SDS-PAGE. Proteins were stained with Coomassie Brilliant Blue R-250 (Sigma-Aldrich). Following electrophoresis, the proteins were transferred electrophoretically to a

Polyvinylidene difluoride (PVDF) membrane for western blotting. The membrane was blocked with 5% skim milk powder in phosphate-buffered saline (PBS) for 2 h at room temperature. It was then washed three times with PBS containing 0.1% tween 20 and incubated with monoclonal mouse anti-His HRP (Horseradish peroxidase, Sigma), diluted 1:1,000 in PBS for 2 h at room temperature. Following three washes with PBST, the specific protein band was visualized by the 3,3'-Diaminobenzidine tetrahydrochloride detection method.

2.6 ELISA assay

The immune enzymatic assay was used to detect specific binding of the recombinant protein rTC to anti-CD antibodies. The wells of the polystyrene plates (Costar) were sensitized with 8.75 ng purified rTC dissolved in 100 μ l 0.1 M sodium carbonate-bicarbonate buffer (pH 9.6). After incubation at 4°C for 16 h, the coated wells were washed with PBST and blocked for 1 h at 37°C with PBS buffer (pH 7.2) containing 5% (w/v) dried skim milk powder and washed again with PBS-T. Subsequently, 100 μ l of a dilution (100 μ l PBST, 5% [w/v] dried skim milk powder, and 1 μ l serum) was placed into the wells resulting in a final dilution of approximately 1:100. After incubation for 30 min at 37°C, the wells were washed with PBST and 100 μ l of peroxidase-labeled goat either anti-human immunoglobulin G conjugate (Sigma) and diluted at 1:40,000 in PBS buffer (pH 7.2) was added following incubation for 30 minutes at 37°C. The wells were again washed, and 100 μ l 3,3,5,5-tetramethyl-benzidine substrate (BD Biosciences) was added and incubated for 15 minutes at room temperature in the dark. Reactions were stopped by the addition of 100 μ l sulfuric acid (0.2 M). The optical densities were read at 450 nm.

2.6 Statistical Analyses

The cut-off values of the *in-house* ELISA assay were established for optimal sensitivity, specificity, and accuracy (area under the curve) using receiver operating characteristic (ROC) curve analysis in GraphPad Prism. The absorbance readings produced by the ELISA assay were compared using the Mann-Whitney U test, and the Kruskal-Wallis test considering a level of significance of $p < 0.05$.

3. Results

3.1. Design, Expression, and Purification of the rTC

To design a multiepitope protein that could be of diagnostic use of conserved epitopes which are known to elicit anti-CD antibodies were selected based on data from the literature. The antigenic epitopes chosen were assembled and connected by flexible glycine-serine linkers. This would allow the epitopes to be freely available for interaction with their cognate antibodies thus contributing to the overall sensitivity and specificity of the diagnostic test. To increase epitope density, the selected epitopes were joined in tandem, resulting in a protein of ~50 kDa called rTC, an SDS-PAGE 12% gel is shown (Fig 1a). After purification (Fig. 2), a western blotting experiment was conducted using an anti-histidine antibody to confirm the presence of rMELEISH protein (Fig 1b).




Figure 1. Time course rTC expression in *E. coli* BL21 (λDE3) and analysis by SDS-PAGE 12%. (a) Lane M, unstained protein molecular mass marker (Thermo Scientific); Lane 1, non-induced control; Lanes 2-4, after 0.5, 1.5 and 2.5 hours of induction, respectively. (b) Western blotting analysis. The monoclonal antibody, anti-polyhistidine conjugated horseradish peroxidase (HRP), was diluted 1:1000 in PBS.




Figure 2. Analysis of the rTC by 12% SDS-PAGE, after the purification procedure. Fractions were collected after Ni-NTA chromatography. Lane M, molecular mass markers: Unstained Protein Weight Marker Standards (Thermo Scientific). Lane 1, flow-through; Lanes 2-4, wash steps; Lanes 5-7 purified rTC after elution with 100 mM imidazole.

3.2. ELISA - Human anti-IgG Chagas Recognized rTC

Since the underlying goal of this study was the development of a recombinant multipeptipeptide protein that could be used in a Chagas diagnosis kit, the ability of rTC to detect anti-CD antibodies was tested in an in-house ELISA. The cut-off value of 0.063 was defined by the analysis of the ROC curve and the area under the curves was 0.97 (95% CI 0.98 to 1.0) (Fig. 3a). The median of the absorbance readings produced by the in-house ELISA presented a statistically significant difference ($p < 0.0001$) between the positive and negative groups (Fig. 3b). rTC1 was able to detect both indeterminate and cardiac forms

with a minimal cross-reaction for LVH and LCH (Fig.3c). The sensitivity and specificity values were 96.67% (57/58) and 97.96% (88/90), respectively, and the positive likelihood ratio presented by assay was 47.37. These results show the ability of rTC to distinguish positive and negative human sera for IgG and its potential for use in diagnosis kits.




Figure 3: (a) ROC curve showing the in-house ELISA's area under the curve (AUC). (b) Boxplots of the absorbance readings from the sera in the positive and negative groups presented statistically significant differences (*Mann-Whitney U: $p < 0.0001$). The median of absorbance readings showed by *in-house* ELISA in the positive group was 0.969, 25% percentile = 0.825, 75% percentile = 1.059. In contrast, the median of absorbance readings showed by *in-house* ELISA in the negative group was 0.012, 25% percentile = 0.004, 75% percentile = 0.024 (Mann-Whitney U: $p < 0.0001$). (c) Reactivity and positivity of the *in-house* ELISA were measured in serum samples of positives groups (Cardiac Form (CARD) n=28 and Indeterminate Form (IND) n=30) and control negatives groups (NI), Healthy individuals (HI) n=30, Human visceral leishmaniasis (LVH) n=30 and Human cutaneous leishmaniasis LCH n=30). The cut-off was determined as 0.063. The sensitivity and specificity were 96.67% (95% CI: 90.65-99.09%) and 97.96% (95% CI: 89.31-99.90%) respectively. Two samples, from the negatives group, showed false-positive results. In the Kruskal-Wallis test, not observe no significant difference between the diagnosis ($p > 0.9999$) of IND and CARD clinical forms, only with NI ($p < 0.0001$).

4. Discussion

The diagnosis of CD is essential to ensuring that the patient will receive adequately chemotherapy. This paper describes the development of a recombinant multiepitope protein and shows its performance in the serological diagnosis of Chagas disease. rTC1 can detect individuals infected with Chagas disease for both IND and CARD forms, presenting 96.67% sensitivity, 97.96% specificity, and 0.975 area over the curve (AUC).

Currently, there is no gold standard for the diagnosis of CD, requiring at least two distinct serological assays. Different ELISA assays for detecting Chagas are commercially available, although variations in their performance when testing samples from different countries have been reported [37,38].

Different factors may be associated with the performance of the test, such as the intrinsic characteristic of each antigen and the use of the assays in countries and regions where the genetic strains differ from those used for the development of the antigen [39–41]. To overcome this problem, a multiepitope-based protein could be used as an alternative to crude antigens improving sensitivity, and specificity and reducing production costs [42,43]. The construction of such recombinant proteins containing high-density epitopes has been a preferred alternative for the diagnosis of diseases given that these proteins exhibit a broad ability to expose their epitopes more efficiently, resulting in improved sensitivity and specificity [42,43].

In this work, a multiepitope protein was developed and evaluated as a strategy for the diagnosis of CD, focusing on antigens and conserved proteins to construct the encoding rTC1. Such antigens were evaluated using information available in the literature, with the main B-cell epitopes being selected based on the following criteria: (i) immunodominance, (ii) specificity for anti-CD antibodies, (iii) straight chain, and (iv) phylogenetic conservation in most genotypes in circulation in different parts of the world [42–45].

Several multiepitope proteins for the diagnosis of different diseases, both human [42–44,46–50] and animal [51–53], have been described and show promising results in terms of increasing and improving sensitivity and specificity and reducing cross-reactivity, indicating that multiepitope technology is efficient for the detection of different pathologies. There are few studies in the literature using recombinant multiepitope to diagnose CD. A search conducted at Pubmed using the keywords: “Multiepitope protein” and “diagnosis” identified 270 published articles. Out of these, only four were related to the diagnosis of CD, showing sensitivity and specificity data [40,54–56]. However, none of them evaluated the potential of their antigens to detect both chagasic forms classified with the clinical form as done in this study.

Houghton et al., (2000) [54] described the use of a single multi-epitope peptide, 2/D/E, formed by sequences of antigens TcD, TcE, and PEP-2. This protein was highly sensitive, detecting 239 out of 240 consensus-positive sera, indicating a sensitivity of 99.6%. In contrast, of the 149-test serum sample from healthy, random donors, 148 were below the designated cutoff, indicating a specificity of 99.33%. Camussone et al., (2009) [55] reported the use of two multiepitope-based proteins, CP1 and CP2, tested separately as antigens to detect CD. The discrimination efficiency values obtained for CP1 and CP2 were 25% and 52% higher than those of their antigens. CP2 was the only multiepitope protein that showed discrimination efficiency between negative and positive samples of CD, showing 98.6 % sensitivity and 99.4% specificity. Duthie et al. (2013) [57] studied the use of multiepitope fusion proteins TcF, TcF43, TcF26 for the diagnosis of CD. The authors demonstrated that fusion proteins TcF26 and TcF43 yielded strong responses against sera that were also strongly detected by TcF, one of the recombinant antigens currently used with tests for CD. Conversely, Peverengo et al. (2018) [56] tested the use of multiepitope proteins CP1 and CP3 to diagnose chronic CD. The sensitivity of CP3 and CP1 was 100% and 90.2%, respectively, and specificity was 92.5% and 100%, respectively, showing that both multiepitope proteins detect chronic *T. cruzi* infection.

Despite not having an established treatment for chronic infections, the evaluation of treatment results depends on serological monitoring [58]. Therefore, the development of

new biomarkers is essential to identify the disease in its acute or chronic phase [34]. Non-parametric analysis was performed for more than two groups to determine whether rTC was able to distinguish between clinical forms, but no significant difference was observed between the diagnosis ($p>0.9999$) of IND and CARD clinical forms. However, the diagnosis of IND individuals is relevant for the management and care of CD. Intervention in the initial phase of Chagas disease (acute and undetermined phase) can favor the reduction of transmission, improve the health condition of infected individuals, and save money on the cost of treatment [36,59].

The results of the present study differ from the one conducted by Longhi et al, (2012) [60], in which a recombinant J7 protein and several peptides were evaluated for their diagnostic capacity considering clinical, electrocardiographic, and radiological results. They were then classified as indeterminate, (HF), digestive (CD), mild cardiomyopathy (MCC), and severe cardiomyopathy (SCC) forms of CD. The authors found that the anti-JL7 Ab concentration (by optical density [OD]) was higher in sera from patients with SCC as compared to those of IC ($P<0.019$), suggesting that this antigen could be used as a prognostic marker for the indeterminate form (HF).

Another important factor for an accurate diagnosis is the absence of cross-reactivity, especially between CD and Leishmaniasis. This study's results showed low cross-reactivity with visceral (1/30) and tegumentary (1/30) Leishmania. A similar result is described by Daltro et al. (2019) [61], in which four chimeric antigens were evaluated to assess cross-reactivity with American tegumentary or visceral leishmaniasis. The chimeric antigens exhibited poor seropositivity for leishmaniasis infection as compared to the evaluated commercial immunoassays.

5. Conclusions

In conclusion, in this preliminary work, rTC showed satisfactory performance in the diagnosis of CD in its chronic phase and individuals in the IND form and proved to be effective for possible application in Chagas diagnostic kits. The development of more accurate tests that consider the genetic variability of *T. cruzi* and present low cross-reactivity may reduce the costs of diagnosis. Nevertheless, further studies using a greater serological panel, including samples from other countries or regions, are essential to evaluating the maintenance of the diagnostic performance of rTC.

6. Patents

One patent resulting from the work reported in this manuscript is under protection at Instituto Nacional de Propriedade Industrial (INPI): Brazilian patent No. BR1020220183139.

Author Contributions: JMM, IAGP, IBG, AJR, JOS, MS, AAMG, MCP, ATC, EAFC and ASG wrote the manuscript. JMM, IAGP and ATC prepared the figures. JMM, IAGP, ACGM, MFCF, LMN, IBG, AJR, KAS, CAAR, JOS, MS, GSVT and MCP carried out experiments. JMM, IAGP, GSVT, MACF, ATC, EAFC and ASG revised and edited the manuscript. JMM, RCG, GSVT, MOCR, ATC, EAFC and ASG conceptualized the rational of experiments and supervised the students. JMM and IAGP contributed equally to this work.

Funding: This work was supported by grants from CNPq (APQ-408675/2018-7, APQ 428962/2018-1 and 303507/2021-7).

Institutional Review Board Statement: The study was conducted under the Declaration of Helsinki and approved by the Human Research Ethics Committee of UFMG, with protocol number CAAE-32343114.9.0000.5149 / CAAE-48354315.8.0000.5149.

Informed Consent Statement: "Informed consent was obtained from all subjects involved in the study."

Data Availability Statement: "Not applicable" here.

Acknowledgments: The authors would like to thank CAPES (Finance Code 001), CNPq, and FAPEMIG for the scholarships. ASG, RCG, MOCR, and EAFC would like to thank CNPq for their research fellowship. The authors are grateful to UFSJ, UFMG, CAPES, CNPq and FAPEMIG.

Conflicts of Interest: "The authors declare no conflict of interest."

References

1. de Fuentes-Vicente, J.A.; Vidal-López, D.G.; Flores-Villegas, A.L.; Moreno-Rodríguez, A.; de Alba-Alvarado, M.C.; Salazar-Schettino, P.M.; Rodríguez-López, M.H.; Gutiérrez-Cabrera, A.E. Trypanosoma Cruzi: A Review of Biological and Methodological Factors in Mexican Strains. *Acta Trop* 2019, 195, 51–57, doi:10.1016/J.ACTATROPICA.2019.04.024.
2. Chagas, C. Nova Tripanozomiae Humana: Estudos Sobre a Morfolojia e o Ciclo Evolutivo Do Schizotrypanum Cruzi n. Gen., n. Sp., Ajente Etiolojico de Nova Entidade Morbida Do Homem. *Mem Inst Oswaldo Cruz* 1909, 1, 159–218, doi:10.1590/S0074-02761909000200008.
3. Nunes, M.C.P.; Dones, W.; Morillo, C.A.; Encina, J.J.; Ribeiro, A.L. Chagas Disease: An Overview of Clinical and Epidemiological Aspects. *J Am Coll Cardiol* 2013, 62, 767–776, doi:10.1016/J.JACC.2013.05.046.
4. Acosta-Herrera, M.; Strauss, M.; Casares-Marfil, D.; Martín, J. Genomic Medicine in Chagas Disease. *Acta Trop* 2019, 197, 105062, doi:10.1016/J.ACTATROPICA.2019.105062.
5. PAHO Chagas Disease Available online: <https://www.paho.org/en/topics/chagas-disease> (accessed on 12 October 2022).
6. Antinori, S.; Galimberti, L.; Bianco, R.; Grande, R.; Galli, M.; Corbellino, M. Chagas Disease in Europe: A Review for the Internist in the Globalized World. *Eur J Intern Med* 2017, 43, 6–15, doi:10.1016/J.EJIM.2017.05.001.
7. Santos, F.A.C. dos; Braga, A.P.C.; Ferreira, I.S.; Arêde, M. da P.S.P.; Baetas, A.L.F.D.; Silva, M.V.A.; Lima, L.S.A. Avaliação Da Prevalência e Do Perfil Epidemiológico Da Doença de Chagas Aguda Entre 2014 e 2017 No Estado Do Pará, Brasil. *Brazilian Journal of Health Review* 2020, 3, 8974–8982, doi:10.34119/bjhrv3n4-142.
8. Echavarriá, N.G.; Echeverría, L.E.; Stewart, M.; Gallego, C.; Saldarriaga, C. Chagas Disease: Chronic Chagas Cardiomyopathy. *Curr Probl Cardiol* 2021, 46, 100507, doi:10.1016/J.CPCARDIOL.2019.100507.
9. Bern, C. Chagas' Disease. <https://doi.org/10.1056/NEJMra1410150> 2015, 373, 456–466, doi:10.1056/NEJMRA1410150.
10. Hughes, J.M.; Wilson, M.E.; Shikanai-Yasuda, M.A.; Carvalho, N.B. Oral Transmission of Chagas Disease. *Clinical Infectious Diseases* 2012, 54, 845–852, doi:10.1093/CID/CIR956.
11. Bern, C.; Messenger, L.A.; Whitman, J.D.; Maguire, J.H. Chagas Disease in the United States: A Public Health Approach. *Clin Microbiol Rev* 2020, 33, doi:10.1128/CMR.00023-19.
12. Tanowitz, H.B.; Kirchhoff, L. v.; Simon, D.; Morris, S.A.; Weiss, L.M.; Wittner, M. Chagas' Disease. *Clin Microbiol Rev* 1992, 5, 400, doi:10.1128/CMR.5.4.400.
13. Candia-Puma, M.A.; Machaca-Luque, L.Y.; Roque-Pumahuanca, B.M.; Galdino, A.S.; Giunchetti, R.C.; Coelho, E.A.F.; Chávez-Fumagalli, M.A. Accuracy of Diagnostic Tests for the Detection of Chagas Disease: A Systematic Review and Meta-Analysis. *Diagnostics* 2022, 12, 2752, doi:10.3390/DIAGNOSTICS12112752/S1.
14. Machado, F.S.; Dutra, W.O.; Esper, L.; Gollob, K.J.; Teixeira, M.M.; Factor, S.M.; Weiss, L.M.; Nagajyothi, F.; Tanowitz, H.B.; Garg, N.J. Current Understanding of Immunity to Trypanosoma Cruzi Infection and Pathogenesis of Chagas Disease. *Seminars in Immunopathology* 2012 34:6 2012, 34, 753–770, doi:10.1007/S00281-012-0351-7.
15. Abras, A.; Ballart, C.; Fernández-Arévalo, A.; Llovet, T.; Gállego, M.; Muñoz, C. ARCHITECT Chagas® as a Single Test Candidate for Chagas Disease Diagnosis: Evaluation of Two Algorithms Implemented in a Non-Endemic Setting. *Clinical Microbiology and Infection* 2021, 27, 782.e1–782.e6, doi:10.1016/J.CMI.2020.07.002.
16. Flores-Chávez, M.; Cruz, I.; Rodríguez, M.; Nieto, J.; Franco, E.; Gárate, T.; Cañavate, C. Comparación de Técnicas Serológicas Convencionales y No Convencionales Para El Diagnóstico de La Enfermedad de Chagas Importada En España. *Enferm Infect Microbiol Clin* 2010, 28, 284–293, doi:10.1016/j.eimc.2009.07.005.
17. Bern, C. Chagas' Disease. *New England Journal of Medicine* 2015, 373, 456–466, doi:10.1056/NEJMra1410150.
18. Alonso-Padilla, J.; Cortés-Serra, N.; Pinazo, M.J.; Bottazzi, M.E.; Abril, M.; Barreira, F.; Sosa-Estani, S.; Hotez, P.J.; Gascón, J. Strategies to Enhance Access to Diagnosis and Treatment for Chagas Disease Patients in Latin America. <https://doi.org/10.1080/14787210.2019.1577731> 2019, 17, 145–157, doi:10.1080/14787210.2019.1577731.

19. Alonso-Padilla, J.; Gallego, M.; Schijman, A.G.; Gascon, J. Molecular Diagnostics for Chagas Disease: Up to Date and Novel Methodologies. <https://doi.org.ez32.periodicos.capes.gov.br/10.1080/14737159.2017.1338566> 2017, 17, 699–710, doi:10.1080/14737159.2017.1338566.

20. Suárez, C.; Nolder, D.; García-Mingo, A.; Moore, D.A.J.; Chiodini, P.L. Diagnosis and Clinical Management of Chagas Disease: An Increasing Challenge in Non-Endemic Areas. *Res Rep Trop Med* 2022, 13, 25, doi:10.2147/RRTM.S278135.

21. Cisneros, J.S.; Chain, C.Y.; Daza Millone, M.A.; Labriola, C.A.; Scollo, K.; Ruiz, A.M.; Estrela, P.; Vela, M.E. Electrochemical Impedance Biosensor for Chagas Disease Diagnosis in Clinical Samples. *Biosens Bioelectron X* 2022, 12, 100261, doi:10.1016/J.BI-OSX.2022.100261.

22. Guzmán-Gómez, D.; López-Monteon, A.; de La Soledad Lagunes-Castro, M.; Álvarez-Martínez, C.; Hernández-Lutzon, M.J.; Dumonteil, E.; Ramos-Ligonio, A. Highly Discordant Serology against Trypanosoma Cruzi in Central Veracruz, Mexico: Role of the Antigen Used for Diagnostic. *Parasit Vectors* 2015, 8, 1–8, doi:10.1186/S13071-015-1072-2/FIGURES/2.

23. Pérez-Molina, J.A.; Molina, I. Chagas Disease. *The Lancet* 2018, 391, 82–94, doi:10.1016/S0140-6736(17)31612-4.

24. Ferreira-Silva, M.M.; Pereira, G. de A.; Rodrigues-Júnior, V.; Meira, W.S.; Basques, F. v; Langhi-Júnior, D.M.; Romanelli, M.; Umezawa, E.S.; Késpér-Júnior, N.; Louzada-Neto, F.; et al. Chagas Disease: Performance Analysis of Immunodiagnostic Tests Anti-Trypanosoma Cruzi in Blood Donors with Inconclusive Screening Results. *Hematol Transfus Cell Ther* 2021, 43, 410–416, doi:10.1016/j.hctc.2020.06.016.

25. Imai, K.; Murakami, T.; Misawa, K.; Fujikura, Y.; Kawana, A.; Tarumoto, N.; Maesaki, S.; Maeda, T. Optimization and Evaluation of the ARCHITECT Chagas Assay and In-House ELISA for Chagas Disease in Clinical Settings in Japan. *Parasitol Int* 2021, 80, 102221, doi:10.1016/J.PARINT.2020.102221.

26. Angheben, A. Air in the Tires: Towards an Achievable, Efficacious and Timely Diagnosis for Chagas Disease. *Mem Inst Oswaldo Cruz* 2022, 117, doi:10.1590/0074-02760210444CHGSA.

27. PAHO Guidelines for the Diagnosis and Treatment of Chagas Disease; 2019;

28. Echeverria, L.E.; Morillo, C.A. American Trypanosomiasis (Chagas Disease). *Infect Dis Clin North Am* 2019, 33, 119–134, doi:10.1016/j.idc.2018.10.015.

29. Chadalawada, S.; Sillau, S.; Archuleta, S.; Mundo, W.; Bandali, M.; Parra-Henao, G.; Rodriguez-Morales, A.J.; Villamil-Gomez, W.E.; Suárez, J.A.; Shapiro, L.; et al. Risk of Chronic Cardiomyopathy Among Patients With the Acute Phase or Indeterminate Form of Chagas Disease. *JAMA Netw Open* 2020, 3, e2015072, doi:10.1001/jamanetworkopen.2020.15072.

30. World Health Organization Control of Chagas Disease: Second Report of the WHO Expert Committee. WHO Technical Report Series 905; Geneva, 2002;

31. Rocha, M.O.; Teixeira, M.M.; Ribeiro, A.L. An Update on the Management of Chagas Cardiomyopathy. *Expert Rev Anti Infect Ther* 2007, 5, 727–743, doi:10.1586/14787210.5.4.727.

32. Longhi, S.A.; Brandariz, S.B.; Lafon, S.O.; Niborski, L.L.; Luquetti, A.O.; Schijman, A.G.; Levin, M.J.; Gómez, K.A. Evaluation of In-House ELISA Using Trypanosoma Cruzi Lysate and Recombinant Antigens for Diagnosis of Chagas Disease and Discrimination of Its Clinical Forms. *Am J Trop Med Hyg* 2012, 87, 267–271, doi:10.4269/ajtmh.2012.11-0533.

33. Abras, A.; Gállego, M.; Llovet, T.; Tebar, S.; Herrero, M.; Berenguer, P.; Ballart, C.; Martí, C.; Muñoz, C. Serological Diagnosis of Chronic Chagas Disease: Is It Time for a Change? *J Clin Microbiol* 2016, 54, 1566–1572, doi:10.1128/JCM.00142-16.

34. Balouz, V.; Agüero, F.; Buscaglia, C.A. Chagas Disease Diagnostic Applications. In; 2017; pp. 1–45.

35. Candia-Puma, M.A.; Machaca-Luque, L.Y.; Roque-Pumahuanca, B.M.; Galdino, A.S.; Giunchetti, R.C.; Coelho, E.A.F.; Chávez-Fumagalli, M.A. Accuracy of Diagnostic Tests for the Detection of Chagas Disease: A Systematic Review and Meta-Analysis. *Diagnostics* 2022, 12, 2752, doi:10.3390/diagnostics12112752.

36. Rocha, M.O.C.; Ribeiro, A.L.P.; Teixeira, M.M. Clinical Management of Chronic Chagas Cardiomyopathy. *Frontiers in Bioscience* 2003, 8, 926, doi:10.2741/926.

37. Whitman, J.D.; Bulman, C.A.; Gunderson, E.L.; Irish, A.M.; Townsend, R.L.; Stramer, S.L.; Sakanari, J.A.; Bern, C. Chagas Disease Serological Test Performance in U.S. Blood Donor Specimens. *J Clin Microbiol* 2019, 57, doi:10.1128/JCM.01217-19.

38. Kelly, E.A.; Bulman, C.A.; Gunderson, E.L.; Irish, A.M.; Townsend, R.L.; Sakanari, J.A.; Stramer, S.L.; Bern, C.; Whitman, J.D. Comparative Performance of Latest-Generation and FDA-Cleared Serology Tests for the Diagnosis of Chagas Disease. *J Clin Microbiol* 2021, 59, doi:10.1128/JCM.00158-21.

39. Truyens, C.; Dumonteil, E.; Alger, J.; Cafferata, M.L.; Ciganda, A.; Gibbons, L.; Herrera, C.; Sosa-Estani, S.; Buekens, P. Geographic Variations in Test Reactivity for the Serological Diagnosis of Trypanosoma Cruzi Infection. *J Clin Microbiol* 2021, 59, doi:10.1128/JCM.01062-21.

40. Duthie, M.S.; Guderian, J.A.; Vallur, A.C.; Misquith, A.; Liang, H.; Mohamath, R.; Luquetti, A.O.; Carter, D.; Tavares, S.N.B.; Reed, S.G. Multi-Epitope Proteins for Improved Serological Detection of Trypanosoma Cruzi Infection and Chagas Disease. *Diagn Microbiol Infect Dis* 2016, 84, 191–196, doi:10.1016/j.diagnmicrobio.2015.11.006.

41. Freitas, N.E.M.; Santos, E.F.; Leony, L.M.; Silva, Á.A.O.; Daltro, R.T.; Vasconcelos, L. de C.M.; Duarte, G.A.; Oliveira da Mota, C.; Silva, E.D.; Celedon, P.A.F.; et al. Double-Antigen Sandwich ELISA Based on Chimeric Antigens for Detection of Antibodies to Trypanosoma Cruzi in Human Sera. *PLoS Negl Trop Dis* 2022, 16, e0010290, doi:10.1371/journal.pntd.0010290.

42. Dipti, C.A.; Jain, S.K.; Navin, K. A Novel Recombinant Multiepitope Protein as a Hepatitis C Diagnostic Intermediate of High Sensitivity and Specificity. *Protein Expr Purif* 2006, 47, 319–328, doi:10.1016/j.pep.2005.12.012.

43. Ribeiro, P.A.F.; Souza, M.Q.; Dias, D.S.; Álvares, A.C.M.; Nogueira, L.M.; Machado, J.M.; dos Santos, J.C.; Godoi, R.R.; Nobrega, Y.K.M.; Campos-da-Paz, M.; et al. A Custom-Designed Recombinant Multiepitope Protein for Human Cytomegalovirus Diagnosis. *Recent Pat Biotechnol* 2019, 13, 316–328, doi:10.2174/1872208313666190716093911.

44. de Souza, M.Q.; Galdino, A.S.; dos Santos, J.C.; Soares, M.V.; Nóbrega, Y.C. de; Álvares, A. da C.M.; de Freitas, S.M.; Torres, F.A.G.; Felipe, M.S.S. A Recombinant Multiepitope Protein for Hepatitis B Diagnosis. *Biomed Res Int* 2013, 2013, 1–7, doi:10.1155/2013/148317.

45. Galdino, A.S.; Santos, J.C.; Souza, M.Q.; Nóbrega, Y.K.M.; Xavier, M.-A.E.; Felipe, M.S.S.; Freitas, S.M.; Torres, F.A.G. A Novel Structurally Stable Multiepitope Protein for Detection of HCV. *Hepat Res Treat* 2016, 2016, 6592143, doi:10.1155/2016/6592143.

46. Anandarao, R.; Swaminathan, S.; Fernando, S.; Jana, A.M.; Khanna, N. Recombinant Multiepitope Protein for Early Detection of Dengue Infections Recombinant Multiepitope Protein for Early Detection of Dengue Infections. 2006, 13, 59–67, doi:10.1128/CVI.13.1.59.

47. Galdino, A.S.; Santos, J.C.; Souza, M.Q.; Nóbrega, Y.K.M.; Xavier, M.-A.E.; Felipe, M.S.S.; Freitas, S.M.; Torres, F.A.G. A Novel Structurally Stable Multiepitope Protein for Detection of HCV. *Hepat Res Treat* 2016, 2016, 1–9, doi:10.1155/2016/6592143.

48. Taherkhani, R.; Farshadpour, F.; Makvandi, M. Design and Production of a Multiepitope Construct Derived from Hepatitis E Virus Capsid Protein. *J Med Virol* 2015, 87, 1225–1234, doi:10.1002/jmv.24171.

49. Thomasini, R.L.; Souza, H.G.A.; Bruna-Romero, O.; Totola, A.H.; Gonçales, N.S.L.; Lima, C.X.; Teixeira, M.M. Evaluation of a Recombinant Multiepitope Antigen for Diagnosis of Hepatitis C Virus: A Lower Cost Alternative for Antigen Production. *J Clin Lab Anal* 2018, 32, e22410, doi:10.1002/jcla.22410.

50. Yengo, B.N.; Shintou, C.M.; Hotterbeekx, A.; Yaah, N.E.; Shey, R.A.; Quanico, J.; Baggerman, G.; Ayong, L.; Vanhamme, L.; Njemini, R.; et al. Immunoinformatics Design and Assessment of a Multiepitope Antigen (OvMCBL02) for Onchocerciasis Diagnosis and Monitoring. *Diagnostics* 2022, 12, 1440, doi:10.3390/diagnostics12061440.

51. Faria, A.R.; de Castro Veloso, L.; Coura-Vital, W.; Reis, A.B.; Damasceno, L.M.; Gazzinelli, R.T.; Andrade, H.M. Novel Recombinant Multiepitope Proteins for the Diagnosis of Asymptomatic *Leishmania infantum*-Infected Dogs. *PLoS Negl Trop Dis* 2015, 9, e3429, doi:10.1371/journal.pntd.0003429.

52. Jameie, F.; Dalimi, A.; Pirestani, M.; Mohebali, M. Detection of *Leishmania infantum* Infection in Reservoir Dogs Using a Multiepitope Recombinant Protein (PQ10). *Arch Razi Inst* 2020, 75, 327–338, doi:10.22092/ari.2019.126524.1346.

53. Lica, A.; Faria, R.; de Castro Veloso, L.; Coura-Vital, W.; Reis, A.B.; Damasceno, L.M.; Gazzinelli, R.T.; Lida, H.; Andrade, M. Novel Recombinant Multiepitope Proteins for the Diagnosis of Asymptomatic *Leishmania infantum*-Infected Dogs. *PLoS Negl Trop Dis* 2015, 9, doi:10.1371/journal.pntd.0003429.

54. Houghton, R.L.; Benson, D.R.; Reynolds, L.; McNeill, P.; Sleath, P.; Lodes, M.; Skeiky, Y.A.W.; Badaro, R.; Krettli, A.U.; Reed, S.G. Multiepitope Synthetic Peptide and Recombinant Protein for the Detection of Antibodies to *Trypanosoma cruzi* in Patients with Treated or Untreated Chagas' Disease. *J Infect Dis* 2000, 181, 325–330, doi:10.1086/315165.

55. Camussone, C.; Gonzalez, V.; Belluzo, M.S.; Pujato, N.; Ribone, M.E.; Lagier, C.M.; Marcipar, I.S. Comparison of Recombinant *Trypanosoma cruzi* Peptide Mixtures versus Multiepitope Chimeric Proteins as Sensitizing Antigens for Immunodiagnosis. *Clinical and Vaccine Immunology* 2009, 16, 899–905, doi:10.1128/CVI.00005-09.

56. Peverengo, L.M.; Garcia, V.; Rodeles, L.M.; Mendicino, D.; Vicco, M.; Lagier, C.; Gonzalez, V.; Gugliotta, L.; Marcipar, I. Development and Assessment of an Improved Recombinant Multiepitope Antigen-Based Immunoassay to Diagnose Chronic Chagas Disease. *Parasitology* 2018, 145, 1594–1599, doi:10.1017/S0031182018000458.

57. Duthie, M.S.; Guderian, J.A.; Vallur, A.C.; Misquith, A.; Liang, H.; Mohamath, R.; Luquetti, A.O.; Carter, D.; Tavares, S.N.B.; Reed, S.G. Multi-Epitope Proteins for Improved Serological Detection of *Trypanosoma cruzi* Infection and Chagas Disease. *Diagn Microbiol Infect Dis* 2016, 84, 191–196, doi:10.1016/j.diagmicrobio.2015.11.006.

58. Jackson, Y.; Chatelain, E.; Mauris, A.; Holst, M.; Miao, Q.; Chappuis, F.; Nda, M. Serological and Parasitological Response in Chronic Chagas Patients 3 Years after Nifurtimox Treatment. *BMC Infect Dis* 2013, 13, 85, doi:10.1186/1471-2334-13-85.

59. Bartsch, S.M.; Avelis, C.M.; Asti, L.; Hertenstein, D.L.; Ndeffo-Mbah, M.; Galvani, A.; Lee, B.Y. The Economic Value of Identifying and Treating Chagas Disease Patients Earlier and the Impact on *Trypanosoma cruzi* Transmission. *PLoS Negl Trop Dis* 2018, 12, e0006809, doi:10.1371/journal.pntd.0006809.

60. Longhi, D.A.; Dalcanton, F.; Aragão, G.M.F. de; Carciofi, B.A.M.; Laurindo, J.B. Assessing the Prediction Ability of Different Mathematical Models for the Growth of *Lactobacillus plantarum* under Non-Isothermal Conditions. *J Theor Biol* 2013, 335, 88–96, doi:10.1016/j.jtbi.2013.06.030.

61. Daltro, R.T.; Leony, L.M.; Freitas, N.E.M.; Silva, Â.A.O.; Santos, E.F.; Del-Rei, R.P.; Brito, M.E.F.; Brandão-Filho, S.P.; Gomes, Y.M.; Silva, M.S.; et al. Cross-Reactivity Using Chimeric *Trypanosoma cruzi* Antigens: Diagnostic Performance in Settings Where Chagas Disease and American Cutaneous or Visceral Leishmaniasis Are Coendemic. *J Clin Microbiol* 2019, 57, doi:10.1128/JCM.00762-19.