

Low Prevalence of SARS-CoV-2 Antibodies in Canine and Feline Serum Samples Collected during the Covid-19 Pandemic in Hong Kong and Korea

[Yun Young Go](#) , [Maura Carrai](#) , [Yan Ru Choi](#) , Christopher J. Brackman , Karina W.S. Tam , Pierra Y.T. Law , Fiona Woodhouse , Jane Gray , Ji Hun Kim , Joohyung Park , Chae Won Jeon , Hyomi Jang , Ioannis Magouras , [Nicola Decaro](#) , Samuel MS Cheng , Malik Peiris , [Julia A. Beatty](#) , [Vanessa R. Barrs](#) *

Posted Date: 13 January 2023

doi: 10.20944/preprints202301.0250.v1

Keywords: SARS-CoV-2; COVID-19; Feline; Canine; seroprevalence; zoonosis



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Low Prevalence of SARS-CoV-2 Antibodies in Canine and Feline Serum Samples Collected during the COVID-19 Pandemic in Hong Kong and Korea

Yun Young Go ^{1,†}, Maura Carrai ^{2,†}, Yan Ru Choi ^{2,†}, Christopher J. Brackman ³, Karina W.S. Tam ³, Pierra Y.T. Law ³, Fiona Woodhouse ⁴, Jane Gray ⁴, Ji Hun Kim ⁵, Joohyung Park ⁶, Chae Won Jeon ⁷, Hyomi Jang ⁷, Ioannis Magouras ¹, Nicola Decaro ⁸, Samuel MS Cheng ⁹, Malik Peiris ⁹, Julia A. Beatty ² and Vanessa R. Barrs ^{2,*}

¹ Department of Infectious Diseases and Public Health, Jockey Club College of Veterinary Medicine and Life Sciences, City University of Hong Kong, Hong Kong SAR, China

² Department of Veterinary Clinical Sciences, Jockey Club College of Veterinary Medicine and Life Sciences; and Centre for Animal Health and Welfare, City University of Hong Kong, Hong Kong SAR, China

³ Agriculture, Fisheries and Conservation Department, Government of the Hong Kong Special Administrative Region, Hong Kong SAR, China

⁴ Society for the Prevention of Cruelty to Animals (Hong Kong), 5 Wan Shing St, Wan Chai, Hong Kong

⁵ 24 h Jamsil On Animal Medical Center, Seoul, Republic of Korea

⁶ VIP Animal Medical Center, Seoul, Republic of Korea

⁷ Neodin Biovet Laboratory, Seoul, Republic of Korea

⁸ Department of Veterinary Medicine, University of Bari Aldo Moro, 70010 Valenzano, Italy

⁹ School of Public Health, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong

* Correspondence: vanessa.barrs@cityu.edu.hk

† These authors contributed equally to the manuscript.

Abstract: Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has affected millions of people worldwide since its emergence in 2019. The current global pandemic was driven by human-to-human transmission. Knowing the zoonotic origin of the disease and the potential capacity of the virus to adapt to other species, it is important to understand the extent of natural SARS-CoV-2 infection of animals, in particular cats and dogs in households that are in direct contact with their owners. Hong Kong and Seoul are two of the most densely-populated urban cities in Asia, where companion animals often live in close contact with humans. In this study, we screened sera from 1,040 cats and 855 dogs during the early phase of the pandemic in Hong Kong and Seoul for SARS-CoV-2 antibodies by using an ELISA that detects antibodies against the receptor binding domain (RBD) of the viral spike protein. Sera testing positive on ELISA were also tested for the presence of neutralizing antibodies using a surrogate virus neutralization (sVNT) and plaque reduction neutralization test (PRNT). Among feline sera, 4.51% and 2.54% of samples from Korea and Hong Kong, respectively, tested ELISA positive. However only 1.64% of samples from Korea and 0.18% from Hong Kong tested positive by sVNT, while only 0.41% of samples from Korea tested positive by PRNT. Among canine samples, 4.94% and 6.46% from Korea and Hong Kong, respectively, tested positive by ELISA, while only 0.29% sera from Korea were positive on sVNT and no canine sera tested positive by PRNT. These results confirm a low seroprevalence of SARS-CoV-2 exposure in companion animals in Korea and Hong Kong. The discordance between RBD-ELISA and neutralization tests may indicate possible ELISA cross-reactivity with other coronaviruses, especially in canine sera.

Keywords: SARS-CoV-2; COVID-19; Feline; Canine; seroprevalence; zoonosis

1. Introduction

Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), the causative agent of the ongoing coronavirus disease (COVID-19) pandemic, belongs to the genus of *Betacoronaviruses* in the family *Coronaviridae*. Coronaviruses can infect a wide range of birds and mammals, and manifest high genetic diversity due to mutations and recombination of their genomes [1-3]. Several members of the

family *Coronaviridae* circulate continuously in human and animal populations and mostly cause mild symptoms in humans, such as the common cold or mild enteric disease [4-7]. In contrast, the SARS-CoV and the Middle East respiratory syndrome-associated coronavirus (MERS-CoV) are highly pathogenic in humans and cause severe lower respiratory tract infections that can further develop into acute respiratory distress syndrome (ARDS) and extrapulmonary manifestations [4, 8-11]. SARS-CoV and MERS-CoV which originated in bats and infected humans via intermediate hosts are well-known for cross-species transmission. Similarly, it is believed that the novel SARS-CoV-2 that emerged in late 2019 in Wuhan, China, has a zoonotic origin from bats, while the intermediate host has not yet been identified [12-14].

While the current global pandemic is driven by human-to-human transmission, concern has arisen regarding disease transmission between humans and susceptible animals as evidenced by sporadic cases of SARS-CoV-2 infection in companion, farmed, wild and captive zoological animals that have been in close contact with infected humans [15-20]. Knowing the zoonotic origin of the disease and the potential capacity of the virus to adapt to other species, it is important to understand the extent of natural SARS-CoV-2 infection of animals, in particular cats and dogs in households that are in direct contact with their owners and other household members. To date, natural SARS-CoV-2 infection has been sporadically detected in companion animals where human-to-animal transmission was suspected in almost all cases [16, 21-24]. In line with this, serological investigations in cats and dogs have been conducted in different countries since the early phases of the COVID-19 pandemic. Molecular and antibody prevalence studies revealed that dogs and cats living in areas of high human infection had been occasionally infected by SARS-CoV-2, thus confirming previous reports of natural infections of companion animals [25-27]. Specifically, the seroprevalence of SARS-CoV-2 in cats with an unknown history of exposure to the virus was 0.69% (6/920) in Germany [28], 0.76% (1/131) in Croatia [29], 0.4% (2/500) in the Netherlands [30], and 3.3% (15/451) in Italy [31]. In dogs, the prevalence of neutralizing antibodies was as low as 0.31% (2/654) in Croatia [29], 0.2% (1/500) in the Netherlands [30], and 3.3% (15/451) in Italy [31]. Not surprisingly, the seroprevalence of animals from COVID-19-affected households was higher, ranging from 10% to 12.8% in dogs and 4.5% to 43.8% in cats [25-27, 31].

Hong Kong and Seoul are two of the most densely-populated urban cities in Asia, where companion animals often live in close contact with humans. It is estimated that 241,900 (9.4%) households in Hong Kong keep dogs and/or cats [32], whereas 1 in 3 South Korean households have companion animals, with the number of pet dogs and cats reaching 8.6 million (6.02 million dogs and 2.58 million cats) [33]. In this study, we used an ELISA that detects the SARS-CoV-2 receptor binding domain (RBD) to assess the prevalence of anti-SARS-CoV-2 antibodies in 1,040 cats and 855 dogs with an unknown history of exposure to the disease during the early phase of the pandemic in Hong Kong and Seoul. Positive results were confirmed using a surrogate virus neutralization (sVNT) and plaque reduction neutralization test (PRNT). In addition, we investigated the prevalence of SARS-CoV-2 RNA shedding in oronasal swabs from a subset of 301 healthy free-roaming cats presented to a trap-neuter release desexing clinic in Hong Kong using RT-PCR.

2. Materials and Methods

2.1. Sample description

Seoul, Korea

Residual diagnostic sera from 344 dogs and 488 cats collected from April to July of 2020 were obtained from three veterinary practices in different districts of Seoul, Republic of Korea. All animals were sampled by veterinarians during routine healthcare visits for various reasons. For each sample, the following information was available: species, gender, age at sampling, and date of sampling.

Hong Kong, China

Residual diagnostic sera were collected from 511 dogs (from January 2021 until August 2021) and from 251 cats (from March 2020 to April 2021) presenting for routine healthcare visits at a large veterinary primary care and referral hospital in Sham Shui Po, Hong Kong. In addition, serum

samples, nasal swabs and oropharyngeal were collected from 301 healthy free-roaming colony cats desexed in a trap-neuter release (TNR) program at a large animal shelter in Wan Chai, Hong Kong, from January 2021 to May 2021.

Positive control sera were obtained from a dog and a cat admitted to the Agriculture, Fisheries, and Conservation Department (AFCD) quarantine facility because their owners were diagnosed with COVID-19. These two animals had been confirmed to be shedding SARS-CoV-2 in respiratory and fecal secretions and tested positive for serum-neutralizing antibodies using plaque reduction neutralization 90 (PRNT₉₀) test, with titers of 1:160 for the dog and $\geq 1:320$ for the cat [16, 34].

For use as negative controls and ELISA optical density (OD) cut-off determination, 46 feline sera and 23 canine sera were collected before COVID-19 outbreaks, before January 2020, that had been stored at -80°C were used. Animal ethics approval for this study was granted by the Animal Ethics Committee of the City University of Hong Kong, approval no. A-0478. Research licenses for sampling colony cats were granted by The Government of the Hong Kong SAR, Department of Health (20-164 to 20-179).

2.2. Assay description and optimisation

All serum samples were heat-inactivated at 56°C for 30 min before use. The sera were tested using an ELISA to detect anti-SARS-CoV-2 spike glycoprotein receptor binding domain (RBD) antibodies. The protocol was based on a previously described assay with modifications [35]. Briefly, 96-well plates (Nunc MaxiSorp, Thermo Fisher Scientific, cat. no. 44-2404-21) were coated with 100 ng per well of SARS-CoV-2 Spike RBD-mFc Recombinant Protein (Sino Biological, Beijing, China, Sinobiological, cat. no. 40592-V05H) diluted in PBS buffer and refrigerated overnight at 4°C. The next day, coated plates were blocked using 100 μ l of Chonblock blocking/sample dilution ELISA buffer (Chondrex Inc, Redmon, US, cat. no. 9068) for 1 h at room temperature. Heat-inactivated serum samples were diluted 1:100 in Chonblock blocking/sample dilution ELISA buffer, with a 15 min incubation at room temperature. After 3 rounds of washing the plate with PBS containing 0.2 % Tween 20, the diluted sera samples were transferred to the coated plate and incubated in a humidified chamber at 37°C for 2 h. Washing was performed before adding horseradish peroxidase (HRP)-conjugated goat anti-feline or canine IgG antibodies (Thermo Fisher Scientific, cat. no PA1-84673 or A18763 respectively) diluted 1:2000 in PBS. The plates were incubated in a humidified chamber for 1 h at 37°C, then washed. Subsequently, 100 μ l of 3,3',5,5' tetramethylbenzidine (TMB) substrate (Ncm TMB One; New Cell and Molecular Biotech Co. Ltd, Suzhou, China, cat. no. M30500) was added into each well. After a 6 min (feline) or 5 min incubation (canine), the reaction was stopped by adding 50 μ l of 2 M H₂SO₄ solution and analyzed on an absorbance microplate reader at 450 nm wavelength (SpectraMax ABS, Molecular Devices). Test samples were run in duplicate; each plate contained a duplicate positive control, negative control, and blank control (PBS only). For cut-off determination, the mean of the OD₄₅₀ of negative sera collected pre-COVID-19 plus 3 \times and 6 \times the SD of the negative control readings for cats and dogs were used, respectively.

2.3. Confirmatory Serological Tests: Surrogate Virus Neutralization and Plaque Reduction Neutralization tests

Canine and feline sera that tested positive on ELISA were evaluated for SARS-CoV-2 specific-neutralizing antibodies using a commercially available (sVNT; GeneScript Inc. Piscataway, USA) according to the manufacturer's instructions. The OD values were read on a spectrophotometer (Molecular Devices SpectraMax ABS, San Jose, CA USA) at 450 nm. Positive and negative controls provided by the kit were included in duplicate in every run. The quality control and validation of the results were based on the OD values for positive and negative controls falling in the recommended values. The results were interpreted as positive for SARS-CoV-2 neutralizing activity when the sVNT inhibition was calculated to be >20%, while <20% was regarded as a negative result.

Subsequently, canine and feline serum samples with positive results in sVNT were further tested with a plaque reduction neutralization test (PRNT) in a BSL-3 laboratory at the University of Hong Kong, as previously described [34].

2.4. Detection of SARS-CoV-2 RNA shedding in oronasal secretions of healthy-free roaming cats in Hong Kong by RT-PCR

To investigate the prevalence of active shedding of SARS-CoV-2 RNA in oronasal secretions collected from the 301 healthy free-roaming cats in Hong Kong, RT-PCR was performed to detect the *E* gene and *RdRp* genes as previously described [16].

2.5. Statistical Analyses

Data analysis was performed using GraphPad Prism software v.9.3.1 (GraphPad Software, LLC, CA, USA). The coefficient of variation (CV) was calculated as the ratio of the standard deviation to the mean multiplied by 100. Intra-assay CV $\leq 10\%$ and inter-assay CV $\leq 15\%$ were considered acceptable [36]. A Pearson correlation coefficient was calculated for ELISA-positive serum samples that were further analyzed by sVNT.

3. Results

3.1. Intra- and inter-assay coefficients of variation (CV)

The in-house SARS-CoV-2 RBD ELISA showed intra-assay CV values within acceptable limits, with CV $< 10\%$ for the canine and feline sera. The inter-assay CV of the canine ELISA was 5.66 % for the positive control serum and 10.94 % for the negative control serum. The inter-assay CV of the feline ELISA was 9.93 % for the positive control serum and 13.28 % for the negative control serum.

3.2. SARS-CoV-2 seroprevalence in domestic cats

The positive control cat serum (PRNT₉₀ $\geq 1:320$) showed a mean OD₄₅₀ value of 1.918 by RBD-ELISA. The mean OD₄₅₀ value of 46 pre-COVID-19 feline sera was 0.196 ± 0.123 , resulting in a cut-off OD₄₅₀ of 0.564 for positive samples. Of the 488 feline serum samples from Korea, 22 samples (4.51%) were deemed to be positive for anti-S RBD antibodies by ELISA with OD₄₅₀ values ranging from 0.606 to 2.336, while of the 552 feline sera collected in Hong Kong, 14 samples (2.54 %) were positive (OD₄₅₀ range 0.580 to 1.388). Specifically, 13 out of 251 sera (5.18%) and only one out of 301 sera (0.33%) sampled in veterinary primary care hospitals and animal shelters in Hong Kong, respectively, tested positive. In addition, nasal and oropharyngeal swabs collected from 301 cats of the animal shelter tested negative for the presence of SARS-CoV-2 RNA by RT-PCR. The 36 feline sera that tested ELISA-positive were evaluated using the sVNT. Nine samples, including eight from Korea (1.64 %) and one from Hong Kong (0.18%), tested positive, with an inhibition ranging from 30.46 to 38.75% (Figure 1). One feline serum sample obtained from Hong Kong tested ELISA negative (OD₄₅₀ 0.512) but was positive by sVNT with an 35.71% inhibition. The Pearson correlation coefficient analysis confirmed the lack of correlation between RBD ELISA and sVNT.




Figure 1. Comparison of serological results of feline samples tested by ELISA detecting antibodies to the receptor binding domain (RBD) of the SARS-CoV2 spike protein, with virus neutralization (sVNT). The dotted lines show the positive cut-off levels. Cat serum samples ($n = 37$) are indicated in circles, whereas semi-closed circle indicates ELISA-negative and sVNT-positive ($n=1$) and closed circles indicate positive samples by both assays ($n=8$). Pearson correlation coefficient was calculated to determine the correlation between the reactivities of RBD ELISA vs. sVNT.

Subsequently, the nine positive samples by sVNT were further evaluated for the presence of neutralizing antibodies by PRNT₉₀ and PRNT₅₀, of which two from Korea (H-C262 and H-C266) were positive. Specifically, H-C262 had a mean OD₄₅₀ of 2.336 and 32.35% inhibition, whereas H-C266 showed a mean OD₄₅₀ of 0.7659 and 30.10% inhibition by RBD-ELISA and sVNT, respectively. One sample (ID 155961) obtained from a veterinary primary care hospital in Hong Kong with an OD₄₅₀ value of 0.5121 tested positive by sVNT with 35.71 % inhibition. However, this sample was determined negative by PRNT₉₀. Both PRNT₉₀ and PRNT₉₀ gave comparable results. A summary of the SARS-CoV-2-specific antibodies detected from feline sera by three serological assays is shown in Table 1.

Table 1. Seropositivity of SARS-CoV-2 detected with RBD-ELISA, sVNT, and PRNT₉₀ in feline sera from Hong Kong and Korea.

Species	Total Sera	Origin	Collection Date	No. of sera	No. of ELISA positive (%)	No. of sVNT positive (%)	No. of PRNT positive (%)
Cats	1040	Korea	04/2020 to 07/2020	488	22 (4.51 %)	8 (1.64 %)	2 (0.41 %)
		Hong Kong		552	14 (2.54 %)	1 (0.18 %)	0 (0 %)
		- Veterinary clinic	03/2020 to 04/2021	251	13	1	0
		- Animal shelter	01/2021 to 05/2021	301	1	0	0

3.3. SARS-CoV-2 seroprevalence in domestic dogs

The positive control dog serum (PRNT₉₀ 1:160) showed a mean OD₄₅₀ value of 1.762 by RBD-ELISA. The mean OD value of the negative control pre-COVID sera was 0.223 ± 0.065 , yielding a cut-off of 0.610 for positive samples. Of the 344 sera from Korea, 17 (4.94 %) were positive with OD values

ranging from 0.632 to 2.479, while of the 511 sera collected in Hong Kong, 33 (6.46 %) were positive with an OD range of 0.611 to 2.135. When the positive sera by RBD-ELISA were tested with sVNT, only one sample from Korea (H-D303; OD₄₅₀ 0.8879) tested positive with a 23.92 % inhibition (Figure 2).




Figure 2. Comparison of serological results of canine samples tested by RBD ELISA and sVNT. The dotted lines show the positive cut-off levels. Dog serum samples (n = 36) are indicated in circles, whereas closed circles indicate positive samples by both assays.

A summary of the SARS-CoV-2-specific antibodies detected from canine sera by three serological assays is shown in Table 2.

Table 2. Seropositivity of SARS-CoV-2 detected with RBD-ELISA, sVNT, and PRNT₉₀ in canine sera from Hong Kong and Korea.

Species	Total Sera	Origin	Collection Date	No. of sera	No. of ELISA positive (%)	No. of sVNT positive (%)	No. of PRNT positive (%)
Dogs	855	Korea	04/20 – 05/20	344	17 (4.94%)	1 (0.29%)	0
		Hong Kong	01/21 - 08/21	511	33 (6.46 %)	0	0

3.4. Prevalence of SARS-CoV-2 RNA shedding in oronasal secretions of healthy-free roaming cats in Hong Kong-

SARS-CoV-2 RNA was not detected in any of the samples from the 301 healthy free-roaming cats in Hong Kong, using RT-PCR to detect the E gene and RdRp genes.

Discussion

In this study, we performed a large-scale serological investigation in cats and dogs during the emergence of the COVID-19 pandemic in Seoul and Hong Kong, regions geographically adjacent to Mainland China, from where SARS-CoV-2 originated. Our results showed that the seroprevalence of SARS-CoV-2 exposure in cats and dogs in Korea was 4.51% and 4.94%, respectively, whereas SARS-CoV-2 seropositive cats and dogs in Hong Kong were 2.54% and 6.46%, respectively, as determined by RBD-ELISA. For cats, our data indicated that the seroprevalence in Hong Kong was relatively lower than in Korea, which likely reflects the difference in human incidence of SARS-CoV-2 between the two regions during the sample collection period. In Korea, pet cat samples were collected from

three veterinary clinics located in different districts of Seoul from April 2020 to July 2020, where the human incidence rate was 1,150 cases in Seoul during this period (total cumulative human incidence of 5,707,477 cases in the city by the end of July 2020), according to the data released by the Ministry of Health and Welfare of South Korea (available on <https://ncov.kdca.go.kr/en/>). The pet cat sera from Hong Kong were collected over a more extended period (March 2020 to April 2021 inclusive), encompassing second to fourth COVID-19 waves with an average of only 28 human cases per day [37, 38]. Since both Hong Kong and South Korea used aggressive public health control measures to contain COVID-19 transmission, cumulative human infection rates were low during 2020-2021. Nevertheless, the seroprevalence among cats in both cities was comparable to other regions, including Germany (4.2%), the United Kingdom (3.3%), Italy (4.2%), and Spain (6.4%) during the first COVID-19 wave in Europe [39].

Similarly, the low seroprevalence (0.33%, 1/301) and absence of SARS-CoV-2 RNA shedding found in free-roaming colony cats in Hong Kong might be explained by the lower human incidence during the collection period, January 2021 to May 2021, which coincides with the end of the fourth wave in Hong Kong, but also by less human-cat interactions among these free-roaming cats. Interestingly, among 22 RBD-ELISA-positive feline sera from Korea, eight exhibited neutralizing activity by sVNT, while only one out of 14 RBD-ELISA-positive sera from Hong Kong tested positive on sVNT. Serum collection from the animals in this study was carried out during 2020-21, a period where antigenic diversity of circulating strains was close to the wild-type virus used for the serological tests reported here. Thus, antigenic diversity is unlikely to have led to false negative results.

Although virus neutralization assays are considered the gold standard, little is known regarding their sensitivity compared to RBD-ELISAs in identifying SARS-CoV-2 infections in dogs and cats. The discrepant results between these two assays could be explained by the detection of different types of anti-SARS-CoV-2 antibodies by the two assays. The SARS-CoV-2 sVNT detects antibodies with neutralizing activity against RBD of SARS-CoV-2 spike protein [40, 41]. Previously, monoclonal antibody studies demonstrated that the RBD-sVNT measures functional neutralizing antibodies, whereas the RBD-ELISA detects both binding and neutralizing antibodies [42]. In addition, it should be considered that not all neutralizing antibodies necessarily bind to the RBD, as reported by previous SARS-CoV studies that indicated antibodies raised against other regions in the S1 or S2 protein could also play a role in virus neutralization [43]. Furthermore, the difference observed in measuring neutralizing antibodies between sVNT and RBD ELISA also suggested that the level of RBD-binding IgG antibodies used alone is not reliable for evaluating the level of neutralizing capacity in companion animals. Dileepan et al. showed that the neutralizing antibodies in RBD-seropositive samples exhibited generally low titers and, in some instances, even lower than some of the RBD-seronegative samples, suggesting SARS-CoV-2 neutralizing activity in pet cats targets non-RBD regions of the S protein [44]. Nevertheless, possible cross-reaction with other feline as yet unidentified betacoronaviruses, as described in other studies [41, 45], cannot be ruled out, which warrants further investigation regarding the specificity of the RBD-ELISA.

Similar to cat samples, a modified RBD-ELISA used in human epidemiologic studies was established in the laboratory to detect SARS-CoV-2 RBD antibodies in dog sera. However, the positive cut-off for dogs could not be determined using the pre-COVID-19 cohort sample values plus a three-fold standard deviation (SD), as described for the cat samples. The pre-pandemic samples exhibited low reactivity, which led to using a six-fold SD value for the positive cut-off [40]. The low reactivity in the pre-COVID-19 samples in RBD-ELISA was also described previously [46]. Despite using a high positive cut-off, a high antibody-positive rate was observed in dog sera regardless of the origin, highlighting the possible cross-reactivity with other canine coronaviruses, probably canine respiratory coronavirus (CRCoV) [37, 47]. Unfortunately, the lack of negative control sera for CRCoV collected during the pandemic limited the capacity to determine the specificity of the RBD-ELISA. In this study, only one out of 17 RBD-ELISA-positive samples from dogs in Korea was confirmed positive by sVNT, indicating a low SARS-CoV-2 infection rate in dogs, consistent with previous publications.

The data presented in this study supports previous studies showing that dogs and cats can develop neutralizing antibodies against SARS-CoV-2 [30, 31, 40, 46]. Samples used in this study were collected from companion animals presented to veterinary clinics. However, except for samples collected from cats in an animal shelter in Hong Kong that were tested by RT-PCR, the status of SARS-CoV-2 shedding in pet animals was not determined, nor was the owners' health status in association with COVID-19 known at the time of sampling. Therefore, it was not possible to determine associations between antibody levels and possible exposure to SARS-CoV-2 from infected humans, nor any perceived association of COVID-19 disease in these patients.

In summary, our data confirmed that cats and dogs in close contact with people can develop an antibody response against SARS-CoV-2 infection, and similar to other studies, the seroprevalence of SARS-CoV-2 among cats and dogs belonging to owners of unknown COVID-19 status was low. [30, 31, 39, 40, 44, 46, 48-50]. This study also highlights the need to establish species-specific serological assays that can assess neutralizing antibodies in companion animals to provide accurate measurements of active infection in households and communities to ensure that all transmission opportunities are prevented.

Author Contributions: Approval for this study was granted by the Animal Ethics Committee of the City University of Hong Kong, approval no. A-0478. Research licenses for sampling colony cats were granted by The Government of the Hong Kong SAR, Department of Health (20-164 to 20-179).

Funding: This study was funded by a grant from the Health and Medical Research Fund (HMRP) of the Hong Kong Government Food and Health Bureau.

Conflict of Interest: The authors declare no conflict of interest.

References

- Ludwig, S. and A. Zarbock, *Coronaviruses and SARS-CoV-2: A Brief Overview*. Anesth Analg, 2020. **131**(1): p. 93-96.
- Voskarides, K., Animal-to-Human Viral Transitions: Is SARS-CoV-2 an Evolutionarily Successful One? J Mol Evol, 2020. **88**(5): p. 421-423.
- Wang, H., et al., *The genetic sequence, origin, and diagnosis of SARS-CoV-2*. Eur J Clin Microbiol Infect Dis, 2020. **39**(9): p. 1629-1635.
- Guan, Y., et al., Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. Science, 2003. **302**(5643): p. 276-8.
- Cavanagh, D., Nidovirales: a new order comprising Coronaviridae and Arteriviridae. Arch Virol, 1997. **142**(3): p. 629-33.
- Graham, R.L. and R.S. Baric, Recombination, reservoirs, and the modular spike: mechanisms of coronavirus cross-species transmission. J Virol, 2010. **84**(7): p. 3134-46.
- Fung, T.S. and D.X. Liu, *Human Coronavirus: Host-Pathogen Interaction*. Annu Rev Microbiol, 2019. **73**: p. 529-557.
- de Wit, E., et al., *SARS and MERS: recent insights into emerging coronaviruses*. Nat Rev Microbiol, 2016. **14**(8): p. 523-34.
- Wong, G., et al., Zoonotic origins of human coronavirus 2019 (HCoV-19 / SARS-CoV-2): why is this work important? Zool Res, 2020. **41**(3): p. 213-219.
- Azhar, E.I., et al., *Evidence for camel-to-human transmission of MERS coronavirus*. N Engl J Med, 2014. **370**(26): p. 2499-505.
- Reusken, C.B., et al., Middle East respiratory syndrome coronavirus neutralising serum antibodies in dromedary camels: a comparative serological study. Lancet Infect Dis, 2013. **13**(10): p. 859-66.
- Lu, R., et al., Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. Lancet, 2020. **395**(10224): p. 565-574.
- Latif, A.A. and S. Mukaratirwa, Zoonotic origins and animal hosts of coronaviruses causing human disease pandemics: A review. Onderstepoort J Vet Res, 2020. **87**(1): p. e1-e9.
- Zhou, P., et al., A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature, 2020. **579**(7798): p. 270-273.
- Sit, T.H.C., et al., *Infection of dogs with SARS-CoV-2*. Nature, 2020. **586**(7831): p. 776-778.
- Barrs, V.R., et al., SARS-CoV-2 in Quarantined Domestic Cats from COVID-19 Households or Close Contacts, Hong Kong, China. Emerg Infect Dis, 2020. **26**(12): p. 3071-3074.
- Newman, A., et al., First Reported Cases of SARS-CoV-2 Infection in Companion Animals - New York, March-April 2020. MMWR Morb Mortal Wkly Rep, 2020. **69**(23): p. 710-713.

18. Prince, T., et al., SARS-CoV-2 Infections in Animals: Reservoirs for Reverse Zoonosis and Models for Study. *Viruses*, 2021. **13**(3).
19. Kok, K.H., et al., Co-circulation of two SARS-CoV-2 variant strains within imported pet hamsters in Hong Kong. *Emerg Microbes Infect*, 2022. **11**(1): p. 689-698.
20. Pickering, B., et al., Divergent SARS-CoV-2 variant emerges in white-tailed deer with deer-to-human transmission. *Nat Microbiol*, 2022. **7**(12): p. 2011-2024.
21. Pagani, G., et al., Human-to-Cat SARS-CoV-2 Transmission: Case Report and Full-Genome Sequencing from an Infected Pet and Its Owner in Northern Italy. *Pathogens*, 2021. **10**(2).
22. Hosie, M.J., et al., Detection of SARS-CoV-2 in respiratory samples from cats in the UK associated with human-to-cat transmission. *Vet Rec*, 2021. **188**(8): p. e247.
23. Garigliany, M., et al., SARS-CoV-2 Natural Transmission from Human to Cat, Belgium, March 2020. *Emerg Infect Dis*, 2020. **26**(12): p. 3069-3071.
24. Piewbang, C., et al., SARS-CoV-2 Transmission from Human to Pet and Suspected Transmission from Pet to Human, Thailand. *J Clin Microbiol*, 2022. **60**(11): p. e0105822.
25. Hamer, S.A., et al., SARS-CoV-2 Infections and Viral Isolations among Serially Tested Cats and Dogs in Households with Infected Owners in Texas, USA. *Viruses*, 2021. **13**(5).
26. Goryoka, G.W., et al., One Health Investigation of SARS-CoV-2 Infection and Seropositivity among Pets in Households with Confirmed Human COVID-19 Cases-Utah and Wisconsin, 2020. *Viruses*, 2021. **13**(9).
27. Fritz, M., et al., High prevalence of SARS-CoV-2 antibodies in pets from COVID-19+ households. *One Health*, 2021. **11**: p. 100192.
28. Michelitsch, A., et al., Occurrence of Antibodies against SARS-CoV-2 in the Domestic Cat Population of Germany. *Vaccines (Basel)*, 2020. **8**(4).
29. Stevanovic, V., et al., Seroprevalence of SARS-CoV-2 infection among pet animals in Croatia and potential public health impact. *Transbound Emerg Dis*, 2021. **68**(4): p. 1767-1773.
30. Zhao, S., et al., Serologic Screening of Severe Acute Respiratory Syndrome Coronavirus 2 Infection in Cats and Dogs during First Coronavirus Disease Wave, the Netherlands. *Emerg Infect Dis*, 2021. **27**(5): p. 1362-1370.
31. Patterson, E.I., et al., Evidence of exposure to SARS-CoV-2 in cats and dogs from households in Italy. *Nat Commun*, 2020. **11**(1): p. 6231.
32. Thematic Household Survey Report No. 66 – Keeping of Dogs and Cats. 2019, Census and Statistics Department, Hong Kong SAR.
33. Ministry of Agriculture, F., and Rural Affairs. *8.60 million companion animals are raised in 6.38 million households nationwide 2021* [cited 2022 Nov. 01]; Available from: <https://www.mafra.go.kr/mafra/293/subview.do?enc=Zm5jdDF8QEB8JTJGYmJzJTJGbWFmcmElMkY2OCUyRjMyNjk3MCUyRmFydGNsVmllY5kbyUzRg%3D%3D> (in Korean).
34. Sit, T.H.C., et al., *Infection of dogs with SARS-CoV-2*. *Nature*, 2020.
35. Perera, R.A., et al., Serological assays for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), March 2020. *Euro Surveill*, 2020. **25**(16).
36. Deshpande, S.S., *Enzyme Immunoassays: From Concept to Product Development*. 1996, New York: Chapman & Hall.
37. Donnik, I.M., et al., *Coronavirus Infections of Animals: Future Risks to Humans*. *Biol Bull Russ Acad Sci*, 2021. **48**(1): p. 26-37.
38. SAR, T.G.o.H.K. *Local situation dashboard*. Available from: <https://www.coronavirus.gov.hk/eng/index.html>.
39. Schulz, C., et al., SARS-CoV-2-Specific Antibodies in Domestic Cats during First COVID-19 Wave, Europe. *Emerg Infect Dis*, 2021. **27**(12): p. 3115-3118.
40. Klaus, J., et al., Detection and Genome Sequencing of SARS-CoV-2 in a Domestic Cat with Respiratory Signs in Switzerland. *Viruses*, 2021. **13**(3).
41. Perera, R., et al., Evaluation of a SARS-CoV-2 Surrogate Virus Neutralization Test for Detection of Antibody in Human, Canine, Cat, and Hamster Sera. *J Clin Microbiol*, 2021. **59**(2).
42. Tan, C.W., et al., A SARS-CoV-2 surrogate virus neutralization test based on antibody-mediated blockage of ACE2-spike protein-protein interaction. *Nat Biotechnol*, 2020. **38**(9): p. 1073-1078.
43. Jiang, S., C. Hillyer, and L. Du, Neutralizing Antibodies against SARS-CoV-2 and Other Human Coronaviruses: (Trends in Immunology 41, 355-359; 2020). *Trends Immunol*, 2020. **41**(6): p. 545.
44. Dileepan, M., et al., Seroprevalence of SARS-CoV-2 (COVID-19) exposure in pet cats and dogs in Minnesota, USA. *Virulence*, 2021. **12**(1): p. 1597-1609.
45. Felten, S., et al., Correlation of Feline Coronavirus Shedding in Feces with Coronavirus Antibody Titer. *Pathogens*, 2020. **9**(8).
46. Klaus, J., et al., SARS-CoV-2 Infection in Dogs and Cats from Southern Germany and Northern Italy during the First Wave of the COVID-19 Pandemic. *Viruses*, 2021. **13**(8).

47. Ito, G., et al., Seroprevalence of antibodies against severe acute respiratory coronavirus 2 (SARS-CoV-2) in household dogs in Japan. *J Vet Med Sci*, 2021. **83**(11): p. 1722-1725.
48. Barroso-Arevalo, S., et al., Large-scale study on virological and serological prevalence of SARS-CoV-2 in cats and dogs in Spain. *Transbound Emerg Dis*, 2022. **69**(4): p. e759-e774.
49. Pomorska-Mol, M., et al., A cross-sectional retrospective study of SARS-CoV-2 seroprevalence in domestic cats, dogs and rabbits in Poland. *BMC Vet Res*, 2021. **17**(1): p. 322.
50. Udom, K., et al., Serological survey of antibodies against SARS-CoV-2 in dogs and cats, Thailand. *Transbound Emerg Dis*, 2022. **69**(4): p. 2140-2147.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.