

Review

Not peer-reviewed version

Freshwater Cyanobacterial Toxins, Cyanopeptides and Neurodegenerative Diseases

[Galina Nugumanova](#) , Eugene D. Ponomarev , [Sholpan Askarova](#) , [Elizaveta Fasler-Kan](#) ,
[Natasha S. Barteneva](#) *

Posted Date: 19 January 2023

doi: 10.20944/preprints202301.0357.v1

Keywords: cyanotoxins; cyanobacteria; harmful algae bloom; neurodegenerative disease; microcystin; BMAA; non-proteogenic amino acids; mistranslation; amyotrophic lateral sclerosis; Alzheimer's disease



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

Freshwater Cyanobacterial Toxins, Cyanopeptides and Neurodegenerative Diseases

Galina Nugumanova ¹, Eugene Ponomarev ¹, Sholpan Askarova ², Elizaveta Fasler-Kan ³ and Natasha S. Barteneva ^{1,4,*}

¹ School of Sciences and Humanities, Nazarbayev University, 010000, Kazakhstan

² Center for Life Sciences, National Laboratory Astana, Nazarbayev University, 010000, Kazakhstan

³ Department of Pediatric Surgery, Children's Hospital, Inselspital Bern, University of Bern, 3010 Bern, Switzerland

⁴ The Environment & Resource Efficiency Cluster (EREC), Nazarbayev University, Astana 010000, Kazakhstan

* Correspondence: natalie.barteneva@nu.edu.kz

Abstract: Cyanobacteria produce a wide range of structurally diverse cyanotoxins and bioactive cyanopeptides in freshwater, marine, and terrestrial ecosystems. The health significance of these metabolites, which include genotoxic- and neurotoxic agents, is confirmed by continued associations between the occurrence of animal and human acute toxic events and, in the long term, by associations between cyanobacteria and neurodegenerative diseases. One of the implicated mechanisms includes a misincorporation of cyanobacterial non-proteogenic amino acids leading to mistranslation and protein misfolding. A better understanding of the interaction between the cyanopeptide metabolism and the nervous system will be crucial to target or to prevent pathogenic response.

Keywords: cyanotoxins; cyanobacteria; harmful algae bloom; neurodegenerative disease; microcystin; BMAA; non-proteogenic amino acids; mistranslation; amyotrophic lateral sclerosis; Alzheimer's disease

1. Introduction

Cyanobacteria are photosynthetic gram-negative bacterial common inhabitants of diverse aquatic freshwater, marine, and terrestrial environments. They are ancient prokaryotic life forms on Earth having photosynthesis which contributed oxygen to our atmosphere over 3.5 billion years ago, and can survive in different, sometimes frequently changing environmental conditions. Cyanotoxin production is considered to be an ancient trade exceeding 2.5 billion years of age [1].

The ability of cyanobacteria to produce cyanotoxins and their ubiquity in freshwater ecosystems with increasing demands upon water resources require better detection and a more comprehensive understanding of the cyanobacterial distribution and its impact on animals and humans. Although cyanobacterial cell numbers change seasonally, the toxins can persist in water for several months, extending low-dose exposure [2]. Aquatic organisms (e.g., grazers/herbivores, fish, mammals) may consume aquatic plants with high concentrations of cyanotoxins, and trophic transfer may happen to higher-order organisms though, for some toxins, biodilution, and not biomagnification may be a predominant process in the food webs [3]. The detectable presence of cyanobacterial toxins in animal tissues has been found to be associated with mass mortalities of animals, including cows, dogs, and sea mammals [4–12], birds [13], and some human cases [14,15]. The terrestrial vertebrates affected by cyanotoxins are more diverse than was thought before [16,17]. Moreover, cyanotoxins demonstrate sublethal effects, including growth inhibition in zooplankton and aquatic plants, macroinvertebrates, and aquatic plants [18].

The predicted climate changes favor increased water temperatures, anthropogenic nutrient loadings, and freshwater cyanobacterial frequency, duration, and size of algal blooms [19–21]. The cyanobacterial abundance has increased disproportionally relative to other phytoplankton since 1945

[22]. Though our knowledge about cyanotoxins mostly comes from temperate and tropic areas, Arctic regions undergo the most pronounced and rapid climate changes, and high-latitude lakes support cyanobacteria blooming and cyanotoxins production [23]. Moreover, the occurrence and intensity of near-surface phytoplankton harmful algal blooms (HABs) have been increasing across the world [24,25] due to the eutrophication or nutrient enrichment of water bodies [26,27]. The potential presence of low doses of cyanobacterial toxins in drinking water is likely to be a continuing problem.

2. Cyanobacterial Toxins

Cyanobacteria produce a variety of toxins. Traditionally, they are divided based on functional properties into main groups: hepatotoxins, neurotoxins, dermatotoxins, and cytotoxins. Lipopolysaccharides (components of the cyanobacteria cell wall), due to their toxic effects, are classified as a separate group called – endotoxins [21,28]. The toxins found in cyanoHABs include microcystins and nodularins, and neurotoxins such as anatoxins (anatoxin-a, homoanatoxin-a, guanitoxin), ciguatoxins, saxitoxins, β -methylamino-L-alanine (BMAA) and its isomers (2,4-diaminobutyric acid (2,4-DAB) and N-(2-aminoethyl)-glycine (AEG) [21]. More than 80 cyanobacteria species are known to be toxigenic, and assays for the detection and toxicity of cyanotoxins continue to develop.

The major source of cyanotoxins for humans is drinking water. However, guidelines for water quality monitoring are limited to microcystins [29] and very few neurotoxins. Furthermore, cyanobacteria may be present in fields used for livestock grazing or can be fed to livestock directly [30–33], which aggregates in cow's milk [34,35] or bird eggs and increase human exposure [36]. There is an urgent need to detect other cyanobacterial toxins in drinking water and food and to understand how they are involved in the pathogenesis of chronic diseases such as neurodegenerative disorders.

Many cyanobacterial toxins still have to be discovered. For major toxins groups, new variants can be found and characterized. Thus, a structural variant of anatoxin-a, dihydro-anatoxin-a has been recently identified in many samples of benthic cyanobacteria, even exceeding the abundance of anatoxin-a [37]. Vacuolar spongiform myelopathy has recently been linked to aetokthonotoxin (AETX) from epiphytic cyanobacterium *Aetokthonos hydrillicola* that is growing in man-made water bodies of the southeastern United States [38]. This finding warrants further research into the potential toxins produced by epiphytic and benthic cyanobacteria species.

2.1. Microcystins (MC) Family

A full structural chemical analysis of MCs was achieved in the 1980s through a combination of spectroscopy, nuclear magnetic resonance, mass spectrometry, and amino acid analysis and demonstrated that the chemical structure of microcystins consists of a cyclic heptapeptide with two variable and five relatively conservative amino acids biosynthesized non-ribosomally via an MC synthetase gene cluster [39]. A universal nomenclature system was suggested based on the positions of amino acid residues 2 and 4 (denoted as X and Z in the original structure of microcystin i.e, microcystin -XZ) [40]. By 2019 the identification of at least 279 different MC congeners was reported in the literature [39].

Between cyanobacterial toxins, MCs are the most diverse group and the best described, though MC-LR and MC-RR—are the only two widely researched. Minority MC congeners demonstrate different toxicokinetic and toxicodynamic features [41]. Variations *in vivo* toxicity between MC congeners can be attributed to the differences in their uptake by organic anion transporting polypeptides (OATP) transport versus serine/threonine protein phosphatases (PP) 1 and 2a inhibition [42]. The toxicity of MCs depends on variations in their chemical structure and ranges over six orders of magnitude [43].

MC toxicity affects not only the liver but also the brain [44] and other organs. Multiple neurotoxic effects of MC-LR were demonstrated using multiple biological models, including birds, fishes, and mammals [45–47]. For example, using murine brain cell line as a model, congener-dependent pronounced neurodegenerative effects were identified (MC-LF>>MC-LW>MC-LR) [48].

2.2. BMAA (β -N-methylamino-L-alanine) and Isomers

BMAA is a non-proteogenic amino acid produced by all known groups of free-living and symbiotic cyanobacteria [49]. The isomeric forms of BMAA, such as 2,4-diaminobutyric acid (2,4-DAB) and N-(2-aminoethyl) glycine (AEG), can also be found in different species of cyanobacteria, including *Anabaena*, *Leptolyngbya* sp., *Oscillatoria* sp., *Merismopedia* sp., and *Microcystis aeruginosa* [50]. The isomers are detected in nature along with BMAA but are less studied. BMAA isomers are neurotoxic [51]. Recent research using larval zebrafish as a biological model identified 2,4 DAB as a more potent neurotoxin than AEG and BMAA [52].

BMAA has been shown to contribute to protein misfolding, enzyme inhibition, and neuroinflammation [53]. BMAA toxic effects were found to be related to the misincorporation of serine in multiple human proteins [54,55] which can lead to the formation of inclusion bodies in neurons [56]. Since some serine sites, such as tau serine residue 422 have been identified of key importance in neuropathologies [57], misincorporation in such sites can be particularly detrimental. Even a low level of misincorporation rate (1 per 10,000 codons) can lead to neurodegeneration in a rodent model [58]. Several groups found in their in vitro and in vivo studies that BMAA leads to the overexpression TDP-43 (TAR DNA-binding protein 43) encoded by the TARDBP gene [59–61].

Most animal models' BMAA studies were concentrated on investigating BMAA effects in the brain and other organs. Using mice Xie and co-authors reported that less than 1% of total BMAA from adult mice plasma was taken in the brain [62], i.e., BBB is not easily permeable for BMAA. The mechanism of neurotoxicity may involve a direct action on the NMDA receptor, activation of glutamate receptor 5, and induction of oxidative stress.

Recently, Han and co-authors [63] found that BMAA can serve as a substrate for human alanyl-tRNA synthetase (AlaRS), avoiding the intrinsic editing activity of AlaRS, acting as a competitive inhibitor, and comprising the editing ability of AlaRS. Terminally differentiated cells, such as neurons, are particularly susceptible to mistranslation and accumulation and forming of misfolded and aggregated proteins [58,64].

2.3 Other Cyanobacterial Neurotoxins

Traditional neurotoxins from cyanobacteria with acute effects include alkaloid or organophosphorus compounds such as (a) anatoxin-a and homologs which affect nicotinic acetylcholine alkaloid toxins and muscarinic acetylcholine receptors [65,66]; (b) saxitoxins can be produced by both dinoflagellates and by cyanobacteria from several genera including *Aphanizomenon*, *Cylindrospermopsis* and *Dolichospermum*; (c) guanitoxins which are similar in structure to organophosphates and able to irreversibly inhibit acetylcholinesterase [67]; their presence was also found in desert assemblages [68,69]. Cylindrospermopsin toxin- another frequent finding during fish kills is a highly biologically active alkaloid consisting of a tricyclic guanidine moiety combined with hydroxymethyluracil [70] interferes with cellular metabolism and causes hepatotoxic and genotoxic effects, as well as neurotoxic effects [71]. There are an increasing number of research on different groups of cyanotoxins from cyanobacteria genera—Nostocales (*Nostoc*, *Hapalosiphon*, *Fischerella*, *Anabaenopsis*, *Aphanizomenon*, *Gloeotrichia*, *Cylindrospermopsis*, *Scytonema*, *Raphidiopsis*, *Cuspidothrix*, *Nodularia*, *Stigonema*, *Calothrix*, *Cylindrospermum*, and *Desmonostoc* species) and others [72].

It is known that cyanobacteria are capable of producing different toxins which can be present during the same HABs [73]. It is not clear, however, how the environmental factors regulate the abundance of different MC congeners and isoforms of other toxins in a bloom [39]. In natural lake cyanobacteria, species composition can be complex, and cyanotoxins can be represented by a structural variety of toxins and cyanopeptides. Thus, BMAA can co-occur with its isomers (2-DAB and AEG) and have synergistic neurotoxic effects, as demonstrated in in vitro cell line experiments [74]. Moreover, the joint presence of MC-LR and BMAA leads to their interaction in vivo and to the neurotoxic effect enhancement [75]. The development of methods allowing for the assessment of multiple toxins during algal blooms is needed [76].

2.4. Cyanopeptides

The majority of cyanobacterial secondary metabolites are peptides or include peptidic substructures. Cyanopeptides are non-ribosomal peptides rich with posttranslational modifications and non-proteinogenic amino acids and consist of linear, cyclic or multicyclic molecules with basic, depsipeptidic, or lipopeptidic structures. More than 500 cyanopeptides range from app. 300 to 2000 Da have been structurally identified by 2019 [77]. Natural selection did not minimize the pool of peptides but favors the production of a wide array of different peptide structures. The biosynthetic non-ribosomal pathways of peptide synthesis are evolutionarily ancient and precede synthetic pathways of higher plants and animals. During HABs, cyanobacteria produce a tremendous amount of diverse cyanopeptides; however, their ecological significance is unclear. They can happen at surface waters in the same nanomolar concentration as MCs, exhibit toxicity towards grazers in the same micromolar range as MCs, and their production is synchronized with *Microcystis* sp. While the abundance of MCs can be monitored successfully, more studies on cyanopeptides appearance and persistence during blooms [77,78] and their potential for chronic toxicity are needed. The bioactive compounds produced by cyanobacteria are not limited by peptides and also include alkaloids, cyclophanes, terpenes, lactones, etc. Cyanobacterial compounds have a broad bioactive spectrum, with many acting as serine protease inhibitors, trypsin and chymotrypsin inhibitors, and anti-cancer compounds, capable of modulating infectious diseases [79].

2.5. Chronic Effects of Cyanobacterial Toxins

The epidemiological studies of human health impacts of chronic cyanobacterial toxins exposure are nascent. They have been associated with neurodegenerative diseases, including Alzheimer’s disease (AD), Parkinson’s disease (PD), and amyotrophic lateral sclerosis (ALS) [80–84]. Clusters of ALS and ALS-like diseases have been reported in relation to cyanobacteria in Guam, France, Japan, New Hampshire, and Wisconsin (summarized in Table 1). The slow onset of neurodegenerative diseases (time distance between exposure and possible outcome) and problems with the assessment of environmental exposure interfere with our understanding of the role and significance of cyanotoxins in neurodegenerative diseases.

Table 1. ALS/PD clusters related to environmental factors and cyanobacteria.

Location	Period	ALS/PD Cases	Water Quality	Toxic Food or Dietary Components	Key Findings	Reference
Guam	1945–1969	492	Mn↑	Cycad flour	Cycad toxic effect; Biochemical and neuropathologic abnormalities in ALS/PD diagnosed locals	[85]
Guam	1940s–60s	-	-	Cycad flour, flying foxes; food containing phytotoxins	Accumulation of cycad neurotoxins (BMAA, cycasin) in flying foxes; Flying foxes consumption → ALS-PDC	[86]
Guam and other Mariana Islands	1956–1980	39	-	-	Similar genotypic composition of Chamorros on all the Mariana Islands but different mortality rates of ALS/PD on Saipan than on Guam; Environmental factors of ALS > genetic	[87]
Guam, Canada	–	23	HABs: BMAA	Cycad flour, flying foxes (for Guam)	BMAA in tissues from frontal cortex; BMAA-containing food relates to ALS/PDC; HABs → cyanobacterial contamination water supplies →BMAA biomagnification	[88,89]

Western New Guinea	1950s–1984	-	Ca, Mg ↓	-	Environmental factors of ALS > genetic	[90]
Kii Peninsula, Japan	1961	>4	Ca, Mg, Na, KHCO ₃ , Cl ↓	-	Low mineral content in water supplies possibly leads to ND; > ALS - Mitogawa area	[91]
KII Peninsula, Japan	1972	40	Mn↑	-	Possible association of Mn to ALS.	[92]
Skaraborg, Sweden	1973–1984	70 males	-	-	Cluster of MND in Skaraborg; Agricultural occupation → MND risk.	[93]
Two Rivers, Small Wisconsin, USA	1975–1983	6	-	fish	Polychlorinated biphenyl Contaminated fish consumption → ALS risk.	[94]
France	1975–1999	18	-	-	ALS cluster in south-eastern France; Infections or environmental factors of ALS > genetic.	[95]
Italy	1980–2001	634	-	-	16 ALS clusters; Low efficiency in detoxification systems; Environmental factors of ALS (toxins)	[96]
Finland	1985–1995	576	Pb, Cd, Zn↑	-	Two ALS clusters; Environmental factors of ALS.	[97]
Enfield, NH, northeastern USA	1990–2007	278	HABs: BMAA, MC	Fish, shellfish	High ALS incidence near Lake Mascoma; Chronic exposure to cyanotoxins → ALS; Combined impact of multiple cyanotoxins.	[82]
Iraq, Saudi Arabia	1991–2001	48	BMAA	-	48 ALS cases in Persian Gulf war veterans linked to desert's crust contains BMAA; Aerosolization of cyanobacteria → inhalation of dust → BMAA exposure	[98–100]
Southern France, Hérault district	1994–2009	381	HABs: BMAA	shellfish	ALS cluster in Thau lagoon; Association with high concentrations of BMAA in mussels and oysters	[101]
Northern New England, USA	1997–2009	688	HABs, [CH ₃ Hg] ⁺	-	11 clusters of ALS grouped in 4 regions; Location of ALS cases are close to water bodies where HABs occurs; Environmental factors → ALS risk	[102]
Northern New England, USA	1997–2009	>800	HABs	-	HABs → water-quality → ALS risk	[103]
Northern New England, USA	1999–2009	-	HABs: BMAA	-	Mapping cyanobacterial HABs for northern New England lakes; Cyanotoxins increase ALS risk.	[84]
Western NH, USA	-	-	HABs: BMAA	fish	High concentrations of BMAA and DAB were found in the Lake Mascoma fish; BMAA, DAB, AEG in the air filters; ALS linked to BMAA.	[83]
France	2003–2011	72	HABs: BMAA	-	Nine ALS clusters; ALS linked to BMAA.	[104]
South Korea	2005–2017	-	HABs: BMAA and other	-	HABs exposure → ND occurrence; HABs → long-term impacts on human health	[105]
Guadeloupe	1996–2011	63	-	-	The highest incidence of ALS - Marie-Galante island; Environmental factor(s) → ALS risk	[106]

Northern and Southern Italy	2002– 2012	95	-	dietary supplements	Private wells using → ALS risk↑; Amino acid supplements → ALS risk	[107]
Annapolis, Maryland, USA	2013	3	HABs: BMAA	blue crab	High concentrations of BMAA in the crabs originated Chesapeake Bay HABs exposure → ALS occurrence	[108]

The link between BMAA and neurodegenerative diseases is yet to be further elucidated. Several studies have reported the presence of BMAA post-mortem in the brain tissues of patients who die from ALS/PD [89,109]. However, in ALS/Alzheimer’s disease [110], and other studies were not able to identify BMAA presence [111,112]. The ALS/PD neurodegenerative disorder, formerly hyperendemic in Guam-USA, Kii-Japan, and Papua-Indonesia associated with several cycad food toxins, including BMAA [113], has now been identified in both aquatic and terrestrial eco-systems in North America [82,83], The Baltic Sea [114], France [101], Sweden [115], Peru [116], and Qatar [117]; and is produced by several different cyanobacteria [49], diatoms [118], and dinoflagellates [119].

Residential exposure to environmental pollutants may play an essential role in the etiology of ALS, which is supported by non-random distribution by addresses of ALS patients [120].

2.6. Stability of Cyanotoxins

Many cyanotoxins possessing cyclic peptide structure are resistant to chemical degradation [121], highly stable, and may persist in aquatic ecosystems for weeks and months [2,122]. Thus, MCs can be retained in mussels (*Mytilus californians*) for up to eight weeks [123]. The high stability of MCs, cylindrospermopsin [124,125], and other cyanotoxins over a wide range of pH and temperature might have significant consequences for aquatic ecosystems and contribute to bioaccumulation of toxins to higher levels of food chains. These peptides are synthesized non-ribosomally and may contain non-proteinogenic amino acids [126–128].

Toxins undergoing attenuation via photodegradation may vary depending on the type of toxin, HABs timing, and environmental conditions [129]. Though cyanotoxins are resistant to chemical degradation, recent advances in molecular microbial communities research have found that toxic cyanoHABs favor the specific members of bacterioplankton with degrading abilities towards cyanotoxins [130,131]. The strains of the bacterial genera *Sphingomonas* (majority of MC-degrading bacteria), *Rhodococcus*, *Brevibacterium*, *Burkholderia*, *Mycobacterium*, *Pseudomonas*, *Novosphingobium* and others can degrade MCs in time scale from hours to days [132–134]. The genomes of some MCs-degrading bacteria are sequenced [135,136], and *mlr* gene cluster have been implicated in playing a prominent role in the sequential hydrolysis of MCs peptide bonds [137,138].

Similar to MCs, cylindrospermopsin toxin can also be degraded by bacteria isolated from cyanobacterial blooms (*Bacillus* sp., *Aeromonas* sp.) [139,140] and by some probiotic bacteria [141]. Not only bacteria but fungi demonstrate algicidal activities such as *Trichoderma citrinoviride* degrading MCs [142]. Furthermore, Mohamed and co-authors [143] summarized data on six fungal species with biodegrading activities against MCs. Between zooplankton grazers, metazoans, such as *Daphnia* and *Cyclops*, are also susceptible to cyanotoxins [144,145]. Protozoa, on the other hand, are highly resistant to cyanotoxins and demonstrate great potential in controlling harmful cyanobacteria and improving phytoplankton composition in eutrophic waters [146,147].

2.7. Data gaps in Cyanotoxins Analytical Methods

Analytical techniques (selectivity and sensitivity, fraction analysis, quality control) play a critical role in assessing the cyanotoxins effects [148,149]. There are significant data gaps in analytical methods, including (a) the absence of all the relevant standards [150]; (b) the need for validated methodologies for cyanotoxins outside the water samples; (c) the need for standardization of cyanotoxins in multi-center monitoring programs [151]; (d) the need for new technologies allowing simultaneous identification of as many toxins as possible; (e) and the need to improve robustness and

a detection limit of detection methods Another critical challenge is analyze cyanotoxins faster and feasible in situ [152]. Unsuitable analytical methods may partly explain the lack of consensus over the widespread presence of some cyanotoxins (BMAA) in aquatic ecosystems [153].

Recently, high-resolution mass spectrometry (HRMS/MS) has become more available for researchers. Current methods rely on defined cyanotoxins and cyanopeptides targets and are generally inappropriate for detecting and identifying of emerging novel compounds. The recent approach of the non-targeted analysis of pollutants and toxins in water focus on a comprehensive workflow for the acquisition and treatment of the data generated after liquid chromatography coupled with high-resolution mass spectrometry (LC-HRMS) analysis [154–157]. So-called suspect screening identifies novel compounds, including cyanotoxins and cyanopeptides based on the exact mass (m/z), presence of one or more charge states ($z = 1, 2$, etc.), expected isotope pattern and common adducts) as well as secondary fragmentation (MS2) even without reference chemicals [158–161].

Since 1957, when the first MC was detected, many methods have been developed to analyze environmental samples for cyanotoxins [162]. Nowadays, the number of studied cyanotoxins and analytical methods for their qualitative and quantitative detections have increased. Detection approaches vary in terms of accuracy, sensitivity, and specificity. The most commonly used techniques for cyanotoxins detection are enzyme-linked immunosorbent assays (ELISA), protein phosphatase inhibition assay (PPIA), molecular assays - polymerase chain reaction (PCR), and quantitative real-time PCR (qPCR) for toxins producing genotypes for cyanobacteria identification, liquid chromatography (LC) and high-performance liquid chromatography (HPLC) combined with different detectors. Among these, liquid chromatography-mass spectrometry (LC-MS) takes a special place because it identifies the target cyanotoxins with high accuracy at a significantly low detection level [163,164]. Moreover, despite the structural diversity of cyanotoxins, LC-MS allows for determining groups of toxins simultaneously. That is one extra advantage of using this detection technique integrated with bioassays and molecular assays in complex environmental samples for complete water quality assessment [165,166].

Table 2. Chemical detection methods for cyanotoxins in water samples.

№	Cyanotoxins	Detection Techniques	Sensitivity		Reference
			LOD	LOQ	
1.	MC-LR and 2 congeners	UHPLC-MS/MS	0.02–0.04 ng/mL	–	[167]
2.	MC-LR and 11 congeners	UHPLC-MS/MS	–	0.2 µg/L	[168]
3.	MC-LR and 4 congeners	LC-MS/MS	0.005–0.0817 µg/L	0.005–0.0817 µg/L	[169]
	Nodularin		0.0048 µg/L	0.0048 µg/L	
	Anatoxin-a		0.0001 µg/L	0.0004 µg/L	
	Cylindrospermopsin		0.0001 µg/L	0.0004 µg/L	
4.	BMAA	UHPLC-MS/MS	0.02 pg/µL	0.05 pg/µL	[170]
	2,4-DAB		0.04 pg/µL	0.13 pg/µL	
5.	MC-LR and 7 congeners	LC-MS/MS	–	0.04–0.5 µg/L	[171]
	Anatoxin-a		–	0.02 µg/L	
	Cylindrospermopsin (and deoxyCYN)		–	0.01–0.02 µg/L	
	Saxitoxins (4 congeners), GTX (5 congeners), decarbamoylgonyautoxin, N-sulfogonyautoxins-1 and -2		–	0.1–2 µg/L	
6.	MC-LR and 11 congeners	HPLC-MS/MS	0.01±0.01–0.19±0.2 µg/L	0.04±0.04–0.64±0.65 µg/L	[172]
	Nodularin		0.04 ± 0.02 µg/L	0.13 ± 0.06 µg/L	
7.	MC-LR and 2 congeners	HPLC-UV/PDA	3–4 µg/L	9–13 µg/L	[173]
8.	MC-LR and 2 congeners	HPLC-HRMS	0.002 µg/L	–	[174]
	Anatoxin-a,		0.004 ng/mL	0.01 ng/mL	
9.	Cylindrospermopsin	HILIC-MS/MS	0.07 ng/mL	0.23 ng/mL	[175]
	Saxitoxin		0.01 ng/mL	0.04 ng/mL	

	MC-LR and 4 congeners Nodularin	RPLC- MS/MS	0.02–0.08 ng/mL 0.05 ng/mL	0.07–0.28 ng/mL 0.18 ng/mL	
10.	MC-LR and 5 congeners Nodularin Cylindrospermopsin	UHPLC-MS/MS	– – –	0.046–0.052 µg/L 0.049 µg/L 0.052 µg/L	[176]
11.	BMAA DAB AEG BAMA	UHPLC-HRMS	5 ng/L 3 ng/L 2 ng/L 5 ng/L	10 ng/L 5 ng/L 5 ng/L 10 ng/L	[177]
12.	MC-LR and 5 congeners Cylindrospermopsin Saxitoxin, dcSTX	HPLC-MS/MS	0.0003–0.0009 µg/L 0.0005 µg/L 0.0009–0.0013 µg/L	– – –	[178]
13.	BMAA	LC-MS/MS	10 ng/L	–	[179]
14.	MC-LR and 5 congeners Nodularin Anatoxin-a Cylindrospermopsin	LC-MS/MS	0.04–0.8 µg/L 0.3 µg/L 0.27 µg/L 0.14 µg/L	0.1–2.3 µg/L 0.9 µg/L 0.81 µg/L 0.4 µg/L	[180]
15.	MC-LR and 2 congeners	HPLC-DAD	0.08–0.15 µg/L	–	[181]
16.	MC-LR and 11 congeners Nodularin Anatoxin-a Cylindrospermopsin	LC-MS/MS	0.001–0.007 µg/L 0.002 µg/L 0.001 µg/L 0.001 µg/L	0.003–0.020 µg/L 0.006 µg/L 0.003 µg/L 0.003 µg/L	[182]
17.	MCs Anatoxin-a Cylindrospermopsin Saxitoxin BMAA	UPLC-MS/MS	0.005 µg/L 0.02 µg/L 0.02 µg/L 0.8 µg/L 0.03 µg/L	– – – – –	[183]
18.	BMAA	LC-MS/MS	0.030 µg/L	0.096 µg/L	[184]
19.	MC-LR and 11 congeners Anatoxin-a Homoanatoxin-a Cylindrospermopsin	on-line SPE – UHPLC-HRMS	5–37 ng/L 15–18 ng/L 11–12 ng/L 41–53 ng/L	15–130 ng/L 50–60 ng/L 30–40 ng/L 130–170 ng/L	[185]
20.	MC-LR and 1 congener Anatoxin-a	HPLC-DAD	0.2–0.3 µg/L 1.1 ng/L	– 2.5 ng/L	[186]
21.	Cylindrospermopsin Saxitoxins (4 congeners) GTx (7 congeners)	UHPLC-MS/MS	10.9 ng/L 3.5–9 ng/L 18.5–54.5 ng/L	21.7 ng/L 7.1–26.9 ng/L 42.2–227.6 ng/L	[187]
22.	MC-LR and 7 congeners Nodularin	UHPLC-MS/MS	0.1 µg/L 0.1 µg/L	0.5 µg/L 0.5 µg/L	[164]
23.	MC-LR and 2 congeners	UHPLC-MS/MS	0.1 µg/L	24 µg/L	[188]
24.	Cylindrospermopsin	UHPLC-MS/MS	0.029 µg/L	0.091 µg/L	[189]
25.	Saxitoxins (4 congeners)	on-line SPE– HILIC-HRMS	0.72 –3.9 ng/L	2.4–13 ng/L	[190]
26.	MC-LR and 1 congener Nodularin Anatoxin-a Cylindrospermopsin BMAA DAB AEG	tandem-SPE- HILIC-MS/MS	0.0012–0.0021 µg/L 0.0021 µg/L 0.03 µg/L 0.0012 µg/L 0.015 µg/L 0.009 µg/L 0.006 µg/L	0.004–0.007 µg/L 0.007 µg/L 0.1 µg/L 0.004 µg/L 0.05 µg/L 0.03 µg/L 0.02 µg/L	[191]
27.	BMAA DAB	LC-MS/MS	2.8 ng/mL 1.7 ng/mL	– –	[192]
28.	BMAA BAMA DAB AEG	on-line SPE- UHPLC-HRMS	10 ng/L 10 ng/L 10 ng/L 5 ng/L	– – – –	[153]

29.	BMA	UHPLC-MS/MS	–	2.5 µg/L	[193]
	AEG		–	2.5 µg/L	
	DABA		–	5 µg/L	
30.	MC-LR and 7 congeners	UHPLC-MS/MS (ESI)	0.02–0.2 µg/L	0.05–0.5 µg/L	[194]

LOD—limit of detection; LOQ—limit of quantification; UHPLC—ultra high-performance liquid chromatography; HILIC-MS/MS—hydrophilic interaction liquid chromatography-tandem mass spectrometry; RPLC-MS/MS—reverse phase chromatography tandem mass spectrometry; UV/PDA—ultraviolet/photodiode array detection; DAD—diode array detector; ESI—electrospray ionization; SPE—solid phase extraction; BAMA—β-amino-N-methylalanine; GTX—gonyautoxins; dcSTX—decarbamoylsaxitoxin.

As separation instruments, HPLC and UHPLC are usually used. UHPLC, is faster due to the higher pressure applied, and the online SPE-procedure provides reduced sample time processing [2,181]. HPLC-UV/PDA [173], HPLC-DAD [186] have less LOD values (3–4 µg/L, 0.2–0.3 µg/L, respectively) for MCs than LC-MS/MS where minimum LOD values vary within 0.0003–0.1 µg/L [164,178,188]. Concerning other cyanotoxins, MS detection techniques also provide relatively low values of LOD and LOQ. The LC-MS method requires expensive instruments and thorough sample preparation, which makes it a time-consuming procedure. That limits LC-MS techniques' application as ubiquitous [195]. Nevertheless, this method remains preferable for precise quantitative analysis of cyanotoxins in water samples.

3. Toxin Exposure Pathways

Major cyanotoxins exposure routes include ingestion through drinking water or dermal contact with recreational waters [196], also through food, and inhalation since cyanotoxins were identified in aerosols generated by HABs [197]. Historically, cyanoHABs were considered a public health threat to freshwater lakes, rivers, and reservoirs. However, freshwater-sourced MCs can accumulate in marine mollusks in concentrations 100-fold greater than in surrounding water [10,123].

3.1. Transport of Cyanotoxins in Freshwater and Marine Systems

Recent studies demonstrated that cyanotoxins could persist during transport into estuarine and marine waters and can directly affect marine ecosystems [10,123,198–200]. MCs and other toxins produced by freshwater cyanobacteria can enter the marine ecosystem via freshwater channels and outflows [12]. This changes HABs management approach, requiring monitoring of multiple toxins across the freshwater-to-marine continuum and including cyanotoxins in marine and estuarine monitoring [201].

3.2. Toxin exposure pathways: Oral (Drinking water)

When drinking water is impacted by cyanobacterial toxins resulting from HABs and not treated adequately to reduce the cyanotoxin levels, it can cause serious effects on the entire region [197].

The causes of cyanobacteria proliferation in urban environments are mainly the disposal of untreated domestic sewage in water reservoirs and surface runoff water from soils. In analyzing sewage disposal systems in the main cities of Kazakhstan—Almaty and Astana, the efficiency of biogenic compounds removal remains unsatisfactory, reaching only 30–40%. This eutrophication is due to the increase of nutrients, such as phosphorus and nitrogen, arising from human action, representing a serious risk to the health of living beings and drastically reducing water quality. To cope with this problem, the possibility of intensifying nitrogen and phosphorus removal using zeolite as a biofilm carrier in an activated sludge tank is examined [202].

3.3. Toxin Exposure Pathways: Oral (Food)

Food is an important source of cyanotoxin exposure [203]. Worryingly, exposure of crop plants to cyanotoxins through irrigation was already demonstrated [204–206]. For centuries some species of Nostoc—the symbiotic colonial cyanobacteria *N. flagelliforme*, *N. commune*, and *N. sphaeroides* –

have been wild-harvested and consumed as a part of the traditional diet by indigenous people in different countries, including Peru, China, Ecuador, Mexico, Fiji, Philippines, Mongolia [209,210].

Chronic dietary exposure to BMAA present in the traditional Chamorro diet was associated with the formation of both β -amyloid deposits and neurofibrils tangles (NFT) found in brain tissues of Chamorros who died with ALS/ Parkinson's dementia complex (ALS/PD). BMAA occurs not only as a free amino acid at different levels of the trophic chain (cyanobacteria *Nostoc* sp., root symbioses, cycad seeds, flying foxes, and brain tissues of Chamorro people who passed away from ALS/PD) but can also be released by acid hydrolysis increasing in concentrations 10- to 240-fold [207]. Vervet monkeys fed for only six months with BMAA-dosed fruit developed β -amyloid deposits and NFT in the brain. Increasing the amount of L-serine in the vervets diet reduced the density of NFT and the risk of neurodegenerative pathological brain findings [56]. Recently, Downing and co-authors [208] revealed that human liver hepatocyte and intestinal epithelial cultures could not metabolize BMAA, demonstrating that BMAA detoxication is impossible and BMAA will likely accumulate in these cells [53].

3.4. Toxin Exposure Pathways: Air (Aerosolization)

The algae can be dispersed by air [211], and aerosol can be created from algae during HABs [212]. The increase in the salinity of freshwater streams is likely to influence the abundance and diversity of aerosolized bacteria [213]. The cyanotoxins may be transported in aerosols from lakes with high concentrations of toxigenic cyanobacteria [214–217]. Recent findings with rat models confirmed that BMAA exposure was insufficient in producing gross toxic effects; however, it still leaves the possibility of lifelong exposure via inhalation [218].

MC-LR exposure in the existing rodent models increases lung infiltration with granulocytes [219,220] and increases proinflammatory cytokine expression [221,222]. Recently, Breidenbach and co-authors [223] reported that human airway epithelium response to MC-LR is represented by proinflammatory phenotype, including chemokines.

The aerosolization of cyanobacteria was proposed as a risk factor for ALS [100]. Aerial link of exposure was investigated with ALS/PD. BMAA and its isomers (DAB and AEG) were measured in air filters around lake Mascoma [83]. Moreover, Facciponte and co-authors [224] found that humans routinely inhale aerosolized cyanobacteria. Using PCR, authors identified cyanobacteria at high frequencies in the upper respiratory tract (93.20%) and central airway (79.31%). They concluded that cyanobacteria exposure might be a prevalent and chronic phenomenon and not necessarily restricted to water bodies.

Autoradiographic imaging in mice showed a distinct localization of radioactivity in olfactory mucosa and bulb following intranasal instillation of radiolabelled BMAA, confirming a direct transfer of BMAA via olfactory pathways to mice brain circumventing the blood-brain barrier [225].

3.6. Natural Model of Toxin Exposure

The complexity of neurodegenerative diseases requires a deep understanding of the disease biology and makes it challenging to develop a model of cyanotoxin exposure close to neurodegenerative findings in humans due to the species-specific variations in the phosphorylation and cleavage of the tau protein [226]. Natural animal models should recapitulate two major features of human neurodegenerative diseases: A β deposition and NFT formation. Chronic low BMAA concentrations induce neurodegenerative changes in non-human primates [56,227]. BMAA can bioaccumulate in marine apex predators such as dolphins and sharks, and in humans [207,228,229]. It was detected in the brains of stranded dolphins with pathological hallmarks of AD at concentrations higher than those found post-mortem in individuals with ALS and AD [230]. Chronic low BMAA concentrations induce neurodegenerative changes in non-human primates [56,227]. There are increased numbers of β -amyloid+ and dystrophic neurites in the auditory cortex compared to the visual cortex and brainstem [230].

3.7. Cyanotoxins and Infections

BMAA can facilitate most of the mechanisms related to neurodegeneration [231]. Thus, Lobner and co-authors [232] demonstrated that BMAA at the concentration range of 10–100 μmol potentiates neurotoxicity induced by amyloid- β and NMDA.

STX doubled the quantity of ZIKV-induced neural cell death in progenitor areas of human brain organoids, while the chronic ingestion of water contaminated with STX before and during gestation caused brain abnormalities in offspring of ZIKV-infected immunocompetent C57BL/6J mice. These results raise a public health concern regarding the consequences of arbovirus outbreaks in areas with droughts and/or frequent freshwater cyanobacterial blooms [233].

The outbreak of Zika syndrome coincided with a major drought in the region between 2012 and 2016. Characteristic of dry seasons, the concentration of nutrients from untreated effluents and lower volume of water, and an increase in atmospheric temperature allow greater blooming of cyanobacteria. Consequently, the concentration of cyanotoxins, such as saxitoxins, increases. This led authors to formulate the hypothesis that cyanobacteria in the water supply would be a causal cofactor of zika-associated microcephaly.

4. Mechanisms of Brain Toxicity

Well-studied neurotoxins of algal origin are alkaloids saxitoxins (STXs) that have been identified in dinoflagellates and several cyanobacterial genera, including *Anabaena*, *Aphanizomenon*, *Planktothrix*, *Cylindrospermopsis*, and *Scytonema* [234,235]. STXs are represented by more than 50 structural analogs commonly known as paralytic shellfish toxins (PSTs). They block the passage of sodium across a biological membrane and interfere with potassium and calcium-mediated ion channels [72].

While the pathophysiology of some toxins (STXs, anatoxins, etc.) are relatively well studied, others, such as ciguatera, are not clear. Recently, the neurotoxic effects of cyanopeptides attracted more attention [77,236–238]. Some cyanopeptides exhibit anti-proliferative effects on tubulin and microtubules essential for neurons. Thus, the anti-proliferative toxic cyclodepsipeptides cryptophycins are 100–1000 fold compared with paclitaxel and vinblastin [236]. Further research needs to improve analytical methods and assess potentially toxic cyanopeptides.

4.1. Neurodevelopmental Effects

The link between neurodegeneration and neonatal BMAA exposure, dose-dependent neuronal loss, beta-amyloid deposition, and behavioral deficits was recently demonstrated in a rat model [239]. Autoradiographic imaging confirmed transplacental research of radiolabelled BMAA and specific uptake in mouse fetal [240]. Furthermore, in neonatal rats, the free BMAA concentration was higher in the neonatal brain than in peripheral tissues such as the thymus, pancreas, and spleen, except for the liver. The level of protein-associated BMAA was significantly higher in the hippocampus than in other brain regions [241]. The BMAA exposure to neural stem cells decreased neurite outgrowth, and a number of neurites in neural stem cells (NSC) [242]. The authors conclude that BMAA acts as a developmental toxin. BMAA can negatively impact NSC homeostasis, increasing susceptibility to neurodegenerative disease later in life [242]. Perinatal exposure in mice, even with low doses of BMAA, leads to neurobehavioral disturbances during the postnatal period and adulthood [243].

4.2. Blood-Brain Barrier (BBB)

The blood-brain barrier and the blood-CSF barrier separate CNS from blood and include the endothelial lining of the brain capillaries associated with astrocytes, pericytes, and neurons. The pericytes and astrocytes are closely associated with the endothelial cells and are required for capillary maturation (pericytes) and the maintenance of the permeability-barrier functions (astrocytes). The basement membrane (contains laminin, proteoglycans, fibronectin, collagen IV, nidogen, and entactin) and is essential for blood-barrier differentiation. BBB separates neurons from the circulating blood and maintains the internal chemical composition of the brain "milieu" responsible

for the proper functioning of neuronal circuits, neurogenesis, angiogenesis, synaptic transmission, etc. BBB breakdown due to disruption of the tight junctions may result in synaptic and neuronal dysfunction and contribute to neurodegenerative disorders such as ALS, Alzheimer's disease, Parkinson's disease, and multiple sclerosis [244].

Berntzon and co-authors found BMAA in the CSF of a patient with ALS and some controls, though they did not confirm a prevalence of BMAA findings in ALS patients [109]. Significant amounts of BMAA were found in brain tissues of American ALS and Alzheimer's disease patients, confirming the ability of BMAA to cross the BBB [110]. Alternative findings regarding AD patients were reported by Meneely and co-authors [112]. The other possible route of entry to the CNS is through the olfactory epithelium and the nasal passage or via the blood. The cyanobacterial neurotoxin BMAA can be directly transferred through olfactory pathways circumventing the BBB in mice and directly affecting olfactory neurons [225].

Microcystin-LR (MC-LR) has been confirmed to cause blood-brain barrier disruption and enter the brain tissue, resulting in non-negligible toxic effects. However, the neurotoxicity of MC-LR is mainly unknown. This study revealed that MC-LR disrupted the function of the ubiquitin-proteasome system in neurons, which inhibited the degradation of α -synuclein (α -syn), leading to its release from neurons for transport into microglia. α -Syn is the main component of Lewy bodies, which has been identified as one of the main pathological features of Parkinson's disease (PD). In vitro, we observed that α -syn mediated by MC-LR activated HMC3 cells and polarized them towards M1 type. In addition, we confirmed that α -syn was transported into HMC3 cells through TLR4 receptors and activated the NLRP3 inflammasome, which in turn enhanced the maturation and release of IL-18 and IL-1 β [245].

4.3. Glia

Microglial activation and neuroinflammation are common to many neurodegenerative diseases. Glial cells, including microglia, have long been suspected of playing a role in Alzheimer's disease but only because of their ability to react to neuronal dysfunctions (e.g., amyloid and Tau aggregates). This neurocentric view, which considered glial cells as secondary, has been challenged recently by the results of genetic association studies identifying genetic loci associated with the risk of Alzheimer's that are associated with genes preferentially or exclusively expressed in glial cells [246].

The research on cyanopeptides effects on glia is limited. Chiu and co-authors [247,248] demonstrated a gliotoxicity of BMAA using the olfactory ensheathing cell as in vitro model. A study conducted by Bubic and co-workers [249] showed that depsipeptide planktopeptin and anabaenopeptins impair the metabolic activities of normal human astrocytes via membrane perforation, oxidative stress, and changes in mitochondrial metabolism. Later, Mello and co-authors showed cytotoxic effects of BMAA and MC-LR on primary astrocytes isolated from mixed adult brain cell cultures [250], and Soto, with co-workers, demonstrated damaging BMAA effects on Muller's glial cells [251]. Both glial cells and neurons can uptake and accumulate BMAA, as demonstrated using a specific, polyclonal antibody against BMAA [252].

The role of dysfunctional astrocytes in the pathogenesis of ALS and other neurodegenerative diseases indicates that astrocytes may be targeted with strategies for their revival. These strategies may include direct intervention on astrocytes with modulatory medicines, exosomes and miRNA-based therapies, or their replacement.

5. Cyanotoxins, Cyanopeptides and Neurodegenerative Diseases

A central dogma of age-related neurodegenerative diseases claims that the accumulation and propagation of aggregated proteins cause neurodegeneration [253]. Recently, a mechanism that does not involve a specific neuropathogenic protein but is mediated by error-prone translation leading to stochastic near-cognate missense substitutions was suggested. Drummond and Wilke proposed in 2008 that tolerance to translation errors of certain proteins provides a new mechanism to explain their propensity to misfold pathologically. mistranslation destabilizes the proteome by leading to misfolding and accumulation in the cells of potentially toxic protein aggregates [58,254]. The finding

that translational error increases with age in some biological models (*Drosophila*) [255] may suggest the possibility that the rate of translation leading to aging-related proteostasis failure may be a key event in early ND diseases [256]. The mistranslating cells exhibit severely inhibited protein synthesis and formation of protein aggregates in the cellular ND model [257]. Aminoacyl-tRNA synthetases (AARSs) catalyze covalent binding tRNA with their cognate amino acids and are 2–3 orders of magnitude more selective than other amino acid-utilizing [258].

Hundreds of non-proteinogenic amino acids produced by cyanobacteria include BMAA and can, in principle, enter human protein synthesis through foods and drinking water. Earlier studies support the ability of BMAA to be incorporated into the proteins [54,259]. It has been suggested that BMAA is misincorporated at serine codons during protein synthesis [54]. However, recently Han and co-authors [63] demonstrated that BMAA is not a substrate for human seryl-tRNA synthetase (SerRS) but a substrate for human alanyl-tRNA synthetase (AlaRS) and can form BMAA-tRNA^{Ala} by escaping from the intrinsic AlaRS proofreading activity. Furthermore, BMAA inhibits the cognate amino acid activation, the editing functions of AlaRS, and the deacylation activity of HsAlaRS on Ser-tRNA^{Ala} [63]. The AlaRS possess canonical and non-canonical cellular functions and are predominantly linked to neurodegenerative disorders in human and mouse models [260]. Furthermore, using transcriptomic analysis, Wang and co-authors [261] confirmed that BMAA could alter the expression of major genes encoding components related to translation in prokaryotes (diazotrophic algae *Anabaena*). The ability of BMAA to affect protein hemostasis can be evolutionarily ancient and initially directed to inhibit the growth of neighboring microalgae. The inhibition of cell growth and progression in the cell cycle of eukaryotic cells was demonstrated in *in vitro* experiments [262]. The production of cyclic peptides, including non-proteinogenic amino acids, leads to the lysis of cyanobacteria and may be an effective control mechanism of cyanobacterial density during algal blooms [126].

6. Conclusions

The structural variety of cyanotoxins and cyanopeptides is produced during cyanobacterial blooms. Many structural aspects of key metabolites involved in the cyanotoxins pathways have yet to be elucidated. However, it is becoming clear that non-proteinogenic amino acids, free-existing or initially a part of cyanopeptides, may affect protein hemostasis and lead to mistranslation and misfolding of proteins in eukaryotic cells, building a link to neurodegenerative diseases development. There are many aspects pertaining to the regulation, role, and function of these compounds that also require the development of novel detection approaches. This knowledge may be harnessed to identify novel biomarkers for neurodegenerative diseases and new targets for interventions.

Author Contributions: G.N. and N.S.B. wrote an original draft, E.P., E.F.-K., S.A. edited the manuscript. S.A., E.F.-K., and N.S.B. contributed to the funding of the research.

Funding: This research received funding from Nazarbayev University (PURE ID: 16482715) and Kazakhstan MS grant # SSH2022019.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We are thankful to Dmitry Malashenkov, Nazarbayev University for helpful discussion.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Murray, S.A.; Mihali, T.K.; Neilan, B.A. Extraordinary conservation, gene loss, and positive selection in the evolution of an ancient neurotoxin. *Mol. Biol. Evol.* **2011**, *28*, 1173–1182. doi:10.1093/molbev/msq295
2. Chorus, I.; Welker, M. *Toxic cyanobacteria in water: a guide to their public health consequences, monitoring and management*, 2nd ed.; Taylor & Francis: Boca Raton (FL), on behalf of the World Health Organization, Geneva, CH, 2021. doi: 10.1201/9781003081449

3. Kozłowski-Suzuki, B.; Wilson, A.E.; da Silva Ferrao-Filho, A. Biomagnification or biodilution of microcystins in aquatic foodwebs? Meta-analyses of laboratory and field studies. *Harmful Algae* **2012**, *18*, 47-55. doi:10.1016/j.hal.2012.04.002
4. Pybus, M.; Hobson, D.; Onderka, D. Mass mortality of bats due to probable blue-green algal toxicity. *J. Wildl. Dis.* **1986**, *22*, 449-450. doi:10.7589/0090-3558-22.3.449
5. Mahmood, N.A.; Carmichael, W.W.; Pfahler, D. Anticholinesterase poisonings in dogs from a cyanobacterial (blue-green algae) bloom dominated by *Anabaena flos-aquae*. *Am. J. Vet. Res.* **1988**, *49*, 500-503. PMID: 3132068
6. Henriksen, P.; Carmichael, W.W.; An, J.; Moestrup, O. Detection of an anatoxin-a (s)-like anticholinesterase in natural blooms and cultures of cyanobacteria/blue-green algae from Danish lakes and in the stomach contents of poisoned birds. *Toxicon* **1997**, *35*, 901-913. doi: 10.1016/s0041-0101(96)00190-0
7. Mez, K.; Beattie, K.A.; Codd, G.A.; Hanselmann, K.; Hauser, B.; Naegeli, H.; Preisig, H.R. Identification of a microcystin in benthic cyanobacteria linked to cattle deaths on alpine pastures in Switzerland. *Eur. J. Phycol.* **1997**, *32*, 111-117. doi:10.1080/09670269710001737029
8. Saker, M.; Thomas, A.; Norton, J. Cattle mortality attributed to the toxic cyanobacterium *Cylindrospermopsis raciborskii* in an outback region of north Queensland. *Environ. Toxicol.* **1999**, *14*, 179-182. doi:10.1002/(SICI)1522-7278(199902)14:1<179::AID-TOX23>3.0.CO;2-G
9. Codd, G.A.; Lindsay, J.; Young, F.M.; Morrison, L.F.; Metcalf, J.S. Harmful cyanobacteria: from mass mortalities to management measures. In *Harmful Cyanobacteria*; Huisman, J., Matthijs, H.C.P., Visser, P.M., Eds.; Springer: Dordrecht, The Netherlands, 2005. pp. 1-23. doi: 10.1007/1-4020-3022-3_1
10. Miller, M.A.; Kudela, R.M.; Mekebri, A.; Crane, D.; Oates, S.C.; Tinker, M.T.; Staedler, M.; Miller, W.A.; Toy-Choutka, S.; Dominik, C. Evidence for a novel marine harmful algal bloom: cyanotoxin (microcystin) transfer from land to sea otters. *PLoS One* **2010**, *5*, e12576. doi:10.1371/journal.pone.0012576
11. Chatziefthimiou, A.D.; Richer, R.; Rowles, H.; Powell, J.T.; Metcalf, J.S. Cyanotoxins as a potential cause of dog poisonings in desert environments. *Vet. Rec.*, **2014**, *174*, 484-485. doi:10.1136/vr.g3176
12. Danil, K.; Berman, M.; Frame, E.; Preti, A.; Fire, S.E.; Leighfield, T.; Carretta, J.; Carter, M.L.; Lefebvre, K. Marine algal toxins and their vectors in southern California cetaceans. *Harmful Algae* **2021**, *103*, 102000. doi:10.1016/j.hal.2021.102000
13. Rattner, B.A.; Wazniak, C.E.; Lankton, J.S.; McGowan, P.C.; Drovetski, S.V.; Egerton, T.A. Review of harmful algal bloom effects on birds with implications for avian wildlife in the Chesapeake Bay region. *Harmful Algae* **2022**, 102319. doi:10.1016/j.hal.2022.102319
14. Pouria, S.; de Andrade, A.; Barbosa, J.; Cavalcanti, R.; Barreto, V.; Ward, C.; Preiser, W.; Poon, G.K.; Neild, G.; Codd, G. Fatal microcystin intoxication in haemodialysis unit in Caruaru, Brazil. *The Lancet* **1998**, *352*, 21-26. doi:10.1016/S0140-6736(97)12285-1
15. Carmichael, W.W.; Azevedo, S.; An, J.S.; Molica, R.; Jochimsen, E.M.; Lau, S.; Rinehart, K.L.; Shaw, G.R.; Eaglesham, G.K. Human fatalities from cyanobacteria: chemical and biological evidence for cyanotoxins. *Environ. Health Perspect.* **2001**, *109*, 663-668. doi:10.1289/ehp.01109663
16. Briand, J.-F.; Jacquet, S.; Bernard, C.; Humbert, J.-F. Health hazards for terrestrial vertebrates from toxic cyanobacteria in surface water ecosystems. *Vet. Res.* **2003**, *34*, 361-377. doi: 10.1051/vetres:2003019
17. McLellan, N.L.; Manderville, R.A. Toxic mechanisms of microcystins in mammals. *Toxicol. Res.* **2017**, *6*, 391-405. doi:10.1039/c7tx00043j
18. White, S.H.; Duivenvoorden, L.J.; Fabbro, L.D. A decision-making framework for ecological impacts associated with the accumulation of cyanotoxins (cylindrospermopsin and microcystin). *Lakes Reserv.: Res. Manag.* **2005**, *10*, 25-37. doi:10.1111/j.1440-1770.2005.00258.x
19. Paerl, H.W.; Huisman, J. Blooms like it hot. *Science* **2008**, *320*, 57-58. doi: 10.1126/science.1155398
20. Moe, S.J.; Haande, S.; Couture, R.-M. Climate change, cyanobacteria blooms and ecological status of lakes: A Bayesian network approach. *Ecol. Modell.* **2016**, *337*, 330-347. doi:10.1016/j.ecolmodel.2016.07.004
21. Huisman, J.; Codd, G.A.; Paerl, H.W.; Ibelings, B.W.; Verspagen, J.M.; Visser, P.M. Cyanobacterial blooms. *Nat. Rev. Microbiol.* **2018**, *16*, 471-483. doi: 10.1038/s41579-018-0040-1
22. Taranu, Z.E.; Gregory-Eaves, I.; Leavitt, P.R.; Bunting, L.; Buchaca, T.; Catalan, J.; Domaizon, I.; Guilizzoni, P.; Lami, A.; McGowan, S. Acceleration of cyanobacterial dominance in north temperate-subarctic lakes during the Anthropocene. *Ecol. Lett.* **2015**, *18*, 375-384. doi:10.1111/ele.12420
23. Trout-Haney, J.V.; Wood, Z.T.; Cottingham, K.L. Presence of the cyanotoxin microcystin in Arctic Lakes of Southwestern Greenland. *Toxins* **2016**, *8*, 256. doi:10.3390/toxins8090256
24. Ho, J.C.; Michalak, A.M.; Pahlevan, N. Widespread global increase in intense lake phytoplankton blooms since the 1980s. *Nature* **2019**, *574*, 667-670. doi:10.1038/s41586-019-1648-7
25. Gobler, C.J. Climate change and harmful algal blooms: insights and perspective. *Harmful algae* **2020**, *91*, 101731. doi:10.1016/j.hal.2019.101731
26. Bowling, L.; Baker, P. Major cyanobacterial bloom in the Barwon-Darling River, Australia, in 1991, and underlying limnological conditions. *Mar. Freshw. Res.* **1996**, *47*, 643-657. doi:10.1071/MF9960643

27. Heisler, J.; Glibert, P.M.; Burkholder, J.M.; Anderson, D.M.; Cochlan, W.; Dennison, W.C.; Dortch, Q.; Gobler, C.J.; Heil, C.A.; Humphries, E.; et al. Eutrophication and harmful algal blooms: A scientific consensus. *Harmful Algae* **2008**, *8*, 3-13. doi:10.1016/j.hal.2008.08.006
28. Sivonen, K.; Jones, G. Cyanobacterial toxins. In *Encyclopedia of microbiology*; Elsevier Scientific Publ. Co., Amsterdam, 2009; pp. 290-307. doi:10.1016/B978-012373944-5.00005-5
29. World Health Organization. Cyanobacterial toxins: microcystin-LR in drinking-water. No. WHO/SDE/WSH/03.04/57. World Health Organization, 2020.
30. Simkus, A.; Oberauskas, V.; Laugalis, J.; Želvytė, R.; Monkevičienė, I.; Sederevičius, A.; Šimkienė, A.; Pauliukas, K. The effect of weed *Spirulina platensis* on the milk production in cows. *Vet. Zootech.* **2007**, *38*, 74-77.
31. Kulpys, J. ; Paulauskas, E. ; Pilipavicius V. ; Stankevicius, R. Influence of cyanobacteria *Arthrospira (Spirulina) platensis* biomass additives towards the body condition of lactation cows and biochemical milk indexes. *Agric. Res.* **2009**, *7*, 823–835
32. Christaki, E.; Florou-Paneri, P.; Bonos, E. Microalgae: a novel ingredient in nutrition. *Int. J. Food Sci.* **2011**, *62*, 794-799. doi:10.3109/09637486.2011.582460
33. McGorum, B.C.; Pirie, R.S.; Glendinning, L.; McLachlan, G.; Metcalf, J.S.; Banack, S.A.; Cox, P.A.; Codd, G.A. Grazing livestock are exposed to terrestrial cyanobacteria. *Vet. Res.* **2015**, *46*, 1-10. doi:10.1186/s13567-015-0143-x
34. Andersson, M.; Karlsson, O.; Bergström, U.; Brittebo, E.B.; Brandt, I. Maternal transfer of the cyanobacterial neurotoxin β -N-methylamino-L-alanine (BMAA) via milk to suckling offspring. *PLoS One* **2013**, *8*, e78133. doi: 10.1371/journal.pone.0078133
35. Andersson, M.; Karlsson, O.; Banack, S.A.; Brandt, I. Transfer of developmental neurotoxin β -N-methylamino-L-alanine (BMAA) via milk to nursed offspring: Studies by mass spectrometry and image analysis. *Toxicol. Lett.* **2016**, *258*, 108-114. doi:10.1016/j.toxlet.2016.06.015
36. Andersson, M.; Karlsson, O.; Brandt, I. The environmental neurotoxin β -N-methylamino-L-alanine (l-BMAA) is deposited into birds' eggs. *Ecotoxicol. Environ. Saf.* **2018**, *147*, 720-724. doi:10.1016/j.ecoenv.2017.09.032
37. Puddick, J.; van Ginkel, R.; Page, C.D.; Murray, J.S.; Greenhough, H.E.; Bowater, J.; Selwood, A.I.; Wood, S.A.; Prinsep, M.R.; Truman, P. Acute toxicity of dihydroanatoxin-a from *Microcoleus autumnalis* in comparison to anatoxin-a. *Chemosphere* **2021**, *263*, 127937. doi:10.1016/j.chemosphere.2020.127937
38. Breinlinger, S.; Phillips, T.J.; Haram, B.N.; Mareš, J.; Martínez Yerena, J.A.; Hrouzek, P.; Sobotka, R.; Henderson, W.M.; Schmieder, P.; Williams, S.M. Hunting the eagle killer: a cyanobacterial neurotoxin causes vacuolar myelinopathy. *Science* **2021**, *371*, eaax9050. doi: 10.1126/science.aax9050
39. Bouaïcha, N.; Miles, C.O.; Beach, D.G.; Labidi, Z.; Djabri, A.; Benayache, N.Y.; Nguyen-Quang, T. Structural diversity, characterization and toxicology of microcystins. *Toxins* **2019**, *11*, 714. doi:10.3390/toxins11120714
40. Carmichael, W.W. Naming of cyclic heptapeptide toxins of cyanobacteria (blue-green algae). *Toxicon* **1988**, *26*, 971-973. doi:10.1016/0041-0101(88)90195-x
41. Díez-Quijada, L.; Prieto, A.I.; Guzmán-Guillén, R.; Jos, A.; Cameán, A.M. Occurrence and toxicity of microcystin congeners other than MC-LR and MC-RR: A review. *Food Chem. Toxicol.* **2019**, *125*, 106-132. doi:10.1016/j.fct.2018.12.042
42. MacKintosh, C.; Beattie, K.A.; Klumpp, S.; Cohen, P.; Codd, G.A. Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2A from both mammals and higher plants. *FEBS letters* **1990**, *264*, 187-192. doi:10.1016/0014-5793(90)80245-E
43. Altaner, S.; Jaeger, S.; Fotler, R.; Zemskov, I.; Wittmann, V.; Schreiber, F.; Dietrich, D.R. Machine learning prediction of cyanobacterial toxin (microcystin) toxicodynamics in humans. *ALTEX* **2020**, *37*. doi:10.14573/altex.1904031
44. Hu, Y.; Chen, J.; Fan, H.; Xie, P.; He, J. A review of neurotoxicity of microcystins. *Environ. Sci. Pollut. Res.* **2016**, *23*, 7211-7219. doi:10.1007/s11356-016-6073-y
45. Tzima, E.; Serifi, I.; Tsikari, I.; Alzualde, A.; Leonardos, I.; Papamarcaki, T. Transcriptional and behavioral responses of zebrafish larvae to microcystin-LR exposure. *Int. J. Mol. Sci.* **2017**, *18*, 365. doi:10.3390/ijms18020365
46. Pašková, V.; Adamovský, O.; Pikula, J.; Skočovská, B.; Band'ouchová, H.; Horáková, J.; Babica, P.; Maršálek, B.; Hilscherová, K. Detoxification and oxidative stress responses along with microcystins accumulation in Japanese quail exposed to cyanobacterial biomass. *Sci. Total Environ.* **2008**, *398*, 34-47. doi:10.1016/j.scitotenv.2008.03.001
47. Zhao, S.; Li, G.; Chen, J. A proteomic analysis of prenatal transfer of microcystin-LR induced neurotoxicity in rat offspring. *J. Proteomics* **2015**, *114*, 197-213. doi:10.1016/j.jprot.2014.11.015
48. Feurstein, D.; Stemmer, K.; Kleinteich, J.; Speicher, T.; Dietrich, D.R. Microcystin congener- and concentration-dependent induction of murine neuron apoptosis and neurite degeneration. *Toxicol. Sci.* **2011**, *124*, 424-431. doi:10.1093/toxsci/kfr243

49. Cox, P.A.; Banack, S.A.; Murch, S.J.; Rasmussen, U.; Tien, G.; Bidigare, R.R.; Metcalf, J.S.; Morrison, L.F.; Codd, G.A.; Bergman, B. Diverse taxa of cyanobacteria produce β -N-methylamino-L-alanine, a neurotoxic amino acid. *PNAS* **2005**, *102*, 5074-5078. doi:10.1073/pnas.0501526102
50. Violi, J.P.; Mitrovic, S.M.; Colville, A.; Main, B.J.; Rodgers, K.J. Prevalence of β -methylamino-L-alanine (BMAA) and its isomers in freshwater cyanobacteria isolated from eastern Australia. *Ecotoxicol. Environ. Saf.* **2019**, *172*, 72-81. doi:10.1016/j.ecoenv.2019.01.046
51. Schneider, T.; Simpson, C.; Desai, P.; Tucker, M.; Lobner, D. Neurotoxicity of isomers of the environmental toxin L-BMAA. *Toxicon* **2020**, *184*, 175-179. doi:10.1016/j.toxicon.2020.06.014
52. Martin, R.M.; Bereman, M.S.; Marsden, K.C. The cyanotoxin 2, 4-DAB reduces viability and causes behavioral and molecular dysfunctions associated with neurodegeneration in larval zebrafish. *Neurotox. Res.* **2022**, *40*, 347-364. doi:10.1007/s12640-021-00465-4
53. Nunes-Costa, D.; Magalhães, J.D.; G-Fernandes, M.; Cardoso, S.M.; Empadinhas, N. Microbial BMAA and the pathway for Parkinson's disease neurodegeneration. *Front. Aging Neurosci.* **2020**, *12*, 26. doi:10.3389/fnagi.2020.00026
54. Dunlop, R.A.; Cox, P.A.; Banack, S.A.; Rodgers, K.J. The non-protein amino acid BMAA is misincorporated into human proteins in place of L-serine causing protein misfolding and aggregation. *PloS one* **2013**, *8*, e75376. doi:10.1371/journal.pone.0075376
55. Frøyset, A.K.; Khan, E.A.; Fladmark, K.E. Quantitative proteomics analysis of zebrafish exposed to sub-lethal dosages of β -methyl-amino-L-alanine (BMAA). *Sci. Rep.* **2016**, *6*, 1-9. doi:10.1038/srep29631
56. Cox, P.A.; Davis, D.A.; Mash, D.C.; Metcalf, J.S.; Banack, S.A. Dietary exposure to an environmental toxin triggers neurofibrillary tangles and amyloid deposits in the brain. *Proc. Royal Soc. B.: Biol. Sci.* **2016**, *283*, 20152397. doi:10.1098/rspb.2015.2397
57. Haase, C.; Stieler, J.; Arendt, T.; Holzer, M. Pseudophosphorylation of tau protein alters its ability for self-aggregation. *J. Neurochem.* **2004**, *88*, 1509-1520. doi:10.1046/j.1471-4159.2003.02287.x
58. Lee, J.W.; Beebe, K.; Nangle, L.A.; Jang, J.; Longo-Guess, C.M.; Cook, S.A.; Davisson, M.T.; Sundberg, J.P.; Schimmel, P.; Ackerman, S.L. Editing-defective tRNA synthetase causes protein misfolding and neurodegeneration. *Nature* **2006**, *443*, 50-55. doi:10.1038/nature05096
59. De Munck, E.; Munoz-Saez, E.; Miguel, B.G.; Solas, M.T.; Ojeda, I.; Martinez, A.; Gil, C.; Arahuetes, R.M. β -N-methylamino-L-alanine causes neurological and pathological phenotypes mimicking Amyotrophic Lateral Sclerosis (ALS): the first step towards an experimental model for sporadic ALS. *Environ. Toxicol. Pharmacol.* **2013**, *36*, 243-255. doi:10.1016/j.etap.2013.04.007
60. Muñoz-Saez, E.; de Munck, E.; Arahuetes, R.M.; Solas, M.T.; Martínez, A.M.; Miguel, B.G. β -N-methylamino-L-alanine induces changes in both GSK3 and TDP-43 in human neuroblastoma. *J. Toxicol. Sci.* **2013**, *38*, 425-430. doi:10.2131/jts.38.425
61. Karlsson, O.; Jiang, L.; Ersson, L.; Malmström, T.; Ilag, L.L.; Brittebo, E.B. Environmental neurotoxin interaction with proteins: dose-dependent increase of free and protein-associated BMAA (β -N-methylamino-L-alanine) in neonatal rat brain. *Sci. Rep.* **2015a**, *5*, 1-9. doi:10.1038/srep15570
62. Xie, X.; Basile, M.; Mash, D.C. Cerebral uptake and protein incorporation of cyanobacterial toxin β -N-methylamino-L-alanine. *Neuroreport* **2013**, *24*, 779-784. doi: 10.1097/WNR.0b013e328363fd89
63. Han, N.-C.; Bullwinkle, T.J.; Loeb, K.F.; Faull, K.F.; Mohler, K.; Rinehart, J.; Ibba, M. The mechanism of β -N-methylamino-L-alanine inhibition of tRNA aminoacylation and its impact on misincorporation. *J. Biol. Chem.* **2020**, *295*, 1402-1410. doi: 10.1016/S0021-9258(17)49898-X
64. Rochet, J.-C. Errors in translation cause selective neurodegeneration. *ACS Chem. Biol.* **2006**, *1*, 562-566. doi:10.1021/cb6004068
65. Carmichael, W.W.; Biggs, D.F.; Peterson, M.A. Pharmacology of anatoxin-a, produced by the freshwater cyanophyte *Anabaena flos-aquae* NRC-44-1. *Toxicon* **1979**, *17*, 229-236. doi:10.1016/0041-0101(79)90212-5
66. Christensen, V.G.; Khan, E. Freshwater neurotoxins and concerns for human, animal, and ecosystem health: A review of anatoxin-a and saxitoxin. *Sci. Total Environ.* **2020**, *736*, 139515. doi:10.1016/j.scitotenv.2020.139515
67. Fiore, M.F.; de Lima, S.T.; Carmichael, W.W.; McKinnie, S.M.; Chekan, J.R.; Moore, B.S. Guanitoxin, re-naming a cyanobacterial organophosphate toxin. *Harmful Algae* **2020**, *92*, 101737. doi:10.1016/j.hal.2019.101737
68. Metcalf, J.; Richer, R.; Cox, P.; Codd, G. Cyanotoxins in desert environments may present a risk to human health. *Sci. Total Environ.* **2012**, *421*, 118-123. doi:10.1016/j.scitotenv.2012.01.053
69. Metcalf, J.; Banack, S.; Richer, R.; Cox, P. Neurotoxic amino acids and their isomers in desert environments. *J. Arid Environ.* **2015**, *112*, 140-144. doi:10.1016/j.jaridenv.2014.08.002
70. Ohtani, I.; Moore, R.E.; Runnegar, M.T. Cylindrospermopsin: a potent hepatotoxin from the blue-green alga *Cylindrospermopsis raciborskii*. *J. Amer. Chem. Soc.* **1992**, *114*, 7941-7942. doi:10.1021/ja00046a067
71. Rzymiski, P.; Poniedziałek, B. In search of environmental role of cylindrospermopsin: A review on global distribution and ecology of its producers. *Wat. Res.* **2014**, *66*, 320-337. doi:10.1016/j.watres.2014.08.029

72. Nowruzi, B.; Porzani, S.J. Toxic compounds produced by cyanobacteria belonging to several species of the order Nostocales: A review. *J. Appl. Toxicol.* **2021**, *41*, 510-548. doi:10.1002/jat.4088
73. Christensen, V.G.; Olds, H.T.; Norland, J.; Khan, E. Phytoplankton community interactions and cyanotoxin mixtures in three recurring surface blooms within one lake. *J. Hazard. Mater.* **2022**, *427*, 128142. doi:10.1016/j.jhazmat.2021.128142
74. Martin, R.M.; Stallrich, J.; Bereman, M.S. Mixture designs to investigate adverse effects upon co-exposure to environmental cyanotoxins. *Toxicology* **2019**, *421*, 74-83. doi:10.1016/j.tox.2019.04.013
75. Martin, R.M.; Bereman, M.S.; Marsden, K.C. BMAA and MCLR interact to modulate behavior and exacerbate molecular changes related to neurodegeneration in larval zebrafish. *Toxicol. Sci.* **2021**, *179*, 251-261. doi: 10.1093/toxsci/kfaa178
76. Roy-Lachapelle, A.; Duy, S.V.; Munoz, G.; Dinh, Q.T.; Bahl, E.; Simon, D.F.; Sauvé, S. Analysis of multiclass cyanotoxins (microcystins, anabaenopeptins, cylindrospermopsin and anatoxins) in lake waters using on-line SPE liquid chromatography high-resolution orbitrap mass spectrometry. *Anal. Methods* **2019**, *11*, 5289-5300. doi: 10.1039/C9AY01132C
77. Janssen, E.M.-L. Cyanobacterial peptides beyond microcystins—a review on co-occurrence, toxicity, and challenges for risk assessment. *Water Res.* **2019**, *151*, 488-499. doi:10.1016/j.watres.2018.12.048
78. Natumi, R.; Janssen, E.M.-L. Cyanopeptide co-production dynamics beyond microcystins and effects of growth stages and nutrient availability. *Environ. Sci. Technol.* **2020**, *54*, 6063-6072. doi:10.1021/acs.est.9b07334
79. Nagarajan, M.; Maruthanayagam, V.; Sundararaman, M. SAR analysis and bioactive potentials of freshwater and terrestrial cyanobacterial compounds: a review. *J. Appl. Toxicol.* **2013**, *33*, 313-349. doi:10.1002/jat.2833
80. Caban-Holt, A.; Mattingly, M.; Cooper, G.; Schmitt, F.A. Neurodegenerative memory disorders: a potential role of environmental toxins. *Neurol. Clin.* **2005**, *23*, 485-521. doi:10.1016/j.ncl.2004.12.005
81. Shaw, C.A.; Höglinger, G.U. Neurodegenerative diseases: neurotoxins as sufficient etiologic agents? *Neuromolecular Med.* **2008**, *10*, 1-9. doi: 10.1007/s12017-007-8016-8
82. Caller, T.A.; Doolin, J.W.; Haney, J.F.; Murby, A.J.; West, K.G.; Farrar, H.E.; Ball, A.; Harris, B.T.; Stommel, E.W. A cluster of amyotrophic lateral sclerosis in New Hampshire: a possible role for toxic cyanobacteria blooms. *Amyotroph. Lateral Scler.* **2009**, *10*, 101-108. doi:10.3109/17482960903278485
83. Banack, S.A.; Caller, T.; Henegan, P.; Haney, J.; Murby, A.; Metcalf, J.S.; Powell, J.; Alan, P.; Stommel, E. Detection of cyanotoxins, β -N-methylamino-L-alanine and microcystins, from a lake surrounded by cases of amyotrophic lateral sclerosis. *Toxins (Basel)* **2015**, *7*, 322-336, doi:10.3390/toxins7020322.
84. Torbick, N.; Ziniti, B.; Stommel, E.; Linder, E.; Andrew, A.; Caller, T.; Haney, J.; Bradley, W.; Henegan, P.L.; Shi, X. Assessing cyanobacterial harmful algal blooms as risk factors for amyotrophic lateral sclerosis. *Neurotox. Res.* **2018**, *33*, 199-212, doi:10.1007/s12640-017-9740-y
85. Stanhope, J.M.; Brody, J.A.; Morris, C.E. Epidemiologic features of amyotrophic lateral sclerosis and parkinsonism-dementia in Guam, Mariana Islands. *Int. J. Epidemiol.* **1972**, *1*, 199-210. doi:10.1093/ije/1.3.199
86. Cox, P.A.; Sacks, O.W. Cycad neurotoxins, consumption of flying foxes, and ALS-PDC disease in Guam. *Neurology* **2002**, *58*, 956-959, doi:10.1212/WNL.58.6.956.
87. Yanagihara, R.T.; Garruto, R.M.; Gajdusek, D.C. Epidemiological surveillance of amyotrophic lateral sclerosis and parkinsonism-dementia in the commonwealth of the Northern Mariana Islands. *Ann. Neurol.* **1983**, *13*, 79-86. doi:10.1002/ana.410130117
88. Cox, P.A.; Anne Banack, S.; Murch, S.J. Biomagnification of cyanobacterial neurotoxins and neurodegenerative disease among the Chamorro People of Guam. *PNAS* **2003**, *100*, 13380-13383. doi: 10.1073/pnas.2235808100
89. Murch, S.J.; Cox, P.A.; Banack, S.A.; Steele, J.C.; Sacks, O.W. Occurrence of β -methylamino-L-alanine (BMAA) in ALS/PDC patients from Guam. *Acta Neurol. Scand.* **2004a**, *110*, 267-269, doi:10.1111/j.1600-0404.2004.00320.x
90. Gajdusek, D.C. *Calcium deficiency induced secondary hyperparathyroidism and resultant CNS deposition of calcium and other metallic cations as the cause of ALS and PD in high incidence among the Auyu and Jakai people in west New Guinea*; National Institute of Neurological and Communicative Disorders and Stroke: 1984; pp. 145-171.
91. Kimura, K.; Yase, Y.; Higashi, Y.; Yamamoto, K.; Sugiura, M.; Tsumoto, T.; Uno, S.; Yoshimura, S.; Namikawa, K.; Kumura, J. Endemiological and geomedical studies on amyotrophic lateral sclerosis and allied diseases in Kii Peninsula, Japan (preliminary report). *Psychiatry Clin. Neurosci.* **1961**, *15*, 175-181. doi:10.1111/j.1440-1819.1961.tb00645.x
92. Yase, Y. The pathogenesis of amyotrophic lateral sclerosis. *The Lancet* **1972**, *300*, 292-296. doi:10.1016/S0140-6736(72)92903-0
93. Gunnarsson, L.-G.; Lygner, P.-E.; Veiga-Cabo, J.; de Pedro-Cuesta, J. An epidemic-like cluster of motor neuron disease in a Swedish county during the period 1973-1984. *Neuroepidemiology* **1996**, *15*, 142-152. doi:10.1159/000109901

94. Sienko, D.G.; Davis, J.P.; Taylor, J.A.; Brooks, B.R. Amyotrophic lateral sclerosis: a case-control study following detection of a cluster in a small Wisconsin community. *Arch. Neurol.* **1990**, *47*, 38-41. doi:10.1001/archneur.1990.00530010046017
95. Corcia, P.; Jafari-Schluep, H.-F.; Lardillier, D.; Mazyad, H.; Giraud, P.; Clavelou, P.; Pouget, J.; Camu, W. A clustering of conjugal amyotrophic lateral sclerosis in southeastern France. *Arch. Neurol.* **2003**, *60*, 553–557. doi:10.1001/archneur.60.4.553
96. Uccelli, R.; Binazzi, A.; Altavista, P.; Belli, S.; Comba, P.; Mastrantonio, M.; Vanacore, N. Geographic distribution of amyotrophic lateral sclerosis through motor neuron disease mortality data. *Eur. J. Epidemiol.* **2007**, *22*, 781–790, doi:10.1007/s10654-007-9173-7
97. Sabel, C.E.; Boyle, P.J.; Löytönen, M.; Gatrell, A.C.; Jokelainen, M.; Flowerdew, R.; Maasilta, P. Spatial clustering of amyotrophic lateral sclerosis in Finland at place of birth and place of death. *Am. J. Epidemiol.* **2003**, *157*, 898–905, doi:10.1093/aje/kwg090
98. Horner, R.D.; Grambow, S.C.; Coffman, C.J.; Lindquist, J.H.; Oddone, E.Z.; Allen, K.D.; Kasarskis, E.J. Amyotrophic lateral sclerosis among 1991 Gulf War veterans: evidence for a time-limited outbreak. *Neuroepidemiology* **2008**, *31*, 28–32, doi:10.1159/000136648
99. Miranda, M.L.; Alicia Overstreet Galeano, M.; Tassone, E.; Allen, K.D.; Horner, R.D. Spatial analysis of the etiology of amyotrophic lateral sclerosis among 1991 Gulf War veterans. *Neurotoxicology* **2008**, *29*, 964–970, doi:10.1016/j.neuro.2008.05.005
100. Stommel, E.W.; Field, N.C.; Caller, T.A. Aerosolization of cyanobacteria as a risk factor for amyotrophic lateral sclerosis. *Med. Hypotheses* **2013**, *80*, 142–145, doi:10.1016/j.mehy.2012.11.012
101. Masseret, E.; Banack, S.; Boumédiène, F.; Abadie, E.; Briant, L.; Pernet, F.; Juntas-Morales, R.; Pageot, N.; Metcalf, J.; Cox, P. Dietary BMAA exposure in an amyotrophic lateral sclerosis cluster from southern France. *PloS one* **2013**, *8*, e83406. doi: 10.1371/journal.pone.0083406
102. Caller, T.A.; Chipman, J.W.; Field, N.C.; Stommel, E.W. Spatial analysis of amyotrophic lateral sclerosis in Northern New England, USA, 1997-2009. *Muscle Nerve* **2013**, *48*, 235–241, doi:10.1002/mus.23761.
103. Torbick, N.; Hession, S.; Stommel, E.; Caller, T. Mapping amyotrophic lateral sclerosis lake risk factors across northern New England. *Int. J. Health Geogr.* **2014**, *13*, 1-15. doi:10.1186/1476-072X-13-1
104. Delzor, A.; Couratier, P.; Boumédiène, F.; Nicol, M.; Druet-Cabanac, M.; Paraf, F.; Méjean, A.; Ploux, O.; Leleu, J.-P.; Briant, L. Searching for a link between the L-BMAA neurotoxin and amyotrophic lateral sclerosis: a study protocol of the French BMAA-LS programme. *BMJ open* **2014**, *4*, e005528. doi:10.1136/bmjopen-2014-005528
105. Lee, S.; Choi, B.; Kim, S.J.; Kim, J.; Kang, D.; Lee, J. Relationship between freshwater harmful algal blooms and neurodegenerative disease incidence rates in South Korea. *Environ. Health* **2022**, *21*, 1-11. doi:10.1186/s12940-022-00935-y
106. Lannuzel, A.; Mecharles, S.; Tressières, B.; Demoly, A.; Alhend, R.; Hédreille-Tablon, M.-A.; Alecu, C. Clinical varieties and epidemiological aspects of amyotrophic lateral sclerosis in the Caribbean island of Guadeloupe: A new focus of ALS associated with Parkinsonism. *Amyotroph. Lateral. Scler. Frontotemporal Degener.* **2015**, *16*, 216-223, doi:10.3109/21678421.2014.992026
107. Filippini, T.; Fiore, M.; Tesauro, M.; Malagoli, C.; Consonni, M.; Violi, F.; Arcolin, E.; Iacuzio, L.; Conti, G.O.; Cristaldi, A.; et al. Clinical and lifestyle factors and risk of amyotrophic lateral sclerosis: a population-based case-control study. *Int. J. Environ. Res. Public Health* **2020**, *17*, doi:10.3390/ijerph17030857
108. Field, N.C.; Metcalf, J.S.; Caller, T.A.; Banack, S.A.; Cox, P.A.; Stommel, E.W. Linking β -methylamino-L-alanine exposure to sporadic amyotrophic lateral sclerosis in Annapolis, MD. *Toxicon* **2013**, *70*, 179-183. doi:10.1016/j.toxicon.2013.04.010
109. Berntzon, L.; Ronnevi, L.; Bergman, B.; Eriksson, J. Detection of BMAA in the human central nervous system. *Neuroscience* **2015**, *292*, 137-147. doi:10.1016/j.neuroscience.2015.02.032
110. Pablo, J.; Banack, S.; Cox, P.; Johnson, T.; Papapetropoulos, S.; Bradley, W.; Buck, A.; Mash, D. Cyanobacterial neurotoxin BMAA in ALS and Alzheimer's disease. *Acta Psychiatr. Scand.* **2009**, *120*, 216-225. doi:10.1111/j.1600-0404.2008.01150.x
111. Montine, T.J.; Li, K.; Perl, D.P.; Galasko, D. Lack of β -methylamino-L-alanine in brain from controls, AD, or Chamorro with PDC. *Neurology* **2005**, *65*, 768-769. doi:10.1212/01.wnl.0000174523.62022.52
112. Meneely, J.P.; Chevallier, O.P.; Graham, S.; Greer, B.; Green, B.D.; Elliott, C.T. β -methylamino-L-alanine (BMAA) is not found in the brains of patients with confirmed Alzheimer's disease. *Sci. Rep.* **2016**, *6*, 1-9. doi:10.1038/srep36363
113. Spencer, P.S.; Palmer, V.S.; Kisby, G.E. Western Pacific ALS-PDC: Evidence implicating cycad genotoxins. *J. Neurol. Sci.* **2020**, *419*, 117185. doi:10.1016/j.jns.2020.117185
114. Jonasson, S.; Eriksson, J.; Berntzon, L.; Spáčil, Z.; Ilag, L.L.; Ronnevi, L.-O.; Rasmussen, U.; Bergman, B. Transfer of a cyanobacterial neurotoxin within a temperate aquatic ecosystem suggests pathways for human exposure. *PNAS* **2010**, *107*, 9252-9257. doi:10.1073/pnas.0914417107

115. Jiang, L.; Eriksson, J.; Lage, S.; Jonasson, S.; Shams, S.; Mehine, M.; Ilag, L.L.; Rasmussen, U. Diatoms: a novel source for the neurotoxin BMAA in aquatic environments. *PloS one* **2014a**, *9*, e84578. doi: 10.1371/journal.pone.0084578
116. Johnson, H.E.; King, S.R.; Banack, S.A.; Webster, C.; Callanaupa, W.J.; Cox, P.A. Cyanobacteria (*Nostoc commune*) used as a dietary item in the Peruvian highlands produce the neurotoxic amino acid BMAA. *J.Ethnopharmacol.* **2008**, *118*, 159-165. doi:10.1016/j.jep.2008.04.008
117. Cox, P.A.; Richer, R.; Metcalf, J.S.; Banack, S.A.; Codd, G.A.; Bradley, W.G. Cyanobacteria and BMAA exposure from desert dust: a possible link to sporadic ALS among Gulf War veterans. *Amyotroph. Lateral Scler.* **2009**, *10*, 109-117. doi:10.3109/17482960903286066
118. Jiang, L.; Kiselova, N.; Rosén, J.; Ilag, L.L. Quantification of neurotoxin BMAA (β -N-methylamino-L-alanine) in seafood from Swedish markets. *Sci. Rep.* **2014b**, *4*, 1-7. doi:10.1038/srep06931
119. Lage, S.; Costa, P.R.; Moita, T.; Eriksson, J.; Rasmussen, U.; Rydberg, S.J. BMAA in shellfish from two Portuguese transitional water bodies suggests the marine dinoflagellate *Gymnodinium catenatum* as a potential BMAA source. *Aquat. Toxicol.* **2014**, *152*, 131-138. doi:10.1016/j.aquatox.2014.03.029
120. Bradley, W.G.; Miller, R.; Levine, T.; Stommel, E.; Cox, P. Studies of environmental risk factors in amyotrophic lateral sclerosis (ALS) and a phase I clinical trial of L-serine. *Neurotox. Res.* **2018**, *33*, 192-198. doi:10.1007/s12640-017-9741-x
121. Saitou, T.; Sugiura, N.; Itayama, T.; Inamori, Y.; Matsumura, M. Degradation characteristics of microcystins by isolated bacteria from Lake Kasumigaura. *J. Water Supply: Res. Technol. AQUA* **2003**, *52*, 13-18. doi:10.2166/aqua.2003.0002
122. Edwards, C.; Graham, D.; Fowler, N.; Lawton, L.A. Biodegradation of microcystins and nodularin in freshwaters. *Chemosphere* **2008**, *73*, 1315-1321. doi:10.1016/j.chemosphere.2008.07.015
123. Gible, C.M.; Peacock, M.B.; Kudela, R.M. Evidence of freshwater algal toxins in marine shellfish: implications for human and aquatic health. *Harmful Algae* **2016**, *59*, 59-66. doi:10.1016/j.hal.2016.09.007
124. Chiswell, R.K.; Shaw, G.R.; Eaglesham, G.; Smith, M.J.; Norris, R.L.; Seawright, A.A.; Moore, M.R. Stability of cylindrospermopsin, the toxin from the cyanobacterium, *Cylindrospermopsis raciborskii*: Effect of pH, temperature, and sunlight on decomposition. *Environ. Toxicol.* **1999**, *14*, 155-161. doi:10.1002/(SICI)1522-7278(199902)14:1<155::AID-TOX20>3.0.CO;2-Z
125. Adamski, M.; Żmudski, P.; Chrapusta, E.; Bober, B.; Kaminski, A.; Zabaglo, K.; Latkowska, E.; Bialczyk, J. Effect of pH and temperature on the stability of cylindrospermopsin. Characterization of decomposition products. *Algal Res.* **2016**, *15*, 129-134. doi:10.1016/j.algal.2016.02.020
126. Sedmak, B.; Carmeli, S.; Eleršek, T. "Non-toxic" cyclic peptides induce lysis of cyanobacteria – an effective cell population density control mechanism in cyanobacterial blooms. *Microb. Ecol.* **2008**, *56*, 201-209. doi:10.1007/s00248-007-9336-9
127. Sedmak, B.; Sukenik, A.; Eleršek, T.; Kosi, G. The biological role of cyclic hepatotoxic and non-hepatotoxic cyanopeptides and its ecological consequences. In *Ecotoxicology research developments*; Santos, Eduardo B. Ed.; Nova Science Pub Inc; UK, 2009; pp. 169-300.
128. Bionda, N. Cyclic lipodepsipeptides as lead structures for the discovery of new antibiotics. Ph.D., Florida Atlantic University, Ann Arbor, 2013.
129. Kurtz, T.; Zeng, T.; Rosario-Ortiz, F.L. Photodegradation of cyanotoxins in surface waters. *Water Res.* **2021**, *192*, 116804. doi:10.1016/j.watres.2021.116804
130. Li, H.; Xing, P.; Wu, Q.L. The high resilience of the bacterioplankton community in the face of a catastrophic disturbance by a heavy *Microcystis* bloom. *FEMS Microbiol. Ecol.* **2012**, *82*, 192-201. doi:10.1111/j.1574-6941.2012.01417.x
131. Kormas, K.A.; Lymperopoulou, D.S. Cyanobacterial toxin degrading bacteria: who are they? *Biomed Res. Int.* **2013**, *2013*, 463894. doi:10.1155/2013/463894
132. Mazur-Marzec, H.; Plinski, M. Do toxic cyanobacteria blooms pose a threat to the Baltic ecosystem? *Oceanologia* **2009**, *51*, 293-319. doi:10.5697/oc.51-3.293
133. Stolz, A. Molecular characteristics of xenobiotic-degrading sphingomonads. *Appl. Microbiol. Biotechnol.* **2009**, *81*, 793-811. doi:10.1007/s00253-008-1752-3
134. Kato, H.; Imanishi, S.Y.; Tsuji, K.; Harada, K.-i. Microbial degradation of cyanobacterial cyclic peptides. *Water Res.* **2007**, *41*, 1754-1762. doi:10.1016/j.watres.2007.01.003
135. Okano, K.; Shimizu, K.; Maseda, H.; Kawauchi, Y.; Utsumi, M.; Itayama, T.; Zhang, Z.; Sugiura, N. Whole-genome sequence of the microcystin-degrading bacterium *Sphingopyxis* sp. strain C-1. *Genome Announc.* **2015**, *3*, e00838-00815. doi:10.1128/genomeA.00838-15
136. Qin, L.; Zhang, X.; Chen, X.; Wang, K.; Shen, Y.; Li, D. Isolation of a novel microcystin-degrading bacterium and the evolutionary origin of *mlr* gene cluster. *Toxins* **2019**, *11*, 269. doi:10.3390/toxins11050269
137. Imanishi, S.; Kato, H.; Mizuno, M.; Tsuji, K.; Harada, K.-i. Bacterial degradation of microcystins and nodularin. *Chem. Res. Toxicol.* **2005**, *18*, 591-598. doi:10.1021/tx049677g

138. Maghsoudi, E.; Fortin, N.; Greer, C.; Maynard, C.; Pagé, A.; Duy, S.V.; Sauvé, S.; Prévost, M.; Dorner, S. Cyanotoxin degradation activity and *mlr* gene expression profiles of a *Sphingopyxis* sp. isolated from Lake Champlain, Canada. *Environ. Sci.: Process. Impacts*. **2016**, *18*, 1417-1426. doi: 10.1039/C6EM00001K
139. Mohamed, Z.A.; Alamri, S.A. Biodegradation of cylindrospermopsin toxin by microcystin-degrading bacteria isolated from cyanobacterial blooms. *Toxicon* **2012**, *60*, 1390-1395. doi:10.1016/j.toxicon.2012.10.004
140. Dziga, D.; Kokocinski, M.; Maksylewicz, A.; Czaja-Prokop, U.; Barylski, J. Cylindrospermopsin biodegradation abilities of *Aeromonas* sp. isolated from Rusałka Lake. *Toxins* **2016**, *8*, 55. doi:10.3390/toxins8030055
141. Nybom, S.M.; Salminen, S.J.; Meriluoto, J.A. Specific strains of probiotic bacteria are efficient in removal of several different cyanobacterial toxins from solution. *Toxicon* **2008**, *52*, 214-220. doi: 10.1016/j.toxicon.2008.04.169
142. Mohamed, Z.A.; Hashem, M.; Alamri, S.A. Growth inhibition of the cyanobacterium *Microcystis aeruginosa* and degradation of its microcystin toxins by the fungus *Trichoderma citrinoviride*. *Toxicon* **2014**, *86*, 51-58. doi:10.1016/j.toxicon.2014.05.008
143. Mohamed, Z.A.; Hashem, M.; Alamri, S.; Campos, A.; Vasconcelos, V. Fungal biodegradation and removal of cyanobacteria and microcystins: potential applications and research needs. *Environ. Sci. Pollut. Res.* **2021**, *28*, 37041-37050. doi: 10.1007/s11356-021-14623-w
144. Kurmayer, R.; Jüttner, F. Strategies for the co-existence of zooplankton with the toxic cyanobacterium *Planktothrix rubescens* in Lake Zurich. *Journal of Plankton Research* **1999**, *21*. doi:10.1093/plankt/21.4.659
145. Akbar, S.; Huang, J.; Zhou, Q.; Gu, L.; Sun, Y.; Zhang, L.; Lyu, K.; Yang, Z. Elevated temperature and toxic *Microcystis* reduce *Daphnia* fitness and modulate gut microbiota. *Environ. Pollut.* **2021**, *271*, 116409. doi: 10.1016/j.envpol.2020.116409
146. Combes, A.; Dellinger, M.; Cadel-six, S.; Amand, S.; Comte, K. Ciliate *Nassula* sp. grazing on a microcystin-producing cyanobacterium (*Planktothrix agardhii*): impact on cell growth and in the microcystin fractions. *Aquatic toxicol.* **2013**, *126*, 435-441. doi:10.1016/j.aquatox.2012.08.018
147. Zhang, L.; Wang, Z.; Wang, N.; Gu, L.; Sun, Y.; Huang, Y.; Chen, Y.; Yang, Z. Mixotrophic *Ochromonas* addition improves the harmful *Microcystis*-dominated phytoplankton community in *in situ* microcosms. *Environ. Sci. Tech.* **2020**, *54*, 4609-4620. doi:10.1021/acs.est.9b06438
148. Duncan, M.W. Good mass spectrometry and its place in good science. *J. Mass Spectrom.* **2012**, *47*, 795-809. doi:10.1002/jms.3038
149. Faassen, E.J.; Antoniou, M.G.; Beekman-Lukassen, W.; Blahova, L.; Chernova, E.; Christophoridis, C.; Combes, A.; Edwards, C.; Fastner, J.; Harmsen, J. A collaborative evaluation of LC-MS/MS based methods for BMAA analysis: Soluble bound BMAA found to be an important fraction. *Marine drugs* **2016**, *14*, 45. doi:10.3390/md14030045
150. Van Apeldoorn, M.E.; Van Egmond, H.P.; Speijers, G.J.; Bakker, G.J. Toxins of cyanobacteria. *Mol. Nutr. Food Res.* **2007**, *51*, 7-60. doi:10.1002/mnfr.200600185
151. Testai, E.; Buratti, F.M.; Funari, E.; Manganelli, M.; Vichi, S.; Arnich, N.; Biré, R.; Fessard, V.; Sialehaamo, A. Review and analysis of occurrence, exposure and toxicity of cyanobacteria toxins in food. *EFSA Support. Publ.* **2016**, *13*, 998E. doi:10.2903/sp.efsa.2016.EN-998
152. Merel, S.; Walker, D.; Chicana, R.; Snyder, S.; Baurès, E.; Thomas, O. State of knowledge and concerns on cyanobacterial blooms and cyanotoxins. *Environ. Int.* **2013**, *59*, 303-327. doi:10.1016/j.envint.2013.06.013
153. Abbes, S.; Vo Duy, S.; Munoz, G.; Dinh, Q.T.; Simon, D.F.; Husk, B.; Baulch, H.M.; Vinçon-Leite, B.; Fortin, N.; Greer, C.W. Occurrence of BMAA isomers in bloom-impacted lakes and reservoirs of Brazil, Canada, France, Mexico, and the United Kingdom. *Toxins* **2022**, *14*, 251. doi:10.3390/toxins14040251
154. Altenburger, R.; Ait-Aissa, S.; Antczak, P.; Backhaus, T.; Barceló, D.; Seiler, T.-B.; Brion, F.; Busch, W.; Chipman, K.; de Alda, M.L. Future water quality monitoring—adapting tools to deal with mixtures of pollutants in water resource management. *Sci. Total Environ.* **2015**, *512*, 540-551. doi:10.1016/j.scitotenv.2014.12.057
155. Hollender, J.; Schymanski, E.L.; Singer, H.P.; Ferguson, P.L. Nontarget screening with high resolution mass spectrometry in the environment: ready to go? *Environ. Sci. Technol.* **2017**, *51*, 11505-11512. doi:10.1021/acs.est.7b02184
156. Escher, B.I.; Stapleton, H.M.; Schymanski, E.L. Tracking complex mixtures of chemicals in our changing environment. *Science* **2020**, *367*, 388-392. doi:10.1126/science.aay6636
157. Dom, I.; Biré, R.; Hort, V.; Lavison-Bompard, G.; Nicolas, M.; Guérin, T. Extended targeted and non-targeted strategies for the analysis of marine toxins in mussels and oysters by (LC-HRMS). *Toxins* **2018**, *10*, 375. doi:10.3390/toxins10090375
158. Krauss, M.; Singer, H.; Hollender, J. LC-high resolution MS in environmental analysis: from target screening to the identification of unknowns. *Anal. Bioanal. Chem.* **2010**, *397*, 943-951. doi:10.1016/j.hal.2012.04.002

159. Bogialli, S.; Bortolini, C.; Di Gangi, I.M.; Di Gregorio, F.N.; Lucentini, L.; Favaro, G.; Pastore, P. Liquid chromatography-high resolution mass spectrometric methods for the surveillance monitoring of cyanotoxins in freshwaters. *Talanta* **2017**, *170*, 322-330. doi:10.1016/j.talanta.2017.04.033
160. Gonzalez-Gaya, B.; Lopez-Herguedas, N.; Bilbao, D.; Mijangos, L.; Iker, A.; Etxebarria, N.; Irazola, M.; Prieto, A.; Olivares, M.; Zuloaga, O. Suspect and non-target screening: the last frontier in environmental analysis. *Anal. Methods* **2021**, *13*, 1876-1904. doi:10.1039/D1AY00111F
161. Picardo, M.; Sanchís, J.; Núñez, O.; Farré, M. Suspect screening of natural toxins in surface and drinking water by high performance liquid chromatography and high-resolution mass spectrometry. *Chemosphere* **2020**, *261*, 127888. doi:10.1016/j.chemosphere.2020.127888
162. He, Q.; Wang, W.; Xu, Q.; Liu, Z.; Teng, J.; Yan, H.; Liu, X. Microcystins in water: detection, microbial degradation strategies, and mechanisms. *Int. J. Environ. Res. Public Health*. **2022**, *19*, 13175. doi:10.3390/ijerph192013175
163. Sanseverino, I.; António, D.C.; Loos, R.; Lettieri, T. *Cyanotoxins: methods and approaches for their analysis and detection*; Publications Office of the European Union: Luxembourg, 2017; pp. doi:10.2760/36186
164. Van Hassel, W.H.R.; Huybrechts, B.; Masquelier, J.; Wilmotte, A.; Andjelkovic, M. Development, validation and application of a targeted LC-MS method for quantification of microcystins and nodularin: towards a better characterization of drinking water. *Water* **2022a**, *14*, doi:10.3390/w14081195
165. Panda, D.; Dash, B.P.; Manickam, S.; Boczkaj, G. Recent advancements in LC-MS based analysis of biotoxins: Present and future challenges. *Mass Spectrom. Rev.* **2022**, *41*, 766-803. doi:10.1002/mas.21689
166. Triantis, T.; Tsimeli, K.; Kaloudis, T.; Thanassoulas, N.; Lytras, E.; Hiskia, A. Development of an integrated laboratory system for the monitoring of cyanotoxins in surface and drinking waters. *Toxicon* **2010**, *55*, 979-989. doi:10.1016/j.toxicon.2009.07.012
167. Zhang, L.; Liu, J.; Zhang, D.; Luo, L.; Liao, Q.; Yuan, L.; Wu, N. Seasonal and spatial variations of microcystins in Poyang Lake, the largest freshwater lake in China. *Environ Sci Pollut Res Int* **2018**, *25*, 6300-6307, doi:10.1007/s11356-017-0967-1
168. Turner, A.D.; Dhanji-Rapkova, M.; O'Neill, A.; Coates, L.; Lewis, A.; Lewis, K. Analysis of microcystins in cyanobacterial blooms from freshwater bodies in England. *Toxins (Basel)* **2018a**, *10*, doi:10.3390/toxins10010039
169. Mantzouki, E.; Lurling, M.; Fastner, J.; de Senerpont Domis, L.; Wilk-Wozniak, E.; Koreiviene, J.; Seelen, L.; Teurlinx, S.; Verstijnen, Y.; Krzton, W.; et al. Temperature effects explain continental scale distribution of cyanobacterial toxins. *Toxins (Basel)* **2018**, *10*, doi:10.3390/toxins10040156
170. Main, B.J.; Bowling, L.C.; Padula, M.P.; Bishop, D.P.; Mitrovic, S.M.; Guillemin, G.J.; Rodgers, K.J. Detection of the suspected neurotoxin beta-methylamino-L-alanine (BMAA) in cyanobacterial blooms from multiple water bodies in Eastern Australia. *Harmful Algae* **2018**, *74*, 10-18, doi:10.1016/j.hal.2018.03.004
171. Pitois, F.; Fastner, J.; Pagotto, C.; Dechesne, M. Multi-toxin occurrences in ten french water resource reservoirs. *Toxins (Basel)* **2018**, *10*, 283. doi:10.3390/toxins10070283
172. Turner, A.D.; Waack, J.; Lewis, A.; Edwards, C.; Lawton, L. Development and single-laboratory validation of a UHPLC-MS/MS method for quantitation of microcystins and nodularin in natural water, cyanobacteria, shellfish and algal supplement tablet powders. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **2018b**, *1074-1075*, 111-123, doi:10.1016/j.jchromb.2017.12.032
173. Thuret-Benoist, H.; Pallier, V.; Feuillade-Cathalifaud, G. Quantification of microcystins in natural waters by HPLC-UV after a pre-concentration step: validation of the analytical performances and study of the interferences. *Environ. Toxicol. Pharmacol.* **2019**, *72*, 103223, doi:10.1016/j.etap.2019.103223
174. Botha, C.J.; Laver, P.; Singo, A.; Venter, E.; Ferreira, G.C.H.; Rösemann, M.; Myburgh, J.G. Evaluation of a Norwegian-developed ELISA to determine microcystin concentrations in fresh water. *Water Supply* **2019**, *19*, 743-752. doi:10.2166/ws.2018.118
175. Haddad, S.P.; Bobbitt, J.M.; Taylor, R.B.; Lovin, L.M.; Conkle, J.L.; Chambliss, C.K.; Brooks, B.W. Determination of microcystins, nodularin, anatoxin-a, cylindrospermopsin, and saxitoxin in water and fish tissue using isotope dilution liquid chromatography tandem mass spectrometry. *J. Chromatogr. A* **2019**, *1599*, 66-74, doi:10.1016/j.chroma.2019.03.066
176. Leon, C.; Penuela, G.A. Detected cyanotoxins by UHPLC MS/MS technique in tropical reservoirs of northeastern Colombia. *Toxicon* **2019**, *167*, 38-48, doi:10.1016/j.toxicon.2019.06.010
177. Vo Duy, S.; Munoz, G.; Dinh, Q.T.; Tien Do, D.; Simon, D.F.; Sauve, S. Analysis of the neurotoxin beta-N-methylamino-L-alanine (BMAA) and isomers in surface water by FMOC derivatization liquid chromatography high resolution mass spectrometry. *PLoS One* **2019**, *14*, e0220698, doi:10.1371/journal.pone.0220698
178. McKindles, K.M.; Zimba, P.V.; Chiu, A.S.; Watson, S.B.; Gutierrez, D.B.; Westrick, J.; Kling, H.; Davis, T.W. A multiplex analysis of potentially toxic cyanobacteria in Lake Winnipeg during the 2013 bloom season. *Toxins (Basel)* **2019**, *11*, doi:10.3390/toxins11100587

179. Yan, B.; Liu, Z.; Huang, R.; Xu, Y.; Liu, D.; Wang, W.; Zhao, Z.; Cui, F.; Shi, W. Impact factors on the production of beta-methylamino-L-alanine (BMAA) by cyanobacteria. *Chemosphere* **2020**, *243*, 125355, doi:10.1016/j.chemosphere.2019.125355
180. Abbas, F.; Porojan, C.; Mowe, M.A.D.; Lehane, M.; Mitrovic, S.M.; Lim, R.P.; Yeo, D.C.J.; Furey, A. Sample extraction and liquid chromatography–tandem mass spectrometry (LC-MS/MS) method development and validation for the quantitative detection of cyanobacterial hepatotoxins and neurotoxins in Singapore's reservoirs. *Mar. Freshw. Res.* **2020**, *71*, doi:10.1071/mf19157
181. Stoyneva-Gärtner, M.; Stefanova, K.; Descy, J.-P.; Uzunov, B.; Radkova, M.; Pavlova, V.; Mitreva, M.; Gärtner, G. *Microcystis aeruginosa* and *M. wesenbergii* were the primary planktonic microcystin producers in several Bulgarian waterbodies (August 2019). *Appl. Sci.* **2020**, *11*, doi:10.3390/app11010357
182. Zervou, S.K.; Moschandrea, K.; Paraskevopoulou, A.; Christophoridis, C.; Grigoriadou, E.; Kaloudis, T.; Triantis, T.M.; Tsiaoussi, V.; Hiskia, A. Cyanobacterial toxins and peptides in Lake Vegoritis, Greece. *Toxins (Basel)* **2021**, *13*, doi:10.3390/toxins13060394
183. Zhang, Y.; Husk, B.R.; Duy, S.V.; Dinh, Q.T.; Sanchez, J.S.; Sauve, S.; Whalen, J.K. Quantitative screening for cyanotoxins in soil and groundwater of agricultural watersheds in Quebec, Canada. *Chemosphere* **2021**, *274*, 129781, doi:10.1016/j.chemosphere.2021.129781
184. Choi, J.-W.; Jang, J.-H.; Kim, A.-K.; Lee, S.-H. Determination of β -N-Methylamino-L-Alanine (BMAA) levels in surface water and treated water by liquid chromatography-tandem mass spectrometry. *J. Environ. Anal. Health Toxicol.* **2021**, *24*, 171-175, doi:10.36278/jeht.24.4.171
185. Dinh, Q.T.; Munoz, G.; Simon, D.F.; Vo Duy, S.; Husk, B.; Sauve, S. Stability issues of microcystins, anabaenopeptins, anatoxins, and cylindrospermopsin during short-term and long-term storage of surface water and drinking water samples. *Harmful Algae* **2021**, *101*, 101955, doi:10.1016/j.hal.2020.101955
186. Wang, T.; Xie, H.; Cao, Y.; Xu, Q.; Gan, N. Magnetic solid phase extraction coupled with high-performance liquid chromatography-diode array detection based on assembled magnetic covalent organic frameworks for selective extraction and detection of microcystins in aquatic foods. *J. Chromatogr. A* **2022**, *1685*, 463614, doi:10.1016/j.chroma.2022.463614
187. Pan, L.; Huang, J.J.; Chen, J.; He, X.; Wang, Y.; Wang, J.; Wang, B. Trace determination of multiple hydrophilic cyanotoxins in freshwater by off- and on-line solid phase extraction coupled to liquid chromatography-tandem mass spectrometry. *Sci. Total Environ.* **2022**, *853*, 158545, doi:10.1016/j.scitotenv.2022.158545
188. Van Hassel, W.H.R.; Andjelkovic, M.; Durieu, B.; Marroquin, V.A.; Masquelier, J.; Huybrechts, B.; Wilmotte, A. A summer of cyanobacterial blooms in Belgian waterbodies: microcystin quantification and molecular characterizations. *Toxins (Basel)* **2022b**, *14*, doi:10.3390/toxins14010061
189. Choi, J.-w.; Jang, J.-h.; Lee, S.-h.; Yoon, M.-a. Determination of cylindrospermopsin in surface and treated water using liquid chromatography-tandem mass spectrometry. *Environ. Anal. Health. Toxicol.* **2022**, *25*, 71-76. doi:10.36278/jeht.25.2.71
190. Vo Duy, S.; Munoz, G.; Dinh, Q.T.; Zhang, Y.; Simon, D.F.; Sauve, S. Fast screening of saxitoxin, neosaxitoxin, and decarbamoyl analogues in fresh and brackish surface waters by on-line enrichment coupled to HILIC-HRMS. *Talanta* **2022**, *241*, 123267, doi:10.1016/j.talanta.2022.123267
191. Aparicio-Muriana, M.M.; Carmona-Molero, R.; Lara, F.J.; Garcia-Campana, A.M.; Del Olmo-Iruela, M. Multiclass cyanotoxin analysis in reservoir waters: Tandem solid-phase extraction followed by zwitterionic hydrophilic interaction liquid chromatography-mass spectrometry. *Talanta* **2022**, *237*, 122929, doi:10.1016/j.talanta.2021.122929.
192. Zhao, P.; Qiu, J.; Li, A.; Yan, G.; Li, M.; Ji, Y. Matrix Effect of Diverse Biological Samples Extracted with Different Extraction Ratios on the Detection of beta-N-Methylamino-L-Alanine by Two Common LC-MS/MS Analysis Methods. *Toxins (Basel)* **2022**, *14*, doi:10.3390/toxins14060387
193. Emmons, R.V.; Karaj, E.; Cudjoe, E.; Bell, D.S.; Tillekeratne, L.M.V.; Gionfriddo, E. Leveraging multi-mode microextraction and liquid chromatography stationary phases for quantitative analysis of neurotoxin beta-N-methylamino-L-alanine and other non-proteinogenic amino acids. *J. Chromatogr. A* **2022**, *1685*, 463636, doi:10.1016/j.chroma.2022.463636
194. Zhang, H.; Li, Y.; Abdallah, M.F.; Tan, H.; Li, J.; Liu, S.; Zhang, R.; Sun, F.; Li, Y.; Yang, S. Novel one-point calibration strategy for high-throughput quantitation of microcystins in freshwater using LC-MS/MS. *Sci Total Environ* **2023**, *858*, 159345, doi:10.1016/j.scitotenv.2022.159345
195. Kulabhusan, P.K.; Campbell, K. Recent trends in the detection of freshwater cyanotoxins with a critical note on their occurrence in Asia. *Trends Environ. Anal. Chem.* **2021**, *32*, e00150. doi:10.1016/j.teac.2021.e00150
196. Chorus, I.; Bartram, J. *Toxic Cyanobacteria in Water: A Guide to their Public Health Consequences, Monitoring and Management*, 1st ed.; Chorus, I., Bartram, J., Eds.; CRC Press: London, United Kingdom, 1999; pp. 440. doi:10.1201/9781482295061
197. Lee, J.; Lee, S.; Jiang, X. Cyanobacterial toxins in freshwater and food: important sources of exposure to humans. *Annu. Rev. Food Sci. Technol.* **2017**, *8*, 281-304. doi:10.1146/annurev-food-030216-030116

198. Tatters, A.O.; Howard, M.D.; Nagoda, C.; Busse, L.; Gellene, A.G.; Caron, D.A. Multiple stressors at the land-sea interface: Cyanotoxins at the land-sea interface in the Southern California Bight. *Toxins* **2017**, *9*, 95. doi:10.3390/toxins9030095
199. Tatters, A.O.; Howard, M.D.; Nagoda, C.; Fetscher, A.E.; Kudela, R.M.; Caron, D.A. Heterogeneity of toxin-producing cyanobacteria and cyanotoxins in coastal watersheds of southern California. *Estuaries Coasts* **2019**, *42*, 958-975. doi : doi:10.1007/s12237-019-00546-w
200. Tatters, A.O.; Smith, J.; Kudela, R.M.; Hayashi, K.; Howard, M.D.; Donovan, A.R.; Loftin, K.A.; Caron, D.A. The tide turns: Episodic and localized cross-contamination of a California coastline with cyanotoxins. *Harmful Algae* **2021**, *103*, 102003. doi:10.1016/j.hal.2021.102003
201. Howard, M.D.; Smith, J.; Caron, D.A.; Kudela, R.M.; Loftin, K.; Hayashi, K.; Fadness, R.; Fricke, S.; Kann, J.; Roethler, M. Integrative monitoring strategy for marine and freshwater harmful algal blooms and toxins across the freshwater-to-marine continuum. *Integr. Environ. Assess. Manag.* **2022**, *1*, 1-19. doi:10.1002/ieam.4651
202. Andraka, D.; Ospanov, K.; Myrzakhmetov, M. Current state of communal sewage treatment in the republic of Kazakhstan. *J. Ecol. Eng.* **2015**, *16*. doi:10.12911/22998993/60463
203. Gutiérrez-Praena, D.; Jos, Á.; Pichardo, S.; Moreno, I.M.; Cameán, A.M. Presence and bioaccumulation of microcystins and cylindrospermopsin in food and the effectiveness of some cooking techniques at decreasing their concentrations: A review. *Food Chem. Toxicol.* **2013**, *53*, 139-152. doi:10.1016/j.fct.2012.10.062
204. Crush, J.; Briggs, L.; Sprosen, J.; Nichols, S. Effect of irrigation with lake water containing microcystins on microcystin content and growth of ryegrass, clover, rape, and lettuce. *Environ. Toxicol.* **2008**, *23*, 246-252. doi:10.1002/tox.20331
205. Mohamed, Z.A.; Al Shehri, A.M. Microcystins in groundwater wells and their accumulation in vegetable plants irrigated with contaminated waters in Saudi Arabia. *J. Hazard. Mater.* **2009**, *172*, 310-315. doi:10.1016/j.jhazmat.2009.07.010
206. Kittler, K.; Schreiner, M.; Krumbein, A.; Manzei, S.; Koch, M.; Rohn, S.; Maul, R. Uptake of the cyanobacterial toxin cylindrospermopsin in Brassica vegetables. *Food Chem.* **2012**, *133*, 875-879. doi:10.1016/j.foodchem.2012.01.107
207. Roney, B.R.; Renhui, L.; Banack, S.A.; Murch, S.; Honegger, R.; Cox, P.A. Consumption of fa cai Nostoc soup: a potential for BMAA exposure from Nostoc cyanobacteria in China? *Amyotroph. Lateral Scler.* **2009**, *10*, 44-49. doi:10.3109/17482960903273031
208. Han, D.; Deng, Z.; Lu, F.; Hu, Z. Biology and biotechnology of edible Nostoc. In *Handbook of microalgal culture: Applied phycology and biotechnology*, 2nd ed.; Richmond, A.; Hu, Q. Eds.; John Wiley & Sons: 2013, pp. 433-444. doi:10.1002/9781118567166.ch23
209. Murch, S.J.; Cox, P.A.; Banack, S.A. A mechanism for slow release of biomagnified cyanobacterial neurotoxins and neuro-degenerative disease in Guam. *Proceed. National Acad. Sci.* **2004b**, *101*, 12228-12231. doi:10.1073/pnas.0404926101
210. Downing, S.; Van Onselen, R.; Kemp, G.; Downing, T.G. Metabolism of the neurotoxic amino acid β -N-methylamino-L-alanine in human cell culture models. *Toxicon* **2019**, *168*, 131-139. doi:10.1016/j.toxicon.2019.07.007
211. Brown Jr, R.M.; Larson, D.A.; Bold, H.C. Airborne algae: their abundance and heterogeneity. *Science* **1964**, *143*, 583-585. doi:10.1126/science.143.3606.583
212. Olson, N.E.; Cooke, M.E.; Shi, J.H.; Birbeck, J.A.; Westrick, J.A.; Ault, A.P. Harmful algal bloom toxins in aerosol generated from inland lake water. *Environ. Sci. Technol.* **2020**, *54*, 4769-4780. doi:10.1021/acs.est.9b07727
213. Harb, C.; Pan, J.; DeVilbiss, S.; Badgley, B.; Marr, L.C.; Schmale III, D.G.; Foroutan, H. Increasing freshwater salinity impacts aerosolized bacteria. *Environ. Sci. Technol.* **2021**, *55*, 5731-5741. doi:10.1021/acs.est.0c08558
214. Brodie, E.L.; DeSantis, T.Z.; Parker, J.P.M.; Zubietta, I.X.; Piceno, Y.M.; Andersen, G.L. Urban aerosols harbor diverse and dynamic bacterial populations. *PNAS* **2007**, *104*, 299-304. doi:10.1073/pnas.0608255104
215. Wood, S.A.; Dietrich, D.R. Quantitative assessment of aerosolized cyanobacterial toxins at two New Zealand lakes. *J. Environ. Monit.* **2011**, *13*, 1617-1624. doi:10.1039/C1EM10102A
216. Genitsaris, S.; Kormas, K.A.; Moustaka-Gouni, M. Airborne algae and cyanobacteria: occurrence and related health effects. *Front. Biosci. — Elite* **2011**, *3*, 772-787. doi:10.2741/e285
217. Wisniewska, K.A.; Śliwińska-Wilczewska, S.; Lewandowska, A.U. Airborne microalgal and cyanobacterial diversity and composition during rain events in the southern Baltic Sea region. *Sci. Rep.* **2022**, *12*, 1-9. doi:10.1038/s41598-022-06107-9
218. Scott, L.L.; Downing, S.; Downing, T.G. The evaluation of BMAA inhalation as a potential exposure route using a rat model. *Neurotox. Res.* **2018**, *33*, 6-14. doi:10.1007/s12640-017-9742-9
219. Oliveira, V.R.; Mancin, V.G.; Pinto, E.F.; Soares, R.M.; Azevedo, S.M.; Macchione, M.; Carvalho, A.R.; Zin, W.A. Repeated intranasal exposure to microcystin-LR affects lungs but not nasal epithelium in mice. *Toxicon* **2015**, *104*, 14-18. doi:10.1016/j.toxicon.2015.07.331

220. Li, X.; Xu, L.; Zhou, W.; Zhao, Q.; Wang, Y. Chronic exposure to microcystin-LR affected mitochondrial DNA maintenance and caused pathological changes of lung tissue in mice. *Environ. Pollut.* **2016**, *210*, 48-56. doi:10.1016/j.envpol.2015.12.001
221. Carvalho, G.M.; Oliveira, V.R.; Soares, R.M.; Azevedo, S.M.; Lima, L.M.; Barreiro, E.J.; Valença, S.S.; Saldiva, P.H.; Faffe, D.S.; Zin, W.A. Can LASSBio 596 and dexamethasone treat acute lung and liver inflammation induced by microcystin-LR? *Toxicon* **2010**, *56*, 604-612. doi:10.1016/j.toxicon.2010.06.005
222. Zhao, S.; Sun, H.; Yan, W.; Xu, D.; Shen, T. A proteomic study of the pulmonary injury induced by microcystin-LR in mice. *Toxicon* **2018**, *150*, 304-314. doi:10.1016/j.toxicon.2018.06.072
223. Breidenbach, J.D.; French, B.W.; Gordon, T.T.; Kleinhenz, A.L.; Khalaf, F.K.; Willey, J.C.; Hammersley, J.R.; Wooten, R.M.; Crawford, E.L.; Modyanov, N.N. Microcystin-LR aerosol induces inflammatory responses in healthy human primary airway epithelium. *Environ. Int.* **2022**, *169*, 107531. doi:10.1016/j.envint.2022.107531
224. Facciponte, D.N.; Bough, M.W.; Seidler, D.; Carroll, J.L.; Ashare, A.; Andrew, A.S.; Tsongalis, G.J.; Vaickus, L.J.; Henegan, P.L.; Butt, T.H. Identifying aerosolized cyanobacteria in the human respiratory tract: A proposed mechanism for cyanotoxin-associated diseases. *Sci. Total Environ.* **2018**, *645*, 1003-1013. doi:10.1016/j.scitotenv.2018.07.226
225. Pierozan, P.; Piras, E.; Brittebo, E.; Karlsson, O. The cyanobacterial neurotoxin β -N-methylamino-L-alanine (BMAA) targets the olfactory bulb region. *Arch. Toxicol.* **2020**, *94*, 2799-2808. doi:10.1007/s00204-020-02775-6
226. Sarasa, M.; Pesini, P. Natural non-transgenic animal models for research in Alzheimer's disease. *Curr. Alzheimer Res.* **2009**, *6*, 171-178. doi:10.2174/156720509787602834
227. Davis, D.A.; Cox, P.A.; Banack, S.A.; Lecusay, P.D.; Garamszegi, S.P.; Hagan, M.J.; Powell, J.T.; Metcalf, J.S.; Palmour, R.M.; Beierschmitt, A. L-serine reduces spinal cord pathology in a vervet model of preclinical ALS/MND. *J. Neuropathol. Exp. Neurol.* **2020**, *79*, 393-406. doi:10.1093/jnen/nlaa002
228. Mondo, K.; Hammerschlag, N.; Basile, M.; Pablo, J.; Banack, S.A.; Mash, D.C. Cyanobacterial neurotoxin β -N-methylamino-L-alanine (BMAA) in shark fins. *Marine drugs* **2012**, *10*, 509-520. doi:10.3390/md10020509
229. Davis, D.A.; Garamszegi, S.P.; Banack, S.A.; Dooley, P.D.; Coyne, T.M.; McLean, D.W.; Rotstein, D.S.; Mash, D.C.; Cox, P.A. BMAA, Methylmercury, and mechanisms of neurodegeneration in dolphins: A natural model of toxin exposure. *Toxins* **2021**, *13*, 697. doi:10.3390/toxins13100697
230. Davis, D.A.; Mondo, K.; Stern, E.; Annor, A.K.; Murch, S.J.; Coyne, T.M.; Brand, L.E.; Niemeyer, M.E.; Sharp, S.; Bradley, W.G. Cyanobacterial neurotoxin BMAA and brain pathology in stranded dolphins. *PLoS One* **2019**, *14*, e0213346. doi:10.1371/journal.pone.0213346
231. Lopicic, S.; Svirčev, Z.; Palanački Malešević, T.; Kopitović, A.; Ivanovska, A.; Meriluoto, J. Environmental neurotoxin β -N-Methylamino-L-alanine (BMAA) as a widely occurring putative pathogenic factor in neurodegenerative diseases. *Microorganisms* **2022**, *10*, 2418. doi:10.3390/microorganisms10122418
232. Lobner, D.; Piana, P.M.T.; Salous, A.K.; Peoples, R.W. β -N-methylamino-L-alanine enhances neurotoxicity through multiple mechanisms. *Neurobiol. Dis.* **2007**, *25*, 360-366. doi:10.1016/j.nbd.2006.10.002
233. Pedrosa, C.d.S.; Souza, L.R.; Gomes, T.A.; de Lima, C.V.; Ledur, P.F.; Karmirian, K.; Barbeito-Andres, J.; Costa, M.d.N.; Higa, L.M.; Rossi, Á.D. The cyanobacterial saxitoxin exacerbates neural cell death and brain malformations induced by Zika virus. *PLOS Negl. Trop. Dis.* **2020**, *14*, e0008060. doi:10.1371/journal.pntd.0008060
234. Wiese, M.; D'agostino, P.M.; Mihali, T.K.; Moffitt, M.C.; Neilan, B.A. Neurotoxic alkaloids: saxitoxin and its analogs. *Marine drugs* **2010**, *8*, 2185-2211. doi:10.3390/md8072185
235. Smith, F.M.; Wood, S.A.; van Ginkel, R.; Broady, P.A.; Gaw, S. First report of saxitoxin production by a species of the freshwater benthic cyanobacterium, *Scytonema* Agardh. *Toxicon* **2011**, *57*, 566-573. doi:10.1016/j.toxicon.2010.12.020
236. Weiss, C.; Figueras, E.; Borbely, A.N.; Sewald, N. Cryptophycins: cytotoxic cyclodepsipeptides with potential for tumor targeting. *J. Pept. Sci.* **2017**, *23*, 514-531. doi:10.1002/psc.3015
237. Thuan, N.H.; An, T.T.; Shrestha, A.; Canh, N.X.; Sohng, J.K.; Dhakal, D.; Recent advances in exploration and biotechnological production of bioactive compounds in three cyanobacterial genera: *Nostoc*, *Lyngbya*, and *Microcystis*. *Front. Chem.* **2019**, *7*, 604. doi:10.3389/fchem.2019.00604.
238. Monteiro, P.R.; do Amaral, S.C.; Siqueira, A.S.; Xavier, L.P.; Santos, A.V. Anabaenopeptins: What We Know So Far. *Toxins* **2021**, *13*, 522. doi:10.3390/toxins13080522
239. Scott, L.; Downing, T. Dose-dependent adult neurodegeneration in a rat model after neonatal exposure to β -n-methylamino-L-alanine. *Neurotox. Res.* **2019**, *35*, 711-723. doi: 10.1007/s12640-019-9996-5
240. Karlsson, O.; Lindquist, N.G.; Brittebo, E.B.; Roman, E. Selective brain uptake and behavioral effects of the cyanobacterial toxin BMAA (β -N-methylamino-L-alanine) following neonatal administration to rodents. *Toxicol. Sci.* **2009**, *109*, 286-295. doi:10.1093/toxsci/kfp062
241. Karlsson, O.; Berg, A.-L.; Hanrieder, J.; Arnerup, G.; Lindström, A.-K.; Brittebo, E.B. Intracellular fibril formation, calcification, and enrichment of chaperones, cytoskeletal, and intermediate filament proteins in

- the adult hippocampus CA1 following neonatal exposure to the nonprotein amino acid BMAA. *Arch. Toxicol.* **2015b**, *89*, 423-436. doi:10.1007/s00204-014-1262-2
242. Pierozan, P.; Karlsson, O. Mitotically heritable effects of BMAA on striatal neural stem cell proliferation and differentiation. *Cell Death Dis.* **2019**, *10*, 1-14. doi:10.1038/s41419-019-1710-2
 243. Laugeray, A.; Oummadi, A.; Jourdain, C.; Feat, J.; Meyer-Dilhet, G.; Menuet, A.; Plé, K.; Gay, M.; Routier, S.; Mortaud, S. Perinatal exposure to the cyanotoxin β -N-méthylamino-L-alanine (BMAA) results in long-lasting behavioral changes in offspring—potential involvement of DNA damage and oxidative stress. *Neurotox. Res.* **2018**, *33*, 87-112. doi:10.1007/s12640-017-9802-1
 244. Zlokovic, B. Alzheimer's disease: a matter of blood-brain barrier dysfunction? *Alzheimers Dement.* **2018**, *14*, P205-P205. doi:10.1016/j.jalz.2018.06.2314
 245. Yan, M.; Jin, H.; Pan, C.; Han, X. Chronic microcystin-LR-induced α -synuclein promotes neuroinflammation through activation of the NLRP3 inflammasome in microglia. *Mol. Neurobiol.* **2022**, *1*-17. doi:10.1007/s12035-022-03134-5
 246. Verheijen, J.; Sleegers, K. Understanding Alzheimer disease at the interface between genetics and transcriptomics. *Trends Genet.* **2018**, *34*, 434-447. doi:10.1016/j.tig.2018.02.007
 247. Chiu, A.S.; Gehringer, M.M.; Braid, N.; Guillemain, G.J.; Welch, J.H.; Neilan, B.A. Gliotoxicity of the cyanotoxin, β -methyl-amino-L-alanine (BMAA). *Sci. Rep.* **2013**, *3*, 1-5. doi:10.1038/srep01482
 248. Chiu, A.S.; Braid, N.; Marçal, H.; Welch, J.H.; Gehringer, M.M.; Guillemain, G.J.; Neilan, B.A. Global cellular responses to β -methyl-amino-L-alanine (BMAA) by olfactory ensheathing glial cells (OEC). *Toxicon* **2015**, *99*, 136-145. doi:10.1016/j.toxicon.2015.03.009
 249. Bubik, A.; Sedmak, B.; Novinec, M.; Lenarčič, B.; Lah, T.T. Cytotoxic and peptidase inhibitory activities of selected non-hepatotoxic cyclic peptides from cyanobacteria. *Biol. Chem.* **2008**, *389*, 1339-1346. doi:10.1515/BC.2008.153
 250. Mello, F.D.; Braid, N.; Marçal, H.; Guillemain, G.; Nabavi, S.M.; Neilan, B.A. Mechanisms and effects posed by neurotoxic products of cyanobacteria/microbial eukaryotes/dinoflagellates in algae blooms: A review. *Neurotox. Res.* **2018**, *33*, 153-167. doi:10.1007/s12640-017-9780-3
 251. Soto, T.; Buzzi, E.D.; Rotstein, N.P.; German, O.L.; Politi, L.E. Damaging effects of BMAA on retina neurons and Müller glial cells. *Exp. Eye Res.* **2021**, *202*, 108342. doi:10.1016/j.exer.2020.108342
 252. Tan, V.X.; Mazzocco, C.; Varney, B.; Bodet, D.; Guillemain, T.A.; Bessede, A.; Guillemain, G.J. Detection of the cyanotoxins L-BMAA uptake and accumulation in primary neurons and astrocytes. *Neurotox. Res.* **2018**, *33*, 55-61. doi:10.1007/s12640-017-9787-9
 253. Forum on Neuroscience and Nervous System Disorders; Board on Health Sciences Policy; Institute of Medicine. *Neurodegeneration: Exploring Commonalities Across Diseases: Workshop Summary*; National Academies Press: Washington (DC): US; December 12, 2013. doi:10.17226/18341
 254. Drummond, D.A.; Wilke, C.O. Mistranslation-induced protein misfolding as a dominant constraint on coding-sequence evolution. *Cell* **2008**, *134*, 341-352. doi:10.1016/j.cell.2008.05.042
 255. Martinez-Miguel, V.E.; Lujan, C.; Espie, T.; Martinez-Martinez, D.; Moore, S.; Backes, C.; Gonzalez, S.; Galimov, E.R.; Brown, A.E.; Halic, M. Increased fidelity of protein synthesis extends lifespan. *Cell metabolism* **2021**, *33*, 2288-2300. doi:10.1016/j.cmet.2021.08.017
 256. Brilkova, M.; Nigri, M.; Kumar, H.S.; Moore, J.; Mantovani, M.; Keller, C.; Grimm, A.; Eckert, A.; Shcherbakov, D.; Akbergenov, R. Error-prone protein synthesis recapitulates early symptoms of Alzheimer disease in aging mice. *Cell Rep.* **2022**, *40*, 111433. doi:10.1016/j.celrep.2022.111433
 257. Lant, J.T.; Kiri, R.; Duennwald, M.L.; O'Donoghue, P. Formation and persistence of polyglutamine aggregates in mistranslating cells. *Nucleic Acids Res.* **2021**, *49*, 11883-11899. doi:10.1093/nar/gkab898
 258. Tawfik, D.S.; Gruic-Sovulj, I. How evolution shapes enzyme selectivity—lessons from aminoacyl-tRNA synthetases and other amino acid utilizing enzymes. *FEBS J.* **2020**, *287*, 1284-1305. doi:10.1111/febs.15199
 259. Glover, W.; Mash, D.C.; Murch, S.J. The natural non-protein amino acid N- β -methylamino-L-alanine (BMAA) is incorporated into protein during synthesis. *Amino Acids* **2014**, *46*, 2553-2559. doi:10.1007/s00726-014-1812-1
 260. Zhang, H.; Yang, X.-L.; Sun, L. The uniqueness of AlaRS and its human disease connections. *RNA Biol.* **2021**, *18*, 1501-1511. doi:10.1080/15476286.2020.1861803
 261. Wang, Z.-Q.; Zhang, C.-C. A tRNA t6A modification system contributes to the sensitivity towards the toxin β -N-methylamino-L-alanine (BMAA) in the cyanobacterium *Anabaena* sp. PCC 7120. *Aquat. Toxicol.* **2022**, *245*, 106121. doi:10.1016/j.aquatox.2022.106121
 262. Okamoto, S.; Esumi, S.; Hamaguchi-Hamada, K.; Hamada, S. β -N-methylamino-L-alanine (BMAA) suppresses cell cycle progression of non-neuronal cells. *Scientific reports* **2018**, *8*, 1-10. doi:10.1038/s41598-018-36418-9.