

## Article

# Long Covid and Neurodegenerative Disease

Patrick Chambers

Department of Pathology, Torrance Memorial Medical Center, Torrance, California  
Email: pwc@gte.net

**Abstract:** Brain fog with compromised ability to concentrate has been the most frequent Long Covid (LC) complaint. This is due to an increased transforming growth factor (TGF)  $\beta$ /interferon (IFN)- $\gamma$  ratio with consequently increased bradykinin (BKN), especially in Caucasian females. Brain and lung blood vessels "leak." This same ratio is increased in Alzheimer's disease (AD), but decreased in Parkinson's disease (PD), because CD4<sup>+</sup> and CD8<sup>+</sup> T cells are differentially affected by the invading associated viruses, e.g., SARS CoV2, HIV, EBV,.... In Covid-19 CD147 receptors on immune cells are critical in generating the increased TGF- $\beta$ /IFN- $\gamma$  and those on endothelial cells, platelets, and erythrocytes are critical to the abnormal microvascular blood flow. ACE2 receptors on pneumocytes and enterocytes enable pulmonary and GI entry, initiating gut dysbiosis. Epigenetics, methylation, magnesium, vitamin D, the B vitamins, and antioxidants suggest that these issues can be surmounted. Biochemical, physiologic, and epidemiologic data are analyzed to answer these questions. An LC model is presented and discussed in the context of the most recent research. Suggestions to avoid these and other worrisome concerns are included. Other topics discussed include estrogen, the gut microbiome, type 2 diabetes (T2D), and homocysteine.

**Keywords** homocysteine; estrogen; bradykinin; magnesium; vitamin D; CD147

## 1. Introduction

Long Covid (LC) has replaced Covid-19 as the topic du jour. Long term LC risks are unknown but have stoked growing concern. The neurodegenerative and tumorigenic implications are at the top of this list. Unfortunately the wide spectrum of LC symptoms has defied mechanistic attempts to link their pathogenesis. There are clearly multiple factors involved, complicating these attempts. The male dominated Covid-19 stands in stark contrast to the female dominated LC. "Evidence based" efforts investigating such issues have traditionally relied on Random Clinical Trials (RCTs) and meta-analyses - the top down approach. Two inherent problems are loss of timeliness (RCTs) and diluted results (meta-analyses). A bottom up approach based on biochemistry, physiology, and epidemiology may be more advantageous, given the urgency of and universal interest in LC. This is an opinion piece aided by the deluge of recent research on this burgeoning problem.

## 2. LC Model and Implications

### Model

- CD147 receptors on T cells bind CD147 epitopes (the falciparum antigen) on the spike protein S<sup>1,2</sup> (no ACE2 receptors on circulating immune cells<sup>3,4,5</sup> or on erythrocytes)
- Subsequently SARS CoV2 overwhelms and exhausts CD4<sup>+</sup> and CD8<sup>+</sup> T cells and natural killer cells (NKs)
- Persistent chronic lymphopenia after Covid-19<sup>6</sup> lowers secretion of IFN- $\gamma$  (type II IFN),<sup>7,8</sup> produced only by T cells and NK cells, but especially by CD4<sup>+</sup> T cells.
- Decreased secretion of IFN- $\gamma$  lowers hepatic synthesis of complement component 1 inhibitor (C1-INH)<sup>9</sup>
- Uninhibited C1 triggers the Classic Complement Pathway (CCP) and crosstalk with the Kallikrein Kinin System (KKS)<sup>10</sup>

- The KKS triggers an increase in BKN, which is normally catabolized by angiotensin converting enzyme (ACE)
- Estrogen downregulates ACE and prolongs BKN half life<sup>11</sup> This makes estrogen an ACE inhibitor of sorts and increases the risk of some cancers<sup>12,13,14</sup>
- BKN enhances vascular permeability creating “leaks” primarily in lungs<sup>15</sup> and brain<sup>16</sup>, linking brain fog and dyspnea/post exertional malaise
- IL-1 $\beta$ , prominent in LC<sup>17</sup>, potentiates the BKN induced microvascular leakage<sup>18</sup> and brain fog primarily in Caucasian females.
- The ACE DD genotype in African Americans, an evolutionary adaptation to falciparum malaria, downregulates this leakage (tighter endothelial junctions) and elevates the relative frequency of LC in Caucasian females
- IFN- $\gamma$  and TGF- $\beta$  counterbalance each other<sup>19,20</sup> and the loss of IFN- $\gamma$  secreted by CD4+ and CD8+ T cells leaves TGF- $\beta$  unopposed.
- Chronic low grade IL-1 $\beta$  and TNF- $\alpha$  redirect pleiotropic TGF- $\beta$  from wound healing fibrosis to endothelial mesenchymal transition (EndMT)/epithelial mesenchymal transition (EMT)<sup>21,22</sup> and from tumor suppressor to tumor promoter<sup>23</sup>
- The switch of pleiotropic TGF- $\beta$  from anti-inflammatory to proinflammatory appears to be more organ specific, e.g., neurovascular pericytes<sup>24</sup>

### Implications

- Long term LC may drive an increase in sporadic/late onset AD due to an elevated TGF- $\beta$ /IFN- $\gamma$  <sup>25,26,27</sup>.
- Late onset AD may appear earlier
- AD frequency in Caucasian females, especially in those also on hormone replacement therapy (HRT)<sup>28,29</sup>, may approach that in African American females.
- Cancer risk/progression and fibrosis may also increase in Caucasian females
- CD147 is the primary receptor involved in the pathogenesis of ASCVD<sup>30,31</sup> and LC long term may increase its incidence
- The presence of CD147 receptors (but not ACE2 receptors) on platelets (and erythrocytes) creates platelet aggregates, further complicating the microcirculation<sup>32</sup> (elevated mean platelet volume (MPV)).
- The presence of the CD147 epitope on the spike protein S poses significant risks involving microvascular thrombosis in the short term for all exposed to the spike protein S.
- The preference of SARS CoV2 for CD4+ T cells (laden with CD147 receptors) may cause growing cognitive impairment (without actual dementia) and reactivation of EBV (further exhausts CD4+ T cells) in LC and post vaccination.
- The incidence of PD in diabetics post Covid-19 should also increase due to a depressed TGF- $\beta$ /IFN- $\gamma$  in PD.

## 3. Discussion

### 3.1. To LC and Beyond

Viral load and TGF- $\beta$ /IFN- $\gamma$  ratio determine Covid-19 symptoms (or not). This ratio decreases notably from the control group, passing through asymptomatic, up to symptomatic SARS-CoV-2 individuals<sup>33</sup>.

But as IFN- $\gamma$  secreting T cells are lost to the invading virus (see figure 1), this ratio inverts (increases). A depressed TGF- $\beta$ /IFN- $\gamma$  is affiliated with PD<sup>34,35</sup>. An elevated ratio is affiliated with AD<sup>25,26</sup>. This suggests that TGF- $\beta$  in those with LC may trigger fibrosis in multiple organs<sup>36,37</sup> (see section 3.3).




Figure 1. SARS CoV2 destroys IFN-γ producing cells<sup>38</sup>.





Figure 2. Low levels of IFN-γ persist into the LC phase<sup>39</sup>.

The cytokines of LC provide insight to possible future complications. TGF-β and IFN-γ are pleiotropic LC linked cytokines that can work in either direction, i.e., anti- to

proinflammatory, suppressing to promoting EMT or tumor for TGF- $\beta$  and anti- to proliferative, pro-apoptotic to necrotic, antitumor to tumor for IFN- $\gamma$ . However, it appears that the deleterious effects of the switch are more significant for TGF- $\beta$ . IFN- $\gamma$  appears to retain a net positive effect<sup>40</sup>.

TGF- $\beta$  functions initially as an anti-inflammatory, keeping the inflammatory response of IFN- $\gamma$  under control. Under chronic inflammatory conditions TNF- $\alpha$  is elevated and can upregulate TGF- $\beta$ , which opposes IFN- $\gamma$ . Chronic low doses of TGF- $\beta$  when combined with chronic low doses of TNF- $\alpha$  facilitate the switch of TGF- $\beta$  from suppressing to promoting tumor<sup>23</sup>. This also appears to be the case for switching TGF- $\beta$  from wound healing fibrosis to endothelial or epithelial mesenchymal transition (EndMT<sup>41</sup> or EMT<sup>42</sup>).

Therefore, it seems reasonable to assume that TNF- $\alpha$  (chronic inflammation) might redirect TGF- $\beta$  from anti-inflammatory to pro-inflammatory and open the door to neuro-degenerative disease. The constant stimulus to chronic inflammation posed by residual spike protein S could easily trigger this (see figure 3)



**Figure 3.** LC is characterized by persistence of the spike protein S<sup>43</sup>.

3.2. AD, PD, and LBD

The three primary forms of accelerated pre-senile dementia are AD, PD, and Lewy Body Dementia (LBD). AD, the predominant form of pre-senile dementia, is more common in females while PD is more common in males. LBD is intermediate<sup>44</sup>. AD is characterized by an elevated TGF- $\beta$ /IFN- $\gamma$ <sup>26,27</sup>. Onset of PD in females is later and less severe than that in men<sup>45</sup> and might be due to benefits from premenopausal estrogen or hormone replacement therapy<sup>46</sup>.


Although protective against PD, estrogen possesses ACE inhibitor properties that elevate BKN. Lung and brain BKN induced leakage contributes to LC and AD. African American females appear to be less affected by LC but suffer the highest incidence of AD (by gender or race). At first this seems contradictory for two reasons. First, in African

American but not in Caucasian females estrogen levels decrease with increasing premenopausal age and BMI<sup>47</sup>, decreasing their risk of LC. Secondly, the incidence of the ACE DD genotype (tighter endothelial junctions) is higher, also decreasing their risk of LC. However, according to a recent NHANES survey the incidence of obesity in African American females was 50% greater than that in Caucasian males, Caucasian females, or African-American males. This is presumed to be due to dietary factors, possibly monosodium glutamate (MSG)<sup>48</sup>. Perhaps escalating dietary MSG, induced by obesity and diabetes<sup>49,50</sup>, overwhelms the protective properties of the ACE DD genotype, yielding more AD in African-American females. Obesity and diabetes also up regulate TGF- $\beta$ , increasing the risk of AD<sup>51</sup>.

Individuals with diabetes are up to four times more likely to develop LC.

Although homocysteine is elevated and contributes to the development of AD, LBD, and PD, PD is different. It is in some ways the opposite of AD, e.g., TGF- $\beta$ /IFN- $\gamma$  is depressed, not elevated<sup>34,35</sup>. AD, LBD, and PD all feature extracellular plaques - amyloid  $\beta$  in AD,  $\alpha$ -synuclein in LBD and PD. IFN- $\gamma$ , elevated in PD, triggers microglial removal of amyloid  $\beta$ <sup>52,53</sup>.

PD also exhibits abnormal tryptophan metabolism due to increased IFN- $\gamma$  and perhaps pyridoxal-5-phosphate (P5P) deficiency<sup>54</sup> (see figure 4).



**Figure 4.** IFN- $\gamma$  shunts tryptophan away from serotonin synthesis and production of melatonin<sup>55,56</sup>.

P5P is a required cofactor for aromatic amino acid decarboxylase (AADC) (see figure 5), which produces DOPAMINE and serotonin. Additionally magnesium is a required cofactor for the synthesis of melatonin from serotonin. All three hormones are deficient in PD.

CD4<sup>+</sup> (superior producers of IFN- $\gamma$ <sup>57</sup>) and not CD8<sup>+</sup> T cells appear to be responsible for the increased IFN- $\gamma$  in PD<sup>58</sup>. SARS CoV2 preferentially attacks CD4<sup>+</sup> T cells without involvement of ACE2 receptors<sup>59</sup>.

### 3.3. TGF and IFN

CD4 T cells can mount a much stronger IFN- $\gamma$  response than CD8 T cells<sup>57</sup>. A high CD4<sup>+</sup>/CD8<sup>+</sup> characterizes PD<sup>60</sup>. IFN- $\gamma$  induces loss of dopamine neurons and nigrostriatal degeneration<sup>61</sup>.

There are some interesting parallels between PD, HIV/AIDS, and T2D that involve IFN- $\gamma$ . HIV preferentially destroys CD8<sup>+</sup> T cells<sup>62</sup>, accentuating the IFN- $\gamma$  response from

CD4<sup>+</sup> T cells<sup>63</sup>. The HIV induced IFN- $\gamma$  increases the risk of PD<sup>64</sup> and autoimmune disease<sup>65</sup>. T2D also increases the risk of PD<sup>66,67</sup>. HIV increases the incidence of T2D<sup>68</sup> and PD. TNF- $\alpha$  is chronically elevated in T2D<sup>69</sup>, AD, LBD, and PD. The most recent research suggests that Parkinson's is an autoimmune disease, which conforms to the well known linkage between IFN- $\gamma$  and autoimmune disease. Abnormal tryptophan metabolism exhibited in Parkinson's disease, due to an IFN- $\gamma$  imbalance, is also seen in T2D<sup>70</sup> and HIV<sup>71</sup>. TNF- $\alpha$  is elevated in AD, LBD, and PD<sup>72</sup>.

In summary TGF beta is a key player in AD; likewise for IFN gamma in PD. TGF beta and IFN gamma are both pleiotropic and the direction of each appears to be determined by the chronic cytokine milieu, including pleiotropic IL-1beta, pleiotropic TNF alpha, and IL-6. Perhaps chronic inflammation (post viral infection that targets CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells) in those with marginal onboard antioxidants are predisposed to AD and PD.

TGF- $\beta$ /Smad signaling pathway in renal, hepatic, pulmonary and cardiac fibrosis has been well documented<sup>73</sup>. Recent studies show this to include the brain as well<sup>74,75,76</sup>. Magnesium possesses the capacity to down regulate this SMAD pathway in the liver<sup>77</sup> and the lungs<sup>78</sup>. IFN- $\gamma$  also appears capable in this regard<sup>79</sup>. Exercise also helps by increasing IFN- $\gamma$ <sup>80</sup>. This is because exercise upregulates angiotensin II<sup>81</sup> and angiotensin II upregulates IFN- $\gamma$ <sup>82</sup>.

### 3.4. ACE and BKN

Angiotensin converting enzyme (ACE) produces angiotensin II and degrades BKN. Estrogen downregulates ACE, up regulating BKN. BKN upregulates tyrosine hydroxylase, the rate limiting step in dopamine synthesis<sup>83</sup>. This might help explain the protective effects of estrogen in avoiding PD.

The frequency of the ACE II genotype in AD is 1.4x higher than that in controls v 0.4x for the ACE DD genotype<sup>84</sup>. ACE levels are up to 70% higher in the DD genotype<sup>85</sup>. Endothelial cell junctions are tighter and less permeable in the DD ACE genotype<sup>86</sup>.

However after menopause tight junction permeability increases due to endothelial dysfunction initiated by oxidative stress, microthrombosis (loss of RBC deformability), immune complexes (endothelial CD147 and perhaps ACE2 receptors linking to spike S epitopes), ..... The ACE DD genotype is protective against AD not only due to tighter endothelial junctions but also to increased ACE and less BKN<sup>87,88</sup>.

BKN induced endothelial permeability not only produces perivascular angioedema but also leads to increased fibrosis<sup>89</sup>. Either mechanism may contribute to the brain fog, dyspnea, and post exertional malaise of LC.

Because ACE is higher and BKN levels are commensurately lower in those with the ACE DD genotype, African American females should be less likely to develop LC versus their Caucasian counterparts. Some studies purport to show a decreased incidence of cancer in those with Alzheimer's disease. However, African-American females are 40% more likely to die of breast cancer than Caucasian females<sup>90</sup>. African-Americans have the highest death rate and shortest survival of any racial/ethnic group in the US for most cancers and have a greater incidence of Alzheimer's disease than any other racial group in America<sup>91</sup>.

On the other hand, overall cancer risk is lower in people with PD, compared to the general population. This difference in cancer risk between AD and PD speaks to further linkage between TGF- $\beta$  in cancer causation and IFN- $\gamma$  in cancer avoidance<sup>92</sup>. Estrogen and HRT for more than 10 years have been linked to a slight increase in cancer risk<sup>93</sup>. Estrogen downregulates ACE and is an ACE inhibitor of sorts. ACE inhibitors have been linked to an increase in lung cancer<sup>94</sup>. Not surprisingly, BKN has been linked to aggressive prostate cancer<sup>95</sup>.



### 3.5. Homocysteine and B Vitamins

Homocysteine plays a prominent role in all forms of dementia<sup>96</sup>. Asians have a lower Ca:Mg diet and a lower incidence of AD<sup>97</sup>. P5P figures prominently in homocysteine recycling and is a required cofactor for aromatic amino acid decarboxylase (AADC)(see figure 5). PD patients are frequently B6 deficient<sup>98</sup> and exhibit abnormalities in both dopamine and serotonin synthesis<sup>99</sup>. The dopamine synthesis shortfall in PD appears to be primarily driven by IFN- $\gamma$  and its effect on tryptophan metabolism (see figure 4).




Figure 5. AADC requires the cofactor P5P.





Figure 6. Vitamins B2, B3, B6, B9, and B12 are all heavily involved in both the folate cycle and the methionine cycle. All require  $Mg^{++}$  for activation. B6 is also required as a cofactor to recycle homocysteine<sup>100</sup>.

Homocysteine induces oxidative stress<sup>101</sup> and is a marker for COVID-19 severity<sup>102</sup>, LC<sup>103</sup>, and dementia<sup>104,105</sup>. As shown in figure 6, vitamins B2,3,6,9,12<sup>106</sup> are prominent cofactors in both the folate and methionine cycles, which are integral in the metabolism of homocysteine. These B vitamin deficiencies are precisely those associated with cognitive

impairment and AD<sup>107</sup>. B complex supplements usually provide methylated B12, but pyridoxine (B6), and folate (B9) are not their active forms. Indeed, the active forms of B1,2,3,9,12 all require magnesium<sup>100</sup> and some require activated B6 as a cofactor.

B1,3, and 6 must be phosphorylated (ATP and magnesium as chelate); B2,9,12 must be methylated (magnesium as cofactor)<sup>108</sup>. ATP and methylation both require  $Mg^{++}$ .


The active form of B6 (P5P) requires ATP and chelated Mg (phosphorylation) and its activated form as a cofactor, creating a catch 22 situation. Most B6 supplements contain pyridoxine (PNP), which in excess can lead to peripheral neuropathy<sup>109</sup>. This can be avoided by substituting P5P. Rxn 2 (PNPO) in figure 7 mediated by P5P oxidase is the rate limiting step.



**Figure 7.** Note the need for ATP (and  $Mg^{++}$ ) to convert pyridoxal (and PNP, PMP) to P5P. PNPO requires P5P as cofactor.

P5P is critical to its own synthesis, to that of the active forms of other B vitamins, and to the recycling of homocysteine to glutathione (see figure 8). Its critical role in cognition is well known<sup>110,111</sup>. Magnesium is also critical to the synthesis of all endogenous and most exogenous antioxidants<sup>112</sup>.





**Figure 8.** Magnesium and P5P are required to recycle homocysteine.

Vitamin B5 (pantothenate) deficiency is also associated with both AD<sup>113</sup> and PD<sup>114</sup>. In order for pantothenate to reach its active form, three phosphorylations must occur. Each requires ATP and magnesium<sup>115</sup>.

### 3.6. Vitamin D and Ca:Mg

An elevated  $\text{Ca}^{++}$  and a little depressed  $\text{Mg}^{++}$  (high Ca:Mg) are linked to AD<sup>116,117</sup> and PD<sup>118,119</sup>. A recent 2023 article reported that vitamin D, folic acid and vitamin B12 could reverse the cognitive decline leading to AD. Total benefit exceeded that from any single supplement<sup>120</sup>. Adding magnesium and P5P to this regimen would improve the results immeasurably.

As Ca:Mg increases, Vitamin D loses its efficacy for colorectal cancer, prostate cancer, esophageal cancer, cardiovascular disease, metabolic syndrome, total mortality, and cognitive function<sup>121</sup>.

The Western diet is high in processed foods with high monosodium glutamate (MSG)/calcium and low fiber/magnesium. The typical Asian diet offers more magnesium

but less calcium. The target Ca:Mg for both is 2.0. A fiber rich diet that includes fermented vegetables is inversely proportional to Covid-19 mortality in Europe<sup>122,123</sup>, as demonstrated in Germany (sauerkraut) and South Korea ((kimchi)<sup>124</sup>.

Ionized serum Ca:Mg<sup>125</sup> and 25(OH)D<sub>3</sub> are measures of general health that reflect CRP and HRV<sup>126</sup>. Serum magnesium also reflects gut microbiota diversity and gut health<sup>127,128</sup>.

The correlation between the serum magnesium and dementia depends on the Ca:Mg. On a Western diet the healthy upper limit is 2.6. On an Eastern diet the lower limit is 1.7<sup>129</sup>.

### 3.7. APOE and Methylation


Reducing the Ca:Mg ratio reduces the risk of dementia<sup>130</sup>. Methylation of DNA for stability retards the onset of both neurodegenerative disease, e.g., the APOE gene, and cancer<sup>131,132</sup>, e.g., basigin (CD147) gene<sup>133</sup>.

Hypomethylation increases with age and leads to an unstable genome, with activation of some tumor promoter genes<sup>134,135</sup>. Hypomethylation of CpG islands (cytosine-guanine pairs) promotes AD<sup>136</sup>, LBD<sup>137,138</sup>, and PD<sup>139</sup>.

There are three alleles for APOE and APOE4 is the major risk factor for Alzheimer's disease<sup>140</sup>. Hypomethylation of APOE4 is a major determinant in this. 15% of the US population is heterozygous for this allele and 5% are homozygous. Chris Hemsworth recently announced a pullback in his schedule to spend more time with family. It was also reported that while working on a nature film he learned that he was homozygous for APOE4. But as worrisome as that might seem, AD appears to be less genetic and more epigenetic. The native American Indian population presents plenty of APOE4 but very little Alzheimer's disease<sup>141</sup>. The Paleolithic diet provides an excellent Ca:Mg balance. Dairy is not included (eggs 5, milk 10) . Perhaps this dementia gene (APOE4) itself is not the problem but an elevated Ca:Mg, crowding out the Mg<sup>++</sup>. Ca<sup>++</sup> and Mg<sup>++</sup> share the same receptor - CaSR (calcium sensing receptor). DNA methylation occurs via SAmE and magnesium (see figure 8).

### 3.8. Treatment

1. Magnesium and vitamin D (50 ng/mL target) are at the top of the list for both prevention and treatment. Magnesium is a critical mineral in the human body and is involved in ~80% of known metabolic functions<sup>142</sup>. Vitamin D possesses invaluable antioxidant and anti-inflammatory properties (see figure 9). Approximately 75% of human immune system functions depend on maintaining a healthy, physiological



**Figure 9.** Vitamin D provides anti-inflammatory protection upstream and downstream of the AT1R<sup>144</sup>.

2. The target Ca:Mg is 2.0, but any ratio greater than 2.6 or less than 1.7 compromises the efficacy of vitamin D<sup>121</sup>. Covid-19 makes AD<sup>145</sup> worse.  $\text{Ca}^{++}$  dysregulation plays a prominent role in both AD and amyloid  $\beta$  deposition<sup>130</sup>.
3. Antioxidants are vital in the defense of COVID-19 infection. However, if the onboard supply is suboptimal, the vast numbers of ROS generated may overwhelm mitochondria and markedly compromise ATP production. Most endogenously produced and some exogenous antioxidants require ATP (and magnesium) to attain activated status. Vitamin C (water-soluble), vitamins A, D<sub>3</sub>, E, K (all fat-soluble), Zn, D-ribose, selenium, and many others require no processing<sup>112</sup>. Furthermore, hydroxylation of C1 of 25(OH)D in the synthesis of active vitamin D occurs in the mitochondria and is suppressed by calcium<sup>146</sup>. Loss of mitochondria due to oxidative stress compromises vitamin D efficacy in addition to the elevated Ca:Mg.
4. P5P aka PLP (pyridoxal phosphate) is the active form of B6, which is required for activation of many of the B vitamins associated with homocysteine metabolism. P5P is the required cofactor for its own synthesis.
5. A sedentary lifestyle risks eventual obesity and diabetes. Exercise also facilitates a better IFN- $\gamma$ :TGF- $\beta$  balance by increasing IFN- $\gamma$  levels<sup>80</sup>.
6. Probiotics, especially after antibiotic therapy, improves the diversity of the gut microbiome<sup>147,148,149</sup>.
7. Dehydration triggers the renal resorption of  $\text{Na}^{+}$  and water. This also means loss of  $\text{Mg}^{++}$  (and  $\text{K}^{+}$ ) to maintain electroneutrality. Also, the thirst reflex diminishes with age. What good is increased dietary/supplemental  $\text{Mg}^{++}$  in the face of a magnesuric drain. Hydration maintenance, easily overlooked, potentially trumps all the other suggestions.


**4. Conclusion**

The most recent research presented here suggests that AD, PD, LBD, and T2D might all be secondary to an imbalanced TGF- $\beta$ /IFN- $\gamma$  perhaps induced by some previous viral infection that preferentially targeted CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells. These T cells secrete IFN- $\gamma$  with the lion's share coming from CD4<sup>+</sup>. HIV preferentially attacks CD8<sup>+</sup> T cells, leaving

the heavy IFN- $\gamma$  producing CD4<sup>+</sup> T cells. HIV mimics PD in many ways, including susceptibility to de novo T2M. TGF- $\beta$ /IFN- $\gamma$  is elevated in AD (and EBV) and depressed in PD, LBD, HIV, and T2M.

Estrogen protects against and delays PD, but predisposes to LC and postmenopausal AD, especially if on HRT for more than 10 years. Increased BKN mediates LC and AD. The ACE DD genotype is protective against LC and AD. Estrogen down regulates ACE, causing an increase in BKN. Obesity and diabetes predispose to AD, LBD, and PD. Magnesium and P5P, the active form of vitamin B6, are deficient in most with AD, LBD, and PD.

CD147 and BKN in the proposed model play vital roles both in the development of LC and in its long term consequences. But an imbalance between TGF- $\beta$  and IFN- $\gamma$  due to SARS CoV2 induced lymphopenia supercharges the roles of CD147 and BKN. The implications of the proposed hypothesis with respect to LC and its possible long term consequences extend not only to the unvaccinated but also to the vaccinated, who might be even more susceptible to recurrent SARS CoV2 (see figure 10), unless preventative measures are taken.



**Figure 10.** Recurrent COVID-19 is directly related to the number of boosters<sup>150</sup>.

ACE2 receptors are not present on immune cells (CD4<sup>+</sup>, CD8<sup>+</sup> T cells, NK cells), the primary combatants against SARS CoV2. The presence of the ACE2 receptor on endothelial cells is controversial<sup>151,152</sup>. Indeed the pathogenesis of microvascular thrombosis, lymphopenia, and TGF- $\beta$  predominance in Covid-19 cannot be explained without acknowledging the presence of the CD147 epitope on the spike protein S, first reported in a Chinese study<sup>1</sup>. This finding was quickly challenged, but those challenges were later debunked<sup>2</sup>.

However, the worrisome implications of this model are not irreversible. These preventative measures include serum Ca:Mg near 2.0, serum 25(OH)D<sub>3</sub> near 50 ng/mL, and an abundance of micronutrient antioxidants, especially P5P<sup>153</sup>. The efficacy of vitamin D is compromised in the face of an elevated Ca:Mg<sup>121</sup>. But D<sub>3</sub> may be even better for endothelial health<sup>154</sup>. Endothelial competence is not only at the center of LC, AD, LBD, PD, and many cancers but also in the progression of cardiovascular disease, arthritis, multiple sclerosis, and sepsis<sup>154</sup>. So, D<sub>3</sub> may be of benefit even in those with an elevated Ca:Mg.

Vitamin D (1,25(OH)<sub>2</sub>D, the active form, is much more reliant on Mg<sup>++</sup> for its efficacy. It requires Mg<sup>++</sup> as a cofactor for three steps in its synthesis. Even parathormone synthesis requires Mg<sup>++</sup>.

Ultimately exercise, diet including some supplementation, and hydration are the primary determinants of epigenetically determined health over and above those associated with LC. HRV, the fifth vital sign, connects all the vital players - gut microbiome diversity<sup>155</sup>, dietary micronutrients, e.g., magnesium<sup>156</sup>, balanced Ca:Mg<sup>125</sup>, balanced TGF-β<sup>157</sup> and TNF-α<sup>158</sup>, and D<sub>3</sub><sup>125</sup>. It's all about balance.

## References

1. Wang K, Chen W, Zhang Z, Deng Y, Lian J-Q, Du P.. CD147-Spike Protein Is a Novel Route for SARS-CoV-2 Infection to Host Cells *Signal Transduction and Targeted Therapy* (2020) 5:283. <https://doi.org/10.1038/s41392-020-00426-x>,
2. Chambers P, Basigin Binds Spike S on SARS-CoV2. *Open Access Library Journal* (2021) 9:1-19. <https://doi.org/10.4236/oalib.1108064>
3. Behl T, Kaur I, Aleya L, Sehgal A, Singh S, Sharma N. CD147-spike protein interaction in COVID-19: Get the ball rolling with a novel receptor and therapeutic target. *Science of The Total Environment* (2022) 808:2022, 152072. <https://doi.org/10.1016/j.scitotenv.2021.152072>
4. Fenizia C, Galbiati S, Vanetti C, Vago R, Clerici M, Tacchetti C, Daniele T. SARS-CoV-2 Entry: At the Crossroads of CD147 and ACE2. *Cells* (2021) 10(6):1434. <https://doi.org/10.3390/cells10061434>
5. Song X, Hu W, Yu H, Zhao L, Zhao Y, Zhao X. Little to no expression of angiotensin-converting enzyme-2 on most human peripheral blood immune cells but highly expressed on tissue macrophages. *Cytometry* (2020) 1–10. <https://doi.org/10.1002/cyto.a.24285>
6. Manne BK, Denorme F, Middleton EA, Portier I, Rowley JW, Stubben C. Platelet gene expressions and function in patients with COVID-19. *Blood* (2020) 136(11):1317–1329 <https://doi.org/10.1182/blood.2020007214>
7. Varghese J, Sandmann S, Schrempf I-M, Frömmel C, Dugas M. Persistent symptoms and lab abnormalities in patients who recovered from COVID-19 (2021) *Sci Rep* 11, 12775. <https://doi.org/10.1038/s41598-021-91270-8>
8. Kim M-O, Salloum S, Wang JY, Wong LP, Regan J. Type I, II, and III Interferon Signatures Correspond to Coronavirus Disease 2019 Severity. *The Journal of Infectious Diseases* (2021) 224(5):777–782 <https://doi.org/10.1093/infdis/jiab288>
9. Zahedi K, Prada AE, Davis AE 3rd. Transcriptional regulation of the C1 inhibitor gene by gamma-interferon. *J Biol Chem* (1994) 269(13):9669-74. <https://pubmed.ncbi.nlm.nih.gov/8144555/>
10. Bossi, F., Peerschke, E.I., Ghebrehwet, B. and Tedesco, F. (2011) Cross-Talk between the Complement and the Kinin System in Vascular Permeability. *Immunology Letters* (2011) 140, 7-13. <https://doi.org/10.1016/j.imlet.2011.06.006>
11. Gallagher, P.E., Li, P., Lenhart, J.R., Chappell, M.C. and Brosnihan, K.B. Estrogen Regulation of Angiotensin-Converting Enzyme mRNA. *Hypertension* (1999) 33:323-328. <https://doi.org/10.1161/01.HYP.33.1.323>
12. Wang W, Zhou Y, Wei R, Jiang G, Li F, Chen X. Bradykinin promotes proliferation, migration, and invasion of cervical cancer cells through STAT3 signaling pathways. *Oncol Rep* (2019) 42: 2521-27 <https://doi.org/10.3892%2For.2019.7380>
13. Greco S, Elia MG, Muscella A, Romano S, Storelli C, Marsigliante S. Bradykinin stimulates cell proliferation through an extracellular-regulated kinase 1 and 2-dependent mechanism in breast cancer cells in primary culture, *Journal of Endocrinology* (2023) 186(2):291-301. <https://doi.org/10.1677/joe.1.06052>



14. Sun D-P, Lee Y-W, Chen J-T, Lin Y-W, Chen R-M. The Bradykinin-BDKRB1 Axis Regulates Aquaporin 4 Gene Expression and Consequential Migration and Invasion of Malignant Glioblastoma Cells via a Ca<sup>2+</sup>-MEK1-ERK1/2-NF- $\kappa$ B Mechanism. *Cancers* (2020) 12(3):667. <https://doi.org/10.3390/cancers12030667>
15. Ichinose M, Barnes PJ. Bradykinin-induced airway microvascular leakage and bronchoconstriction are mediated via a bradykinin B2 receptor. *Am Rev Respir Dis* (1990) 142(5):1104-7 <https://doi.org/10.1164/ajrccm/142.5.1104>
16. Sarker MH, Hu DE, Fraser PA. Acute effects of bradykinin on cerebral microvascular permeability in the anaesthetized rat. *The Journal of Physiology* (2000) 528: 177-187. <https://doi.org/10.1111/j.1469-7793.2000.00177.x>
17. Schultheiß C, Willscher E, Paschold L, Gottschick C, Klee B, Henkes SS. The IL-1 $\beta$ , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19. *Cell Rep Med.* (2022) 3(6):100663. <https://doi.org/10.1016/j.xcrm.2022.100663>
18. Freitas F, Tibiriçá E, Singh M, Fraser PA, Mann GE. Redox Regulation of Microvascular Permeability: IL-1 $\beta$  Potentiation of Bradykinin-Induced Permeability Is Prevented by Simvastatin. *Antioxidants* (2020) 9(12):1269. <https://doi.org/10.3390/antiox9121269>
19. Delvig AA, Lee JJ, Chrzanowska-Lightowlers ZM, Robinson JH. TGF-beta1 and IFN-gamma cross-regulate antigen presentation to CD4 T cells by macrophages. *J Leukoc Biol.* (2002) 72(1):163-6. <https://pubmed.ncbi.nlm.nih.gov/12101276/>
20. Sobral LM, Montan PF, Martelli-Junior H, Graner E, Coletta RD. Opposite effects of TGF- $\beta$ 1 and IFN- $\gamma$  on transdifferentiation of myofibroblast in human gingival cell cultures. *Journal of Clinical Periodontology* (2007) 34: 397-406. <https://doi.org/10.1111/j.1600-051X.2007.01063.X>
21. Yoshimatsu Y, Kimuro S, Pauty J, Takagaki K, Nomiyama S, Inagawa A. TGF-beta and TNF-alpha cooperatively induce mesenchymal transition of lymphatic endothelial cells via activation of Activin signals. *PLoS ONE* (2020) 15 (5): e0232356. <https://doi.org/10.1371/journal.pone.0232356>
22. Masola V, Carraro A, Granata S, Signorini L, Bellin G. In vitro effects of interleukin (IL)-1 beta inhibition on the epithelial-to-mesenchymal transition (EMT) of renal tubular and hepatic stellate cells *J Transl Med* (2019) 17:12. <https://doi.org/10.1186/s12967-019-1770-1>
23. Liu ZW, Zhang YM, Zhang LY, Zhou T, Li YY, Zhou GC. Duality of Interactions Between TGF- $\beta$  and TNF- $\alpha$  During Tumor Formation. *Front Immunol* (2022) 12:810286. <https://doi.org/10.3389/fimmu.2021.810286>
24. Rustenhoven J, Aalderink M, Scotter EL, Oldfield RL, Bergen PS. TGF-beta1 regulates human brain pericyte inflammatory processes involved in neurovasculature function. *J Neuroinflammation* (2016) 13:37 <https://doi.org/10.1186/s12974-016-0503-0>
25. Rota E, Bellone G, Rocca P. Increased intrathecal TGF- $\beta$ 1, but not IL-12, IFN- $\gamma$  and IL-10 levels in Alzheimer's disease patients (2006) *Neurol Sci* 27:33-39. <https://doi.org/10.1007/s10072-006-0562-6>
26. Taipan R, das Neves SP, Sousa AL, Fernandez J Pinto, Correia AP. Proinflammatory and anti-inflammatory cytokines in the CSF of patients with Alzheimer's disease and their correlation with cognitive decline *Neurobiology of Aging* (2019) 76:125-132 <https://doi.org/10.1016/j.neurobiolaging.2018.12.019>
27. Zhang X, Huang WJ, Chen WW. TGF- $\beta$ 1 factor in the cerebrovascular diseases of Alzheimer's disease. *Eur Rev Med Pharmacol Sci.* (2016) 20(24):5178-5185. <https://pubmed.ncbi.nlm.nih.gov/28051272/>



28. Maki PM, Girard LM, Manson JE. Menopausal hormone therapy and cognition *BMJ* (2019) 364:l877  
<https://doi.org/10.1136/bmj.l877>
29. Savolainen-Peltonen H, Rahkola-Soisalo P, Hoti F, Vattulainen P, Gissler M, Ylikorkala O. Use of postmenopausal hormone therapy and risk of Alzheimer's disease in Finland: nationwide case-control study *BMJ* (2019)364:l665.  
<https://doi.org/10.1136/bmj.l665>
30. Chen PY, Qin L, Li G. Endothelial TGF- $\beta$  signalling drives vascular inflammation and atherosclerosis. *Nat Metab* (2019) 1:912–926 <https://doi.org/10.1038/s42255-019-0102-3>
31. Wang C, Jin R, Zhu X. Function of CD147 in Atherosclerosis and Atherothrombosis. *J. of Cardiovasc Trans. Res.* (2015) 8:59–66 <https://doi.org/10.1007/s12265-015-9608-6>
32. Nishikawa M, Kanno H, Zhou Y. Massive image-based single-cell profiling reveals high levels of circulating platelet aggregates in patients with COVID-19. *Nat Commun* (2021) 12, 7135. <https://doi.org/10.1038/s41467-021-27378-2>
33. Villalba MCM, Ramírez OV, Jiménez MM, Garcia AA, Alfonso JM, Baéz GG. Interferon gamma, TGF- $\beta$ 1 and RANTES expression in upper airway samples from SARS-CoV-2 infected patients. *Clinical Immunology* (2020) 220:108576  
<https://doi.org/10.1016/j.clim.2020.108576>
34. Tesseur I, Nguyen A, Chang B, Li L, Woodling NS, Wyss-Coray T. Deficiency in Neuronal TGF- $\beta$  Signaling Leads to Nigrostriatal Degeneration and Activation of TGF- $\beta$  Signaling Protects against MPTP Neurotoxicity in Mice. *J Neurosci.* (2017) 37(17):4584–92. <https://doi.org/10.1523/JNEUROSCI.2952-16.2017>
35. Mohamed A, Maha A, Hanan H, Samaa T. Association of Serum Interferon Gamma Level with Parkinson's Disease in Egyptian Patients *EJMM-Egyptian Journal of Medical Microbiology [The]* (2014) 23 (1):1-5 <http://dx.doi.org/10.12816/0024255>
36. Hu ZJ, Xu J, Yin JM, Li L, Hou W, Zhang LL. Lower Circulating Interferon-Gamma Is a Risk Factor for Lung Fibrosis in COVID-19 Patients. *Front Immunol.* (2020) 11:585647. <https://doi.org/10.3389/fimmu.2020.585647>
37. Weng H, Mertens PR, Gressner AM, Dooley S. IFN-gamma abrogates profibrogenic TGF-beta signaling in liver by targeting expression of inhibitory and receptor Smads. *J Hepatol* (2007) 46(2):295–303.  
<https://doi.org/10.1016/j.jhep.2006.09.014>
38. Dhanda AD, Felmler D, Boeira P, Moodley P, Tan H, Scalioni LDP. Patients with moderate to severe COVID-19 have an impaired cytokine response with an exhausted and senescent immune phenotype. *Immunobiology* (2022) 227(2):152185  
<https://doi.org/10.1016/j.imbio.2022.152185>
39. Elizabeth SC, Williams P, Martins TB, Hill HR, Coiras M, Shah KS. Plasma cytokine levels reveal deficiencies in IL-8 and gamma interferon in Long-COVID (2022) *medRxiv* (preprint Oct 5 2022) <https://doi.org/10.1101/2022.10.03.22280661>
40. Castro F, Cardoso AP, Gonçalves RM, Serre K, Oliveira MJ, Interferon-Gamma at the Crossroads of Tumor Immune Surveillance or Evasion *Frontiers in Immunology* (2018) 9. <https://doi.org/10.3389/fimmu.2018.00847>
41. Yoshimatsu Y, Wakabayashi I, Kimuro S. TNF- $\alpha$  enhances TGF- $\beta$ -induced endothelial-to-mesenchymal transition via TGF- $\beta$  signal augmentation. *Cancer Sci.* (2020) 111: 2385–99. <https://doi.org/10.1111/cas.14455>
42. Borthwick LA, Gardner A, De Soyza A. Transforming Growth Factor- $\beta$ 1 (TGF- $\beta$ 1) Driven Epithelial to Mesenchymal Transition (EMT) is Accentuated by Tumour Necrosis Factor  $\alpha$  (TNF $\alpha$ ) via Crosstalk Between the SMAD and NF- $\kappa$ B Pathways. *Cancer Microenvironment* (2012) 5:45–57 <http://dx.doi.org/10.1007/s12307-011-0080-9>

43. Swank Z, Senussi Y, Manickas-Hill Z, Yu XG, Li JZ, Alter G. Persistent Circulating Severe Acute Respiratory Syndrome Coronavirus 2 Spike Is Associated With Post-acute Coronavirus Disease 2019 Sequelae, *Clinical Infectious Diseases* (2022), ciac722 <https://doi.org/10.1093/cid/ciac722>
44. Mouton A, Blanc F, Gros A. Sex ratio in dementia with Lewy bodies balanced between Alzheimer's disease and Parkinson's disease dementia: a cross-sectional study. *Alz Res Therapy* (2018) 10:92. <https://doi.org/10.1186/s13195-018-0417-4>.
45. Reekes TH, Higginson CI, Ledbetter CR. Sex specific cognitive differences in Parkinson disease. *npj Parkinsons Dis.* (2020) 6:7. <https://doi.org/10.1038/s41531-020-0109-1>
46. de Souza HDN, Oliveira A, Ulrich H. Bradykinin reverts contralateral rotation and restores dopamine levels in an animal model of Parkinson disease. *11th World Congress on Neurology and Therapeutics* March 27-29, 2017 Madrid, Spain <https://www.iomcworld.org/conference-abstracts-files/2155-9562-C1-047-007.pdf>
47. Manson JM, Sammel MD, Freeman EW, Grisso JA. Racial differences in sex hormone levels in women approaching the transition to menopause. *Fertil Steril.* (2001) 75(2):297-304. [https://doi.org/10.1016/s0015-0282\(00\)01723-4](https://doi.org/10.1016/s0015-0282(00)01723-4)
48. Pepino MY, Finkbeiner S, Beauchamp GK, MMennella JA., Obese Women Have Lower Monosodium Glutamate Taste Sensitivity and Prefer Higher Concentrations Than Do Normal-weight Women. *Obesity* (2010) 18: 959-965. <https://doi.org/10.1038/oby.2009.493>
49. Bautista RJH, Mahmoud AM, Königsberg M, López NE, Guerrero D. Obesity: Pathophysiology, monosodium glutamate-induced model and anti-obesity medicinal plants, *Biomedicine & Pharmacotherapy* (2019) 111:503-516 <https://doi.org/10.1016/j.biopha.2018.12.108>
50. Bahadoran Z, Mirmiran P, Ghasemi A. Monosodium Glutamate (MSG)-Induced Animal Model of Type 2 Diabetes. In: Guest, P. (eds) Pre-Clinical Models. *Methods in Molecular Biology* (2019) vol 1916. Humana Press, New York, NY. [https://doi.org/10.1007/978-1-4939-8994-2\\_3](https://doi.org/10.1007/978-1-4939-8994-2_3)
51. Tann C, Chong H, Tan E. Getting 'Smad' about obesity and diabetes. *Nutr & Diabetes* (2012)/2, e29. <https://doi.org/10.1038/nutd.2012.1>
52. He Z, Yang Y, Xing Z. Intraperitoneal injection of IFN- $\gamma$  restores microglial autophagy, promotes amyloid- $\beta$  clearance and improves cognition in APP/PS1 mice. *Cell Death Dis* (2020) 11:440 <https://doi.org/10.1038/s41419-020-2644-4>
53. Zheng C, Zhou XW, Wang JZ. The dual roles of cytokines in Alzheimer's disease: update on interleukins, TNF- $\alpha$ , TGF- $\beta$  and IFN- $\gamma$ . *Transl Neurodegener* (2016) 5:7. <https://doi.org/10.1186/s40035-016-0054-4>
54. Shen L. Associations between B Vitamins and Parkinson's Disease. *Nutrients* (2015) 7(9):7197-7208. <https://doi.org/10.3390/nu7095333>
55. Ma H, Yan J, Sun W, Jiang M, Zhang Y. Melatonin Treatment for Sleep Disorders in Parkinson's Disease: A Meta-Analysis and Systematic Review. *Front Aging Neurosci* (2022) 4;14:784314. <https://www.frontiersin.org/articles/10.3389/fnagi.2022.784314/full>
56. Müller N, Myint A-M, Schwarz MJ. The impact of neuroimmune dysregulation on neuroprotection and neurotoxicity in psychiatric disorders - relation to drug treatment. *Dialogues in Clinical Neuroscience* (2009) 11(3):319-332. <http://dx.doi.org/10.31887/DCNS.2009.11.3/nmueller>

57. Ngai P, McCormick S, Small C, Zhang X, Zganiacz A, Aoki N, Gamma interferon responses of CD4 and CD8 T-cell subsets are quantitatively different and independent of each other during pulmonary Mycobacterium bovis BCG infection. *Infect Immun* (2007) 75(5):2244-52. <https://doi.org/10.1128/IAI.00024-07>
58. Williams GP, Schonhoff AM, Jurkuvenaite A, Gallups NJ, Standaert DG, Harms AS, CD4 T cells mediate brain inflammation and neurodegeneration in a mouse model of Parkinson's disease. *Brain* (2021) 144(7):2047–2059 <https://doi.org/10.1093/brain/awab103>
59. Shen, XR., Geng, R., Li, Q. ACE2-independent infection of T lymphocytes by SARS-CoV-2. *Sig Transduct Target Ther* (2022) 7:83 <https://doi.org/10.1038/s41392-022-00919-x>
60. Contaldi E, Magistrelli L, Comi C. T Lymphocytes in Parkinson's Disease. *Journal of Parkinson's Disease* (2022) 12(s1):S65 – S74. <https://doi.org/10.3233/JPD-223152>
61. Ferrari DP, Bortolanza NM, Del Bel E. Interferon- $\gamma$  Involvement in the Neuroinflammation Associated with Parkinson's Disease and L-DOPA-Induced Dyskinesia *Neurotox Res* (2022) 39:705–719. <https://doi.org/10.1007/s12640-021-00345-x>
62. Gulzar N, Copeland KF. CD8+ T-cells: function and response to HIV infection (2004) *Curr HIV Res* 2(1):23-37. <https://doi.org/10.2174/1570162043485077>
63. Roff SR, Noon-Song EN, Yamamoto JK. The Significance of Interferon- $\gamma$  in HIV-1 Pathogenesis, Therapy, and Prophylaxis. (2014) *Front Immunol* 4:498. <https://doi.org/10.3389/fimmu.2013.00498>
64. Koutsilieri, E., Sopper, S., Scheller, C. et al. Parkinsonism in HIV dementia (2002) *J Neural Transm* 109:767–775 <https://doi.org/10.1007/s007020200063>
65. Lees JR. Interferon gamma in autoimmunity: A complicated player on a complex stage. *Cytokine* (2015) 74(1):18-26. <https://doi.org/10.1016/j.cyto.2014.10.014>
66. Chohan H, Senkevich K, Patel RK, Bestwick JP, Jacobs BM. Type 2 Diabetes as a Determinant of Parkinson's Disease Risk and Progression. *Mov Disord* (2021) 36(6):1420-1429. <https://doi.org/10.1002/mds.28551>
67. De Pablo-Fernández E, Warner T. Association between diabetes and subsequent Parkinson disease: A record-linkage cohort study. *Neurology* (2019) 92(19):925-926 <https://doi.org/10.1212/WNL.0000000000005771>
68. Kalra S, Kalra B, Agrawal N. Understanding diabetes in patients with HIV/AIDS. *Diabetol Metab Syndr* (2011) 3, 2. <https://doi.org/10.1186/1758-5996-3-2>
69. Swaroop J, Rajarajeswari D, Naidu J. Association of TNF- $\alpha$  with insulin resistance in Type 2 diabetes mellitus. *The Indian journal of medical research* (2012) 135. 127-30. <https://doi.org/10.4103%2F0971-5916.93435>
70. Sudar-Milovanovic E, Gluvic Z, Obradovic M, Zaric B, Isenovic ER. Tryptophan Metabolism in Atherosclerosis and Diabetes. *Curr Med Chem* (2022) 29(1):99-113 <https://doi.org/10.2174/0929867328666210714153649>
71. Murray MF. Tryptophan depletion and HIV infection: a metabolic link to pathogenesis. *Lancet Infect Dis* (2003) 3(10):644-52 [https://doi.org/10.1016/s1473-3099\(03\)00773-4](https://doi.org/10.1016/s1473-3099(03)00773-4)
72. Sawada M, Imamura K, Nagatsu T. Role of cytokines in inflammatory process in Parkinson's disease. *J Neural Transm Suppl.* 2006;(70):373-81. [https://doi.org/10.1007/978-3-211-45295-0\\_57](https://doi.org/10.1007/978-3-211-45295-0_57)
73. Hu HH, Chen DQ, Wang YN, Feng YL, Cao G, Vaziri ND. New insights into TGF- $\beta$ /Smad signaling in tissue fibrosis. *Chem Biol Interact* (2018) 292:76-83. <https://doi.org/10.1016/j.cbi.2018.07.008>

74. Town T, Laouarn Y, Pittenger C. Blocking TGF- $\beta$ -Smad2/3 innate immune signaling mitigates Alzheimer-like pathology. *Nat Med* (2008) 681–687. <https://doi.org/10.1038/nm1781>
75. Yang C, Xu P. The role of transforming growth factor  $\beta$ 1 /Smad pathway in Alzheimer's disease inflammation pathology. *Mol Biol Rep* (2023) 50:777–788. <https://doi.org/10.1007/s11033-022-07951-8>
76. Xu L, Pan CL, Wu XH, Song JJ, Meng P, Li L. Inhibition of Smad3 in macrophages promotes A $\beta$  efflux from the brain and thereby ameliorates Alzheimer's pathology. *Brain Behav Immun* (2021) 95:154-167. <https://doi.org/10.1016/j.bbi.2021.03.013>
77. Tee JK, Peng F, Tan YL, Yu B, Ho HK. Magnesium Isoglycyrrhizinate Ameliorates Fibrosis and Disrupts TGF- $\beta$ -Mediated SMAD Pathway in Activated Hepatic Stellate Cell Line LX2. *Front Pharmacol* (2018) 9:1018. <https://doi.org/10.3389/fphar.2018.01018>
78. Luo X, Deng Q, Xue Y, Zhang T, Wu Z, Peng H. Anti-Fibrosis Effects of Magnesium Lithospermate B in Experimental Pulmonary Fibrosis: By Inhibiting TGF- $\beta$ RI/Smad Signaling. *Molecules* (2021) 26(6):1715. <https://doi.org/10.3390/molecules26061715>
79. Ulloa L, Doody J, Massagué J. Inhibition of transforming growth factor- $\beta$ /SMAD signalling by the interferon- $\gamma$ /STAT pathway. *Nature* (1999) 397, 710–713. <https://doi.org/10.1038/17826>
80. Vijayaraghavaavaava AKR. Alteration of Interferon Gamma (IFN- $\gamma$ ) in Human Plasma with Graded Physical Activity. *J Clin of Diagn Res* (2014) 8(6):BC05-BC07. <https://doi.org/10.7860/2FJCDR%2F2014%2F9502.4440>
81. Li X, Wang K. Effects of moderate-intensity endurance exercise on angiotensin II and angiotensin II type I receptors in the rat heart. *Mol Med Rep* (2017) 16(3):2439-2444. <https://doi.org/10.3892%2Fmmr.2017.6864>
82. Fernandez-Castelo S, Arzt ES, Pesce A, Criscuolo ME, Diaz A, Finkielman S. Angiotensin II regulates interferon-gamma production. *J Interferon Res* (1987) 7(3):261-8. <https://doi.org/10.1089/jir.1987.7.261>
83. Gebreyesus K, Kilbourne EJ, Sabban EL. Bradykinin elevates tyrosine hydroxylase and dopamine beta-hydroxylase mRNA levels in PC12 cells. *Brain Research* (1993) 608(2):345-348. [https://doi.org/10.1016/0006-8993\(93\)91477-a](https://doi.org/10.1016/0006-8993(93)91477-a)
84. Hu J, Igarashi A, Kamata M, Nakagawa H. Angiotensin-converting enzyme degrades Alzheimer amyloid beta-peptide (A $\beta$ ); retards A $\beta$  aggregation, deposition, fibril formation; and inhibits cytotoxicity. *J Biol Chem* (2001) 276(51):47863-8. <https://doi.org/10.1074/jbc.M104068200>
85. Rigat B, Hubert C, Alhenc-Gelas F, Cambien F, Corvol P, Soubrier F. An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *J Clin Invest* (1990) 86(4):1343-6. <https://doi.org/10.1172/JCI114844>
86. Narain Y, Yip A, Murphy T, Brayne C, Easton D, Evans JG. The ACE gene and Alzheimer's disease susceptibility. *J Med Genet* (2000) 37(9):695-7. <https://jmg.bmj.com/content/37/9/695>
87. Brown NJ, Blais Jr C, Gandhi SK, Albert A. ACE Insertion/Deletion Genotype Affects Bradykinin Metabolism. *Journal of Cardiovascular Pharmacology* (1998) 32(3):p 373-377 <https://doi.org/10.1097/00005344-199809000-00006>
88. Singh PK, Chen ZL, Ghosh D, Strickland S, Norris EH. Increased plasma bradykinin level is associated with cognitive impairment in Alzheimer's patients. *Neurobiol Dis* (2020) 139:104833. <https://doi.org/10.1016/j.nbd.2020.104833>
89. Douillet CD, Velarde V, Christopher JT, Mayfield RK, Trojanowska ME, Jaffa AA. Mechanisms by which bradykinin promotes fibrosis in vascular smooth muscle cells: role of TGF-beta and MAPK. *Am J Physiol Heart Circ Physiol* (2000) 279(6):H2829-37. <https://doi.org/10.1152/ajpheart.2000.279.6.h2829>

90. Bandera EV, Qin B, Lin Y, Zeinomar N, Xu B, Chanumolu D. Association of Body Mass Index, Central Obesity, and Body Composition With Mortality Among Black Breast Cancer Survivors. *JAMA Oncol* (2021) 7(8):1–10.  
<https://doi.org/10.1001/jamaoncol.2021.1499>
91. Giaquinto AN, Miller KD, Tossas KY, Winn RA, Jemal A, Siegel RL. Cancer statistics for African American/Black People 2022. *CA Cancer J Clin* (2022) 72(3):202–229. <https://doi.org/10.3322/caac.21718>
92. Jorgovanovic, D., Song, M., Wang, L. et al. Roles of IFN- $\gamma$  in tumor progression and regression: a review. *Biomark Res* (2020) 8:49. <https://doi.org/10.1186/s40364-020-00228-x>
93. Liang J, Shang Y. Estrogen and cancer. *Annu Rev Physiol.* (2013) 75:225–40. <https://doi.org/10.1146/annurev-physiol-030212-183708>
94. Hicks BM, Filion KB, Yin H, Sakr L, Udell JA, Azoulay L. Angiotensin converting enzyme inhibitors and risk of lung cancer: population based cohort study. *BMJ* (2018) 363:k4209. <https://doi.org/10.1136%2Fbmj.k4209>
95. Yu HS, Lin TH, Tang CH. Bradykinin enhances cell migration in human prostate cancer cells through B2 receptor/PKC $\delta$ /c-Src dependent signaling pathway. *Prostate* (2013) 73(1):89–100. <https://doi.org/10.1002/pros.22544>
96. Smith AD, Refsum H, Bottiglieri T, Fenech M, Hooshmand B, McCaddon A. Homocysteine and Dementia: An International Consensus Statement. *J Alzheimers Dis* (2018) 62(2):561–570. <https://doi.org/10.3233/jad-171042>
97. Jang Y, Yoon H, Park NS, Rhee MK, Chiriboga DA. Asian Americans' concerns and plans about Alzheimer's disease: The role of exposure, literacy and cultural beliefs. *Health Soc Care Community* (2018) 26(2):199–206.  
<https://doi.org/10.1111/hsc.12509>
98. Rojo-Sebastián A, González-Robles C, García de Yébenes J. Vitamin B6 Deficiency in Patients With Parkinson Disease Treated With Levodopa/Carbidopa. *Clin Neuropharmacol.* (2020) 43(5):151–157.  
<https://doi.org/10.1097/wnf.0000000000000408>
99. Politis M, Loane C. Serotonergic dysfunction in Parkinson's disease and its relevance to disability. *ScientificWorldJournal.* (2011) 11:1726–34. <https://doi.org/10.1100/2011/172893>
100. Kennedy DO. B Vitamins and the Brain: Mechanisms, Dose and Efficacy--A Review. *Nutrients* (2016) 8(2):68.  
<https://doi.org/10.3390/nu8020068>
101. Tyagi N, Sedoris KC, Steed M. Mechanisms of homocysteine-induced oxidative stress. *American Journal of physiology. Heart and Circulatory Physiology* (2005) 289(6):H2649–56. <https://doi.org/10.1152/ajpheart.00548.2005>
102. Keskin A, U Ustun G, Aci R, Duran U. Homocysteine as a marker for predicting disease severity in patients with COVID-19. *Biomark Med* (2022) 16(7):559–568. <https://doi.org/10.2217/bmm-2021-0688>
103. McCaddon A, Regland B. COVID-19: A methyl-group assault? *Med Hypotheses* (2021) 149:110543.  
<https://doi.org/10.1016/j.mehy.2021.110543>
104. Fan X, Zhang L, Li H, Chen G, Qi G, Ma X. Role of homocysteine in the development and progression of Parkinson's disease. *Ann Clin Transl Neurol* (2020) 7(11):2332–2338. <https://doi.org/10.1002/acn3.51227>
105. Farkas M, Keskitalo S, Smith DE, Bain N, Semmler A, Ineichen B. Hyperhomocysteinemia in Alzheimer's disease: the hen and the egg? *J Alzheimers Dis* (2013) 33(4):1097–104. <http://dx.doi.org/10.3233/JAD-2012-121378>
106. Gilfix BM. Vitamin B12 and homocysteine. *CMAJ* (2005) 173 (11):1360 <https://doi.org/10.1503/cmaj.1050170>



107. Morris MC, Schneider JA, Tangney CC. Thoughts on B-vitamins and dementia. *J Alzheimers* (2006) 9(4):429-33.  
<https://doi.org/10.3233/jad-2006-9409>
108. Łoboś P, Regulska-Iłow B. Link between methyl nutrients and the DNA methylation process in the course of selected diseases in adults. *Rocz Panstw Zakl Hig* (2021) 72(2):123-136. <https://pubmed.ncbi.nlm.nih.gov/34114759/>
109. Hadtstein F, Vrolijk M. Vitamin B-6-Induced Neuropathy: Exploring the Mechanisms of Pyridoxine Toxicity. *Adv Nutr* (2021) 12(5):1911-1929. <https://doi.org/10.1093/advances/nmab033>
110. Noah L, Dye L, Bois De Fer B, Mazur A, Pickering G, Pouteau E. Effect of magnesium and vitamin B6 supplementation on mental health and quality of life in stressed healthy adults: Post-hoc analysis of a randomised controlled trial. *Stress Health* (2021) 37(5):1000-1009. <https://doi.org/10.1002/smi.3051>
111. Malouf R, Grimley Evans J. The effect of vitamin B6 on cognition. *Cochrane Database Syst Rev*. 2003;(4):CD004393.  
<https://doi.org/10.1002/14651858.cd004393>
112. Chambers P, Antioxidants and Long Covid. *Open Access Library Journal* (2022) 9:e9414.  
<https://www.scirp.org/journal/paperinformation.aspx?paperid=120821>
113. Xu J, Patassini S, Begley P, Church S, Waldvogel HJ, Faull RLM. Cerebral deficiency of vitamin B5 (d-pantothenic acid; pantothenate) as a potentially-reversible cause of neurodegeneration and dementia in sporadic Alzheimer's disease. *Biochem Biophys Res Commun* (2020) 527(3):676-681. <https://doi.org/10.1016/j.bbrc.2020.05.015>
114. Scholefield M, Church SJ, Xu J, Patassini S, Hooper NM, Unwin RD. Substantively Lowered Levels of Pantothenic Acid (Vitamin B5) in Several Regions of the Human Brain in Parkinson's Disease Dementia. *Metabolites* (2021) 11(9):569.  
<https://doi.org/10.3390/metabo11090569>
115. Leonardi R, Jackowski S. Biosynthesis of Pantothenic Acid and Coenzyme A. *EcoSal Plus* (2007) 2(2):10.1128/ecosalplus.3.6.3.4. <https://doi.org/10.1128/ecosalplus.3.6.3.4>
116. Alzheimer's Association Calcium Hypothesis Workgroup. Calcium Hypothesis of Alzheimer's disease and brain aging: A framework for integrating new evidence into a comprehensive theory of pathogenesis. *Alzheimers Dement* (2017) 13(2):178-182.e17. <https://doi.org/10.1016/j.jalz.2016.12.006>
117. Magnesium in Alzheimer's disease Section 3 of the book Magnesium in the Central Nervous System (2011). Vink R, Nechifor M, editors. Adelaide (AU): University of Adelaide Press <https://www.ncbi.nlm.nih.gov/books/NBK507256/>
118. Surmeier DJ, Schumacker PT, Guzman JD, Ilijic E, Yang B, Zampese E. Calcium and Parkinson's disease. *Biochem Biophys Res Commun* (2017) 483(4):1013-1019. <https://doi.org/10.1016/j.bbrc.2016.08.168>
119. Miyake Y, Tanaka K, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y. Fukuoka Kinki Parkinson's Disease Study Group. Dietary intake of metals and risk of Parkinson's disease: a case-control study in Japan. *J Neurol Sci.* (2011)306(1-2):98-102.  
<https://doi.org/10.1016/j.jns.2011.03.035>
120. Wang L, Zhou C, Yu H, Hao L, Ju M, Feng W. Vitamin D, Folic Acid and Vitamin B12 Can Reverse Vitamin D Deficiency-Induced Learning and Memory Impairment by Altering 27-Hydroxycholesterol and S-Adenosylmethionine. *Nutrients* (2023) 15(1):132. <https://doi.org/10.3390/nu15010132>
121. Chambers P. Ca:Mg + D, the Shield that Interdicts the Crown Viruses and Vaccines. *Open Access Library Journal* (2022) 9(9):1-24. <https://www.scirp.org/journal/paperinformation.aspx?paperid=119926>



122. Fonseca SC, Rivas I, Romaguera D, et al. Association between consumption of fermented vegetables and COVID-19 mortality at a country level in Europe. *medRxiv* (2020 preprint) <https://doi.org/10.1101/2020.07.06.20147025>
123. Kiana TV, Nikan Z, Armin E, Mohammad MS, Parnia M, Pari M. Association of fruits, vegetables, and fiber intake with COVID-19 severity and symptoms in hospitalized patients: A cross-sectional study *Frontiers in Nutrition* (2022) vol 9 2022 <https://doi.org/10.3389/fnut.2022.934568>
124. Wastyk HC, Fragiadakis GK, Perelman D, Dahan D, Merrill BD, Yu FB. Gut-microbiota-targeted diets modulate human immune status. *Cell* (2021) 184(16):4137-4153.e14. <https://doi.org/10.1016/j.cell.2021.06.019>
125. Kim YH, Jung KI, Song CH. Effects of Serum Calcium and Magnesium on Heart Rate Variability in Adult females. *Biological Trace Element Research* (2012) 150:116-122. <https://doi.org/10.1007/s12011-012-9518-2>
126. Haensel, A., Mills, P.J., Nelesen, R.A., Ziegler, M.J. and Dimsdale, J.E. The Relationship between Heart Rate Variability and Inflammatory Markers in Cardiovascular Diseases. *Psychoneuroendocrinology* (2008) 33:1305-1312. <https://doi.org/10.1016/j.psyneuen.2008.08.007>.
127. Schiopu C, Ștefănescu G, Diaconescu S, Bălan GG, Gimiga N, Rusu E. Magnesium Orotate and the Microbiome–Gut–Brain Axis Modulation: New Approaches in Psychological Comorbidities of Gastrointestinal Functional Disorders. *Nutrients* (2022) 14(8):1567. <https://doi.org/10.3390%2Fnu14081567>
128. Winther G, Pyndt Jørgensen B, ElfvingmB, Nielsen D, Kihl P, Lund S. Dietary magnesium deficiency alters gut microbiota and leads to depressive-like behaviour. *Acta Neuropsychiatrica* (2015) 27(3):168-176. <https://doi.org/10.1017/neu.2015.7>
129. Luo J, Zhang C, Zhao Q. Dietary calcium and magnesium intake and risk for incident dementia: The Shanghai Aging Study. *Alzheimer's & Dementia* (2022) 8(1):e12362. <https://doi.org/10.1002/trc2.12362>
130. Zhu X. Ca:Mg Ratio, APOE Cytosine Modifications, and Cognitive Function: Results from a Randomized Trial. *Journal of Alzheimer's Disease* (2020) 75(1):85-98. <https://doi.org/10.3233/jad-191223>
131. Kong LM, Liao CG, Chen L, Yang HS, Zhang SH, Zhang Z. Promoter hypomethylation up-regulates CD147 expression through increasing Sp1 binding and associates with poor prognosis in human hepatocellular carcinoma. *J Cell Mol Med* (2011)15(6):1415-28. <https://doi.org/10.1111%2Fj.1582-4934.2010.01124.x>
132. Liao CG, Liang XH, Ke Y. Active demethylation upregulates CD147 expression promoting non-small cell lung cancer invasion and metastasis. *Oncogene* (2022) 41:1780-1794. <https://doi.org/10.1038/s41388-022-02213-0>
133. Ehrlich M. DNA hypomethylation in cancer cells. *Epigenomics* (2009) 1(2):239-59. <https://doi.org/10.2217%2Fepi.09.33>
134. Kulis M, Esteller M. DNA methylation and cancer. *Adv Genet* (2010) 70:27-56. <https://doi.org/10.1016/B978-0-12-380866-0.60002-2>.
135. Johnson AA, Akman K, Calimport SR, Wuttke D, Stolzing A, de Magalhães JP. The role of DNA methylation in aging, rejuvenation, and age-related disease. *Rejuvenation Res* (2012) 15(5):483-94. <https://doi.org/10.1089%2Ffrej.2012.1324>
136. Song H, Yang J, Yu W. Promoter Hypomethylation of TGFBR3 as a Risk Factor of Alzheimer's Disease: An Integrated Epigenomic-Transcriptomic Analysis. *Front Cell Dev Biol* (2022) 9:825729. <https://doi.org/10.3389/fcell.2021.825729>
137. Chouliaras L, Kumar GS, Thomas AJ, Lunnon K, Chinnery PF, O'Brien JT. Epigenetic regulation in the pathophysiology of Lewy body dementia. *Prog Neurobiol* (2020) 192:101822. <https://doi.org/10.1016/j.pneurobio.2020.101822>
138. Tulloch J, Leong L, Chen S, Keene CD, Millard SP, Shutes-David A. APOE DNA methylation is altered in Lewy body dementia. *Alzheimers Dement* (2018) 14(7):889-894. <https://doi.org/10.1016/j.jalz.2018.02.005>

139. Harhangi BS, de Rijk MC, van Duijn CM, Van Broeckhoven C, Hofman A, Breteler MM. APOE and the risk of PD with or without dementia in a population-based study. *Neurology* (2000) 54(6):1272-6.  
<https://doi.org/10.1212/wnl.54.6.1272>
140. Foraker J, Millard SP, Leong L, Thomson Z, Chen S, Keene CD, Bekris LM, Yu CE. The APOE Gene is Differentially Methylated in Alzheimer's Disease. *J Alzheimers Dis* (2015) 48(3):745-55. <https://doi.org/10.3233/jad-143060>
141. Suchy-Dicey A, Howard B, Longstreth WT Jr, Reiman EM, Buchwald D. APOE genotype, hippocampus, and cognitive markers of Alzheimer's disease in American Indians: Data from the Strong Heart Study. *Alzheimers Dement* (2022) 18(12):2518-2526. <https://doi.org/10.1002/alz.12573>
142. Workinger, J.L., Doyle, R.P. and Bortz, J. Challenges in the Diagnosis of Magnesium Status. *Nutrients* (2018) 10(1202).  
<https://doi.org/10.3390/nu10091202>
143. Nakayama Y, Ueda S, Okuda S. Molecular mechanism underlying the renoprotective action of vitamin D. *Circ J* (2014) 78(3):599-600. <https://doi.org/10.1253/circj.cj-14-0058>.
144. Wimalawansa SJ. Rapidly Increasing Serum 25(OH)D Boosts the Immune System, against Infections—Sepsis and COVID-19. *Nutrients* (2022) 14(14):2997. <https://doi.org/10.3390/nu14142997>
145. Xian X, Wang Y, Zheng J. COVID-19 and Alzheimer's disease: how one crisis worsens the other. *Transl Neurodegener* (2021) 10:15. <https://doi.org/10.1186/s40035-021-00237-2>
146. Bikle DD. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem Biol* (2014) 21(3):319-29.  
<https://doi.org/10.1016%2Fj.chembiol.2013.12.016>
147. Thye A Y-Y, Tan L T-H, Law J W-F, Pusparajah P, Letchumanan V. Long COVID-19: psychological symptoms in COVID-19 and probiotics as an adjunct therapy. *Progress In Microbes & Molecular Biology* (2022).  
<https://doi.org/10.36877/pmmb.a0000267>
148. Gutiérrez-Castrellón P, Gandara-Martí T, Abreu Y Abreu AT, Nieto-Rufino CD, López-Orduña E. Probiotic improves symptomatic and viral clearance in Covid19 outpatients: a randomized, quadruple-blinded, placebo-controlled trial. *Gut Microbes* (2022) 14(1):2018899. <https://doi.org/10.1080/19490976.2021.2018899>
149. Yong SJ, Tong T, Chew J, Lim WL. Antidepressive Mechanisms of Probiotics and Their Therapeutic Potential. *Front Neurosci* (2020) 13:1361. <https://doi.org/10.3389%2Ffnins.2019.01361>
150. Shrestha NK, Burke PC, Nowacki AS, Simon JF, Hagen A, Gordon SM. Effectiveness of the Coronavirus Disease 2019 (COVID-19) Bivalent Vaccine *medRxiv* (2023 preprint) <https://www.medrxiv.org/content/10.1101/2022.12.17.22283625v1>
151. McCracken IR, Saginc G, He L, Huseynov A, Daniels A, Fletcher S. Lack of Evidence of Angiotensin-Converting Enzyme 2 Expression and Replicative Infection by SARS-CoV-2 in Human Endothelial Cells. *Circulation* (2021) 143(8):865-868.  
<https://doi.org/10.1161/CIRCULATIONAHA.120.052824>
152. Ganier C, Du-Harpur X, Harun N, Wan B, Arthurs C, Luscombe NM. CD147 (BSG) but not ACE2 expression is detectable in vascular endothelial cells within single cell RNA sequencing datasets derived from multiple tissues in healthy individuals *bioRxiv* (2020 preprint) <https://doi.org/10.1101/2020.05.29.123513>
153. Hsu CC, Cheng CH, Hsu CL, Lee WJ, Huang SC, Huang YC. Role of vitamin B6 status on antioxidant defenses, glutathione, and related enzyme activities in mice with homocysteine-induced oxidative stress. *Food Nutr Res* (2015) 59:25702. <https://doi.org/10.3402%2Ffnr.v59.25702> .

- 
154. Gibson CC, Davis CT, Zhu W, Bowman-Kirigin JA, Walker AE, Tai Z. Dietary Vitamin D and Its Metabolites Non-Genomically Stabilize the Endothelium. *PLoS One* (2015) 10(10):e0140370. <https://doi.org/10.1371/journal.pone.0140370>
  155. Tsubokawa M, Nishimura M, Mikami T, Ishida M, Hisada T, Tamada Y. Association of Gut Microbial Genera with Heart Rate Variability in the General Japanese Population: The Iwaki Cross-Sectional Research Study. *Metabolites* (2022) 12(8):730. <https://doi.org/10.3390/metabo12080730>
  156. Brilla L, Teichler L, Hahn T, Freeman J, LiNY. Effect of Magnesium on Heart Rate Variability in Healthy Subjects. *FASEB J* (2010) 24: 917.3-917.3. [https://doi.org/10.1096/fasebj.24.1\\_supplement.917.3](https://doi.org/10.1096/fasebj.24.1_supplement.917.3)
  157. Wyller VB, Nguyen CB, Ludviksen JA, Mollnes TE. Transforming growth factor beta (TGF- $\beta$ ) in adolescent chronic fatigue syndrome. *J Transl Med* (2017) 15(1):245. <https://doi.org/10.1186/s12967-017-1350-1>
  158. Thanou A, Stavrakis S, Dyer JW, Munroe ME, James JA, Merrill JT. Impact of heart rate variability, a marker for cardiac health, on lupus disease activity. *Arthritis Res Ther* (2016) 18(1):197. <https://doi.org/10.1186/s13075-016-1087-x>