

LIVER FIBROSIS IN INDIVIDUALS WITH HEPATITIS C VIRUS GENOTYPE 1B FROM BRAZILIAN EASTERN AMAZON

Running Title: Liver Fibrosis and Hepatitis C virus in Eastern Amazon

Tatiane Lobato da Silva¹
Apio Ricardo Nazareth Dias¹
Juarez Antônio Simões Quaresma^{1,3,†}
Luiz Fábio Magno Falcão^{1,2,†*}

Author affiliations

¹ Health and Biological Centre, State of Pará University, 2623 Perebeui St, Belem, Pará 66087-670, Brazil.

²University of São Paulo, 455 Dr Arnaldo Ave, São Paulo, São Paulo 01246-903, Brazil.

³Tropical Medicine Centre, Federal University of Pará, 92 Generalissimo Deodoro Ave, Belém, Pará 66055-240, Brazil.

*Corresponding author: fabiofalcao@uepa.br

° These authors contributed equally for this study

† These authors share Senior authorship

* Corresponding author:

Dr. Luiz Fábio Magno Falcão.

2623 Perebeui St, Belem, PA 66087-670, Brazil

+55 (91) 3131-1761

E-mail: fabiofalcao@uepa.br

All data related to this study is available with correspondent author

ABSTRACT

INTRODUCTION Hepatitis C virus (HCV) has a high genetic diversity, with seven genotypes with 86 subtypes. This genetic variability confers persistence in the infection and escape of the immune system with evolution to cirrhosis and cancer. Environmental factors can contribute to different disease progression, being essential to assess the viral genotype in the infection and discuss the environmental particularities of Eastern Amazon, and the frequencies of Liver fibrosis between different HCV genotypes in patients living in a region of the Brazilian Eastern Brazilian Amazon. **METHODS** Consists in an observational cross-sectional study. Sociodemographic and clinical data of 76 individuals diagnosed with Hepatitis C between 2019 and 2020 in public health services were selected. Data collected was tabulated in Microsoft Excel 2010™ spreadsheets and analysed in GraphPad Prism 5.0™. Liver fibrosis was associated with genetic subtypes. **RESULTS** Subtype 1b was predominant (42.1%), followed by 1a (13%) and 3a (1.3%). 69.7% of participants had chronic hepatitis, with mild fibrosis (F1/F2) being the most prevalent (38.1%). Severe fibrosis was detected in 75% of individuals infected with the subtype 1b, that is associated with more severe disease. **CONCLUSIONS** We suggest further studies, to assess other communities in the region, as well as the monitoring of these patients with Liver Elastography to determine the disease evolution and its better management.

Keywords: Hepatitis C virus, Genetic, Chronic Hepatitis C, Liver, Elastography

INTRODUCTION

Hepatitis C virus (HCV) is a virus with a global infection incidence of 70 million worldwide¹. Brazil's northern region shows highest prevalence rate of this disease, and the states of Pará (Eastern Amazon) and Acre (Western Amazon) have the highest rates of this region, with 2% and 5.9%, respectively¹⁴. Complications will be developed by 55 - 85% of infected people. Liver Cirrhosis and hepatocellular carcinoma were responsible for approximately 290,000 deaths associated with the disease in the year of 2019^{2,3,4}. Chronic HCV is one of the main health problems globally, with highest rate of complications in comparison to other viral hepatitis.^{5, 6,7}.

HCV exhibits a high genetic diversity, with seven genotypes being described, subdivided into 86 subtypes. This genetic variability provides viral persistence, its mechanisms of immune escape, and increase the risk of evolution to cirrhosis and cancer. The determination of the virus genotype is important because it is related to clinical presentation and prognostic^{8,9,10}. The most prevalent HCV genotypes in Amazon region are related with development of Liver fibrosis that are associated with disease complications.

The Amazon population is exposed to environmental factors that would influence the diseases evolution. The effects of rainforest fires or agrobusiness actions directly impact in ecosystem structure, favouring the endemic diseases¹¹; vulnerability of populations with lack access of health services, basic sanitation, and adequate nutrition¹²; and genetic factors related to miscegenation of brazilian population must be considered. There is epidemiological evidence that the genotype of this virus associated with these environmental, and individual behaviours influence in the disease progression¹³.

There are few studies associating clinical data of Liver Fibrosis with the viral genotypic classification in Eastern Amazon. This study aimed to identify the degree of liver fibrosis and HCV genotypes in patients from a community located in the Eastern Brazilian Amazon.

MATERIAL AND METHODS

The study was approved by the Research Ethics Committee of Biologic and Health Centre, State of Pará University (opinion nº 4.478,618), being conducted in accordance with Helsinki Declaration for Research with human-beings, the data of participants was accessed from databases after authorization of Health authorities.

This was an observational cross-sectional study with quantitative, descriptive, and analytical characteristics. The sample consisted of patients residing in the city of Abaetetuba, Pará, Eastern Brazilian Amazon, evaluated in the public health services between years 2019 and 2020.

Data were collected from 103 patients aged ≥ 18 years. Twenty-seven patients were excluded from the study due to non-performance of viral load and/or genotyping tests (n=24) and co-infection (n=3): Human Immunodeficiency Virus (HIV) (n=1) and Hepatitis B Virus (HBV) (n=2); resulting in a final sample of 76 clinical records analysed.

The clinical and epidemiological data were extracted from the patients' clinical records of viral hepatitis investigation from Notifiable Diseases Notification System (SINAN) and Sexual transmitted infectious diseases department (CTA/SAE). This database contains (i) the Notification Form for viral hepatitis which contains the sociodemographic data, (ii) the results of viral load tests, (iii)

the results of viral genotyping, and (iv) the report of Liver Elastography exam (Fibroscan™, Echosens SA, Paris, France).

The Fibroscan™ transient elastography consists in a non-invasive exam that evaluates the hepatic parenchyma at 25-65 mm of the skin in the same intercostal space where the liver biopsy is performed. The patients were classified in the following fibrosis stages: F0 < 7.1 kPa (normal liver), F1 (mild fibrosis), F2 (mild to moderate fibrosis), F3 (severe fibrosis), and F4 (cirrhosis)^{31, 32, 33}.

The data was organized in Microsoft Excel 2010™ (Microsoft Software Inc., Richmond, USA) spreadsheets and analysed using GraphPad Prism 5.0 software (GraphPad Software Inc., San Diego, USA). Data with normal distribution were analysed using Student's T-test, while data without normal distribution were evaluated using Chi square or Fisher's exact test. An alpha-level of 0.05 was adopted to reject the null hypothesis.

RESULTS

Sociodemographic profile of study participants shows an predominance of male patients (67.1%) and those aged ≥ 60 years (53.9%), who are married or in stable union (69.7%), self-declared Dark-skinned (73.7%), residing in the urban area (85.5%). with more than nine years of education (59.2%) (**Table 1**).

Clinical profiles of the cohort indicated that the degree of fibrosis measured were classified as mild /mild to moderate fibrosis in 38.1% of cases (F1[14, 18.4%] /F2[15, 19.7%]). There are a high prevalence of fibrosis degree classified as cirrhosis [F4(13, 17.1%)]. The most prevalent genotype was genotype 1 (55.3%), followed by genotype 3 (28.9%). The most expressive sub genotype was 1b (42.1%) (**Table 2**).

Genotypes and sub genotypes of the participants were associated with the degree of fibrosis. There is a difference between the different degrees of fibrosis related to their sub genotypes. Participants without fibrosis had type 1a sub genotype predominantly (3, 60%), while those with severe fibrosis (F3)/ cirrhosis (F4) (15, 75%), and mild (F1)/ mild to moderate (F2) fibrosis (9, 60%) had 1b sub genotype (**Figure**). There was no significant difference in the genotype of patients according to the different degrees of fibrosis; thus, patients without fibrosis (F0) (5; 62.5%), with mild (F1) / mild to moderate (F2) fibrosis (15; 55.6%), and with severe fibrosis (F3) / cirrhosis (F4) (16; 80%) had mostly type 1 genotype, between individuals infected with subtype 1b (15; 75%) are diagnosed with severe fibrosis (F3)/ cirrhosis (F4) (**Table 3**).

DISCUSSION

In this study, most individuals with HCV were male and aged ≥ 60 years. 59.2% of the participants had 9 years or more of education. There was a predominance of dark-skinned (73.7%). With regard to transmission, 60.5% of patients stated that it occurred by blood transfusion. Genotype 1 had the highest prevalence rate (55.3%), type 3 was the second most prevalent type (28.9%), followed by type 2 (2.6%).

Men are more vulnerable to the risks of hepatitis C, their biological, social, behavioural, and cultural issues, with turn them more vulnerable to HCV infection. They have a greater intensity of sexual activity and seek for health services with less frequency than women¹⁵. In a 2014 study, 50% of the patients had a complete high school education, which was not protective in terms of preventing HCV infection¹⁶.

The genetic heterogeneity of population may impact the frequency and distribution of polymorphisms¹⁷. Latin American population was composed by native americans, europeans, mostly from Spain and Portugal, and africans, that conferred a predominance of dark-skinned or black people in Brazilian and Amazonian population¹⁸. About 30 million people are living in Amazon region, including quilombolas, indigenous people and riverine communities, as well as people of urban centers¹¹. In this region, several people do not have adequate housing and basic sanitation conditions, and access to health services and adequate nutrition¹², it creates the conditions to maintenance and spread of infectious diseases, that can potentially modify the development of HCV infection.

The majority of study participants developed the chronic form of the disease. It is known that infection occurs mainly through contact with contaminated blood. Transfusion of blood and blood products was the main risk factor for hepatitis C transmission in the past, which resulted in a 10% risk of acquiring infection¹⁶, and the chronic form of the disease are the main clinical form of cases of hepatitis C, with > 60% of cases in almost all age groups analysed⁴. Furthermore, another factor that may be related to the chronicity of the disease is the occurrence in people with less access to information and health services, as this vulnerability is a risk factor for infection and not early diagnosis¹⁹.

The environmental and socioeconomic reality of Amazon region, linked to agrarian economy of region have an influence in evolution of infectious diseases¹¹. The contamination of water, soil, air, and food by mining activity, deforestation, and rainforest fires causes loss of biodiversity, alteration in the structure and function of ecosystem, contributing for climate changes, and impacts in the behaviour of disease-transmitting vectors²⁰. The interaction of

population with other endemic infectious diseases, such as Malaria, Dengue, Zika, Chikungunya, Yellow Fever, Chagas Disease, HIV, HTLV, and others, as well as the relationship between these pathogens and HCV would be related with the disease evolution in this population (**Figure**).

Genotype 1 had the highest prevalence rate (55.3%). Type 3 was the second most prevalent type (28.9%), followed by type 2 (2.6%) in the studied region. This result corroborates other studies conducted in the Brazilian Amazon and Latin American countries. Regarding sub-genotypes, the most prevalent subtype was 1b (42.1%). Studies carried out in other locations corroborate the results of this research, in the Eastern Amazon^{16, 21,22, 23}, Paraguay and Colombia^{24,25}, and recent studies from Northern, Southern and Eastern Europe have also shown the predominance of genotype 1^{26,27,28}.

Studies of genotypic distribution is important for a better understanding of the epidemiological profile and the clinical, diagnostic, and therapeutic measures to be adopted. The subtypes 1b and 3a are associated with more severe forms of the disease, including a major risk for development of hepatocellular carcinoma^{29,30}.

80% of the Individuals infected with HCV genotype 1 who had their degree of fibrosis measured were diagnosed with severe fibrosis/cirrhosis (F3/F4), as well as, 75% of the infected with sbtype 1b. It is known that the weighting of the fibrotic degree of the liver in patients with hepatitis C has a high value in terms of diagnosis and prognosis, since approximately 20% of patients with chronic disease can have progression to liver cirrhosis³¹. Liver fibrosis grade according to the liver elastography test, which is a non-invasive, validated diagnostic method used to identify and classify liver fibrosis, was classified as follows. F0

(without fibrosis) < 7.1 kPa; F1/F2 (mild/ mild to moderate fibrosis) 8.8–9.5; and F3/F4 (severe fibrosis/ cirrhosis) 9.6 to ≥ 12.5 ^{32, 33, 34}.

However, 18.4% (n=14) of the participants did not undergo an examination to verify the degree of fibrosis. These data are worrying, as it is of fundamental importance to determine the degree of fibrosis so that the patient can receive the suitable therapeutic measures and the most appropriate follow-up for their clinical condition. It is known that, even today, the gold standard for this determination is liver biopsy; and despite Liver transient elastography be an efficient non-invasive technique, have a high cost and is less available in public health centres^{32,35}.

CONCLUSION

The study revealed the high predominance of genotypes 1 and 3 and subtypes 1a, 1b, and 3a among the studied population, with a higher prevalence of severe liver fibrosis (F3) and cirrhosis (F4) between individuals infected with genotype 1, subtype 1b. These HCV genotypes are the most prevalent in the national territory, being associated with more aggressive clinical forms. The results of this study allow us to elucidate the genotypic behaviour of HCV and its association with liver fibrosis and cirrhosis in the Brazilian Eastern Amazon, which is known to be a region of high prevalence for the virus, and with environmental factors that would impacts in the evolution of diseases.

ACKNOWLEDGMENT

The authors thank all patients who agreed to voluntarily participate in the study and PAPQ program (UFPA).

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflict of interest.

AUTHORS' CONTRIBUTION

TLS, ARND, JASQ and LFMF conceived the project, wrote and reviewed the manuscript, performed the analysis of data and images. All authors reviewed and approved the article.

Tatiane Lobato da Silva.

Health and Biological Centre, State of Pará University, 2623 Perebeui St, Belem, Pará 66087-670, Brazil.

tatianelobato34@gmail.com

Apio Ricardo Nazareth Dias.

Health and Biological Centre, State of Pará University, 2623 Perebeui St, Belem, Pará 66087-670, Brazil.

apio.dias@uepa.br

Juarez Antônio Simões Quaresma.

Health and Biological Centre, State of Pará University, 2623 Perebeui St, Belem, Pará 66087-670, Brazil.

Tropical Medicine Centre, Federal University of Pará, 92 Generalissimo Deodoro Ave, Belém, Pará 66055-240, Brazil

juarez@ufpa.br

Luiz Fábio Magno Falcão

Health and Biological Centre, State of Pará University, 2623 Perebeui St, Belem, Pará 66087-670, Brazil.

University of São Paulo, 455 Dr Arnaldo Ave, São Paulo, São Paulo 01246-903, Brazil

fabiofalcao@uepa.br

REFERENCES

1. Mbituyumuremyi A, Van Nuil JI, Umuhire J et al. Controlling hepatitis C in Rwanda: a framework for a national response. *Bull World Health Organ.* 2018;96(1):51-8.
2. Piselli P, Serraino D, Fusco M *et al.* Hepatitis C virus infection and risk of liver-related and non-liver-related deaths: a population-based cohort study in Naples, southern Italy. *BMC Infect Dis.* 2021; 21:667. <https://doi.org/10.1186/s12879-021-06336-9>
3. World Health Organization, Hepatitis C, Hepatitis C key facts, World Health Organization Media Centre, Geneva, Switzerland, 2022, <https://www.who.int/news-room/fact-sheets/detail/hepatitis-c>
4. Brazil Ministry of Health, 2021 Epidemiological Bulletin on Viral Hepatitis, Health Surveillance Secretariat, Department of STD, AIDS and Viral Hepatitis, Ministry of Health, Brasília, Brazil, 2021 (Portuguese), <http://www.aids.gov.br/pt-br/pub/2021/boletim-epidemiologico-hepatites-virais-2021>
5. Westbrook RH, Dusheiko G. "Natural History of Hepatitis C." *Journal of Hepatology.* 2014; 61: S58-68.
6. Motawi TMK, Sabry D, Shehata NI, et al. Impact of FOXP1 rs2687201 genetic variant on the susceptibility to HCV-related hepatocellular carcinoma in Egyptians. *J Biochem Mol Toxicol.* 2021;16: e22965.
7. Yousaf A, Ghafoor A, Fátima N, et al. Gender-specific frequency distribution of hepatitis C virus genotypes in Punjab Province, Pakistan: a clinically significant cross-sectional study. *Cureus.* 2021; 13: e17480.
8. Wang B, Krüger L, Machnowska P, et al. Characterization of a hepatitis C virus genotype 1 divergent isolate from an HIV-1 coinfecting individual in Germany assigned to a new subtype 1o. *Viol J.* 2019;16: 2803-04.
9. Strauss E. Natural History of Hepatitis C - Progression Factors. Prognostic assessment of chronic hepatitis C. *Treatise on Viral Hepatitis and Associated Diseases.* São Paulo: Atheneu. 2013: 453-470.
10. Cheng XD, Xu HF, Wei F, et al. Genotype analysis of hepatitis C virus in Heilongjiang Province, China. *Medicine.* 2021;100: e25203.
11. Codeço CT, Dal'Asta AP, Rorato AC. et al. Epidemiology, Biodiversity, and Technological Trajectories in the Brazilian Amazon: From Malaria to COVID-19. *Front Public Health.* 2021; 13: 647754. doi: 10.3389/fpubh.2021.647754.
12. Alves de Oliveira, BF, Bottino, MJ, Nobre, P. et al. Deforestation and climate change are projected to increase heat stress risk in the Brazilian Amazon. *Commun Earth Environ.* 2021; 2. Doi: 10.1038/s43247-021-00275-8

13. Azhar MJ, Khalid N, Azhar S, et al. Study of the Effect of Different Hepatitis C Virus Genotypes on Splenomegaly. *Cureus*. 2020; 3112: e10164.
14. Fonseca JCF, Brazil LM. Hepatitis C virus infection in the Brazilian Amazon region. *Rev. Soc. Bras. Med Trop*. 2004; 37: 1-8.
15. Nicolau S, Medeiros AS, Santos MCA, Montarroyos JS. Epidemiological profile of hepatitis B in a regional health center in Recife, *Rev. Saúde Col. UEFS*. 2017; 7: 30-35.
16. Moia LJMP, Amaral ISA, Farias AJL, Silva MMA. Hepatitis C virus infection in a referral hospital in the Brazilian Amazon. *Rev Para Med*. 2014; 28:19-25
17. Suarez-Kurtz G, Paula DP, Struchiner CJ. Pharmacogenomic implications of population admixture: Brazil as a model case. *Pharmacogenomics*. 2014; 15: 209-19. doi: 10.2217/pgs.13.238.
18. Salzano FM and Sans M. Interethnic admixture and the evolution of Latin American populations. *Genetics and Molecular Biology* [online]. 2014; 37 suppl 1:151-170. Doi:10.1590/S1415-47572014000200003.
19. Sá LC, Araújo TME, Griep RH, Campelo V, Monteiro CFS. Hepatitis C seroprevalence and associated factors in crack users. *Rev Latinoam Enferm*. 2013; 21:1195-202.
20. Confalonieri UEC, Margonari C, Quintão AF. Environmental change and the dynamics of parasitic diseases in the Amazon. *Acta Tropica*. 2014; 129: 33-41. Doi:10.1016/j.actatropica.2013.09.013.
21. Sawada L, Pinheiro ACC, Locks D, et al. Distribution of hepatitis C virus genotypes among different exposure categories in the State of Pará, Brazilian Amazon. *Rev. Soc. Bras. Med. Trop*. 2011;44: 8-12.
22. Guimaraes VS, Melo TG, Ferreira RCD, Almeida SF, Martins LC. Prevalence of hepatitis C virus genotypes in the State of Pará, Brazil. *Rev. Soc. Bras. Med. Trop. Uberaba*, 2018; 51: 508-512.
23. Barbosa KMV, Moreira LVL, Oliveira CMA, et al. Hepatitis C in the 1980s: rescue of old "non-A and non-B" hepatitis cases from a hepatology service in the Amazon, Brazil. 2019. *Rev Pan Amaz Saude*, 2019; 10:1-11.
24. Olmedo G, Nagai M, Sugastti AL, et al. Epidemiological and molecular characterization of Hepatitis C in the population that went to the Central Laboratory of Public Health, Paraguay 2013-2018. *Rev. public salud Parag*. 2019; 9: 73-80.
25. Santos Ó, Gómez A, Vizcaíno V, Casas MC, Ramírez MDP, Olaya P. [Hepatitis C virus genotypes circulating in Colombia]. *Biomedica*. 2017 Jan 24; 37:22-27.

26. Duran A. Ç., Çetinaka ÖK, Sayiner AA, et al. Changes on Hepatitis C virus genotype distribution in Western Turkey: Evaluation of twelve-year data. *Turk J Gastroenterol*; 2020, 31: 128-135.
27. Døssing A, Røge BT, Madsen LG, et al. Hepatitis C virus elimination in Denmark. *Ugeskr Laeger*. 2020; 182: v08190481.
28. Petruzzello A, Sabatino R, Loquercio G, et al. Nine-year distribution pattern of hepatitis C virus (HCV) genotypes in Southern Italy. *PLoS One*. 2019; 14: e0212033.
29. Bruno S, Crosignani A, Maisonneuve P, Rossi S, Silini E, Mondelli MU. Hepatitis C virus genotype 1b as a major risk factor associated with hepatocellular carcinoma in patients with cirrhosis: a seventeen-year prospective cohort study. *Hepatology*. 2007 Nov;46(5):1350-6. doi: 10.1002/hep.21826. PMID: 17680653.
30. Kermani FR, Sharifi Z, Ferdowsian F, Paz Z, Zamanian M. Distribution of hepatitis C virus genotypes among chronic infected injecting drug users in Tehran, Iran. *Jundishapur J Microbiol*. 2013; 6: 265-8.
31. Carnaúba LAB, Pontes LR, Flavio LF, et al. Hepatitis C staging: elastography compared with liver biopsy. *Cad. Med. UNIFESO*. 2018; 1(1): 111-123.
32. Franco, KMV, Vieira WB, Dias ARN, et al. Doppler ultrasound: a non-invasive method used to diagnose and monitor patients with chronic hepatitis C. *Journal of Gastroenterology and Hepatology*. 2019; 35.
33. Lutz H, Gassler N, Tischendorf F, Trautwein C, Tischendorf J. Doppler ultrasound of liver blood flow for noninvasive assessment of liver fibrosis compared with liver biopsy and transient elastography. *Excavation. Dis. Sci*. 2012; 57: 2222–30.
34. Fahmy MI, Badran HM. Comparison of transient elastography with Doppler indices in predicting hepatitis C-induced liver fibrosis and cirrhosis. *The Egyptian J. Radio. Nucl. Med*. 2011; 42:111–7.
35. Mello FSF, Lima JMDC, Hyppolito EB, et al. Comparison of hepatic fibrosis degrees in chronic hepatitis C (HCC) measured by elastography and serology methods: ARFI and Fibroscanvs APRI and FIB4. *Rev. Med. UFC*. 2020; 60: 18-25.

Table 1 – Sociodemographic profile of patients with hepatitis C, Abaetetuba, Pará, Brazil (2019- 2020).

SOCIODEMOGRAPHIC PROFILE		n	%	P-Valor ⁽¹⁾
AGE				
Up to 59 years old	35	46.1%	0.491 ^{ns}	
> 60 years old	41	53.9%		
Sex				
Men	51	67.1%	0.004*	
Women	25	32.9%		
COLOR/RACE				
White	12	15.8%	0.000*	
Black	8	10.5%		
Dark-skinned	56	73.7%		
SCHOOLING				
Up to 9 years of study	31	40.8%	0.136ns	
> 9 years of study	45	59.2%		
Incomplete/complete higher education	0	0.0%		
MUNICIPALITY OF RESIDENCE				
Abaetetuba	71	93.4%	0.000*	
Others	5	6.6%		
AREA OF RESIDENCE				
Rural area	11	14.5%	0.000*	
Urban area	65	85.5%		
OCCUPATION				
Retired	29	38.2%	0.051ns	
Active work	47	61.8%		
MARITAL STATUS				
Married / Stable union	53	69.7%	0.001*	
Single/ Widower	23	30.3%		
POSSIBLE FORM OF TRANSMISSION				
Surgical	43	56.6%	0.000*	
Blood transfusion (Parenteral route)	46	60.5%		
Sexual Route	3	3.9%		
Not informed	20	26%		
CLINICAL FORM				
Acute	4	5.3%	0.000*	
Chronicle	53	69.7%		
Not filled	19	25.0%		

(1) Pearson's Chi-square (Wilks' G²) test for independence (p-value < 0.05).

*Significant Values; ns - Non-Significant Values.

Table 2 – Clinical profile of patients with hepatitis C according to the degree of liver fibrosis and the genotype of the hepatitis C virus, Abaetetuba, Pará, Brazil (2019-2020).

EOTIC 2020%			
CLINICAL PROFILE	n	%	p-value ⁽¹⁾
FIBROSIS CLASSIFICATION			
F0	8	10.5%	0.140ns
F1	14	18.4%	
F2	15	19.7%	
F3	8	10.5%	
F4	13	17.1%	
Exam not performed	14	18.4%	
Exam performed/not valid measures	4	5.3%	
ELASTOGRAPHY (RESULT IN KPA)			
1.2-6.2	11	14.5%	0.000*
6.2-11.2	32	42.1%	
11.2-16.2	6	7.9%	
16.2 ou mais	9	11.8%	
GENOTYPE OF HEPATITIS C VIRUS			
1	42	55.3%	0.001*
2	2	2.6%	
3	22	28.9%	
HEPATITIS C VIRUS SUBGENOTYPE			
1a	10	13.2%	0.000*
1b	32	42.1%	
3a	1	1.3%	
Not Subtyped	10	13.2%	

(1) Pearson's Chi-square (Wilks' G²) test for independence (p-value < 0.05).

*Significant Values; ns - Non-Significant Values.

Table 3 - Distribution of patients with hepatitis C according to the genotypes, sub-genotypes, and liver elastography test (Kpa), Abaetetuba, Pará, Brazil (2019-2020).

Genotypes/Subgenotypes			F0 (n = 8)		F1/F2 (n = 27)		F3/F4 (n = 20)		p-value (1)
Hepatitis C virus genotype	C	virus	n	%	n	%	n	%	
1			5	62.5%	15	55.6%	16	80.0%	
2			0	0.0%	1*	3.7%	0	0.0%	0.607 ^{ns}
3			3	37.5%	11	40.7%	4	20.0%	
Hepatitis C virus subgenotype	C	virus	F0 (n = 5)		F1/F2 (n = 15)		F3/F4 (n = 20)		
1a			3	60.0%	4	26.7%	3	15.0%	
1b			2	40.0%	9	60.0%	15	75.0%	0.002*
3a			0	0.0%	0	0.0%	1	5.0%	
Not subtyped			0	0.0%	2	13.3%	1	5.0%	

(1) Pearson's Chi-square (Wilks' G²) test for independence (p-value < 0.05).

*Significant Values; ns- Non-Significant Values.

FIGURE LEGEND

Figure: Hepatitis C virus genotypes, environmental factors and Liver fibrosis in Eastern Amazon

Legends: Subtype 1b, 1a and 3a are the most prevalent in Eastern Amazon, Subtype 1b is related to Advanced and Mild Liver fibrosis, the individuals with fibrosis have a risk to development of Liver cirrhosis of 20%. Associated to HCV infection, the environmental factors as ethnic diversity, social vulnerability, diversity of pathogens in Amazon region, and the risk behaviors increase the risk of worse prognostic.