

Article

Not peer-reviewed version

Passive dynamics of the head and forelimb in equine foetuses- an observational study.

[Carla M Lusi](#) * and [Helen M.S. Davies](#)

Posted Date: 28 February 2023

doi: 10.20944/preprints202302.0484.v1

Keywords: passive dynamics; equine; forelimb; head; lameness



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Passive Dynamics of the Head and Forelimb in Equine Foetuses- an Observational Study

Carla M Lusi * and Helen MS Davies

Department of Veterinary BioSciences, The University of Melbourne, Parkville, Victoria 3010, Australia

* Correspondence: h.davies@unimelb.edu.au

Abstract: Passive dynamics is an aspect of locomotion which is entirely dependent on the mechanical configuration and linkages of adjacent body segments. Tension distribution along mechanical linkages enables execution of movement patterns with reduced need for complex neurological pathways and may play a role in reestablishing postural stability following external disturbances. Here we demonstrate a uni-directional mechanical relationship between the equine forelimb and head, which may have implications for balance and forelimb loading in the horse. These observations suggest that forelimb and head movement coordination (observed in the horse during unrestrained locomotion) is largely controlled by the mechanical linkages between body segments, rather than being entirely dependent on neurological input as previously thought. This highlights the potential significance of research directed at investigating passively induced movements in understanding common locomotory patterns. Additionally, it suggests a mode of postural control which may provide instantaneous adjustments to postural disturbances, thus promoting rapid and efficient locomotion.

Keywords: passive dynamics; equine; forelimb; head; lameness

1. Introduction

In the existing literature, two main modes of postural stability and motor control are established, including reflexive neural feedback and anticipatory systemic changes of posture (Deliagina et al., 2006; Horak and Macpherson, 2010; Moritz and Farley, 2004). However, more recently, a third mode which places emphasis on the mechanical reactions and passive dynamics of linked segments in the body has begun to filter into the literature (Eng et al., 1996; Moritz and Farley, 2004; Patla and Prentice, 1995). It has been suggested that complex and skilled movements are simply and effectively controlled via the utilization of passive forces (Bernshteĩn, 1967) and that distribution of these forces (through anatomical or mechanical coupling) further simplifies postural and movement control by reducing the degrees of freedom the nervous system has to consider (Bernshteĩn, 1967; Stahl, 2010; Turvey, 1990).

In the horse, research into the role of passive dynamics in maintaining balance and locomotory patterns has mostly been approached from one of two directions. The first of these considers the passive interaction of joint segments through the storage and release of elastic energy (Alexander, 1984, 2002; Biewener, 1998; Cavagna et al., 1977; McGuigan & Wilson, 2003; Wilson et al., 2003). This has been most clearly demonstrated in the biomechanics of the forelimb (Swanstrom et al., 2004; Swanstrom et al., 2005; Wilson et al., 2003). The second approach relates to the role of visual, vestibular and somatosensory inputs in enabling segmental movement and contributing to whole body balance and spatial orientation (Dunbar, 2008; Fung & Macpherson, 1995; Gómez Álvarez, 2006; Roberts, 1978; Weishaupt et al., 2006). Ultimately, it has been demonstrated that there are mechanisms of motor control, which rely on a sophisticated interaction between labyrinth and neck reflexes and serve to stabilize positions of the head, neck, trunk and limbs in order to compensate for asymmetrical loads or forces.

Although both research directions have been useful in understanding forelimb and whole-body biomechanics in the horse, it is noted here that a functional link exists between the two research areas that is yet to be explored. Investigation of energy storage and release through biological spring systems is essentially limited to the mechanical connections of the musculotendinous unit. Hence, passive movements enabled through elastic energy storage are confined to the segments across which the musculotendinous unit spans. On the other hand, segmental mechanics through nervous input does focus on the movement and coordination across several body segments (such as the head, neck and trunk); yet it does not investigate the mechanical link between these segments.

Hence, it is the purpose of this present study to investigate the mechanical interaction between body segments in the horse which may contribute to the efficiency of locomotion. The relationship between head and forelimb movement presents as a prominent non-volitional movement pattern where passive forces between mechanically linked segments may dominate and exist independently of neurological input. Studies looking at head and limb movement synchronization in other species suggest that neural reflex mechanisms play a large part (Daanje, 1951; Hancock, 2010), in coordination of body segments; however, several other recent studies suggest that intersegmental dynamics may dominate in this role (Eng et al., 1996; Kubow and Full, 1999; McGeer, 1990a; Patla and Prentice, 1995; Valero-Cuevas et al., 2007). If this is the case, mechanical linkages would need to be independent of feeding and other neural based responses instituted by the main exteroceptors like the eyes and ears in order to avoid interference with body coordination and locomotion from such movements of the head and neck. Therefore, it is hypothesized here that a uni-directional mechanical relationship exists between the head and limb which may provide a simplified mechanism of movement coordination and therefore makes redundant the need for complex neurological reflex pathways from the distal forelimbs for such coordination.

2. Materials and methods

Sixteen foetal foal cadavers (estimated ages 80-340 days post coitum (Njaa, 2012)) of various breeds were collected from pregnant mares that had been euthanized for reasons not associated with this study. A foal that had died within 24 hours post partum was also collected and investigated as part of the study.

Foal cadavers that were not investigated immediately after collection were stored at -20°C until needed. In preparation for dissection, they were thawed at -4°C for 4-7 days depending on their size. Once thawed, foals were placed in lateral recumbency on a smooth, metal surface with their limbs positioned to approximate a standing position and their head placed in a natural position as would be observed when the head and neck are unrestrained. The surface under the head and neck of the horse was wet sufficiently with water to reduce friction and to allow freedom of movement prior to the commencement of video recording. A Sony NEX-5 digital camera was positioned on a tripod at a distance of approximately 1 metre above the foal to photograph and record observed movements between the forelimb and head. This camera remained in the same position for the entire investigation.

The primary investigator protracted and retracted the uppermost forelimb within its normal physiological range. Maximum protraction and retraction was taken as the most cranial and caudal points respectively to which the forelimb could be moved without influencing the position of the thorax and hindlimbs. This was done first with a natural degree of flexion allowed at the carpus (i.e. no force applied to the carpus), and then with the carpus maintained in maximum extension (obtained by placing pressure on the dorsal aspect of the carpus). Following this, the limbs were returned to the natural position described above and the head was moved repeatedly between a flexed and extended position (of various degrees) to observe any related movements produced in either the limbs or the trunk. The direction of each movement was noted and the presence or absence of effects distributing to other body segments with each movement of the head or limb.

Eleven of the seventeen subjects investigated were then skinned and the same sequence of forelimb and head movements were carried out. In three foetuses, the left forelimb and the cervical musculature on the left side was completely removed to reveal the skeleton and to observe any

thoracic and spinal movement accompanying protraction and retraction of the remaining right forelimb (Figure 1).



Figure 1. Foal placed in right lateral recumbency with the left forelimb and cervical musculature removed.

3. Results

Despite differences in ages, breeds and genders, a fundamental and consistent uni-directional mechanical relationship was observed between forelimb and head movement.

Movement of the head between a flexed and extended position did not produce any observable movement of the trunk or limbs. However, retraction of the forelimb caused simultaneous extension of the head in all foals, which was most noticeable with pressure applied to the dorsal aspect of the carpus (to maintain straightness of the limb). In subjects that were skinned and/or had the left forelimb and cervical musculature removed, forelimb retraction was also observed to cause substantial movement of the thoracic cage in the dorsocaudal direction (Figure 2). This, in turn, produced dorsiflexion of the thoracic and cervical spine.




Figure 2. Caudodorsal movement of the thorax and extension of the head with forelimb retraction observed in equine foetuses (highlighted in green. Grey outline illustrates skeletal positioning in natural unrestrained position).

In contrast, protraction of the forelimb resulted in simultaneous head flexion in all foals. Effects on the position of the thoracic cage were not as clearly evident as they were with forelimb retraction, however cervical-thoracic ventriflextion was clearly observed in specimens where cervical musculature had been removed (see supplementary video). The greatest range of axial skeletal movement related to protraction or retraction of the forelimb was related to the caudal cervical vertebrae and the cranial thorax.

There were no observable differences in any of these movement patterns when placing foetuses in either left or right lateral recumbency, nor were there any grossly observable differences noted after removal of skin (in foetuses where this was applicable).

4. Discussion

This study set out to investigate the hypothesis that the coordinated movement observed between the head and forelimb in the horse during locomotion is largely dependent on the mechanical connectivity between segments rather than being entirely dependent on neurological input. Results from this study confirmed this hypothesis, identifying synchronous uni-directional locomotory patterns between the forelimb and head in equine foetal foal cadavers. To our knowledge, this is the first example of anatomically related movements which cannot be directly explained by reflexive neural feedback mechanisms. Instead, it is concluded here that coordinated forelimb and head movement in the equine foetus is, to a large extent, achieved through passive dynamics.

The concept of intersegmental passive dynamics has most clearly been demonstrated in the study of robotics (Cham et al., 2002; Ikemata et al., 2009; McGeer, 1990a; McGeer, 1990b; Poulakakis et al., 2006). Mechanical bipedal robots (lacking any active control system) have been shown to settle into a steady state of walking which is maintained solely by the passive forces acting between segments (Ikemata et al., 2009; McGeer, 1990a; McGeer, 1990b). This implies a transferal of forces (ground reaction and gravitational) acting on the limb, which is dependent solely on the mechanical configuration of the robot.

Observations from this present study demonstrate a similar mechanically-dependent passive interaction between the head and forelimb of foetal foal cadavers, which correlates with synchronous head and forelimb movement patterns observed in the live horse at a walk. Application of an external force to the forelimb to replicate forelimb protraction (or retraction) produced simultaneous head flexion (or extension) without any other external forces being applied to the head, neck or any other body part. This suggests that, in equine foetal foal cadavers, forces applied to the distal forelimb in the sagittal plane are distributed through the fascial and musculoskeletal elements of the forelimb to the shoulder girdle, neck and head.

The findings from this study provide support for previous studies on humans, which have presented hypotheses on the role of passive dynamics in human locomotion but have been unable to irrefutably attribute their observations exclusively to the mechanical interaction of segments (Eng et al., 1996; Moritz and Farley, 2004; Patla and Prentice, 1995; Valero-Cuevas et al., 2007). By using foal cadavers, it has been possible in this study to identify the relative degree of synchronous head and forelimb movement, which is executed independently of neural input in the equine foetus or neonate. Additionally, it has enabled a clear demonstration of a passively generated uni-directional movement pattern that is functionally significant in the live adult horse and suggests that the study of passively related movements, which occur independently of any neural input, may assist in understanding equine biomechanics.

With this said, the importance of nervous system control is not undervalued. Research has demonstrated an interaction between labyrinth and neck reflexes which influences limb stance and muscle tone in order to stabilize the position of the trunk and maintain the animal's centre of mass (COM) (Roberts, 1973). In the horse, the mechanical relationship between the head and forelimbs may further contribute to the maintenance or repositioning of their COM. Dagg (1962) suggested that forward and backward head movement in horses throughout locomotion adjusts the horizontal COM position and mediates forward movement as a result. Our findings demonstrated that movement of the forelimbs can instigate this forward and backward head movement, suggesting that the passive relationship between the forelimb and head mediate weight transfer and perhaps minimize the muscular effort required for propulsive forward movement. In response to this, it is proposed that labyrinth and neck reflexes, originating from the central control system in the head, instigate postural changes, which serve to re-stabilize the COM with movement. Hence, it appears that a complex and dynamic coupling exists between the nervous and passive dynamic systems of the body.

This then brings us to the next question raised from the observations of this present study, which relates to the directional control of each of these systems. Consistent external forces applied to the forelimb caused consistent coordinated movements of the head, yet consistent external forces applied to the head produced no observable effects on the limbs. Thus, it is hypothesized here that the nervous system implements control through connections, which develop outwardly from the head,

whereas the mechanical system develops functional connections to the head, which originate from the musculoskeletal elements of the limb and trunk. It is understood that the peripheral nervous pathways employed with reflexive postural adjustments are direct extensions of the neural crest derived structures in the head and neck, thereby supporting the first part of this hypothesis. However, more detailed studies looking at the development of the head and limbs in various quadrupedal species is necessary to confirm the latter part of the hypothesis.

Finally, it is suggested here that findings and hypotheses formed from this study on foetal foal cadavers can be extrapolated to live adult horses. Research on the development of connective tissue in utero suggests that the anatomical relationships observed in the foetuses of this study are indeed applicable to adult horses, as the collagenous fascial and muscular connections responsible for connected movements arise from a structural network developed very early on in utero (Parry et al., 1978; Myers, 1997; Gaivão et al., 2014; Myers, 1997). Additionally, in cases where the foals underwent a freeze-thaw cycle, it was assumed that the passive movements observed in this study were not significantly altered by this. Past research investigating the effects of multiple freeze-thaw cycles on joint movement and tensile behaviour in soft connective tissue structures have demonstrated single freeze-thaw cycles to have no significant effect on the movement or biomechanical properties of these structures (Jung et al., 2011; Woo et al., 1986; Hongo et al., 2008). Further, the connected movements occurred in all foetuses and were completely consistent between foetuses of different stages of development and breeds. Hence, the mechanical relationship between the head and forelimb that are obvious in foetal foal cadavers is suggested here to be a valid representation of a passive dynamic system in the live adult horse, which likely has significant implications for the efficiency and control of movement.

Supplementary Information A video is linked to the online version of the paper. Three clips of equine foetuses are included which show: 1) no forelimb movement with forced head movement, 2) passive head movement with forced forelimb movement 3) passive movement of the thorax and head with forced right forelimb movement after removing the left side forelimb, hindlimb and cervical musculature.

Author Contributions: HMS Davies supervised the project and assisted with foetal movements and manuscript revision. CM Lusi carried out observations on foetal movements and wrote the manuscript. Correspondence and requests for materials should be addressed to clusi28@gmail.com

Acknowledgements: The authors would like to thank Brendan Kehoe for assistance with collection of foals.

Conflict of interest: The authors declare no conflicts of interest.

References

- Alexander, R. (1984). Elastic Energy Stores in Running Vertebrates. *Am. Zool.*, **24**(1), 85-94.
- Alexander, R. (2002). Tendon elasticity and muscle function. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, **133**(4), 1001-1011.
- Biewener, A.A. (1998). Muscle-tendon stresses and elastic energy storage during locomotion in the horse. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, **120**(1), 73-87.
- Bernshstein, N. (1967). *The co-ordination and regulation of movements*, Oxford, New York, Pergamon Press.
- Cavagna, G.A., Heglund, N.C., & Taylor C.R. (1977). Mechanical work in terrestrial locomotion, two basic mechanisms for minimising energy expenditure. *Am. J. Physiol.* **233**(5): 243-261
- Cham, J., Bailey, S., Clark, J., Full, R. & Cutkosky, M. (2002). Fast and robust: Hexapedal robots via shape deposition manufacturing. *Int. J. Robot. Res.*, **21**(10-11): 869-882.
- Daanje, A. (1951). On locomotory movements in birds and the intention movements derived from them. *Behaviour*, **3**, 48-98.
- Dagg, A. (1962). The role of the neck in the movements of the giraffe. *J. Mammol.*, **43**, 88-97.
- Deliagina, T., Orlovsky, G., Zelenin, P. & Beloozerova, I. (2006). Neural bases of postural control. *Physiology*, **21**, 216-225.
- Dunbar, D.C., Macpherson, J.M., Simmons, R.W., Zarcades, A. (2008). Stabilization and mobility of the head, neck and trunk in horses during overground locomotion: comparisons with humans and other primates. *J. Exp. Biol.*, **211**(24): 3889-3907.

- Eng, J., Winter, D. & Patla, A. (1996). Intralimb dynamics simplify recative control strategies during locomotion. *J. Biomech.*, **30**, 581-588.
- Fung, J. & Macpherson, JM. (1995). Determinants of postural orientation in quadrupedal stance. *J. Neuroscience* **15**(2): 1121-1131.
- Gaivão, M., Rambags, B., & Stout, T. (2014). Gastrulation and the establishment of the three germ layers in the early horse conceptus. *Theriogenology*, **82**(2): 354-365.
- Gómez Álvarez, C. B., Rhodin, M., Bobbert, M. F., Meyer, H., Weishaupt, M. A., Johnston, C., & van Weeren, P. R. (2006). The effect of head and neck position on the thoracolumbar kinematics in the unriden horse. *Equine Vet. J. Supp.*, **38**(36): 445-451.
- Hancock, J. (2010). *The mechanics of terrestrial locomotion and the function and evolutionary history of head-bobbing in birds*. PhD thesis, Ohio University.
- Hongo, M., Gay, R.E., Hsu, J-T., Zhao, K.D., Ilharreborde, B., Berglund, L.J., & An, K-N. (2008). Effect of multiple freeze-thaw cycles on intervertebral dynamic motion characteristics in the porcine lumbar spine. *J. Biomech.*, **41**(4): 916-920.
- Horak, F. & Macpherson, J. (2010). Postural Orientation and Equilibrium. *Comprehensive Physiology*. John Wiley & Sons, Inc.
- Ikemata, Y., Sano, A., Yasuhara, K. & Fujimoto, H. (2009). Dynamic effects of arc feet on the leg motion of passive walker. IEEE International Conference on Robotics and Automation, Kobe, Japan.
- Jung, H.J., Vangipuram, G., Fisher, M.B., Yang, G., Hsu, S., Bianchi, J., Ronholdt, C., & Woo, S.L.Y. (2011). The effects of multiple freeze-thaw cycles on the biomechanical properties of the human bone-patellar tendon-bone allograft. *J. Ortho. Res.*, **29**(8): 1193-1198.
- Kubow, T. & Full, R. (1999). The role of the mechanical system in control: a hypothesis of self-stabilization in hexapedal runners. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, **354**(1385): 854-862.
- McGeer, T. (1990a). Passive dynamic walking. *J. Robot. Res.*, **9**(2): 62-82.
- McGeer, T. (1990b). Passive walking with knees. Proc. of IEEE Robotics and Automation Conference Cincinnati, OH. 1640-1645.
- McGuigan, M., & Wilson, A. (2003). The effect of gait and digital flexor muscle activation on limb compliance in the forelimb of the horse *Equus caballus*. *J Exp Biol*, **206**(8): 1325-1336.
- Moritz, C. & Farley, C. (2004). Passive dynamics change leg mechanics for an unexpected surface during human hopping. *J. Appl. Physiol.*, **97**(4): 1313-1322.
- Myers, T. (1997). The 'anatomy trains'. *J. Bodyw. Mov. Ther.*, **1**(2): 91-101.
- Njaa, B.L. (2012). *Kirkbride's Diagnosis of Abortion and Neonatal Loss in Animals*. John Wiley & Sons Ltd.
- Parry, D. A. D., Craig, A. S., & Barnes, G. R. G. (1978). Tendon and ligament from the horse: an ultrastructural study of collagen fibrils and elastic fibres as a function of age. *Proc. R. Soc. Lond. B.*, **203**(1152): 293-303.
- Patla, A. & Prentice, S. (1995). The role of active forces and intersegmental dynamics in the control of limb trajectory over obstacles during locomotion in humans. *Exp. Brain Res.*, **106**(3): 499-504.
- Poulakakis, I., Papadopoulos, E. & Buehler, M. (2006). On the stability of passive dynamics of quadrupedal running with a bounding gait. *Int. J. Robot. Res.*, **25**(7): 669-687.
- Roberts, T. (1978). *Neurophysiology of Postural Mechanisms*, London, Butterworth & Co Ltd.
- Stahl, V. (2010). *A biomechanical analysis of the role of the crural fascia in the cat hindlimb*. PhD thesis, Georgia Institute of Technology and Emory University.
- Swanstrom, M., Stover, S., Hubbard, M., & Hawkins, D. (2004). Determination of passive mechanical properties of the superficial and deep digital flexor muscle-ligament-tendon complexes in the forelimbs of horses. *Am. J. Vet. Res.*, **65**(2), 188-197.
- Swanstrom, M., Zarucco, L., Stover, S., Hubbard, M., Hawkins, D., Driessen, B., & Steffey, E. (2005). Passive and active mechanical properties of the superficial and deep digital flexor muscles in the forelimbs of anesthetized Thoroughbred horses. *J. Biomech.*, **38**(3), 579-586.
- Turvey, M. (1990). Coordination. *Am. Psych.*, **45**(8): 938-953.
- Weishaupt, M., Wiestner, T., von Peinen, K., et al. (2006). Effect of head and neck position on vertical ground reaction forces and interlimb coordination in the dressage horse ridden at walk and trot on a treadmill. *Equine Vet. J. Supp.*, **38**(36): 387-392.
- Wilson, A., Watson, J., & Lichtwark, G. (2003). Biomechanics: A catapult action for rapid limb protraction. *Nature*, **421**(6918): 35-36.

Woo, S.L.Y., Orlando, C.A., Camp, J.F. & Akeson, W.H. (1986). Effects of postmortem storage by freezing on ligament tensile behaviour, *J. Biomech.*, **19**(5): 399-404.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.