

Article

Not peer-reviewed version

A Common Approach to Three Open Problems in Number Theory

[Apoloniusz Tyszk](#) *

Posted Date: 29 March 2023

doi: 10.20944/preprints202303.0420.v2

Keywords: Brocard's problem; Brocard-Ramanujan equation $x!+1=y^2$; composite Fermat numbers; composite numbers of the form $2^{2^n}+1$; Erdős' equation $x(x+1)=y!$



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

A Common Approach to Three Open Problems in Number Theory

Apoloniusz Tyszk

Faculty of Production and Power Engineering, University of Agriculture, Balicka 116B, 30-149 Kraków, Poland; rttyszka@cyf-kr.edu.pl

Abstract: The following system of equations $\{x_1 \cdot x_1 = x_2, x_2 \cdot x_2 = x_3, 2^{2^{x_1}} = x_3, x_4 \cdot x_5 = x_2, x_6 \cdot x_7 = x_2\}$ has exactly one solution in $(\mathbb{N} \setminus \{0, 1\})^7$, namely $(2, 4, 16, 2, 2, 2, 2)$. Hypothesis 1 states that if a system of equations $\mathcal{S} \subseteq \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, 7\}\} \cup \{2^{2^{x_j}} = x_k : j, k \in \{1, \dots, 7\}\}$ has at most five equations and at most finitely many solutions in $(\mathbb{N} \setminus \{0, 1\})^7$, then each such solution (x_1, \dots, x_7) satisfies $x_1, \dots, x_7 \leq 16$. Hypothesis 1 implies that there are infinitely many composite numbers of the form $2^{2^n} + 1$. Hypotheses 2 and 3 are of similar kind. Hypothesis 2 implies that if the equation $x! + 1 = y^2$ has at most finitely many solutions in positive integers x and y , then each such solution (x, y) belongs to the set $\{(4, 5), (5, 11), (7, 71)\}$. Hypothesis 3 implies that if the equation $x(x + 1) = y!$ has at most finitely many solutions in positive integers x and y , then each such solution (x, y) belongs to the set $\{(1, 2), (2, 3)\}$.

Keywords: Brocard's problem; Brocard-Ramanujan equation $x! + 1 = y^2$; composite Fermat numbers; composite numbers of the form $2^{2^n} + 1$; Erdős' equation $x(x + 1) = y!$


MSC: 11D61; 11D85

1. Composite numbers of the form $2^{2^n} + 1$

Let \mathcal{A} denote the following system of equations:

$$\{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, 7\}\} \cup \{2^{2^{x_j}} = x_k : j, k \in \{1, \dots, 7\}\}$$

The following subsystem of \mathcal{A}



has exactly one solution in $(\mathbb{N} \setminus \{0, 1\})^7$, namely $(2, 4, 16, 2, 2, 2, 2)$.

Hypothesis 1. If a system of equations $\mathcal{S} \subseteq \mathcal{A}$ has at most five equations and at most finitely many solutions in $(\mathbb{N} \setminus \{0, 1\})^7$, then each such solution (x_1, \dots, x_7) satisfies $x_1, \dots, x_7 \leq 16$.

Lemma 1. ([7], p. 109). For every non-negative integers x and y , $x + 1 = y$ if and only if $2^{2^x} \cdot 2^{2^x} = 2^{2^y}$.

Theorem 1. Hypothesis 1 implies that $2^{2^{x_1}} + 1$ is composite for infinitely many integers x_1 greater than 1.

Proof. Assume, on the contrary, that Hypothesis 1 holds and $2^{2^{x_1}} + 1$ is composite for at most finitely many integers x_1 greater than 1. Then, the equation

$$x_2 \cdot x_3 = 2^{2^{x_1}} + 1$$

has at most finitely many solutions in $(\mathbb{N} \setminus \{0, 1\})^3$. By Lemma 1, in positive integers greater than 1, the following subsystem of \mathcal{A}

$$\left\{ \begin{array}{l} 2^{2^{x_1}} = x_5 \\ 2^{2^{x_5}} = x_6 \\ 2^{2^{x_4}} = x_7 \\ x_6 \cdot x_6 = x_7 \\ x_2 \cdot x_3 = x_4 \end{array} \right. \quad \begin{array}{ccccc} x_1 & \xrightarrow{2^{2^{(\cdot)}}} & x_5 & \xrightarrow{+1} & x_4 = x_2 \cdot x_3 \\ & & \downarrow 2^{2^{(\cdot)}} & & \downarrow 2^{2^{(\cdot)}} \\ & & x_6 & \xrightarrow{\text{squaring}} & x_7 \end{array}$$

has at most finitely many solutions in $(\mathbb{N} \setminus \{0, 1\})^7$ and expresses that

$$\left\{ \begin{array}{lcl} x_2 \cdot x_3 & = & 2^{2^{x_1}} + 1 \\ x_4 & = & 2^{2^{x_1}} + 1 \\ x_5 & = & 2^{2^{x_1}} \\ x_6 & = & 2^{2^{2^{x_1}}} \\ x_7 & = & 2^{2^{2^{x_1}}} + 1 \end{array} \right.$$

Since $641 \cdot 6700417 = 2^{2^5} + 1 > 16$, we get a contradiction. \square

Most mathematicians believe that $2^{2^n} + 1$ is composite for every integer $n \geq 5$, see [2], p. 23.

Open Problem 1. ([3], p. 159). Are there infinitely many composite numbers of the form $2^{2^n} + 1$?

Primes of the form $2^{2^n} + 1$ are called Fermat primes, as Fermat conjectured that every integer of the form $2^{2^n} + 1$ is prime, see [3], p. 1. Fermat remarked that $2^{2^0} + 1 = 3$, $2^{2^1} + 1 = 5$, $2^{2^2} + 1 = 17$, $2^{2^3} + 1 = 257$, and $2^{2^4} + 1 = 65537$ are all prime, see [3], p. 1.

Open Problem 2. ([3], p. 158). Are there infinitely many prime numbers of the form $2^{2^n} + 1$?

2. The Brocard-Ramanujan equation $x! + 1 = y^2$

Let \mathcal{B} denote the following system of equations:

$$\{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, 6\}\} \cup \{x_j! = x_k : (j, k \in \{1, \dots, 6\}) \wedge (j \neq k)\}$$

The following subsystem of \mathcal{B}

$$\left\{ \begin{array}{l} x_1! = x_2 \\ x_1 \cdot x_1 = x_3 \\ x_2 \cdot x_2 = x_3 \\ x_3! = x_4 \\ x_4! = x_5 \\ x_5! = x_6 \end{array} \right. \quad \begin{array}{ccccccc} x_1 & & & & & & \\ \downarrow ! & \searrow \text{squaring} & & & & & \\ x_2 & \xrightarrow{\text{squaring}} & x_3 & \xrightarrow{!} & x_4 & \xrightarrow{!} & x_5 & \xrightarrow{!} & x_6 \end{array}$$

has exactly two solutions in positive integers, namely $(1, \dots, 1)$ and $(2, 2, 4, 24, 24!, (24!)!)$.


Hypothesis 2. If a system of equations $\mathcal{S} \subseteq \mathcal{B}$ has at most finitely many solutions in positive integers x_1, \dots, x_6 , then each such solution (x_1, \dots, x_6) satisfies $x_1, \dots, x_6 \leq (24!)!$.

Lemma 2. For every positive integers x and y , $x! \cdot y = y!$ if and only if

$$(x + 1 = y) \vee (x = y = 1)$$

Theorem 2. Hypothesis 2 implies that if the equation $x_1! + 1 = x_2^2$ has at most finitely many solutions in positive integers x_1 and x_2 , then each such solution (x_1, x_2) belongs to the set $\{(4, 5), (5, 11), (7, 71)\}$.

Proof. The following system of equations \mathcal{B}_1



is a subsystem of \mathcal{B} . By Lemma 2, in positive integers, the system \mathcal{B}_1 expresses that $x_1 = \dots = x_6 = 1$ or

$$\begin{cases} x_1! + 1 = x_2^2 \\ x_3 = x_1! \\ x_4 = (x_1!)! \\ x_5 = x_1! + 1 \\ x_6 = (x_1! + 1)! \end{cases}$$

If the equation $x_1! + 1 = x_2^2$ has at most finitely many solutions in positive integers x_1 and x_2 , then \mathcal{B}_1 has at most finitely many solutions in positive integers x_1, \dots, x_6 and Hypothesis 2 implies that every tuple (x_1, \dots, x_6) of positive integers that solves \mathcal{B}_1 satisfies $(x_1! + 1)! = x_6 \leq (24!)!$. Hence, $x_1 \in \{1, \dots, 23\}$. If $x_1 \in \{1, \dots, 23\}$, then $x_1! + 1$ is a square only for $x_1 \in \{4, 5, 7\}$. \square


It is conjectured that $x! + 1$ is a square only for $x \in \{4, 5, 7\}$, see [10], p. 297. A weak form of Szpiro's conjecture implies that the equation $x! + 1 = y^2$ has only finitely many solutions in positive integers, see [6].

3. Erdős' equation $x(x + 1) = y!$

Let \mathcal{C} denote the following system of equations:

$$\{x_i \cdot x_j = x_k : (i, j, k \in \{1, \dots, 6\}) \wedge (i \neq j)\} \cup \{x_j! = x_k : (j, k \in \{1, \dots, 6\}) \wedge (j \neq k)\}$$

The following subsystem of \mathcal{C}



has exactly three solutions in positive integers, namely $(1, \dots, 1)$, $(1, 1, 2, 2, 2, 2)$, and $(2, 2, 3, 6, 720, 720!)$.

Hypothesis 3. If a system of equations $\mathcal{S} \subseteq \mathcal{C}$ has at most finitely many solutions in positive integers x_1, \dots, x_6 , then each such solution (x_1, \dots, x_6) satisfies $x_1, \dots, x_6 \leq 720!$.

Theorem 3. Hypothesis 3 implies that if the equation $x_1(x_1 + 1) = x_2!$ has at most finitely many solutions in positive integers x_1 and x_2 , then each such solution (x_1, x_2) belongs to the set $\{(1, 2), (2, 3)\}$.

Proof. The following system of equations \mathcal{C}_1

$$\left\{ \begin{array}{l} x_1! = x_4 \\ x_5! = x_6 \\ x_4 \cdot x_5 = x_6 \\ x_2! = x_3 \\ x_1 \cdot x_5 = x_3 \end{array} \right. \quad \begin{array}{c} x_1 \xrightarrow{+1} x_5 \\ \text{or } x_1 = x_5 = 1 \end{array} \quad \begin{array}{c} x_1 \cdot x_5 = x_3 \\ \uparrow \\ x_2 \end{array}$$

$x_4 \xrightarrow{!} x_6$

is a subsystem of \mathcal{C} . By Lemma 2, in positive integers, the system \mathcal{C}_1 expresses that $x_1 = \dots = x_6 = 1$ or

$$\left\{ \begin{array}{l} x_1 \cdot (x_1 + 1) = x_2! \\ x_3 = x_1 \cdot (x_1 + 1) \\ x_4 = x_1! \\ x_5 = x_1 + 1 \\ x_6 = (x_1 + 1)! \end{array} \right.$$

If the equation $x_1(x_1 + 1) = x_2!$ has at most finitely many solutions in positive integers x_1 and x_2 , then \mathcal{C}_1 has at most finitely many solutions in positive integers x_1, \dots, x_6 and Hypothesis 3 implies that every tuple (x_1, \dots, x_6) of positive integers that solves \mathcal{C}_1 satisfies $x_2! = x_3 \leq 720!$. Hence, $x_2 \in \{1, \dots, 720\}$. If $x_2 \in \{1, \dots, 720\}$, then $x_2!$ is a product of two consecutive positive integers only for $x_2 \in \{2, 3\}$ because the following MuPAD program

```
for x2 from 1 to 720 do
x1:=round(sqrt(x2!+(1/4))-(1/2)):
if x1*(x1+1)=x2! then print(x2) end_if:
end_for:
```

returns 2 and 3. \square

The question of solving the equation $x(x + 1) = y!$ was posed by P. Erdős, see [1]. F. Luca proved that the *abc* conjecture implies that the equation $x(x + 1) = y!$ has only finitely many solutions in positive integers, see [4].

4. There is no hope for a hypothesis that is similar to Hypothesis 2 or 3 and holds for an arbitrary number of variables

Let $f(1) = 2$, $f(2) = 4$, and let $f(n + 1) = f(n)!$ for every integer $n \geq 2$. Let \mathcal{U}_1 denote the system of equations $\{x_1! = x_1\}$. For an integer $n \geq 2$, let \mathcal{U}_n denote the following system of equations:

$$\left\{ \begin{array}{l} x_1! = x_1 \\ x_1 \cdot x_1 = x_2 \\ \forall i \in \{2, \dots, n-1\} \ x_i! = x_{i+1} \end{array} \right. \quad \begin{array}{c} x_1 \text{ squaring} \\ \downarrow \\ x_2 \end{array} \xrightarrow{!} x_3 \cdots \xrightarrow{!} x_{n-1} \xrightarrow{!} x_n$$

For every positive integer n , the system \mathcal{U}_n has exactly two solutions in positive integers x_1, \dots, x_n , namely $(1, \dots, 1)$ and $(f(1), \dots, f(n))$. For a positive integer n , let Ψ_n denote the following statement: if a system of equations

$$\mathcal{S} \subseteq \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\} \cup \{x_j! = x_k : j, k \in \{1, \dots, n\}\}$$

has at most finitely many solutions in positive integers x_1, \dots, x_n , then each such solution (x_1, \dots, x_n) satisfies $x_1, \dots, x_n \leq f(n)$. The statements Ψ_n are discussed in [8,9].

Theorem 4. Every factorial Diophantine equation can be algorithmically transformed into an equivalent system of equations of the forms $x_i \cdot x_j = x_k$ and $x_j! = x_k$. It means that this system of equations satisfies a modified version of Lemma 4 in [7].

Proof. It follows from Lemmas 2–4 in [7] and Lemma 2. \square

The statement $\forall n \in \mathbb{N} \setminus \{0\} \Psi_n$ is dubious. By Theorem 4, this statement implies that there is an algorithm which takes as input a factorial Diophantine equation and returns an integer which is greater than the solutions in positive integers, if these solutions form a finite set. This conclusion is strange because properties of factorial Diophantine equations are similar to properties of exponential Diophantine equations and a computable upper bound on non-negative integer solutions does not exist for exponential Diophantine equations with a finite number of solutions, see [5].

References

1. D. Berend and J. E. Harmse, *On polynomial-factorial Diophantine equations*, Trans. Amer. Math. Soc. 358 (2006), no. 4, 1741–1779.
2. J.-M. De Koninck and F. Luca, *Analytic number theory: Exploring the anatomy of integers*, American Mathematical Society, Providence, RI, 2012.
3. M. Křížek, F. Luca, L. Somer, *17 lectures on Fermat numbers: from number theory to geometry*, Springer, New York, 2001.
4. F. Luca, *The Diophantine equation $P(x) = n!$ and a result of M. Overholt*, Glas. Mat. Ser. III 37 (57) (2002), no. 2, 269–273.
5. Yu. Matiyasevich, *Existence of noneffectivizable estimates in the theory of exponential Diophantine equations*, J. Sov. Math. vol. 8, no. 3, 1977, 299–311, <http://dx.doi.org/10.1007/bf01091549>.
6. M. Overholt, *The Diophantine equation $n! + 1 = m^2$* , Bull. London Math. Soc. 25 (1993), no. 2, 104.
7. A. Tyszka, *A hypothetical upper bound on the heights of the solutions of a Diophantine equation with a finite number of solutions*, Open Comput. Sci. 8 (2018), no. 1, 109–114, <http://doi.org/10.1515/comp-2018-0012>.
8. A. Tyszka, *Statements and open problems on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that contain informal notions and refer to the current knowledge on \mathcal{X}* , <http://arxiv.org/abs/1506.08655>, to appear in Creative Mathematics and Informatics 32 (2023), no. 2.
9. A. Tyszka, *Statements and open problems on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that refer to the current knowledge on \mathcal{X}* , Journal of Applied Computer Science & Mathematics 16 (2022), no. 2, 31–35, <http://doi.org/10.4316/JACSM.202202005>.
10. E. W. Weisstein, *CRC Concise Encyclopedia of Mathematics*, 2nd ed., Chapman & Hall/CRC, Boca Raton, FL, 2002.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.