

Article

Not peer-reviewed version

Developed Method. Interactions and Their Quantum Picture

[Piotr Ogonowski](#) *

Posted Date: 25 July 2023

doi: 10.20944/preprints202304.0011.v4

Keywords: field theory; Lagrangian mechanics; quantum mechanics; General Relativity; unification of interactions



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Developed Method. Interactions and Their Quantum Picture

Piotr Ogonowski

Kozminski University, Jagiellonska 57/59, Warsaw, 03-301, Poland; piotrogonowski@kozminski.edu.pl

Abstract: By developing the previously proposed method of combining continuum mechanics with Einstein Field Equations, it has been shown that the classic relativistic description, curvilinear description, and quantum description of the physical system may be reconciled. For a system with an electromagnetic field, the Lagrangian density equal to the invariant of the electromagnetic field was obtained, vanishing four-divergence of canonical four-momentum appears to be consequence of the Poynting theorem, and explicit form of one of gauges of the electromagnetic four-potential was introduced. The proposed method allows for further development with additional fields.

Keywords: field theory; Lagrangian mechanics; quantum mechanics; General Relativity; unification of interactions

1. Introduction

Over the past decades, great strides have been made in attempts to combine quantum description of interactions with General Relativity [1]. There are currently many promising approaches to connecting the Quantum Mechanics and General Relativity, including perhaps the most promising ones: Loop Quantum Gravity [2–4], String Theory [5–7] and Noncommutative Spacetime Theory [8,9].

A lot of work has also been done to clear up some challenges related to General Relativity and Λ CMD model [10]. An explanation for the problem of dark energy [11] and dark matter [12] is still being sought, and efforts are still being made to explain the origin of the cosmological constant [13–15].

The author also tries to bring his own contribution to the explanation of the above physics challenges, based on a recently discovered method, described in [16]. As this article will show, this method seems very promising and may help clarify at least some of the issues mentioned above. The author's method, similar to the approach presented in [17], also points to the essential connections between electromagnetism and General Relativity, however, the postulated relationship is of a different nature.

According to conclusions from [16], the description of motion in curved spacetime and its description in flat Minkowski spacetime with fields are equivalent, and the transformation between curved spacetime and Minkowski spacetime is known. This allows for a significant simplification of research, because the results obtained in flat Minkowski spacetime can be easily transformed into curved spacetime. The last missing link seems to be the quantum description.

In this article, the author will focus on developing the method proposed in [16] in such a way, as to obtain the convergence of the description with the description of quantum mechanics. In the first chapter, the Lagrangian density for the system will be derived, allowing to obtain the tensor described in [16]. These conclusions will be used later in the article to propose quantum description of the system.

The author uses the Einstein summation convention, metric signature $(+, -, -, -)$ and commonly used notations. In order to facilitate the analysis of the article, the key conclusions from [16] are quoted in the subsection below.

1.1. Short summary of the method

According to [16], stress-energy tensor $T^{\alpha\beta}$ for a system with electromagnetic field in a given spacetime, described by a metric tensor $g^{\alpha\beta}$ is equal to

$$T^{\alpha\beta} = \varrho U^\alpha U^\beta - (c^2 \varrho + \Lambda_\rho) (g^{\alpha\beta} - \xi h^{\alpha\beta}) \quad (1)$$

where ϱ_o is for rest mass density and

$$\varrho \equiv \varrho_o \gamma \quad (2)$$

$$\frac{1}{\xi} \equiv \frac{1}{4} g_{\mu\nu} h^{\mu\nu} \quad (3)$$

$$\Lambda_\rho \equiv \frac{1}{4\mu_o} \mathbb{F}^{\alpha\mu} g_{\mu\gamma} \mathbb{F}^{\beta\gamma} g_{\alpha\beta} \quad (4)$$

$$h^{\alpha\beta} \equiv 2 \frac{\mathbb{F}^{\alpha\delta} g_{\delta\gamma} \mathbb{F}^{\beta\gamma}}{\sqrt{\mathbb{F}^{\alpha\delta} g_{\delta\gamma} \mathbb{F}^{\beta\gamma} g_{\mu\beta} \mathbb{F}^{\alpha\eta} g^{\eta\zeta} \mathbb{F}^{\mu}_{\zeta}}} \quad (5)$$

where $\mathbb{F}^{\alpha\beta}$ represents electromagnetic field tensor, and where the stress-energy tensor for electromagnetic field, denoted as $Y^{\alpha\beta}$ may be presented as follows

$$Y^{\alpha\beta} \equiv \Lambda_\rho (g^{\alpha\beta} - \xi h^{\alpha\beta}) = \Lambda_\rho g^{\alpha\beta} - \frac{1}{\mu_o} \mathbb{F}^{\alpha\delta} g_{\delta\gamma} \mathbb{F}^{\beta\gamma} \quad (6)$$

Thanks to the proposed amendment to the continuum mechanics, in flat Minkowski spacetime occurs

$$\partial_\alpha U^\alpha = -\frac{d\gamma}{dt} \rightarrow \partial_\alpha \varrho U^\alpha = 0 \quad (7)$$

thus denoting four-momentum density as $\varrho U^\mu = \varrho_o \gamma U^\mu$, total four-force density f^μ acting in the system is

$$f^\mu \equiv \varrho A^\mu = \partial_\alpha \varrho U^\alpha U^\mu \quad (8)$$

Denoting rest charge density in the system as ρ_o and

$$\rho \equiv \rho_o \gamma \quad (9)$$

electromagnetic four-current J^α is equal to

$$J^\alpha \equiv \rho U^\alpha = \rho_o \gamma U^\alpha \quad (10)$$

The pressure p in the system is equal to

$$p \equiv c^2 \varrho + \Lambda_\rho \quad (11)$$

In the flat Minkowski spacetime, total four-force density f^α acting in the system calculated from $\partial_\beta T^{\alpha\beta} = 0$ is the sum of electromagnetic (f_{EM}^α), gravitational (f_{gr}^α) and the sum of remaining (f_{oth}^α) four-force densities

$$f^\alpha = \begin{cases} f_{EM}^\alpha \equiv -\Lambda_\rho \partial_\beta \xi h^{\alpha\beta} & (\text{electromagnetic}) \\ + \\ f_{gr}^\alpha \equiv (g^{\alpha\beta} - \xi h^{\alpha\beta}) \partial_\beta p & (\text{gravitational}) \\ + \\ f_{oth}^\alpha \equiv \frac{\varrho c^2}{\Lambda_\rho} f_{EM}^\alpha & (\text{sum of remaining forces}) \end{cases} \quad (12)$$

As was shown in [16], in curved spacetime ($g_{\alpha\beta} = h_{\alpha\beta}$) presented method reproduces Einstein Field Equations with an accuracy of $\frac{4\pi G}{c^4}$ constant and with cosmological constant Λ dependent on invariant of electromagnetic field tensor $\mathbb{F}^{\alpha\gamma}$

$$\Lambda = -\frac{\pi G}{c^4 \mu_0} \mathbb{F}^{\alpha\mu} h_{\mu\gamma} \mathbb{F}^{\beta\gamma} h_{\alpha\beta} = -\frac{4\pi G}{c^4} \Lambda_\rho \quad (13)$$

where $h_{\alpha\beta}$ appears to be metric tensor of the spacetime in which all motion occurs along geodesics and where Λ_ρ describes vacuum energy density.

It was also shown, that in this solution, Einstein tensor describes the spacetime curvature related to vanishing in curved spacetime four-force densities $f_{gr}^\alpha + f_{oth}^\alpha$.

The proposed solution allows to add additional fields while maintaining its properties.

2. Lagrangian density for the system

Since for the considered method the transition to curved spacetime is known (based on electromagnetic field tensor), the rest of the article will focus on the calculations in the Minkowski spacetime with presence of electromagnetic field, where $\eta^{\alpha\beta}$ represents Minkowski metric tensor.

Using a simplified notation

$$\frac{d \ln(p)}{d\tau} = U_\mu \partial^\mu \ln(p) = U_\mu \frac{\partial^\mu p}{p} \quad (14)$$

it can be seen that the four-force densities resulting from the obtained stress-energy tensor (12) in flat Minkowski spacetime can be written as follows

$$\begin{cases} f_{gr}^\alpha = (\eta^{\alpha\beta} - \xi h^{\alpha\beta}) \partial_\beta p = \frac{d \ln(p)}{d\tau} q U^\alpha - T^{\alpha\beta} \partial_\beta \ln(p) \\ f_{EM}^\alpha = \frac{\Lambda_\rho}{p} (f^\alpha - f_{gr}^\alpha) \\ f_{oth}^\alpha = \frac{qc^2}{p} (f^\alpha - f_{gr}^\alpha) \end{cases} \quad (15)$$

where f_{EM}^μ can also be represented in terms of electromagnetic four-potential and four-current. This means that to fully describe the system and derive the Lagrangian density, it is enough to find an explicit equation for the gravitational force or some gauge of electromagnetic four potential.

Referring to definitions from section 1.1 one may notice, that by proposing following electromagnetic four-potential $\mathbb{A}^\mu$

$$\mathbb{A}^\mu \equiv -\frac{\Lambda_\rho}{p} \frac{q_0}{\rho_0} U^\mu \quad (16)$$

one obtains electromagnetic four-force density f_{EM}^α in form of

$$f_{EM}^\alpha = J_\beta (\partial^\alpha \mathbb{A}^\beta - \partial^\beta \mathbb{A}^\alpha) = \frac{\Lambda_\rho}{p} \left(f^\alpha - \frac{d \ln(p)}{d\tau} q U^\alpha + qc^2 \partial^\alpha \ln(p) \right) \quad (17)$$

where J_β is electromagnetic four-current and where Minkowski metric property was utilized

$$U_\beta U^\beta = c^2 \quad \rightarrow \quad U_\beta \partial^\alpha U^\beta = \frac{1}{2} \partial^\alpha (U_\beta U^\beta) = 0 \quad (18)$$

and where the forces in the system can be described by the following equality

$$J_\beta (\partial^\alpha \mathbb{A}^\beta - \partial^\beta \mathbb{A}^\alpha) + q U_\beta \left(\partial^\beta \frac{qc^2}{p} U^\alpha - \partial^\alpha \frac{qc^2}{p} U^\beta \right) = q U_\beta (\partial^\beta U^\alpha - \partial^\alpha U^\beta) = f^\alpha \quad (19)$$

Comparing (15) and (17) it is seen, that introduced electromagnetic four-potential yields

$$0 = \left(T^{\alpha\beta} - qc^2 \eta^{\alpha\beta} \right) \partial_\beta \ln(p) \quad (20)$$

which is equivalent to imposing following condition on normalized stress-energy tensor

$$0 = \partial_\beta \left(\frac{T^{\alpha\beta}}{\eta_{\mu\gamma} T^{\mu\gamma}} \right) + \partial^\alpha \ln(\eta_{\mu\gamma} T^{\mu\gamma}) \quad (21)$$

and yields gravitational four-force density in Minkowski spacetime in form of

$$f_{gr}^\mu = q \left(\frac{d \ln(p)}{d\tau} U^\mu - c^2 \partial^\mu \ln(p) \right) \quad (22)$$

Now, one may show, that proposed electromagnetic four-potential leads to correct solutions.

At first, recalling the classical Lagrangian density [18] for electromagnetism one may show why, in the light of the conclusions from [16] and above, it does not seem to be correct and thus makes it difficult to create a symmetric stress-energy tensor [19]. The classical value of the Lagrangian density for electromagnetic field, written with the notation used in the article, is

$$-\mathcal{L}_{EMclassic} = \Lambda_\rho + \mathbb{A}^\mu J_\mu \quad (23)$$

In addition to the obvious doubt that by taking the different gauge of the four-potential $\mathbb{A}^\mu$ one changes the value of the Lagrangian density, one may notice, that with considered electromagnetic four-potential, such Lagrangian density is equal to

$$-\mathcal{L}_{EMclassic} = \Lambda_\rho - \frac{\Lambda_\rho q_0}{p \rho_0} U^\mu U_\mu \rho_0 \gamma = \Lambda_\rho - \frac{\Lambda_\rho qc^2}{p} = \frac{\Lambda_\rho^2}{p} \quad (24)$$

As it is seen, above Lagrangian density is not invariant under gradient over four-position and $\mathbb{A}^\mu$ and J_μ are dependent, what is not taken into account in classical calculation

$$\frac{\mathbb{A}^\alpha}{\mathbb{A}^\mu \mathbb{A}_\mu} = \frac{J^\alpha}{\mathbb{A}^\mu J_\mu} \quad (25)$$

Above yields

$$\frac{\partial \ln \left(\frac{1}{\sqrt{\mathbb{A}^\mu \mathbb{A}_\mu}} \right)}{\partial \mathbb{A}_\alpha} = - \frac{J^\alpha}{\mathbb{A}^\mu J_\mu} = \frac{p}{qc^2} \frac{J^\alpha}{\Lambda_\rho} \quad (26)$$

One may decompose

$$\ln \left(\frac{1}{\sqrt{\mathbb{A}^\mu \mathbb{A}_\mu}} \right) = \ln \left(\frac{p \rho_0}{\Lambda_\rho q_0 c} \right) = \ln(p) - \ln(\Lambda_\rho) - \ln \left(\frac{q_0 c}{\rho_0} \right) \quad (27)$$

and since $\frac{q_0 c}{\rho_0}$ are constants, one may simplify (26) to

$$\frac{\partial \ln(p)}{\partial \mathbb{A}_\alpha} - \frac{\partial \ln(\Lambda_\rho)}{\partial \mathbb{A}_\alpha} = \frac{p}{qc^2} \frac{J^\alpha}{\Lambda_\rho} = \frac{J^\alpha}{qc^2} + \frac{J^\alpha}{\Lambda_\rho} \quad (28)$$

Above yields

$$\frac{\partial \ln(\Lambda_\rho)}{\partial \mathbb{A}_\alpha} = - \frac{J^\alpha}{\Lambda_\rho} \quad (29)$$

which leads to the conclusion that Λ_ρ acts as the Lagrangian density for the system

$$\frac{\partial \Lambda_\rho}{\partial \mathbb{A}_\alpha} = \partial_\nu \left(\frac{\partial \Lambda_\rho}{\partial (\partial_\nu \mathbb{A}_\alpha)} \right) = -J^\alpha \quad (30)$$

thus stress-energy tensor for the system is equal to

$$T^{\alpha\beta} = \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial^\beta \mathbb{A}_\gamma - \Lambda_\rho \eta^{\alpha\beta} \quad (31)$$

In fact, the proof of correctness of the electromagnetic field tensor (noted as $Y^{\alpha\beta}$) allows to see this solution

$$f_{EM}^\beta = \partial_\alpha Y^{\alpha\beta} = J^\gamma \mathbb{F}^\beta_\gamma - \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial_\alpha \mathbb{F}^\beta_\gamma \quad (32)$$

what yields following property of electromagnetic field tensor

$$\mathbb{F}^{\alpha\gamma} \partial_\alpha \partial_\gamma \mathbb{A}^\beta = \mathbb{F}^{\alpha\gamma} \partial^\beta \partial_\alpha \mathbb{A}_\gamma \quad (33)$$

Using the above substitution, one may note

$$\partial_\alpha Y^{\alpha\beta} = \partial_\alpha \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial_\gamma \mathbb{A}^\beta - \partial_\alpha \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial^\beta \mathbb{A}_\gamma = \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial^\beta \partial_\alpha \mathbb{A}_\gamma - J^\gamma \partial_\gamma \mathbb{A}^\beta - \partial_\alpha \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial^\beta \mathbb{A}_\gamma \quad (34)$$

Therefore the invariance of Λ_ρ with respect to $\mathbb{A}_\alpha$ and $\partial_\nu \mathbb{A}_\alpha$ is both a condition on the correctness of the electromagnetic stress-energy tensor and on Λ_ρ in the role of Lagrangian density

$$0 = \frac{\partial \Lambda_\rho}{\partial (\partial_\alpha \mathbb{A}_\gamma)} \partial^\beta (\partial_\alpha \mathbb{A}_\gamma) + \frac{\partial \Lambda_\rho}{\partial \mathbb{A}_\gamma} \partial^\beta \mathbb{A}_\gamma = \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial^\beta \partial_\alpha \mathbb{A}_\gamma - J^\gamma \partial^\beta \mathbb{A}_\gamma = \partial_\alpha \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial^\beta \mathbb{A}_\gamma \quad (35)$$

what yields for (34) that

$$\partial_\alpha Y^{\alpha\beta} = J^\gamma \partial^\beta \mathbb{A}_\gamma - J^\gamma \partial_\gamma \mathbb{A}^\beta = f_{EM}^\beta \quad (36)$$

Equations (1), (6) and (31) yield

$$\frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial_\gamma \mathbb{A}^\beta = q U^\alpha U^\beta - \frac{c^2 q}{\Lambda_\rho} Y^{\alpha\beta} \quad (37)$$

what yields second representation of the stress-energy tensor

$$T^{\alpha\beta} = \frac{p}{qc^2} \cdot \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial_\gamma \mathbb{A}^\beta - \frac{\Lambda_\rho}{c^2} U^\alpha U^\beta = \frac{p}{qc^2} \partial_\gamma \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \mathbb{A}^\beta \quad (38)$$

After four-divergence, it gives additional expression for relation between forces and gives useful clues about the behavior of the system when transitioning to the description in curved spacetime.

3. Hamiltonian density and quantum picture

Since

$$\frac{\partial \Lambda_\rho}{\partial (\partial \mathbb{A}_\gamma / \partial x_\alpha)} = \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \rightarrow \frac{\partial \Lambda_\rho}{\partial (\partial \mathbb{A}_\gamma / \partial x_0)} = \frac{1}{\mu_0} \mathbb{F}^{0\gamma} \quad (39)$$

thus noting Hamiltonian density in terms of the Lagrangian density [20] (and also from (31)) one gets

$$\mathcal{H} = \left(\frac{\partial \Lambda_\rho}{\partial (\partial \mathbb{A}_\gamma / \partial x_0)} \right) \frac{\partial \mathbb{A}_\gamma}{\partial x_0} - \Lambda_\rho = \frac{1}{\mu_0 c} \mathbb{F}^{0\gamma} \frac{\partial \mathbb{A}_\gamma}{\partial t} - \Lambda_\rho \quad (40)$$

where conjugate momentum field Π^γ is

$$\Pi^\gamma \equiv \frac{1}{\mu_o} \mathbb{F}^{0\gamma} \quad (41)$$

As it is seen, above Hamiltonian density agrees with the classical Hamiltonian density for electromagnetic field [20] except that this Hamiltonian density was currently mainly considered for sourceless regions. According to the result above, this Hamiltonian density describes also gravitational interaction and other forces resulting from electromagnetism. Above, therefore, significantly simplifies quantum field theory equations [21–23].

At first one may notice, that in transformed (31)

$$-T^{\alpha 0} = -\frac{1}{\mu_o} \mathbb{F}^{\alpha\gamma} \partial_\gamma \mathbb{A}^0 + Y^{\alpha 0} \quad (42)$$

first row of electromagnetic stress-energy tensor $Y^{\alpha 0}$ is a four-vector, representing energy density of electromagnetic field and Poynting vector [24] - the Poynting four-vector. Therefore vanishing four-divergence of the $T^{\alpha 0}$ must represent Poynting theorem. Indeed, properties (33) and (35) provide such equality

$$0 = -\partial_\alpha T^{\alpha 0} = J_\gamma \mathbb{F}^{0\gamma} + \partial_\alpha Y^{\alpha 0} \quad (43)$$

which also shows, that for the constant total energy of the system, $T^{\alpha 0}$ is independent of the four-position.

Next, one may introduce auxiliary variable ε with the energy density dimension, defined as follows

$$c\varepsilon \equiv -\frac{1}{\mu_o} \mathbb{F}^{0\mu} \frac{d\mathbb{A}_\mu}{d\tau} \quad (44)$$

and comparing the result

$$-U_\beta T^{0\beta} = c\varepsilon + c\gamma \Lambda_\rho \quad (45)$$

between the two tensor definitions (31), (38) one may notice, that it must hold

$$-\frac{p}{qc^2} \cdot \frac{1}{\mu_o} \mathbb{F}^{0\mu} \partial_\mu \mathbb{A}^\beta = -\frac{p}{qc^2 \mu_o} \cdot \left(U^\beta \mathbb{F}^{0\mu} \partial_\mu \frac{\mathbb{A}^0}{c\gamma} + \frac{\mathbb{A}^0}{c\gamma} \mathbb{F}^{0\mu} \partial_\mu U^\beta \right) = \frac{\varepsilon}{c} U^\beta - \frac{p}{qc^2} \frac{\mathbb{A}^0}{\gamma c \mu_o} \mathbb{F}^{0\mu} \partial_\mu U^\beta \quad (46)$$

because the second component of above vanishes contracted with U_β , due to the property of the Minkowski metric (18). Therefore (31) also yields

$$-T^{0\beta} = \varepsilon \frac{qc}{p} U^\beta - \frac{c\varepsilon_o \mathbb{A}^0}{\gamma} \mathbb{F}^{0\mu} \partial_\mu U^\beta + Y^{0\beta} \quad (47)$$

where ε_o is electric vacuum permittivity.

It now may be concluded, that after integration of the $\frac{1}{c} T^{0\beta}$ with respect to the volume, the total energy transported in the isolated system should be the sum of the four-momentum and four-vectors describing energy transport related to fields. Therefore, by analogy with the Poynting four-vector $\frac{1}{c} Y^{0\beta}$, one may introduce a four-vector Z^β understood as its equivalent for the remaining interactions and rewrite (47) as

$$-\frac{1}{c} T^{0\beta} = \varrho_o U^\beta + Z^\beta + \frac{1}{c} Y^{0\beta} \quad (48)$$

where

$$Z^\beta \equiv \left(\frac{\varepsilon\gamma}{p} - 1 \right) \varrho_o U^\beta - \frac{\varepsilon_o \mathbb{A}^0}{\gamma} \mathbb{F}^{0\mu} \partial_\mu U^\beta \quad (49)$$

Above also drives to conclusion, that

$$\frac{1}{c\gamma}U_{\beta}T^{0\beta} = -\frac{\varepsilon}{\gamma} - \Lambda_{\rho} \quad (50)$$

may act as a Lagrangian density, used in the classic relativistic description based on four-vectors.

Based on the knowledge of the existing Lagrangians, a probable assumption can be made at this point that

$$\varepsilon = \varrho_0 c^2 \quad (51)$$

what would yield

$$-\frac{1}{c}U_{\beta}T^{0\beta} = \varrho_0 c^2 + \gamma\Lambda_{\rho} \quad (52)$$

$$Z^{\beta} = \rho_0 \mathbb{A}^{\beta} - \frac{\epsilon_0 \mathbb{A}^0}{\gamma} \mathbb{F}^{0\mu} \partial_{\mu} U^{\beta} \quad (53)$$

where $-\frac{\epsilon_0 \mathbb{A}^0}{\gamma} \mathbb{F}^{0\mu} \partial_{\mu} U^{\beta}$, due to its properties, may be associated with some description of the spin.

The above result should not be surprising, because considering the system with electromagnetic field and canonical four-momentum density as part of the stress-energy tensor, actually one of components of canonical four-momentum density should be related to electromagnetic four-potential and charge density.

Finally, one may define another gauge $\bar{\mathbb{A}}_{\gamma}$ of electromagnetic four-potential $\mathbb{A}_{\gamma}$ in following way

$$\bar{\mathbb{A}}^{\gamma} \equiv \mathbb{A}^{\gamma} - \partial^{\gamma} \mathbb{A}^{\beta} X_{\beta} = -X_{\beta} \partial^{\gamma} \mathbb{A}^{\beta} \quad (54)$$

and note, that

$$-X_{\beta} T^{0\beta} = \frac{1}{\mu_0} \mathbb{F}^{0\gamma} \bar{\mathbb{A}}_{\gamma} + X_{\beta} Y^{0\beta} \quad (55)$$

Since for constant total energy of the system

$$\partial^{\alpha} X_{\beta} T^{0\beta} = T^{0\alpha} \quad (56)$$

this brings two more important insights:

- $\frac{1}{c} X_{\beta} T^{0\beta}$ may play a role of the density of Hamilton's principal function,
- Hamilton's principal function may be expressed based on the electromagnetic field only, so in the absence of the electromagnetic field it disappears.

One may now summarize the above and propose a method for quantizing the system and for the description of the system with the use of classical field theory.

At first, it should be noted, that the above reasoning changes the interpretation of what the relativistic principle of least action means. As one may conclude from above, there is no inertial system in which no fields act, and in the absence of fields, the Lagrangian, the Hamiltonian and Hamilton's principal function vanish. Since the metric tensor (5) for description in curved spacetime depends on the electromagnetic field tensor only, it seems clear, that in the considered system, the absence of the electromagnetic field means actually the disappearance of spacetime and the absence of any action.

One may then introduce generalized, canonical four-momentum H^{μ} as four-gradient on Hamilton's principal function S

$$H^{\mu} \equiv -\frac{1}{c} \int T^{0\mu} d^3x \equiv -\partial^{\mu} S \quad (57)$$

where

$$-S \equiv H^{\mu} X_{\mu} \quad (58)$$

and where H^μ does not depend on four-position. One may also conclude from previous findings, that canonical four-momentum should be in form of

$$H^\mu = P^\mu + V^\mu \quad (59)$$

where

$$V^\mu \equiv \int Z^\mu + \frac{1}{c} Y^{0\mu} d^3x \quad (60)$$

and where four-momentum P^μ may be now considered as just "other gauge" of $-V^\mu$

$$\partial^\alpha P^\mu = -\partial^\alpha V^\mu \quad (61)$$

Since in the limit of the inertial system one gets $P^\mu X_\mu = mc^2\tau$, therefore, to ensure vanishing Hamilton's principal function in the inertial system, one may expect that

$$V^\mu X_\mu \equiv -mc^2\tau \quad (62)$$

what also yields vanishing in the inertial system Lagrangian L in form of

$$-\gamma L = U_\mu H^\mu = F^\mu X_\mu \quad (63)$$

where F^μ is four-force, and where all above yields

$$V^\mu U_\mu - F^\mu X_\mu = -mc^2 \quad (64)$$

thus

$$H^\mu H_\mu + m^2 c^2 = 2mF^\mu X_\mu + V^\mu V_\mu \quad (65)$$

In this picture, the Hamilton's principal function, generalized canonical four-momentum and Lagrangian vanish for inertial system as expected. Assuming that equation (52) is indeed correct and denoting

$$E_\Lambda \equiv - \int \Lambda_\rho d^3x \quad (66)$$

one obtains from (65)

$$H^\mu H_\mu - m^2 c^2 = V^\mu V_\mu - 2m\gamma E_\Lambda \quad (67)$$

To ensure compatibility with the equations of quantum mechanics it suffices to assume that right side of above equation vanishes

$$\gamma E_\Lambda = \frac{V^\mu V_\mu}{2m} \quad (68)$$

Thanks to above, by introducing quantum wave function Ψ in form of

$$\Psi \equiv e^{\pm iK^\mu X_\mu} \quad (69)$$

where K^μ is wave four-vector related to canonical four-momentum

$$\hbar K^\mu \equiv H^\mu \quad (70)$$

from (67) one obtains Klein-Gordon equation

$$\left(\square + \frac{m^2 c^2}{\hbar^2} \right) \Psi = 0 \quad (71)$$

which allows for further analysis of the system in the quantum approach, eliminating the problem of negative energy appearing in solutions [25].

The above representation allows the analysis of the system in the quantum approach, classical approach based on (42) and the introduction of a field-dependent metric in (5) for curved spacetime, which connects previously divergent descriptions of physical systems.

4. Conclusions and Discussion

As shown above, the proposed method summarized in section 1.1 seems to be very promising area of farther research. In addition to the earlier agreement with Einstein Field Equations in curved spacetime, by imposing condition (20) on normalized stress-energy tensor in flat Minkowski spacetime, one obtains consistent results, developing the knowledge of the physical system with electromagnetic field:

- Lagrangian density for the systems appears to be equal to $\mathcal{L} = \Lambda_\rho = \frac{1}{4\mu_0} \mathbb{F}^{\alpha\beta} \mathbb{F}_{\alpha\beta}$
- Stress-energy tensor may be simplified to familiar form: $T^{\alpha\beta} = \frac{1}{\mu_0} \mathbb{F}^{\alpha\gamma} \partial^\beta \mathbb{A}_\gamma - \Lambda_\rho \eta^{\alpha\beta}$
- $H^\beta \equiv -\frac{1}{c} \int T^{0\beta} d^3x$ acts as canonical four-momentum for the point-like particle
- The vanishing four-divergence of H^β turns out to be the consequence of Poynting theorem
- Some gauge of electromagnetic four-potential may be expressed as $\mathbb{A}^\mu = -\frac{\Lambda_\rho}{p} \frac{q_0}{\rho_0} U^\mu$
- Gravitational, electromagnetic and sum of other forces acting in the system may be expressed as shown in (15) where gravitational four-force is dependent on the pressure p in the system as shown in (22)

Obtained canonical four-momentum H^μ is equal to

$$H^\mu = P^\mu + V^\mu \quad (72)$$

where P^μ is four-momentum and V^μ describes the transport of energy due to the field. It is calculated as

$$V^\mu = q\mathbb{A}^\mu + \mathbb{S}^\mu + Y^\mu \quad (73)$$

where $\mathbb{A}^\mu$ is electromagnetic four-potential, $\mathbb{S}^\mu$ due to its properties, seems to be some description of the spin, and where Y^μ is the volume integral of the Poynting four-vector.

It also seems, that this approach allows to combine previously divergent methods of curvilinear and classic description with the quantum description. The proposed method should facilitate further research on the quantum picture of individual fields, significantly simplifying equations of the Quantum Field Theory and leading in natural way to Klein-Gordon equation (71).

Further analysis using the variational method may also lead to next discoveries regarding both the theoretical description of quantum fields and elementary particles associated with them, and the possibility of experimental verification of the obtained results.

Funding: The author did not receive support from any organization for the submitted work.

Data Availability Statement: Data sharing is not applicable to this article, as no datasets were generated or analyzed during the current study.

Conflicts of Interest: The author has no relevant financial or non-financial interests to disclose.

References

1. Padmanabhan, T. Gravity and quantum theory: Domains of conflict and contact. *International Journal of Modern Physics D* **2020**, *29*, 2030001.
2. Ashtekar, A.; Bianchi, E. A short review of loop quantum gravity. *Reports on Progress in Physics* **2021**, *84*, 042001.
3. Gambini, R.; Olmedo, J.; Pullin, J. Spherically symmetric loop quantum gravity: analysis of improved dynamics. *Classical and Quantum Gravity* **2020**, *37*, 205012.
4. Lewandowski, J.; Mäkinen, I. Scalar curvature operator for models of loop quantum gravity on a cubical graph. *Physical Review D* **2022**, *106*, 046013.

5. Manoukian, E.; Manoukian, E. String Theory. *100 Years of Fundamental Theoretical Physics in the Palm of Your Hand: Integrated Technical Treatment* **2020**, pp. 285–289.
6. Cano, P.A.; Ruipérez, A. String gravity in $D=4$. *Physical Review D* **2022**, *105*, 044022.
7. Guerrieri, A.; Penedones, J.; Vieira, P. Where is string theory in the space of scattering amplitudes? *Physical Review Letters* **2021**, *127*, 081601.
8. Novikov, O.O. P T-symmetric quantum field theory on the noncommutative spacetime. *Modern Physics Letters A* **2020**, *35*, 2050012.
9. Kupriyanov, V.G.; Vitale, P. A novel approach to non-commutative gauge theory. *Journal of High Energy Physics* **2020**, *2020*, 1–15.
10. Melia, F. A Candid Assessment of Standard Cosmology. *Publications of the Astronomical Society of the Pacific* **2022**, *134*, 121001.
11. Frusciante, N.; Perenon, L. Effective field theory of dark energy: A review. *Physics Reports* **2020**, *857*, 1–63.
12. Oks, E. Brief review of recent advances in understanding dark matter and dark energy. *New Astronomy Reviews* **2021**, *93*, 101632.
13. Demirtas, M.; Kim, M.; McAllister, L.; Moritz, J.; Rios-Tascon, A. Exponentially small cosmological constant in string theory. *Physical Review Letters* **2022**, *128*, 011602.
14. Firouzjahi, H. Cosmological constant problem on the horizon. *Physical Review D* **2022**, *106*, 083510.
15. Dymnikova, I. The Higgs Mechanism and Cosmological Constant Today. *Universe* **2022**, *8*, 305.
16. Ogonowski, P. Proposed method of combining continuum mechanics with Einstein Field Equations. *International Journal of Modern Physics D* **2023**, *2350010*, 15.
17. Danehkar, A. Electric-magnetic duality in gravity and higher-spin fields. *Frontiers in Physics* **2019**, *6*, 146.
18. Helrich, C.S. *The classical theory of fields: electromagnetism*; Springer Science & Business Media, 2012.
19. Brau, C.A. *Modern problems in classical electrodynamics*; Oxford univ. press, 2004.
20. Popławski, N. Classical physics: spacetime and fields. *arXiv preprint arXiv:0911.0334* **2009**.
21. Casini, H.; Magán, J.M. On completeness and generalized symmetries in quantum field theory. *Modern Physics Letters A* **2021**, *36*, 2130025.
22. Buchbinder, I.L.; Shapiro, I. *Introduction to quantum field theory with applications to quantum gravity*; Oxford University Press, 2021.
23. Meurice, Y. *Quantum Field Theory*; IOP Publishing, 2021.
24. Wylleman, L.; Costa, L.F.O.; Natário, J. Poynting vector, super-Poynting vector, and principal observers in electromagnetism and general relativity. *Classical and Quantum Gravity* **2021**, *38*, 165009.
25. Bussey, P. Improving our understanding of the Klein-Gordon equation. *arXiv preprint arXiv:2212.06878* **2022**.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.